The developmental changes in intrinsic and synaptic properties of prefrontal neurons enhance

local network activity from the second to the third postnatal week in mice

Katerina Kalemaki[1,2], Xiaxia Xu[3], Angeliki Velli[2,4], Ourania Christodoulou[4,5], Myrto Denaxa[5] Ileana

L. Hanganu-Opatz[3], Domna Karagogeos[1, 2*] and Kyriaki Sidiropoulou[2,4*]

[1]Department of Basic Science, Faculty of Medicine, University of Crete, Heraklion, Greece

[2]Institute of Molecular Biology and Biotechnology (IMBB), FORTH, Heraklion, Greece

[3]Developmental Neurophysiology, Institute of Neuroanatomy, University Medical Center

Hamburg-Eppendorf, Hamburg, Germany

[4]Department of Biology, University of Crete, Heraklion, Greece

[5]BSRC 'Alexander Fleming'

* co-corresponding author and equal senior contribution

Abstract

The prefrontal cortex (PFC) is characterized by protracted maturation that extends until

adulthood. The cellular mechanisms controlling the early development of prefrontal circuits are

still largely unknown. Our study delineates the developmental cellular processes that are on-

going in the mouse medial PFC (mPFC) during the second and third postnatal weeks and

compares them to those in the barrel cortex (BC). We show that from the second to the third

postnatal week in both brain areas due to increased spontaneous inhibitory postsynaptic

currents (sIPSC) and reduced spontaneous excitatory postsynaptic currents (sEPSC) ones. Drug

application suggested that the increased sEPSC frequency at mPFC at P10 is due to depolarizing

GABA$_A$ receptor (GABA$_A$R) function. Evoked IPSC recordings revealed that the GABA$_A$R in mPFC

at P10 exhibits significantly depolarizing reversal potential, compared to mPFC at P20.

Additionally, the K-Cl co-transporter 2 (KCC2) expression is decreased in the neonatal mPFC

compared to the juvenile reinforcing the idea that GABA$_A$R function in the neonatal mPFC is

depolarizing. Furthermore, increasing GABA$_A$R activity leads to increased basal synaptic

response and spontaneous activity of neonatal mPFC, but not BC. Moreover, the GABAergic

interneurons in the neonatal mPFC exhibit immature active properties. The above

developmental changes in the synaptic and intrinsic properties of mPFC neurons contribute to

the enhanced network activity in the juvenile mPFC, compared to neonatal mPFC, using a

computational modeling approach, a result that was then confirmed by in vivo recordings.

Abbreviations

BC: Barrel Cortex

INs: Interneurons

KCC2: K-Cl co-transporter 2

NKCC1: Na^+-K^+-Cl^- cotransporter 1

mPFC: medial Prefrontal Cortex

P10: Postnatal day 10

P20: Postnatal day 20

PNs: Pyramidal neurons

sEPSC: spontaneous excitatory postsynaptic current

sIPSC: spontaneous inhibitory postsynaptic current

YFP: yellow fluorescent protein

PV: parvalbumin

SST: somatostatin

Introduction

During early postnatal development, major events that contribute to cortical circuit

maturation include spatial and temporal patterns of electrical activity, intrinsically determined

cell death of early postnatal cortical interneurons and the depolarizing action of the

neurotransmitter GABA (γ-aminobutyric acid) (Khazipov et al. 2004; Khazipov and Luhmann

2006; Allene et al. 2008; Brockmann et al. 2011; Ben-Ari et al. 2012; Southwell et al. 2012;

Khazipov et al. 2013; Kirmse et al. 2015; Mòdol et al. 2019). The developmental switch of GABA

action from depolarizing to hyperpolarizing results from changes in chloride co-transporter

expression: Na^+-K^+-Cl^- cotransporter 1 (NKCC1), the Cl^- importer, is highly expressed early in

development, while the expression of the co-transporter KCC2, the Cl^- exporter, increases after

the first postnatal week (Ben-Ari 2001; 2002; Ben-Ari et al. 2007; Ben-Ari 2012). In addition, both

intrinsic properties of neurons and synaptic transmission undergo dramatic changes during early

postnatal development in a brain-area specific manner (Kriegstein et al. 1987; McCormick and

Prince 1987; Burgard and Hablitz 1993; Ramoa and McCormick 1994; Bahrey and Moody 2003).

Most studies on these developmental changes of the GABAergic system in the cortex have
focused on the primary somatosensory cortex, visual cortex and hippocampus (Bartolini et al.
2013; Le Magueresse and Monyer 2013; Hensch 2016). The barrel cortex (BC) is part of the
primary somatosensory cortex and is organized vertically in columns of cells associated with
sensory perception (Guo et al. 2014) and horizontally in six layers of distinct cell types. In
contrast, cortical association areas such as the prefrontal cortex (PFC) regulate cognitive
functions and do not directly control sensory information (Fuster 2015). Anatomically, the
mouse medial PFC (mPFC) is defined as the agranular part of frontal lobe, lacking the prominent
granular layer IV and is divided into distinct subregions, namely infralimbic, prelimbic and
cingulate cortex (Heidbreder and Groenewegen 2003; Van De Werd et al. 2010). The timeline of
mPFC development is delayed compared to other sensory cortices, such as BC (Casey et al. 2000;
Best and Miller 2010; Kolb et al. 2012, Rinetti-Vargas et al., 2017). From infancy to adulthood,
the developing mPFC undergoes considerable transcriptional, structural and functional changes
(Diamond 2005; Tsujimoto 2008; Kolb et al. 2012; Schubert et al. 2014; Kroeze et al. 2017).
While adolescent development of prefrontal circuitry and the underlying cellular mechanisms
have been addressed by a large number of studies, few investigations tackled the wiring
processes at earlier stages (Brockmann et al. 2011; Bitzenhofer et al. 2015; 2017), in particular
he mPFC between the second (neonatal) and third (juvenile) postnatal week. Here, we aim to fill
this gap by investigating the synaptic and intrinsic properties of neonatal and juvenile mPFC
neurons and comparing them to those in the BC, with a primary focus on the GABAergic system.
84

Methods
All *in vitro* experiments with mice took place under an experimental protocol approved by the
Research Ethics Committee and by our Institutional Animal Care and Use Committee that has
been approved by the Veterinarian Authorities Office (protocol license no. 93164). Experiments
were carried out by trained scientists and in accordance with the 3R principles. *In vivo*
experiments were performed in compliance with the German laws and the guidelines of the
European Community for the use of animals in research and were approved by the local ethical
committee (015/17, 015/18).
93

94 **Animals**

The *in vitro* experiments were performed on male C57Bl/6J; Lhx6Tg(Cre); R26R-YFP+/+ mice from animal facility of IMBB-FORTH were used. For the *in vivo* experiments, timed-pregnant C57BL/6J mice from the animal facility of the University Medical Center Hamburg-Eppendorf were used. The day of vaginal plug detection was defined as embryonic day (E)0.5, whereas the day of birth was defined as P0. The offspring of both sexes are used for *in vivo* electrophysiology recordings. All procedures were performed according to the European Union ethical standards outlined in the Council Directive 2010/63EU of the European Parliament on the protection of animals used for scientific purposes.

Mice were housed with their mothers and provided with standard mouse chow and water ad libitum, under a 12 h light/dark cycle (light on at 7:00 am) with controlled temperature (21°C). The P10 experimental group includes ages P9-P11 and the P20 group includes ages P19-P21, also referred to as second and third postnatal weeks or neonatal and juvenile, respectively. All efforts were made to minimize both the suffering and the number of animals used.

In vitro extracellular recordings

Slice Preparation*:* Mice (P10 and P20) were decapitated under halothane anesthesia. The brain was removed promptly and placed in ice cold, oxygenated (95% O_2 –5% CO_2) artificial cerebrospinal fluid (aCSF) containing (in mM): 125 NaCl, 3.5 KCl, 26 $NaHCO_3$, 1 $MgCl_2$ and 10 glucose (pH = 7.4, 315 mOsm/l). The brain was blocked and glued onto the stage of a vibratome (Leica, VT1000S). Rostrocaudal coronal slices (400 μm thick) containing either the mPFC (prefrontal cortex) or the BC (barrel cortex) region were selected and transferred to a submerged chamber, which was continuously superfused with oxygenated (95% O_2 –5% CO_2) aCSF containing (in mM): 125 NaCl, 3.5 KCl, 26 $NaHCO_3$, 2 $CaCl_2$, 1 $MgCl_2$ and 10 glucose (pH = 7.4, 315 mOsm/l) at room temperature (RT). The slices were allowed to equilibrate for at least 1 h in this chamber before recordings began. Slices were then transferred to a submerged recording chamber, continuously superfused with oxygenated (95% O_2 –5% CO_2) aCSF (same constitution as the one used for maintenance of brain slices) at RT during recordings.

Data Acquisition: Electrophysiological recordings were performed in all experimental groups under the same conditions as described below. Pulled glass micropipettes were filled with NaCl (2M) and placed in layers II/III of PFC and BC. Platinum/iridium metal microelectrodes (Harvard apparatus United Kingdom, 161 Cambridge, United Kingdom) were placed on layer II/III of the mPFC and the BC, about 300 μm away from the 1MΩ recording electrode, and were used to

evoke field excitatory postsynaptic potentials (fEPSPs). Local field potentials (LFPs) were amplified using a headstage with selectable high pass filter of 30 Hz to remove any offsets coupled to a Dagan BVC-700A amplifier, amplified 100 times and low-pass filtered at 1-kHz. A notch filter was used to eliminate line frequency noise at 30 Hz. Signals were digitized using the ITC-18 board (InstruTech, Inc.) on a PC with custom-made procedures in IgorPro (Wavemetrics, Inc.) and stored on a PC hard drive. All voltage signals were collected at a sampling frequency of 100 kHz (fs = 100 kHz).

For evoked fEPSPs, the electrical stimulus consisted of a single square waveform of 100 µs duration given at intensities of 0.1– 0.3 mA (current was increased from 0.1 mA to 0.3 mA, with 0.1 mA steps) generated by a stimulator equipped with a stimulus isolation unit (World Precision Instruments, Inc.). The effect of $GABA_AR$ activation was investigated by bath application of 2 µM Diazepam ($GABA_AR$ agonist). Diazepam was acquired from the Pharmacy of the University General Hospital in Heraklion as a 5 mg/ml solution and was diluted in aCSF during recordings. Other drugs used include CNQX (10µM), AP5 (50µM) and bumetanide (10µM) (Tocris). For spontaneous activity recordings, 20 5-sec recordings were acquired without any stimulation.

Data Analysis: Data were analyzed using custom-written procedures in IgorPro software (Wavemetrics, Inc.). No additional high-pass filters were applied to the raw data. For evoked recordings, the peak values of the fEPSP were measured using the minimum value of the synaptic response (4–5 ms following stimulation) and were compared to the baseline value prior to stimulation. Both parameters were monitored in real- time in every experiment. A stimulus–response curve was then plotted using stimulation intensities between 0.1 and 0.8 mA, in 0.1 mA steps. For each different intensity level, two traces were acquired and averaged.

To identify spontaneous activity events, the acquired spontaneous activity voltage signals of 5 seconds' duration were decimated (down-sampled) by a factor of 10 and the standard deviation σ_b of background signal was calculated in the 'quiet' part of each voltage response trace. To identify the 'quiet' period, each 5sec trace was split into 100msec increments and the range of voltage deflection was computed in each increment. The 'quiet' part of the LFP trace was the 100msec increment with the smallest σ_b value. As a spontaneous event, any voltage response larger than $4 \cdot \sigma_b$ was identified. We calculated the frequency of spontaneous events by measuring the number of spontaneous events divided by the duration of the trace (5s). The frequency was calculated in 20 consecutive 5-sec traces and then averaged for each condition

animal. The spontaneous events do not correspond to spiking of individual neurons, they rather reflect population spikes.

In vitro patch-clamp recordings

Slice Preparation: Mice were decapitated under halothane anesthesia. The brain was removed immediately and coronal slices of mPFC and BC (300–350 μm thick), using a vibratome (Leica, VT1000S, Leica Biosystems) were prepared from mice at the ages of P10 and P20 in ice-cold oxygenated (95% O_2 - 5% CO_2) modified choline-based aCSF (in mM) 0.5 $CaCl_2$, 7 mM $MgSO_4$; NaCl replaced by an equimolar concentration of choline). Slices were incubated for 30min at 32°C in an oxygenated normal aCSF containing (in mM): 126 NaCl, 3.5 KCl, 1.2 NaH_2PO_4, 26 $NaHCO_3$, 1.3 $MgCl_2$, 2.0 $CaCl_2$, and 10 D-glucose, pH 7.4, 315 mOsm/l. Slices were allowed to equilibrate for at least 30 min at RT before being transferred to the recording chamber. During recordings, slices were perfused at a rate of 4 ml/min with continuously aerated (95% O_2-5% CO_2) normal aCSF at RT.

Data Acquisition: Neurons were impaled with patch pipettes (5–7 MΩ) and recorded in the whole-cell configuration, either in the current-clamp or voltage-clamp mode. For current-clamp experiments, the composition of the intracellular solution was: 130 mM K-MeSO$_4$, 5 mM KCl, 5 mM NaCl, 10 mM HEPES, 2.5 mM Mg-ATP, and 0.3 mM GTP, 265–275 mOsm, pH 7.3. For voltage-clamp experiments, the composition of the intracellular solution was: 120 mM Cs-gluconate, 20mM CsCl, 0.1 mM $CaCl_2$, 1 mM EGTA, 0.4 mM Na-guanosine triphosphate, 2mM Mg-adenosine triphosphate, 10 mM HEPES. No correction from liquid junction potential was applied between the pipette and the aCSF. Whole-cell measurements were low-pass filtered at 5 kHz using an Axopatch 200B amplifier (Molecular Devices, Inc). Recordings were digitized with the ITC-18 board (Instrutech, Inc) on a PC using custom-made codes in IgorPro (Wavemetrics, Inc). All signals were collected at a sampling frequency of 20kHz.

Data Analysis: Data were analyzed using custom-written codes in IgorPro software (Wavemetrics, Inc.). For passive membrane properties, the resting membrane potential (RMP, mV) was measured within 3 min after establishing the whole-cell configuration, and monitored throughout the experiment. To measure input resistance, a 500ms step-pulse protocol was used with current stimulation from -200pA to +50pA. The input resistance (R_{in}, MΩ) was measured by plotting the steady-state voltage deflection in an I-V plot and calculating the slope of the best fit line curve ($R_{in}=V/I$). The τ_m (membrane time constant, ms) was obtained by fitting a single

exponential curve to the voltage deflection at -50pA, and the membrane capacitance (C_m) was

calculated using the formula $C_m = \tau_m/R_{in}$. In addition, the number of spikes generated in response

to a 500ms step-pulse range from +100pA to +300pA was measured.

To measure action potentials (APs) properties, we applied small supra-threshold 5ms step-pulse

currents to the cell from -65mV. The active properties were measured at the minimum current

stimulation (Rheobase, pA) that generated an AP. The AP threshold (mV) was calculated by

taking the first derivative of the voltage trace, defining a threshold and identifying the voltage

level at that time point. The rate of rise of the AP (dV/dt, mV/ms) was the maximum value of

that first derivative trace. The AP amplitude (mV) was defined as the voltage difference between

AP threshold and AP peak. The AP duration (ms) was calculated by the full width of the waveform

at the half maximal amplitude (half-width). The fast afterhyperpolarization (fAHP) minimum

(mV) was defined as the minimum voltage right after the AP. The fAHP amplitude (mV) was

calculated as the difference between the AHP minimum and the AP threshold. The fAHP time

(ms) was defined as the time duration from the time point of AP threshold to the fAHP minimum.

Voltage-clamp/spontaneous IPSC and EPSCs

The composition of our intracellular solution resulted in chloride reversal potential of -60mV and

Na^+/K^+ reversal potential of +10mV. This allowed for measurements of spontaneous excitatory

postsynaptic currents (sEPSCs) to be recorded at -60mV and of spontaneous inhibitory

postsynaptic currents (sIPSCs) to be recorded at +10mV. Automatically selected events were

subsequently visually monitored to discard erroneously included noise. The events showing only

single peaks were selected for kinetics analysis. All currents detected from every single neuron

were averaged. The peak amplitude was calculated as the maximum current value. The time

constant of the decay phase was detected by curve fitting with a single exponential decay

function.

Evoked IPSC recordings: To investigate eIPSCs and the reversal potential, a stimulating electrode

was positioned 100μm away from the recording electrode at the border between layer I and

layer II. The current used for stimulation was adjusted to give a 100-300pA response at -70mV.

For p10 recordings, the current used was 0.3mA while for p20 recordings the current used was

0.1mA. Recordings were taken at different voltage steps, from -80mV to +40mV in the presence

of CNQX (20μM) and AP5 (50μM).

***In vivo* extracellular recordings**

Surgery: Multisite extracellular recordings were performed in the PFC of P8–P10 (n=13) and P20–
P23 (n=14) C57/BL mice with both sexes. Mice were under urethane anesthesia (intraperitoneal
injection, 1 mg/g body weight; Sigma-Aldrich) before surgery. The bone over the mPFC (0.8 mm
anterior to bregma, 0.1–0.5 mm right to the midline) was carefully removed. One-shank
electrodes with 4 recording sites (0.4–0.8 MΩ impedance, 100 μm spacing, NeuroNexus) was
inserted into PFC at a depth of 1.9 mm from the skull surface. Electrodes were labelled with DiI
(1,1'-dioctadecyl-3,3,3',3'-tetramethyl indocarbocyanine; Invitrogen) to confirm their position
after histological assessment post-mortem. One silver wire was inserted into the cerebellum to
serve as ground and reference electrode.

Data Acquisition: A recovery period of 10 min following the insertion of electrodes before
acquisition of data was provided. Data acquired during the first 30 min of recording were used
for analysis to ensure similar state of anesthesia in all investigated pups. Extracellular signals
were bandpass filtered (0.1 Hz to 5 kHz) and digitized (32 kHz) with a multichannel extracellular
amplifier (Digital Lynx SX, Neuralynx) and the Cheetah acquisition software (Neuralynx).

Data Analysis: Data were imported and analyzed off-line using custom-written tools in MATLAB
software version 7.7 (MathWorks). Multiple unit activity (MUA) is detected when negative
deflections exceeding five times the SD of the bandpass filtered (500–5000 Hz) signals. Single
unit activity (USA)- was detected and clustered using klusta (Rossant et al., 2016) and manually
curated using phy (https://github.com/cortex-lab/phy). Data were imported and analyzed using
custom-written tools in the MATLAB. Burst activity was detected by using the method from
(Gorin, Tsitoura et al. 2016). First, for each unit, Poisson distribution, which assumed random
independent spiking, was defined. The single parameter λ of Poisson distribution is the average
firing rate of the unit. Second, the median interspike interval (ISI) value of the Poisson
distribution was derived. Third, we defined a burst activity as a sequence of ≥4 consecutive
spikes separated by intervals smaller than the median ISI. Last, the percentage of the burst
activity of a single unit is calculated by 100* (the number of the burst activity / the number of
the spikes). A unit was classified as "burst unit" if >50% of all spikes occurred in bursts.

Immunohistochemistry

Mice at the age of P10 and P20 were perfused with 4% paraformaldehyde, followed by fixation
with the same solution for 1h at 4∘C, followed by cryoprotection and preparation of 12 μm
cryostat sections as previously described[88]. Primary antibodies used were rat monoclonal anti-

GFP (Nacalai Tesque, Kyoto, Japan, 1:5000), rabbit polyclonal anti-GFP (1:500; Minotech Biotechnology, Heraklion, Greece) and rabbit polyclonal anti-parvalbumin (PV) (Swant, Bellinzona, Switzerland; 1:2000. Secondary antibodies used were goat anti-rat-Alexa Fluor-488, goat anti-rabbit Alexa Fluor-488, and goat anti-rabbit-Alexa Fluor-555 (Molecular Probes, Eugene, OR, United States, 1:800). Images were obtained with a confocal microscope (Leica TCS SP2, Leica, Nussloch, Germany). For each age group (P10, P20), 2-4 10μm-thick sections from each mouse brain were selected, all including the mPFC and BC.

RNA *In Situ* Hybridization

Non-radioactive *in situ* hybridization experiments were performed on cryostat sections (12μm thick, see immunochemistry) according to the protocol described (Schaeren-Wiemers and Gerfin-Moser 1993). Riboprobe was prepared by *in vitro* transcription and was specific Somatostatin (SST) (Liodis et al. 2007).

Nissl Staining

Cryostat sections (12μm thick, see immunochemistry) were incubated in 1:1 100% ethanol:chloroform overnight at RT. Then, sections were rehydrated for 1 min in 100%, 95% ethanol solutions and dH$_2$O at RT, followed by a 10-min incubation in 0.1% cresyl violet solution at 50°C. Sections were then dehydrated with dH$_2$O, 95%, 100% ethanol and xylene for 5 min and coverslipped with permount. Images from whole sections were obtained in 5× magnification of a light microscope (Axioskop 2FS, Carl Zeiss AG, 268 Oberkochen, Germany) and merged using Adobe Photoshop CC 2015, Adobe Systems, Inc.

Analysis for Immunochemistry, *in situ* hybridization and Nissl staining

Images taken from Nissl staining slices were analyzed with Matlab, using a custom-made algorithm, which was double checked with hand-counting (Konstantoudaki et al. 2016; Chalkiadaki et, 2019). Images from immunochemistry and in situ hybridization experiments, were hand counted using ImageJ.

Western blots

Mice were decapitated following cervical dislocation, the brain was quickly removed, placed in ice cold PBS (phosphate-buffered saline) and then positioned on a brain mould, where 1.5 mm

slices were taken containing the mPFC and BC. The slices were placed on dry ice, and the prelimbic area of mPFC was dissected out and stored at -80°C. The BC was also isolated from the corresponding slices and stored at -80°C. Frozen tissue blocks were lysed in a solution containing (in mM) HEPES 50, NaCl 150, MgCl2 1.5, EGTA 5, Glycerol 1%, Triton-X100 1%, 1:1000 protease inhibitors cocktail. Proteins ran on 8.5% bis-acrylamide gel and were transferred onto a nitrocellulose membrane (Whatman GmbH, Dassel, Germany). The membrane was blocked, incubated in rabbit polyclonal anti-K+/Cl-Cotransporter (KCC2) (Merck KGaA, Darmstadt, Germany, 1:1000) or rabbit monoclonal anti-GAPDH (Cell Signaling Technology Europe BV, Leiden, Netherlands, 1:1000), washed, incubated in secondary goat anti-rabbit IgG Horseradish Peroxidase Conjugate antibody (Invitrogen, 1:5000), and digitally exposed using the Molecular Imaging system ChemiDoc (BioRad Laboratories, Inc, California, U.S.A.). Analysis of KCC2 and GAPDH expression was performed with ImageJ software, and the raw values of KCC2 from each sample were normalized to their respective GAPDH values.

Statistical analysis

Statistical analyses were performed in Microsoft Office Excel 2007 and GraphPad Prism Software 7.0. Data are presented as mean ± standard error of mean (SEM). Normality distribution and equality of variances of dataset were tested with the Kolmogorov-Smirnov test normality test. The null hypothesis was rejected for a >5%. When four experimental groups (P10 mPFC, P20 mPFC, P10 BC and P20 BC) were assessed and two variables were taken into consideration (age and brain area), data were analyzed with a two-way ANOVA with Fisher LSD, Sidak's or Tukey's multiple comparisons (electrophysiological recordings and cell counting). When three groups (P10 mPFC, P20 mPFC and P10 BC) data were analyzed with one-way ANOVA (electrophysiological recordings). For the comparison of *in vivo* spiking activity between P10 and P20, statistical analyses were performed with MATLAB. Significant differences were detected by one-way ANOVA. Significance levels of *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$ or ****$p < 0.0001$ were tested. For comparison of Western blot analysis, the significant effect of each developmental age group from mPFC and BC was assessed using Student's t-test depending on the experiment.

Modeling

We adapted the PFC microcircuit model we had developed previously (Konstantoudaki et al., 2014) to fit our experimental data with regards to intrinsic properties of pyramidal neurons and fast-spiking interneurons. We generated two model networks: a) a neonatal mPFC model network and b) a juvenile mPFC model network. In the neonatal mPFC model network the following adjustments were made: a) the pyramidal model neuron was adjusted to fit the passive and active properties of the neurons recorded in this paper, primarily the neonatal pyramidal model neuron exhibited increased input resistance and reduced spike amplitude, b) the fast-spiking interneuron model neuron was adjusted to have reduced fast afterhyperpolarization, as found in our experiments and c) the $GABA_AR$ receptor model had a reversal potential of -40mV. In the juvenile mPFC model network, the following adjustments were made: a) the $GABA_AR$ receptor model had a reversal potential of -60mV. All other properties were maintained as in the original model. The network was stimulated with spontaneous synaptic activation on pyramidal model neurons with characteristics similar to the sEPSC and sIPSC properties we recorded. Each condition was simulated in 10 different trials. In each trial, the location of activated synapses varied.

Data availability

Data presented in the figures in this paper are available upon request.

Results

Mice belonging to two age groups were investigated: (i) neonatal mice included pups of postnatal days (P) 9-11 and are defined as P10 while (ii) juvenile mice defined as P20 animals included pups of P19-P21. Due to the high density of intra-cortical synapses in the superficial cortical layers (DeFelipe and Fariñas 1992; Clancy et al. 2001) and their specific involvement in neurodevelopmental disorders (Chini and Hanganu-Opatz, 2020; Bitzenhofer eta l., 2017) we focused on the superficial layers of the mPFC and BC. From each mouse brain, both the mPFC and the BC were studied. All analyses that had four groups (mPFC P10 and P20, BC P10 and P20) were conducted using two-way ANOVA, with the two factors being the brain area (mPFC and BC) and age (P10 and P20).

Synaptic transmission adaptations in mPFC and BC across development

To examine spontaneous synaptic transmission, we performed patch-clamp recordings from

layer II/III pyramidal neurons in mPFC and BC from P10 and P20 mice. We recorded spontaneous

inhibitory postsynaptic currents (sIPSCs, at +10mV) and spontaneous excitatory postsynaptic

currents (sEPSCs, at -60mV) and we measured the frequency, amplitude and decay time

constant.

In mPFC, the frequency of sIPSCs was significantly augmented at P20 compared to P10 (**Figure**

1a,b), while the sIPSC amplitude and decay-time constant did not significantly change over the

investigated time window (**Figure 1a,c,d**). Similarly, the sIPSC frequency and amplitude were

significantly increased, at P20 compared to P10, in BC (**Figure 1a,b,c**), while the decay-time

constant was not altered (**Figure 1a,d**). On a different note, the sIPSC in BC at P20 was

significantly increased compared to P10 in BC but also compared to P20 in mPFC.

The sEPSC frequency was significantly decreased at P20 compared to P10, in both areas (**Figure**

1e,f), while the amplitude and decay time constant were unaltered (**Figure 1g,h**). Upon

comparing the two brain areas, the sEPSC frequency and amplitude were found significantly

decreased in mPFC, compared to BC, at P10 (**Figure 1f**). At P20, the sEPSC frequency was similar

between the two cortical areas, while the amplitude remained significantly smaller in mPFC

compared to BC in both ages (**Figure 1g**). The decay time constant was not different between

areas at both ages (**Figure 1h**).

We further investigated the changes in sIPSC and sEPSC frequency changes in the mPFC, in

particular, in the presence of AMPA and NMDA receptor antagonists, CNQX and AP5, respectively

and a GABA$_A$ receptor antagonist, bicuculine. The sIPSC frequency remained increased at P20,

compared to P10, CNQX and AP5 (**Figure 2a,c**). On the other hand, bicuculine blocked the sIPSCs

both in P10 and P20 mPFC (**Figure 2 b, c**). Application of CNQX eliminated the sEPSCs in mPFC

slices at P20 and significantly reduced, but did not eliminate sEPSCs, in mPFC slices at P10 (**Figure**

2e,g). Application of bicuculine (10uM) did not affect the sEPSC frequency in mPFC slices at P20

but significantly reduced sEPSC frequency at mPFC P10 (**Figure 2f,g**). These results suggest that

depolarizing GABA$_A$R currents could contribute to the increased sEPSC frequency observed in

layer II/III mPFC pyramidal neurons at P10.

GABA is depolarizing in the neonatal mPFC but not BC

Our data so far suggests that GABA$_A$R function is still depolarizing at P10 in the mPFC. To further

investigate this, we performed evoked IPSCs (eIPSCs) at P10 and P20 mPFC in the presence of

CNQX and AP5 at different voltage steps. A line curve was fit across the eIPSC currents recorded from -60mV to +10mV, because there was a linear V-I relationship at this range. From the graphs, it is evident that reversal potential at P10 is more positive than -30mV, while the reversal potential at P20 is around -45mV (**Figure 3a,b**).

We further investigated the GABA$_A$R activity by isolating a GABA$_A$R response in fEPSP recordings by using CNQX and AP5. The addition of diazepam (2µM) (a GABA$_A$R agonist) increased the fEPSP amplitude in mPFC at P10, but not at P20 (**Figure 4a-b**). In BC, there was no effect of diazepam, either at P10 or P20 (**Supplemental Figure 1**). The switch in the GABA$_A$R function from depolarizing to hyperpolarizing occurs due to the increased expression of the K$^+$-Cl$^-$ co-transporter 2 (KCC2) (Rivera et al. 1999). To determine whether modulating chloride transporters could alter the diazepam-induced enhancement of the fEPSP at P10 mPFC, we recorded the fEPSP in the presence of bumetanide (10uM), which blocks the NKCC1 transporter, and tested the effect of diazepam. We find that in the presence of bumetanide, diazepam did not result in an increase of the fEPSP (**Figure 4c-d**). In addition, we measured KCC2 protein levels and demonstrated that they were significantly increased at P20 compared to P10 in the mPFC but not in the BC (**Figure 4e-g**). The above results collectively suggest that the GABA$_A$R function is depolarizing in the mPFC at P10.

Passive and active membrane properties of MGE-derived interneurons are altered in the mPFC across development

To investigate whether the change in GABA$_A$R function in mPFC is also associated with changes in interneuron properties, we performed current-clamp recordings from layer II/III mPFC and BC of Lhx6$^+$ interneurons. For this reason, Lhx6-cre;ROSA26fl-STOP-fl-YFP mice were used in which Lhx6$^+$ interneurons express YFP. Lhx6 is expressed by all post-mitotic and mature MGE-derived interneurons (Liodis et al. 2007), therefore, YFP is expressed in MGE-derived interneurons, which include interneurons that express parvalbumin (PV$^+$) and somatostatin (SST$^+$).

Upon analysis of the passive properties, we found a significant increase in the input resistance and membrane time constant, as well as a significant decrease in the membrane capacitance in the mPFC at P10 compared to P20 (**Figure 5**). In addition, the input resistance and the membrane time constant were higher at P10 mPFC, compared to BC (P10 and P20). There was no difference in the resting membrane potential (RMP) between ages and brain areas (**Figure 5, Supplementary Table 1**).

Regarding the active properties, there was no significant difference between ages and brain areas in the AP amplitude, AP threshold and rheobase, fAHP time (duration) (**Figure 5a, d, e, g; Supplementary Table 1**). The AP rate of rise (dv/dt) was significantly increased while the AP duration (half-width) was significantly reduced at P20 compared to P10 in both the mPFC and BC (**Figure 6b,c, Supplementary Table 1**). In addition, the fAHP amplitude was significantly lower in the mPFC (**Figure 6f, Supplementary Table 1**), compared to BC. The increased rate of rise and the decreased AP duration are possibly linked with the up-regulation of voltage-dependent sodium channels during development (Huguenard et al. 1988), and in combination with the reduced fAHP amplitude suggest that the mPFC MGE-interneurons at P10 are still quite immature, when compared to adult PV[+]/SST[+] interneurons in mPFC (Yang et al. 2013; Pan et al. 2017).

Overall, these data indicate that some intrinsic properties of interneurons in mPFC change with age (from P10 to P20), reaching values that closer resemble adult MGE-derived interneurons (Yang et al. 2013; Pan et al. 2017). The increased sIPSC frequency of mPFC pyramidal neurons observed at P20, compared to P10 could partly be explained by these altered properties of presynaptic interneurons.

Decreased PV interneurons in mPFC compared to BC

An additional explanation for the adaptations in inhibitory transmission could come from alterations in interneuron cell densities. To test this, we quantified the number of interneurons per area in cryosections at P10 and P20 mPFC and BC coronal brain slices of Lhx6[+]-expressing mice. The YFP[+] positive cells per area (i.e. Lhx6[+] cell density) in mPFC and BC was similar between ages, but was significantly lower in the mPFC, compared to BC (**Figure 7a**). But we found no contribution of cell death in the mPFC interneuron population, as the percentage of cell death in Lhx6[+] neurons is very low in both P10 mPFC and BC (**Supplementary Figure 2**).

The transcription factor Lhx6 is required for the specification and maintenance of main MGE-derived interneurons, PV and SST-positive interneuron subtypes, at postnatal ages (Liodis et al. 2007). The neuropeptide SST (both mRNA and protein) is progressively expressed from embryonic to postnatal levels (Bendotti et al. 1990; Forloni et al. 1990). We found that the SST mRNA levels were similar between areas and ages (**Figure 7b**). On the other hand, the emergence of PV immunoreactivity in the mouse cortex shows a delayed development, starting from early postnatal period to adult, with marked area-specific differences(Del Rio et al. 1992).

We found that PV was only immunoreactive in BC, and not in mPFC, at P10 (**Figure 7c and Supplementary Figure 3**). At P20, PV was immunoreactive in both mPFC and BC, but PV$^+$ cell density was significantly lower in the mPFC, compared to BC (**Figure 7c**).

We also counted the total cell density of mPFC and BC from neonatal and juvenile mice (**Supplementary Figure 4a**) using Nissl staining. In the mPFC, the cell density significantly decreased at P20 compared to P10 (**Supplementary Figure 4b**). On the contrary, in BC, the total cell density significantly increased at P20 compared to P10 (**Supplementary Figure 4b**). When the two brain areas were compared, no difference was found at P10, while the mPFC cell density was significantly lower compared to BC at P20 (**Supplementary Figure 4b**).

We further examined whether the alterations in total cell density are derived from alterations in cell density of interneurons by measuring the Lhx6$^+$ neurons over the Nissl-positive cells. No differences were detected between areas and ages (**Supplementary Figure 4c).** These results suggest that the changes in total cell density in mPFC and BC respectively are probably due to changes in other neuronal or glial populations.

No significant changes in pyramidal neuron excitability

To determine whether the reduced sEPSC frequency can be explained by changes in pyramidal neuron excitability, we investigated their intrinsic properties. The passive and active properties of these neurons were measured using current-clamp recordings from layer II/III mPFC and BC pyramidal neurons. With regards to passive properties, no significant differences were observed in the RMP, the input resistance and the membrane time constant between brain regions and ages (**Supplementary Figure 5, Supplementary Table 1**). Only the membrane capacitance was significantly increased at P20 compared to P10 (**Supplementary Figure 5d, Supplementary Table 1**), in both brain areas. In addition, the number of spikes generated with increasing current stimulation was not significantly different between ages and regions (**Supplementary Figure 6**). In terms of active properties, the AP amplitude and rate of rise were increased at P20 compared to P10 mPFC, while the AP half-width, rheobase and threshold were not significantly different (**Figure 8, Supplementary Table 1**). The AP amplitude was also significantly increased at P20, compared to P10 in BC, while the other properties did not change (**Figure 8, Supplementary Table 1**). Comparing the two regions at the two ages, we found no significant differences of AP properties of pyramidal neurons (**Figure 8, Supplementary Table 1**). The developmental increase of AP amplitude and rate of rise in the mPFC could be due to the on-going maturation of sodium

channels in pyramidal neurons. However, these changes could not account for the reduced sEPSCs in the neonatal, compared to juvenile, mPFC and BC.

Effects of mPFC changes in intrinsic and synaptic properties on PFC network activity

In an effort to understand the circuit effects of the aforementioned differences in intrinsic and synaptic properties between p10 and P20 mPFC, we adapted an already validated PFC network model (Konstantoudaki et al., 2014) to our current data to generate a neonatal (P10) mPFC model network and juvenile (P20) mPFC model network. Both model networks were stimulated with spontaneous excitatory and inhibitory inputs based on our results for sEPSC and sIPSC frequency at P10 and P20 (**Figure 1 and Supplementary Table 1**). Our simulation results predict that the intrinsic and synaptic changes from P10 to P20 in mPFC result in enhanced mPFC network activity at P20, compared to P10 (**Figure 9a,e**), as indicated by single-cell activity (**Figure 9a,b**) and the filtered signal (**Figure 9c,d**). Furthermore, we investigated which adaptations have a significant contribution to the network activity properties observed in each model network. Therefore, we generated neonatal model networks in which the pyramidal model neuron used had P20 properties (N2 network), the fast-spiking interneuron model used had P20 properties (N3 network) and the GABA$_A$R reversal potential was that of P20 (N4 network). We identify that in all the above networks, the resulting neuronal activity generated was significantly reduced compared to the control P10 network (N1) (**Figure 9f**). Similarly, we generated juvenile model networks in which the pyramidal model neuron used had P10 properties (J2 network), the fast-spiking interneuron model used had P10 properties (J3 network) and the GABA$_A$R reversal potential was that of P10 (J4 network). We find that the resulting neuronal activity generated was significantly increased compared to the control P20 network (J1) (**Figure 9g**).

Increased firing activity *in vivo* in the mPFC between the second and third postnatal weeks

To determine whether our modeling prediction on network activity changes can be validated *in vivo*, multisite recordings of the LFP and multi-unit activity (MUA) were performed in layers II/III of mPFC at P8-10 and P20-23. A significant increase of MUA was identified at P22 compared to P9 mice, indicating a developmental increased spiking activity in layers II/III of mPFC on the third compared to the second postnatal week (**Figure 10**). Similarly, the spiking activity in the BC augmented with age (**Supplementary Figure 7**). To characterize the firing dynamics along development, we used MUA for clustering single-unit activity. We detected bursts of spikes using

the previously developed method (Gorin et al. 2016). The burst occurrence in the mPFC

increased with age (33.62 ± 1.91 %, 161 single units at neonatal age vs. 50.12 ± 4.91 %, 150 single

units at juvenile age; $p<0.0001$, One-way ANOVA, $F_{(1, 309)}=272.26$). Correspondingly, the number

of units organized in bursts increased with age, as well, from 21 out of 161 single units at

neonatal age to 104 out of 150 single units at juvenile age (**Figure 10d**). Moreover, we analyzed

the mPFC and BC units in their amplitude, dv/dt and half-width and detected, similarly to the in

vitro conditions, a significant increase in the AP amplitude and rate of rise but no difference in

the half-width (**Supplementary Figure 8**). These results provide further evidence that the

decreased excitatory synaptic activity of mPFC cannot be attributed to increased spiking activity

of mPFC neurons at P10.

Discussion

Our study has identified significant developmental events in the mPFC and the BC between the

second and third postnatal weeks. Specifically, we have shown that spontaneous inhibitory

transmission, measured as sIPSCs, is increased in mPFC from P10 to P20. Moreover, our data

support a depolarizing action of GABA$_A$R in the second postnatal week, in the mPFC, as indicated

by the presence of non-AMPAR-mediated sEPSCs, depolarizing reversal potential of evoked

GABA$_A$R responses, increased basal fEPSPs following GABA$_A$R activation, which is blocked by

concurrent bumetanide application, and decreased protein levels of KCC2. In parallel,

differences the intrinsic properties of GABAergic interneurons of mice in the juvenile period

resemble are more mature, compared to the neonatal period in mPFC. The above

developmental adaptations, along with increased AP amplitude and AP rate-of-rise in pyramidal

cells age result in augmented network activity in the juvenile mPFC, compared to the neonatal

mPFC.

Depolarizing action of GABA in the immature cortex

GABA plays a crucial role in inhibiting adult neurons, acting primarily via the chloride-permeable

GABA$_A$R and resulting in hyperpolarization of the membrane potential (Kaila and Voipio 1987).

However, GABA action leads to depolarization of immature neurons (i.e. during the first

postnatal week in mice), due to an initially higher intracellular chloride concentration $[Cl^-]_{in}$ (Ben-

Ari 2001; Ben-Ari et al. 2007; Ben-Ari 2012). The developmental switch of GABA action from

depolarizing to hyperpolarizing results from changes in cation-chloride co-transporter

expression: NKCC1, a cation-Cl$^-$ importer, is highly expressed in neuronal precursor cells during

early brain development (Plotkin et al. 1997; Yamada et al. 2004), while the expression of the

K$^+$-Cl$^-$ cotransporter 2 (KCC2), a cation-Cl$^-$ exporter, increases after the first postnatal week (Ben-

Ari 2001; Ben-Ari et al. 2007; Ben-Ari 2012). This increased KCC2 transporter expression might

provide a central mechanism for the depolarization to hyperpolarization switch of GABAergic

transmission via progressive reduction of $[Cl^-]_{in}$ (Lu et al. 1999; Rivera et al. 1999; Ganguly et al.

2001; Ben-Ari 2002; Dzhala et al. 2005; Fiumelli et al. 2005).

The GABA$_A$R switch from depolarizing to hyperpolarizing occurs at P7 in the hippocampus,

cortex, amygdala (Ben-Ari et al. 1989; Luhmann and Prince 1991; LoTurco et al. 1995; Owens et

al. 1996; Martina et al. 2001; Gulledge and Stuart 2003; Ben-Ari et al. 2007). The most

developmentally delayed switch in the GABAergic polarity has identified in the mPFC. In Rinetti-

Viargas et al., 2017, they showed that the GABA$_A$ signaling matures around birth at dendrites

and after preadolescent period (~P20) at axon initial segment of prefrontal pyramidal neuros.

They showed the changes in local chloride transporters function using gramicidin-based

perforated patch recordings and applied GABA via an iontophoretic pipette. We reinforce the

GABA$_A$ signaling change using evoked IPSC recordings in the soma if prefrontal pyramidal

neurons.

Our study suggests that this switch is delayed in the mPFC compared to primary somatosensory

cortex and it takes place between P10 and P20. Specifically, we show that increased GABA$_A$R

activity leads to enhanced fEPSPs in neonatal mPFC (P10), suggesting that the GABA$_A$R function

is depolarizing in the mPFC at P10. This enhancement is prevented in the presence of the NKCC1

blocker, bumetanide. Furthermore, diazepam increases the number of spontaneous activity

events in the brain slice, which is also prevented in the presence of bumetanide. This hypothesis

is further supported by decreased levels of KCC2 transporter in the neonatal mPFC and it is

conceded the age- dependent differences in membrane NKCC1 and KCC2 protein level and

phosphorylation (Rinetti-Viargas et al., 2017) Our results could have implications for

understanding the protracted maturation of mPFC compared to other cortical areas, which may

depend on a combination of a delayed switch from depolarizing-to-hyperpolarizing function of

GABA$_A$R and maturation of interneurons.

Interneurons and mPFC development

Recordings of Lhx6[+]- interneurons indicate that both passive and active properties are regulated
by age and reach values that better resemble adult MGE-derived interneurons. Specifically, we
have found that the input resistance and AP width decrease while the AP rate of rise increases
in the mPFC at P20 compared to P10. In part, similar findings have been identified for PV[+] cells
in the hippocampus (Doischer et al. 2008; Miyamae et al. 2017) and SST[+] cells in the anterior
cingulate cortex (Pan et al. 2017). On the other hand, the AHP amplitude is still quite immature
in the mPFC at P20, compared to PV[+], SST[+] interneurons in primary sensory areas or the
hippocampus and compared to adult mPFC (Doischer et al. 2008; Yang et al. 2013; Pan et al.
2017). Therefore, it is likely that the physiological properties of PV+ and SST+ interneurons in
the mPFC continue to change past the third postnatal week.

Our knowledge on the neonatal physiology of mPFC GABAergic interneurons is very limited. It
has been shown that PV expression is lowest in juveniles and increases during adolescence to
levels similar to those observed in adulthood (Caballero et al. 2014). Furthermore, PV expression
is not evident in the neonatal period and emerges during the juvenile period in the mPFC (del
Rio et al. 1994; de Lecea et al. 1995; Zheng et al. 2011; Spampanato and Sullivan 2016). Our
results agree with these findings, as PV expression was detected in very low amounts during the
juvenile period in the mPFC.

In addition, our study has identified decreased excitatory and increased inhibitory synaptic
function between the second and third postnatal weeks. We show that the frequency of sIPSCs
in layer II/III pyramidal cells of mPFC increases from neonatal to juvenile period, consistent with
the developmental changes of IPSCs in layer III pyramidal neurons of monkey PFC (González-
Burgos et al. 2015) and mouse mPFC (Kroon et al. 2019).

Pyramidal neurons and network activity
It has been suggested that spontaneous network activity changes from local, highly synchronized
to more diffuse from the second to the third postnatal weeks, in the primary sensory cortices
(Golshani et al. 2009; Frye and MacLean 2016). Oscillatory activity in the mPFC first emerges at
P15 (Bitzenhofer et al. 2019). In this study, we have found increased spiking activity in the mPFC
during the third, compared to the second postnatal week. This occurred despite the decreased
excitatory and increased inhibitory synaptic function, but could be explained partly by the
developmental increase of AP amplitude and rate of rise in the mPFC layer II/III pyramidal

neurons, which could be due to the on-going maturation of sodium channels in pyramidal

neurons.

Studies in developing mPFC pyramidal neurons have proposed that there is a unique sensitive

time window for synaptic maturation of these neurons from individual cortical layers. During rat

mPFC layer V development, the intrinsic properties, synaptic inputs and morphology of

pyramidal neurons develop together during early postnatal life. While the greatest changes were

reported during the first ten days after birth, the adult-like properties emerged after the end of

the third week (P21) (Zhang et al. 2011). This study confirms that the second postnatal week is

a period of rapid growth, similar to that in other neocortical regions by combining functional and

structural measurements of developing pyramidal neurons in mouse mPFC (Zhu 2000; Romand

et al. 2011).

Developmental PFC malformation leads to cognitive disorders in adulthood

The neonatal functional maturation of GABAergic circuits and E/I (excitation to inhibition)

balance are critical for PFC-dependent behaviours and plasticity in the adult while their

malfunction leads to many psychiatric disorders (Benes 1991; Kilb 2012; Ferguson and Gao

2018). From the prenatal period to late adolescence, the PFC network is highly vulnerable to

genetic and environmental factors (Andersen 2003), since the mPFC is one of the latest cortical

regions to develop (Huttenlocher 1990). While many studies have focused on understanding

several developmental processes during adolescence (Caballero et al. 2016), our knowledge

regarding the ongoing cellular and network developmental processes during the perinatal period

is notably limited, despite significant evidence showing that environmental manipulations

during this period manifest as complex psychiatric and neurologic disorders in adulthood

(Weinberger 1986).

The delayed developmental shift of GABA action in various mouse models mimicking human

brain disorders have been investigated, including the maternal immune activation model

(Corradini et al. 2018; Fernandez et al. 2018), the Scn1a and Scn1b mouse models of Dravet

syndrome (Yuan et al., 2019), the 22q11.2 deletion syndrome (Amin et al. 2017) and the Fmr1

deficient model of fragile X syndrome (He et al. 2018). In the latter study, early postnatal

correction of GABA depolarization (bumetanide-treated) led to sufficient normalization of the

mature BC network (He et al. 2018). The impaired KCC2 has been proposed as a potential

therapeutic target of epilepsies by many studies in animal models and human patients (Moore

et al. 2017).

Our study focuses in understanding the early developmental cellular and physiological mechanisms of mPFC circuits, before adolescence, and proposes that the neonatal mPFC compared to BC exhibits a delayed switch from depolarization to hyperpolarization function of $GABA_AR$. Our results raise the possibility that the delayed maturation of mPFC compared to other cortical areas depends on a combination of a delayed switch from depolarization to hyperpolarization function of the $GABA_AR$ and delayed maturation of interneurons.

Acknowledgements:

Authors are grateful to Emmanuella Foinikianaki for her help with Matlab analysis and histology and to Giasemi Eptaminitaki for her help in *in situ* hybridization experiments. They also would like to thank all the members of Karagogeos and Sidiropoulou Labs and the animal facility of the IMBB for help with experiments. This study was co-financed through the Operational Program "Education and Lifelong Learning" of the National Strategic Reference Framework – Research Funding Program (EDBM34) by a grant to DK (10040) and through the BIOIMAGING-GR, National Roadmap for Research Infrastructures from the European Union (European Social Fund-ESF) and Greek National Funds. KK has been a recipient of the Manasaki fellowship and a Medical School fellowship of the University of Crete and a poster award at the 27th Hellenic Society for Neuroscience Meeting.

Author Contributions:

All experiments were conceived and designed by K.K., K.S., and D.K. All experiments performed by K.K., A.V., O.C. and X.X. Computational simulations were performed by K.S. Data were analyzed by K.K., A.V., O.C., M.D. K.S. and discussed with D.K. X.X. and I.L.H.-O. X.X. carried out and analyzed the *in vivo* experiments. Manuscript was written by K.K., D.K. and K.S. All authors discussed and commented on the manuscript.

References:

Allene C, Cattani A, Ackman JB, Bonifazi P, Aniksztejn L, Ben-Ari Y, Cossart R. 2008. Sequential
Generation of Two Distinct Synapse-Driven Network Patterns in Developing Neocortex. J
Neurosci. 28:12851–12863.

Amin H, Marinaro F, De Pietri Tonelli D, Berdondini L. 2017. Developmental excitatory-to-
inhibitory GABA-polarity switch is disrupted in 22q11.2 deletion syndrome: a potential
target for clinical therapeutics. Scientific Reports. 1–18.

Andersen SL. 2003. Trajectories of brain development: point of vulnerability or window of
opportunity? Neuroscience & Biobehavioral Reviews. 27:3–18.

Bahrey HLP, Moody WJ. 2003. Early Development of Voltage-Gated Ion Currents and Firing
Properties in Neurons of the Mouse Cerebral Cortex. Journal of Neurophysiology. 89:1761–
1773.

Bartolini G, Ciceri G, Marin O. 2013. Integration of GABAergic Interneurons into Cortical Cell
Assemblies: Lessons from Embryos and Adults. 79:849–864.

Ben-Ari Y. 2001. Developing networks play a similar melody. Trends in Neurosciences. 24:353–
360.

Ben-Ari Y. 2002. Excitatory actions of gaba during development: the nature of the nurture. Nat
Rev Neurosci. 3:728–739.

Ben-Ari Y. 2012. Refuting the challenges of the developmental shift of polarity of GABA actions:
GABA more exciting than ever! 1–18.

Ben-Ari Y, Cherubini E, Corradetti R, Gaiarsa JL. 1989. Giant synaptic potentials in immature rat
CA3 hippocampal neurones. The Journal of Physiology. 416:303–325.

Ben-Ari Y, Gaiarsa JL, Tyzio R, Khazipov R. 2007. GABA: A Pioneer Transmitter That Excites
Immature Neurons and Generates Primitive Oscillations. Physiological Reviews. 87:1215–
1284.

Ben-Ari Y, Khalilov I, Kahle KT, Cherubini E. 2012. The GABA Excitatory/Inhibitory Shift in Brain
Maturation and Neurological Disorders. The Neuroscientist. 18:467–486.

Bendotti C, Hohmann C, Forloni G, Brain RRD, 1990. 1990. Developmental expression of
somatostatin in mouse brain. II. In situ hybridization. Elsevier. 53:26–39.

Benes FM. 1991. Deficits in Small Interneurons in Prefrontal and Cingulate Cortices of
Schizophrenic and Schizoaffective Patients. Archives of General Psychiatry. 48:996–1001.

Best JR, Miller PH. 2010. A Developmental Perspective on Executive Function. Child
Development. 81:1641–1660.

Bitzenhofer SH, Ahlbeck J, Wolff A, Wiegert JS, Gee CE, Oertner TG, Hanganu-Opatz IL. 2017.

Layer-specific optogenetic activation of pyramidal neurons causes beta-gamma entrainment

of neonatal networks. Nature Communications. 8:14563.

Bitzenhofer SH, Poepplau JA, Chini M, Marquardt A, Hanganu-Opatz I. 2019. Activity-dependent

maturation of prefrontal gamma oscillations sculpts cognitive performance. bioRxiv. 1–39.

Bitzenhofer SH, Sieben K, Siebert KD, Spehr M, Hanganu-Opatz IL. 2015. Oscillatory Activity in

Developing Prefrontal Networks Results from Theta-Gamma-Modulated Synaptic Inputs.

CellReports. 11:486–497.

Brockmann MD, Pöschel B, Cichon N, Hanganu-Opatz IL. 2011. Coupled Oscillations Mediate

Directed Interactions between Prefrontal Cortex and Hippocampus of the Neonatal Rat.

Neuron. 71:332–347.

Burgard EC, Hablitz JJ. 1993. Developmental changes in NMDA and non-NMDA receptor-

mediated synaptic potentials in rat neocortex. Journal of Neurophysiology. 69:230–240.

Caballero A, Flores-Barrera E, Cass DK, Tseng KY. 2014. Differential regulation of parvalbumin

and calretinin interneurons in the prefrontal cortex during adolescence. Brain Struct Funct.

219:395–406.

Caballero A, Granberg R, Tseng KY. 2016. Mechanisms contributing to prefrontal cortex

maturation during adolescence. Neuroscience & Biobehavioral Reviews. 1–29.

Casey BJ, Giedd JN, Thomas KM. 2000. Structural and functional brain development and its

relation to cognitive development. Biological Psychology. 54:241–257.

Chalkiadaki K, Velli A, Kyriazidis E, Stavroulaki V, Vouvoutsis V, Chatzaki E, Aivaliotis M,

Sidiropoulou K. 2019. Development of the MAM model of schizophrenia in mice: Sex

similarities and differences of hippocampal and prefrontal cortical function.

Neuropharmacology 144, 193–207.

Chini, M. & Hanganu-Opatz, I. L. 2020. Prefrontal Cortex Development in Health and Disease:

Lessons from Rodents and Humans. *Trends Neurosci*

Clancy B, Darlington RB, Neuroscience BF, 2001. 2001. Translating developmental time across

mammalian species. Elsevier. 105:7–17.

Corradini I, Focchi E, Rasile M, Morini R, Desiato G, Tomasoni R, Lizier M, Ghirardini E, Fesce R,

Morone D, Barajon I, Antonucci F, Pozzi D, Matteoli M. 2018. Maternal Immune Activation

Delays Excitatory-to-Inhibitory Gamma-Aminobutyric Acid Switch in Offspring. Biological

Psychiatry. 83:680–691.

de Lecea L, del Ri o JA, Soriano E. 1995. Developmental expression of parvalbumin mRNA in the
cerebral cortex and hippocampus of the rat. Molecular Brain Research. 32:1–13.

DeFelipe J, Fariñas I. 1992. The pyramidal neuron of the cerebral cortex: Morphological and
chemical characteristics of the synaptic inputs. Progress in Neurobiology. 39:563–607.

del Rio J, De Lecea L, Ferrer I, Soriano E. 1994. The development of parvalbumin-
immunoreactivity in the neocortex of the mouse. 81:247–259.

Del Rio JA, Soriano E, Ferrer I. 1992. Development of GABA-immunoreactivity in the neocortex
of the mouse. J Comp Neurol. 326:501–526.

Diamond A. 2005. Normal Development of Prefrontal Cortex from Birth to Young Adulthood:
Cognitive Functions, Anatomy, and Biochemistry. In: Principles of Frontal Lobe Function.
Oxford University Press. p. 466–503.

Doischer D, Aurel Hosp J, Yanagawa Y, Obata K, Jonas P, Vida I, Bartos M. 2008. Postnatal
Differentiation of Basket Cells from Slow to Fast Signaling Devices. J Neurosci. 28:12956–
12968.

Dzhala VI, Talos DM, Sdrulla DA, Brumback AC, Mathews GC, Benke TA, Delpire E, Jensen FE,
Staley KJ. 2005. NKCC1 transporter facilitates seizures in the developing brain. Nature
Medicine 2005 11:11. 11:1205–1213.

Ferguson BR, Gao W-J. 2018. PV Interneurons: Critical Regulators of E/I Balance for Prefrontal
Cortex-Dependent Behavior and Psychiatric Disorders. Front Neural Circuits. 12:479–13.

Fernandez A, Dumon C, Guimond D, Tyzio R, Bonifazi P, Lozovaya N, Burnashev N, Ferrari DC,
Ben-Ari Y. 2018. The GABA Developmental Shift Is Abolished by Maternal Immune Activation
Already at Birth. Cerebral Cortex. 130:e1447–11.

Fiumelli H, Cancedda L, Poo M-M. 2005. Modulation of GABAergic Transmission by Activity via
Postsynaptic Ca2+-Dependent Regulation of KCC2 Function. Neuron. 48:773–786.

Forloni G, Hohmann C, Research JCDB, 1990. 1990. Developmental expression of somatostatin
in mouse brain. I. Immunocytochemical studies. Elsevier. 53:6–25.

Frye CG, MacLean JN. 2016. Spontaneous activations follow a common developmental course
across primary sensory areas in mouse neocortex. Journal of Neurophysiology. 116:431–
437.

Fuster J. 2015. The Prefrontal Cortex. Academic Press.

Ganguly K, Schinder AF, Wong ST, Poo M-M. 2001. GABA Itself Promotes the Developmental
Switch of Neuronal GABAergic Responses from Excitation to Inhibition. Cell. 105:521–532.

Golshani P, Goncalves JT, Khoshkhoo S, Mostany R, Smirnakis S, Portera-Cailliau C. 2009.
Internally Mediated Developmental Desynchronization of Neocortical Network Activity. J
Neurosci. 29:10890–10899.

Gorin, M., C. Tsitoura, A. Kahan, K. Watznauer, D. R. Drose, M. Arts, R. Mathar, S. O'Connor, I. L.
Hanganu-Opatz, Y. Ben-Shaul and M. Spehr (2016). "Interdependent Conductances Drive
Infraslow Intrinsic Rhythmogenesis in a Subset of Accessory Olfactory Bulb Projection
Neurons." The Journal of Neuroscience 36(11): 3127-3144.

González-Burgos G, Miyamae T, Pafundo DE, Yoshino H, Rotaru DC, Hoftman G, Datta D, Zhang
Y, Hammond M, Sampson AR, Fish KN, Bard Ermentrout G, Lewis DA. 2015. Functional
Maturation of GABA Synapses During Postnatal Development of the Monkey Dorsolateral
Prefrontal Cortex. Cerebral Cortex. 25:4076–4093.

Gulledge AT, Stuart GJ. 2003. Excitatory Actions of GABA in the Cortex. Neuron. 37:299–309.

Guo ZV, Li N, Huber D, Ophir E, Gutnisky D, Ting JT, Feng G, Svoboda K. 2014. Flow of Cortical
Activity Underlying a Tactile Decision in Mice. Neuron. 81:179–194.

He Q, Arroyo ED, Smukowski SN, Xu J, Piochon C, Savas JN, Portera-Cailliau C, Contractor A. 2018.
Critical period inhibition of NKCC1 rectifies synapse plasticity in the somatosensory cortex
and restores adult tactile response maps in fragile X mice. Mol Psychiatry. 8:109–116.

Heidbreder CA, Groenewegen HJ. 2003. The medial prefrontal cortex in the rat: evidence for a
dorso-ventral distinction based upon functional and anatomical characteristics.
Neuroscience & Biobehavioral Reviews. 27:555–579.

Hensch TK. 2016. The Power of the Infant Brain. Nature Publishing Group. 314:64–69.

Huguenard JR, Hamill OP, Prince DA. 1988. Developmental changes in Na+ conductances in rat
neocortical neurons: appearance of a slowly inactivating component. Journal of
Neurophysiology. 59:778–795.

Huttenlocher PR. 1990. Morphometric study of human cerebral cortex development.
Neuropsychologia. 28:517–527.

Kaila K, Voipio J. 1987. Postsynaptic fall in intracellular p H induced by GABA-activated
bicarbonate conductance. Nature. 330:163–165.

Khazipov R, Luhmann HJ. 2006. Early patterns of electrical activity in the developing cerebral
cortex of humans and rodents. Trends in Neurosciences. 29:414–418.

Khazipov R, Minlebaev M, Valeeva G. 2013. Early gamma oscillations. Neuroscience. 250:240–
252.

Khazipov R, Sirota A, Leinekugel X, Holmes GL, Ben-Ari Y, Buzsáki G. 2004. Early motor activity
drives spindle bursts in the developing somatosensory cortex. Nature. 432:758–761.

Kilb W. 2012. Development of the GABAergic System from Birth to Adolescence. The
Neuroscientist. 18:613–630.

Kirmse K, Kummer M, Kovalchuk Y, Witte OW, Garaschuk O, Holthoff K. 2015. GABA depolarizes
immature neurons and inhibits network activity in the neonatal neocortex in vivo. Nature
Communications. 6:7750.

Kolb B, Mychasiuk R, Muhammad A, Li Y, Frost DO, Gibb R. 2012. Experience and the developing
prefrontal cortex. Proc Natl Acad Sci USA. 109:17186–17193.

Konstantoudaki X, Papoutsi A, Chalkiadaki K, Poirazi P, Sidiropoulou K. Modulatory effects of
inhibition on persistent activity in a cortical microcircuit model. Front Neural Circuits. 2014
Jan 31;8:7.

Konstantoudaki X, Chalkiadaki K, Tivodar S, Karagogeos D, Sidiropoulou K. 2016. Impaired
synaptic plasticity in the prefrontal cortex of mice with developmentally decreased number
of interneurons. Neuroscience. 1–13.

Kriegstein AR, Suppes T, Prince DA. 1987. Cellular and synaptic physiology and epileptogenesis
of developing rat neocortical neurons in vitro. Developmental Brain Research. 34:161–171.

Kroeze Y, Oti M, van Beusekom E, Cooijmans RHM, van Bokhoven H, Kolk SM, Homberg JR, Zhou
H. 2017. Transcriptome Analysis Identifies Multifaceted Regulatory Mechanisms Dictating a
Genetic Switch from Neuronal Network Establishment to Maintenance During Postnatal
Prefrontal Cortex Development. Cerebral Cortex. 1–19.

Kroon T, van Hugte E, van Linge L, Mansvelder HD, Meredith RM. 2019. Early postnatal
development of pyramidal neurons across layers of the mouse medial prefrontal cortex.
Scientific Reports. 9:5037–16.

Le Magueresse C, Monyer H. 2013. GABAergic Interneurons Shape the Functional Maturation of
the Cortex. Neuron. 77:388–405.

Liodis P, Denaxa M, Grigoriou M, Akufo-Addo C, Yanagawa Y, Pachnis V. 2007. Lhx6 Activity Is
Required for the Normal Migration and Specification of Cortical Interneuron Subtypes.
Journal of Neuroscience. 27:3078–3089.

LoTurco JJ, Owens DF, Heath MJS, Davis MBE, Kriegstein AR. 1995. GABA and glutamate
depolarize cortical progenitor cells and inhibit DNA synthesis. Neuron. 15:1287–1298.

Lu J, Karadsheh M, Delpire E. 1999. Developmental regulation of the neuronal-specific isoform
of K-CL cotransporter KCC2 in postnatal rat brains. J Neurobiol. 39:558–568.

Luhmann HJ, Prince DA. 1991. Postnatal maturation of the GABAergic system in rat neocortex.
Journal of Neurophysiology. 65:247–263.

Martina M, Royer S, Paré D. 2001. Cell-Type-Specific GABA Responses and Chloride Homeostasis
in the Cortex and Amygdala. Journal of Neurophysiology. 86:2887–2895.

McCormick DA, Prince DA. 1987. Post-natal development of electrophysiological properties of
rat cerebral cortical pyramidal neurones. The Journal of Physiology. 393:743–762.

Miyamae T, Chen K, Lewis DA, González-Burgos G. 2017. Distinct physiological maturation of
parvalbumin-positive neuron subtypes in mouse prefrontal cortex. J Neurosci. 3325–16–48.

Moore YE, Kelley MR, Brandon NJ, Deeb TZ, Moss SJ. 2017. Seizing Control of KCC2: A New
Therapeutic Target for Epilepsy. Trends in Neurosciences. 40:555–571.

Mòdol L, Bollmann Y, Tressard T, Baude A, Che A, Duan ZRS, Babij R, De Marco García NV, Cossart
R. 2019. Assemblies of Perisomatic GABAergic Neurons in the Developing Barrel Cortex.
Neuron. 1–18.

Owens DF, Boyce LH, Davis MBE, Kriegstein AR. 1996. Excitatory GABA Responses in Embryonic
and Neonatal Cortical Slices Demonstrated by Gramicidin Perforated-Patch Recordings and
Calcium Imaging. J Neurosci. 16:6414–6423.

Pan G, Yang J-M, Hu X-Y, Li X-M. 2017. Postnatal development of the electrophysiological
properties of somatostatin interneurons in the anterior cingulate cortex of mice. Nature
Publishing Group. 1–12.

Plotkin MD, Snyder EY, Hebert SC, Delpire E. 1997. Expression of the Na-K-2Cl cotransporter is
developmentally regulated in postnatal rat brains: A possible mechanism underlying GABA's
excitatory role in immature brain. J Neurobiol. 33:781–795.

Ramoa AS, McCormick DA. 1994. Developmental changes in electrophysiological properties of
LGNd neurons during reorganization of retinogeniculate connections. J Neurosci. 14:2089–
2097.

Rinetti-Vargas, Gina, Khanhky Phamluong, Dorit Ron, and Kevin J. Bender. 2017. "Periadolescent
Maturation of GABAergic Hyperpolarization at the Axon Initial Segment." *CellReports*
20(1):21–29.

Rivera C, Voipio J, Payne JA, Ruusuvuori E, Lahtinen H, Lamsa K, Pirvola U, Saarma M, Kaila K.
1999. The K+/Cl- co-transporter KCC2 renders GABA hyperpolarizing during neuronal
maturation. Nature. 397:251–255.

Romand S, Wang Y, Toledo-Rodriguez M, Markram H. 2011. Morphological Development of
Thick-Tufted Layer V Pyramidal Cells in the Rat Somatosensory Cortex. Front Neuroanat. 5.

Rossant C, Kadir SN, Goodman DFM, et al. Spike sorting for large, dense electrode arrays. *Nat*
Neurosci. 2016;19(4):634-641. doi:10.1038/nn.4268

Schaeren-Wiemers N, Gerfin-Moser A. 1993. A single protocol to detect transcripts of various
types and expression levels in neural tissue and cultured cells: in situ hybridization using
digoxigenin-labelled cRNA probes. Histochemistry. 100:431–440.

Schubert D, Martens GJM, Kolk SM. 2014. Molecular underpinnings of prefrontal cortex
development in rodents provide insights into the etiology of neurodevelopmental disorders.
20:795–809.

Southwell DG, Paredes MF, Galvao RP, Jones DL, Froemke RC, Sebe JY, Alfaro-Cervello C, Tang Y,
García-Verdugo JM, Rubenstein JL, Baraban SC, Alvarez-Buylla A. 2012. Intrinsically
determined cell death of developing cortical interneurons. Nature. 491:109–113.

Spampanato J, Sullivan R. 2016. Development and physiology of GABAergic feedback excitation
in parvalbumin expressing interneurons of the mouse basolateral amygdala. Physiological
…. 4:e12664–15.

Tsujimoto S. 2008. The prefrontal cortex: functional neural development during early childhood.
The Neuroscientist. 14:345–358.

Van De Werd HJJM, Rajkowska G, Evers P, Uylings HBM. 2010. Cytoarchitectonic and
chemoarchitectonic characterization of the prefrontal cortical areas in the mouse. Brain
Struct Funct. 214:339–353.

Weinberger DR. 1986. Physiologic Dysfunction of Dorsolateral Prefrontal Cortex in
Schizophrenia. Archives of General Psychiatry. 43:114.

Yamada J, Okabe A, Toyoda H, Kilb W, Luhmann HJ, Fukuda A. 2004. Cl− uptake promoting
depolarizing GABA actions in immature rat neocortical neurones is mediated by NKCC1. The
Journal of Physiology. 557:829–841.

Yang JM, Zhang J, Yu YQ, Duan S, Li XM. 2013. Postnatal Development of 2 Microcircuits Involving
Fast-Spiking Interneurons in the Mouse Prefrontal Cortex. Cerebral Cortex. 24:98–109.

Yuan Y, Malley HAOX, Smaldino MA, Bouza AA, Hull JM, Isom LL. 2019. Delayed maturation of
GABAergic signaling in the Scn1a and Scn1b mouse models of Dravet Syndrome. Scientific
Reports. 1–16.

Zhang Z, Jiao Y-Y, Sun Q-Q. 2011. Developmental maturation of excitation and inhibition balance
in principal neurons across four layers of somatosensory cortex. Neuroscience. 174:10–25.

Zheng K, An JJ, Yang F, Xu W. 2011. TrkB signaling in parvalbumin-positive interneurons is critical
for gamma-band network synchronization in hippocampus. Proc Natl Acad Sci USA; 108
(41):17201-17206;

Zhu JJ. 2000. Maturation of layer 5 neocortical pyramidal neurons: amplifying salient layer 1 and
layer 4 inputs by Ca2+ action potentials in adult rat tuft dendrites. The Journal of Physiology.
526:571–587.

Figures and figure legends

Figure 1

Figure 1. Properties of sIPSCs and sEPSCs at P10 and P20 of layer II/III mPFC and BC pyramidal neurons

(a) Representative traces of spontaneous inhibitory postsynaptic currents (sIPSCs) from layer II/III mPFC (left) and BC (right) pyramidal neurons at P10 (green) and P20 (black).

(b) Bar graph showing the sIPSC frequency (Hz) at P10 and P20 mPFC and BC pyramidal neurons. Two-way ANOVA analyses showed a significant effect of age ($F_{(1,26)}$=19.286, p<0.001) and brain

area ($F_{(1,26)}=12.554$, p=0.002) and a trend towards significance of age*brain interaction ($F_{(1,26)}=3.826$, p=0.063).

(c) Bar graph showing the sIPSC peak amplitude at P10 and P20 of mPFC and BC pyramidal neurons. Two-way ANOVA analyses showed a trend towards significance in the brain area factor ($F_{(1,26)}=3.420$, p=0.077) and no effect of age ($F_{(1,26)}=1.981$ p=0.177) or area*age interaction ($F_{(1,26)}=0.050$, p=0.826).

(d) Bar graph showing the sIPSC decay time constant (τm) at P10 to P20 of mPFC and BC pyramidal neurons. Two-way ANOVA analyses did not show any significant effect of age ($F_{(1,26)}=1.129$, p=0.299), brain area ($F_{(1, 26)}=0.211$, p=0.651) or area*age interaction ($F_{(1, 26)}=1.052$, p=0.316) was found.

(e) Representative traces of spontaneous excitatory postsynaptic currents (sEPSCs) from layer II/III mPFC (left) and BC (right) pyramidal neurons at P10 (green) and P20 (black).

(f) Bar graph showing the sEPSC frequency at P10 to P20 of mPFC and BC pyramidal neurons. Two-way ANOVA analyses showed a significant effect of age ($F_{(1,25)}=5.273$, p=0.032) and brain area ($F_{(1,25)}=7.388$, p=0.013), but not a significant interaction age*brain area ($F_{(1,25)}=0.591$, p=0.450)

(g) Bar graph showing the sEPSCs peak amplitude at P10 to P20 of mPFC and BC pyramidal neurons. Two-way ANOVA analyses did not reveal significant effect of brain area ($F_{(1,25)}=0.853$, p=0.366), age ($F_{(1,25)}=0.144$, p=0.708) or interaction ($F_{(1,25)}=0.712$, p=0.408).

(h) Bar graph showing the sEPSCs decay time constant (τm) at P10 to P20 of mPFC and BC pyramidal neurons. Two-way ANOVA analyses showed no significant effect of age ($F_{(1,25)}=0.557$, p=0.463), brain area ($F_{(1,25)}=0.720$, p=0.405) or interaction ($F_{(1,25)}=1.128$, p=0.300).

*n=9-13 cells from 5-9 mice/age group.

Figure 2

Figure 2- sIPSCs and sEPSCs frequency at P10 and P20 of layer II/III mPFC and BC pyramidal

neurons

(a) Representative traces showing that CNQX+AP5 did not alter the sIPSC frequency either at

P10 or P20 mPFC (recorded at +10mV).

(b) Representative traces showing that bicuculine blocks sIPSCs either at P10 or P20 mPFC

(recorded at +10mV).

(c) Graphs showing that bicuculine blocks the sIPSCs in P10 and P20 mPFC (t-test, p=0.001), while

CNQX+AP5 does not have an effect (t-test, p=0.5).

(d) Representative traces showing that CNQX reduces sEPSC frequency at p10 and blocks sEPSCs

at P20 mPFC (recorded at -70mV).

(e) Representative traces showing that bicuculine reduces sEPSC frequency at p10 and does not

affect sEPSCs at P20 mPFC (recorded at -70mV).

(f) Graphs showing that at P10 mPFC both CNQX+AP5 and bicuculine reduces the sEPSC frequency in P10 mPFC (t-test, p=0.03 for both drugs), while in P20 mPFC only CNQX+AP5 blocks sEPSCs.

Figure 3

Figure 3 – The reversal potential for evoked IPSCs is depolarized at p10 mPFC compared to p20

(a) Representative traces of evoked IPSCs at p10 and p20 mPFC

(b) Graph showing the average of eIPSCs at different holding potentials.

Figure 4

Figure 4. Changes in the chloride transporters mediate the depolarizing action of GABA$_A$R at p10 mPFC

fEPSPs were recorded were recorded in layer II/III mPFC in response to current pulses of increasing stimulus strength of layer II/III, during two experimental treatments, before and after application of 2μM diazepam (GABA$_A$R agonist) at P10 and P20.

(a) Representative traces (left) and graph (right) showing the fEPSPs amplitude before (green) and after (dark green) diazepam bath application (in the presence of CNQX and AP5), in mPFC at P10. Two-way repeated measures ANOVA analyses of evoked fEPSPs revealed significant effect of stimulus strength ($F_{(1,7)}$ = 20.64, p=0.004) and experimental treatments ($F_{(1,7)}$ = 5.025, p=0.036) (n=2 brain slices from 3-4 mice).

(b) Graph (right) and representative traces (left) showing that diazepam bath application does

not have any effect on the fEPSP amplitude in mPFC at P20 (in the presence of CNQX and AP5).

Two-way repeated measures ANOVA analyses of evoked fEPSPs revealed a significant effect of

stimulus strength ($F_{(1,7)}$ = 10.36, $p<0.0001$) but not experimental treatment ($F_{(1,7)}$=0.03,

p=0.9382), (n=2 brain slices from 3-4 mice).

(c) Graph showing that the effect of diazepam on fEPSP is occluded in the presence of

bumetanide (plus CNQX and AP5) at P10. Two-way repeated measures ANOVA analyses of

evoked fEPSPs did not reveal a significant effect of experimental treatment ($F_{(1,7)}$=0.08, p=0.752),

(n=2 brain slices from 3-4 mice).

(d) Graph showing the effect of diazepam on fEPSP in the presence of bumetanide (plus CNQX

and AP5) at P20. Two-way repeated measures ANOVA analyses of evoked fEPSPs did not reveal

a significant effect of experimental treatment ($F_{(1,7)}$=0.06, p=0.831), (n=2 brain slices from 3-4

mice).

(e) Representative blots showing changes of the K-Cl co-transporter (KCC2) levels, relative to

GAPDH at P10 and P20 in mPFC and BC.

(f) Graph showing the normalized protein level (KCC2/GAPDH) in mPFC at P10 and P20. The KCC2

protein levels was significantly increased at P20 compared to P10 in mPFC (two-tailed t-test, p=

0.01) but not in BC (two-tailed t-test, p= 0.97) (n=3-4 mice).

1001

1002

1003

1004

1005

1006

1007

1008

1009

1010

1011

1012

1013

1014

Figure 5

Figure 5. Poor development of active membrane properties of Lhx6[+] interneurons in mPFC.

(a) Bar graph showing the action potential (AP) amplitude of interneurons at P10 and P20 in mPFC and BC. Two-way ANOVA analyses did not show any significant effect of age ($F_{(1,22)}$=2.46, p=0.13) or brain area ($F_{(1, 22)}$=2.13, p=0.15) was found., (n=6-9 cells from 5-6 mice/age group).

(b) Bar graph showing the AP rate of rise (dv/dt) of interneurons at P10 and P20 in mPFC and at P10 in BC. Two-way ANOVA analyses showed a significant effect between ages ($F_{(1,20)}$=58.96, p<0.0001) but not brain area ($F_{(1,20)}$=0.16 p=0.69). Post-hoc analysis showed that the AP rate of rise significantly increased at P20 compared to P10 in mPFC (Tukey's test, p=0.0034) and at P20 compared to P10 in BC (Tukey's test, p<0.001), (n=6-9 cells from 5-6 mice/age group).

(c) Bar graph showing the AP duration (half-width) of interneurons at P10 and P20 in mPFC and BC. Two-way ANOVA analyses showed a significant effect between ages ($F_{(1,21)}$=39.16, p<0.0001) but not brain area ($F_{(1,21)}$=0.16 p=0.73). Post-hoc analysis showed that the AP duration significantly decreased at P20 compared to P10 in mPFC (Tukey's test, p= 0.0093) and at P20 compared to P10 in BC (Tukey's test, p=0.0002), (n=6-9 cells from 5-6 mice/age group).

(d) Bar graph showing the AP rheobase of interneurons at P10 and P20 in mPFC and BC. Two-way ANOVA analyses did not show any significant effect of age ($F_{(1,20)}$=1.60, p=0.22) or brain area ($F_{(1, 20)}$=0.40, p=0.53) was found.

(e) Bar graph showing the AP threshold of interneurons at P10 and P20 in mPFC and BC. . Two-way ANOVA analyses showed significant effect of age ($F_{(1,22)}$=5.048, p=0.035) and brain area ($F_{(1,22)}$=8.00, p=0.009) was found. Post-hoc analysis showed that the AP threshold was not significantly different at P20 compared to P10 in mPFC (Tukey's test, p=0.1673) and in BC (Tukey's test, p=0.72009) or at P10 in mPFC compared to P10 in BC (Tukey's test, p=0.067) and at P20 in mPFC compared to P20 in BC (Tukey's test, p=0.72).

(f) Bar graph showing the AHP (afterhypolarization) amplitude of interneurons at P10 and P20 in mPFC and BC. Two-way ANOVA analyses showed significant effect of age ($F_{(1,18)}$=7.35, p=0.0143) and brain area ($F_{(1,18)}$=63.72, p<0.0001) was found. Post-hoc analysis showed that the AHP amplitude was not significantly different at P20 compared to P10 in mPFC (Tukey's test, p=0.7187) and was significantly decreased in mPFC compered to BC, at P10 (Tukey's test, p=0.0005) and in mPFC compered to BC at P20 (Tukey's test, p<0.00001), (n=6-9 cells from 5-6 mice/age group).

(g) Bar graph showing the AHP time of interneurons at P10 and P20 in mPFC and BC. Two-way ANOVA analyses did not show any significant effect of age ($F_{(1,19)}$=0.009, p=0.92) or brain area ($F_{(1,19)}$=1.074, p=0.31) was found.

(h) Representative traces of APs of layer II/III Lhx6+ interneurons in mPFC (left) and BC (right) at P10 (green) and P20 (black).

Figure 6

1053

Figure 6. Passive membrane properties of Lhx6+ interneurons at P10 and P20 mPFC and P10 BC.

(a) Representative voltage responses (top traces) to 500ms positive and negative current pulses (bottom traces, +50, -50, -70, -100, -150, -200 pA) in mPFC at P10 and P20 and BC of Lhx6+ florescent interneurons from layer II/III.

(b) Bar graph showing the resting membrane potential (RMP) of interneurons at P10 and P20 in mPFC and BC. Two-way ANOVA analyses did not show any significant effect of age ($F_{(1,25)}$=1.55, p=0.22) or brain area ($F_{(1, 25)}$=0.50, p=0.48) was found., (n=6-9 cells from 5-6 mice/age group).

(c) Bar graph showing the input resistance of interneurons at P10 and P20 in mPFC and BC. Two-way ANOVA analyses showed a significant effect of age ($F_{(1,27)}$=10.94, p=0.0027) and brain area ($F_{(1,27)}$=6.65 p=0.0157). Post-hoc analysis showed that the input resistance significantly decreased at P20 compared to P10 in mPFC (Tukey's test, p=0.0006) and was significantly higher in mPFC compared with BC, at P10 (Tukey's test, p=0.0017), (n=8-9 cells from 5-6 mice/age group).

(d) Bar graph showing the membrane time constant (τ_m) of interneurons at P10 and P20 in

mPFC and BC. Two-way ANOVA analyses showed a significant effect of age ($F_{(1,24)}$=14.71,

p=0.0008) and brain area ($F_{(1,24)}$=6.92 p=0.0147). Post-hoc analysis showed that τ_m was

significantly higher at P10 compared to P20 in mPFC (Tukey's test, p=0.0006) while it was

significantly higher in mPFC compared to BC, at P10 (Tukey's test, p=0.0023), (n=8-9 cells from

5-6 mice/age group).

(e) Bar graph showing the membrane capacitance (C_m) of interneurons at P10 and P20 in mPFC

and BC. Two-way ANOVA analyses showed a significant effect between brain areas ($F_{(1,21)}$=6.82,

p=0.00163) and not between ages ($F_{(1,21)}$=2.60 p=0.1219). Post-hoc analysis showed that Cm was

significantly higher at P10 compared with P20 in mPFC (Tukey's test, p=0.0291) and was not

significantly different between mPFC and BC, at P10 (Tukey test, p=0.97) while it was significantly

higher at P20 in mPFC compared to P20 in BC (Tukey test, p=0.0202), (n=6-9 cells from 5-6

mice/age group).

1081

1082

1083

1084

1085

Figure 7

Figure 7. Significant differences in cellular density of Lhx6$^+$ interneurons in mPFC and BC at P10 and P20.

(a) A representative immunostaining with GFP for Lhx6$^+$ interneurons in Lhx6-cre;ROSA26fl-STOPfl-YFP mice in mPFC and BC at P20 is showing on the left. Scale bars: 150 µm. On the right, bar graph comparing Lhx6$^+$ interneurons cell density (per mm^2) at P10 and P20 in mPFC and BC. Two-way ANOVA analyses of the cell density revealed a significant effect of brain area ($F_{(1, 18)}$ = 13.11, p=0.0020), but not of age ($F_{(1,18)}$=0.7185, p=0.4078). Post-hoc analysis showed that the Lhx6$^+$ cell density was not significant different at P20 compared to P10 in mPFC and BC (LSD test, p=0.77 and p= 0.38, respectively). The Lhx6$^+$ cell density was significantly lower in mPFC compared to BC at P10 and P20, respectively (LSD test, p=0.0077 and p= 0.0440, respectively), (P10 in mPFC and BC:n= 5 mice, P20 in mPFC and BC: n=4).

(b) A representative in situ hybridization staining for somatostatin positive cells (SST$^+$) using wild type animals in mPFC and BC at P20 is showing on the left. Scale bar: 200µm. Bar graph comparing cell density based on SST$^+$ expression at P10 and P20 in mPFC and BC. Two-way ANOVA analyses of the cell density showed no significant effect of age ($F_{(1, 13)}$= 2.78, p=0.12) and brain area ($F_{(1,13)}$=071, p=0.41) was found, (P10 in mPFC: n= 5 mice, P10 in BC and P20 in mPFC and BC: n=4).

(c) A representative double immunostaining for GFP; PV (PV: parvalbumin) in mPFC and BC at P20 is showing on the left. Scale bars: 150 µm. On the right, bar graph comparing cell density based on PV$^+$ expression at P10 and P20 in mPFC and BC. Two-way ANOVA analyses of the cell density revealed a significant effect of age ($F_{(1, 14)}$ = 45.49, p<0.0001) and brain area ($F_{(1, 14)}$ = 170.2, p<0.0001). PV$^+$ cells were not found in mPFC but were identified in BC, at P10. Post-hoc analysis showed that the PV$^+$ cell density was not significantly different at P20 compared to P10 in BC (LSD test, p= 0.1089), but was significantly lower in mPFC compared to BC at P20 (LSD test, p<0.0001), (P10 and P20 in mPFC: n= 5 mice, P10 and P20 in BC: n=4).

1119

Figure 8

1120
1121

Figure 8. Active properties of mPFC and BC pyramidal neurons.

(a) Representative traces of action potentials (APs) of layer II/III pyramidal neurons in mPFC (left) and BC (right) at P10 (green) and P20 (black), respectively.

(b) Bar graph showing the AP amplitude of pyramidal neurons at P10 and P20 in mPFC and BC. Two-way ANOVA analyses showed a significant effect of age ($F_{(1, 31)}$ = 18.74, p=0.0001) but not on brain area ($F_{(1, 31)}$ = 0.99, p=0.32) was found. Post-hoc analysis showed that the AP amplitude significantly increased at P20 compared to P10 in mPFC and BC (Tukey's test, p=0.0386 and p= 0.0131, respectively) (n=9-14 cells from 6-10 mice/age group).

(c) Bar graph showing the AP rate of rise (dv/dt) of pyramidal neuron at P10 and P20 in mPFC and BC. Two-way ANOVA analyses showed a significant effect of age ($F_{(1,30)}$= 13.53, p=0.0009) but not on brain area ($F_{(1, 30)}$ = 0.36, p=0.55) was found. Post-hoc analysis showed that the AP rate of rise significantly increased at P20 compared to P10 in mPFC (Tukey's test, p= 0.0095), but not in BC (Tukey's test, p= 0.25) (n=8-14 cells from 6-10 mice/age group).

(d) Bar graph showing the AP duration (half-width) of pyramidal neuron at P10 and P20 in mPFC and BC. Two-way ANOVA analyses showed no significant effect of age ($F(1, 33) = 0.52$, $p=0.47$) or brain area ($F_{(1, 33)} = 0.43$, $p=0.51$) was found (n=9-14 cells from 6-10 mice/age group).

(e) Bar graph showing the AP rheobase of pyramidal neuron at P10 and P20 in mPFC and BC. Two-way ANOVA analyses showed no significant effect of age ($F_{(1, 36)} = 0.66$, $p=0.41$) or brain area ($F_{(1, 36)} = 0.16$, $p=0.69$) was found (n=9-14 cells from 6-10 mice/age group).

(f) Bar graph showing the AP threshold of pyramidal neuron at P10 and P20 in mPFC and BC. Two-way ANOVA analyses showed no significant effect of age ($F_{(1,31)} = 1.90$, $p=0.17$) or brain area ($F_{(1,31)}=0.55$, $p=0.46$) was found (n=9-14 cells from 6-10 mice/age group).

Figure 9. Modeling p10 and p20 PFC network activity

(a,b) Neuronal response of a single neuron within the PFC model network at p10 and p20

(c, d) The filtered voltage response of the p10 and p20 model networks

(e) Average frequency of the P10 and P20 model networks.

(f) Graph showing how the responses change in modified P10 model networks. N1: control P10 model network, N2: pyramidal model neuron of P20 model network used, N3: fast-spiking interneuron model network of P20 model network used, N4: GABAAR reversal potential was set to -60mV (that of P20 model network).

(g) Graph showing how the responses change in modified P20 model networks. J1: control P20 model network, J2: pyramidal model neuron of P10 model network used, J3: fast-spiking

1160   interneuron model network of P10 model network used, J4: GABAAR reversal potential was set

1161   to -40mV (that of P10 model network).

1162

Figure 10

Figure 10. Spike activity in PFC of neonatal and juvenile mice.

(a) Extracellular LFP recordings of oscillatory activity in PFC from a P9 mouse displayed after bandpass (1-100 Hz) filtering (top) and the corresponding MUA after bandpass (500-5000 Hz) filtering (bottom). Traces are accompanied by the color-coded wavelet spectra of the LFP at identical time scale.

(b) the same display as (**a**), but in one P22 mouse.

(c) Bar diagram displaying the mean MUA of neurons in PFC of neonatal and juvenile mice. During development, significant increase of MUA in PFC in juvenile mice (n=14) compared with in neonatal mice (n=13) (1.71±0.16 vs. -2.29±0.45, p<0.0001, One-way ANOVA, $F_{(1, 25)}$=80.19).

(d) Bar diagram displaying the percentage of the burst activity of single units in PFC of neonatal and juvenile mice. During development, significant more burst activity per single unit in PFC in juvenile mice (150 single unit) than in neonatal mice (161 single unit) (55.73±0.92 % vs. 27.90±1.41 %, p<0.0001, One-way ANOVA, F(1, 309)=272.26). In neonatal mice, 21 out 161 single

1178 units were classified as "burst unit". In juvenile mice, 104 out of 150 single units were classified

1179 as "burst unit".