Geometric models for robust encoding of dynamical information into embryonic patterns

Laurent Jutras-Dubé [1], Ezzat El-Sherif [2] *, Paul François [1] *

1 Department of Physics, McGill University, 3600 rue University, H3A 2T8, Montreal, QC, Canada

2 Division of Developmental Biology, Department of Biology, Friedrich-Alexander-Universität Erlangen-Nürnberg, Erlangen 91058, Germany

* Corresponding Authors

Abstract

During development, cells gradually assume specialized fates via changes of transcriptional dynamics, sometimes even within the same developmental stage. For anterior-posterior (AP) patterning in metazoans, it has been suggested that the gradual transition from a dynamic genetic regime to a static one is encoded by different transcriptional modules. In that case, the static regime has an essential role in pattern formation in addition to its maintenance function. In this work, we introduce a geometric approach to study such transition. We exhibit two types of genetic regime transitions, respectively arising through local or global bifurcations. We find that the global bifurcation type is more generic, more robust, and better preserves dynamical information. This could parsimoniously explain common features of metazoan segmentation, such as changes of periods leading to waves of gene expressions, "speed/frequency-gradient" dynamics, and changes of wave patterns. Geometric approaches appear as possible alternatives to gene regulatory networks to understand development.

Introduction

Development from one zygote to a viable animal is a complex process (Wolpert et al., 2006), comprising multiple dynamical sub-processes, including cell movements, tissue morphogenesis, dynamical gene expressions, and cellular differentiations. Eventually, cell identities are fixed by various mechanisms, such as multistable gene regulatory networks and epigenetic markers. Little is known about how this transition from a dynamic/initiation phase to a static/maintenance one is mediated. Are there general characteristics that should be matched between dynamic and static phases to mediate a robust transition?

In dynamical systems theory, a transition between different regimes is called a 'bifurcation', defined as a qualitative change in the dynamics of a system driven by a so-called 'control parameter' (Strogatz, 2015). Bifurcations are of many types but can be systematically classified. For instance, generic families of potentials driving the dynamics have been identified as different "catastrophes" (Poston & Stewart, 2012). While such mathematical descriptions are highly technical, they are reminiscent of the theory of epigenetic landscapes pushed forward by Waddington (Waddington, 1957). It is thus natural to ask if such classifications can be done for development. Could dynamical systems theory help us in this pursuit, and in studying development in general? Here, the main issue is to find a way to frame the problem to derive general results.

In recent years, numerous experimental studies have revealed that quantitative changes of gene expressions during development often followed standard predictions

from dynamical systems theory (Huang et al., 2007). The Waddington landscape's analogy (Jaeger & Monk, 2014) has led to many insights in cell differentiation (Graf & Enver, 2009), and recent data on cell reprogramming quantitatively validated the associated "landscape picture" (Pusuluri et al., 2018). Geometric models of development have been developed in particular cases, precisely predicting the general phenotypes of wildtype and mutants (e.g. the development of *C. elegans* vulva (Corson & Siggia, 2012) and *Drosophila* bristle patterns (Corson et al., 2017)).

The Clock-and-Wavefront model (Cooke & Zeeman, 1976), accounting for the observed dynamical somite (vertebrae precursors) formation, was inspired by catastrophe theory. The model predicted that a retracting wavefront translates the periodic expression of a genetic clock into a spatial pattern via "catastrophic" transitions demarcating the positions of the somites (Figure 1A). Identification of the predicted clock in 1997 (Palmeirim et al., 1997) has since led to many subsequent theoretical and experimental works, including observation of similar clocks in multiple arthropods (El-Sherif et al., 2012; Sarrazin et al., 2012). Cooke and Zeeman originally assumed that the clock is an external process, blind to the subsequent segmentation process it directs (Cooke & Zeeman, 1976). However, it has been very clear from the early experiments in (Palmeirim et al., 1997) that cellular oscillators increase their period prior to segmentation, leading to traveling waves of various signalling pathways such as Notch (Giudicelli et al., 2007; Morelli et al., 2009) (Figure 1A). Importantly, Notch waves eventually stabilize into a pattern of *delta* ligand stripes (Giudicelli & Lewis, 2004; Jiang et al., 2000), with a functional continuity between the dynamic and the static regime. Indeed, it has been shown that the dynamical phase of the clock is encoded into static

rostro-caudal identities (Oginuma et al., 2010). This suggests that the observed

oscillation is not a simple external pacemaker for segment formation: rather, clocks,

associated waves and the resulting stripes combine into an emergent process leading

to proper fate encoding. Segmentation thus possibly appears as the canonical example

of transition from a dynamical gene expression regime to a static functional one.

Two broad scenarios have been proposed to model this process (see Figure 1). In the

first scenario, the period of the individual oscillators is diverging to infinity as they

become more anterior (or similarly, the frequency of the clock is approaching zero),

automatically giving rise to a fixed pattern (Figure 1B-F). This model corresponds to

Julian Lewis' model for *c-hairy 1* expression pattern in somitogenesis (appendix of

(Palmeirim et al., 1997)), and it is possible to experimentally quantify the period

divergence within this model (Giudicelli et al., 2007). This also corresponds to the

implicit scenario of many theoretical models assuming that the frequency of the clock

approaches zero as cells get more anterior, such as the models in (Ares et al., 2012;

Morelli & Jülicher, 2007), possibly with a sharp discontinuity suppressing period

divergence (Jörg et al., 2015). Those models are appealing owing to their simplicity,

since all behaviour is encoded in a dynamical frequency gradient (possibly mediated by

FGF (Dubrulle & Pourquié, 2004)). However it is unclear what happens from a

dynamical systems theory standpoint (a noteworthy exception being the proposal that

the gradient emerges through a singularity in phase similar to the Burger's equation

(Murray et al., 2013)). In particular, the pattern in this scenario literally corresponds to a

frozen clock, such that there is an infinite number of local steady states corresponding

to the frozen phases of the oscillators.

A second scenario grounded in dynamical systems theory has been proposed (François & Siggia, 2012). In this scenario, a genetic network transits from an oscillatory state to an ensemble of (stable) epigenetic states (in Waddington's sense) fixing the pattern. Possible examples include the initial reaction-diffusion based model by Meinhardt (Meinhardt, 1986), or the cell-autonomous model under morphogen control evolved in (François et al., 2007) (Figure 1G). Based on geometric arguments, if bifurcations are local, the most generic model of this transition is expected to present two steps as explained in (François & Siggia, 2012). As a steep control parameter (possibly controlled by a morphogen such as FGF) decreases, the oscillation dies out through a Hopf bifurcation, leading to a single transient intermediate state. Then, for even lower values of the morphogen, one or several new (stable) states appear (technically through saddle-node bifurcations, see Figure 1–figure supplement 1). If the system translates rapidly enough from the oscillatory regime to the multistable regime, a pattern can be fixed (Figure 1H-K). Contrary to the previous scenario where the period of the clock goes to infinity, a Hopf bifurcation is associated to a finite period when the clock stops. The pattern of gene expression itself is laid thanks to multiple expression states discretizing the phase of the clock (Figure 1—figure supplement 1). Importantly, a finite number of states are observed, e.g. anterior and posterior fates within one somite (as first pointed out by Meinhardt (Meinhardt, 1982)).

In this paper, we revisit those ideas with explicit modelling to characterize the behaviour of systems transitioning from a dynamical regime (such as an oscillation) to a static multistable regime. We introduce two new assumptions: 1. the two different phases of developmental expression (dynamic and static) can be separated into two independent

sets of transcriptional modules acting on several genes simultaneously, and 2. the

system smoothly switches from one set to the other. This proposal is motivated by the

recent suggestion in insects that different sets of enhancers control waves of gap genes

at different phases of embryonic growth (El-Sherif & Levine, 2016). Such assumptions

explain the so-called "speed-gradient" model suggested to explain the gene expression

wave dynamics observed during AP patterning in the beetle *Tribolium* (Zhu et al., 2017)

(see Figure 1—figure supplement 2) and (with some additional assumptions) the more

subtle gene expression dynamics observed during AP patterning in (Rudolf, Zellner, & El-

Sherif, 2020). Using both gene-network and geometric formalisms, we characterize the

types of bifurcations found in systems transitioning from a dynamic to a static regime.

Surprisingly, we find that if the transition is smooth enough, global bifurcations appear.

This situation is different from the standard scenario (Hopf and saddle-nodes) that we

nevertheless recover if the transition is more non-linear. This is a generic result that is

better studied and understood using geometric models. We further show that the

transition through a global bifurcation is more robust than the sequence of Hopf and

saddle-node bifurcations with respect to several perturbations that we simulate. Finally,

we find that this model can explain many features of metazoan segmentation, such as

"speed-gradient" mechanisms or changes of spatial wave profiles due to relaxation-like

oscillations, and we discuss biological evidence and implications. This geometric

approach thus offers a plausible scenario underlying embryonic patterning with many

associated experimental signatures.

Model

In the following, we consider a class of developmental models based on the combination of (at least) two different transcriptional modules. Biologically, those two modules correspond to two sequential developmental phases. The main assumptions are that those transcriptional modules are globally regulated for multiple genes at the same time (which could be done for instance through chromatin global regulations) and that there is a continuous transition from one to the other. Here we focus on metazoan segmentation and regionalization, but the formalism might be applicable to other patterning processes where both an initiation phase and a maintenance phase have been described.

We use ordinary differential equations to model our system. Calling P a vector encoding the state of all proteins in any given cell (typically P corresponds to concentrations of proteins), a generic single-cell equation describing all models presented in the following is:

$$\dot{P} = \theta_D(g)\, D(P) + \theta_S(g)\, S(P) + C(P) + \eta(g, P) \qquad (1)$$

In Eq. 1, variable g encodes an external control parameter of the developmental transition. For example, g could be an external morphogen concentration driving patterning, but more complex situations with feedback are possible, where g could also be part of the system (e.g. the phase difference between oscillators (Beaupeux & François, 2016; Sonnen et al., 2018)). For simplicity, we rescale variables so that g is constrained between 0 and 1. The terms $D(P)$ and $S(P)$ correspond to different sets of modules, their influence on the dynamics being weighted by functions $\theta_D(g)$ and $\theta_S(g)$,

respectively. The term $\eta(g, P)$ encodes the noise. Finally, $C(P)$ represents dynamical

terms that are independent of the transcriptional modules, such as protein degradation.

We focus here on the simplest two-module case, where $S(P)$ encodes a multistable

system (i.e. with multiple fixed points at steady state) and $D(P)$ a dynamic system (i.e.

oscillatory). In this situation we will assume $\theta_S(0) = 1$, $\theta_S(1) = 0$, $\theta_D(0) = 0$, and

$\theta_D(1) = 1$, meaning that for $g = 1$ the network is in a pure dynamic phase, while for

$g = 0$ the network is multistable. Details on the specific forms of $D(P)$, $S(P)$, $\theta_D(g)$ and

$\theta_S(g)$ are given in the following and in the Appendix. We study two types of models:

gene-network like models where $D(P)$ and $S(P)$ explicitly model biochemical

interactions between genes (such as transcriptional repression), and geometric models

where $D(P)$ and $S(P)$ directly encode flows in an abstract 2D phase space, similarly to

(Corson & Siggia, 2017).

We model an embryo as a line of cells, corresponding to the antero-posterior axis. The

dynamics within each cell (position x) is described by Eq. 1. The only difference

between cells is that the dynamics of g is a prescribed function of x, e.g. we assume

that there is a function $g(x, t)$ describing the dynamics of a morphogen. We focus on the

transition between the two regimes as g continuously changes from 1 to 0 in different

cells as a function of time. We will typically consider a sliding morphogen gradient

moving along the antero-posterior axis with speed v, described by $H(s(x - vt))$ where

the function H encodes the shape of the morphogen, and parameter s is a measure of

the gradient's spatial steepness.

We also include noise in the system with the help of an additive Gaussian white noise.

For gene networks, we follow an approach similar to the τ-leaping algorithm (Gillespie,

2001), where the variance of the noise corresponds to the sum of the production and

the degradation terms (approximating independent Poisson noises). A multiplicative

noise intensity term $\sqrt{1/\Omega}$ is introduced, where Ω can be interpreted as the typical

concentration of the proteins in the system, so that bigger Ω corresponds to lower noise.

In addition, we add diffusion coupling the cells in the stochastic gene network models.

For the geometric model, the variance of the noise is held independent of the position x.

A more detailed description of the noise and diffusion terms is provided in the Appendix.

All source codes and data used for this paper are available at :

https://github.com/laurentjutrasdube/Dual-

Regime_Geometry_for_Embryonic_Patterning

Relation to existing biological networks

The model described above aims at being generic, but one can relate it to existing

developmental networks. A two modules dichotomy was initially proposed in the

Drosophila context, where two enhancers (early and late) were observed for *Krüppel*

and *knirps* (E. El-Sherif & Levine, 2016). An extension of this model has been proposed in

(Zhu et al., 2017) for a gap gene cascade under control of the maternal gene *Caudal*,

which would thus play the role of g (a detailed model is reproduced in the Supplement,

see Figure 1—figure supplement 2). The dynamic module corresponds to a genetic

cascade comprising *hunchback, Krüppel, mille-pates,* and *giant.* Each gene in this

sequence activates the next one, and later genes repress earlier ones. For the static

module, each gene self-activates, and *hunchback* and *Krüppel* repress one another.

The situation is less clear for vertebrates, given the plethora of oscillating genes and

possible redundancy in three different pathways, specifically *Notch, Wnt* and *FGF*

([Dequeant et al., 2006](#)). The two-module system we consider assumes both the dynamic

and static regimes are realized by the same set of genetic components (genes and/or

signalling pathways). *Notch* signalling is an ideal candidate to be a component of both a

dynamic and a static regime that might mediate vertebrate somitogenesis, since Notch

is implicated in both the core segmentation clock of vertebrates (e.g. via *Lfng* ([Dale et al.,](#)

[2003](#)), or the *hes/her* family in zebrafish ([Lewis, 2003](#))) and oscillation stabilization ([Jiang](#)

[et al., 2000](#); [Oginuma et al., 2010](#)). Importantly, several genes of the Notch signalling

pathways (e.g. *DeltaC* in zebrafish) are first expressed in an oscillatory manner then

stabilize in striped patterns, as expected in our model ([F. Giudicelli & Lewis, 2004](#); [Wright](#)

[et al., 2011](#)). There are also multiple genetic interactions between members of the *Notch*

pathway, in particular again *Delta* genes ([Schwendinger-Schreck, Kang, & Holley, 2014](#)) ,

with different roles and changes of regulations in the dynamic vs static phase (see e.g.

([Wright et al., 2011](#))).The oscillation itself could be mediated through one or several

negative feedback loops in this pathway ([Lewis, 2003](#)), and stabilization could be realized

through one of the multiple *Notch* signalling interactions (possibly via cell coupling,

similarly to what is observed in other systems ([Corson, Couturier, Rouault, Mazouni, &](#)

[Schweisguth, 2017](#))).

Results

A model for the transition between two genetic modules: Hopf vs. SNIC.

In (Zhu et al., 2017), it was suggested that the transition from a "wave-like" behaviour to a static pattern during *Tribolium* segmentation was mediated by a smooth transition from one set of modules (corresponding to the oscillatory phase) towards another (corresponding to the fixed pattern). This explained the "speed-gradient" mechanism where the typical time-scale of the dynamical system depends strongly on an external gradient (in this case the concentration of the transcription factor *Caudal*). In the Appendix, we further study the associated bifurcation, and observe that new fixed points corresponding to the stabilization of gap gene expressions appear on the dynamical trajectory of those gap genes (Figure 1—figure supplement 2F). In simple words, the gap gene expression pattern slowly "freezes" without any clear discontinuity in its behaviour from the dynamic to the static phase, which is reminiscent of the "infinite-period" scenario displayed on Figure 1.

We first aim to generalize this observed property. A simple way to generate waves of gene expressions (as in the gap-gene system described above) is to consider an oscillatory process, so that each wave of the oscillation corresponds to a wave of gap genes. We are not saying here that the gap-gene system is an oscillator, but rather that its dynamics can be encompassed into a bigger oscillator (Verd et al., 2018)). The other advantage of considering oscillators is that we can better leverage dynamical systems theory to identify and study the bifurcations. Furthermore, it allows for a better connection with oscillatory segmentation processes in vertebrates and arthropods.

We thus start with an idealized gene regulatory network with 3 genes under the control

of two regulatory modules (Figure 2). In the dynamic phase $D(P)$, we assume that the 3

genes are oscillating with a repressilator dynamics (Elowitz & Leibler, 2000), so that the

system keeps a reference dynamical attractor and an associated period. In the static

phase $S(P)$, we assume that the module encodes a tristable system via mutual

repression (Figure 2A).

We study the dynamics in a simulated embryo under the control of a regressing front of

g (Figure 2B). Transition from the dynamic module to the static module is expected to

form a pattern by translating the phase of the oscillator into different fates, implementing

a clock and wavefront process similar in spirit to the one in (François et al., 2007). We

compare two versions of this model, presenting the two different behaviours that we

found. In Model 1 (Figure 2C-H), the weights of the two modules are non-linear in g:

$\theta_D(g) = g^2$ and $\theta_S(g) = (1-g)^2$ (Figure 2C). In Model 2 (Figure 2I-N), the weights of

the two modules are linear in g: $\theta_D(g) = g$ and $\theta_S(g) = 1-g$ (Figure 2I). We note that

the initial and final attractors of both models are identical. Importantly, only the

transition from one set of modules (and thus one type of dynamics) to the other is

different. This two-module system thus offers a convenient way to compare the

performance of different modes of developmental transition while keeping the same

"boundary conditions" (i.e. the same initial and final attractors).

Figure 2E and Figure 2K show the kymographs for both models without noise, with

behaviours of individual cells in Figure 2D and Figure 2J. While the final patterns of both

models are the same (Figure 2F and Figure 2L), giving rise to a repeated sequence of

three different fates, it is visually clear that the pattern formed with Model 2 is more

precise and sharper along the *entire dynamical trajectory* than the one formed with

Model 1, which goes through a "blurry" transitory phase (compare mid-range values of g

on Figure 2E and Figure 2K).

To better understand this result, we plot the bifurcation diagram of both models as a

function of g in Figure 2G and Figure 2M. As g decreases, Model 1 is the standard case

of a local Hopf bifurcation (Strogatz, 2015), which happens at $g = 0.72$. Three

simultaneous saddle-node bifurcations appear for lower values of g, corresponding to

the appearance of the fixed points defining the three regions of the pattern. The

behaviour of Model 2 is very different: the fixed points form on the dynamical trajectory,

via three simultaneous Saddle Node on Invariant Cycle (or SNIC) bifurcations (Strogatz,

2015). Both models display waves corresponding to the slowing down of the oscillators,

leading to a static regime. In Model 1, the time-scale disappears with a finite value

because of the Hopf bifurcation (Figure 2H). For Model 2, it diverges because of the

SNIC (Figure 2N), suggesting an explicit mechanism for the infinite-period scenario of

Figure 1.

To further quantify the differences of performance between the two models, we

introduce noise (encoded with variable Ω, see the Model section and the Appendix) and

diffusion (Figure 3A-D). We also define a mutual information metric measuring how

precisely the phase of the oscillator is read to form the final pattern (Figure 3E, see the

Appendix for details), consistent with the experimental observation in vertebrate

segmentation that oscillatory phases and pattern are functionally connected (Oginuma

et al., 2010). Intuitively, this metric quantifies in a continuous way the number of fates

encoded by the system at steady state. Ideal mutual information for the three mutually

exclusive genes of Models 1 and 2 gives $log(3) \sim 1.6$ bits of mutual information,

meaning that the pattern deterministically encodes the phase of the cycle into three

static fates with equal weights. While addition of noise decreases this mutual

information as expected (Figure 3E), Model 2 (black curves) always outperforms Model

1 (red curves). For a reasonable level of noise corresponding to a few thousands of

proteins in the system, Model 2 can encode $2^{1.3} \sim 2.5$ fates, close to the optimum 3.

Furthermore, for a given diffusion constant, Model 1 requires a ten times smaller noise

level than Model 2 to encode the same amount of mutual information, which thus

indicates much better noise resistance for Model 2.

Those observations suggest that appearance of stable fixed points through SNIC

bifurcations rather than through Hopf bifurcations generates a more robust pattern. The

superiority of Model 2 can be rationalized in the following way: when there is a Hopf

bifurcation, only one fixed point exists for a range of g values, so that all trajectories are

attracted towards it. This corresponds to the "blurred" zone in the kymographs of Figure

2 and Figure 3. In presence of noise, the effect is to partially erase the memory of the

phase of the oscillation when only one fixed point is present for the dynamics.

Conversely, a SNIC bifurcation directly translates the phase of the oscillation into fixed

points, without any erasure of phase memory, ensuring higher information transfer from

the dynamic to the static phase, and therefore more precise patterning.

The Hopf bifurcation of Model 1 occurs when the weights of the dynamic and static

modules become small compared to the degradation term, which generates an

"intermediate regime" with one single fixed point after the oscillations of the dynamic

module and before the multistability of the static module. The specific form of the

weights is not what determines the bifurcation, but rather the presence or absence of an

intermediate regime. We confirm this observation with similar 3-gene models that used

Hill functions for the weights θ_D and θ_S (Figure 2—figure supplement 1 and Figure 3—

figure supplement 1). Interestingly, we get both Hopf and SNIC bifurcations with the

same shape for the two weights; the Hopf is obtained by shifting the weight of the

dynamic term towards larger values of the control parameter. This effectively generates

the required intermediate regime where both weights are small compared to the

degradation term.

<u>Gene-free models present a similar geometry of transition</u>

Hopf and saddle-node bifurcations are "local" bifurcations, in the sense that changes of

the flow in phase space are confined to an arbitrarily small region of phase space as the

bifurcation is approached. They do not in principle require complex changes of the flow

or fine-tuning of the parameters to happen. As such, they are the most standard cases

in many natural phenomena and in most theoretical studies. Conversely, SNIC

bifurcations are "global" bifurcations (Strogatz, 2015) (Ermentrout, 2008): they are

associated to changes of the flow in large regions of phase space (e.g. when a limit

cycle disappears with a non-zero amplitude) and usually require some special

symmetries or parameter adjustments to occur (e.g. to ensure that a saddle-node

collides with a cycle).

It is therefore a surprise that SNIC bifurcations spontaneously appear in the models

considered here. To better understand how this is possible and if this is a generic

phenomenon, we follow ideas first proposed by Corson and Siggia (Corson & Siggia,

2012) and consider geometric (or gene-free) systems. We aim to see if: 1. SNIC

bifurcations are generically observed, and 2. a model undergoing a SNIC bifurcation is

in general more robust to perturbations than a model undergoing a Hopf bifurcation,

with initial and final attractors being held fixed. We thus build 2D geometric versions of

the system (variables y and z). The dynamic module $D(P)$ is defined by a non-harmonic

oscillator on the unit circle, while the static module $S(P)$ is defined by two stable fixed

points, at $y = \pm 1$, $z = 0$ (see Figure 4A, and the Appendix for the equations). Like

previously, we build a linear interpolation between the two systems as a function of g

and explore the consequence on the bifurcations (Figure 4B-H). Since the flow in the

system is 2D, we can also easily visualize it (Figure 4I and Figure 4—movie supplement

2).

In brief, this geometric approach confirms all the observations made on the gene

network model of the previous section, and further clarifies the origin of the SNIC

bifurcation. Because of the smooth transition between modules, the entire flow in 2D

needs to interpolate from a cycle to a bistable situation. When both modules have close

to equal weights (around $g = 0.5$), the flow and associated cycle concentrate around

two future fixed points. This appears in retrospect as the most natural way to interpolate

between the two situations since both types of attractor (stable limit cycle, and multiple

stable fixed points) are effectively present at the same time around $g = 0.5$. For this

reason, the oscillations are also more similar to relaxation oscillations, rapidly jumping

between two values corresponding to the future fixed points. When g is further lowered,

the weight of the static module dominates and "tears apart" the cycle, forming two fixed

points.

This situation is so generic that in fact, to obtain a Hopf bifurcation, we need to

mathematically reinforce the fixed point at $y = 0$ for intermediate values of g. More

precisely, three things are required to get a Hopf bifurcation. First, we need to add an

extra term, the "intermediate module", characterized by a single fixed point at $y = 0$ and

$z = 0$. Without this module, we consistently get a SNIC bifurcation, for all forms of

coupling tested, including linear and non-linear couplings (Figure 4—figure supplement

1A-D). Second, we need to set the weight of the intermediate module to 0 for $g = 1$ and

$g = 0$, so that we get the oscillations of the dynamic module at the beginning of the

simulation and the bistability of the static module at the end of the simulation. We

achieve this by setting the weight of the intermediate term to $g(1 - g)$, which is of 2^{nd}

order in g. Third, we need to make the weights of the dynamic and static modules

smaller than the weight of the intermediate module for intermediate values of the control

parameter, i.e. for g around 0.5. This is achieved by using cubic weights for the dynamic

and static modules (Figure 4—figure supplement 1G and Figure 4—figure supplement

2). Weights of the 4^{th} order in g also lead to a Hopf bifurcation (Figure 4—figure

supplement 1H), while linear weights lead to a SNIC bifurcation (Figure 4—figure

supplement 1E). Interestingly, quadratic weights lead to simultaneous supercritical Hopf

and pitchfork bifurcations (Figure 4—figure supplement 1F). The non-linearity of the

coupling helps reinforce the relative weight of the intermediate module for values of g

around 0.5, but the exact shape of the non-linearity is not crucial. Furthermore, our

mutual information metric confirms that the pattern is more robustly encoded when the

system goes through a SNIC bifurcation rather than through a sequence of Hopf and

pitchfork bifurcations (Figure 4—figure supplement 5F).

Detailed simulations of the gene-free model with the intermediate module and with cubic

weights for the dynamic and static modules are shown on Figure 4—figure supplement

2. As expected for a Hopf bifurcation, the flow first concentrates on the central fixed

point at $y = 0$, before re-emerging in a bistable pattern for lower g (Figure 4—figure

supplement 2I and Figure 4—movie supplement 1). There are technically two types of

Hopf bifurcations, depending on the stability of the limit cycle. A Hopf bifurcation is

supercritical (resp. subcritical) if a stable (resp. unstable) limit cycle becomes a stable

(resp. unstable) fixed point. All Hopf bifurcations discussed previously are supercritical.

However, by changing slightly the dynamic module of the gene-free model (and

including the intermediate module as well as cubic weights for the dynamic and static

modules) we can also obtain a subcritical Hopf bifurcation (Figure 4—figure supplement

3). During this bifurcation, an unstable limit cycle forms around the origin. This unstable

limit cycle coexists with the stable limit cycle of the dynamic module for some range of g

values before they annihilate each other during a so-called "saddle-node of limit cycles"

bifurcation. Finally, the two fixed points of the static module form during two

simultaneous saddle-node (of fixed points) bifurcations (Figure 4—figure supplement 3I

and Figure 4—movie supplement 3). Again, our mutual information metric confirms that

the pattern is more precise when the system goes through a SNIC bifurcation (Model 2),

rather than through supercritical or subcritical Hopf bifurcations (Models 1 and 3, resp.)

(Figure 4—figure supplement 5E). Taken together, these results suggest that keeping

the static and dynamic attractors fixed, patterning is both more generic and more

robustly encoded through a SNIC bifurcation than through a Hopf bifurcation, at least in

simple low-dimension models.

Robustness and asymmetry in the fixed points

A concern with the results of the previous section might be that those mathematical

models are in fact fine-tuned and too symmetrical, so that in particular when the

transition occurs, both new fixed points appear for the same value of the control

parameter. Furthermore, real biological networks have no reason to be perfectly

symmetrical (although evolution itself might select for more symmetrical dynamics if

needed). We thus relax our hypotheses to study a system where parameters and

trajectories are not symmetrical (Figures 5 and 6).

Going back first to the gene network model, we induce an asymmetry between the fixed

points by changing thresholds of repression in the static phase (Figure 5A). The

bifurcation diagrams of Figure 5B-C indicate that the asymmetry of the fixed points

indeed breaks the simultaneity of appearance of all fixed points in both scenarios. We

nevertheless notice that for those changes of parameters, all bifurcations still happen in

a very narrow range of g for the SNIC model.

Asymmetry of the fixed points might therefore destroy the advantage of SNIC vs Hopf

by creating a transient zone where one of the fixed points always dominates. We thus

perform a comparison between Models 1 and 2 with the same asymmetric static

enhancers (Figure 5, see also Figure 5—figure supplements 1 and 2, and the Appendix

for details). To compare the two cases, we consider different time-scales of the

morphogen gradient. The reasoning is that the slower the decay of g, the more time the

system spends in a region of parameter space without all three final fixed points,

allowing the system to relax and "lose" phase information. Conversely, a faster decay of

g means that less time is spent in a region with few fixed points, and therefore the

patterns are expected to be more robust.

We first decrease the thresholds of repression of gene A by both genes B and C (Figure

5A). Results of these simulations are shown in Figure 5: Model 2 with a SNIC

bifurcation still outperforms Model 1 with Hopf and saddle-node bifurcations. In

particular, it is again visually clear on kymographs that Model 2 produces a robust and

well-defined pattern at any time point of the simulations, while Model 1 gives rise to a

much "fuzzier" pattern before the transition. Model 1 produces a robust static pattern

only for a steep gradient (allowing to quickly move through the "fuzzy" phase) and a

weak asymmetry in the static module (Figure 5E). It is brittle to any change of the

dynamics of g (Figure 5H) or to stronger asymmetry in the static module (Figure 5—

figure supplement 1E,H). Conversely, Model 2 is robust to different shapes of the

morphogen (Figure 5F,I). Only for a strong asymmetry does the system lose one fixed

point (Figure 5—figure supplement 1I), but even in this case transitions through a SNIC

bifurcation appear superior to transitions through a Hopf bifurcation.

The fragility of the Hopf bifurcation to asymmetries in the parameters can be understood

as follows. In the asymmetric versions of Model 1, one of the fixed points of the static

term forms during the Hopf bifurcation, way before the two other fixed points form. It is

therefore the only attractor available for a large range of g values. However, in Model 2

the same asymmetry only favors one of the saddle-nodes for a small range of g values,

generating a robust pattern. Again, we can use the mutual information metric defined

above to quantify the robustness of the pattern and confirm the superiority of Model 2

(Figure 5—figure supplement 2J). We also confirmed these results for the case of

random modifications of the repression thresholds of all interactions in the static term

(Figure 5—figure supplement 2).

The asymmetry introduced in Figure 5 changes the shapes of the basins of attraction

and the positions of the fixed points. The geometric model allows to change those

features independently. The most generic way to introduce an asymmetry in the system

is to fix the positions of the fixed points of the static regime and change only the

positions of the basins of attraction (the reason is that the future fates depend on the

position of the separatrix between different regimes (Corson & Siggia, 2012)). To

replicate this situation in the 2D gene-free models, we move the unstable fixed point of

the static term along the y axis. Results of this procedure are shown on Figure 6 and

confirm our results on the gene-network based models: Model 2 bifurcates via a SNIC

and is always more robust than Model 1. When we change the positions of the fixed

points in the static regime to move them away from the limit cycle (still in an asymmetric

way), interestingly both Models 1 and 2 now bifurcate via SNICs (Figure 6—figure

supplement 1). Furthermore, we see that for Model 1, the amplitude of the limit cycle

decreases before the bifurcation, while for Model 2, the amplitude increases (Figure 6—

figure supplement 1E).

We conclude from all those numerical perturbations that even with asymmetric basins of

attraction and asymmetric parameters, transitions based on SNIC bifurcations are both

more generic and more robust than the ones based on Hopf bifurcations, at least in

simple low-dimension models.

Spatial wave profiles: Hopf vs SNIC

Our theoretical study suggest that SNIC based transitions are both more robust and

more generic than Hopf/saddle-nodes ones. We thus now examine ways to distinguish

between Hopf and SNIC based transitions experimentally. The natural method would be

to modify the value of the control parameter of the bifurcations (in our case parameter g)

and check how the attractors of the system change. A recent experimental example can

be found in the auditory hair-bundle context where it has been suggested that Hopf

bifurcations can be distinguished from SNIC bifurcations by tuning the external driving

force (Salvi, Maoileidigh, & Hudspeth, 2016). However, the situation in development is

different from any sensing system since the actual control parameters are not known

and are likely combinations of various signaling systems (such as *FGF* or *Wnt*). There

could also be many compensatory developmental mechanisms, making it difficult to fully

take control of the system, as well known by developmental geneticists.

In the absence of a direct control, we have to rely on indirect measurements. The most

obvious choice is to use the antero-posterior position along the embryo (or along the

PSM in the case of somite formation) as a proxy for the control parameter. In situs are

informative of what locally happens in insects (Ezzat El-Sherif, Averof, & Brown, 2012) (Zhu

et al., 2017). In vertebrates, it is possible to monitor in real time segmentation oscillators

in embryos (Aulehla et al., 2008) (Webb et al., 2016) (Delaune, François, Shih, & Amacher,

2012) and in tissue cultures (Lauschke, Tsiairis, François, & Aulehla, 2013) (Matsuda et al.,

2020) (Diaz-Cuadros et al., 2020), so that many properties of the cycles can be inferred.

A first relevant metric (used in (Salvi et al., 2016)) is the shape of the limit cycle as a

function of the control parameter. There is a clear contrast between Hopf and SNIC

bifurcations on our simulated kymographs: for Hopf bifurcations, as mentioned above

the transition zone is "blurred" because the damped oscillations relax towards the

unique fixed point, while for SNIC bifurcations the oscillations localize close to the fixed

points, and as a consequence the pattern is gradually reinforced (Figures 2-5; damped

oscillations are especially visible in the geometric model). To better visualize the "blurry"

transition zone characteristic of Hopf bifurcations in the gene-free model (resp. in the 3-

gene model), we compute the distribution of the values of the geometric variables (resp.

of the gene concentrations) for different values of parameter g (Figure 7—figure

supplement 2). In models with Hopf bifurcations, the distribution becomes very narrow

for intermediate values of g, which is a direct consequence of the damped oscillations

past the Hopf bifurcations. This suggests that measuring experimentally the distribution

of gene concentrations across developmental time at fixed antero-posterior positions

can help distinguish SNIC from Hopf bifurcations during actual pattern formation

processes. The experimental picture, both from in situs and live imaging, does not

support a narrowing of the distribution : it shows clear increases of amplitude that

seems more consistent with a gradual reinforcement of the pattern, similar to a SNIC

(see e.g. (Delaune et al., 2012)).

Another metric that could help identify the type of bifurcation is the time evolution of the

period (resp. of the frequency), which presents a discontinuity for both types of Hopf

bifurcations, but continuously goes to infinity (resp. zero) for SNIC bifurcations. The

local spatial wavelength of the pattern provides a continuous measurement related to

the local period (as first used and derived in (Francois Giudicelli, Ozbudak, Wright, & Lewis,

2007)). Calling S the somite size (or the wavelength of anterior/posterior markers within

somites after clock stopping), $T(x)$ and $S(x)$ the respective period and local wavelength

of the pattern at position x (with $x = 0$ being the tail and $x = 1$ the front), and assuming

the cells move towards the anterior with constant speed, we have

$$S(x) = \frac{S}{1 - \frac{T(0)}{T(x)}}$$

In the posterior $T(x) = T(0),$ so that $S(x)$ is infinite: all cells are synchronized locally. As

cells move towards the anterior, $T(x)$ decays and $S(x)$ decreases but stays bigger than

S. If there is a Hopf bifurcation, one expects that the clock will stop with a final period

$T(1)$, corresponding to a critical wavelength of the wave $S(1) = S \frac{T(1)}{T(1)-T(0)}$, which is

strictly greater than S. There is therefore in principle a discontinuity between the

wavelength of the propagating wave in the PSM and the wavelength of the static pattern.

Conversely, if $T(x)$ goes continuously to infinity, $S(x)$ decays continuously to converge

towards S (as measured in (Francois Giudicelli et al., 2007)). This is a simple and intuitive

experimental outcome: in this situation the wavelength of the propagating wave in the

PSM decreases until it exactly matches the pattern of anterior markers. Therefore, a

signature of the finite vs infinite period bifurcation might be visible by comparing the

local wavelength of propagating waves in the PSM to the wavelength of anterior

markers in the pattern. Practically, monitoring the distance between the peaks of the

oscillations does not give a clear difference between Hopf and SNIC, the reason being

that such a discrete measurement can smooth out an abrupt change in the period and

that past the Hopf bifurcation, damped oscillations actually give rise to a "ghost"

frequency which can decrease quite significantly (see Figure 2—figure supplement 2).

Therefore, a measurement of the local wavelength might be less conclusive than a

measurement of the amplitude. Nevertheless, we notice that a SNIC scenario gives rise

to a combination of an increase of the amplitude with a continuous shrinking of the

wavelength, meaning that many refining waves (corresponding to broad variations of

periods) will simultaneously propagate.

Finally, SNIC bifurcations are also accompanied by characteristic changes of the shape

of spatial wave profiles that might be observable in experiments. We further compare a

SNIC-based model to a phase model where an infinite-period behaviour is explicitly

assumed, namely the model of a collection of coupled oscillators from (Morelli et al.,

2009). A kymograph of the spatio-temporal profile of the frequency imposed on the

oscillators is shown in Figure 7A, and the dynamics of the resulting pattern formation

process is shown on the kymograph of Figure 7B, with the final pattern on Figure 7C.

The most striking difference is observed on the shape of the spatial wave profile as it

moves towards the region where the pattern stabilizes. In the infinite-period scenario of

(Morelli et al., 2009), the phase profile is by construction symmetric (albeit stretched in

the posterior compared to the anterior, see Figure 7E). In the SNIC scenario, we see a

clear asymmetry in the peaks of the wave profile (as shown schematically in Figure 7D).

Indeed, following the peaks of oscillations spatially from anterior to posterior (left to

right), we see that the transition from positive to negative values of y occurs in two steps.

During the first step, the system stays close to $y = 1$ for some time, with y values

decreasing slowly, while in the second step the system goes rapidly towards negative y

values and reaches $y = -1$ fast (Figure 7F). This phenomenon, which gives rise to a

"sawtooth" pattern of propagating waves, is observed in all our versions of Model 2 (and

is notably absent from all our versions of Model 1, see Figure 7—figure supplement 1).

Those spatial asymmetries are likely due to the asymmetries of the limit cycles of

relaxation oscillators, where a system jumps between two (or more) steady states in an

asymmetric way (which can also be observed in systems close to criticality, see (Tufcea

& François, 2015)). Our model thus offers a simple explanation of wave asymmetry,

solving the long-standing problem of the asymmetry of AP vs PA transitions, which is

possibly crucial for segment polarity as first suggested by Meinhardt (Meinhardt, 1982).

Discussion

In this work, we have explored the dynamical properties of generic two-module systems, where one module corresponds to a dynamic phase of genetic expression and the other corresponds to a static phase controlling embryonic patterning. The unexpected result is that those models typically present global bifurcations where new fixed points appear on the trajectories in phase space. Such SNIC bifurcations come from the smooth interpolation between a flow defining an oscillator in phase space and a landscape characterized by several fixed points. The oscillating attractor then gets continuously deformed until it breaks into several fixed points, leading to the SNIC. This interpolation is a direct consequence of the assumed two-module control as shown on multiple examples above. Importantly, the overall developmental sequence in this context is emergent, since the dynamics close to the bifurcation cannot be understood independently from the static or dynamic modules only. SNIC bifurcations also provide robustness to various perturbations (since fixed points appearing on cycles better preserve information on the oscillatory phase). Below we detail how this model also recapitulates many observed features of metazoan segmentation.

Experimental Evidence

In absence of a direct handle on the control parameter, comparison of SNIC and Hopf bifurcations in simulated systems reveal three possible signatures of SNIC bifurcations in the propagating antero-posterior waves of genetic expression: 1. oscillations gradually reinforcing the pattern for SNIC (instead of damped oscillations for Hopf) 2.

relaxation-like oscillations leading to some asymmetry in the wave pattern, and 3. a

continuous shrinking of the spatial wavelength during the dynamic phase to match the

static pattern for SNIC, with many well-defined propagating waves.

For prediction 1, decades of in situs of oscillating genes and of experimental monitoring

of *Notch* signalling pathways argue against damped oscillations in somitogenesis. Weak

waves gradually refine into well-defined stripes in the anterior PSM, in most species(see

for example the comparison made in (Gomez et al., 2008)), suggesting an increase of

amplitude in oscillations. Consistent with this, visualising segmentation clock oscillations

in live vertebrate embryos suggests an increase of oscillation amplitude as cells get

more anterior in the PSM (Delaune et al., 2012; Lauschke et al., 2013). A similar

observation was also made for gap genes during short-germ segmentation in *Tribolium*

(Zhu et al., 2017).

For prediction 2, in somitogenesis, there is an asymmetry in the wave pattern before

stabilization. The transition from the high to low phase within one somite is shallower

than the transition between those two phases from one somite to the other (i.e. posterior

of one somite to anterior of the next), as detailed in (Shih et al., 2015). A possible read-

out is the downstream pattern of *Cadherin* controlling segment boundary formation,

which presents a sawtooth profile (McMillen, Chatti, Julich, & Holley, 2016), consistent with

the observed asymmetry in the wave pattern. It has been proposed that the main

reason for the clock control was precisely to generate such periodic sawtooth pattern,

and a SNIC bifurcation offers a plausible mechanism. In insects, it has been proposed

that one of the roles of traveling waves of segmentation genes (e.g. *eve* genes shift) is

indeed to provide segment polarity (Rothschild, Tsimiklis, Siggia, & François, 2016) (Clark,

2017; Clark & Akam, 2016), which is consistent with this proposal, although precise

quantification of wave asymmetry has not been done in this context to our knowledge.

In situs of gap genes in short germ insects show asymmetry between the anterior and

posterior borders of the propagating gene expression waves as well (Zhu et al., 2017).

For prediction 3, the slowdown of oscillations during vertebrate segmentation appears

quite variable between species. Mouse segmentation clock seems at first more

consistent with a relatively sudden Hopf scenario. There is roughly a 2π phase shift

between the posterior and the front, consistent with only a moderate change of the

period/wavelength (Lauschke et al., 2013). Other species have several propagating waves

within the PSM, seemingly more consistent with a broader period regime. In zebrafish,

in vivo real-time imaging suggests a wavelength of propagating waves in the anterior

PSM of roughly twice the segment length, meaning that the period of the clock at the

front is at least half of the period in the tail bud (Shih, François, Delaune, & Amacher, 2015).

However, this measurement likely overestimates the wavelength and thus

underestimates the period at the front since it is based on a peak to peak measurement

between the last two waves. Thus the period of the clock might further increase in the

anterior, consistent with a period divergence and with measurements of (Francois

Giudicelli et al., 2007) on in situs, who proposed that the period indeed diverges (Francois

Giudicelli et al., 2007). Importantly, as said above, while the period increases the

amplitude increases too (Shih et al., 2015), which is more consistent with a SNIC

bifurcation. In situs in snake PSM clearly show a decrease of the wavelength that

seems to continuously match the pattern in the anterior, possibly more consistent with

an infinite-period bifurcation as well (Gomez et al., 2008).

A direct experimental measure of the wavelength of the pattern for the last oscillations

can be found with *Mesp2* ([Koseki et al., 2000](#)) ([Saga & Takeda, 2001](#)). The consensus on

Mesp2 is that it is expressed in the last few waves of oscillations, and that its pattern

continuously regresses to reach exactly the wavelength of the anterior pattern. This is

consistent with an infinite-period bifurcation. In (Oginuma et al., 2010) it is argued that

Mesp helps setting the segment boundary in mouse by precisely reading the Notch

wave, explaining both its biological role and why it is a convenient marker of the

wavelength. We notice that the model used to explain the experiments in (Oginuma et

al., 2010) is Julian Lewis' model from the appendix of (Palmeirim et al., 1997), thus

presenting an infinite-period divergence.

One cannot completely exclude from all those experimental observations a Hopf

scenario where waves would not be very damped (or would be amplified for some

reason) past the bifurcation, while the period of damped oscillations would become long

until the saddle-node bifurcation is reached. A way to experimentally falsify this scenario

is to induce a smoother transition and see if damped oscillations appear. Interestingly,

both in vertebrates and in short germ insects, changes of *Wnt* signalling indeed give rise

to smoother transitions from the dynamic to the static regime (suggesting that *Wnt* could

be related to the control parameter g). In mouse, *beta-catenin* gain of function mutants

give rise to considerably extended PSM towards the anterior (Aulehla et al., 2008), with

up to 5 waves of oscillating *Lfng* (compared to only one in WT). Thus, this extended

PSM qualitatively looks much more like a zebrafish or a snake PSM, with reinforcement

of *Lfng* in situs from anterior to posterior (suggesting a refinement of the pattern and not

damped oscillations, see (Aulehla et al., 2008, see Figure 4g,h)). In the anterior PSM of

such mutants, the wavelength of the oscillations decreases to match exactly the *Mesp2*

wavelength (Aulehla et al., 2008, see e.g. Fig 3), and the very last waves move

extremely slowly towards the anterior (Alexander Aulehla, private communication),

consistent with a divergence of the time-scale. In *Tribolium*, *axin* RNAi similarly changes

the wave pattern, where smoother and more spatially extended propagation of gap

gene expressions are observed in those mutants, with reduced wavelengths compared

to the WT (see Fig. 4 of (Zhu et al., 2017)). Those phenotypes are exactly what is

expected from an infinite-period bifurcation, again consistent with the SNIC scenario.

Lastly, damped oscillations would also be absent from Hopf scenarios where the

bifurcation that destroys the limit cycle occurs at a value of the control parameter that is

almost exactly the value at which the system becomes multistable. This happens for

example when the supercritical Hopf bifurcation that kills the limit cycle happens at the

same value of parameter g than the pitchfork bifurcation that generates bistability

(Figure 4—figure supplement 1F). A similar phenomenon occurs in the model presented

on Figure 4—figure supplement 4: the two simultaneous subcritical Hopf bifurcations

that generate the bistability and the saddle-node of limit cycles that destroys the stable

limit cycle happen in a very narrow range of g values. We cannot fully exclude those

cases from the experimental data, but we notice they are intrinsically less generic since

multiple bifurcations have to occur at the same time, which happens only in our models

because of specific parameter choices, symmetries in the equations or specific choices

of non-linearity for the weights of the modules.

 Model Predictions

The most straightforward prediction of the model proposed here is the presence of

several global transcriptional modules between strongly interacting genes, directly

controlling the smooth changes of developmental time-scale (in a similar way to the

"speed-gradient" model in (Zhu et al., 2017)). Many developmental genes are regulated

by multiple "shadow" enhancers (Cannavò et al., 2016). A smooth transition between

different enhancers has even been observed for gap genes in *Drosophila* (El-Sherif &

Levine, 2016). Global regulation of transcriptional modules could be biologically

achieved through "super enhancers" regulating many genes at the same time (Hnisz et

al., 2017). A non-trivial prediction of our model is that the intrinsic time-scale of the

system is a function of the relative balance of transcriptional activities of the modules.

The transcriptional control described here naturally allows for infinite-period bifurcations,

an implicit mechanism in several models of metazoan segmentation. This is to be

contrasted with classical models of negative feedback oscillators such as the Goodwin

model, where the time-scale is entirely controlled by degradation and is independent

from transcription and translation rates (Forger, 2011), and with models of delayed

oscillators, where the time-scale is essentially controlled by transcriptional delays (Lewis,

2003).

Our model is controlled by an external parameter g. The most natural hypothesis would

be that g corresponds to an actual morphogen gradient. As said above, *Wnt* is a natural

candidate, but feedbacks clearly exist with *Caudal* in *Tribolium* (Zhu et al., 2017) and

FGF in vertebrates. However, in the spirit of the initial wavefront proposal by Cooke and

Zeeman, g could also be in some context a temporal variable, e.g. an effective timer.

Recent works on somitogenesis have suggested that the segmentation front could also

be coupled to the slowing down of two oscillators (Lauschke et al., 2013), possibly one

corresponding to *Notch/FGF* and the other to *Wnt* (Sonnen et al, 2018), so that the

oscillation could feedback on itself to define g. One could also imagine that g is related

to more biophysical variables (density, elasticity) ([Hubaud, Regev, Mahadevan, & Pourquie,](

[2017](). It is important to point out that in our framework the nature of the bifurcation does

not depend on the nature of g. While it might be difficult to experimentally disentangle

feedbacks between the bifurcations and the control parameter from actual properties of

the bifurcations themselves, our predictions on the nature of the bifurcations would not

change.

An assumption of this two-module framework is that the same genes interact to control

the system in *both* the dynamic and static regimes, giving rise to a smooth dynamical

transition during development. This is consistent with what is observed for gap gene

dynamics in short germ insects (Zhu et al., 2017). For vertebrate segmentation, we do

not know yet mechanistically how both regimes are controlled, but the *Notch* signalling

pathway is implicated in all steps of somitogenesis and in particular is known to gate

information from the oscillatory to the segmented regime (Oginuma et al., 2010). An

opposite view would be that the transition from dynamic to static regime is *de facto*

sudden (even if it appears smooth for other reasons). Such scenario could be realized

in different ways. For instance, different enhancers could regulate completely different

sets of genes in the dynamic vs static regimes. The "static" genes would then interact

with the "dynamic" genes only briefly during development, ensuring transmission of

positional information between the static and dynamic regimes in a very localized region

of time and space. In somitogenesis, as said above specific genes are indeed

expressed at the so-called "front" (such as *Mesp2*) and could act like gating processes

transferring the information from the clock to an independent patterning system. In this

case, we would be back to a sequential point of view where different regimes of

development live in different regions of phase space, and the local bifurcation scenario

would then be more plausible.

Evolution and developmental plasticity

Evolutionary simulations for the evolution of patterning have favored a scenario based

on Hopf and saddle-node bifurcations (François et al., 2007). Those simulations did not

include multiple enhancers like here, and all transcriptional regulations had essentially

to evolve from scratch, possibly suggesting a "Ur-segmentation clock" based on Hopf

and saddle-node bifurcations. This scenario is not excluded by our model: in particular

there would be no difference between Hopf and SNIC bifurcations under the control of

steep gradients of g, which shortcut the region where only one fixed point is present. So

Hopf/saddle node bifurcations under control of a steep gradient are likely the easiest

solution found by (simulated) evollution. However, adding a more complex, enhancer-

based evolutionary grammar might allow for more combinatorial use of dynamical

attractors, and associated robustness to external perturbations (such as the shape of

the g gradient). A SNIC bifurcation might then plausibly evolve from the modularization

(West-Eberhard, 2003) of a system based on Hopf and saddle-node bifurcations (both

in vivo and *in silico*). First, since the SNIC bifurcation is the generic scenario that we

observe in our framework, it should be easy to discover by evolution once a multi-

enhancer system combining an oscillator and a bistable system has evolved. Second,

SNIC-controlled segmentation is plastic in the sense that changes of dynamics of the

control parameter would change the transient phenotypes (such as the number of

oscillating waves) but would still generate a pattern. Modularizations leading to

developmental plasticity has been suggested to be an important engine for evo-devo

(West-Eberhard, 2003), since it allows for intra-specific variability without impinging on

the most important phenotypes (here, segmentation).

Importantly, such plasticity is indeed observed experimentally both for short germ and

vertebrate segmentation. For instance, in *Tribolium* one can considerably modify *Caudal*

gradient dynamics and still see proper patterning (Zhu et al., 2017). It could thus explain

how and why there is so much quantitative variability in segmentation mechanisms,

such as short vs intermediate germ band segmentation (as suggested in (Zhu et al.,

2017)), In somitogenesis, there is a lot of interspecies quantitative variability in the

numbers of waves and rescaled periods (Gomez et al., 2008)) while the qualitative

dynamics itself appears very conserved see e.g. (Krol et al., 2011) . In other words, a

two-module mechanism makes the dynamics both more robust – a generic bifurcation

scenario gives precise phase encoding – and more evolvable – one can vary many

features of the system (e.g. basins of attractions, dynamics of the control parameter)

and still get proper patterning, again a hallmark of developmental plasticity.

In brief, we have discussed a two-module based model of the smooth transition in

development from a dynamical regime to a static one. This model explains time-scale

divergence, as well as robustness to changes of morphogen dynamics (Zhu et al., 2017)

and to noise. It provides a possible explanation for smooth robust transitions in

metazoan segmentation, with a non-trivial (global) bifurcation. Further experimental and

theoretical studies are required to assess the importance of smooth transitions for

encoding dynamic information into spatial patterns of genetic expressions.

Acknowledgements

We thank members of the François and El-Sherif groups for insightful discussions, as well as referees for their insightful comments.

This work was supported by a CREATE : Complex Dynamics fellowship to L. Jutras-Dubé, a DFG grant EL 870/2-1 to E. El-Sherif, a NSERC Discovery Grant and a Simons Investigator in Mathematical Modelling of Living Systems to P. François.

Competing Interests

The authors declare that no competing interests exist.

References

Ares, S., Morelli, L. G., Jörg, D. J., Oates, A. C., & Jülicher, F. F. (2012). Collective modes of coupled phase oscillators with delayed coupling. *Physical Review Letters*, *108*(20), 204101.

Aulehla, A., Wiegraebe, W., Baubet, V., Wahl, M. B., Deng, C., Taketo, M., . . . Pourquié, O. (2008). A beta-catenin gradient links the clock and wavefront systems in mouse embryo segmentation. *Nat Cell Biol, 10*(2), 186-193.

Beaupeux, M., & François, P. (2016). Positional information from oscillatory phase

shifts : insights from in silico evolution. *Physical Biology*, *13*(3), 1–14.

Cannavò, E., Khoueiry, P., Garfield, D. A., Geeleher, P., Zichner, T., Gustafson, E. H.,

Ciglar, L., Korbel, J. O., & Furlong, E. E. M. (2016). Shadow enhancers are

pervasive features of developmental regulatory networks. *Current Biology*, *26*(1),

38–51.

Clark, E. (2017). Dynamic patterning by the Drosophila pair-rule network reconciles
long-germ and short-germ segmentation. *PLoS Biol, 15*(9), e2002439.
810
811   Clark, E., & Akam, M. (2016). Odd-paired controls frequency doubling in Drosophila
812   segmentation by altering the pair-rule gene regulatory network. *eLife, 5*.
813

Cooke, J., & Zeeman, E. C. (1976). A clock and wavefront model for control of the

number of repeated structures during animal morphogenesis. *Journal of Theoretical*

Biology, *58*(2), 455–476.

Corson, F., Couturier, L., Rouault, H., Mazouni, K., & Schweisguth, F. (2017). Self-

organized Notch dynamics generate stereotyped sensory organ patterns in

Drosophila. *Science*, *356*(6337), eaai7407.

Corson, F., & Siggia, E. D. (2012). Geometry, epistasis, and developmental patterning.

Proceedings of the National Academy of Sciences, *109*(15), 5568–5575.

Corson, F., & Siggia, E. D. (2017). Gene free methodology for cell fate dynamics during

development. *ELife*, *6*, e30743.

Dale, J. K., Maroto, M., Dequeant, M. L., Malapert, P., McGrew, M., & Pourquie, O.
(2003). Periodic notch inhibition by lunatic fringe underlies the chick
segmentation clock. *Nature, 421*(6920), 275-278.
827
Delaune, E. A., François, P., Shih, N. P., & Amacher, S. L. (2012). Single-Cell-

Resolution Imaging of the Impact of Notch Signaling and Mitosis on Segmentation

Clock Dynamics. *Developmental Cell, 23*(5), 995-1005

Dequeant, M.-L., Glynn, E., Gaudenz, K., Wahl, M., Chen, J., Mushegian, A., &
Pourquié, O. (2006). A complex oscillating network of signaling genes underlies
the mouse segmentation clock. *Science, 314*(5805), 1595-1598.
834

835
836 Diaz-Cuadros, M., Wagner, D. E., Budjan, C., Hubaud, A., Tarazona, O. A., Donelly,

837   S., . . . Pourquie, O. (2020). In vitro characterization of the human segmentation

838   clock. *Nature, 580*(7801), 113-118.

839
Dubrulle, J., & Pourquié, O. (2004). fgf8 mRNA decay establishes a gradient that

couples axial elongation to patterning in the vertebrate embryo. *Nature*, 419–422.

El-Sherif, E., Averof, M., & Brown, S. J. (2012). A segmentation clock operating in

blastoderm and germband stages of Tribolium development. *Development*, *139*(23),

4341–4346.

El-Sherif, E., & Levine, M. (2016). Shadow enhancers mediate dynamic shifts of gap

gene expression in the drosophila embryo. *Current Biology*, *26*(9), 1164–1169.

Elowitz, M. B., & Leibler, S. (2000). A synthetic oscillatory network of transcriptional

regulators. *Nature*, *403*(6767), 335–338.

Ermentrout, B. (2008). Ermentrout-Kopell canonical model. *Scholarpedia, 3*(3), 1398.

Forger, D. B. (2011). Signal processing in cellular clocks. *Proceedings of the National*

Academy of Sciences, *108*(11), 4281–4285.

François, P., Hakim, V., & Siggia, E. D. (2007). Deriving structure from evolution:

metazoan segmentation. *Molecular Systems Biology*, *3*, 9.

François, P., & Siggia, E. D. (2012). Phenotypic models of evolution and development:

geometry as destiny. *Current Opinion in Genetics & Development*, *22*(6), 627–633.

Gillespie, D. T. (2001). Approximate accelerated stochastic simulation of chemically

reacting systems. *The Journal of Chemical Physics*, *115*(4), 1716.

Giudicelli, François, & Lewis, J. (2004). The vertebrate segmentation clock. *Current*

Opinion in Genetics & Development, *14*(4), 407–414.

Giudicelli, F., Ozbudak, E. M., Wright, G. J., & Lewis, J. (2007). Setting the tempo in

development: an investigation of the zebrafish somite clock mechanism. *PLoS*

Biology, *5*(6), e150.

Gomez, C., Ozbudak, E. M., Wunderlich, J., Baumann, D., Lewis, J., & Pourquié, O.

(2008). Control of segment number in vertebrate embryos. *Nature*, *454*(7202), 335–

339.

Graf, T., & Enver, T. (2009). Forcing cells to change lineages. *Nature*, *462*(7273), 587–

594.

Hnisz, D., Shrinivas, K., Young, R. A., Chakraborty, A. K., & Sharp, P. A. (2017). A

phase separation model for transcriptional control. *Cell, 169*(1), 13–23.

Huang, S., Guo, Y., May, G., & Enver, T. (2007). Bifurcation dynamics in lineage-

commitment in bipotent progenitor cells. *Developmental Biology*, *305*(2), 695–713.

Hubaud, A., Regev, I., Mahadevan, L., & Pourquie, O. (2017). Excitable Dynamics and

Yap-Dependent Mechanical Cues Drive the Segmentation Clock. *Cell, 171*(3), 668-

682 e611.

Jaeger, J., & Monk, N. (2014). Bioattractors: dynamical systems theory and the

evolution of regulatory processes. *The Journal of Physiology*, *592*(11), 2267–2281.

Jiang, Y. J., Aerne, B. L., Smithers, L., Haddon, C., Ish-Horowicz, D., & Lewis, J. (2000).

Notch signalling and the synchronization of the somite segmentation clock. *Nature*,

408(6811), 475–479.

Jörg, D. J., Morelli, L. G., Soroldoni, D., Oates, A. C., & Jülicher, F. (2015). Continuum
theory of gene expression waves during vertebrate segmentation. *New Journal of
Physics, 17*(9), 093042.

Koseki, H., Saga, Y., Takahashi, Y., Koizumi, K., Takagi, A., Kitajima, S., & Inoue, T.
(2000). Mesp2 initiates somite segmentation through the Notch signalling pathway.
Nature Genetics, *25*(4), 390–396.

Krol, A. J., Roellig, D., Dequéant, M.-L., Tassy, O., Glynn, E., Hattem, G., Mushegian,
A., Oates, A. C., & Pourquié, O. (2011). Evolutionary plasticity of segmentation
clock networks. *Development*, *138*(13), 2783–2792.

Lauschke, V. M., Tsiairis, C. D., François, P., & Aulehla, A. (2013). Scaling of embryonic
patterning based on phase-gradient encoding. *Nature*, *493*(7430), 101–105.

Lewis, J. (2003). Autoinhibition with transcriptional delay: a simple mechanism for the
zebrafish somitogenesis oscillator. *Current Biology*, *13*(16), 1398–1408.

Matsuda, M., Yamanaka, Y., Uemura, M., Osawa, M., Saito, M. K., Nagahashi, A., . . .
Alev, C. (2020). Recapitulating the human segmentation clock with pluripotent
stem cells. *Nature, 580*(7801), 124-129.

McMillen, P., Chatti, V., Julich, D., & Holley, S. A. (2016). A Sawtooth Pattern of
Cadherin 2 Stability Mechanically Regulates Somite Morphogenesis. *Curr Biol,
26*(4), 542-549.

Meinhardt, H. (1982). *Models of biological pattern formation*. New York, NY: Academic
Press.

Meinhardt, H. (1986). Models of segmentation. In *Somites in developing embryos* (pp.
179–189). Boston, MA: Springer.

Morelli, L. G., Ares, S., Herrgen, L., Schröter, C., Jülicher, F. F., & Oates, A. C. (2009).

Delayed coupling theory of vertebrate segmentation. *HFSP Journal*, *3*(1), 55–66.

Morelli, L. G., & Jülicher, F. F. (2007). Precision of genetic oscillators and clocks.

Physical Review Letters, *98*(22), 228101.

Murray, P. J., Maini, P. K., & Baker, R. E. (2013). Modelling Delta-Notch perturbations

during zebrafish somitogenesis. *Developmental Biology*, *373*(2), 407–421.

Oginuma, M., Takahashi, Y., Kitajima, S., Kiso, M., Kanno, J., Kimura, A., & Saga, Y.

(2010). The oscillation of Notch activation, but not its boundary, is required for

somite border formation and rostral-caudal patterning within a somite. *Development*,

137(9), 1515–1522.

Palmeirim, I., Henrique, D., Ish-Horowicz, D., & Pourquié, O. (1997). Avian hairy gene

expression identifies a molecular clock linked to vertebrate segmentation and

somitogenesis. *Cell*, *91*(5), 639–648.

Poston, T., & Stewart, I. (2012). *Catastrophe theory and its applications*. London: Dover

Publications.

Pusuluri, S. T., Lang, A. H., Mehta, P., & Castillo, H. E. (2018). Cellular reprogramming

dynamics follow a simple 1D reaction coordinate. *Physical Biology*, *15*(1), 16001.

Rothschild, J. B., Tsimiklis, P., Siggia, E. D., & François, P. (2016). Predicting Ancestral

Segmentation Phenotypes from Drosophila to Anopheles Using In Silico Evolution.

PLoS Genetics, 12(5), e1006052-1006019.

Rudolf, H., Zellner, C., & El-Sherif, E. (2020). Speeding up anterior-posterior patterning
of insects by differential initialization of the gap gene cascade. *Dev Biol, 460*(1),
20-31.
928

Saga, Y., & Takeda, H. (2001). The making of the somite: molecular events in

vertebrate segmentation. *Nat Rev Genet, 2*(11), 835-845.

Salvi, J. D., Maoileidigh D. O., & Hudspeth, A. J. (2016). Identification of Bifurcations

from Observations of Noisy Biological Oscillators. *Biophys J, 111*(4), 798-812.

Sarrazin, A. F., Peel, A. D., & Averof, M. (2012). A segmentation clock with two-

segment periodicity in insects. *Science, 336*(6079), 338–341.

Schwendinger-Schreck, J., Kang, Y., & Holley, S. A. (2014). Modeling the Zebrafish

Segmentation Clock's Gene Regulatory Network Constrained by Expression Data

Suggests Evolutionary Transitions Between Oscillating and Nonoscillating

Transcription. *Genetics, 197*(2), 725-738.

Shih, N. P., François, P., Delaune, E. A., & Amacher, S. L. (2015). Dynamics of the

slowing segmentation clock reveal alternating two-segment periodicity.

Development, 142(10), 1785–1793.

Sonnen, K. F., Lauschke, V. M., Uraji, J., Falk, H. J., Petersen, Y., Funk, M. C.,

Beaupeux, M., François, P., Merten, C. A., & Aulehla, A. (2018). Modulation of

phase shift between Wnt and Notch signaling oscillations controls mesoderm

segmentation. *Cell, 172*(5), 1079–1090.e12.

Strogatz, S. H. (2015). *Nonlinear dynamics and chaos: with applications to physics,*

biology, chemistry, and engineering. Boca Raton, FL: CRC Press.

Tufcea, D. E., & François, P. (2015). Critical Timing without a Timer for Embryonic

Development. *Biophysj, 109*(8), 1724-1734.

Verd, B., Clark, E., Wotton, K. R., Janssens, H., Jiménez-Guri, E., Crombach, A., &

Jaeger, J. (2018). A damped oscillator imposes temporal order on posterior gap

gene expression in Drosophila. *PLoS Biology*, *16*(2), e2003174.

Waddington, C. H. (1957). *The strategy of the genes*. London: Routledge.

West-Eberhard, M-J (2003). *Developmental Plasticity and Evolution* . Oxford University

Press

Webb, A. B., Lengyel, I. M., Jörg, D. J., Valentin, G., Jülicher, F. F., Morelli, L. G., . . .
Whitfield, T. T. (2016). Persistence, period and precision of autonomous cellular
oscillators from the zebrafish segmentation clock. *eLife, 5*, e08438.
961

962 Wolpert, L., Smith, J., Jessel, T., Lawrence, P., Robertson, E., & Meyerowitz, E. (2006).

963   *Principles of development*. Oxford University Press.

964 Wright, G. J., Giudicelli, F., Soza-Ried, C., Hanisch, A., Ariza-McNaughton, L., & Lewis,
965   J. (2011). DeltaC and DeltaD interact as Notch ligands in the zebrafish
966   segmentation clock. *Development, 138*(14), 2947-2956.
967

968 Zhu, X., Rudolf, H., Healey, L., François, P., Brown, S. J., Klingler, M., & El-Sherif, E.

969   (2017). Speed regulation of genetic cascades allows for evolvability in the body

970   plan specification of insects. *Proceedings of the National Academy of Sciences*,

971   *114*(41), E8646–E8655.

Figures

High resolution versions of all figures can be found at

https://github.com/laurentjutrasdube/Dual-

Regime_Geometry_for_Embryonic_Patterning/tree/master/Figures

Figure 1: Scenarios for segment formation. (*A*) General phenomenology of

segment or somite formation. The white to blue gradient represents the oscillating system (e.g. some Notch signaling pathway gene). The determination front (red vertical line) sweeps the embryo in the posterior direction (red arrow) and translates the periodic expression of a genetic clock into a spatial pattern. (*B-F*) Pattern formation with the infinite-period scenario. (*B*) Period divergence is imposed as control parameter g decreases from 1 to 0. (*C*) Two simulated cells with the same dynamics of g end up with different final values of the phase. (*D-E*) Kymographs showing respectively the dynamics of parameter g used in the simulated embryo and the dynamics of the genetic clock. (*F*) Schematic of the final pattern. (*G-K*) Pattern formation with the Hopf scenario. (*G*) Schematic of the gene regulatory network. (*H*) Depending on the dynamics of g, simulated cells can end up with either a high or a low concentration of protein E. (*I-J*) Kymograph showing respectively the dynamics of parameter g used in the simulated embryo and the dynamics of protein E. (*K*) Schematic of the final pattern. The boundary between two segments ("S_i") is set arbitrarily at the transition from high to low concentrations of protein E.

Figure 2: 3-gene models for pattern formation. (*A*) Schematic of the gene regulatory networks encoded by the dynamic term (dotted line) and the static term (solid line). (*B*) Kymograph showing the dynamics of parameter g used in the simulated embryos for both Models 1 and 2. (*C-H*) Simulation results for Model 1. (*C*) Weights of the dynamic (dotted line) and static (solid line) modules as a function of parameter g. (*D*) Gene concentration and value of parameter g inside a representative simulated cell as a function of time. (*E*) Kymograph showing the dynamics of gene expression in the simulated embryo. Transparent colors are used to represent the concentration of the 3 genes, so that mixes of the 3 genes can be easily perceived. Genes *A*, *B*, and *C* are shown in transparent white, blue and purple, respectively. Simulated cells with intermediate concentrations of all genes appear grey. (*F*) Schematic of the final pattern. (*G*) Bifurcation diagram showing the types of dynamics available to the simulated embryo as a function of parameter g. The maximum and minimum concentrations of gene *A* on the stable limit cycles are shown in black. Stable and unstable fixed points are shown in green and red, respectively. "SN" stands for

1041   saddle-node bifurcation. (*H*) Period (grey line) and

1042   amplitude (red line) of the oscillations along the stable

1043   limit cycle. (*I-N*) Simulation results for Model 2.

Figure 3: Stochastic simulations of the 3-gene models. (*A-D*) Kymographs showing the stochastic dynamics of gene expression in simulated embryos. The specific values of the typical concentration Ω and of the diffusion constant *D* used to generate each kymograph are indicated on the panels. The concentration of the three genes at the last simulated time point is shown schematically in the lower part of each panel. (*E*) Mutual information as a function of typical concentration Ω for Model 1 (red lines) and Model 2 (black lines). Paler colors correspond to lower values of the diffusion constant *D.* The thick horizontal black line indicates the ideal mutual information for 3 mutually exclusive genes. Note that higher values of Ω correspond to lower noise levels.

Figure 4: Gene-free geometric model for

pattern formation (Model 2). (*A*) Schematic of

the flow encoded by the dynamic and static terms.

The grey circle represents oscillations on the unit

circle. Green and red dots represent unstable and

stable fixed points, respectively. (*B*) Weights of the

dynamic (dotted line) and static (solid line) modules

as a function of parameter g. (*C*) Values of geometric

coordinates y and z and of parameter g in a simulated

cell as a function of time. (*D-E*) Kymographs showing

respectively the dynamics of parameter g used in the

simulated embryo and the dynamics of coordinate y.

(*F*) Schematic of the final pattern. (*G*) Bifurcation

diagram showing the types of dynamics available to

the simulated embryo as a function of parameter g.

The maximum and minimum values of coordinate y

on the stable limit cycles are shown in black. Stable

and unstable fixed points are shown in green and red,

respectively. (*H*) Period and amplitude of the

oscillations. (*I*) Flow in phase space for different

values of parameter g. The same color scheme than

panel *A* is used to represent the cycles and the fixed

points. Positions along the limit cycle at time points

separated by a fixed time interval are indicated with

black dots, so that variations of the speed of the

oscillations along the limit cycle can be visualized.

The yellow and orange lines represent the y and z

nullclines, respectively.

Figure 5: Perturbation of the morphogen gradient steepness in asymmetric 3-gene models. (*A*) Schematic of the gene regulatory networks encoded by the dynamic term (dotted line) and the static term (solid line). The thick red lines indicate stronger repression than the black lines (see the parameters in the Appendix). (*B-C*) Bifurcation diagram showing the types of dynamics available in Models 1 and 2. The maximum and minimum concentrations of gene *A* on the stable limit cycles are shown in black. Stable and unstable fixed points are shown in green and red, respectively. The main bifurcations are identified with vertical lines. "SN" stands for saddle-node bifurcation. (*D-F*) Simulation

1106   results for a steep gradient of parameter g. (*D*) Kymograph showing the dynamics of parameter g used in

1107   the simulated embryos for both Models 1 and 2. (*E-F*) Kymograph showing the dynamics of gene

1108   expression in the simulated embryo of Models 1 and 2. The concentration of the three genes at the last

1109   simulated time point is shown schematically in the lower part of the panels. (*H-J*) Simulation results for a

1110   shallow gradient of parameter g.

1111

1112

Figure 6: Perturbation of the morphogen gradient steepness in the gene-free geometric models. (*A*) Flow plots showing the changes of geometry of the static module. (*B-C*) Corresponding kymographs and final patterns for Model 1. (*D*) Associated bifurcations diagrams. "SN" stands for saddle-node bifurcation. (*E-F*) Corresponding kymographs and final patterns for Model 2. (*G*) Associated bifurcation diagrams.

1118

1119
Figure 7: Wave pattern in different models for the infinite-period scenario.

(*A*) Frequency profile for the simulation of the model of coupled oscillators from (Morelli et al., 2009).

(*B-C*) Kymograph showing the dynamics of the phase of the oscillators and the corresponding final

pattern. (*D*) Two schematic examples of possible wave patterns (symmetrical vs asymmetrical). The

symmetric wave is obtained with a sine function. The asymmetric wave is retrieved from simulations of a

Van der Pol oscillator. (*E*) Wave pattern for the model of Panels (*A-C*) for two different time points. (*F*)

Wave pattern for Model 2 of Figure 4 for two different time points.

 Supplementary Figures

 Figure 1—figure supplement 1:

 Bifurcation analysis of the Hopf

 scenario of Fig. 1. (*A*)

Bifurcation diagram showing the types of

dynamics available to the system as a

function of morphogen *g* concentration.

The maximum and minimum

concentrations of gene *E* on the stable

limit cycle are shown in black. Stable

fixed points are show in green. The main

bifurcation events are identified with

vertical lines. "SN" stands for saddle-

node bifurcation. (*B*) Period of the

oscillations along the limit cycle. (*C*)

Same bifurcation diagram as in *A*, but as

a function of time instead. For any given

time, the diagram shows the stable fixed

points and the extrema of the stable limit

cycles for the concentration of

morphogen *g* that is found in a given

simulated cell at that time. The specific

cell used in this diagram is the first cell of

Fig. 1H.

Figure 1—figure supplement 2: Two-enhancer model for *Tribolium* segmentation.

(*A*) Schematic of the gene regulatory networks encoded by the dynamic term (dotted line) and the static

term (solid line). (*B*) Weights of the dynamic (dotted line) and static (solid line) enhancers. (*C*) Kymograph

showing the dynamics of the concentration of morphogen *caudal* (*cad*) used in the simulated embryo. (*D*)

Kymographs showing the dynamics of the concentration of proteins *hunchback* (*hb*) and *Krüppel* (*Kr*). (*E*)

Final pattern of protein expression. The vertical lines identify the positions of the two cells whose

trajectories are shown on the bottom subpanels of panel F. (*F*) Flow in the phase space defined by *hb*

and *Kr* for different concentrations of morphogen *cad.* The green disk represents the stable fixed point

corresponding to the fate with high concentration of *hb*. (*Top subpanels*) Projection of the trajectories of

all cells in the *hb-Kr* phase space. The trajectories of cells that end up with high *hb*, *Kr*, and *X*

concentrations are represented with transparent blue, red and black lines, respectively. (*Bottom subpanels*) Projection of the trajectories of the two cells identified on panel E. For a given cell, the part of the trajectory that is shown is from the initial time point until the time point when *cad* reaches the concentration used to compute the flow. (G) Speed in phase space of all cells as a function of the time since the formation of the fixed point. The top, middle, and bottom subpanels show the speed of the cells that end up with high *hb*, *Kr*, and *X* concentrations at the end of the simulation, respectively. The thick blue and red lines correspond to the speed of cells 1 and 2, respectively.

Figure 2—figure supplement 1: 3-
gene models for pattern formation
with Hill functions for the weights. (*A*)
Schematic of the gene regulatory networks
encoded by the dynamic term (dotted line) and
the static term (solid line). (*B*) Kymograph
showing the dynamics of parameter g used in
the simulated embryos for both Models 1 and 2.
(*C-H*) Simulation results for Model 1. (*C*)
Weights of the dynamic (dotted line) and static
(solid line) enhancers as a function of
parameter g. (*D*) Gene concentration and value
of parameter g inside a representative
simulated cell as a function of time. (*E*)
Kymograph showing the dynamics of gene
expression in the simulated embryo. (*F*)
Schematic of the final pattern. (*G*) Bifurcation
diagram showing the types of dynamics
available to the simulated embryo as a function
of parameter g. "SN" stands for saddle-node
bifurcation. (*H*) Period (grey line) and amplitude
(red line) of the oscillations along the stable limit
cycle. (*I-N*) Simulation results for Model 2.

Figure 2—figure supplement 2: Peak-to-

peak frequency in the 3-gene models.

The red and black dots represent the normalized

peak-to-peak frequency as a function of

the position along the antero-posterior

(AP) axis for Models 1 and 2,

respectively. These data points were

computed numerically by using

equation 2 of (Giudicelli et al., 2007).

The transparent red and black lines are

the theoretical normalized frequencies

of Models 1 and 2, respectively,

obtained via bifurcation analysis. Note

that after the Hopf bifurcation in Model 1, the system

performs damped oscillations. It is therefore possible

to extract a numerical peak-to-peak frequency even

after the stable oscillations die during the Hopf

bifurcation.

Figure 3—figure supplement 1: Stochastic simulations of the 3-gene models with Hill functions for the weights. (*A-D*) Kymographs showing the stochastic dynamics of gene expression in simulated embryos. The concentration of the three genes at the last simulated time point is shown schematically in the lower part of each panel. (*E*) Mutual information as a function of typical concentration Ω for Model 1 (red lines) and Model 2 (black lines). Paler colors correspond to lower values of the diffusion constant *D*. The thick horizontal black line indicates the ideal mutual information for three mutually exclusive genes.

1229

Figure 4—figure supplement 1: Gene-free models with different weights for the

dynamic and static modules (*A-D*) Bifurcation analysis of gene-free models with linear weights

(Model 2), quadratic weights, cubic weights and weights to the power 4. (*E-H*) Bifurcation analysis of

gene-free models with linear weights, quadratic weights, cubic weights (Model 1) and weights to the

power 4.

Figure 4—figure supplement 2: supercritical Hopf scenario in the gene-free model (Model 1). (*A*) Schematic of the flow encoded by the dynamic, static and intermediate terms. (*B*) Weights of the dynamic (dotted black line), static (solid black line) and intermediate (solid red line) enhancers as a function of parameter g. (*C*) Values of geometric coordinates y and z, and of parameter g in a simulated cell as a function of time. (*D*) Kymograph showing the dynamics of parameter g used in the simulated embryo. (*E*) Kymograph showing the dynamics of geometric coordinate y. (*F*) Schematic of the final pattern. (*G*) Bifurcation diagram showing the types of dynamics available to the simulated embryo as a function of parameter g. (*H*) Period (grey line) and amplitude (red line) of the oscillations. (*I*) Flow in phase space for different values of parameter g. The limit cycles are represented by thick grey lines. Positions along the limit cycle at time points separated by a fixed time interval are indicated with black dots, such that the (absence of) variations of the speed along the limit cycles can be visualized. The yellow and orange lines represent the y and z nullclines, respectively. The green and red dots represent stable and unstable fixed points, respectively.

Figure 4—figure supplement 3: subcritical Hopf scenario in the gene-free model (Model 3). We can modify the dynamic module (making it more non linear) to obtain subcritical Hopf scenario (see equations in appendix) (*A*) Weights of the dynamic (dotted black line), static (solid black line) and intermediate (solid red line) enhancers as a function of parameter g. (*B*) Values of geometric coordinates y and z, and of parameter g in a simulated cell as a function of time. (*C*) Kymograph showing the dynamics of parameter g used in the simulated embryo. (*D*) Kymograph showing the dynamics of geometric coordinate y. (*E*) Schematic of the final pattern. (*F*) Bifurcation diagram showing the types of dynamics available to the simulated embryo as a function of parameter g. "SN" stands for saddle-node bifurcation. (*G*) Period (grey line) and amplitude (red line) of the oscillations. (*H*) Flow in phase space for different values of parameter g. The limit cycles are represented by thick grey lines. Positions along the limit cycle at time points separated by a fixed time interval are indicated with black dots, such that the variations of the speed along the limit cycles can be visualized. The yellow and orange lines represent the y and z nullclines, respectively. The green and red dots represent stable and unstable fixed points, respectively. "SNOLC" stands for saddle-node of limit

1287	cycles	bifurcation.

1288

 Figure 4—figure supplement 4: Gene-free model with two subcritical Hopf

 bifurcations (Model 4) We can modify the dynamic module (making it more non linear) to obtain

subcritical Hopf scenario (see equations in appendix) (*A*) Weights of the dynamic and static enhancers as

a function of parameter g. (*B*) Values of geometric coordinates y and z, and of parameter g in a simulated

cell as a function of time. (*C*) Kymograph showing the dynamics of parameter g used in the simulated

embryo. (*D*) Kymograph showing the dynamics of geometric coordinate y. (*E*) Schematic of the final

pattern. (*F*) Bifurcation diagram showing the types of dynamics available to the simulated embryo as a

function of parameter g. "SN" stands for saddle-node bifurcation. (*G*) Period and amplitude of the

oscillations. (*H*) Flow in phase space for different values of parameter g. The limit cycles are represented

by thick grey lines. Positions along the limit cycle at time points separated by a fixed time interval are

indicated with black dots, such that the (absence of) variations of the speed along the limit cycles can be

visualized. The yellow and orange lines represent the y and z nullclines, respectively. The green and red

dots represent stable and unstable fixed points, respectively. (*I-J*) Bifurcation diagram and flow plots for

values of g close to the subcritical Hopf and saddle-node of limit cycles bifurcations. "SNOLC" stands for

saddle-node of limit cycles bifurcation.

Figure 4—figure supplement 5: Stochastic simulations of the gene-free models. (*A-D*) Kymographs showing the stochastic dynamics of variable y in simulated embryos. The specific value of parameter Ω used to generate each kymograph is indicated on the panels. (*E*) Mutual information as a function of typical concentration Ω for Models 1, 2, 3 and 4. The thick horizontal black line indicates the ideal mutual information for a pattern with two symmetric regions. (*F*) Mutual information as a function of typical concentration Ω for the gene-free models of Figure 4—figure supplement 4.

Figure 5—figure supplement 1: Perturbations of the morphogen gradient

steepness in strongly asymmetric 3-gene models. (*A*) Schematic of the gene regulatory

networks encoded by the dynamic (dotted line) and static (solid line) terms. For each interaction, the color

indicates the strength of the repression, with darker shades of red corresponding to weaker repression.

(*B-C*) Bifurcation diagram showing the types of dynamics available in Models 1 and 2. "SN" stands for

saddle-node bifurcation. (*D-F*) Simulation results for a steep gradient of parameter g. (*D*) Kymograph

showing the dynamics of parameter g used in the simulated embryos of both Models 1 and 2. (*E*)

Kymograph showing the dynamics of gene expression in the simulated embryos of Model 1. The

concentration of the three genes at the last simulated time point is shown schematically in the lower part

of the panel. (*F*) Kymograph showing the dynamics of gene expression in the simulated embryos of

Model 2. (*G-I*) Simulation results for a shallow gradient of parameter g.

Figure 5—

 figure supplement 2: Perturbations of the morphogen gradient steepness in

 randomly asymmetric 3-gene models. (*A*) Schematic of the gene regulatory networks encoded

 by the dynamic (dotted line) and static (solid line) terms. For each interaction, the color indicates the

 strength of the repression, with darker shades of red corresponding to weaker repression. (*B-C*)

Bifurcation diagram showing the types of dynamics available in Models 1 and 2. "SN" stands for saddle-node bifurcation. (*D-F*) Simulation results for a steep gradient of parameter g. (*D*) Kymograph showing the dynamics of parameter g used in the simulated embryos of both Models 1 and 2. (*E*) Kymograph showing the dynamics of gene expression in the simulated embryos of Model 1. The concentration of the three genes at the last simulated time point is shown schematically in the lower part of the panel. (*F*) Kymograph showing the dynamics of gene expression in the simulated embryos of Model 2. (*G-I*) Simulation results for a shallow gradient of parameter g. (*J*) Mutual information as a function of the steepness of the g gradient for Models 1 (red line) and 2 (black line). The thick horizontal black line indicates the theoretical upper bound. The left, center and right subpanels show respectively the mutual information for models with a weak asymmetry in the fixed points of the static term (Fig. 5), with a strong asymmetry (Supp. Fig. 7) and with a random asymmetry (this figure).

Figure 6—figure

supplement 1: Model 1 becomes a SNIC when fixed points are outside the limit

cycle. (*A*) Flow plots showing the change of geometry of the static module. (*B-C*) Corresponding

kymographs and final patterns for Models 1 and 2. (*D-E*) Associated bifurcations diagrams and plots of

the period and amplitude of the oscillations as a function of parameter g.

1351

1352 **Figure 7—figure supplement 1: Wave pattern in different versions of the 2D gene-**

1353 **free model.** (*A*) Wave patterns of the gene-free Model 1 with a Hopf bifurcation. The top and bottom

1354 subpanels show the wave pattern for symmetric and asymmetric fixed points in the static term,

1355 respectively. (*B*) Wave patterns of the gene-free Model 2 with a SNIC bifurcation.

1356

Figure 7—figure supplement 2:
Distribution of the values of the
geometric variables (gene-free
models) and of the gene
concentrations (3-gene models).

(*A*) Histograms showing the distributions of the values of the y (dark blue) and z (light blue) variables in gene-free Models 1, 2, 3 and 4 for different values of parameter g. (*B*) Histograms showing the distributions of the concentrations of genes A (grey), B (blue) and C (pink) in 3-gene Models 1 and 2 for different values of parameter g.

A

Gene-free Model 1

Gene-free Model 2

Gene-free Model 3

Gene-free Model 4

B

3-gene Model 1

3-gene Model 2

3-gene Model 1 (Hill functions)

3-gene Model 2 (Hill functions)

Supplementary Movie Legends

Figure 4—video 1: Flow of the gene-free with a supercritical Hopf bifurcation (Model 1). Flow in phase space as parameter g goes from 1 to 0. The stable limit cycles are represented by thick grey lines. The yellow and orange lines represent the y and z nullclines, respectively. The green and red dots represent stable and unstable fixed points, respectively.

Figure 4— video 2: Flow of the gene-free with a SNIC bifurcation (Model 2). Flow in phase space as parameter g goes from 1 to 0. The stable limit cycles are represented by thick grey lines. The yellow and orange lines represent the y and z nullclines, respectively. The green and red dots represent stable and unstable fixed points, respectively.

Figure 4— video 3: Flow of the gene-free with a subcritical Hopf bifurcation (Model 3). Flow in phase space as parameter g goes from 1 to 0. The stable and unstable limit cycles are represented by thick grey and blue lines, respectively. The yellow and orange lines represent the y and z nullclines, respectively. The green and red dots represent stable and unstable fixed points, respectively.

Figure 4— video 4: Flow of the gene-free with two subcritical Hopf bifurcations (Model 4). Flow in phase space as parameter g goes from 1 to 0. The stable limit cycles

1393 are represented by thick grey lines. The yellow and orange lines represent the y and z

1394 nullclines, respectively. The green and red dots represent stable and unstable fixed

1395 points, respectively.

1396

1397 **Figure 7— video 1: Comparison of pattern formation dynamics in different models.**

1398 Dynamics of the spatial wave patterns in four models: a phase model with diverging

1399 period similar to the infinite-period scenario of Figure 1, the symmetric gene-free Model

1400 1, the symmetric gene-free Model 2, and the asymmetric gene-free Model 2.

1401

1402

1403