Bcl6 Preserves the Suppressive Function of Regulatory T Cells during Tumorigenesis

Yiding Li[1], Zhiming Wang[1], Huayu Lin[1], Lisha Wang[1], Xiangyu Chen[1], Qiao Liu[1], Qianfei
Zuo[1], Jianjun Hu[2], Haoqiang Wang[3], Junyi Guo[4], Luoyingzi Xie[1], Jianfang Tang[1], Zhirong Li[1],
Li Hu[1], Lilin Ye[1*], Qizhao Huang[5*], Lifan Xu[1*]

[1]Institute of Immunology, Third Military Medical University, Chongqing, China

[2]Department of Oncology, Southwest Hospital, Third Military Medical University, Chongqing, China

[3]Cancer Center, The General Hospital of Southern Theater Command, Guangzhou, China

[4]Guanghua School of Stomatology, Guangdong Provincial Key Laboratory of Stomatology,
Stomatological Hospital, Sun Yat-Sen University, Guangzhou, China

[5]Cancer Center, The General Hospital of Western Theater Command, Chengdu, China

* Correspondence:
Lifan Xu
xlftofu@sina.com

Qizhao Huang
huangqizhao1988@163.com

Lilin Ye
yelilinlcmv@tmmu.edu.cn

Keywords: Bcl6, Treg, Tumorigenesis, Checkpoint blockade therapy, Anti-tumor immunity.

Abstract

During tumorigenesis, tumor infiltrating regulatory T (Treg) cells restrict the function of effector T
cells in tumor microenvironment and thereby promoting tumor growth. The anti-tumor activity of
effector T cells can be therapeutically unleashed, and is now being exploited for the treatment of
various types of human cancers. However, the immune suppressive function of Treg cells remains a
major hurdle to broader effectiveness of tumor immunotherapy. In this article, we reported that the
deletion of Bcl6 specifically in Treg cells led to stunted tumor growth, which was caused by impaired
Treg cell responses. Notably, Bcl6 is essential in maintaining the lineage stability of Treg cells in
tumor microenvironment. Meanwhile, we found that the absence of follicular regulatory T (Tfr) cells,
which is a result of Bcl6 deletion in Foxp3[+] cells, was dispensable for tumor control. Importantly, the
increased Bcl6 expression in Treg cells is associated with poor prognosis of human colorectal cancer
and lymph node metastasis of skin melanoma. Furthermore, Bcl6 deletion in Treg cells exhibits
synergistic effects with immune checkpoint blockade therapy. Collectively, these results indicate that
Bcl6 actively participates in regulating Treg cell immune responses during tumorigenesis and can be
exploited as a therapeutic target of anti-tumor immunity.

1   Introduction

Regulatory T (Treg) cells, with the master regulator Foxp3 (forkhead box P3), represent a functional
distinct subset of CD4 T cells, which are endowed with the ability of immune suppression and play a
pivotal role in maintaining immune-homeostasis and autoimmune diseases (1–3). Foxp3[+] Treg cells
exert their effector functions through a variety of molecular mechanisms. Firstly, Treg cells
constitutively express the high-affinity heterotrimeric interleukin 2 (IL2) receptor, also known as
CD25, which further bind to and consume IL2 from their surroundings, thus compromising its effects
on non-Foxp3 effector T cells (Teff) (4,5). Treg cells also express high level of cytotoxic T
lymphocyte antigen 4 (CTLA4), which can bind to CD80/CD86 on antigen presenting cells (APCs)
and thereby transmitting suppressive signals to these cells and reducing their capacity to activate Teff
cells (6). Besides, CTLA4 exhibits a higher affinity to CD80/CD86 than that of CD28, thus
competing with this co-stimulatory receptor, which further disrupts the priming and/or activation of
Teff cells (7). Additionally, Treg cells can produce immunosuppressive cytokines, such as TGFβ,
IL10, and IL35 (8), which will downregulate the activity of APCs and Teff cells, and secrete
granzymes and perforin that can directly kill these cells (9).

Akin to CD8 memory T cells consisting of effector memory T cells (TEM), central memory T cells
(TCM) and tissue-resident memory T cells (TRM) (10), thymus derived Treg (tTreg) cells can also
be divided as central (cTreg) and effector (eTreg) Treg cells based on the expression of trafficking
receptors (11). cTreg cells are programmed to recirculate through secondary lymphoid organs (SLOs)
by expressing CD62L as well as CCR7, while eTreg cells capable of entering non-lymphoid tissues
by virtue of expressing chemokine receptors such as CXCR3, CCR4, CCR6, CCR2, and CCR5, etc.
(12) Adoptive transfer studies have shown that cTreg cells could convert into more highly
proliferative eTreg cells in response to tissue self-antigens (13). Notably, eTreg cells have also been
observed in increased numbers within diverse experimental mouse tumors and human cancers which
suggest the involvement of Treg cells in anti-tumor immunity (14,15).

The depletion of Treg cells, which results in tumor rejection and retardation of tumor growth, has
been reported in mouse models (16–18). For several types of human cancers, including melanoma,
non-small-cell lung, gastric and ovarian cancers, Treg cells account for 10–50% of tumor infiltrated
CD4 T cells compared with 2–5% of CD4 T cells in the peripheral blood of healthy individuals
(19,20). Furthermore, a relatively high infiltration abundance of Treg cells versus non-Treg cells in
TME is associated with poor prognosis in cancer patients (21). Using single cell sequencing, it has
been found that the majority of tumor infiltrating Treg cells are uniformly highly activated comparing
with those in periphery tissues (22), with lower level of CD45RA and CCR7 expression, and highly
enriched for a range of co-stimulatory molecules such as CD27, ICOS, OX40, 41BB, GITR, and co-
inhibitory molecules such as CTLA4, PD1, LAG3, and TIGIT. Remarkably, a number of cytokines
and chemokine receptor genes, most notably CCR8, were upregulated in tumor-resident Treg cells in
comparison to normal tissue-resident ones (23). These information supports the notion that tumor
infiltrating Treg cells hold a distinct transcriptional profiling however, the underlying mechanisms
regulating Treg cells within TME still remain obscure (22).

The transcription factor B cell lymphoma 6 (Bcl6) is intensively investigated during the development
of several types of T cell subpopulations and its fine-tuned expression is modulated by a net of
cytokines (e.g., IL6, IL12, IFNγ, type I IFN signaling, TGFβ, and TNFα) in a variety of cell types
and repressed by IL2-STAT5 signaling (24–27). It controls T follicular helper (Tfh) cell development
and is expressed by Foxp3[+] T follicular regulatory (Tfr) cells (28–30). The specific deletion of Bcl6
in Treg cells in mouse models do not result in spontaneous inflammatory disease, however, it did
enhance Th2-mediated airway inflammation following immunization (31). Bcl6-deficient Treg cells
express higher level of GATA3 when compared with WT (wild-type) Treg cells, and it seems that

Bcl6 controls the Th2 inflammatory activity of Treg cells by repressing GATA3 (32). We revealed
that together with TCF1, Bcl6 is critical in transducing mTORC1 signaling in regulating Tfr
differentiation during protein immunization or viral infection (33). However, whether Bcl6 possesses
a role in regulating Treg cells during tumorigenesis has not been explored.

In this study, we found that Bcl6 expression is enhanced in Treg cells under TME, indicating that
Bcl6 potentially serves to regulating the suppressive function of Treg cells on Teff cells in TME.
Accordingly, depleting Bcl6 in Treg cells enhanced anti-tumor capability and retarded tumor
progression. Moreover, this anti-tumor effect can synergize with anti-CTLA4 or anti-PD1 therapy,
thus may serve as a therapeutic target to further improve the efficacy of immune checkpoint blockade
responders.

2 Materials and methods

2.1 Mice

Bcl6-RFP ($Bcl6^{rfp/+}$ or $Bcl6^{rfp/rfp}$) mice were generously provided by Dr. Xindong Liu (Southwest
Hospital, Chongqing, China). $Foxp3^{DTR}$ were generously provided by Dr. Hua Tang (Institute of
Immunology, Shandong First Medical University, Jinan, China). CXCR5-GFP knock-in mice have
been described previously (34). $Bcl6^{fl/fl}$, $Foxp3^{YFP-Cre}$ knock-in, Cxcr5$^{-/-}$, C57BL/6J (CD45.1 and
CD45.2) mice were purchased from Jackson Laboratory. $Bcl6^{fl/fl}$ mice were bred with $Foxp3^{YFP-Cre}$
knock-in mice to generate $Bcl6^{fl/fl}Foxp3^{Cre}$ mice. All these strains are C57BL/6 background. All the
mice used were analyzed at 6-10 weeks of age (indicated in diagram as "Sac"), and both genders
were included without randomization or "binding". Bone marrow (BM) chimeras were used after 8-
10 weeks of reconstitution. LCMV virus (Armstrong strain) was provided by R. Ahmed (Emory
University) and propagated in our laboratory as previously described (35). And 2×10^5 plaque-
forming units of this strain were used to establish acute infection in mice. For all the phenotypic
analysis, at least 3 animals of each genotype with matched age and gender were analyzed.

2.2 Tissue Preparation

Spleens were surgically removed with sterilized surgical equipment and crushed with the blunt of
1mL syringe on Petri dishes containing 3mL of red blood cell lysis buffer. The spleen mixtures were
separately filtered through a 70µM filter into a 15mL conical centrifuge tube, centrifuged at 1800rpm
for 6min at 4℃. After wash, cell pellets were resuspended in 5mL of R2 media (RPMI-1640
(SIGMA Cat. RNBH7001) + 2% fetal bovine serum (FBS; gibco Cat. 10270-106)). Draining lymph
nodes (dLNs) were extracted with sterilized surgical equipment and crushed between the frosted
surfaces of super-frosted microscope slides into wells containing R2. Cell mixtures were then filtered
through a 70µM filter into a 15mL conical centrifuge tube, centrifuged at 1800rpm for 6min at 4℃.
After wash, cell pellets were resuspended in 0.5mL of R2 media. Tumors were removed from mice
with sterile surgical instruments, pictured and weighted then shredded with ophthalmic scissors.
Tumor tissue mixtures were transferred into 15mL conical tubes and filled with collagenase digest
media (R2+Collagenase). B16-F10 Lung tumor tissue were treated with type2 collagenase (Sangon
Biotech Cat. A004174-0001) and MC-38 solid tumor tissues were treated with type1 collagenase
(Sangon Biotech Cat. A004194-0001). Samples were subsequently placed on a 37℃ shaker for 1
hour, then filtered through 100µM filters into 50mL conical tubes and washed with R2 before
centrifugation. B16-F10 tumor cells were further fractionated 2000rpm for 30 min at 4℃ on a two-
step gradient consisting 44% and 67% Percoll solutions (GE Cat. 17-0891-09). The T cell fraction
was recovered from the inter-face between the 2 layers.

2.3   Flow Cytometry and Antibodies

Flow cytometry data were acquired with a FACSCanto Ⅱ (BD Biosciences) and were analyzed with
FlowJo software (Tree Star). The antibodies and reagents used for flow cytometry staining are listed
in Supplementary Table 1. Surface staining was performed in PBS containing 2% BSA or FBS (w/v).
Tfh cell staining has been described (36). Staining of Bcl6, Bcl2, Tbet and Foxp3 were performed
with the Foxp3/Transcription Factor Staining Buffer Set (00-5523; eBioscience). For in vivo
incorporation of the thymidine analog BrdU, mice were given BrdU (1.5mg BrdU (5-
bromodeoxyuridine) in 0.5ml PBS) intraperitoneally 3h before mice were sacrificed. BrdU in T cells
was stained with a BrdU Flow Kit (552598; BD Biosciences) according to the manufacturer's
instructions. For the detection of cytokine production, lymphocytes were stimulated for 5h in the
presence of PMA (50ng/ml), ionomycin (1μg/ml), Golgi Plug, Golgi Stop, anti-CD107a and anti-
CD107b antibodies (BD Bioscience). Intracellular cytokine staining for CD107, granzyme B and
Ki67 were performed with the Cytofix/Cytoperm Fixation/Permeabilization Kit (554714, BD
Biosciences).

2.4   Adoptive Transfer and Generation of Bone Marrow Chimeras

A total of approximately 1×10^6 splenocytes with WT Treg cells 1:1 mixed with KO Treg cells were
adoptively transferred into cyclophosphamide (CTX, Sigma) treated (single dose at 200mg/kg, 12-24
hours before T cell transfer) CD4$^{-/-}$ mice, which were inoculated with B16-F10 cells intravenously or
MC-38 intraperitoneally on the following day. Bone marrow was collected from $Foxp3^{DTR}$ and
$Cxcr5^{-/-}$ mice. Approximately 5×10^6 cells of a 1:1 mixture of $Foxp3^{DTR}$ and $Cxcr5^{-/-}$ bone marrow
cells were transferred intravenously into sub-lethally irradiated (two doses of 4.5 Gy each) CD4$^{-/-}$
recipients. Recipient mice were reconstituted for 8-10 weeks before tumor transplantation.

2.5   Cell culture and In vitro suppression assay

Tumor cell lines, B16-F10, B-Luciferase B16-F10 (BIOCYTOGEN Cat. B-MCL-002) and MC-38
were cultured for cell injection into C57BL/6J mice. B16-F10, B-Luciferase B16-F10 and MC-38
cell medium was composed of DMEM (gibco Cat. C11995500BT), 10% FBS), 1% L-Glutamine
(Solarbio Cat. G0200), and 1% penicillin/streptomycin (gibco Cat. 15070-063). Prior to injection,
cells were adjusted with RPMI, each needle containing $100\mu L/1\times10^6$ (in situ model) or $500\mu L/5\times10^5$
tumor cells. (metastasis model).

CD4$^+$CD25$^+$GITR$^+$CXCR5$^-$ and CD4$^+$CD25$^+$GITR$^+$CXCR5$^+$ T cells from WT or $Bcl6^{fl/fl}Foxp3^{Cre}$
mice were sorted from tumor tissues or dLNs. CD8$^+$CD44$^+$ T cells from WT mice were sorted from
dLNs. 100,000 CD8$^+$ Teffs were incubated with different numbers of CD4$^+$CD25$^+$GITR$^+$CXCR5$^-$ T
cells from tumor bearing WT, $Bcl6^{fl/fl}Foxp3^{Cre}$ or naive WT, $Bcl6^{fl/fl}Foxp3^{Cre}$ mice at 37℃ for 72
hours and were assessed by flow cytometry.

2.6   Quantitative RT-PCR

For comparison of gene expression in $Bcl6^{-/-}$ Treg cells and wild-type Treg cells both in naive state
and tumor bearing state, Treg cells (CD4$^+$CD25$^+$GITR$^+$CXCR5$^-$) were sorted from naive or tumor
implanted mice and were subsequently lysed in TRIzol LS reagent (10296; Life Technologies). Total
RNA was extracted and reverse-transcribed with a RevertAid H Minus First Strand cDNA Synthesis
Kit (K1632; Thermo Scientific). The resulting cDNA was analyzed for the expression of various
target genes with a QuantiTest SYBR Green PCR Kit (204143; Qiagen) and the corresponding
primers on a CFX96 Touch Real-Time System (Bio-Rad) (Supplementary Table 2).

2.7   Analysis of TCGA data

The TCGA normalized RNA-seq data of human colorectal cancer and skin melanoma patients were
downloaded from https://genome-cancer.ucsc.edu/. None of the patients had any record of
immunotherapy treatment. 352 human colorectal patients with primary tumor were included into
analysis. Samples/patients were split based on gene expression levels analyzed by receiver operating
characteristic curve and grouped into Foxp3^{lo} (66.76%) and Foxp3^{hi} (33.24%). Foxp3^{hi} ones were
further divided into Bcl6^{lo} (50.64%) and Bcl6^{hi} (49.36%) subpopulations for analysis. Survival curves
were then generated, and differences were evaluated by Log-rank test. 84 primary skin melanoma
patients and 213 patients with lymph node metastasis were included. Samples/patients were split
based on gene expression levels analyzed by receiver operating characteristic curve.

2.8   Statistical Analysis

Statistical analysis was conducted with Prism 7 software (GraphPad). An unpaired two-tailed t-test
with 95% confidence interval was used to calculate P-values. For bone marrow and splenic chimera
experiments, a paired two-tailed t-test with 95% confidence interval was used to calculate P values.
Tumor growth curves at different time points were plotted by two-way ANOVA with a Turkey post
hoc test comparison among groups.

3   Results

3.1   The deficiency of Bcl6 in Treg cells leads to delayed tumor progression

To investigate the role of Bcl6 in Treg cells immune response during tumorigenesis, we first
evaluated the expression pattern of Bcl6 in Treg cells in TME and tumor dLNs of wild-type (WT)
mice intravenously implanted with B16-F10 melanoma cells (metastasis model). Using real-time
PCR, we observed an upregulation of Bcl6 transcripts in tumor infiltrating Treg cells when compared
with dLNs derived Treg cells (TdLN Treg) and CD44$^-$CD4$^+$ T cells (naive CD4) (Figure 1A). Using
flow cytometric analysis, we confirmed that tumor infiltrating Treg cells expressed higher level of
Bcl6 than Treg cells derived from tumor dLNs and spleens (Figure 1B) in mice challenged with MC-
38 cells subcutaneously (in situ model). The enhanced expression of Bcl6 in tumor infiltrating Treg
cells indicates an important but unsolved role of Bcl6 within TME. To this end, we crossed *Bcl6*$^{fl/fl}$
mice with *Foxp3*$^{YFP-Cre}$ (*Foxp3*^{Cre}) mice to generate mice in which the *Bcl6* alleles are conditionally
deleted in Foxp3$^+$ Treg cells (hereafter referred to as *Bcl6*$^{fl/fl}$*Foxp3*^{Cre} mice). Then we challenged
these mice with MC-38 cells or B16-F10 melanoma cells. We observed that tumor growth was
significantly repressed in knockout (KO) mice when compared to that in WT mice of both models
(Figure 1C; Supplementary Figure 1A). And this phenotype was further corroborated by decreased
tumor weight at the endpoint of each experiment (Figure 1D; Supplementary Figure 1B).

Next, we sought to investigate the role of Bcl6 during tumor metastasis. We challenged
Bcl6$^{fl/fl}$*Foxp3*^{Cre} and WT mice with B16-F10 melanoma cells intravenously to generate lung
metastasis model. In order to continuously monitor tumor metastasis, we introduced the B-Luciferase
B16-F10 cell line, which carries a luciferase-coding sequence in genome. At day 11 (D11) post
tumor inoculation, we noted weaker fluorescent dots in KO mice with in-vivo imaging system
(Figure 1E), indicating less metastatic foci in KO group than in control group. Consistently, we also
observed less metastatic foci in KO mice compared to WT mice at day 18 post tumor inoculation
(Figure 1F). In line with this result, *Bcl6*$^{fl/fl}$*Foxp3*^{Cre} mice also demonstrated better survival rate
when compared to WT counterparts (Figure 1G). Moreover, when proliferation-indicating dye
labeled B16-F10 cells were intravenously transferred into WT and KO mice respectively, we

harvested the same number of violet positive B16-F10 cells from lung tissues of both groups two
hours later, indicating the comparable migration capacity of tumor cells within both groups.
However, 20 hours later, a significant decrease in the number of pulmonary B16-F10 cells suggested
that the lungs of KO mice represent a hostile environment for tumor cell engraftment (Supplementary
Figure 1C).

3.2   The deficiency of Bcl6 impaired the suppressive capacity of Treg cells in TME

To better understand the role of Bcl6 in Treg cells during tumorigenesis, we examined tumor
infiltrated effector T cell response after challenging $Bcl6^{fl/fl}Foxp3^{Cre}$ and WT mice with MC-38 cells
subcutaneously. At D16 post tumor implantation, using flow cytometry analysis, we found that the
deficiency of Bcl6 results in an impairment of Treg cells, evidenced by the decreased proportion and
absolute number of Foxp3$^+$CD4$^+$ T cells (Figure 2A). Correspondingly, we observed increased tumor
infiltrated CD4$^+$CD44$^+$ and CD8$^+$CD44$^+$ T cells upon Bcl6 deletion in Treg cells (Figure 2B). In
intracellular cytokine staining, we found more tumor infiltrating CD8$^+$CD44$^+$ T cells generating
CD107 and Interferon-gama (IFNγ) in KO mice compared to WT mice when stimulated with PMA
plus Ionomycin (Figure 2C). Consistently, in melanoma lung metastasis model, we found a higher
ratio of effector CD4 T cells to Treg cells in tumor compared with WT counterparts (Supplementary
Figure 1D). Although the total number of CD8 T cells remain comparable between these two groups
(Supplementary Figure 1D), CD8 T cells derived from KO mice did show better effector function
than that of WT mice, shown by increased proportion and absolute number of CD107$^+$, TNFα$^+$ and
KLRG1$^+$ CD8$^+$CD44$^+$ T cells in KO mice (Supplementary Figure 1E). We also observed increased
number of total CD4$^+$CD44$^+$ T cells and cytokine producing CD4$^+$CD44$^+$Foxp3$^-$ T (CD107$^+$ and
TNFα$^+$) cells (Supplementary Figure 1F). Thus, these data indicate the critical role of Bcl6 in
regulating Treg cells' suppression on effector T cells during tumorigenesis.

To test whether Bcl6 play an intrinsic role in Treg cells, we generate a splenic chimera mice by
mixing splenocytes derived from WT mice (CD45.1$^+$, 50%) and KO mice ($Bcl6^{fl/fl}Foxp3^{Cre}$, CD45.2$^+$,
50%) and injecting the mixtures into CD4$^{-/-}$ recipients (Figure 2D). We compared the immune
response of Treg cells derived from WT and KO donors in a same host of MC-38 primary tumor
model. At the endpoint of tumor implantation, we observed comparable proportion of Treg cells
originated from both donors (Figure 2E). However, KO mice derived Treg cells exhibited remarkably
reduced expression of Foxp3 and CTLA4 in both tumor tissue (Figure 2F) and dLNs (Supplementary
Figure 2A), suggesting a compromised stability and function of Bcl6-dificient Treg cells. And these
phenomena were further corroborated in lung metastasis model (Supplementary Figure 2B).
Consistently, we conducted in vitro suppression assay by co-culturing WT or Bcl6-deficient tumor
infiltrating Treg cells with CFSE-labelled naive CD8 T cells that were followed by anti-CD3 and
CD28 stimulation, and the results showed that Bcl6-deficient Treg cells exhibited impaired
suppressive function compared with that of WT Treg cells. Contrarily, when isolating cells from
naive mice, WT or Bcl6-deficient Treg cells exhibited comparable suppressive function (Figure 2G).
Collectively, these data indicated that the abrogation of Bcl6 in Treg cells severely dampened the
suppressive function of Treg cells in a cell intrinsic manner in TME, without profoundly influencing
the absolute number of Treg cells.

3.3   Tfr cells are dispensable for tumor control

It has been reported that Bcl6 participates in regulating the differentiation of follicular regulatory T
cells (Tfr) (33). which is a subset of Treg cells endowed with the capacity to inhibit germinal center
reaction. To test whether tumor repression in KO mice was contributed by compromised Tfr
response, we first examined Tfr population within tumor using CXCR5-GFP reporter mice (34).

Notably, GFP$^+$Foxp3$^+$CD4$^+$ T cells were barely detected in tumor, while a substantial population of
Tfr was observed in dLNs (Supplementary Figure 2C), suggesting the absence of Tfr population in
tumor mass. Consistent with published data (33), the deficiency of Bcl6 resulted in significantly
decreased proportion and absolute number of Tfr cells in dLNs, accompanied with upregulated
effector CD4 T cells including Th1 and Tfh cells (Supplementary Figure 2D). To investigate whether
dLNs derived Tfr cells were involved in tumor control, we set up BM chimera mice by mixing bone
marrow cells derived from *Foxp3*^{DTR} mice (50%) and *Cxcr5*$^{-/-}$ mice (50%) and injecting the mixtures
into sub-lethally irradiated CD4$^{-/-}$ recipients (Figure 3A). In this setting, Tfr cells were selectively
depleted after diphtheria (DT) administration (37), while Treg cells remained intact (Figure 3C). At
D16 after tumor implantation (lung metastasis model), we found comparable tumor control ability
between Tfr-deficient mice (DT treatment group) and control mice, indicated by similar lung
metastasis foci between two groups (Figure 3B). Besides, DT induced deletion of Tfr cells did not
influence tumor infiltrating Treg cells as well as effector CD4 T and CD8 T cells, evidenced by
comparable number of Treg cells, Foxp3$^-$CD44$^+$CD4$^+$ T (Figure 3D) and CD44$^+$CD8$^+$ T cells (Figure
3E). Thus, these data demonstrated that Tfr cells were dispensable for tumor control.

3.4   Bcl6 is essential in maintaining the lineage stability of Treg cells in TME

In order to illustrate the underlying mechanisms of Bcl6 in regulating Treg cell response during
tumorigenesis, we sorted antigen experienced Treg, Th1 and Tfh cells as well as naive Treg cells
from WT and KO mice at 16 days after tumor implantation to extract total RNA and conduct an array
of quantitative real-time PCR (RT-qPCR) to profile the transcriptional signatures of the indicated
subsets of CD4 T cells. Consistent with FACS data (shown in Figure 2F), the deficiency of Bcl6
resulted in reduced expression of Foxp3 in Treg cells in tumor model (Figure 4A). However, it had
no effects on Foxp3 expression in naive state (Figure 4B), indicating that the identity of Bcl6-
deficient Treg cells might be preferentially altered within TME. Accordingly, compared with WT
counterparts, KO derived tumor infiltrating Treg cells upregulated the expression level of *Gata3*,
which encodes the master transcriptional factor (TF) of Th2, as well as Th2 related cytokine IL4
(Figure 4C). Meanwhile, the expression of Th17 related cytokine IL17 also increased in KO mice
despite similar level of retinoic acid receptor-related orphan receptor (ROR)γt between WT and KO
group (Figure 4D). During naive state, however, KO Treg cells exhibited normal stability indicated
by comparable level of *Gata3*, *Il17* and *Rorc* expression compared with WT mice (Figure 4E). These
data suggested that the altered identity of Treg cells in KO mice was specifically confined to tumor
micro-environment. Moreover, by analyzing Treg cells derived from different donors in spleen
chimera mice (lung metastasis model), we further confirmed the enhanced expression of Tbet and
GATA3 in KO Treg cells compared with WT ones in both tumor tissue (Figure 4F) and dLNs
(Supplementary Figure 2E). All together, these data suggested the important role of Bcl6 in
maintaining the lineage stability of Treg cells during tumorigenesis.

3.5   Bcl6 expression in Treg cells may predict the prognosis of human colorectal cancer

Given that the deficiency of Bcl6 in Treg cells resulted in repressed tumor growth in mouse models,
we further checked whether Bcl6 expression could be exploited as an indicator for the clinical
prognosis of human cancer patients. Interestingly, analysis of the independent human colorectal
cancer cohort from TCGA (The Cancer Genome Atlas) database demonstrated that patients with
higher level of Bcl6 (in Foxp3$^+$CD4$^+$ cells) showed significantly poorer overall survival compared to
those with a lower level (Figure 5A). Moreover, the expression level of Bcl6 in Treg cells negatively
correlates with the pathological grading of colorectal cancer, indicated by progressively increased
proportion of Bcl6^{hi} Treg cells from pathological stage I to stage IV (TNM stage) colorectal cancer

patients (Figure 5B). In a skin melanoma cohort from TCGA database, we observed increased
proportion of lymph node metastasis in patients with higher Bcl6 expression in Treg cells
(Supplementary Figure 2F), indicating that high level of Bcl6 in Treg cells may correlate with
increased risk of metastasis of melanoma.

Finally, we evaluated the synergetic effects of Bcl6 deletion with immune check point blockade
(ICB). Consistent with published data (38–40), anti-CTLA4 or anti-PD1 alone significantly repressed
tumor growth, which is comparable to that in KO group. Although Bcl6 deletion alone greatly
improved the therapeutic efficacy, the combination of Bcl6 deletion and anti-CTLA4 or anti-PD1 did
exhibit additional effects at delaying tumor growth (Figure 5C-E), indicating that Bcl6 can be a
potential therapeutic target combined with ICB in cancer immunotherapy.

4　　Discussion

Treg cells readily infiltrate into the TME and dampen anti-tumor immune responses, thereby
becoming a barrier to effective cancer immunotherapy. In the past few years, Bcl6 has emerged as a
central regulator of Tfh cell differentiation (35,41,42). However, the role of Bcl6 in Treg cells
function has not been completely elucidated, particularly in terms of anti-tumor immune response. In
this study, to specifically assess the role of Bcl6 in tumor infiltrating Treg cells, we challenged
$Bcl6^{fl/fl}Foxp3^{Cre}$ mice with tumor cells and found that tumor growth was retarded in $Bcl6^{fl/fl}Foxp3^{Cre}$
mice with decreased tumor weight at the end point when compared to WT counterparts. And KO
mice survive longer than WT mice in melanoma lung metastasis model. Besides, the deletion of Bcl6
in Treg cells resulted in impaired suppressive function, evidenced by enhanced functionality of CD8
T cells and non-Foxp3 CD4 T cells, as well as lowed expression of critical function markers of Treg
cells including CTLA4 and CD25. Moreover, Bcl6 is essential in maintaining Treg cell lineage
identity within tumor microenvironment. And tumor infiltrating $Bcl6^{-/-}$ Treg cells exhibited markedly
decreased expression of Foxp3, while upregulating the expression of TFs and cytokines related to
other lineages, including Tbet, RORγt, GATA3, IL4 and IL17, suggesting a weakened lineage
stability. Besides, Bcl6 expression in Treg cells reversely correlates with the prognosis of human
colorectal cancer, while positively correlated with lymph node metastasis of skin melanoma.
moreover, the deletion of Bcl6 in Treg cells acts coordinately with anti-CTLA4 or anti-PD1 to
promote ICB therapeutic efficacy.

It is worth of noting that the impairment of the suppressive function of Treg cells induced by Bcl6
deficiency was only observed in tumor, while undetectable in naive state or non-draining lymph
nodes (data not shown), which may be explained by very low expression levels of Bcl6 in Treg cells
in this condition. Bcl6 conditional KO mice did not show uncontrolled inflammatory or autoimmune
diseases at steady state. Additionally, compared to WT Treg cells, Bcl6-deficient ones in naive state
exhibited normal expression of Foxp3 and comparable capacity in controlling T cell proliferation.
Hence, it is important to investigate the specific role of Bcl6 in Treg cells during tumorigenesis. We
hypothesize that the functional state and stability of tumor infiltrating Treg cells has been altered to a
state in which Bcl6 is essentially required. Recent studies further distinguish peripheral Treg cells
into resting Treg cells (TCF1$^+$LEF1$^+$CD62L$^+$CD44^{lo}), activated Treg cells (TCF1$^+$LEF1$^{+/lo}$CD62L$^-$
CD44$^{int/hi}$) and effector Treg cells (TCF1$^-$LEF1$^-$CD62L$^-$CD44^{hi}) (43). Notably, Bcl6 was selectively
enriched in activated Treg cells (43). In naive state, Treg cells contain a large fraction of resting Treg
subsets, while only a small proportion of Treg cells are active or effector subsets. In tumor
microenvironment, however, the highly suppressive signatures of Treg cells suggest an enrichment of
activated or effector Treg subsets. Thus, Bcl6 may have a more pronounced effect in Treg cells
within tumors.

Treg cells function is highly orchestrated, such that specific transcription factors regulate the ability
of Treg cells to inhibit discrete types of T cell responses. It has been reported that T-bet uniquely
controls the ability of Treg cells to suppress Th1 responses (44), IRF4 regulates the ability of Treg
cells to suppress Th2 responses (45), and Stat3 directs the ability of Treg cells to suppress Th17
responses (46). Accordingly, Bcl6, which is the master regulator of Tfh cells, regulates the
differentiation of Treg cells to Tfr and thus influence the ability of treg cells to control Tfh response
(33). In this study, we found Bcl6 can also regulate Treg cell's function in regulating the antitumor
effects of other T cell lineages, which was shown in a Tfr independent pathway.

Bcl6 deletion significantly down regulate the expression of Foxp3 in tumor infiltrating Treg cells,
which critically specifies the Treg lineage and maintains its functional program (47). But how Bcl6
regulate Foxp3 expression and function of tumor infiltrating Treg cells remains unsolved in our
study. In Th cells, Bcl6 was reported to inhibit Th1, Th2, and Th17 development by suppressing
Tbet, GATA3, and RORγt expression and activities (29,41,48). Meanwhile, it was reported that Bcl6
represses GATA3 expression in Treg cells and thus prevents Treg cells from acquiring Th2 effector-
like characteristics. Consistently, we found an upregulation of Tbet, GATA3, and Th17 related
cytokine (IL17) in KO Treg cells during tumorigenesis. However, whether Bcl6 uniquely targets
these genes in Treg cells in this scenario requires further investigation.

Given the initial clinical successes achieved using PD1/PDL1 or CTLA4 antibody treatments in
patients with cancer, the number of immunotherapy agents in clinical development is expanding
rapidly with goals of improving the limited response rate and generating more durable responses
(49,50). However, the clinical efficacy of ICBs is limited to only a small portion of patients,
suggesting the importance to design new combinatory strategies (51). Currently, Treg cell targeted
anti-cancer therapy have been tested clinically and/or preclinically. Some of the direct strategies
involve targeting of those molecules specifically expressed by Treg cells, such as CTLA4, GITR,
CD25 or ICOS. Several Treg cell-targeted therapies are under investigation alone or in combination
with ICBs (1,52,53). Indeed, the anti-CTLA4 mAb ipilimumab can reinvigorate exhausted CD8 T
cells while reportedly depleting Treg cells from the TME, which might contribute to the clinical
benefits of this agent (54–57). Yet, one possible concern with Treg cell-targeted therapy is that
systemic Treg cell depletion might considerably increase the risk of immune-related adverse events,
such as autoimmunity owing to systemic disruption of immune tolerance (1,52). And several groups
did observe an increased risk of colitis, rash and liver toxicity during the combined therapy of
nivolumab with ipilimumab (an agent that can reportedly deplete or suppress Treg cells) (58,59).
Mogamulizumab has also been associated with the development of rash and liver toxicity (60). For
this reason, targeting Treg cells locally within the tumor could be a better approach that might avoid
adverse events. In our study, we found that the induction of Bcl6 expression was strictly confined to
tumor-infiltrating Treg cells, and the deletion of Bcl6 in Treg cells specifically repressed the function
of this population, leaving systemic Treg cells (outside of TME) that do not express Bcl6 intact.
Moreover, we demonstrated that targeting Bcl6 in Treg cells is sufficient to inhibit tumor growth and
further improve the efficacy of ICB. This result reminds us that Bcl6 might be an optimal therapeutic
candidate against cancer while reducing the side effects. Hence, an extensive and comprehensive
understanding of Treg cell development, maintenance and function, especially in the scenario of
tumorigenesis, could potentially lead to increases in the efficacy of Treg cell-targeted therapies and
reduce the risk of adverse effects of such treatments.

Altogether, we found that Bcl6 is a negative regulator of the suppressive function of Treg cells
during anti-tumor responses. The deletion of Bcl6 in Treg cells greatly improved the accumulation

and function of effector T cells, unleashing a potent and synergistic therapeutic efficacy for
immunotherapy.

5   Ethics Statement

All mouse experiments were performed in accordance with the guidelines of the Institutional Animal
Care and Use Committees of the Third Military Medical University. This study was specifically
reviewed and approved by the Laboratory Animal Welfare and Ethics Committee of the Third
Military Medical University.

6   Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial
relationships that could be construed as a potential conflict of interest.

7   Author Contributions

YL, ZW, HL, LW, QL, QZ, JH, HW, JG, LX, JT, ZL, LH, QH, LX performed the experiments; XC
analyzed TCGA data; LY designed the study, analyzed the data, and wrote the paper with QH, LX
and YL; and LY supervised the study.

8   Funding

This study was supported by grants from the National Natural Science Fund for Distinguished Young
Scholars (No. 31825011 to LY) and the National Natural Science Foundation of China (No.
31700774 to LX and No. 31900643 to QH).

9   Acknowledgments

We thank Dr. Xindong Liu for generously providing us Bcl6-RFP mice. We thank Dr. Hua Tang for
generously providing us *Foxp3*^{DTR} mice, and we thank the core facility center of Third Military
Medical University for cell sorting.

10   References

1.   Togashi Y, Shitara K, Nishikawa H. Regulatory T cells in cancer immunosuppression —
implications for anticancer therapy. *Nat Rev Clin Oncol* (2019) **16**:356–371.
doi:10.1038/s41571-019-0175-7

2.   Sawant D V., Vignali DAA. Once a Treg, always a Treg? *Immunol Rev* (2014) **259**:173–191.
doi:10.1111/imr.12173

3.   Wan YY, Flavell RA. Regulatory T-cell functions are subverted and converted owing to
attenuated Foxp3 expression. *Nature* (2007) **445**:766–770. doi:10.1038/nature05479

4.   Hori S. Control of Regulatory T Cell Development by the Transcription Factor Foxp3. *Science
(80-)* (2003) **299**:1057–1061. doi:10.1126/science.1079490

5.   Sakaguchi S, Wing K, Onishi Y, Prieto-Martin P, Yamaguchi T. Regulatory T cells: How do
they suppress immune responses? *Int Immunol* (2009) **21**:1105–1111.
doi:10.1093/intimm/dxp095

6.   Shevach EM. Mechanisms of Foxp3+ T Regulatory Cell-Mediated Suppression. *Immunity*
(2009) **30**:636–645. doi:10.1016/j.immuni.2009.04.010

7.   Salomon B, Lenschow DJ, Rhee L, Ashourian N, Singh B, Sharpe A, Bluestone JA. B7/CD28
costimulation is essential for the homeostasis of the CD4+CD25+ immunoregulatory T cells
that control autoimmune diabetes. *Immunity* (2000) **12**:431–440. doi:10.1016/S1074-
7613(00)80195-8

8.   Collison LW, Chaturvedi V, Henderson AL, Giacomin PR, Guy C, Bankoti J, Finkelstein D,
Forbes K, Workman CJ, Brown SA, et al. IL-35-mediated induction of a potent regulatory T
cell population. *Nat Immunol* (2010) **11**:1093–1101. doi:10.1038/ni.1952

9.   Tanaka A, Sakaguchi S. Regulatory T cells in cancer immunotherapy. *Cell Res* (2017) **27**:109–
118. doi:10.1038/cr.2016.151

10.   Jameson SC, Masopust D. Understanding Subset Diversity in T Cell Memory. *Immunity*
(2018) **48**:214–226. doi:10.1016/j.immuni.2018.02.010

11.   Smigiel KS, Richards E, Srivastava S, Thomas KR, Dudda JC, Klonowski KD, Campbell DJ.
CCR7 provides localized access to IL-2 and defines homeostatically distinct regulatory T cell
subsets. *J Exp Med* (2014) **211**:121–136. doi:10.1084/jem.20131142

12.   Mempel TR, Marangoni F. Guidance factors orchestrating regulatory T cell positioning in
tissues during development, homeostasis, and response. *Immunol Rev* (2019) **289**:129–141.
doi:10.1111/imr.12761

13.   Wyss L, Stadinski BD, King CG, Schallenberg S, Mccarthy NI, Lee JY, Kretschmer K,
Terracciano LM, Anderson G, Surh CD, et al. Affinity for self antigen selects Treg cells with
distinct functional properties. *Nat Immunol* (2016) **17**:1093–1101. doi:10.1038/ni.3522

14.   Togashi Y, Nishikawa H. "Regulatory T Cells: Molecular and Cellular Basis for
Immunoregulation," in *Assessment & Evaluation in Higher Education*, 3–27.
doi:10.1007/82_2017_58

15.   Sugiyama D, Nishikawa H, Maeda Y, Nishioka M, Tanemura A, Katayama I, Ezoe S,
Kanakura Y, Sato E, Fukumori Y, et al. Anti-CCR4 mAb selectively depletes effector-type
FoxP3+CD4+ regulatory T cells, evoking antitumor immune responses in humans. *Proc Natl
Acad Sci* (2013) **110**:17945–17950. doi:10.1073/pnas.1316796110

16.   Ghiringhelli F, Menard C, Puig PE, Ladoire S, Roux S, Martin F, Solary E, Le Cesne A,
Zitvogel L, Chauffert B. Metronomic cyclophosphamide regimen selectively depletes
CD4+CD25+ regulatory T cells and restores T and NK effector functions in end stage cancer
patients. *Cancer Immunol Immunother* (2007) **56**:641–648. doi:10.1007/s00262-006-0225-8

17.   Wing K, Onishi Y, Prieto-Martin P, Yamaguchi T, Miyara M, Fehervari Z, Nomura T,
Sakaguchi S. CTLA-4 control over Foxp3+ regulatory T cell function. *Science (80-)* (2008)
322:271–275. doi:10.1126/science.1160062

18.   Ge Y, Domschke C, Stoiber N, Schott S, Heil J, Rom J, Blumenstein M, Thum J, Sohn C,
Schneeweiss A, et al. Metronomic cyclophosphamide treatment in metastasized breast cancer

patients: immunological effects and clinical outcome. *Cancer Immunol Immunother* (2012) **61**:353–362. doi:10.1007/s00262-011-1106-3

19. Tada Y, Togashi Y, Kotani D, Kuwata T, Sato E, Kawazoe A, Doi T, Wada H, Nishikawa H, Shitara K. Targeting VEGFR2 with Ramucirumab strongly impacts effector/ activated regulatory T cells and CD8+ T cells in the tumor microenvironment. *J Immunother Cancer* (2018) **6**:1–14. doi:10.1186/s40425-018-0403-1

20. Saito T, Nishikawa H, Wada H, Nagano Y, Sugiyama D, Atarashi K, Maeda Y, Hamaguchi M, Ohkura N, Sato E, et al. Two FOXP3+CD4+ T cell subpopulations distinctly control the prognosis of colorectal cancers. *Nat Med* (2016) **22**:679–684. doi:10.1038/nm.4086

21. Curiel TJ, Coukos G, Zou L, Alvarez X, Cheng P, Mottram P, Evdemon-Hogan M, Conejo-Garcia JR, Zhang L, Burow M, et al. Specific recruitment of regulatory T cells in ovarian carcinoma fosters immune privilege and predicts reduced survival. *Nat Med* (2004) **10**:942–949. doi:10.1038/nm1093

22. De Simone M, Arrigoni A, Rossetti G, Gruarin P, Ranzani V, Politano C, Bonnal RJP, Provasi E, Sarnicola ML, Panzeri I, et al. Transcriptional Landscape of Human Tissue Lymphocytes Unveils Uniqueness of Tumor-Infiltrating T Regulatory Cells. *Immunity* (2016) **45**:1135–1147. doi:10.1016/j.immuni.2016.10.021

23. Plitas G, Konopacki C, Wu K, Bos PD, Morrow M, Putintseva E V., Chudakov DM, Rudensky AY. Regulatory T Cells Exhibit Distinct Features in Human Breast Cancer. *Immunity* (2016) **45**:1122–1134. doi:10.1016/j.immuni.2016.10.032

24. Czerwinska P, Rucinski M, Wlodarczyk N, Jaworska A, Grzadzielewska I, Gryska K, Galus L, Mackiewicz J, Mackiewicz A. Therapeutic melanoma vaccine with cancer stem cell phenotype represses exhaustion and maintains antigen-specific T cell stemness by up-regulating BCL6. *Oncoimmunology* (2020) **9**:1710063. doi:10.1080/2162402X.2019.1710063

25. Huang Q, Xu L, Ye L. *T cell immune response within B-cell follicles*. 1st ed. Elsevier Inc. (2019). doi:10.1016/bs.ai.2019.08.008

26. Sage PT, Sharpe AH. T follicular regulatory cells. *Immunol Rev* (2016) **271**:246–259. doi:10.1111/imr.12411

27. Vinuesa CG, Linterman MA, Yu D, MacLennan ICM. Follicular Helper T Cells. *Annu Rev Immunol* (2016) **34**:335–368. doi:10.1146/annurev-immunol-041015-055605

28. Liu X, Yan X, Zhong B, Nurieva RI, Wang A, Wang X, Martin-Orozco N, Wang Y, Chang SH, Esplugues E, et al. Bcl6 expression specifies the T follicular helper cell program in vivo. *J Exp Med* (2012) **209**:1841–1852. doi:10.1084/jem.20120219

29. Nurieva RI, Chung Y, Martinez GJ, Yang XO, Tanaka S, Matskevitch TD, Wang Y-H, Dong C. Bcl6 Mediates the Development of T Follicular Helper Cells. *Science (80-)* (2009) **325**:1001–1005. doi:10.1126/science.1176676

30. Rawal S, Fan H, Liu Z, Lim H, Tanaka S, Neelapu SS, Chu F, Wang Y-H, Nurieva RI,
Martinez GJ, et al. Follicular regulatory T cells expressing Foxp3 and Bcl-6 suppress germinal
center reactions. *Nat Med* (2011) **17**:983–988. doi:10.1038/nm.2426

31. Sawant D V., Wu H, Yao W, Sehra S, Kaplan MH, Dent AL. The transcriptional repressor
Bcl6 controls the stability of regulatory T cells by intrinsic and extrinsic pathways.
Immunology (2015) **145**:11–23. doi:10.1111/imm.12393

32. Sawant D V., Sehra S, Nguyen ET, Jadhav R, Englert K, Shinnakasu R, Hangoc G,
Broxmeyer HE, Nakayama T, Perumal NB, et al. Bcl6 Controls the Th2 Inflammatory
Activity of Regulatory T Cells by Repressing Gata3 Function. *J Immunol* (2012) **189**:4759–
4769. doi:10.4049/jimmunol.1201794

33. Xu L, Huang Q, Wang H, Hao Y, Bai Q, Hu J, Li Y, Wang P, Chen X, He R, et al. The Kinase
mTORC1 Promotes the Generation and Suppressive Function of Follicular Regulatory T Cells.
Immunity (2017) **47**:538-551.e5. doi:10.1016/j.immuni.2017.08.011

34. He R, Hou S, Liu C, Zhang A, Bai Q, Han M, Yang Y, Wei G, Shen T, Yang X, et al.
Follicular CXCR5-expressing CD8+ T cells curtail chronic viral infection. *Nature* (2016)
537:412–428. doi:10.1038/nature19317

35. Xu L, Cao Y, Xie Z, Huang Q, Bai Q, Yang X, He R, Hao Y, Wang H, Zhao T, et al. The
transcription factor TCF-1 initiates the differentiation of TFH cells during acute viral
infection. *Nat Immunol* (2015) **16**:991–999. doi:10.1038/ni.3229

36. Johnston RJ, Poholek AC, DiToro D, Yusuf I, Eto D, Barnett B, Dent AL, Craft J, Crotty S.
Bcl6 and Blimp-1 Are Reciprocal and Antagonistic Regulators of T Follicular Helper Cell
Differentiation. *Science (80-)* (2009) **325**:1006–1010. doi:10.1126/science.1175870

37. Botta D, Fuller MJ, Marquez-Lago TT, Bachus H, Bradley JE, Weinmann AS, Zajac AJ,
Randall TD, Lund FE, León B, et al. Dynamic regulation of T follicular regulatory cell
responses by interleukin 2 during influenza infection. *Nat Immunol* (2017) **18**:1249–1260.
doi:10.1038/ni.3837

38. Ngiow SF, Young A, Jacquelot N, Yamazaki T, Enot D, Zitvogel L, Smyth MJ. A Threshold
Level of Intratumor CD8 + T-cell PD1 Expression Dictates Therapeutic Response to Anti-
PD1. *Cancer Res* (2015) **75**:3800–3811. doi:10.1158/0008-5472.CAN-15-1082

39. Ngwa W, Irabor OC, Schoenfeld JD, Hesser J, Demaria S, Formenti SC. Using
immunotherapy to boost the abscopal effect. *Nat Rev Cancer* (2018) **18**:313–322.
doi:10.1038/nrc.2018.6

40. Selby MJ, Engelhardt JJ, Quigley M, Henning KA, Chen T, Srinivasan M, Korman AJ. Anti-
CTLA-4 antibodies of IgG2a isotype enhance antitumor activity through reduction of
intratumoral regulatory T cells. *Cancer Immunol Res* (2013) **1**:32–42. doi:10.1158/2326-
6066.CIR-13-0013

41. Yu D, Rao S, Tsai LM, Lee SK, He Y, Sutcliffe EL, Srivastava M, Linterman M, Zheng L,
Simpson N, et al. The Transcriptional Repressor Bcl-6 Directs T Follicular Helper Cell
Lineage Commitment. *Immunity* (2009) **31**:457–468. doi:10.1016/j.immuni.2009.07.002

42. Nurieva RI, Chung Y, Martinez GJ, Yang XO, Tanaka S, Matskevitch TD, Wang Y-H, Dong
C. Bcl6 mediates the development of T follicular helper cells. *Science* (2009) **325**:1001–5.
doi:10.1126/science.1176676

43. Yang B-H, Wang K, Wan S, Liang Y, Yuan X, Dong Y, Cho S, Xu W, Jepsen K, Feng G-S, et
al. TCF1 and LEF1 Control Treg Competitive Survival and Tfr Development to Prevent
Autoimmune Diseases. *Cell Rep* (2019) **27**:3629-3645.e6. doi:10.1016/j.celrep.2019.05.061

44. Koch MA, Tucker-Heard G, Perdue NR, Killebrew JR, Urdahl KB, Campbell DJ. The
transcription factor T-bet controls regulatory T cell homeostasis and function during type 1
inflammation. *Nat Immunol* (2009) **10**:595–602. doi:10.1038/ni.1731

45. Zheng Y, Chaudhry A, Kas A, DeRoos P, Kim JM, Chu T-T, Corcoran L, Treuting P, Klein U,
Rudensky AY. Regulatory T-cell suppressor program co-opts transcription factor IRF4 to
control TH2 responses. *Nature* (2009) **458**:351–356. doi:10.1038/nature07674

46. Chaudhry A, Rudra D, Treuting P, Samstein RM, Liang Y, Kas A, Rudensky AY. CD4+
Regulatory T Cells Control TH17 Responses in a Stat3-Dependent Manner. *Science (80-)*
(2009) **326**:986–991. doi:10.1126/science.1172702

47. Fontenot JD, Rasmussen JP, Williams LM, Dooley JL, Farr AG, Rudensky AY. Regulatory T
Cell Lineage Specification by the Forkhead Transcription Factor Foxp3. *Immunity* (2005)
22:329–341. doi:10.1016/j.immuni.2005.01.016

48. Mondal A, Sawant D, Dent AL. Transcriptional Repressor BCL6 Controls Th17 Responses by
Controlling Gene Expression in Both T Cells and Macrophages. *J Immunol* (2010) **184**:4123–
4132. doi:10.4049/jimmunol.0901242

49. McLane LM, Abdel-Hakeem MS, Wherry EJ. CD8 T Cell Exhaustion During Chronic Viral
Infection and Cancer. *Annu Rev Immunol* (2019) **37**:457–495. doi:10.1146/annurev-immunol-
041015-055318

50. Patel SA, Minn AJ. Combination Cancer Therapy with Immune Checkpoint Blockade:
Mechanisms and Strategies. *Immunity* (2018) **48**:417–433. doi:10.1016/j.immuni.2018.03.007

51. Hashimoto M, Kamphorst AO, Im SJ, Kissick HT, Pillai RN, Ramalingam SS, Araki K,
Ahmed R. CD8 T Cell Exhaustion in Chronic Infection and Cancer : Opportunities for
Interventions. doi:10.1146/annurev-med-012017-043208

52. Raffin C, Vo LT, Bluestone JA. Treg cell-based therapies: challenges and perspectives. *Nat
Rev Immunol* (2019) doi:10.1038/s41577-019-0232-6

53. Zou W. Regulatory T cells, tumour immunity and immunotherapy. *Nat Rev Immunol* (2006)
6:295–307. doi:10.1038/nri1806

54. Vargas FA, Furness AJS, Litchfield K, Joshi K, Rosenthal R, Ghorani E, Solomon I, Lesko
MH, Ruef N, Roddie C, et al. Fc Effector Function Contributes to the Activity of Human Anti-
CTLA-4 Antibodies. *Cancer Cell* (2018) **33**:649-663.e4. doi:10.1016/j.ccell.2018.02.010

55.   Hodi FS, O'Day SJ, McDermott DF, Weber RW, Sosman JA, Haanen JB, Gonzalez R, Robert
C, Schadendorf D, Hassel JC, et al. Improved Survival with Ipilimumab in Patients with
Metastatic Melanoma. *N Engl J Med* (2010) **363**:711–723. doi:10.1056/NEJMoa1003466

56.   Bulliard Y, Jolicoeur R, Windman M, Rue SM, Ettenberg S, Knee DA, Wilson NS, Dranoff G,
Brogdon JL. Activating fc γ receptors contribute to the antitumor activities of
immunoregulatory receptor-targeting antibodies. *J Exp Med* (2013) **210**:1685–1693.
doi:10.1084/jem.20130573

57.   Simpson TR, Li F, Montalvo-Ortiz W, Sepulveda MA, Bergerhoff K, Arce F, Roddie C,
Henry JY, Yagita H, Wolchok JD, et al. Fc-dependent depletion of tumor-infiltrating
regulatory t cells co-defines the efficacy of anti-CTLA-4 therapy against melanoma. *J Exp
Med* (2013) **210**:1695–1710. doi:10.1084/jem.20130579

58.   Hellmann MD, Ciuleanu TE, Pluzanski A, Lee JS, Otterson GA, Audigier-Valette C, Minenza
E, Linardou H, Burgers S, Salman P, et al. Nivolumab plus ipilimumab in lung cancer with a
high tumor mutational burden. *N Engl J Med* (2018) **378**:2093–2104.
doi:10.1056/NEJMoa1801946

59.   Larkin J, Chiarion-Sileni V, Gonzalez R, Grob JJ, Cowey CL, Lao CD, Schadendorf D,
Dummer R, Smylie M, Rutkowski P, et al. Combined nivolumab and ipilimumab or
monotherapy in untreated Melanoma. *N Engl J Med* (2015) **373**:23–34.
doi:10.1056/NEJMoa1504030

60.   Kurose K, Ohue Y, Wada H, Iida S, Ishida T, Kojima T, Doi T, Suzuki S, Isobe M, Funakoshi
T, et al. Phase Ia Study of FoxP3+ CD4 Treg Depletion by Infusion of a Humanized Anti-
CCR4 Antibody, KW-0761, in Cancer Patients. *Clin Cancer Res* (2015) **21**:4327–4336.
doi:10.1158/1078-0432.CCR-15-0357

1   Data Availability Statement

Publicly available datasets were analyzed in this study. These data can be found here:

https://genome-cancer.ucsc.edu/

Figure Legends

Figure 1. The deficiency of Bcl6 in Treg cells leads to delayed tumor progression

(**A**) RT-qPCR of Bcl6 expression in tumor infiltrating Treg (CD4$^+$CD25$^+$GITR$^+$CXCR5$^-$, noted as
TIL Treg) cells, Tfh (gated in CD4$^+$CD44$^+$Foxp3$^-$CXCR5$^+$) and Th1 (gated in CD4$^+$CD44$^+$Foxp3$^-$
CXCR5$^-$) cells sorted from draining lymph node of WT tumor bearing mice. Naive CD4 (gated in
CD4$^+$CD44$^-$) sorted from naive WT mice are included as control. Mice were inoculated with 5×10^5
B16-F10 cells per mice intravenously, and were sacrificed (indicated as "Sac" in figure) at day 16 for
analysis. The data presented are representative of two technical replicates pooled from at least 3 mice
per group. (**B**) Bcl6 expression in tumor infiltrating Treg, dLNs Treg and spleen Treg (CD4$^+$CD25$^+$)

cells. Bcl6-RFP reporter mice were inoculated with 1×10^6 MC-38 subcutaneously, and were
sacrificed at day 10 for flow cytometry analysis. Statistical analysis shown on the right. The data
presented are representative of two independent experiments with at least 3 mice per group. Error
bars indicate the mean$\pm$SEM. *p < 0.05, ***p < 0.001. (**C-D**) Pictures of MC-38 tumor samples
harvested at day 17 after tumor implantation (1×10^6 MC-38 cells per mice subcutaneously, C), with
tumor growth curve and weight (D). (**E**) Tumor progression screened by bioluminescent imaging at
D11. Mice were inoculated with 5×10^5 B16-F10 with luciferase sequence intravenously per mice. (**F**)
Pictures of lung samples harvested from metastasis model, in which mice were inoculated with 1×10^6
B16-F10 cells per mice intravenously and sacrificed at day 16, with foci numbers calculated. (**G**)
Survival curve of $Bcl6^{fl/fl}Foxp3^{Cre}$ and WT mice inoculated with 5×10^5 B16-F10 per mice
intravenously. Data were collected from 8-14 mice per group with two independent experiments.
Statistical differences were calculated by unpaired t test (tumor weight), Log-rank test (survival
curve) or two-way ANOVA with a post hoc Turkey test (tumor growth curve). ***p < 0.001, ****p
< 0.0001. Data are presented as mean$\pm$SEM. See also Supplementary Figure 1

Figure 2. The deficiency of Bcl6 impaired the suppressive capacity of Treg cells

(**A-C**) Flow cytometry analysis, assessed at day 16 after MC-38 inoculation subcutaneously. The
proportion and total number of Treg cells (A) and cytokine producing CD8 subsets (C). Summary of
the total number of $CD4^+CD44^+$ and $CD8^+CD44^+$ T cells (B). The data (A-C) presented are
representative of two independent experiments with 8-14 mice per group. # represents the absolute
number and % represents the proportion of indicating population. Data are presented as mean$\pm$SEM.
Unpaired t test. (**D**) Set up of splenic chimera mice, in situ model. Lymphocytes from WT and
$Bcl6^{fl/fl}Foxp3^{Cre}$ mice were collected, mixed and transfered 1×10^6 cells per mice to $CD4^{-/-}$ recipients
(Treg cells in a ratio of 1:1). On the following day, 1×10^6 MC-38 cells were injected subcutaneously
into recipient mice and mice were sacrificed at day 18 post tumor inoculation. (**E, F**) Flow cytometry
of splenic chimera mice assay, analyzing the ratio of total $CD4^+$T cells in KO versus WT mice, Treg
cell proportion in $CD4^+$ T cells (E), expression level of Foxp3, CTLA4, CD25 and ICOS in Treg
cells (F). The data presented are representative of two independent experiments with at least 4 mice
per group. Paired t test. *p < 0.05, ***p < 0.001, ns, no significant. (**G**) In vitro suppression assay.
$CD8^+$ T cell proliferation were assessed after incubated with tumor infiltrating Treg cells
($CD4^+CD25^+GITR^+$, left) or naive spleen Treg cells ($CD4^+CD25^+GITR^+$, right) sorted from
$Bcl6^{fl/fl}Foxp3^{Cre}$ or WT mice in the ratio of 1:1 to 1:64 for 72 hours. See also Supplementary Figure 1
and S2.

Figure 3. Tfr cells are dispensable for tumor control

(**A**) Set up of Tfr deficient Bone Marrow Chimera mice. Bone marrow cells were isolated from
$Foxp3^{DTR}$ and $Cxcr5^{-/-}$ mice were mixed 1:1 and then transferred 8×10^6 per mice to irradiated $CD4^{-/-}$
mice. After reconstruction, mice were inoculated with 5×10^5 B16-F10 cells intravenously and were
treated with DT or vehicle, then sacrificed at D16 post tumor inoculation. (**B**) Pictures of lung
samples harvested from metastasis model of Tfr deficient mice. (**C-E**) Flow cytometry analyzing
proportion and total number of Treg, Tfr, Tfh, Th1, $CD8^+CXCR5^+$ subpopulation from dLNs (C) and
Treg, $CD4^+CD44^-$ (D), $CD8^+$ T cells (E) from tumor tissue. The data presented are representative of
two independent experiments with at least 4 mice per group. # represents the absolute number and %

represents the proportion of indicating population. Statistical differences were calculated by unpaired
t test. Center values indicate mean. See also Supplementary Figure 2.

Figure 4. Bcl6 is essential in maintaining the lineage stability of Treg cells in TME

(**A**) RT-qPCR analysis of *Foxp3* expression, normalized to its expression in Bcl6-deficient tumor
infiltrating Treg cells (CD4$^+$CD25$^+$GITR$^+$CXCR5$^-$), assessed at day 16 after B16-F10 inoculation
intravenously. (**B**) RT-qPCR analysis of *Foxp3* expression, normalized to its expression in Treg cells
(CD4$^+$CD25$^+$GITR$^+$CXCR5$^-$) from spleen, mesenteric lymph nodes(mLN) or peripheral non-draining
lymph nodes (pLN) of naive *Bcl6*$^{fl/fl}$*Foxp3*^{Cre} and WT mice. (**C, D**) RT-qPCR analysis of *Gata3* and
Il4 (c), *Il17* and *Rorc* (d) normalized to their expression in Bcl6-deficient and WT tumor infiltrating
Treg cells (CD4$^+$CD25$^+$GITR$^+$CXCR5$^-$). Naive CD4 (gated in CD4$^+$CD44$^-$) sorted from naive WT
mice, Tfh (gated in CD4$^+$CD44$^+$Foxp3$^-$CXCR5$^+$) and Th1 (gated in CD4$^+$CD44$^+$Foxp3$^-$CXCR5$^-$)
cells sorted from WT mice infected with Armstrong (Arm) at day 8 are included as control. (**E**) RT-
qPCR analysis of *Il17*, *Rorc* and *Gata3* normalized to their expression in Treg cells
(CD4$^+$CD25$^+$GITR$^+$CXCR5$^-$) from spleen, mesenteric lymph nodes(mLN) or peripheral non-draining
lymph nodes (pLN) of naive *Bcl6*$^{fl/fl}$*Foxp3*^{Cre} and WT mice. The data (A-E) presented are
representative of two technical replicates pooled from at least 3 mice per group. Error bars indicate
the mean$\pm$SEM. (**F**) Flow cytometry analysis of tumor infiltrating Treg cells in splenic chimera
mice assay, assessed at day 18 after B16-F10 inoculation intravenously. Quantification of Tbet and
GATA3 in Bcl6-deficient and WT Treg cells. The data presented are representative of two
independent experiments with at least 4 mice per group. Statistical differences were calculated by
paired t test. **p < 0.01, ****p < 0.0001.

Figure 5. Bcl6 expression in Treg cells may predict the prognosis of human colorectal cancer

(**A**) Survival curves of Bcl6^{hi}Foxp3^{hi} and Bcl6^{lo}Foxp3^{hi} patients in human colorectal cancer TCGA
dataset. (**B**) Proportion of Bcl6^{hi}Foxp3^{hi} and Bcl6^{lo}Foxp3^{hi} patients among Foxp3^{hi} patients in
different stages of human colorectal cancer. (**C**) Set up of anti-CTLA4/anti-PD1 checkpoint blockade
therapy. *Bcl6*$^{fl/fl}$*Foxp3*^{Cre} and WT mice were implanted with MC-38 cells subcutaneously on day 0
and treated with anti-CTLA4 or anti-PD1 antibody at day 7, 9, 12 and 15. (**D, E**) Pictures of tumor
samples harvested at day 18 after tumor implantation, with tumor weight and growth curve. Mice
received either anti-CTLA4 (D) or anti-PD1 (E) therapy. The data were collected from 8-14 mice per
group of two independent experiments. Statistical differences were calculated by unpaired t test
(tumor weight) or two-way ANOVA with a post hoc Turkey test (tumor growth curve). *p < 0.05,
**p < 0.01, ****p < 0.0001, ns, no significance. Data are presented as mean$\pm$SEM. See also
Supplementary Figure 2.

Supplementary Figure 1. Related to Figure 1-2.

(**A-B**) Pictures of B16-F10 tumor samples harvested on day 21 after tumor implantation (1×10^6 B16-
F10 cells per mice subcutaneously, A), with tumor growth curve and weight (B). (**C**) Set up and
analysis of violet labeling assay. *Bcl6*$^{fl/fl}$*Foxp3*^{Cre} and WT mice were inoculated with violet labeled
B16-F10 cells intravenously per mice and were sacrificed 2 or 20 hours post tumor injection.

Summary of violet positive cells 2 or 20 hours post tumor injection are shown on the right. (**D-F**)
Flow cytometry analysis assessed at day 16 after B16-F10 inoculation intravenously. Ratio of
CD4$^+$CD44$^+$ and CD8$^+$CD44$^+$ T cells versus Treg cells, total number of CD4$^+$CD44$^+$ and
CD8$^+$CD44$^+$ T cells (D). Proportion and total number of effecting (KLRG1$^+$) as well as cytokine
producing subsets (CD107$^+$, TNFα$^+$) of CD8$^+$CD44$^+$ T cells (E). Proportion and absolute number of
cytokine-producing subsets of CD4$^+$CD44$^+$Foxp3$^-$ cells (F). The data presented are representative of
two independent experiments with 8-14 mice (D-F) or at least 4 mice (C) per group. # represents the
absolute number and % represents the proportion of indicating population. Center values indicate
mean. Unpaired t test.

Supplementary Figure 2. Related to Figure 2, 3 and 5.

(**A-B, E**) Flow cytometry of splenic chimera mice (B16-F10, metastasis model). Splenocytes from
WT and *Bcl6*$^{\text{fl/fl}}$*Foxp3*$^{\text{Cre}}$ mice were collected and mixed with Treg cells in a ratio of 1:1, and were
adoptively transferred (1×10^6 cells per mice) to CD4$^{-/-}$ recipients. On the following day, 1×10^6 B16-
F10 cells were injected intravenously into recipient mice and mice were sacrificed at day 18 post
tumor inoculation. Quantification of Foxp3, CTLA4, CD25 and ICOS in Treg cells from dLNs (A)
and tumor tissue (B). Quantification of Tbet and GATA3 in Treg cells from dLNs (E). The data
presented are representative of two independent experiments with at least 4 mice per group. Paired t
test. *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001, ns, no significant. (**C**) Flow cytometry
analysis of Tfr (gated in CD4$^+$CD25$^+$GITR$^+$GFP$^+$) cells in dLNs and tumor tissues isolated from
CXCR5-GFP reporter mice, assessed at day 10 after MC-38 inoculation in situ. The data presented
are representative of two independent experiments with at least 3 mice per group. (**D**) Flow
cytometry analyzing total number of lymph node residing Treg cells (gated in
CD4$^+$CD44$^+$Foxp3$^+$CXCR5$^-$), Tfr cells (gated in CD4$^+$CD44$^+$Foxp3$^+$CXCR5$^+$), Tfh cells (gated in
CD4$^+$CD44$^+$Foxp3$^-$CXCR5$^+$) and Th1 cells (gated in CD4$^+$CD44$^+$Foxp3$^-$CXCR5$^-$), assessed at day
16 after B16-F10 inoculation to WT and *Bcl6*$^{\text{fl/fl}}$*Foxp3*$^{\text{Cre}}$ mice intravenously. The data presented are
representative of two independent experiments with 8-14 per group. # represents the absolute
number. Unpaired t test. Center values indicate mean. (**F**) Proportion of Bcl6$^{\text{hi}}$Foxp3$^{\text{hi}}$ and
Bcl6$^{\text{lo}}$Foxp3$^{\text{hi}}$ patients among Foxp3$^{\text{hi}}$ patients in primary and lymph node metastasis patients.