High-throughput microCT scanning of small specimens: preparation, packing,

parameters and post-processing

Christy A. Hipsley[1,2*], Rocio Aguilar[1,2,3], Jay R. Black[4], Scott A. Hocknull[1,5]

[1]School of BioSciences, University of Melbourne, Parkville, VIC, 3010, Australia.

[2]Museums Victoria, GPO Box 666, Melbourne, VIC, 3001, Australia.

[3]School of Biological Sciences, Monash University, Clayton, Victoria, Australia.

[4] School of Earth Sciences, University of Melbourne, Melbourne, Victoria, Australia.

[5]Queensland Museum, Geosciences, 122 Gerler Rd. Hendra, Queensland, 4011, Australia.

*Corresponding author:

Christy A. Hipsley

University of Melbourne

BioSciences 4, Building 147

Parkville, VIC 3010

Australia

christy.hipsley@unimelb.edu.au

Running title: High throughput microCT scanning of small specimens

Abstract

High-resolution X-ray microcomputed tomography, or microCT (μCT), enables the digital

imaging of whole objects in three dimensions. The power of μCT to visualise internal

features without disarticulation makes it particularly valuable for the study of museum

collections, which house millions of physical specimens documenting the spatio-temporal

patterns of life. Despite its potential for comparative analyses, most μCT studies include

limited numbers of museum specimens, due to the challenges of digitising numerous

individuals within a project scope. Here we describe a method for high-throughput μCT

scanning of hundreds of small (< 2 cm) specimens in a single container, followed by

individual labelling and archival storage. We also explore the effects of various packing

materials and multiple specimens per capsule to minimize sample movement that can degrade

image quality, and hence μCT investment. We demonstrate this protocol on vertebrate fossils

from Queensland Museum, Australia, as part of an effort to track community responses to

climate change over evolutionary time. This system can be easily modified for other types of

wet and dry material amenable to X-ray attenuation, including geological, botanical and

zoological samples, providing greater access to large-scale phenotypic data and adding value

to global collections.

Keywords

X-ray microcomputed tomography, microCT, CT scan, 3D imaging, fossils, geometric

morphometrics, high-throughput

Introduction

High-resolution X-ray microcomputed tomography, also known as HRXMT or microCT

(μCT), is an increasingly powerful tool for the non-destructive investigation of whole objects.

Functioning like a microscope with X-ray vision, μCT generates high fidelity 3D models of

solid material from which the outer layers can be virtually dissected or removed, revealing

the inner structure. Starting with radiographs of an object taken over multiple angles, a

computer algorithm is used to digitally reconstruct a stack of 2D X-ray projections, or

tomograms, into a 3D volume. Whereas in human medicine the X-ray source rotates around

the patient (e.g., computerized axial tomography, or CAT scan), in μCT the object is typically

fixed on a rotating stage while the X-ray tube remains stationary. The differential properties

of the object's matter, including thickness and atomic number, interact with the X-ray's

energy beam to determine the number of photons that pass through it to reach the detector on

the other side. This decrease in electromagnetic radiation, termed X-ray attenuation, results in

detector pixels with grayscale values proportional to the radiopacity of the material, meaning

that dense regions such as bone or rock appear white or light gray (radiopaque), while muscle

or skin appears dark (radiolucent). The improved resolution of μCT over standard imaging

techniques can achieve a detail detectability down to 200 nm (0.2 micrometre) – less than the

diameter of a single red blood cell.

Since publication of the first X-ray microtomographic figures nearly four decades

ago[1,2,3], μCT has had profound impacts across scientific disciplines. Studies in biomedicine,

zoology, geology and paleontology now regularly incorporate μCT images[4,5,6], and open

source software for the quantitative analysis of volumetric data is developing rapidly (e.g.,

Fiji[7], Blob3D[8], Dragonfly [Object Research Systems (ORS) Inc., Montreal, Canada]). Within

the biological sciences, μCT has been particularly valuable for the study of museum

collections, which contain millions of often small, delicate and unique specimens not

amenable to traditional (destructive and/or irreversible) preparation. Worldwide, these

collections span taxonomic, geographic and temporal distributions, providing a wealth of

information for understanding the past, present and future of biodiversity[9]. Creation of

cybertypes, or virtual models of type material, is another emerging application of μCT

technology[10,11], allowing researchers around the world to interact with voucher specimens.

When combined with downstream analyses like geometric morphometrics, these models can

be used to test taxonomic hypotheses that could shorten the lagtime between species

discovery and formal description[10,12], thus expanding the taxonomic bottleneck.

Despite the increasing use of μCT in systematic biology[13], a major challenge remains

in the practical imaging of high numbers of museum specimens within a project scope, for

example in the context of large-scale analyses of phenotypic variation. This has proven

difficult because of the need for samples to remain motionless during scan time (minutes to

hours), and because each individual must be digitally labelled to match the physical

specimen's identity and hence retain important information, e.g., material type, locality,

stratigraphic age. These issues have so far limited community-level μCT analyses, which

could provide important insights into evolutionary responses to environmental change in

small, rare, and/or fossilized taxa[6]. While other mass digitisation efforts of museum

collections are already under way, they are typically limited to high resolution photographs of

entire drawers to capture external morphology[14,15]. However internal structures are among the

fastest responders to environmental drivers, suggesting an untapped role for μCT data in

climate change research[6]. Especially in the Anthropocene, μCT offers a promising digital

technique to identify the evolutionary consequences of human activities in natural history

collections at both the internal organismal and ecosystem levels[16,17].

Although some digital morphology studies have included hundreds of[18,19] or over a

thousand[20,21,22,23] 3D models, their images are often aggregated using different sources

(museums, repositories, laboratories), methods (surface scanning, µCT), and parameters (X-

ray energy, voxel size, number of projections), making it difficult to integrate and compare

findings. For smaller specimens in particular, variation in spatial resolution can result in

significant measurement errors, for example in estimated volumes or relationships between

traits[24,25]. Limitations on time and money for µCT scanning also mean that specimen data are

often collected haphazardly, constraining systematic analyses to one or few individuals per

species[6]. This trade-off between data accessibility and data quantity (and quality[26]) precludes

the opportunity for more rigorous investigations of phenotypic diversity in time and space, as

well as focused analyses of variation in a single taxon or locality. Common biodiversity

metrics such as species identity, richness, and evenness are inferable from µCT data, e.g.,

[27,28], especially for cryptic taxa and species complexes with few observable differences[29,23,31].

Therefore, optimisation of the scanning process for high resolution 3D images of multiple

specimens would not only contribute to our understanding of how biodiversity responds to

global change, but also to the value of museum collections and researchers' abilities to access

them.

Here we outline steps for the high-throughput µCT scanning of small (< 2 cm) fossils,

meant to facilitate advanced exploration of museum collections and allow researchers with

limited access to specialized facilities the opportunity to maximize their investments. In

contrast to other high-throughput µCT methods that have focused on clinical evaluations[32,33]

or automated phenotyping of laboratory strains (e.g., mice[34], rice[35], zebrafish[36]), we

concentrate on the practical arrangement of accessioned museum specimens so that each

individual can be identified and labelled in the 3D volume, followed by long-term storage

and archiving. We illustrate this method using vertebrate (reptile, amphibian and mammal)

fossils from the paleontological collections of Queensland Museum, Australia, as part of a

larger effort to identify changes in morphology and species assemblages over geological

time. We succeed in maximizing the quantity of fossils per container to over 200 specimens

to generate high quality 3D models for comparative analyses, while minimising associated

time, handling, and error. These steps should be applicable to any small dry objects

amendable to X-ray attenuation including geological material, plant tissues, and

invertebrates, and wet specimens provided they can be mounted inside small capsules or

other containers[37,38].

Methods

Preparation

The first step is the only time the user is required to handle specimen material directly, we

therefore recommend the use of featherweight spring or feather light forceps and secondary

containment (see Figure 1 for an overview of packing material). Although specimens of

varying size can be scanned together, it is preferable to organise fossils into batches of

roughly similar material properties for X-ray optimization. Thin, unbuffered, neutral pH

tissue is ideal for specimen packing and eventual long-term storage. Other required materials

are small clear two-piece pharmaceutical capsules, paper and/or plastic drinking straws,

archival paper and pen for labelling, a 50 ml plastic centrifuge tube with cap, and medium

density polyethylene foam. Suggested suppliers and estimated costs of material per tube are

listed in Table 1.

Packing procedure

Briefly, each specimen is wrapped in archival tissue and placed inside a capsule with its

label, several of which are inserted lengthwise into a straw. Up to 14 standard paper straws

cut to 9 cm in length fit inside a closed 50 ml Falcon tube, with each straw holding 6 size 4

capsules. For easier tracking of specimens in the 3D volume, we recommend leaving one

capsule empty in different positions along the length of some straws (see post-processing

section below). Following this method, over 80 specimens < 1 cm long can be scanned in a

single tube. This quantity can be doubled or tripled when 2 or 3 fossils (separated by tissue)

are packed inside the same capsule, allowing a maximum of 252 specimens. However, scans

with more than 3 specimens per capsule showed a high (≥ 50%) rate of movement during

rotation, compared to those with fewer (Table 2). We therefore limit packing to 1-2

specimens per capsule, provided they can be distinguished from one another in relative size

and/or shape. Specimens up to 2 cm long can be similarly packed inside size 00 capsules.

Larger diameter paper or smoothie straws sliced lengthwise hold 3 size 00 capsules in a 50 ml

tube (Fig. 1b), which can be scanned together with smaller straws as needed.

Labels for each specimen are written in Indian or archival ink on uncoated acid-free

paper, providing a unique and durable identifier that will be matched in the 3D volume. Our

labels followed a two-part system containing specimen ID (e.g., museum accession number)

and a code for straw and capsule position. Using permanent marker, each straw is labelled

with a letter (A, B, C…N for 14 straws) and each capsule is labelled with its straw's letter

and a number from 1-6, denoting its position in the straw from top to bottom (Fig. 2a). This

system can be amended to suit user needs, for example by including abbreviations for taxon if

known and/or material (e.g., 'il' for ilium, 'max' for maxilla). These codes can also be

extracted from the digital labels and used as variables in downstream analyses, e.g., in the R

statistical package geomorph[39] or MorphoJ[40]. Record labels with capsule and straw positions,

including any empty capsules as a marker, since these will be checked off later during

unpacking.

Once labelling is complete, the specimens, labels, capsules and straws are aligned for

packing. Each specimen is wrapped in a small envelope of Green's Lens tissue (Fig. 2b), and

empty spaces in the capsule are filled with tissue so the specimen does not contact its walls.

This creates more distance between adjacent samples, allowing easier separation (i.e.

segmentation) in the 3D volume. The tissue also prevents movement during image

acquisition (Table 2), which can lead to motion artefacts in the reconstruction like shadows or

streaking. Finally, the label is inserted into the capsule with the ID facing out for long-term

storage, and the capsules are loaded into their respective straws in the correct order. A thinner

cocktail straw can be used to gently push the capsules down without compressing ones below

it (Fig. 2c).

To load the packed straws into the Falcon tube, first fill the conical tip with

polyethylene or other firm material to create a flat surface. Insert the straws with position 1

towards this end, which will be facing up in the μCT machine. The flat surface of the cap on

the other end can be fixed to a glass rod or dowel using a hot glue gun to elevate the tube

from the machine's stage (Fig. 2d). This avoids including the metal clamp in the scan which

could affect X-ray optimization, and centres the tube vertically and horizontally relative to

the detector. Draw a diagram of the straw arrangement with the list of specimen labels, noting

where paper or empty straws are used to separate them for easier identification (Fig. 2e,f).

This material will be visible in reconstructed cross-sections, allowing one to match the 3D

volume to the diagram. Finally, medium density foam can be cut and placed as filler around

the straws and under the cap of the tube before closing. This foam is recommended over

tissue or other supporting material as it acts as an effective shock absorber during rotation.

Secure the tube onto the stage, making sure it can rotate freely without touching the X-ray

source or detector.

X-ray parameters

Scan settings vary depending on the equipment, material size and properties, and desired

results. Here we report parameters as optimised on a Phoenix nanotom m (GE Sensing &

Inspection Technologies GmbH, Wunstorf, Germany) equipped with a 180 kV, 20 W high-

power nanofocus X-ray tube and DXR detector (3072 x 2400 pixels). As with all

microtomographic imaging, initial calibration of the detector is largely software-driven to

reduce potential noise due to defective pixels or incident light. We calibrated the detector by

taking an offset image (dark field) and two sets of gain images (flat field) at the same energy

of the scan and at half the current, where images were averaged over 100 snapshots of the

detector with a skip of 10. For all scans our instrument was fitted with a tungsten target,

although molybdenum may be preferable for amber[4]. Fossils can be radiodense, in such cases

reconstruction artefacts may occur when lower energy X-rays are attenuated at the surface, or

when stronger X-rays hit the detector without passing through the object first (beam

hardening)[4]. A thin metal filter placed in front of the X-ray source can overcome these issues

by removing lower energy photons, although other parameters like current, voltage, or

exposure time should be modified to achieve appropriate contrast.

Ideally the number of images for reconstruction is equal to the maximum width of the

scanned object in pixels $\times \pi/2$[38,41]. For a sample optimized to fill the full width of our

detector, this would result in nearly 5,000 projections at a resolution of 10.6 μm. Such a data

set would be massive in terms of scan time and storage (up to 70 GB before reconstruction),

in addition to the computing power and time needed to reconstruct and render each of the 3D

files individually. To improve the μCT process for smaller specimens without sacrificing

image quality, we divided the Falcon tube into three segments by moving the position of the

stage up or down between scans, while keeping all other settings identical. This produced

separate volumes of equally high quality while avoiding the need to move the detector back

and/or the tube forward to accommodate its length, thereby decreasing resolution. These

volumes can be merged using Avizo (Thermo Fisher Scientific) or other dedicated software

to recreate the entire tube in high resolution. However, for faster processing we aligned each

segment to include two rows of size 4 capsules (or one row of size 00), so they could be

separately visualised and labelled in the 3D volume if preferred.

For all reconstructions we followed the standard protocol of the Phoenix datos|x

software (GE Sensing & Inspection Technologies GmbH, Wunstorf, Germany), which tests

for object movement by comparing the first and last 2D projections. At high resolution, even

minute movements can cause visible artefacts in the reconstructed images[4,38]. Instead of

correcting for movement of single specimens during reconstruction by shifting the axis of

rotation (which would in turn generate artefacts in the rest of the tube), we identified and

labelled those specimens in post-processing so they could be rescanned. Other settings

applied during reconstruction were a 3×3 median filter and an ROI filter.

Computed tomography post-processing

Following reconstruction, digital labelling of the specimens is performed in 3D volume

rendering software. Here we describe steps using VGStudioMax 3.1 (Volume Graphics,

Heidelberg, Germany), although other options for free or commercial licenses are available

(see Table 7 in Keklikoglou et al.[38]). To match the tube orientation to the diagram (Fig. 2f),

the volume can be rotated in the 3D window or by using the registration tool in 2D. Straws

can then be labelled digitally with the indicator tool or by creating a region of interest (ROI)

to name each one by its letter (A, B, C… N). Similar-sized capsules should be aligned across

the straws horizontally, making it easy to identify capsule positions in the 2D and 3D

windows. For example, starting from the top (tip) of the tube in straw A, specimens should be

ordered as A1, A2…A6, with capsules in the neighbouring straw B following the same order.

Verifying the positions of empty capsules is another safeguard to ensure correct orientation.

Using ROIs, the specimens can be segmented and renamed in the rendering software to

match the specimen labels (e.g., museum accession number plus straw/capsule position).

False-colouring straws or capsule rows differently in the 3D volume also helps to track

specimen order. Any specimens with obvious artefacts can be marked with their labels

followed by the word 'REDO', to be set aside during unpacking so they can be scanned again

under improved conditions, i.e, repositioned in the capsule or with better tissue support.

Unpacking and storage

Capsules are unpacked from the straws in the same manner as they were put in, by inserting a

cocktail straw into one end and pushing it through (Fig. 2c). We recommend checking

specimens off the list as they are being unpacked as a final test of correct labelling. These

samples can be stored long-term inside the capsules since archival tissue and paper was used.

Specimens for rescanning can be included in the next batch and labelled digitally with their

ID, new straw and capsule position, and the word 'REDONE' for transparency. This process

can be repeated as many times as necessary to achieve optimal quality scans for all material.

Likewise, all labelled straws and Falcon tubes can be reused in subsequent scans, which will

further ensure consistency in the packing and labelling process.

Results and Discussion

Optimal settings for our fossils in terms of time, resolution and file size were three separate

scans of the Falcon tube at 40 mm focus-to-object distance (FOD) and 225 mm focus-to-

detector distance (FDD), using 40 kV, 300 μA, and 0.5 s exposure for 1400 images, with a

frame average of 3 and image skip of 1. This configuration allowed us to decrease the width

of the detector to 2,000 pixels, meaning that each X-ray image was 9.15 MB. The Y-axis of

the stage was shifted down by 29 mm between scans to allow some overlap between

segments (Fig. 3a). For each segment this resulted in a scan time of 47 minutes and 17.8 μm

voxel size, with reconstructed volumes around 10 GB each (Fig. 3b).

269   This process is in stark contrast to the effort needed to scan each specimen

270 individually at the highest resolution (10.6 μm over 5,000 projections for a single size 4

271 capsule), requiring 167 minutes per scan or over eight hours for a fully packed tube,

272 excluding time to swap samples. Automated sample changers are available for some μCT

273 machines (e.g., Bruker SkyScan, Bruker Biosciences Pty Ltd), although these are still limited

274 to under 20 specimens. Reconstructing and rendering each μCT file separately would also

275 require substantial computing time, making the task unfeasible for individual researchers

276 wishing to work on large data sets. Instead, we stitched the three segments together into a

277 volume of equally high resolution (Fig. 3c), allowing access to multiple specimens by

278 opening a single 30 GB file.

279   The image quality achieved by our method was more than adequate for applications in

280 biological systematics, including geometric morphometrics, functional analyses, and

281 morphological descriptions. Cross-sectional μCT images revealed sharp contrast between the

282 hydroxyapatite of the bone, dentine, and enamel against the clay and crystalline calcium

283 carbonate-rich matrix, while offering precise details of specimen morphology as well as the

284 dividing walls of the capsules, straws and paper (Fig. 3d-f). In few cases artefacts were

285 observed in the reconstruction due to movement (Fig. 3g) or beam hardening (Fig. 3h). These

286 specimens were labelled as 'REDO' in the 3D volume and set aside during unpacking to be

287 rescanned, either with additional tissue in the capsule or using a filter, respectively. In earlier

288 scans we observed some capsules in straws that had been slightly crushed by the one above it.

289 These straws were packed using a solid dowel to push down the capsules, hence why we

290 recommend using a cocktail straw instead.

291   Other examples of material scanned using our high-throughput method are shown in

292 Figure 4. Fossilized frog ilia and varanid osteoderms (bony deposits in the scales of some

293 lizards) yielded excellent results following this protocol (Fig. 4ab), with hundreds of 3D

models generated for minimal time, money and effort. We also applied our system to larger

mammalian specimens, for example fossilized rodent jaws from the Middle Pleistocene Mt

Etna[42]. This material is larger and denser than the herpetological samples, requiring slightly

different parameters for µCT scanning. For larger/denser fossils, a 0.1 mm copper plate was

secured under the collimator to reduce beam hardening and other artefacts, and X-ray voltage

was increased to 50-80 kV. The resulting reconstructed images could be easily segmented

using density-based approaches in VGStudioMax, like the region growing tool to separate

teeth from bone (Fig. 4c).

Stewardship of µCT data is still a major challenge for natural history museums, which

are now tasked with the curation of rapidly expanding digital datasets associated with the

physical collections[43]. There is currently no standard for 3D data storage, meaning that the

fate of µCT images and their metadata is often individual or institution-driven. We retain at

least two copies of the original X-rays and reconstructions in different locations (one cloud-

based, one physical), while working on the prepared volume on a regularly backed-up server.

Davies et al.[5] provide a summary of file types that should be retained for 3D digital

morphology analyses, with best practices recommended when storage space is not a problem.

Online data repositories are another attractive option for long-term data storage, several of

which are free and cater to 3D digital datasets generated for biological research[5], e.g., Zenodo

(zenodo.org), MorphoSource (www.morphosource.org).

For most downstream analyses, µCT data will be converted into polygonal surface

models such as STL (sterolithography) files, which capture the geometric shape of a 3D

object using triangles or vertices to describe its surface (Fig. 4d). These models, also known

as surface meshes, are virtually interactive and can be manipulated, viewed, and measured

using the open source software MeshLab (http://meshlab.sourceforge.net/). They are also

small in terms of data storage, making them easy to distribute online and providing wider

access to museum collections. Finally, surface meshes can be scaled up for displays and

educational programs, for example using 3D printing. Following digital labelling and

unpacking of our specimens, we extracted STL models of each fossil from its ROI in

VGStudioMax, while retaining its original identity.

Many of the fossils included in this project were excavated from cave deposits at

Capricorn Caves in Rockhampton, Queensland, Australia. As part of an annual science

communication event called Capricorn Caves Fossil Open Day[44], we 3D printed select

specimens to engage the public in local biodiversity. There palaeontologists referred to the

enlarged 3D models to assist in the interpretation of microfossils, allowing children to

interact with replicas of local animals from thousands of years in the past (Fig. 4e). In

addition to public outreach, our future work will focus on analysing the generated µCT data

using a growing toolkit of bioinformatic approaches, including machine learning and AI, 3D

landmark-based geometric morphometrics, and high-density point cloud comparisons of

fossil and living forms (Fig. 4f).

Conclusions

Current research on museum-based µCT data encompasses an extraordinary array of material

and topics, including tetrapod origins[45], echolocation in bats[46] and whales[47], limb reduction in

lizards[48,49], mammalian limb development[50,51], primate neuroanatomy[52,53], reproduction in

insects[54] and plants[55], paleoecology of Precambrian biota[56], seafloor biomass estimation[57] and

amber inclusions[58], to name a few. Typically these studies involve few specimens, due to the

challenges of capturing high resolution 3D data for large sample sizes.

Here we demonstrate how simple, inexpensive equipment like drinking straws and

pharmaceutical capsules can be modified for high-throughput µCT scanning of dozens of

fossils in a single container, providing large-scale morphological datasets for comparative

analyses. To date we have scanned over 1,000 accessioned museum specimens following this

technique, including hundreds of frogs, lizards, snakes and mammals. These data will

contribute to ongoing efforts to identify evolutionary responses to climate change in

Australia's fossil record, where reptiles and amphibians constitute half of the terrestrial

vertebrate diversity, and where increasing aridity has transformed large parts of the continent.

This method can also be adapted for other (non-biological) objects, by creating custom-made

holders and stages, and seeking μCT scanners with differing capacities. Some users may wish

to scan individual specimens or holotypes at the highest possible resolution, in which case

our protocol offers an efficient pathway for determining which specimens should be elevated

to type status. At the same time we note that fossils can be difficult to scan, being highly

variable in shape, composition, and density. Thus while μCT is a powerful tool for most

small specimens, in some cases other non-destructive techniques such as magnetic resonance

imaging (MRI), synchrotron, and confocal microscopy may be more appropriate.

Acknowledgements

This work was supported by an Australian Research Council DECRA (DE180100629) to

CAH, with scans performed through the Melbourne TrACEES platform. Danielle Measday,

Conservator of Natural Sciences at Museums Victoria, advised on and provided archival

papers for testing. We thank Kristen Spring, Rochelle Lawrence, Joanne Wilkinson, and

Jonathan Cramb, along with many volunteers at the Queensland Museum, for their assistance

in generating the large quantity of specimens made available for this project.

Author Contributions

C.A.H. and S.H. conceived the project; S.H. selected specimens for μCT scanning; C.A.H.,

R.A., and J.R.B. designed the methodology; J.R.B. operated the μCT machine; C.A.H. and

R.A. collected and analysed the data and led manuscript writing. All authors contributed to

manuscript drafts and gave final approval for publication.

Competing interests

The authors declare no competing interests.

Data Availability

All X-ray and µCT data shown in Fig. 3a-e are available in the MorphoSource repository,

[link provided upon acceptance].

References

1. Sato, T., Ikeda, O., Yamakoshi, Y. & Tsubouchi, M. X-ray tomography for microstructural

objects. *Appl. Opt.* **20**, 3880-3883 (1981).

2. Elliott, J. C. & Dover, S. D. X-ray microtomography. *J. Microsc.* **126**, 211-213 (1982).

3. Elliott, J. C. & Dover, S. D. X-ray microscopy using computerized axial tomography. *J.*

Microsc. **138**, 329-331 (1985).

4. Sutton, M., Rahman, I. & Garwood, R. Techniques for Virtual Palaeontology. 208 p

(Wiley-Blackwell, 2014).

5. Davies, T. G. *et al.* Open data and digital morphology. *Proc. R. Soc. B* **284**, 20170194

(2017).

6. Gutiérrez, Y., Ott, D., Töpperwien, M., Salditt, T. & Scherber, C. X-ray computed

tomography and its potential in ecological research: a review of studies and optimization of

specimen preparation. *Ecol. Evol.* **8**, 7717-7732 (2018).

7. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat.*

Methods **9**, 676-682 (2012).

8. Ketcham, R. A. Computational methods for quantitative analysis of three-dimensional

features in geological specimens. *Geosphere* **1**, 32-41 (2005).

9. Page, L. M., MacFadden, B. J., Fortes, J. A., Soltis, P. S. & Riccardi, G. Digitization of

biodiversity collections reveals biggest data on biodiversity. *BioScience* **65**, 841-842 (2015).

10. Faulwetter, S., Vasileiadou, A., Kouratoras, M., Dailianis, T. & Arvanitidis, C. Micro-

computed tomography: Introducing new dimensions to taxonomy. *ZooKeys* **263**, 1-45 (2013).

11. Akkari, N. *et al*. New avatars for Myriapods: complete 3D morphology of type specimens

transcends conventional species description (Myriapoda, Chilopoda). *PLoS ONE* **13**,

0200158; 10.1371/journal.pone.0200158 (2018).

12. Fontaine, B., Perrard, A. & Bouchet, P. 21 years of shelf life between discovery and

description of new species. *Curr. Biol*. **22**, R943-R944 (2012).

13. Hipsley, C. A. & Sherratt, E. Psychology, not technology, is our biggest challenge to

open digital morphology data. *Sci. Data*. **6**, 41 (2019).

14. Blagoderov, V., Kitching, I. J., Livermore, L., Simonsen, T. J. & Smith, V. S. No

specimen left behind: industrial scale digitization of natural history collections. *Zookeys* **209**,

133-146 (2012).

15. Rogers, N. Museum drawers go digital. *Science* **352**, 762-765 (2016).

16. Meineke, E. K., Davies, T. J., Daru, B. H. & Davis, C. C. Biological collections for

understanding biodiversity in the Anthropocene. *Phil. Trans. R. Soc. B* **374**, 20170386

(2018).

17. Schmitt, C. J., Cook, J. A., Zamudio, K. R. & Edwards, S.V. Museum specimens of

terrestrial vertebrates are sensitive indicators of environmental change in the Anthropocene.

Phil. Trans. R. Soc. B **374**, 20170387 (2018)

18. Sherratt, E., Gower, D. J., Klingenberg, C. P. & Wilkinson, M. Evolution of cranial shape

in caecilians (Amphibia: Gymnophiona). *Evol. Biol.* **41**, 528-545 (2014).

19. Watanabe, A. *et al.* Ecomorphological diversification in squamates from conserved

pattern of cranial integration. *Proc. Natl. Acad. Sci.* **116**, 14688-14697 (2019).

20. Simon, M. N., Machado, F. A. & Marroig, G. High evolutionary constraints limited

adaptive responses to past climate changes in toad skulls. *P. Roy. Soc. B-Biol. Sci.* **283**,

20161783 (2016).

21. Sherratt, E., Serb, J. M. & Adams, D. C. Rates of morphological evolution, asymmetry

and morphological integration of shell shape in scallops. *BMC Evol. Biol.* **17**, 248 (2017).

22. Chira, A. M. *et al.* Correlates of rate heterogeneity in avian ecomorphological traits. *Ecol.*

Lett. **21**, 1505-1514 (2018).

23. Percival, C. J. *et al.* The effect of automated landmark identification on morphometric

analyses. *J. Anat.* **234**, 917-935 (2019).

24. Bouxsein, M. L. *et al.* Guidelines for assessment of bone microstructure in rodents using

micro –computed tomography. *J. Bone Miner. Res.* **25**, 1468-1486 (2010).

25. Broeckhoven, C. & du Plessis, A. X-ray microtomography in herpetological research: a

review. *Amphibia-Reptilia* **39**, 377-401 (2018).

26. Marcy, A. E., Fruciano, C., Phillips, M. J., Mardon, K. & Weisbecker, V. Low resolution

scans can provide a sufficiently accurate, cost- and time-effective alternative to high

resolution scans for 3D shape analyses. *PeerJ* **6**, 5032; 10.7717/peerj.5032 (2018).

27. Gray, J. A., McDowell, M. C., Hutchinson, M. N. & Jones, M. E. Geometric

morphometrics provides an alternative approach for interpreting the affinity of fossil lizard

jaws. *J. Herpetol.* **51**, 375-382 (2017).

28. Thorn, K. M., Hutchinson, M. N., Archer, M. & Lee, M. S. Y. A new scincid lizard from

the Miocene of northern Australia, and the evolutionary history of social skinks (Scincidae:

Egerniinae). *J. Vertebr. Paleontol.* **39**, 1 (2019).

29. Chaplin, K., Sumner, J., Hipsley, C. A. & Melville, J. An integrative approach using

phylogenomics and high-resolution x-ray computed tomography for species delimitation in

cryptic taxa. *Syst. Biol.* **69**, syz048; 10.1093/sysbio/syz048 (2019).

30. Melville, J. *et al.* Integrating phylogeography and high-resolution x-ray CT reveals five

new cryptic species and multiple hybrid zones among Australian earless dragons. *R. Soc.*

Open Sci. **6**, 191166; 10.1098/rsos.191166 (2019).

31. Caro, A., Gómez-Moliner, B. J. & Madeira, M. J. Integrating multilocus DNA data and

3D geometric morphometrics to elucidate species boundaries in the case of *Pyrenaearia*

(Pulmonata: Hygromiidae). *Mol. Phylogenet. Evol.* **132**, 194-206 (2019).

32. Winkelmann, C. T. & Wise, L. D. High-throughput micro-computed tomography imaging

as a method to evaluate rat and rabbit fetal skeletal abnormalities for developmental toxicity

studies. J. Pharmacol. Tox. Met. **59**, 156-165 (2009).

33. Sevilla, R. S. *et al.* Development and optimization of a high-throughput micro-computed

tomography imaging method incorporating a novel analysis technique to evaluate bone

mineral density of arthritic joints in a rodent model of collagen induced arthritis. *Bone* **73**, 32-

41 (2015).

34. Wong, M. D., Maezawa, Y., Lerch, J. P. & Henkelman, R. M. Automated pipeline for

anatomical phenotyping of mouse embryos using micro-CT. *Development* **141**, 2533-2541

(2014).

35. Wu, D. *et al.* Combining high-throughput micro-CT-RGB phenotyping and genome-wide

association study to dissect the genetic architecture of tiller growth in rice. *J. Exp. Bot.* **70**,

545-561 (2019).

36. Ding, Y. *et al.* Computational 3D histological phenotyping of whole zebrafish by X-ray

histotomography. *Elife* **8**, 44898; 10.7554/eLife.44898.001 (2019).

37. Staedtler, Y. M., Masson, D. & Schönenberger, J. Plant tissues in 3D via X-ray

tomography: simple contrasting methods allow high resolution imaging. *PLoS ONE* **8**, 75295;

10.1371/journal.pone.0075295 (2013).

38. Keklikoglou, K. *et al.* Micro-computed tomography for natural history specimens: a

handbook of best practice protocols. *Eur. J. Taxon.* **522**, 1-55 (2019).

39. Adams, D., Collyer, M. & Kaliontzopoulou, A. Geomorph: Software for geometric

morphometric analyses. R package version 3.1.0. https://cran.r-

project.org/package=geomorph (2019).

40. Klingenberg, C. P. MorphoJ: an integrated software package for geometric

morphometrics. *Mol. Ecol. Resour.* **11**, 353-357 (2011).

41. du Plessis, A., Broeckhoven, C., Guelpa, A. & le Roux, S. G. Laboratory x-ray micro-

computed tomography: a user guideline for biological samples. *Gigascience* **6**, 1-11 (2017).

42. Hocknull, S. A., Zhao, J. X., Feng, Y. X. & Webb, G. E. Responses of middle Pleistocene

rainforest vertebrates to climate change in Australia. *Earth Planet. Sc. Lett.* **264**, 317-331

(2007).

43. Hedrick, B. P. *et al.* Digitization and the future of natural history collections. *BioScience*

70, 243-251 (2020).

44. Lawrence, R. A. & Hocknull, S. Engaging the public with small vertebrate fossils and

utilizing citizen science to maximise scientific discovery at Capricorn Caves, Central Eastern

Queensland, Australia. *J. Vertebr. Paleontol. Program and Abstracts* **139** (2019).

45. Long, J. A., Young, G. C., Holland, T., Senden, T. J. & Fitzgerald, E. M. An exceptional

Devonian fish from Australia sheds light on tetrapod origins. *Nature* **444**, 199-202 (2006).

46. Arbour, J. H., Curtis, A. A. & Santana, S. E. Signatures of echolocation and dietary

ecology in the adaptive evolution of skull shape in bats. *Nat. Commun.* **10**, 2036 (2019).

47. Park, T., Fitzgerald, E. M. & Evans, A. R. Ultrasonic hearing and echolocation in the

earliest toothed whales. *Biol. Lett.* **12**, 20160060; 10.1098/rsbl.2016.0060 (2016).

48. Müller, J. *et al.* Eocene lizard from Germany reveals amphisbaenian origins. *Nature* **473**,

364-367 (2011).

49. Miralles, A. *et al.* Distinct patterns of desynchronized limb regression in Malagasy

scincine lizards (Squamata, Scincidae). *PLoS ONE* **10**, 0126074;

10.1371/journal.pone.0126074 (2015).

50. Weisbecker, V. Monotreme ossification sequences and the riddle of mammalian skeletal

development. *Evolution* **65**, 1323-1335 (2011).

51. Newton, A. H. *et al.* Letting the 'cat' out of the bag: pouch young development of the

extinct Tasmanian tiger revealed by X-ray computed tomography. *R. Soc. Open Sci.* **5**,

171914; 10.1098/rsos.171914 (2018).

52. Hublin, J. J. *et al.* New fossils from Jebel Irhoud, Morocco and the pan-African origin of

Homo sapiens. *Nature* **546**, 289-292 (2017).

53. Beaudet, A. & Gilissen, E. Fossil primate endocasts: perspectives from advanced imaging

techniques in *Digital Endocasts: from Skulls to Brains* (eds. Bruner, E., Ogihara, N. &

Tanabe, H.) 47-58 (Springer, 2018).

54. Wulff, N. C., Lehmann, A. W., Hipsley, C. A. & Lehmann, G. U. C. Copulatory courtship

by bushcricket genital titillators revealed by functional morphology, μCT scanning for 3D

reconstruction and female sense structures. *Arthropod Struct. Dev.* **44**, 388-397 (2015).

55. Gee, C. T. Applying microCT and 3D visualization to Jurassic silicified conifer seed

cones: a virtual advantage over thin-sectioning. *Appl. Plant Sci.* **1**, 1300039;

10.3732/apps.1300039 (2013).

56. Meyer, M. *et al.* Three-dimensional microCT analysis of the Ediacara fossil *Pteridinium*

simplex sheds new light on its ecology and phylogenetic affinity. *Precambrian Res.* **249**, 79-

87 (2014).

57. Gooday, A. J., Sykes, D., Goral, T., Zubkov, M. V. & Glover, A.G. Micro-CT 3D

imaging reveals the internal structure of three abyssal xenophyophore species (Protista,

Foraminifera) from the eastern equatorial Pacific Ocean. *Sci. Rep.* **8**, 12103; 10.1038/s41598-

018-30186-2 (2018).

58. Dunlop, J. A. *et al.* Microtomography of the Baltic amber tick *Ixodes succineus* reveals

affinities with the modern Asian disease vector *Ixodes ovatus*. *BMC Evol. Biol.* **16**, 203

(2016).

Figures

Figure 1. Equipment for dense fossil packing prior to μCT scanning: (a) litter tray, (b) paper

and/or plastic straws, (c) clear two-piece pharmaceutical capsules, (d) archival tissue, (e) dry

fossil specimen, (f) supporting material (paper, foam), (g) centrifuge tube with cap, (h)

archival paper for labels shown inside a size 4 capsule, (i) forceps, (j) archival pen. Estimated

costs and material suppliers are listed in Table 1.

Figure 2. Specimen labelling and packing: (a) 50 ml Falcon tube with a labelled straw and

capsules, (b) a single fossil wrapped in Green's Lens tissue to minimise movement during

image acquisition, (c) loading capsules into the paper straw using a thinner cocktail straw to

gently push them down, (d) a packed 50 ml Falcon tube mounted on a glass rod inside the

μCT machine; note that the cap-side is facing down (e) the packed tube from above, and (f)

the diagram of straw arrangements in (e), noting the position of a larger blue straw and paper

used as a divider.

Figure 3. High-throughput μCT results: (a) the first X-ray image from each scanned segment

of the 50 ml tube from top (tip) to bottom (cap), containing 72 frog and lizard specimens. (b)

3D renderings of the individual segments in the same positions, at 17.8 μm voxel size, (c) the

stitched 3D volume at the same resolution, with fossils colour-coded by straw, (d)

reconstructed cross-section of the same tube in side and (e) top views, (f) cross-section of an

agamid lizard jaw showing contrast between the matrix, bone, and teeth, (g) cross-section of

a skink mandible with a shadow motion artefact, (h) an example of beam hardening caused

by a dense surface.

Figure 4. Scanned fossil material and its applications: (a) 3D rendering of 73 fossil frog ilia

scanned in a single tube, (b) a subsample of varanid osteoderms in 3D, (c) mandible of the

rainforest rodent *Pogonomys* from the Riversleigh World Heritage site with molars colour

coded and jaw bone rendered semi-transparent, (d) surface mesh of the same specimen

composed of 109,213 vertices (file size 10.5 MB), (e) photograph of children holding a 3D

print of enlarged fossil skink jaws at Capricorn Caves Fossil Open Day, (f) point cloud

comparison of fossil tree frog (Hylidae) ilia found in Capricorn Caves at 10-20 cm depth

(left) and 40-50 cm depth (right), representing a time difference of 2-3,000 years. Warmer

colours denote regions with greater shape differences, as estimated in CloudCompare

v.2.10.2.

Tables

Table 1. Estimated costs in Australian dollars (AUD) for material to μCT scan a 50 ml

cylinder fully packed with 84 capsules holding 1 small (< 1 cm long) specimen each. Note

that many items (e.g., Falcon tube, straws, forceps, foam, litter tray, pen) can be reused

multiple times, thus decreasing the cost for subsequent scans.

Item	Total cost per scan (AUD)
Centrifuge Falcon Tube 50 mL polypropylene (PP), conical bottom with screw cap[1]	0.60
Hard 2-piece clear gelatine capsules, size 4 (volume 0.23 ml)[2*]	2.10
Standard paper straw (6 x 135 mm)[3]	0.27
Green's Lens Tissue, un-buffered TIS-L Soft, acid-free, long fibred 9gsm repair/wrapping tissue (609 x 914 mm)[4]	0.76
Feather Forcep Sharp E122S, 0.23 mm gauge thick[5]	4.90
Ethafoam 220, medium density, non-cross linked Polyethylene (PE) closed cell foam (2400 x1200 mm)[6]	1.00
Litter tray[7]	4.00
Label paper[8]	0.10
Archival pen [9]	6.54
TOTAL	20.30

Suppliers: [1]https://corning.com.au; [2]https://gelcapsules.com.au; [3]https://www.greenpack.com.au;
[4]http://www.archivalsurvival.com.au; [5]http://www.entosupplies.com.au;
[6]https://www.foamsales.com.au; [7]https://www.kmart.com.au; [8]https://www.talasonline.com;
[9]https://www.newtownartsupplies.com.au/
*Clear 2-piece vegetable capsules are also available, although we did not test their performance in
μCT scans.
584

586   Table 2. Estimated movement among 408 fossils μCT scanned using various packing

587   materials to secure the specimen inside each capsule.

Packing material	Number of specimens per capsule (size 4) that moved during scanning			Archival?	Comments
	1-2	3	> 3		
Green's Lens[1]	0/100 (0%)	2/39 (5.1%)	25/50 (50%)	Yes	Ideal consistency
Kim Tech wipes[2]	0/100 (0%)	4/39 (10.3%)	27/50 (54%)	No	100% fibre, not acid free
Ethafoam 220[3]	10/10 (100%)	10/10 (100%)	10/10 (100%)	No	Slippery and hard
Tyvek[4]	-	-	-	Yes	Not tested, too waxy
Renaissance[5]	-	-	-	Yes	Not tested, too rigid

589   Suppliers: [1]http://www.archivalsurvival.com.au; [2]https://www.kcprofessional.com.au;
590   [3]https://www.foamsales.com.au; [4]http://www.dupont.com.au; [5]http://www.lightimpressionsdirect.com