Failure to maternally reprogram histone methylation causes developmental delay due to germline transcription in somatic tissues

Brandon S. Carpenter[1], Teresa W. Lee[1], Caroline F. Plott[2], Jovan S. Brockett[3], Dexter A. Myrick[1], David J. Katz[1]*

[1]Department of Cell Biology, Emory University School of Medicine, Atlanta GA 30322, USA.

[2]Department of Genetics and Development, Columbia University, New York NY 10032, USA.

[3]Department of Biology, Oglethorpe University, Atlanta GA 30319, USA.

*corresponding author:
David J. Katz
Associate Professor
Department of Cell Biology
Room 443, Whitehead Biomedical Research Building
Emory University School of Medicine
Atlanta, GA 30322, USA
Phone: (404) 727-3403
djkatz@emory.edu

ABSTRACT

In *C. elegans*, the H3K36 methyltransferase, MES-4, helps establish germ cell fate by

maintaining H3K36me2/3 at germline genes between generations. Previously, we showed

that the H3K4me2 demethylase, SPR-5, and the H3K9 methyltransferase, MET-2,

reprogram histone methylation at fertilization to prevent the ectopic expression of

germline genes in somatic tissues. Together, this indicates that SPR-5 and MET-2 maternal

reprogramming may antagonize MES-4 to establish germline versus soma. Here, we show

that *spr-5; met-2* mutant progeny have a severe developmental delay that is associated with

the ectopic maintenance of H3K36me2/3 at MES-4 targeted germline genes in somatic

tissues, and the ectopic expression of these genes. We further show that the developmental

delay and the ectopic expression are dependent upon MES-4. Thus, we propose that SPR-5,

MET-2, and MES-4 balance inherited histone methylation to establish germline versus

soma. Without this balance, the inappropriate transcription of germline genes in somatic

tissues causes developmental delay.

INTRODUCTION

In multicellular organisms, developmental cell fate decisions are established by

tightly controlled spatial and temporal gene expression. This pattern of gene expression

can be regulated by heritable modifications on the N-terminal tails of histone proteins

(Jambhekar et al., 2019). For example, methylation of lysine 4 and 36 on histone 3,

(H3K4me and H3K36me) are generally associated with active transcription, whereas

methylation of lysine 9 on the same histone (H3K9me) is commonly associated with

transcriptional repression. These modifications are dynamically regulated by the specific

and tightly controlled activity of histone modifying enzymes. For instance, mono- and di-

methylation of lysine 4 on histone H3 (H3K4me1/2) are removed by the demethylase

LSD1/KDM1A (Y. Shi et al., 2004; Y.-J. Shi et al., 2005). In the nematode *C. elegans*,

populations of mutants lacking the LSD1 ortholog, SPR-5, become increasingly sterile over

~30 generations (Katz et al., 2009). Failure to erase H3K4me2 at fertilization between

generations in *spr-5* mutants correlates with an accumulation of H3K4me2 and

spermatogenesis gene expression across 30 generations, which leads to increasing sterility

(Katz et al., 2009). These data demonstrate that H3K4me2 can function as an epigenetic

transcriptional memory. More recently, it was demonstrated that SPR-5 synergizes with

the H3K9me2 methyltransferase MET-2, to regulate maternal epigenetic reprogramming

(Kerr et al., 2014). MET-2 is the ortholog of the mammalian SETDB1. Progeny of mutants

lacking both SPR-5 and MET-2 become completely sterile in a single generation and display

synergistic increases in both H3K4me2 and candidate germline gene expression in somatic

tissues (Kerr et al., 2014). Together this work supports a model in which SPR-5 and MET-2

are maternally deposited into the oocyte, where they reprogram histone methylation to

prevent somatic expression of germline genes.

Following fertilization, the *C. elegans* embryo separates germline versus somatic

lineages progressively through a series of asymmetric divisions (Strome, 2005). To

accomplish this, transcription factors coordinate with multiple histone modifications.

Maternal deposition of PIE-1, a germline specific protein that asymmetrically segregates

into germline blastomeres (P lineage cells), maintains the fate of germ cells by inhibiting

POL-II elongation and preventing the ectopic expression of somatic genes (Batchelder et al.,

1999; Mello et al., 1992; Seydoux et al., 1996). In the absence of transcription in the

germline, the maternally provided H3K36me2/3 methyltransferase, MES-4, binds to a

subset of germline genes that were previously expressed in the parental germline

(Furuhashi et al., 2010; Rechtsteiner et al., 2010). These germline genes are recognized by

MES-4 via H3K36me2/3 that was added in the parental germline by the transcription-

coupled H3K36me2/3 methyltransferase, MET-1 (Kreher et al., 2018). MES-4 maintains

H3K36me2/3 at these genes in the early embryo in a transcriptionally independent

manner. Without maternally deposited MES-4, the germline cannot properly proliferate

and animals are sterile (Capowski et al., 1991; Garvin et al., 1998). For the remainder of

this study, we will refer to these MES-4 targeted genes as MES-4 germline genes, and the

process through which MES-4 maintains these genes for re-activation in the primordial

germ cells of the offspring, as bookmarking.

In addition to providing transcriptional memory of germline genes between

generations in the germline, histone methylation has been implicated in repressing

germline transcription in somatic tissues. Recent work in *C. elegans* has shown that a

subset of germline genes have H3K9me2 enrichment at their promoters in somatic tissues

(Rechtsteiner et al., 2019). Loss of LIN15B, a transcriptional repressor that is a putative

member of the DREAM complex, reduces the enrichment of H3K9me2, leading to the

ectopic accumulation of H3K36me3 at gene bodies in somatic tissues (Rechtsteiner et al.,

2019). These data implicate H3K9me2 in germline gene repression in somatic tissues and

raise the possibility that H3K9me2 antagonizes MES-4. As mentioned earlier, previous

work suggests that H3K9me2 also functions antagonistically to H3K4me2 (Kerr et al.,

2014). However, whether the maternal histone modifying enzymes SPR-5 and MET-2

coordinate with MES-4 to ensure the proper inheritance of histone methylation and the

proper specification of germline versus soma remains unclear. To address this question, we

examined somatic development in progeny deficient in SPR-5 and MET-2 maternal

reprogramming. We find that *spr-5; met-2* double mutant progeny (hereafter referred to as

spr-5; met-2 progeny) display a severe developmental delay that is rescued by removing

MES-4. In addition, H3K36me3 ectopically accumulates at germline genes in the somatic

tissues of *spr-5; met-2* progeny. This accumulation correlates with the ectopic expression of

MES-4 germline genes and is rescued by knocking down MES-4 activity. These data

demonstrate that the ectopic expression of MES-4 targeted germline genes in somatic

tissues causes a developmental delay. We propose a model where SPR-5 and MET-2

function together during maternal reprogramming to restrict MES-4 dependent

bookmarking of germline genes in somatic tissues, thereby ensuring proper specification of

germline versus somatic cell fates.

RESULTS

Loss of *spr-5* and *met-2* causes developmental delay

85   Previous work from our lab demonstrated that progeny from *spr-5; met-2* mutants

express germline genes in somatic lineages (Kerr et al., 2014). To examine how

misexpression of germline genes in somatic tissues affects somatic development, we

synchronized embryos laid by N2, *spr-5*, *met-2*, and *spr-5; met-2* mutant hermaphrodites

and monitored their development from hatching to adults. By 72 hours past the

synchronized lay, all N2 progeny (469/469), most of the *spr-5* progeny (363/385), and

many of *met-2* progeny (386/450) were fertile adults (Figure 1 A, B, C, and E, S1A-C). In

contrast, *spr-5; met-2* progeny displayed a severe developmental delay. By 72 hours past

the synchronized lay, none of the progeny (0/463) had reached adulthood (Figure 1 D, E;

and S1D). The majority of *spr-5; met-2* progeny (371/463) resembled L2 larvae at 72 hours,

while a smaller percentage of the population developed to later larval stages (42/463)

(Table S1). By seven days post synchronized lay, a small number of *spr-5; met-2* progeny

(35/876) developed into adults and the majority (31/35) of these adults displayed

protruding vulva (Pvl) (Figure S1E-G). All 35 of the *spr-5; met-2* mutant progeny that

developed to adulthood were sterile.

MES-4 germline genes are ectopically expressed in *spr-5; met-2* mutant soma

Previously we showed that H3K4me2 is synergistically increased in *spr-5; met-2*

progeny compared to *spr-5* and *met-2* single mutant progeny, and that this increase in

H3K4me2 correlates with a synergistic increase in candidate germline gene expression in

somatic tissues (Kerr et al., 2014). To test the extent to which germline genes are

ectopically expressed in somatic tissues, we examined somatic expression genome-wide.

To do this, we performed RNA-seq on *spr-5; met-2* L1 progeny compared to N2 (wild-type)

L1 progeny. We chose to perform this analysis on L1 larvae because this stage immediately

precedes the L2 larval delay that we observe in *spr-5; met-2* progeny (see Figure 1A, D). In

addition, L1 larvae are composed of 550 somatic cells and 2 germ cells. Therefore, L1

larvae are primarily composed of somatic tissue. As a control, we also performed RNA-seq

on L1 progeny from *spr-5* and *met-2* single mutants that were isolated from early

generation animals (within 1-5 generations). These generations are well before the onset

of sterility that we previously reported (Katz et al., 2009; Kerr et al., 2014). We identified

778 differentially expressed transcripts in *spr-5; met-2* progeny compared to N2 (Figure

S2A-B, Figure S3C, F, and Table S2), many of which also overlap with genes differentially

expressed in *spr-5* (159/413) and *met-2* single mutants (113/343) compared to N2.

(Figure S2A-B, Figure S3A, B, S3D, E, and Table S2). Gene Set Enrichment Analysis (GSEA)

did not identify any categories of genes misexpressed in the *spr-5* or *met-2* single mutants.

However, GSEA revealed that genes differentially expressed in *spr-5; met-2* progeny were

significantly enriched (based on Combined Score, (Chen et al., 2013)) for biological

processes and cellular components characteristic of the germline; including meiosis, P-

granules and negative regulation of the cell cycle (Figure S2C-D). Many of the genes

involved in these processes are expressed in the germline of the parental generation,

bound by the H3K36 methyltransferase MES-4 in the early embryo, and marked by

H3K36me2/3, independent of POL-II (Rechtsteiner et al., 2010). As a result, we were

interested in the potential overlap with MES-4 targeted genes (referred to as MES-4

germline genes). Rechtsteiner and colleagues identified 214 MES-4 germline genes

(Rechtsteiner et al., 2010). 17 of these genes are pseudogenes that we were unable to

convert from Ensembl transcript IDs to RefSeq mRNA accession, and another gene was

duplicated, so we removed those genes from our analysis. This leaves 196 MES-4 germline

genes. We reasoned that the absence of SPR-5 and MET-2 reprogramming may cause

germline genes to be aberrantly targeted by MES-4 in the soma, leading to ectopic

expression. To investigate this possibility, we examined the overlap between differentially

expressed genes in *spr-5; met-2* L1 progeny and MES-4 germline genes. Out of 196 MES-4

germline genes, 34 overlapped with genes up-regulated in *spr-5; met-2* progeny compared

to N2 (hypergeometric test (P-value < 4.34e-20, Figure 2A), while zero overlapped with

genes down-regulated in *spr-5; met-2* progeny compared to N2 (Figure 2B). In addition,

when we compared the log2 fold change (FC) in expression of all of the MES-4 genes in *spr-*

5, *met-2*, and *spr-5; met-2* mutant progeny compared to N2, we observed that 108 of the

MES-4 germline genes are synergistically increased in *spr-5; met-2* progeny compared to

single mutant progeny (Figure 2C and Table S3). This analysis included only 176 out of the

196 MES-4 germline genes because the remaining 21 genes do not have an expression

value in one or more of the data sets (*spr-5*, *met-2,* or *spr-5; met-2*).

To confirm that MES-4 germline genes are somatically expressed in *spr-5; met-2* L1

progeny, we performed single molecule fluorescent *in situ* hybridization (smFISH) on two

MES-4 germline targets, *htp-1* and *cpb-1* (Figure 3). Both of these genes were amongst the

genes that were ectopically expressed in *spr-5; met-2* L1 progeny compared to N2 L1

progeny. In N2 L1 larvae, *htp-1* (Figure 3A-C, insets) and *cpb-1* (Figure 3G-I, insets) were

restricted to the two primordial germ cells, Z2 and Z3. In contrast, in *spr-5; met-2* progeny

htp-1 was ectopically expressed throughout the soma (Figure 3D-F), similar to what we

observe with the ubiquitously expressed subunit of RNA polymerase II, *ama-1* (Figure S4A-

L), which was unchanged in our RNA-seq analysis. *cpb-1* wa*s* also ectopically expressed,

though the ectopic expression was not ubiquitous as it is in *htp-1* (Figure 3J, L).

MES-4 dependent germline genes display ectopic H3K36me3 in *spr-5; met-2* mutants

To test whether MES-4 germline genes that are ectopically expressed in the soma of

spr-5; met-2 progeny also ectopically maintain H3K36me3, we performed H3K36me3 ChIP-

seq. Similar numbers of H3K36me3 peaks were called between N2 (replicate 1 = 5507 and

replicate 2 = 3878) and *spr-5; met-2* (replicate 1 = 5886 and replicate 2 = 1938) progeny,

with most of the peaks overlapping between replicates (Figure S5Q-R). MES-4 germline

genes have low levels of H3K36me3 in N2 L1 progeny. However, compared to N2 L1

progeny, *spr-5; met-2* L1 progeny displayed increased enrichment for H3K36me3 across

the gene bodies at MES-4 germline genes (Figure 4A, B; Figure S5A, B). For example, the

MES-4 germline genes *cpb-1*, *T05B9.1, Y18D10A.11*, and *fbxa-101,* that are ectopically

expressed in our RNA-seq analysis, showed increased levels of H3K36me3 in *spr-5; met-2*

progeny (Figure 4J-M; Figure S5J-M) compared to N2 progeny (Figure 4 C-F; Figure S5C-F).

As a control, we examined H3K36me3 enrichment at *ceh-13*, a gene enriched in

hypodermal and ventral nerve chord in L1 progeny (Figure 4G, N; Figure S5G, N), *ama-1*, a

subunit of RNA polymerase II that is expressed ubiquitously (Figure 4H, O; Figure S5H, O),

and *act-1*, a ubiquitously expressed actin related protein (Figure 4I, P; Figure S5I, P). These

control genes displayed similar H3K36me3 enrichment in both *spr-5; met-2* and N2 L1

progeny (compare Figure 4N-P and Figure S5N-P to Figure 4G-I and Figure S5G-I,

respectively).

MES-4 germline genes display H3K9me2 at their promoter peaks

Recent work discovered that a subset of germline specific genes contain H3K9me2

peaks at their promoters in N2 L1 progeny (Rechtsteiner et al., 2019). This finding

implicates H3K9me2 enrichment at promoters of germline genes as being a critical

component for repressing germline genes in somatic tissues. If SPR-5 and MET-2 are

functioning to prevent MES-4 germline genes from being ectopically expressed in somatic

tissues, we would expect MES-4 germline genes that are ectopically expressed in the

somatic tissues of *spr-5; met-2* progeny to be targeted by H3K9 methylation in these

tissues. To examine this possibility, we re-analyzed L1 H3K9me2 Chip-seq data from

Rechsteiner et al. 2019 (Rechtsteiner et al., 2019). This re-analysis shows that many of the

MES-4 germline genes were enriched for H3K9me2 at their promoters (Figure S6A),

including the majority of MES-4 germline genes that are ectopically expressed in the soma

of *spr-5; met-2* progeny (Figure S6B). For example, the MES-4 germline genes *cpb-1*,

T05B9.1, *Y18D10A.11*, and *fbxa-101*, that accumulate ectopic H3K36me3 and were

ectopically expressed in the somatic tissues of *spr-5; met-2* progeny, have H3K9me2 peaks

at their promoters (Figure 6SC-F). In contrast, control genes, *ceh-13*, *ama-1*, and *act-1*, that

are not expressed ectopically in the somatic tissues of *spr-5; met-2* progeny were not

enriched for H3K9me2 at their promoters (Figure 6SG-I).

Knocking down MES-4 rescues ectopic expression of germline genes in *spr-5; met-2*

mutant soma

To test whether the ectopic expression of MES-4 germline genes in *spr-5; met-2*

progeny is dependent on the ectopic H3K36me3, we examined whether the expression of

these genes was dependent upon MES-4. We performed quantitative real-time PCR (qRT-

PCR) on L1 progeny from *spr-5; met-2* hermaphrodites fed control (L4440) RNAi versus

mes-4 RNAi (Figure 5). For this analysis, we selected candidate MES-4 germline genes that

were ectopically expressed and displayed an ectopic H3K36me3 peak in *spr-5; met-2* L1

progeny compared to N2 L1 progeny. Consistent with our RNA-seq analysis, all nine of the

candidate MES-4 germline genes that we examined were ectopically expressed >2 fold in

spr-5; met-2 L1 progeny compared to control (Figure 5). Strikingly, the ectopic expression

of the nine MES-4 candidate germline genes was dependent upon MES-4. Nine out of nine

of these genes were significantly decreased in L1 progeny from *spr-5; met-2*

hermaphrodites treated with *mes-4* RNAi (Figure 5; p-value < 0.001), and all but one

(*T05B9.1*) were reduced to levels that were similar to N2 L1 progeny.

Knocking down MES-4 rescues developmental delay in *spr-5; met-2* progeny

To test whether the developmental delay phenotype that we observe in *spr-5; met-2*

progeny is also dependent on the ectopic expression of MES-4 germline genes, we fed *spr-5;*

met-2 hermaphrodites *mes-4* RNAi and monitored their progeny for 72 hours after a

synchronized lay. If the developmental delay is dependent upon the ectopic expression of

MES-4 germline genes, it should be suppressed when this ectopic expression is eliminated

via *mes-4* RNAi. 72 hours after a synchronized lay, all of the N2 control progeny from

hermaphrodites fed either L4440 control (1089/1089) or *mes-4* (1102/1102) RNAi were

adults (Figure 6A, B, I). Also, consistent with our previous observations, all but one of the

spr-5; met-2 mutant progeny (729/730) from hermaphrodites fed control RNAi remained

in the L2-L3 larval stages (Figure 6E, I). In contrast, most of *spr-5; met-2* progeny

(569/618) from hermaphrodites fed *mes-4* RNAi developed to adults (Figure 6F, I). Though

as expected, these animals remained sterile due to the *mes-4* RNAi preventing any germline

formation.

MET-1 mediates transcription-coupled H3K36me3 in the adult germline and in later

stage transcriptionally active embryos (Furuhashi et al., 2010; Kreher et al., 2018;

Rechtsteiner et al., 2010). If the ectopic somatic expression of MES-4 germline genes is

caused by the transcription independent maintenance of H3K36me3 via MES-4, we would

expect the developmental delay to be independent of MET-1. To determine whether the

developmental delay phenotype in *spr-5; met-2* progeny is dependent on MET-1, we

monitored the development of progeny from *spr-5; met-2* hermaphrodites fed *met-1* RNAi.

In contrast to the suppression that we observed upon *mes-4* RNAi, *met-1* RNAi had no

effect. None of the *spr-5; met-2* mutant progeny (0/573) from hermaphrodites fed *met-1*

RNAi developed to adults, and these animals were indistinguishable from *spr-5; met-2*

progeny fed control RNAi (Figure 6G, I). Additionally, *met-1* RNAi itself had no effect on

developmental delay, as nearly all of the N2 progeny (715/721) from hermaphrodites fed

met-1 RNAi developed to adults (Figure 6C, I).

Knocking down SET-2 rescues *spr-5; met-2* developmental delay

If the developmental delay of *spr-5; met-2* mutants is caused by ectopic H3K4me2

driving the expression MES-4 germline genes in somatic tissues, we would expect the

developmental delay of *spr-5; met-2* progeny to be dependent upon the H3K4

methyltransferase, SET-2. To test this, we monitored the development of progeny of *spr-5;*

met-2 hermaphrodites fed *set-2* RNAi. Identical to N2 progeny from hermaphrodites fed

control RNAi, *set-2* RNAi had no effect on the development of wild-type animals, as all of

the N2 progeny from hermaphrodites fed *set-2* RNAi developed to adults (1114/1114)

(Figure 6D, I). However, in contrast to *spr-5; met-2* progeny fed control RNAi that were

developmentally delayed, most of the progeny from *spr-5; met-2* hermaphrodites fed *set-2*

RNAi developed to adults (347/384) (Figure 6H, I).

***spr-5; met-2* progeny acquire transgene silencing in somatic tissues**

The somatic expression of MES-4 germline genes involved in germline transgene

silencing (Figure S2C-D) raises the possibility that the somatic tissues in *spr-5; met-2*

progeny may acquire the ability to silence transgenes, a function normally restricted to

germline cells. To test this, we examined the somatic expression of an extrachromosomal

multicopy *let-858* transgene that is normally silenced in the germline by both

transcriptional and posttranscriptional germline silencing mechanisms (Kelly and Fire,

1998). This analysis was performed in *spr-5; met-2* mutant L2 larvae that were undergoing

developmental delay. In N2, most of the L2 progeny (117/132) expressed ubiquitous high

levels of LET-858::GFP throughout the entire soma (Figure 7A, B, and K), while the

remaining progeny (15/132) expressed what we describe as a "faint" level of expression

(Figure 7C, D, and K). In contrast, almost none of the *spr-5; met-2* L2 progeny (2/87)

displayed high level of transgene expression that is comparable to the high level seen in

most N2 progeny (Figure 7E, F, and K). Instead, the majority of *spr-5; met-2* progeny

(64/87) had faint LET-858::GFP expression that is comparable to the faint expression

observed in N2 progeny (Figure 7G, H, and K). The remaining 21 *spr-5; met-2* progeny had

no LET-858::GFP expression. Because less than 50% of progeny inherited the *let-858*

transgene, we normalized the percentage of progeny scored as "off" for LET-858::GFP to

presence of the *let-858* transgene based on genotyping for *gfp* (see methods). For N2, all of

the 60 L2 progeny that were scored as "off" failed to inherit the *let-858* transgene,

indicating that the transgene is never silenced in N2 progeny (data not shown). However,

for *spr-5; met-2* L2 progeny, eight out of 50 progeny that were scored as "off" inherited the

let-858 transgene indicating that the *let-858* transgene can be completely silenced in some

spr-5; met-2 progeny. After normalization for the transgene inheritance, we observed that

21/87 of *spr-5; met-2* progeny displayed no visible expression of the LET-858::GFP

transgene (Figure 7I, J, and K).

DISCUSSION

Maternal regulation of inherited histone methylation contributes to the specification

of germline versus soma

SPR-5 and MET-2 are thought to act maternally to reprogram histone methylation

and prevent the transcriptional "baggage" of the parental germline from being

inappropriately transmitted to the offspring (Kerr et al., 2014). In this study, we find that

the lack of SPR-5 and MET-2 maternal reprogramming led to a severe developmental delay.

This developmental delay was associated with the ectopic expression of MES-4 germline

genes in somatic tissues. This finding raises the possibility that SPR-5 and MET-2

reprogramming blocks the ectopic expression of MES-4 germline genes by preventing the

accumulation of MES-4 dependent H3K36me3 at a subset of germline genes in somatic

tissues. Consistent with this, most of the MES-4 germline genes were overexpressed in *spr-*

5; met-2 mutants. Using smFISH, we also defined the somatic expression of two ectopically

expressed MES-4 germline genes, *htp-1* and *cpb-1*. While *htp-1* mRNA was ectopically

detected in many somatic tissues, the ectopic expression of *cpb-1* mRNA was more

restricted, suggesting that the extent of ectopic expression is dependent upon the locus.

To directly test whether SPR-5 and MET-2 reprogramming prevents the

accumulation of MES-4 dependent H3K36me3 at germline genes in the somatic tissues of

spr-5; met-2 progeny, we performed H3K36me3 ChIP-seq. MES-4 germline genes

accumulated ectopic H3K36me3 in the soma of *spr-5; met-2* progeny compared to the soma

of N2 progeny. This suggests that without SPR-5 and MET-2 reprogramming, MES-4

ectopically maintains H3K36me3 at these genes in somatic tissues. Importantly, a low level

of H3K36me3 was detected at germline genes in the somatic tissues of N2 progeny. It is

unclear why there is a low level of H3K36me3 normally in somatic tissues in N2 animals.

Nevertheless, this indicates that an increased level of H3K36me3 is necessary to cause

ectopic transcription.

If the developmental delay of *spr-5; met-2* mutant progeny is caused by the ectopic

maintenance of H3K36me3, and the subsequent expression of MES-4 germline genes, then

the ectopic expression of germline genes and the developmental delay should be

dependent upon MES-4. Indeed, we find that the removal of MES-4 rescued both the ectopic

transcription of MES-4 germline genes in the soma of *spr-5; met-2* progeny, and the

developmental delay. Taken together, these results suggest that the developmental delay in

spr-5; met-2 progeny is caused by the ectopic expression of MES-4 germline genes. In

contrast, removing the transcriptionally coupled H3K36me3 methyltransferase, MET-1, did

not rescue the developmental delay. Thus, the developmental delay that we observe in *spr-*

5; met-2 progeny is only dependent on transcriptionally independent H3K36me3.

Our data support the following model for how maternal reprogramming by SPR-5

and MET-2 facilitates proper specification of germline versus soma. In the germline,

transcriptional elongation is blocked by PIE-1, which segregates to germline blastomeres

during embryogenesis (Batchelder et al., 1999; Mello et al., 1992; Seydoux et al., 1996)

(Figure 8A, B). In the soma of the early embryo, there is also very little transcription,

because the bulk of zygotic transcription does not begin until approximately the 60 cell

stage, just prior to when the primordial germ cells Z2 and Z3 are specified (Sulston et al.,

1983). Thus, in *C. elegans,* germline versus soma is largely specified without transcription.

During normal maternal reprogramming, SPR-5 and MET-2 are deposited into the oocyte,

where at fertilization they cooperate to remove H3K4me2 and add H3K9me2 to genes that

were expressed previously in the parental germline (Kerr et al., 2014) (Figure 8A). This

facilitates the transition from active chromatin to repressed chromatin. Genes

epigenetically reprogrammed by SPR-5 and MET-2 include ubiquitously expressed genes

and germline expressed genes, including MES-4 germline genes. The MES-4 germline genes

are subsequently targeted by the transcription independent H3K36 methyltransferase

MES-4 to maintain H3K36me3 in the germ lineage during embryogenesis (Figure 8A, B).

This H3K36 methylation bookmarking antagonizes the repression caused by the erasure of

H3K4me2 and the addition of H3K9me2. Without the transcription independent

maintenance of inherited H3K36 methylation from the mother to antagonize this

repression, the germline fails to proliferate and animals are sterile (Capowski et al., 1991;

Garvin et al., 1998). This is presumably because germline genes that are targeted by MES-4

fail to reactivate, though this has not yet been demonstrated. In contrast, without SPR-5

and MET-2 repression, the failure to add H3K9me2 as well as the inappropriate retention

of H3K4me2 results in a chromatin environment that is permissive for the ectopic

maintenance of H3K36me3 in the soma (Figure 8B). This causes germline genes to be

inappropriately expressed in somatic tissues and results in a severe developmental delay

(Figure 8B).Thus, we propose that SPR-5, MET-2 and MES-4 carefully balance the

inheritance of histone methylation from the parental germline to ensure the proper

specification of germline versus soma in the progeny.

The model that we have proposed makes the following two predictions. First, MES-4

germline genes should normally be targeted for silencing by H3K9me2 in somatic tissues. It

has recently been shown that a subset of germline specific genes contain H3K9me2 at their

promoters in L1 larvae, and that these peaks are lost or reduced in the absence of LIN15B

(Rechtsteiner et al., 2019). We re-examined the H3K9me2 ChIP-seq dataset from this work

and found that the MES-4 germline genes that are ectopically expressed in the somatic

tissues of *spr-5; met-2* progeny also displayed unique H3K9me2 promoter peaks. This

confirms that these genes are normally repressed by H3K9me2 in somatic tissues.

Additionally, the requirement for LIN15B in maintaining somatic H3K9me2 indicates that

the DREAM complex may function to reinforce the expression pattern established by SPR-5

and MET-2. Interestingly, the MEC complex members MEP-1 and LET-418 have also been

implicated in repressing germline genes in somatic tissues (Passannante et al., 2010;

Unhavaithaya et al., 2002). This raises the possibility that the expression pattern

established by SPR-5 and MET-2 may also be reinforced by the MEC complex. The second

prediction from our model is that the developmental delay in *spr-5; met-2* progeny should

be dependent upon the activity of the H3K4 methyltransferase complex. Consistent with

our model, we find that knocking down SET-2, the H3K4me1/2 methyltransferase

component of the COMPASS complex, also rescued the developmental delay that we

observe in *spr-5; met-2* progeny. These findings suggest that the inheritance of ectopic

H3K4 methylation is required for the ectopic accumulation of MES-4 dependent

H3K36me3, and the subsequent ectopic expression of MES-4 germline genes in somatic

tissues of *spr-5; met-2* progeny.

A new paradigm for how an ectopic transcriptional program interferes with

development

Our model explains how H3K36me3 is inappropriately maintained and drives the

ectopic expression of MES-4 germline genes in somatic tissues, but how does the ectopic

expression of germline genes interfere with somatic tissues to cause developmental delay?

One possibility is that the ectopic expression of MES-4 germline genes causes the soma to

take on germline character. To begin to address this, we asked whether *spr-5; met-2* double

mutants can silence an extrachromosomal array in somatic tissues. The silencing of

extrachromosomal arrays is normally restricted to the germline. However, we find that *spr-*

5; met-2 progeny acquired some ability to silence an extrachromosomal multicopy array in

somatic cells. Intriguingly, in our RNA-seq we detected the ectopic expression of RNA-

dependent RNA polymerase genes (e.g. *rrf-1* and *gld-2*) as well as genes involved in the

RNAi effector complex in *spr-5; met-2* mutants (e.g. *hrde-1* and *ppw-1*) Thus, it is possible

that the somatic silencing of the transgene in *spr-5; met-2* mutants is due to the induction of

the germline small RNA pathway. Regardless, the silencing of the extrachromosomal

multicopy array suggests that the somatic tissues in *spr-5; met-2* progeny make functional

proteins that can perform some germline functions. We propose that either the ectopic

germline transcription, or an ectopic germline function resulting from this ectopic

transcription, interferes with the ability of somatic cells to properly read out the correct

somatic transcriptional program. This background noise delays the proper adoption of cell

fate, leading to an overall delay in the development of the tissue. Normally in L1 larvae, the

primordial germ cells Z2 and Z3 are arrested at the G2/M checkpoint (Fukuyama et al.,

2006). In our RNA-seq, we detected the ectopic expression of genes involved in the

negative regulation of proliferation and the cell cycle, as well as G2/M checkpoint genes.

Thus, it is also possible that ectopic expression of germline genes normally expressed only

in Z2 and Z3 contributes to the developmental delay through the ectopic activation of

germline cell cycle control.

Conservation of maternal epigenetic reprogramming between invertebrates and vertebrates

It is possible that the maternal regulation of histone methylation by SPR-5, MET-2 and MES-4 is conserved in mammals. Along with the Heard lab, we previously demonstrated that progeny from mice that maternally lack the vertebrate SPR-5 homolog, KDM1A/LSD1, ectopically express oocyte genes and die at the two-cell stage (Ancelin et al., 2016; Wasson et al., 2016). Similarly, maternal loss of the MET-2 ortholog SEDB1 or the MES-4 ortholog NSD1 in mice results in early embryonic lethality (J. Kim et al., 2016; Rayasam et al., 2003). Together these results underscore the developmental importance of properly regulating histone methylation between generations and raise the possibility that the mechanism we have uncovered is conserved in mammals. The model that we have proposed may also help explain how patients harboring mutations in histone modifying enzymes suffer from certain symptoms. Recent genome sequencing of patients has revealed that several neurodevelopmental disorders are caused by mutations in histone modifying enzymes (extensively reviewed by (J.-H. Kim et al., 2017). These include: several patients with mutations in the human ortholog of *spr-5, Lsd1* (Chong et al., 2016; Tunovic et al., 2014) , as well as Kabuki Syndrome, caused by mutations in the H3K36 methyltransferase SETD2 (Xu et al., 2019), and the H3K27 demethylase KDM6A (Lederer et al., 2012), and Sotos syndrome, caused by mutations in the H3K36 methyltransferase NSD1 (Kurotaki et al., 2002). Similar to what we observed in *spr-5; met-2* mutant progeny, many of these human patients suffer from global developmental delay. Based on our data, it is possible that the developmental delay in these patients may be caused by the failure to

properly regulate histone methylation during critical developmental transitions. This could

result in the ectopic expression of a developmental program in an inappropriate tissue,

leading to background noise and developmental delay.

MATERIALS AND METHODS

Strains. All *Caenorhabditis elegans strains* were grown and maintained at 20° C under

standard conditions, as previously described (Brenner, 1974). The *C. elegans spr-5*

(by101)(I) strain was provided by R. Baumeister. The N2 Bristol wild-type (WT) strain was

provided by the *Caenorhabditis* Genetics Center*. The *met-2 (n4256)(III) strain* was provided

by R. Horvitz. The *hT2 [bli-4(e937)let-?(q782)qIs48] (I;III)* balancer strain was used to

maintain *spr-5 (by101)(I); met-2 (n4256)(III)* double-mutant animals as heterozygotes.

Because the *hT2 [bli-4(e937)let-?(q782)qIs48](I;III)* balancer allele does not extend

completely to the *spr-5* locus on chromosome I, the F0 animals used to generate F1 *spr-5;*

met-2 progeny were cloned out and genotyped to confirm the presence of the *spr-5*

(by101)(I) allele. For genotyping, single animals were picked into 5-10ul of lysis buffer

(50mM KCl, 10mM Tris-HCl (pH 8.3), 2.5mM $MgCl_2$, 0.45% NP-40, 0.45% Tween-20, 0.01%

gelatin) and incubated at 65°C for 1 hour followed by 95°C for 30 minutes. PCR reactions

were performed with AmpliTaq Gold (Invitrogen) according to the manufacturer's protocol

and reactions were resolved on agarose gels (see Table S4 for primer sequences). Before

completing this study we acquired the FX30208 *tmC27 [unc-75(tmIs1239)]](I)* from the

Caenorhabditis Genetics Center that completely covers the *spr-5* locus on chromosome I.

The *qC1 [qIs26 (lag2::GFP + rol-6(su1006)](III)* strain was obtained from W. Kelly and

crossed to *met-2 (n4256)(III)* to maintain *met-2(n4256)(III)* as heterozygotes. The *spr-5*

(by101)(I)/tmC27[unc-75(tmls1239)](I); met-2 (n4256) (III)/qC1 [qls26 (lag2::gfp+ rol-

6(su1006))](III) strain was then re-created for this study to maintain *spr-5 (by101)(I);*

met-2 (n4256)(III) double-mutant animals as balanced heterozygotes. The LET-858::GFP

(*pha-1*(e2123ts)(III); *let-858::gfp* (ccEx7271)) (Kelly and Fire, 1998) transgenic strain used

in somatic transgene silencing assays was acquired from W. Kelly.

Scoring developmental delay. *C. elegans* adult hermaphrodites were allowed to lay

embryos for 2-4 hours and then removed in order to synchronize the development of

progeny. Progeny were then imaged and scored for development to the adult stage at

either 72 hours or seven days after synchronized lay, depending on the experiment.

RNA sequencing and analysis. Total RNA was isolated using TRIzol reagent (Invitrogen)

from 200-250 starved L1 larvae born at room temperature (21°C - 22°C) overnight in M9

Buffer. Due to difficulty in isolating large numbers of *spr-5; met-2* double-mutant progeny

from the *hT2 [bli-4(e937)let-?(q782)qls48](I;III)* balancer strain, we submitted total RNA to

the Genomic Services Laboratory (GSL) (HudsonAlpha, Huntsville, Alabama) for low input

RNA-seq services. This service utilizes the Ovation RNA-Seq System V2 kit (Nugen) for

initial RNA amplification prior to library preparation and sequencing (Illumina HiSeq v4,

50bp paired-end reads). For each genotype, 2 biological replicates were obtained.

Sequencing reads were checked for quality using FastQC (Wingett and Andrews, 2018),

filtered using Trimmomatic (Bolger et al., 2014), and remapped to the *C. elegans*

transcriptome (ce10, WS220) using HISAT2 (D. Kim et al., 2015). Read count by gene was

obtained by FeatureCounts (Liao et al., 2014) . Differentially expressed transcripts

(significance threshold p-value < 0.05) were determined using DESEQ2 (v.2.11.40.2) (Love

et al., 2014). Transcripts per million (TPM) values were calculated from raw data obtained

from FeatureCounts output. Subsequent downstream analysis was performed using R with

normalized counts and p-values from DESEQ2 (v.2.11.40.2). Heatmaps were produced

using the ComplexHeatmap R Package (Gu et al., 2016). Data was scaled and hierarchical

clustering was performed using the complete linkage algorithm. In the linkage algorithm,

distance was measured by calculating pairwise distance.   Volcano plots were produced

using the EnhancedVolcano package (v.0.99.16). Additionally, Gene Ontology (GO) Pathway

analysis was performed using the online platform WormEnrichr (Chen et al., 2013;

Kuleshov et al., 2016). Custom R-scripts are available upon request. An additional heatmap

comparison of differentially expressed genes between *spr-5*, *met-2*, and *spr-5; met-2*

progeny compared to N2 progeny was generated in Microsoft Excel using Log2 fold change

values from the DESEQ2 analysis. Because transcript isoforms were ignored, we discuss the

data in terms of "genes expressed" rather than "transcripts expressed".

Chromatin immunoprecipitation sequencing and analysis. Chromatin

immunoprecipitation (ChIP) experiments were performed as described by Katz and

colleagues (Katz et al., 2009). Briefly, 600 starved L1 larvae born at room temperature

(21°C - 22°C) overnight in M9 Buffer were collected, frozen in liquid nitrogen, and stored at

-80°C prior to homogenization. Frozen pellets were disrupted by a glass Dounce

homogenizer, fixed with formaldehyde (1% final concentration), and quenched with

glycine. ChIP samples were processed with a Chromatin Immunoprecipitation Assay Kit

(Millipore), according to manufacturer's instructions. Samples were sonicated using a

Diagenode Bioruptor UCD-200 at 4°C on the "high" setting for a total of 30min with a cycle

of 45sec on and 15sec off. A total of 12.5µL (5 ug) H3K36me3 antibody (cat. 61021; Active

motif) was used for immunoprecipitation. The Genomic Services Laboratory (GSL)

(HudsonAlpha, Huntsville, Alabama) performed library preparation and sequencing

(Illumina HiSeq v4, 50bp single-end reads). Reads were checked for quality using FastQC

(Wingett and Andrews, 2018) and remapped to the *C. elegans* transcriptome (ce10, WS220)

using Bowtie2 (Langmead and Salzberg, 2012; Langmead et al., 2009) under default

parameters. bamCoverage in deepTools2 (Ramírez et al., 2016) was used to generate

bigwig coverage tracks in 50bp bins, with blacklisted regions from McMurchy et al. 2017

excluded, using the following parameters: -bs 50, --normalizeUsing RPKM (McMurchy et al.,

2017).

MACS2 (Feng et al., 2012; Zhang et al., 2008) default parameters were used to call

peaks and create bedgraph files for sequenced and mapped H3K36me3 ChIP samples and

input DNA samples with the following adjustments to account for H3K36me3 broader

domains: Broad-cutoff =0.001. Blacklisted regions from McMurchy et al. 2017 were

excluded for this analysis (McMurchy et al., 2017). Using published H3K36me3 modMine

Chip-chip called broad peaks (modENCODE_3555) from N2 L1 larvae as a guide, we then

merged called broad peaks within 1200bp using Bedtools: MergeBED (Quinlan and Hall,

2010). H3K9me2 bedgraph files used in our analysis were from a published dataset

(Rechtsteiner et al., 2019). Integrative Genome Viewer (IGV) was used to visualize

H3K36me3 reads normalized to reads per kilobase millions (RPKM) and H3K9me2 reads

normalized to 15 million reads (genome wide coverage of H3K9me2; Rechtsteiner et al.,

2019).

RNAi methods. RNAi by feeding was carried out using clones from the Ahringer library

(Kamath and Ahringer, 2003). Feeding experiments were performed on RNAi plates (NGM

plates containing 100 ug/ml ampicillin, 0.4mM IPTG, and 12.5ug/ml tetracycline). F0

worms were placed on RNAi plates as L2 larvae and then moved to fresh RNAi plates 48hrs

later where they were allowed to lay embryos for 2-4 hrs. F0 worms were then removed

from plates and sacrificed or placed in M9 buffer overnight so that starved L1 progeny

could be isolated for quantitative PCR (qPCR). F1 progeny were scored 72hrs after the

synchronized lay for developmental progression. For each RNAi experiment, *pos-1* RNAi

was used as a positive control. Each RNAi experiment reported here *pos-1* RNAi resulted in

>95% embryonic lethality, indicating that RNAi plates were optimal.

Real-time expression analysis. Total RNA was isolated using TRIzol reagent (Invitrogen)

from synchronized L1s born at room temperature (21°C – 22°C). cDNA synthesis and qPCR

were carried out as described (Kerr et al. 2014). All experiments were performed in

triplicate and normalized to *ama-1* mRNA expression (see Table S4 for primer sequences).

Differential interference contrast microscopy. Worms were immobilized in 0.1%

levamisole and placed on a 2% agarose pad for imaging at either 10x or 40x magnification.

40x DIC images were overlaid together using Adobe photoshop to generate high resolution

images of whole worms.

Single Molecule Fluorescent *in situ* Hybridization (smFISH). Quasar 570 labeled

smFISH probe sets for *htp-1* and *cpb-1* were designed using Stellaris Probe Designer

(Biosearch) (see Table S5 for probe sequences). The *htp-1* smFISH probes were designed

using the complete 1,059nt *htp-1* protein-coding sequence. Likewise, The *cpb-1* smFISH

probes were designed using the complete 1,683nt *cpb-1* protein-coding sequence. In

addition, an smFISH fluorescent probe set for *ama-1* was purchased from the DesignReady

catalog (cat#: VSMF-6002-5, Biosearch).

Synchronized L1 larvae for smFISH were obtained by bleaching 300-500 gravid

hermaphrodites and allowing embryos to hatch overnight on 6cm NGM plates lightly

seeded OP50 bacteria. L1 larvae were then washed into 1.5ul Eppendorf tubes using

nuclease-free M9 buffer. Fixation and hybridization steps followed the Stellaris RNA FISH

protocol for *C. elegans* adapted from RAJ lab protocol (Raj and Tyagi, 2010). In brief, we

resuspended L1 larvae in fixation buffer (3.7% formaldehyde in 1 x PBS) for 15 minutes at

room temperature then transferred tubes to liquid nitrogen. Samples were thawed in water

and placed on ice for 20 minutes. In our hands, we obtain better fluorescent signal by

freeze cracking L1 larvae. Following fixation, L1 Larvae were resuspended in 70% EtOH

and stored at 4°C for 24-48 hours. For all probe sets, we incubated L1 larvae in 100ul

hybridization buffer (containing 10% formamide and 125nm probe) for 4 hours at 37°C.

After hybridization, samples were washed in wash buffer at 37°C for 30 minutes, incubated

in 50ng/mL DAPI in wash buffer at 30°C for 30 minutes, washed once in 2x SSC for 2

minutes at room temperature, and mounted in Vectashield mounting medium. Mounted

slides were imaged immediately using a 100x objective on a spinning-disk confocal Nikon-

TiE imaging system. Images were captured using NIS-Elements software (Nikon) and

ImageJ (NIH, http://imagej.nih.gov/ij/) was used for viewing. ImageJ maximum projection

was used to project z-stack images to a single plane. The fluorescent intensity of smFISH

dots were > 2-fold above background as expected (Ji and van Oudenaarden, 2012).

LET-858::GFP transgene silencing assay. First generation *spr-5; met-2* hermaphrodites

were crossed to *let-858* transgenic males to generate *spr-5/+*; *met-2/+*; *let-858::gfp* animals.

The let-858 transgene is an extrachromosomal multicopy *let-858* array (Kelly and Fire,

1998). From these animals we generated *spr-5; met-2; let-858::gfp* animals and scored them

for somatic expression of LET-858::GFP using a standard stereoscope. L2 progeny from N2

and *spr-5; met-2* progeny expressing the LET-858::GFP transgene were scored as "bright"

(High level ubiquitous expression) , faint (barely visible and ubiquitous expression), or off

(no expression). Because less than 50% of progeny inherit the *let-858* transgene, we

normalized the percentage of progeny scored as "off" for LET-858::GFP to presence of the

let-858 transgene based on genotyping for *gfp.* For N2, 0 out of 60 progeny that were

scored as "off" failed to inherit the *let-858* transgene indicating that the transgene is never

silenced in N2 progeny (data not shown, see Table S4 for primer sequences). For *spr-5;*

met-2 progeny, 8 out of 50 progeny that were scored as "off" inherited the *let-858*

transgene indicating that the *let-858* transgene is completely silenced in some *spr-5; met-2*

progeny (data not shown).

ACKNOWLEDGEMENTS

We are grateful to members of the Katz lab, as well as T. Caspary and C. Bean, for their

helpful discussion and critical reading of the manuscript; M. J. Rowley for advice on

bioinformatics; S. Strome and L. Petrella for helpful discussion and experimental advice; D.

Lerit, R. Deal, V. Corces, and W. Kelly for the generous sharing of reagents and equipment.

We thank R. Horvitz, W. Kelly, and the *Caenorhabditis* Genetics Center (funded by NIH P40

OD010440) for strains. **Funding:** This work was funded by a grant to D.J.K. (NSF

IOS1931697); T.W.L. and B.S.C. were supported by the Fellowships in Research and Science

Teaching IRACDA postdoctoral program (NIH K12GM00680-15); and B.S.C. was also

supported by NIH F32 GM126734-01. **Author Contributions:** B.S.C. and D.J.K conceived

and designed the study and wrote the manuscript. B.S.C. performed experiments under the

direction of D.J.K. B.S.C and D.J.K. analyzed data and interpreted results, with help from

T.W.L. J.S.B. initially performed *set-2* RNAi rescue experiment and C.F.P. initially performed

the *let-858* transgene silencing experiment. D.A.M. helped with RNA-seq analysis. All

authors discussed the results. **Data availability**: Raw and processed genomic data has

been deposited with the Gene Expression Omnibus (www.ncbi.nlm.nih.gov/geo) under

accession code GSE143839.

Supplementary Material

Figure supplements:

Figure 1: Supplemental Figure 1

Figure 2: Supplemental Figures 2 and 3

Figure 3: Supplemental Figure 4

Figure 4: Supplemental Figures 5 and 6

Supplementary tables 1-5

REFERENCES

Ancelin, K., Syx, L., Borensztein, M., Ranisavljevic, N., Vassilev, I., Briseño-Roa, L., Liu, T.,
Metzger, E., Servant, N., Barillot, E., Chen, C.-J., Schüle, R., Heard, E., 2016. Maternal
LSD1/KDM1A is an essential regulator of chromatin and transcription landscapes
during zygotic genome activation. Elife 5, 4615. doi:10.7554/eLife.08851
Batchelder, C., Dunn, M.A., Choy, B., Suh, Y., Cassie, C., Shim, E.Y., Shin, T.H., Mello, C.,
Seydoux, G., Blackwell, T.K., 1999. Transcriptional repression by the Caenorhabditis
elegans germ-line protein PIE-1. Genes Dev. 13, 202–212. doi:10.1101/gad.13.2.202
Bolger, A.M., Lohse, M., Usadel, B., 2014. Trimmomatic: a flexible trimmer for Illumina
sequence data. Bioinformatics 30, 2114–2120. doi:10.1093/bioinformatics/btu170
Brenner, S., 1974. The genetics of Caenorhabditis elegans. Genetics 77, 71–94.
Capowski, E.E., Martin, P., Garvin, C., Strome, S., 1991. Identification of grandchildless loci
whose products are required for normal germ-line development in the nematode
Caenorhabditis elegans. Genetics 129, 1061–1072.

Chen, E.Y., Tan, C.M., Kou, Y., Duan, Q., Wang, Z., Meirelles, G.V., Clark, N.R., Ma'ayan, A., 2013. Enrichr: interactive and collaborative HTML5 gene list enrichment analysis tool. BMC Bioinformatics 14, 128. doi:10.1186/1471-2105-14-128

Chong, J.X., Yu, J.-H., Lorentzen, P., Park, K.M., Jamal, S.M., Tabor, H.K., Rauch, A., Saenz, M.S., Boltshauser, E., Patterson, K.E., Nickerson, D.A., Bamshad, M.J., 2016. Gene discovery for Mendelian conditions via social networking: de novo variants in KDM1A cause developmental delay and distinctive facial features. Genet. Med. 18, 788–795. doi:10.1038/gim.2015.161

Feng, J., Liu, T., Qin, B., Zhang, Y., Liu, X.S., 2012. Identifying ChIP-seq enrichment using MACS. Nat Protoc 7, 1728–1740. doi:10.1038/nprot.2012.101

Fukuyama, M., Rougvie, A.E., Rothman, J.H., 2006. C. elegans DAF-18/PTEN mediates nutrient-dependent arrest of cell cycle and growth in the germline. Curr. Biol. 16, 773–779. doi:10.1016/j.cub.2006.02.073

Furuhashi, H., Takasaki, T., Rechtsteiner, A., Li, T., Kimura, H., Checchi, P.M., Strome, S., Kelly, W.G., 2010. Trans-generational epigenetic regulation of C. elegans primordial germ cells. Epigenetics Chromatin 3, 15. doi:10.1186/1756-8935-3-15

Garvin, C., Holdeman, R., Strome, S., 1998. The phenotype of mes-2, mes-3, mes-4 and mes-6, maternal-effect genes required for survival of the germline in Caenorhabditis elegans, is sensitive to chromosome dosage. Genetics 148, 167–185.

Gu, Z., Eils, R., Schlesner, M., 2016. Complex heatmaps reveal patterns and correlations in multidimensional genomic data. Bioinformatics 32, 2847–2849. doi:10.1093/bioinformatics/btw313

Jambhekar, A., Dhall, A., Shi, Y., 2019. Roles and regulation of histone methylation in animal development. Nat. Rev. Mol. Cell Biol. 20, 625–641. doi:10.1038/s41580-019-0151-1

Ji, N., van Oudenaarden, A., 2012. Single molecule fluorescent in situ hybridization (smFISH) of C. elegans worms and embryos. WormBook 1–16. doi:10.1895/wormbook.1.153.1

Kamath, R.S., Ahringer, J., 2003. Genome-wide RNAi screening in Caenorhabditis elegans. Methods 30, 313–321. doi:10.1016/s1046-2023(03)00050-1

Katz, D.J., Edwards, T.M., Reinke, V., Kelly, W.G., 2009. A C. elegans LSD1 demethylase contributes to germline immortality by reprogramming epigenetic memory. Cell 137, 308–320. doi:10.1016/j.cell.2009.02.015

Kelly, W.G., Fire, A., 1998. Chromatin silencing and the maintenance of a functional germline in Caenorhabditis elegans. Development 125, 2451–2456.

Kerr, S.C., Ruppersburg, C.C., Francis, J.W., Katz, D.J., 2014. SPR-5 and MET-2 function cooperatively to reestablish an epigenetic ground state during passage through the germ line. Proc. Natl. Acad. Sci. U.S.A. 111, 9509–9514. doi:10.1073/pnas.1321843111

Kim, D., Langmead, B., Salzberg, S.L., 2015. HISAT: a fast spliced aligner with low memory requirements. Nat. Methods 12, 357–360. doi:10.1038/nmeth.3317

Kim, J., Zhao, H., Dan, J., Kim, S., Hardikar, S., Hollowell, D., Lin, K., Lu, Y., Takata, Y., Shen, J., Chen, T., 2016. Maternal Setdb1 Is Required for Meiotic Progression and Preimplantation Development in Mouse. PLoS Genet. 12, e1005970. doi:10.1371/journal.pgen.1005970

Kim, J.-H., Lee, J.H., Lee, I.-S., Lee, S.B., Cho, K.S., 2017. Histone Lysine Methylation and Neurodevelopmental Disorders. Int J Mol Sci 18, 1404. doi:10.3390/ijms18071404

Kreher, J., Takasaki, T., Cockrum, C., Sidoli, S., Garcia, B.A., Jensen, O.N., Strome, S., 2018.
Distinct Roles of Two Histone Methyltransferases in Transmitting H3K36me3-Based
Epigenetic Memory Across Generations in Caenorhabditis elegans. Genetics 210, 969–
982. doi:10.1534/genetics.118.301353

Kuleshov, M.V., Jones, M.R., Rouillard, A.D., Fernandez, N.F., Duan, Q., Wang, Z., Koplev, S.,
Jenkins, S.L., Jagodnik, K.M., Lachmann, A., McDermott, M.G., Monteiro, C.D., Gundersen,
G.W., Ma'ayan, A., 2016. Enrichr: a comprehensive gene set enrichment analysis web
server 2016 update. Nucleic Acids Res. 44, W90–7. doi:10.1093/nar/gkw377

Kurotaki, N., Imaizumi, K., Harada, N., Masuno, M., Kondoh, T., Nagai, T., Ohashi, H.,
Naritomi, K., Tsukahara, M., Makita, Y., Sugimoto, T., Sonoda, T., Hasegawa, T., Chinen,
Y., Tomita Ha, H.-A., Kinoshita, A., Mizuguchi, T., Yoshiura Ki, K.-I., Ohta, T., Kishino, T.,
Fukushima, Y., Niikawa, N., Matsumoto, N., 2002. Haploinsufficiency of NSD1 causes
Sotos syndrome. Nat. Genet. 30, 365–366. doi:10.1038/ng863

Langmead, B., Salzberg, S.L., 2012. Fast gapped-read alignment with Bowtie 2. Nat. Methods
9, 357–359. doi:10.1038/nmeth.1923

Langmead, B., Trapnell, C., Pop, M., Salzberg, S.L., 2009. Ultrafast and memory-efficient
alignment of short DNA sequences to the human genome. Genome Biol. 10, R25.
doi:10.1186/gb-2009-10-3-r25

Lederer, D., Grisart, B., Digilio, M.C., Benoit, V., Crespin, M., Ghariani, S.C., Maystadt, I.,
Dallapiccola, B., Verellen-Dumoulin, C., 2012. Deletion of KDM6A, a histone
demethylase interacting with MLL2, in three patients with Kabuki syndrome. Am. J.
Hum. Genet. 90, 119–124. doi:10.1016/j.ajhg.2011.11.021

Liao, Y., Smyth, G.K., Shi, W., 2014. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. Bioinformatics 30, 923–930.
doi:10.1093/bioinformatics/btt656

Love, M.I., Huber, W., Anders, S., 2014. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. Genome Biol. 15, 550. doi:10.1186/s13059-014-0550-8

McMurchy, A.N., Stempor, P., Gaarenstroom, T., Wysolmerski, B., Dong, Y., Aussianikava, D.,
Appert, A., Huang, N., Kolasinska-Zwierz, P., Sapetschnig, A., Miska, E.A., Ahringer, J.,
2017. A team of heterochromatin factors collaborates with small RNA pathways to
combat repetitive elements and germline stress. Elife 6, 7931. doi:10.7554/eLife.21666

Mello, C.C., Draper, B.W., Krause, M., Weintraub, H., Priess, J.R., 1992. The pie-1 and mex-1
genes and maternal control of blastomere identity in early C. elegans embryos. Cell 70,
163–176. doi:10.1016/0092-8674(92)90542-k

Passannante, M., Marti, C.-O., Pfefferli, C., Moroni, P.S., Kaeser-Pebernard, S., Puoti, A.,
Hunziker, P., Wicky, C., Müller, F., 2010. Different Mi-2 complexes for various
developmental functions in Caenorhabditis elegans. PLoS ONE 5, e13681.
doi:10.1371/journal.pone.0013681

Quinlan, A.R., Hall, I.M., 2010. BEDTools: a flexible suite of utilities for comparing genomic
features. Bioinformatics 26, 841–842. doi:10.1093/bioinformatics/btq033

Raj, A., Tyagi, S., 2010. Detection of individual endogenous RNA transcripts in situ using
multiple singly labeled probes. Meth. Enzymol. 472, 365–386. doi:10.1016/S0076-
6879(10)72004-8

Ramírez, F., Ryan, D.P., Grüning, B., Bhardwaj, V., Kilpert, F., Richter, A.S., Heyne, S., Dündar,
F., Manke, T., 2016. deepTools2: a next generation web server for deep-sequencing data
analysis. Nucleic Acids Res. 44, W160–5. doi:10.1093/nar/gkw257

Rayasam, G.V., Wendling, O., Angrand, P.-O., Mark, M., Niederreither, K., Song, L., Lerouge, T.,
Hager, G.L., Chambon, P., Losson, R., 2003. NSD1 is essential for early post-implantation
development and has a catalytically active SET domain. EMBO J. 22, 3153–3163.
doi:10.1093/emboj/cdg288

Rechtsteiner, A., Costello, M.E., Egelhofer, T.A., Garrigues, J.M., Strome, S., Petrella, L.N.,
2019. Repression of Germline Genes in Caenorhabditis elegans Somatic Tissues by
H3K9 Dimethylation of Their Promoters. Genetics 212, 125–140.
doi:10.1534/genetics.118.301878

Rechtsteiner, A., Ercan, S., Takasaki, T., Phippen, T.M., Egelhofer, T.A., Wang, W., Kimura, H.,
Lieb, J.D., Strome, S., 2010. The histone H3K36 methyltransferase MES-4 acts
epigenetically to transmit the memory of germline gene expression to progeny. PLoS
Genet. 6, e1001091. doi:10.1371/journal.pgen.1001091

Seydoux, G., Mello, C.C., Pettitt, J., Wood, W.B., Priess, J.R., Fire, A., 1996. Repression of gene
expression in the embryonic germ lineage of C. elegans. Nature 382, 713–716.
doi:10.1038/382713a0

Shi, Y., Lan, F., Matson, C., Mulligan, P., Whetstine, J.R., Cole, P.A., Casero, R.A., Shi, Y., 2004.
Histone demethylation mediated by the nuclear amine oxidase homolog LSD1. Cell 119,
941–953. doi:10.1016/j.cell.2004.12.012

Shi, Y.-J., Matson, C., Lan, F., Iwase, S., Baba, T., Shi, Y., 2005. Regulation of LSD1 histone
demethylase activity by its associated factors. Mol. Cell 19, 857–864.
doi:10.1016/j.molcel.2005.08.027

Strome, S., 2005. Specification of the germ line. WormBook 1–10.
doi:10.1895/wormbook.1.9.1

Sulston, J.E., Schierenberg, E., White, J.G., Thomson, J.N., 1983. The embryonic cell lineage of
the nematode Caenorhabditis elegans. Dev. Biol. 100, 64–119. doi:10.1016/0012-
1606(83)90201-4

Tunovic, S., Barkovich, J., Sherr, E.H., Slavotinek, A.M., 2014. De novo ANKRD11 and KDM1A
gene mutations in a male with features of KBG syndrome and Kabuki syndrome. Am. J.
Med. Genet. A 164A, 1744–1749. doi:10.1002/ajmg.a.36450

Unhavaithaya, Y., Shin, T.H., Miliaras, N., Lee, J., Oyama, T., Mello, C.C., 2002. MEP-1 and a
homolog of the NURD complex component Mi-2 act together to maintain germline-
soma distinctions in C. elegans. Cell 111, 991–1002.

Wasson, J.A., Simon, A.K., Myrick, D.A., Wolf, G., Driscoll, S., Pfaff, S.L., Macfarlan, T.S., Katz,
D.J., 2016. Maternally provided LSD1/KDM1A enables the maternal-to-zygotic
transition and prevents defects that manifest postnatally. Elife 5, 296.
doi:10.7554/eLife.08848

Wingett, S.W., Andrews, S., 2018. FastQ Screen: A tool for multi-genome mapping and
quality control. F1000Res 7, 1338. doi:10.12688/f1000research.15931.2

Xu, Q., Xiang, Y., Wang, Q., Wang, L., Brind'Amour, J., Bogutz, A.B., Zhang, Y., Zhang, B., Yu, G.,
Xia, W., Du, Z., Huang, C., Ma, J., Zheng, H., Li, Y., Liu, C., Walker, C.L., Jonasch, E.,
Lefebvre, L., Wu, M., Lorincz, M.C., Li, W., Li, L., Xie, W., 2019. SETD2 regulates the
maternal epigenome, genomic imprinting and embryonic development. Nat. Genet. 51,
844–856. doi:10.1038/s41588-019-0398-7

Zhang, Y., Liu, T., Meyer, C.A., Eeckhoute, J., Johnson, D.S., Bernstein, B.E., Nusbaum, C.,
Myers, R.M., Brown, M., Li, W., Liu, X.S., 2008. Model-based analysis of ChIP-Seq (MACS).
Genome Biol. 9, R137. doi:10.1186/gb-2008-9-9-r137

Figure 1. *spr-5; met-2* mutants display severe developmental delay. 10x Differential Interference Contrast (DIC) images of N2 (A), *spr-5* (B*)*, *met-2* (C), and *spr-5; met-2* progeny (D) 72 hours post synchronized lay. Scale bar: 100μm. (E) Percentage of N2, *spr-5*, *met-2*, and *spr-5; met-2* progeny that reached the adult stage (% Adult Progeny) by 72 hours post synchronized lay. Error bars represent the standard deviation of the mean from three experiments.

Figure 2. MES-4 germline genes are ectopically expressed in *spr-5; met-2* mutant soma. Overlap between MES-4 germline genes and genes up-regulated (A) and down-regulated (B) in *spr-5; met-2* L1 progeny. Significant over-enrichment in A was determined by the hypergeometric test (*P-value < 4.34e-20). (C) Heatmap of log2 fold change (FC) of 176 MES-4 germline genes in *spr-5, met-2* and *spr-5; met-2* mutants compared to N2. log2FC values are represented in a yellow to blue gradient and range from -2 to 5. Yellow represents genes with negative log2FC values and blue represents genes with positive log2FC values compared to N2. The remaining 21 MES-4 germline genes were not included because they do not have an expression value in one or more of the data sets (*spr-5, met-2,* or *spr-5; met-2*)

Figure 3. *spr-5; met-2* L1 progeny ectopically express MES-4 germline genes in multiple somatic tissues. 40x smFISH images of *htp-1* (A, C, D, F) and *cpb-1* (G, I, J, L) endogenous mRNAs in N2 (A-C, G-I) and *spr-5; met-2* (D-F, J-L) L1 progeny. DAPI was used as a nuclear marker (B, C, E, F, H, I, K, L). Insets are high magnification images of the germ cells, Z2 and Z3, in N2 L1 progeny. Arrows (J, L) denote ectopic *cpb-1* mRNA foci in somatic cells. Scale bar 40μm.

Figure 4. MES-4 germline genes display ectopic H3K36me3 in *spr-5; met-2* L1 progeny. Heatmap of H3K36me3 ChIP-seq reads normalized to reads per kilobase million (RPKM) over the gene bodies of 196 MES-4 germline genes in wild-type (N2) (A) versus *spr-5; met-2* (B) L1 progeny (second replicate in Figure S5). Gene bodies were pseudoscaled to 1kb with 500bp borders separated by orange bars that represent the transcriptional start site (TSS) and transcriptional end site (TES). Integrative Genome Viewer (IGV) image of H3K36me3 ChIP-seq reads normalized to RPKM at MES-4 germline genes (C-F, J-M) and control genes (G-I, N-P) in N2 (C-I) versus *spr-5; met-2* (J-P) L1 progeny. RPKM IGV windows were scaled between 0 and 202 RPKM for all genes.

Figure 5. Knocking down MES-4 rescues ectopic expression of MES-4 germline genes in *spr-5; met-2* L1 progeny. Quantitative RT-PCR showing the relative units of expression for nine MES-4 germline genes (*cpb-1, csr-1, fbxa-101*, *htp*-1, *T05B9.1, Y18D10A.11, ftr-1, rmh-1, K12D12.5*) in L1 progeny of *spr-5; met-2* hermaphrodites fed either control L4440 RNAi (grey bars) or *mes-4* RNAi (black bars) versus N2 fed control L4440 RNAi (white bars). Relative units of expression from two biological replicates (R1 and R2) were calculated for each gene by averaging triplicate RT_PCR reactions and normalizing to a control gene, *ama-1*. Error bars represent the standard error of the mean (SEM) for triplicate RT_PCR reactions. For all nine genes, *mes-4* RNAi significantly reduced the relative expression of *spr-5; met-2* compared to *spr-5; met-2* fed L4440 control RNAi (t-test, p-value <0.001).

Figure 6. Knocking down MES-4 rescues developmental delay in *spr-5; met-2* progeny. DIC images of N2 (A-D) or *spr-5; met-2* (E-H) progeny from hermaphrodite parents treated with control (L4440 vector only) RNAi (A, E), *mes-4* RNAi (B, F), *met-1*RNAi (C, G), or *set-2* RNAi (D, H) 72 hours post synchronized lay. Scale bar: 100μm. (I) Quantification of the number of progeny (represented as % Adult Progeny) from A-H that made it to adults by 72 hours. Error bars represent the standard deviation of the mean from two or three experiments. N represents the total number of progeny scored across independent experiments. (t-test, **** represent a p-value <0.0001).

Figure 7. *let-858* transgene silencing in the soma of *spr-5; met-2* mutants. 40x differential contrasting interference (DIC) (A, C, E, G, and I) and immunofluorescent (B, D, F, H, and J) images of N2 (A-D) and *spr-5; met-2* (E-J) L2 progeny. Arrows denote faint expression of LET-858::GFP. Scale bar: 50 μm. Scale bar is the same for all panels. Percentage of animals where the expression level of LET-858::GFP was scored as either bright expressing (bright green, representative shown in panel B and F), faint expressing (dark green, representative shown in panel D and H), or not expressing (black, representative shown in panel J) in N2 (N=132) versus *spr-5; met-2* (N=87) progeny (K). The quantification represents the percentages of LET-858::GFP expressing progeny from two independent experiments. To control for the segregation of the *let-858* transgene, progeny scored as "off" were normalized for the presence of the *let-858* transgene in animals as detected by PCR (see methods).

Figure 8: A model for how maternal reprogramming of inherited histone methylation helps to specify germline versus soma. During development, the COMPASS Complex and MET-1 add transcriptionally coupled H3K4me1/2 and H3K36me2/3 to germline expressed genes in the parental germline. (A) At fertilization, these germline expressed genes undergo maternal epigenetic reprogramming by SPR-5 and MET-2 to remove H3K4me1/2 and add H3K9me1/2. In the germline blastomeres of the embryo, PIE-1 prevents global transcription by inhibiting Pol II. In the absence of transcription, MES-4 maintains H3K36me2/3 at MES-4 germline genes that have acquired transcriptionally coupled H3K36me2/3 in the previous germline. This ensures that these genes remain bookmarked for re-expression once the germline begins to proliferate later in development. (B) Without SPR-5 and MET-2 maternal reprogramming, H3K4me1/2 is inappropriately inherited in somatic tissues, allowing MES-4 to ectopically add H3K36me2/3 at these germline genes. This leads to ectopic expression of MES-4 germline genes in somatic tissues and a severe developmental delay. Orange circles represent somatic cells, Grey circles outlined in blue dashed-lines represent germ cells, and orange circles outlined in dashed-blue lines depict somatic cells that ectopically express MES-4 germline genes. P-lineage, germline blastomeres, are indicated by the letter P, and the primordial germline cells are indicated by Z2 and Z3.

Supplemental Figure 1. *spr-5; met-2* mutants display severe developmental delay and protruding vulva. 40x DIC images of N2 (A), *spr-5* (B), *met-2* (C), and *spr-5; met-2* (D) progeny at 72 hours post synchronized lay. DIC image in (D) was magnified an additional 4x. 40x DIC images of two examples of *spr-5; met-2* adult progeny (E, F) at seven days post synchronized lay. Dashed-line (D) outlines germline and asterisks (E, F) denote protruding vulva. Scale bar: 100μm. (G) Quantification of *spr-5; met-2* progeny that reached the adult stage by seven days post synchronized lay, along with quantification of protruding vulva (Pvl) in these animals that reached the adult stage. The 876 progeny scored came from a total of 25 hermaphrodites across two independent experiments.

G

Total # of progeny scored	Total # of adult progeny	Percent of adult progeny	Total # of adult progeny with Pvl	Percent of adult progeny with Pvl
876	35	3.9%	31	88.5%

Supplemental Figure 2: Differential gene expression in *spr-5, met-2, and spr-5; met-2* progeny *compared* to N2 progeny. (A) Table summary of differentially expressed genes in *spr-5, met-2, and spr-5; met-2* L1 progeny from DeSEQ2 analysis (significance cut-off of p< 0.05). (B) Overlap of differentially expressed genes between *spr-5, met-2, and spr-5; met-2* L1 progeny. Gene Ontology analysis showing Biological Processes (C) and Cellular Components (D) amongst genes that were up-regulated (red) and down-regulated (green) in *spr-5; met-2* L1 progeny compared to N2. Combined Score was computed to determine gene set enrichment (Chen et al., 2013).

A

Genotype	Total # of differentially expressed genes	Up-regulated > 2 fold log2 FC	Down-regulated < 2 fold log2 FC
spr-5	343	142	41
met-2	413	146	67
spr-5; met-2	778	476	294

B

C

D

Supplemental Figure 3. Differential expression and replicate comparison of RNAseq experiments performed on N2, *spr-5*; *met-2*, and *spr-5*; *met-2* L1 progeny. Volcano plot of log2 fold changes in gene expression (x-axis) by statistical significance (-Log$_{10}$ P-value; y-axis) in *spr-5* (A), *met-2* (B), and *spr-5*; *met-2* (C) L1 Progeny compared to N2. Heatmap of differentially expressed RNA-seq transcripts between N2 and *spr-5* (D), *met-2* (E), and *spr-5*; *met-2* (F). Data was scaled and hierarchical clustering was performed using the complete linkage algorithm. Instead of Euclidean distance, distance was measured by calculating pairwise distance. Higher (red) and lower (green) expression is reported as a z-score.

Supplemental Figure 4. smFISH control experiments. smFISH images of *ama-1* mRNA (A, C, D, F) and a no probe control (G, I, J, L) in N2 (A-C, G-I) and *spr-5; met-2* (D-F, J-L) L1 progeny. DAPI was used as a nuclear marker (B, C, E, F, H, I, K, L). Scale bar 40µm.

Supplemental Figure 5: Replicate comparison of H3K36me3 ChIP-seq analysis in *spr-5; met-2* L1 progeny. Heatmap of H3K36me3 ChIP-seq reads from a second replicate (1st replicate in Figure 4) normalized to reads per kilobase million (RPKM) over the gene bodies of 196 MES-4 germline genes in wild-type (N2) (A) versus *spr-5; met-2* (B) L1 progeny. Gene bodies were pseudoscaled to 1kb with 500bp borders separated by orange bars that represent the transcriptional start site (TSS) and transcriptional end site (TES). Integrative Genome Viewer (IGV) image of H3K36me3 ChIP-seq reads from replicate 2 normalized to RPKM at MES-4 germline genes (C-F, J-M) versus control genes (G-I, N-P) in N2 (C-I) and *spr-5; met-2* (J-P) L1 progeny. RPKM IGV windows were scaled between 0 and 107 RPKM for all genes. Venn-diagram displaying the overlap of H3K36me3 ChIP-seq called broad peaks for N2 (Q) and *spr-5; met-2* (R) replicates.

Supplemental Figure 6: MES-4 germline genes display H3K9me2 at their promoters.
Heatmap of H3K9me2 promoter peaks in N2 L1 progeny at all 196 MES-4 germline genes
(A), and at the 34 MES-4 germline genes that were ectopically expressed in the soma of *spr-5; met-2* progeny (B). ChIP-seq reads were scaled to genome wide coverage for H3K9me2
(15 million reads). The transcriptional start site (TSS) is denoted by an orange bar with
500bp flanking regions upstream and downstream of the TSS. Integrative Genome Viewer
(IGV) image of ChIP-seq reads from N2 L1 progeny scaled to genome wide coverage for
H3K9me2 (15 million reads) over the promoters of MES-4 germline genes that were
ectopically expressed in the soma of *spr-5; met-2* progeny (C-F) and control genes (G-I). IGV
windows were scaled between 0 and 255 RPKM. Asterisks (*) denotes H3K9me2 promoter
peaks at MES-4 germline genes.

Supplemental Table S1. Raw scores for developmental delay in N2, *spr-5*, *met-2*, and *spr-5; met-2* progeny (see Figure 1A-E). Progeny were scored 72 hours after a synchronized lay as developing to the L1, L2, L3, L4, or young adult (YA) stage. N represents the total number of progeny scored in each experiment.

Supplemental Table S2. List of differentially expressed transcripts (P<0.05) in *spr-5*, *met-2* and *spr-5; met-2* L1 progeny compared to N2 progeny from DESeq2 analysis.

Supplemental Table S3. List of 176 MES-4 germline gene log2(FC) values used to generate heatmap (see Figure 2C) in *spr-5, met-2, and spr-5; met-2* L1 progeny compared to N2 progeny from DESeq2 analysis.

Supplemental Table S4: List of primer sequences used for genotyping and RT-PCR.

Supplemental Table S5: List of probe sequences for *htp-1* and *cpb-1* used in smFISH experiments.