Title: Role of the medial prefrontal cortex in the effects of rapid acting

antidepressants on affective biases in a rodent decision-making task

Running title: Rapid antidepressant effects on affective bias

Author names: Hales CA[1] (PhD), Bartlett JM[1] (BSc), Arban R[2] (PhD), Hengerer B[2] (PhD),

Robinson ESJ[1] (PhD)

Author Affiliations: [1]School of Physiology, Pharmacology and Neuroscience, Faculty of

Biomedical Sciences, University of Bristol, Bristol, BS8 1TD, UK

[2]CNS Diseases Research, Boehringer Ingelheim GmbH & Co. KG, Biberach an der Riss,

Germany

Corresponding author: Name: Prof. Emma Robinson

Email: emma.s.j.robinson@bristol.ac.uk

Address: School of Physiology, Pharmacology and Neuroscience, Faculty of Biomedical

Sciences, Tankards Close, University of Bristol, Bristol, BS8 1TD, UK.

Telephone: (+44)117 3311449

Abstract

Major Depressive Disorder is a significant and costly cause of global disability. Until the

discovery of the rapid acting antidepressant (RAAD) effects of ketamine, treatments were

limited to drugs that have delayed clinical benefits. The mechanism of action of ketamine is

currently unclear but one hypothesis is that it may involve neuropsychological effects

mediated through modulation of affective biases. Previous work has shown that affective

biases in a rodent decision-making task are differentially altered by ketamine, compared to

conventional, delayed onset antidepressants. This study sought to further investigate these

effects by comparing ketamine with other NMDA antagonists using this decision-making

task. We also investigated the subtype selective GluN2B antagonist, CP-101,606 and

muscarinic antagonist scopolamine which have both been shown to have RAAD effects.

Both CP-101,606 and scopolamine induced similar positive biases in decision-making to

ketamine, but the same effects were not seen with other NMDA antagonists. Using targeted

medial prefrontal cortex (mPFC) infusions, these effects were localised to the mPFC. In

contrast, the $GABA_A$ agonist, muscimol, induced general disruptions to behaviour. These

data suggest that ketamine and other RAADs mediate a specific effect on affective bias

which involves the mPFC. Non-ketamine NMDA antagonists lacked efficacy and we also

found that temporary inactivation of the mPFC did not recapitulate the effects of ketamine,

suggesting a specific mechanism. Furthermore, this operant behavioural task provides a

sensitive method to better understand the neuropsychological effects of RAADs.

Introduction

Major Depressive Disorder (MDD) is a prevalent psychiatric disorder, affecting over 300

million people globally[1]. It is the leading worldwide cause of disability, and, until recently,

pharmacological treatments were limited to drugs that take weeks to improve symptoms and

subjective reporting of mood[2]. The discovery of the rapid acting antidepressant (RAAD)

effects of ketamine, an NMDA receptor antagonist, has rejuvenated the field by

demonstrating that subjective changes in mood in depressed patients can be seen less than

2 hours following administration and are sustained for at least 7 days in some patients[3].

Although this RAAD has been shown repeatedly[4,5, 6,7,8], the mechanism underlying this effect

is unclear, and better understanding could be critical for the development of new, fast-acting

treatments.

Patients with MDD exhibit affective biases, whereby impairments in emotional processing

leads to reduced positive and/or enhanced negative biases in multiple cognitive domains,

including attention, memory, emotional interpretation and decision-making[9,10,11]. In humans,

acute (and chronic) treatment with conventional antidepressants induces positive biases in

emotional memory and recognition in healthy controls[12,13,14] and patients[15], despite a lack of

subjectively reported change in mood. In previous work in rodents, affective biases in a

decision-making task were differentially altered by conventional, delayed acting

antidepressants versus the RAAD ketamine[16]. In this reward-based judgement bias task,

which measures biased processing of ambiguous information associated with more or less

positive outcomes, we found that an acute, low dose of ketamine, but not acute treatment

with another NMDA receptor antagonist, PCP, immediately induced a positive bias, whereas

acute treatment with conventional antidepressants had no effect on bias[16]. However, when

given chronically, the conventional antidepressant fluoxetine did induce a positive bias[16], but

over a timescale that more closely reflects subjective changes in patients.

The aim of this study was to build upon these findings by testing a selection of other drugs

that act via NMDA receptor antagonism: lanicemine, a low-trapping NMDA receptor channel

blocker developed for the treatment of MDD, but failed to show efficacy in clinical trials[17];

memantine, an Alzheimer's medication that is a moderate affinity, non-competitive NMDA

receptor antagonist, but also lacked antidepressant efficacy in clinical trials[5,18]; and MK-801,

a potent, non-competitive NMDA receptor antagonist that has shown RAAD efficacy in

animal models[19]. We also tested other compounds that have been shown to have RAAD in

human clinical trials: the GluN2B subunit selective NMDA receptor antagonist CP-101,606[20],

and the acetylcholine muscarinic receptor antagonist scopolamine[21]. We also tested

additional doses of ketamine and PCP to ensure we had examined effects across a wider

range of receptor occupancy and in line with doses commonly used in preclinical animal

models used to study depression[22]. To investigate the mechanism underlying the rapid

positive change in decision-making bias we tested local administration of drugs shown to

cause this effect directly into the prefrontal cortex (PFC), a brain area thought to be critical in

the mechanism of RAAD of ketamine[23,24] and previously shown to modulate learning biases

in rodents[25].

Materials and Methods

Animals and apparatus

Three cohorts of male Lister Hooded rats (each cohort n=16) were used (Envigo, UK). Rats were pair-housed with environmental enrichment (see Supplementary Materials and Methods) and kept under temperature (19-23°C) and humidity (45-65%) controlled conditions on a 12-h reverse lighting cycle (lights off at 08:00h). Water was available *ad libitum* in the home cage, but rats were maintained at no less than 90% of their free-feeding body weight, matched to a standard growth curve, by restricting access to laboratory chow (LabDiet, PMI Nutrition International) to ~18g per rat per day. All procedures were carried out under local institutional guidelines (University of Bristol Animal Welfare and Ethical Review Board) and in accordance with the UK Animals (Scientific Procedures) Act 1986. During experiments all efforts were made to minimise suffering including using a low stress method of drug administration[26], and at the end of experiments rats were killed by giving an overdose of sodium pentobarbitone (200mg/kg). Behavioural testing was carried out between 0800 and 1800h, using standard rat operant chambers (Med Associates, Sandown Scientific, UK) as previously described[16,27] (and see Supplementary Materials and Methods).

Judgement bias training

Animals were trained and tested using a high versus low reward version of the judgement bias task as previously reported[16]. Rats were first trained to associate one tone (2kHz at 83dB rats, designated high reward) with a high value reward (four 45mg reward pellets; TestDiet, Sandown Scientific, UK) and the other tone (8kHz at 66dB, designated low reward) with a low value reward (one 45mg reward pellet) if they pressed the associated lever (either left or right, counterbalanced across rats) during the 20s tone. Table S1 contains a summary of training stages and criteria used. Rats were considered trained (29/23-25/25 total sessions for cohorts 1/2/3 respectively, see Table S1 for sessions for each training stage) when they maintained stable responding for three consecutive days.

Judgement bias testing

Baseline sessions (100 trials: 50 high and 50 low reward tones; presented pseudorandomly)

were conducted on Monday and Thursday. Probe test sessions (120 trials: 40 high reward,

40 low reward, and 40 ambiguous midpoint tones that were 5kHz at 75dB; pseudorandomly)

were conducted on Tuesday and Friday. The midpoint tone was randomly reinforced so that

a specific outcome could not be learnt, and to maintain responding throughout the

experiments. Cohort 1 were used to test the effect of acute systemic treatments with putative

RAAD and other NMDA receptor antagonists. Cohort 2 were made up of two groups of eight

rats that had previously been used as control animals in another experiment (data not

shown) and were then used for the extension of doses of ketamine and PCP. Cohort 3 were

used for mPFC infusion experiments. For further details see Table S2.

Study 1: the effect of acute, systemic treatments with RAADs and NMDA receptors

antagonists on judgement bias.

Experimental design: Each study used a within-subject fully counterbalanced drug

treatment schedule (see Table S2 for details of individual treatments). All drugs were given

by intraperitoneal injection using a low-stress, non-restrained technique[26]. Ketamine[¥]

(Sigma-Aldrich, UK), scopolamine[§] (Tocris, UK), lanicemine[¥] (Sigma Aldrich, UK),

memantine[¥] (Tocris, UK), MK-801[§] (Tocris, UK) and PCP[¥] (Sigma Aldrich, UK) were

dissolved in 0.9% sterile saline and given 30[§] or 60[¥] minutes prior to testing. CP-101,606

(Experiment 1: Sigma Aldrich, UK; Experiment 2: Boehringer Ingelheim GmbH) was

dissolved in 5% DMSO, 10% cremaphor and 85% sterile saline and given 60 minutes prior

to testing. Drug doses were selected based on previous rodent behavioural studies[16,28].

Doses for ketamine and PCP were chosen to extend the range of doses tested in this task

e.g. higher doses of ketamine and lower doses of PCP were used than previously[16]. For all

studies, the experimenter was blind to drug dose. Each dose-response study was separated

by at least one week (five sessions) of baseline testing. The order of testing is displayed in

Table S2.

Study 2: mPFC cannulation and infusions

To localize the site and mechanism of action of RAAD drugs, rats were implanted with

mPFC guide cannula (see Supplementary Materials and Methods for details). Following

recovery from surgery, one week of probe testing was carried out to check that judgement of

the ambiguous tone had not altered after surgery. Based on this, another two weeks of

probe testing (4 test sessions) was then conducted. Following this, an acute systemic

treatment with ketamine was given as a positive control manipulation to ensure that bias

could still be manipulated in rats post-surgery. Rats were then used for mPFC infusion

experiments. For details of the infusion procedure, see Supplementary Materials and

Methods.

In the first infusion experiment, vehicle (sterile phosphate-buffered saline (PBS); 0.0µg/µl),

ketamine (1.0µg/µl), muscimol (0.1µg/µl) or scopolamine (0.1µg/µl), all dissolved in sterile

PBS, were infused intracerebrally into mPFC 5 minutes before testing. Following this, CP-

101,606 (1.0µg/µl in the first study, 3.0µg/µl in the second study) was dissolved in 10% 2-

hydroxypropyl-cyclodextrin and 90% PBS and tested. All experiments used a within-subject

fully counterbalanced design for drug treatments, with the experimenter blind to treatment.

Drug doses were chosen based on the results from acute, systemic treatments (see Table

S2).

Following the completion of mPFC infusions, rats were killed and brains were fixed and

processed for histology (see Supplementary Materials and Methods). The locations of the

final injector tip positions in the mPFC were mapped onto standardized coronal sections of a

rat brain stereotaxic atlas.

Data and statistical analysis

For sample size justification and exclusion criterion see Supplementary Materials and

Methods. Cognitive bias index (CBI) was used as a measure of judgement bias in response

to the midpoint tone. CBI was calculated by subtracting the proportion of responses made on

the low reward lever from the proportion of responses made on the high reward lever. This

created a score between -1 and 1, where negative values represent a negative bias and

positive values a positive bias. Change from baseline in CBI was then calculated for all

experimental manipulations as follows: vehicle (0.0mg/kg) probe test CBI − drug dose probe

test CBI. This was calculated to take into account individual differences in baseline bias, and

to make directional changes caused by drug treatments clearer. To provide a value for

vehicle probe test sessions for this measure, the population average for the vehicle

(0.0mg/kg) probe test was taken away from each individual rats' CBI score for the same

session. This allowed this measure to be analysed with repeated measures analysis of

variance (rmANOVA) with session as the within-subjects factor for drug studies with more

than two treatments, or paired samples t-test for studies with only two treatments.

Response latency and accuracy, omissions and premature responses were also analysed

(see Table S3 for details of these). These measures were analysed with rmANOVAs with

session and tone as the within-subjects factors. Paired t-tests were performed as post-hoc

tests if significant effects were established. Huynh-Feldt corrections were used to adjust for

violations of the sphericity assumption, and Sidak correction was applied for multiple

comparisons. All statistical tests were conducted using SPSS 24.0.0.2 for Windows (IBM

SPSS Statistics) with $\alpha=0.05$. Results are reported with the ANOVA F-value (degrees of

freedom, error) and p-value as well as any post-hoc p-values. All graphs were made using

Graphpad Prism 7.04 for Windows (Graphpad Software, USA).

Results

Study 1: The effect of acute, systemic treatment with RAADs and selected NMDA receptor antagonists

CP-101,606: One animal was excluded in experiments 1 and 2 as accuracy criteria was not met on the vehicle session. Despite a lack of main effect when comparing all drug doses, CP-101,606 treated animals showed a positive bias for the highest dose (3.0mg/kg; one sample t-test: $p=0.038$; Figure 1A). In the second experiment, CP-101,606 (6.0mg/kg) resulted in a positive bias relative to vehicle treatment (paired samples t-test: $p=0.027$; Figure 1A). In experiment 1, 3.0mg/kg CP-101,606 also caused a decrease in response latency for the midpoint tone (main effect of session: $F_{3,42}=4.858$, $p=0.005$, post-hoc: $p=0.048$; Table 1), with a trend towards the same effect for the low tone (post-hoc: $p=0.076$; Table 1). There were no effects on other behavioural measures in experiment 1 (Table 1). In experiment 2, CP-101,606 (6.0mg/kg) caused response latencies to decrease for all three tones (main effect of session: $F_{1,14}=27.396$, $p<0.001$, post-hoc: $ps≤0.005$; Table 1). This dose had no effect on accuracy for the reference tones (Table 1), but did increase premature responses (paired samples t-test: $p=0.001$), and reduced omissions (main effect of session: $F_{1,14}=10.506$, $p=0.006$; Table 1) for the midpoint ($p=0.008$) and low tones ($p=0.024$; Table 1).

Scopolamine: The highest dose tested (0.3mg/kg) had to be excluded from the analysis as most rats did not complete sufficient trials. Scopolamine (0.1mg/kg) induced a positive bias (main effect of session: $F_{2,30}=6.739$, $p=0.004$, post-hoc: $p=0.035$; Figure 1B). This dose of scopolamine (0.1mg/kg) also increased response latencies for all three tones (main effect of session: $F_{2,30}=17.263$, $p<0.001$, post-hoc: $ps≤0.011$; Table 1), increased premature responding (main effect of session: $F_{1.355,20.330}=4.387$, $p=0.039$, post-hoc: $p=0.047$; Table 1), and increased omissions for all tones (significant session*tone interaction: $F_{2.343,35.150}=4.739$, $p=0.011$, main effect of session: $F_{2,30}=24.257$, $p<0.001$, post-hoc: $ps<0.001$; Table 1). The lower dose also caused response latencies to increase for all tones (post-hoc: $ps≤0.001$; Table 1), accuracy to increase for the low tone only (main effect of session:

$F_{1.605,24.069}$=8.558, p=0.003, post-hoc: p=0.004; Table 1), and omissions to increase for all

tones (post-hoc: ps≤0.019; Table 1).

Ketamine: In the rats who had undergone mPFC cannulation surgery, ketamine (1.0mg/kg)

caused a positive change in CBI (paired samples t-test: p=0.033; Figure 1C), as has been

seen previously[16]. Ketamine did not alter any other behavioural measures (Table 1).

Lanicemine: None of the doses of lanicemine tested caused a change in CBI (Figure 1D).

This drug also had no effect on any other behavioural measures (Table 1).

Memantine: Memantine did not cause any change in CBI at the doses tested (Figure 1E).

There was also no effect on other behavioural measures (Table 1).

MK-801: MK-801 did not change CBI (Figure 1F). The highest dose of MK-801 tested

(0.03mg/kg) decreased response latency for the midpoint tone (main effect of session:

$F_{2,30}$=3.843, p=0.033, post-hoc: p=0.013; Table 1) and also had a tendency to cause

premature responses to increase (trend towards main effect of session: $F_{1.565,23.480}$=2.915,

p=0.085, post-hoc: p=0.039; Table 1). There was no effect on accuracy for the reference

tones or percentage omissions.

High-dose ketamine: In experiment 2 (25mg/kg ketamine) one rat was excluded for failure

to complete sufficient trials. In experiments 1 and 2, ketamine (10mg/kg and 25mg/kg

respectively) did not change CBI (Figure 2A). In both experiments these higher doses did

alter all other behavioural measures. There was an increase in response latency across all

three tones for both 10mg/kg (session*tone interaction: $F_{2,30}$=7.323, p=0.003, post-hoc:

ps<0.001 for all tones; Figure 2B), and 25mg/kg ketamine (session*tone interaction:

$F_{2,28}$=4.686, p=0.018, post-hoc: ps≤0.002 for all tones; Figure 2B). Both doses decreased

premature responses (paired samples t-tests: 10mg/kg – p=0.005, 25mg/kg – p=0.006;

Figure 2C). Ketamine also improved accuracy for the low reward tone only in experiment 1

(10mg/kg: main effect of session: $F_{1,15}$=8.774, p=0.010, and trend towards session*tone

interaction: $F1,15$=3.665, p=0.075, post-hoc: p=0.022; Figure 2D) and experiment 2

(25mg/kg: trend towards main effect of session: $F_{1,14}$=4.500, p=0.052, and session*tone

interaction: $F1,14$ = 5.513, p=0.034, post-hoc: p=0.033; Figure 2D). In both experiments,

there was an increase in omissions for all three tones (experiment 1, 10mg/kg: session*tone

interaction: $F_{1.401,21.021}$=5.662, p=0.018, post-hoc: high reward tone – p=0.015, midpoint tone:

p=0.003, low reward tone: p=0.010; experiment 2, 25mg/kg: session*tone interaction:

$F_{1.368,19.150}$=11.964, p=0.001, post-hoc: high reward tone – p=0.003, midpoint tone – p<0.001,

low reward tone – p=0.001; Figure 2E).

Low dose PCP: Doses of PCP (0.03, 0.1, 0.3mg/kg) that were lower than those previously

tested[16] did not cause any change in CBI (Figure 1G). There was also no effect on any other

behavioural measures (Table 1).

Study 2: mPFC infusions of drugs shown to cause positive judgement biases

Two rats were excluded in cohort 3: one rat did not meet accuracy criteria for any probe (or

baseline) session following the second drug infusion; and after the end of testing another

animal was found to have an incorrect cannula placement. Therefore, both were excluded

retrospectively from the entire study. Compared to pre-surgery performance, the CBI of rats

became more negative after surgery, and this was stable across testing over three weeks

(main effect of week: $F_{3,42}$=6.335, p=0.001, post-hoc: ps≤0.011; Figure 3A). There were no

differences in response latencies, premature responses, accuracies for reference tones or

omissions before compared to after surgery (Table 1).

In the first infusion experiment, ketamine (1.0µg/µl), muscimol (0.1µg/µl) and scopolamine

(0.1µg/µl) all induced positive biases (main effect of session: $F_{3,36}$=7.241, p=0.001; post-hoc:

ketamine – p=0.012, muscimol – p=0.001, scopolamine – p=0.032 Figure 3C). The effect of

PFC infusion of ketamine or scopolamine was specific to CBI, as these drugs had no effect

on other behavioural measures (Figure 3D-G), unlike muscimol infusions which caused

changes to all other behavioural measures. There was an increase in response latency

(session*tone interaction: $F_{6,72}$=4.181, p=0.001) for the high reward (post-hoc: p<0.001) and

midpoint tone (p=0.028; Figure 3D), and a large increase in premature responses to over

100% (main effect of session: $F_{1.151,13.809}$=33.784, p<0.001, post-hoc: p<0.001; Figure 3E).

Muscimol also caused accuracy to decrease for both reference tones (main effect of

session: $F_{1.181,14.172}$=43.775, p<0.001, post-hoc: ps≤0.001; Figure 3F). For the low reward

tone, this reduction was so great that rats were no longer performing any better than chance

(one-sample t-test against a test value of 50%: p=0.197; Figure 3F). Omissions for the high

and low reward tones increased following muscimol infusion (main effect of session:

$F_{1.338,16.057}$=10.418, p=0.003, post-hoc: high tone – p=0.010, low tone – p=0.012; Figure 3G).

In the experiments testing the effect of CP-101,606 mPFC infusion, in experiment 1 the

lower dose (1.0µg/µl) did not alter CBI (Figure 4A), but in experiment 2, the higher dose

(3.0µg/µl) induced a positive bias (paired samples t-test: p=0.043; Figure 4A). In experiment

1, CP-101,606 (1.0µg/µl) caused an increase in response latency for the midpoint tone only

(main effect of session: $F_{1,12}$=5.064, p=0.044, post-hoc: p=0.019; Figure 4B) but had no

other behavioural effects (Figure 4C-E). In experiment 2, 3.0µg/µl CP-101,606 did not have

any effects on other behavioural measures (Figure 4B-E).

Discussion

As previously shown, low dose ketamine (1.0mg/kg) had a specific effect on decision-making

biases, inducing a positive change in CBI following acute administration. This effect of

ketamine was dose dependent, with higher doses having general effects on task

performance without changing CBI. The effects of ketamine were recapitulated to some

extent by the GluN2B antagonist, CP-101,606 and muscarinic antagonist, scopolamine, but

both also had more general effects on other behavioural measures following systemic

administration. All three treatments have previously been reported to have RADD effects in

clinical trials[3,20,21], whilst the other NMDA antagonists tested here did not[5,17,18], and these

also failed to induce a change in bias. The mPFC infusions suggest that this brain region is

central to the effects of ketamine, scopolamine and CP-101,606. Interestingly, mPFC

infusions more specifically altered bias, suggesting other brain regions may contribute to the

systemic effects on other behavioural measures. The importance of the mPFC in modulating

RAAD effects in neuropsychological tasks is consistent with previous findings in our learning

and memory bias assay, the affective bias test[25]. Inactivation of the mPFC with muscimol did

positively change bias but animals also exhibited large changes in other behavioural

measures. This suggests that the RAADs can modulate activity in this brain region in a more

specific way than muscimol, which results in a relatively specific effect on biases in decision-

making.

For lanicemine and memantine, the lack of any behavioural effects means there is a

possibility that the doses tested were too low. However, for both of these the range of doses

tested covers the doses that are equivalent to those used humans in clinical trials

(lanicemine: 50, 100mg[17], equivalent to approximately 0.75, 1.5mg/kg; memantine: 5-20mg[5],

equivalent to approximately 0.07-0.3mg/kg), paralleling our effective dose of ketamine

(1.0mg/kg, similar to the 0.5mg/kg dose used by Zarate et al.[3]). Furthermore, when testing

lower doses of PCP (another NMDA receptor antagonist not known to show RAAD) than

previously used[16], we also failed to see any change in CBI. Conversely, when we tested

higher doses of ketamine than those we had previously, doses that have been shown to

have antidepressant effects in other preclinical models used to study depression[29], we failed

to see any change in bias, instead only seeing non-specific changes in other behavioural

measures. This lends support to our interpretation that this reward-based judgement bias

task can specifically dissociate between drugs that do show RAAD, and those that do not,

despite them having similar pharmacology.

The difference in specificity on behavioural effects, whereby ketamine (1.0mg/kg) only

positively changes decision-making bias, but both CP-101,606 and scopolamine have other

non-specific effects, suggests that 1.0mg/kg ketamine is able to relatively selectively

modulate affective bias. The changes in response latencies, omissions and premature

responses caused by CP-101,606 and scopolamine suggest that these drugs may also be

having effects on other cognitive processes, such as motivation. However, the direction of

changes for these drugs are in opposite directions (decreases in response latency and

omissions for CP-101,606 but increases in these for scopolamine) despite them both

causing positive changes in CBI. This, combined with the lack of change in accuracy for the

reference tones, suggest that these non-specific effects cannot fully explain the change in

decision-making bias.

The neurobiology underlying the relative specificity of ketamine, CP-101,606 and

scopolamine in being able to immediately alter decision-making bias, in contrast to the other

NMDA receptor antagonists tested that have not shown these effects, are likely to be due to

differences in their mechanisms of action. Our findings add weight to the strong body of

evidence suggesting that NMDA receptor antagonism is important for short-term, RAAD

effects of these drugs[30], but suggests that specific modulation of either a specific subtype of

the receptor or a sub-population of neurons may be involved. CP-101,606 is selective for the

GluN2B NMDA receptor subunit, whilst it has been shown that scopolamine, and more

recently ketamine, cause a glutamate burst via blockade of NMDA receptors specifically on

GABA interneurons that leads to increased mechanistic target of rapamycin complex 1

signalling, brain-derived neurotrophic factor release and synaptic changes in the

PFC[31,32,33,34]. Further studies would be required to test whether these mechanisms also drive

these drugs effects on affective bias.

The infusion studies localise the site of action of this rapid change in decision-making bias to

the mPFC, corresponding with brain imaging studies in humans that have also shown

ketamine-dependent changes in prefrontal glutamatergic neurotransmission[23,24]. This also

matches with previous rodent studies, where using the affective bias test, it has been shown

that whilst ketamine does not induce positive biases in learning, it can remediate previously

acquired negative biases, an effect which also localises to the mPFC[25].

Unlike when given systemically, intracerebral mPFC infusion of CP-101,606 or scopolamine

did not cause any non-specific behavioural changes on the task. This could suggest that

these effects are driven by off-target effects of drug binding in other brain areas, or in the

case of scopolamine, the periphery. The pronounced, non-specific behavioural effects seen

following muscimol infusion suggests that mechanism of action of these drugs is more

refined than global inhibition of neurotransmission in the mPFC. Previous work in humans

and rodents has shown that subcortical and limbic brain regions, such as the amygdala, are

important in the neurocircuitry of MDD / depression-related behaviour[35,36,37], and a recent

study suggests that ketamine may play a critical role in restoring dysfunctional connectivity in

these circuits[24]. Furthermore, in rodents, a recent study found that optogenetic activation of

pyramidal mPFC neurons containing dopamine receptor D1 caused RAAD-like responses in

the forced swim test, and that blockade of these receptors prevented the RADD effects of

ketamine[38]. In order to further our understanding of this mechanism, it will be important to

investigate the effects of these drugs on different neuronal subtypes within the mPFC, as

well as investigating the wider circuitry that is altered by these drugs.

Final conclusions

This study adds to the evidence that the neuropsychological effects of ketamine are

potentially important in its RAAD in patients with MDD, and that these effects in altering

affective biases, both in decision-making as demonstrated here, as well as in learning and

memory occur at time points (one hour) before major plastic changes arise. It will be

important to investigate the neurobiological effects of not just the immediate, RAAD of

ketamine, but also the sustained effects by examining how affective biases are altered at

longer time points. Furthermore, investigation of the wider circuits involved in this RAAD

efficacy will be crucial in revealing the mechanism underlying these actions, which will be

important for the development of novel therapeutics. Ketamine (at 1.0 mg/kg) seems to have

very specific effects on affective bias, which we can capitalise on to better understand the

circuits that contribute to these modulations of affective biases that are potentially very

important in the cause, perpetuation and treatment of MDD.

Funding and Disclosure: This research was funded by an Industrial Partnership Award

awarded by BBSRC in collaboration with Boehringer Ingelheim (Grant no: BB/N015762/1)

and carried out with intellectual support from Boehringer Ingelheim. ESJR has current or

previously obtained research grant funding through PhD studentships, collaborative grants

and contract research from Boehringer Ingelheim, Compass Pathways, Eli Lilly, MSD, Pfizer

and SmallPharma.

Author contributions: CAH performed the research, analysed data, wrote and edited the

paper. JMB performed research and analysed data. RA and BH designed the research and

edited the paper. ESJR designed the research and wrote and edited the paper.

References

1. World Health Organization, 2018. Depression fact sheet 2018 [Available from: http://

https://www.who.int/news-room/fact-sheets/detail/depression.] Accessed 05/11/2019.

2. Anderson IM, Nutt DJ, Deakin JFW (2000) Evidence-based guidelines for treating

depressive disorders with antidepressants: a revision of the 1993 British Association for

Psychopharmacology guidelines. Journal of Psychopharmacology 14:3-20.

3. Zarate CA, Jr., Singh JB, Carlson PJ, Brutsche NE, Ameli R, Luckenbaugh DA, Charney

DS, Manji HK (2006b) A randomized trial of an N-methyl-D-aspartate antagonist in

treatment-resistant major depression. Arch Gen Psychiatry 63:856-864.

4. Berman RM, Cappiello A, Anand A, Oren DA, Heninger GR, Charney DS, Krystal JH

(2000) Antidepressant effects of ketamine in depressed patients. Biol Psychiatry 47:351-354.

5. Zarate CA, Jr., Singh JB, Quiroz JA, De Jesus G, Denicoff KK, Luckenbaugh DA, Manji

HK, Charney DS (2006a) A double-blind, placebo-controlled study of memantine in the

treatment of major depression. The American Journal of Psychiatry 163:153-155.

6. DiazGranados N, Ibrahim LA, Brutsche NE, Ameli R, Henter ID, Luckenbaugh DA,

Machado-Vieira R, Zarate CA, Jr. (2010) Rapid resolution of suicidal ideation after a single

infusion of an N-methyl-D-aspartate antagonist in patients with treatment-resistant major

depressive disorder. The Journal of Clinical Psychiatry 71:1605-1611.

7. Lapidus KA, Levitch CF, Perez AM, Brallier JW, Parides MK, Soleimani L, Feder A,

Iosifescu DV, Charney DS, Murrough JW (2014) A randomized controlled trial of intranasal

ketamine in major depressive disorder. Biol Psychiatry 76:970-976.

8. Price RB, Iosifescu DV, Murrough JW, Chang LC, Al Jurdi RK, Iqbal SZ, Soleimani L,

Charney DS, Foulkes AL, Mathew SJ (2014) Effects of ketamine on explicit and implicit

suicidal cognition: a randomized controlled trial in treatment-resistant depression.

Depression and Anxiety 31:335-343.

9. Mathews A, MacLeod C (2005) Cognitive Vulnerability to Emotional Disorders. Annual

Review of Clinical Psychology 1:167-195.

10. Clark L, Chamberlain SR, Sahakian BJ (2009) Neurocognitive Mechanisms in

Depression: Implications for Treatment. Annual Review of Neuroscience 32:57-74.

11. Gotlib IH, Joormann J (2010) Cognition and Depression: Current Status and Future

Directions. Annual Review of Clinical Psychology 6:285-312.

12. Harmer CJ, Bhagwagar Z, Perrett DI, Vollm BA, Cowen PJ, Goodwin GM (2003) Acute

SSRI administration affects the processing of social cues in healthy volunteers.

Neuropsychopharmacology 28:148-152.

13. Harmer CJ, Shelley NC, Cowen PJ, Goodwin GM (2004) Increased positive versus

negative affective perception and memory in healthy volunteers following selective serotonin

and norepinephrine reuptake inhibition. The American Journal of Psychiatry 161:1256-1263.

14. Harmer CJ, de Bodinat C, Dawson GR, Dourish CT, Waldenmaier L, Adams S, Cowen

PJ, Goodwin GM (2011) Agomelatine facilitates positive versus negative affective

processing in healthy volunteer models. Journal of Psychopharmacology (Oxford, England)

25:1159-1167.

15. Harmer CJ, O'Sullivan U, Favaron E, Massey-Chase R, Ayres R, Reinecke A, Goodwin

GM, Cowen PJ (2009) Effect of acute antidepressant administration on negative affective

bias in depressed patients. The American Journal of Psychiatry 166:1178-1184.

16. Hales CA, Houghton CJ, Robinson ESJ (2017) Behavioural and computational methods

reveal differential effects for how delayed and rapid onset antidepressants effect decision

making in rats. European neuropsychopharmacology 27:1268-1280.

17. Sanacora G, Johnson MR, Khan A, Atkinson SD, Riesenberg RR, Schronen JP, Burke

MA, Zajecka JM, Barra L, Su HL, Posener JA, Bui KH, Quirk MC, Piser TM, Mathew SJ,

Pathak S (2017) Adjunctive Lanicemine (AZD6765) in Patients with Major Depressive

Disorder and History of Inadequate Response to Antidepressants: A Randomized, Placebo-

Controlled Study. Neuropsychopharmacology 42:844-853.

18. Smith EG, Deligiannidis KM, Ulbricht CM, Landolin CS, Patel JK, Rothschild AJ (2013)

Antidepressant augmentation using the N-methyl-D-aspartate antagonist memantine: a

randomized, double-blind, placebo-controlled trial. The Journal of Clinical Psychiatry 74:966-

973.

19. Ates-Alagoz Z, Adejare A (2013) NMDA Receptor Antagonists for Treatment of

Depression. Pharmaceuticals (Basel) 6:480-499.

20. Preskorn SH, Baker B, Kolluri S, Menniti FS, Krams M, Landen JW (2008) An innovative

design to establish proof of concept of the antidepressant effects of the NR2B subunit

selective N-methyl-D-aspartate antagonist, CP-101,606, in patients with treatment-refractory

major depressive disorder. Journal of Clinical Psychopharmacology 28:631-637.

21. Furey ML, Drevets WC (2006) Antidepressant efficacy of the antimuscarinic drug

scopolamine: a randomized, placebo-controlled clinical trial. Arch Gen Psychiatry 63:1121-

1129.

22. Browne CA, Lucki I (2013) Antidepressant effects of ketamine: mechanisms underlying

fast-acting novel antidepressants. Frontiers in Pharmacology 4:161.

23. Li CT, Chen MH, Lin WC, Hong CJ, Yang BH, Liu RS, Tu PC, Su TP (2016) The effects

of low-dose ketamine on the prefrontal cortex and amygdala in treatment-resistant

depression: A randomized controlled study. Hum Brain Mapp 37:1080-1090.

24. Abdallah CG, De Feyter HM, Averill LA, Jiang L, Averill CL, Chowdhury GMI, Purohit P,

de Graaf RA, Esterlis I, Juchem C, Pittman BP, Krystal JH, Rothman DL, Sanacora G,

Mason GF (2018) The effects of ketamine on prefrontal glutamate neurotransmission in

healthy and depressed subjects. Neuropsychopharmacology 43:2154-2160.

25. Stuart SA, Butler P, Munafo MR, Nutt DJ, Robinson ES (2015) Distinct

Neuropsychological Mechanisms May Explain Delayed- Versus Rapid-Onset Antidepressant

Efficacy. Neuropsychopharmacology 40:2165-74.

26. Stuart SA, Robinson ESJ (2015) Reducing the stress of drug administration: implications

for the 3Rs. Scientific Reports 5:14288.

27. Hales CA, Robinson ES, Houghton CJ (2016) Diffusion Modelling Reveals the Decision

Making Processes Underlying Negative Judgement Bias in Rats. PloS ONE 11:e0152592.

28. Benn A, Robinson ES (2014) Investigating glutamatergic mechanism in attention and

impulse control using rats in a modified 5-choice serial reaction time task. PloS ONE

9:e115374.

29. Polis AJ, Fitzgerald PJ, Hale PJ, Watson BO (2019) Rodent ketamine depression-related

research: Finding patterns in a literature of variability. Behav Brain Res 376:112153.

30. Drewniany E, Han J, Hancock C, Jones RL, Lim J, Nemat Gorgani N, Sperry JK, 3rd, Yu

HJ, Raffa RB (2015) Rapid-onset antidepressant action of ketamine: potential revolution in

understanding and future pharmacologic treatment of depression. J Clin Pharm Ther 40:125-

130.

31. Li N, Lee B, Liu RJ, Banasr M, Dwyer JM, Iwata M, Li XY, Aghajanian G, Duman RS

(2010) mTOR-dependent synapse formation underlies the rapid antidepressant effects of

NMDA antagonists. Science 329:959-964.

32. Voleti B, Navarria A, Liu RJ, Banasr M, Li N, Terwilliger R, Sanacora G, Eid T,

Aghajanian G, Duman RS (2013) Scopolamine rapidly increases mammalian target of

rapamycin complex 1 signaling, synaptogenesis, and antidepressant behavioral responses.

Biol Psychiatry 74:742-749.

33. Wohleb ES, Wu M, Gerhard DM, Taylor SR, Picciotto MR, Alreja M, Duman RS (2016)

GABA interneurons mediate the rapid antidepressant-like effects of scopolamine. J Clin

Invest 126:2482-2494.

34. Gerhard DM, Pothula S, Liu RJ, Wu M, Li XY, Girgenti MJ, Taylor SR, Duman CH,

Delpire E, Picciotto M, Wohleb ES, Duman RS (2019) GABA interneurons are the cellular

trigger for ketamine's rapid antidepressant actions. J Clin Invest pii: 130808. [Epub ahead of

print]

35. Eshel N, Roiser JP (2010) Reward and punishment processing in depression. Biol

Psychiatry 68:118-124.

36. Murrough JW, Iacoviello B, Neumeister A, Charney DS, Iosifescu DV (2011) Cognitive

dysfunction in depression: neurocircuitry and new therapeutic strategies. Neurobiol Learn

Mem 96:553-563.

37. Biselli T, Lange S, Sablottny L, Steffen J, Walther A (2019) Optogenetic and

chemogenetic insights into the neurocircuitry of depression-like behaviour: A systematic

review. The European Journal of Neuroscience. doi: 10.1111/ejn.14603. [Epub ahead of

print]

38. Hare BD, Shinohara R, Liu RJ, Pothula S, DiLeone RJ, Duman RS (2019) Optogenetic

stimulation of medial prefrontal cortex Drd1 neurons produces rapid and long-lasting

antidepressant effects. Nat Commun 10:223.

  Figure Legends

Figure 1 – *The effect of acute treatment with rapid acting antidepressant drugs and NMDA*

receptor antagonists on judgement bias of the midpoint ambiguous tone.

Ketamine (0.0, 1.0 mg/kg; n = 13), scopolamine (0.0, 0.03, 0.1 mg/kg; n = 16), CP-101,606

(Expt 1: 0.0, 0.3, 1.0, 3.0 mg/kg, n = 15; Expt 2: 0.0, 6.0 mg/kg, n = 15), lanicemine (0.0, 0.3,

1.0, 3.0 mg/kg; n = 16), memantine (0.0, 0.1, 0.3, 1.0 mg/kg; n = 16) and MK-801 (0.0, 0.01,

0.03 mg/kg; n = 16 were administered acutely by intraperitoneal injection prior to testing on

the judgement bias task. (A) Replicating previous studies, ketamine (1.0 mg/kg) positively

changed CBI. (B) Scopolamine (0.1 mg/kg) also caused a positive change from baseline in

CBI. (C) In experiment 1, the 3.0 mg/kg dose caused a positive change in cognitive bias

index (CBI) for the midpoint tone. A positive change was also seen in experiment 2 with a

higher 6.0 mg/kg dose. (D-G) Lanicemine, memantine, MK-801 and low doses of PCP did

not induce a change in CBI for the midpoint tone at the doses tested. Data shown and

represent mean ± SEM (bars and error bars) overlaid with individual data points for each rat.

Dashed line (panel C) indicates separate, counterbalanced experiments. *$p < 0.05$. CP-

101,606, ketamine, lanicemine, memantine, PCP: 60 min pre-treatment; scopolamine, MK-

801: 30 min pre-treatment

Figure 2 – *Behavioural data from the judgement bias task following acute treatment with*

high doses of ketamine.

Acute doses of ketamine (Expt 1: 0.0, 10.0 mg/kg, n = 16; Expt 2: 0.0, 25.0 mg/kg, n = 16)

were administered by intraperitoneal injection to measure their effect on judgement bias. (A)

Neither high dose of ketamine caused a change in interpretation of the midpoint tone. (B)

Both doses of ketamine increased accuracy for the low tone. (C) Both doses of ketamine

increased response latencies across all three tones. (D) Omissions were increased across

all three tones following both ketamine doses. (E) High doses of ketamine (10.0, 25.0 mg/kg)

decreased premature responding. ***$p < 0.001$, **$p < 0.01$, *$p < 0.05$. Data represent mean ±

SEM (panels B-E) with individual data points overlaid for each rat (panel A). Dashed lines

indicate separate, counterbalanced experiments. 60 min pre-treatment. HT - high reward

tone; MT - midpoint tone; LT - low reward tone.

Figure 3 – D*ata from mPFC cannulated rats on the judgement bias task.*

Probe tests with no experimental manipulation were conducted before and after mPFC

cannulation surgery to ensure that the surgery itself did not effect performance in the

judgement bias task. (A) Cognitive bias index became more negative in the probe tests

conducted after surgery. (B) The location of the injector placement was confirmed post-

mortem and black dots represent the location of the cannula tip as assessed from Cresyl

violet-stained brain sections. Coronal sections are +3.7 mm to +2.5mm relative to bregma

(Paxinos and Watson, 1998). (C-G) In the first infusion experiment, ketamine (Ket; 1.0 µg/µl)

muscimol (Mus; 0.1 µg/µl), scopolamine (Sco; 0.1 µg/µl) or vehicle (Veh; 0.0 µg/µl; n = 13),

were administered by intracerebral infusion into the mPFC to measure the effect on

judgement bias. (C) Ketamine, muscimol and scopolamine all caused a positive change in

cognitive bias index (CBI) for the midpoint tone. (D) Muscimol decreased accuracy for both

reference tones. (E) Muscimol increased response latencies for the high and midpoint tones.

(F) For the high and low tones, muscimol increased omissions. (G) Muscimol also increased

premature responding. Data represent mean ± SEM (panels A, C-F) with individual data

points overlaid for each rat (panel A,C). Black dashed line (panel f) represents 50% accuracy

depicting performance at chance. 5 min pre-treatment. ***$p < 0.001$, *$p < 0.05$. HT - high

reward tone; MT - midpoint tone; LT - low reward tone.

Figure 4 – *Behavioural data from the judgement bias task following mPFC infusions of CP-*

101,606.

CP-101,606 (Expt 1: 00.0, 1.0 µg/µl, n = 13; Expt 2: 0.0, 3.0 µg/µl, n = 12) was administered

by intracerebral infusion in the mPFC to measure the effect on judgement bias. (A) The

higher dose of CP-101,606 (3.0 µg/µl) caused a positive change from baseline in CBI. (B)

Accuracy was not altered by either dose of CP-101,606. (C) In experiment 1, CP-101,606

(1.0 µg/µl) increased response latency for the midpoint tone. (D/E) There was no effect of

either dose on omissions or premature responding, *$p < 0.05$. Data represent mean ± SEM

(panels B-E) with individual data points overlaid for each rat (panel A). Dashed lines indicate

separate, counterbalanced experiments. 5 min pre-treatment. HT - high reward tone; MT -

midpoint tone; LT - low reward tone.

Figure 1 – *The effect of acute treatment with rapid acting antidepressant drugs and NMDA receptor antagonists on judgement bias of the midpoint ambiguous tone.* Ketamine (0.0, 1.0 mg/kg; n = 13), scopolamine (0.0, 0.03, 0.1 mg/kg; n = 16), CP-101,606 (Expt 1: 0.0, 0.3, 1.0, 3.0 mg/kg, n = 15; Expt 2: 0.0, 6.0 mg/kg, n = 15), lanicemine (0.0, 0.3, 1.0, 3.0 mg/kg; n = 16), memantine (0.0, 0.1, 0.3, 1.0 mg/kg; n = 16) and MK-801 (0.0, 0.01, 0.03 mg/kg; n = 16 were administered acutely by intraperitoneal injection prior to testing on the judgement bias task. (A) Replicating previous studies, ketamine (1.0 mg/kg) positively changed CBI. (B) Scopolamine (0.1 mg/kg) also caused a positive change from baseline in CBI. (C) In experiment 1, the 3.0 mg/kg dose caused a positive change in cognitive bias index (CBI) for the midpoint tone. A positive change was also seen in experiment 2 with a higher 6.0 mg/kg dose. (D-G) Lanicemine, memantine, MK-801 and low doses of PCP did not induce a change in CBI for the midpoint tone at the doses tested. Data shown and represent mean ± SEM (bars and error bars) overlaid with individual data points for each rat. Dashed line (panel C) indicates separate, counterbalanced experiments. *$p < 0.05$. CP-101,606, ketamine, lanicemine, memantine, PCP: 60 min pre-treatment; scopolamine, MK-801: 30 min pre-treatment

Figure 2 – *Behavioural data from the judgement bias task following acute treatment with high doses of ketamine.*

Acute doses of ketamine (Expt 1: 0.0, 10.0 mg/kg, n = 16; Expt 2: 0.0, 25.0 mg/kg, n = 16) were administered by intraperitoneal injection to measure their effect on judgement bias. (A) Neither high dose of ketamine caused a change in interpretation of the midpoint tone. (B) Both doses of ketamine increased accuracy for the low tone. (C) Both doses of ketamine increased response latencies across all three tones. (D) Omissions were increased across all three tones following both ketamine doses. (E) High doses of ketamine (10.0, 25.0 mg/kg) decreased premature responding. ***$p < 0.001$, **$p < 0.01$, *$p < 0.05$. Data represent mean ± SEM (panels B-E) with individual data points overlaid for each rat (panel A). Dashed lines indicate separate, counterbalanced experiments. 60 min pre-treatment. HT - high reward tone; MT - midpoint tone; LT - low reward tone.

Figure 3 – D*ata from mPFC cannulated rats on the judgement bias task.*
Probe tests with no experimental manipulation were conducted before and after mPFC cannulation surgery
to ensure that the surgery itself did not effect performance in the judgement bias task. (A) Cognitive bias
index became more negative in the probe tests conducted after surgery. (B) The location of the injector
placement was confirmed post-mortem and black dots represent the location of the cannula tip as assessed
from Cresyl violet-stained brain sections. Coronal sections are +3.7 mm to +2.5mm relative to bregma
(Paxinos and Watson, 1998). (C-G) In the first infusion experiment, ketamine (Ket; 1.0 µg/µl) muscimol (Mus;
0.1 µg/µl), scopolamine (Sco; 0.1 µg/µl) or vehicle (Veh; 0.0 µg/µl; n = 13), were administered by intracerebral
infusion into the mPFC to measure the effect on judgement bias. (C) Ketamine, muscimol and scopolamine all
caused a positive change in cognitive bias index (CBI) for the midpoint tone. (D) Muscimol decreased accuracy
for both reference tones. (E) Muscimol increased response latencies for the high and midpoint tones. (F) For
the high and low tones, muscimol increased omissions. (G) Muscimol also increased premature responding.
Data represent mean ± SEM (panels A, C-F) with individual data points overlaid for each rat (panel A,C). Black
dashed line (panel f) represents 50% accuracy depicting performance at chance. 5 min pre-treatment. ***$p <$
0.001, *$p < 0.05$. HT - high reward tone; MT - midpoint tone; LT - low reward tone.

Figure 4 – *Behavioural data from the judgement bias task following mPFC infusions of CP-101,606.*
CP-101,606 (Expt 1: 00.0, 1.0 µg/µl, n = 13; Expt 2: 0.0, 3.0 µg/µl, n = 12) was administered by intracerebral infusion in the mPFC to measure the effect on judgement bias. (A) The higher dose of CP-101,606 (3.0 µg/µl) caused a positive change from baseline in CBI. (B) Accuracy was not altered by either dose of CP-101,606. (C) In experiment 1, CP-101,606 (1.0 µg/µl) increased response latency for the midpoint tone. (D/E) There was no effect of either dose on omissions or premature responding, *p < 0.05. Data represent mean ± SEM (panels B-E) with individual data points overlaid for each rat (panel A). Dashed lines indicate separate, counterbalanced experiments. 5 min pre-treatment. HT - high reward tone; MT - midpoint tone; LT - low reward tone.

Table 1 – *Data for behavioural measures from the JBT.*

Drug / Session	Dose (mg/kg) / Week	Response latency			% Accuracy		% Omissions			% Premature
		HT	MT	LT	HT	LT	HT	MT	LT	
Ketamine	0.0	1.62 ± 0.12	3.23 ± 0.22	3.80 ± 0.36	97.88 ± 0.62	83.06 ± 1.96	0.00 ± 0.00	0.58 ± 0.30	1.15 ± 0.36	12.44 ± 1.94
	1.0	1.77 ± 0.13	3.38 ± 0.29	4.05 ± 0.44	97.69 ± 9.57	83.54 ± 2.48	0.20 ± 0.20	0.38 ± 0.26	0.78 ± 0.34	14.09 ± 3.07
CP-101,606	0.0	2.19 ± 0.18	3.74 ± 0.39	4.26 ± 0.32	97.82 ± 0.77	84.87 ± 1.74	0.38 ± 0.26	0.88 ± 0.41	1.28 ± 0.55	8.34 ± 1.24
	0.3	2.43 ± 0.21	3.85 ± 0.36	4.54 ± 0.34	97.54 ± 0.91	85.44 ± 2.39	0.00 ± 0.00	0.82 ± 0.38	1.15 ± 0.47	7.42 ± 1.23
	1.0	2.21 ± 0.15	3.48 ± 0.26	4.46 ± 0.29	98.33 ± 0.72	83.49 ± 2.16	0.00 ± 0.00	0.69 ± 0.41	0.83 ± 0.47	8.80 ± 1.26
	3.0	1.97 ± 0.17	*2.88 ± 0.23	#3.72 ± 0.27	95.83 ± 1.14	82.45 ± 2.62	0.00 ± 0.00	0.17 ± 0.17	0.33 ± 0.23	12.44 ± 2.58
CP-101,606	0.0	3.05 ± 0.24	4.52 ± 0.27	5.38 ± 0.33	98.25 ± 0.51	86.57 ± 1.70	0.70 ± 0.39	2.50 ± 0.82	2.81 ± 0.80	4.24 ± 2.58
	6.0	**2.32 ± 0.13	***3.52 ± 0.17	***4.10 ± 0.19	97.64 ± 0.87	84.37 ± 1.98	0.00 ± 0.00	**0.00 ± 0.00	*0.72 ± 0.32	**11.75 ± 1.80
Scopolamine	0.0	2.28 ± 0.17	3.84 ± 0.24	4.42 ± 0.31	97.80 ± 0.64	86.56 ± 1.84	0.16 ± 0.16	0.31 ± 0.21	1.09 ± 0.51	6.20 ± 0.70
	0.03	***4.10 ± 0.34	**5.52 ± 0.38	*6.51 ± 0.40	99.50 ± 0.27	**94.01 ± 1.40	*7.17 ± 2.17	***17.18 ± 3.38	**18.02 ± 3.60	7.31 ± 1.55
	0.1	***4.26 ± 0.38	**5.54 ± 0.37	*6.27 ± 0.51	95.00 ± 1.76	85.26 ± 3.93	*22.85 ± 3.79	***27.98 ± 3.91	**35.92 ± 6.05	*14.52 ± 3.85
Lanicemine	0.0	2.42 ± 0.22	3.80 ± 0.31	4.72 ± 0.45	96.60 ± 0.82	86.24 ± 1.81	0.00 ± 0.00	3.16 ± 2.21	2.93 ± 1.89	7.17 ± 1.11
	0.3	2.23 ± 0.17	3.60 ± 0.24	4.40 ± 0.29	98.13 ± 0.63	85.01 ± 1.89	0.00 ± 0.00	0.31 ± 0.21	1.26 ± 0.95	9.48 ± 1.36
	1.0	2.40 ± 0.22	4.20 ± 0.32	4.79 ± 0.40	96.41 ± 1.58	85.45 ± 2.15	0.00 ± 0.00	3.54 ± 1.99	4.03 ± 2.76	6.68 ± 1.03
	3.0	2.55 ± 0.21	3.89 ± 0.28	4.81 ± 0.37	98.58 ± 0.57	86.74 ± 1.71	0.16 ± 0.16	1.56 ± 1.25	2.81 ± 1.72	8.18 ± 1.67
Memantine	0.0	2.38 ± 0.15	4.12 ± 0.21	4.94 ± 0.23	97.77 ± 0.60	83.96 ± 1.52	0.19 ± 0.19	1.16 ± 0.44	2.07 ± 1.03	7.98 ± 0.93
	0.1	2.48 ± 0.18	4.15 ± 0.23	4.65 ± 0.30	97.45 ± 0.82	84.75 ± 1.81	0.16 ± 0.16	2.30 ± 0.85	1.09 ± 0.45	8.45 ± 1.25
	0.3	2.42 ± 0.18	3.84 ± 0.25	4.53 ± 0.30	97.49 ± 0.65	81.21 ± 1.79	0.16 ± 0.16	0.63 ± 0.36	2.34 ± 0.84	9.25 ± 1.40
	1.0	2.40 ± 0.16	3.90 ± 0.23	4.50 ± 0.28	97.66 ± 0.58	84.98 ± 1.64	0.00 ± 0.00	1.56 ± 0.75	2.03 ± 1.25	8.38 ± 1.24
MK-801	0.0	2.51 ± 0.15	4.16 ± 0.28	4.72 ± 0.30	96.39 ± 1.00	85.23 ± 1.78	0.16 ± 0.16	1.12 ± 0.40	1.31 ± 0.54	6.66 ± 0.77
	0.01	2.25 ± 0.14	3.80 ± 0.21	4.39 ± 0.27	96.71 ± 0.78	83.05 ± 1.64	0.16 ± 0.16	0.47 ± 0.25	1.09 ± 0.51	8.39 ± 1.72
	0.03	2.34 ± 0.13	*3.44 ± 0.29	4.07 ± 0.35	97.19 ± 0.88	81.19 ± 2.15	0.00 ± 0.00	1.41 ± 0.56	1.88 ± 0.74	*10.73 ± 1.64
PCP	0.0	2.23 ± 0.17	3.88 ± 0.23	4.96 ± 0.42	98.28 ± 0.50	84.89 ± 1.88	0.00 ± 0.00	0.47 ± 0.25	0.47 ± 0.34	6.46 ± 0.98
	0.03	2.22 ± 0.18	3.99 ± 0.29	5.11 ± 0.42	98.59 ± 0.56	87.44 ± 1.70	0.00 ± 0.00	1.25 ± 0.60	2.03 ± 1.08	5.72 ± 0.75
	0.1	2.25 ± 0.18	3.81 ± 0.29	4.60 ± 0.30	99.07 ± 0.31	86.37 ± 1.87	0.31 ± 0.21	0.94 ± 0.39	1.41 ± 0.94	5.63 ± 0.89
	0.3	2.24 ± 0.20	3.93 ± 0.37	4.99 ± 0.44	98.75 ± 0.51	86.78 ± 2.34	0.57 ± 0.57	1.34 ± 0.73	4.11 ± 1.70	6.77 ± 1.41
Pre-Surgery		2.05 ± 0.14	3.72 ± 0.22	4.47 ± 0.33	95.80 ± 0.69	83.07 ± 2.41	0.51 ±0.30	1.50 ± 0.48	2.54 ± 0.77	14.80 ± 2.30
Post-Surgery	1	1.99 ± 0.15	3.82 ± 0.23	4.64 ± 0.36	95.89 ± 1.00	85.62 ± 1.95	0.00 ± 0.00	0.93 ± 0.31	2.18 ± 0.59	11.88 ± 1.53
	2	1.81 ± 0.12	3.81 ± 0.31	4.29 ± 0.44	97.17 ± 0.71	84.15 ± 1.45	0.17 ± 0.11	0.67 ± 0.24	2.25 ± 0.62	13.33 ± 1.44
	3	1.77 ± 0.16	3.43 ± 0.24	4.19 ± 0.40	96.58 ± 0.59	81.16 ± 2.43	0.08 ± 0.08	1.00 ± 0.28	1.25 ± 0.39	11.97 ± 1.70

Behavioural data are presented as mean ± SEM. Cells marked in bold indicate significant differences (compared to vehicle 0.0 mg/kg session for that drug).
*p<0.05, **p<0.01, ***p<0.001.