Intrinsic fluorescence in non-aromatic peptide structures is induced by collective vibrations, charge reorganisation and short hydrogen bonds, as shown in a new glutamine-related structure

Amberley D. Stephens,*[1] Muhammad Nawaz Qaisrani,*[2,3] Michael T. Ruggiero,[4,1] Saul T.E. Jones,[5] Emiliano Poli,[2] Andrew D. Bond,[6] Philippa J. Woodhams,[1] Elyse M. Kleist,[7] Luca Grisanti,[8] Ralph Gebauer,[2] J. Axel Zeitler,[1] Dan Credgington,[5] Ali Hassanali,[2#] Gabriele S. Kaminski Schierle[1#]

[1] Chemical Engineering and Biotechnology, University of Cambridge, Cambridge CB3 0AS, UK
[2] The Abdus Salam International Centre for Theoretical Physics, Strada Costiera 11, 34151 Trieste, Italy
[3] International School for Advanced Studies, Via Bonomea 265, 34136 Trieste, Italy
[4] Department of Chemistry, University of Vermont, 82 University Place, Burlington, VT, 05405 USA
[5] Cavendish Laboratory, University of Cambridge, JJ Thomson Avenue, Cambridge, CB3 0HE, U.K
[6] Department of Chemistry, University of Cambridge, Lensfield Road, Cambridge CB2 1EW, United Kingdom
[7] Department of Chemistry, Syracuse University, Syracuse, New York 13244, United States
[8] Division of Theoretical Physics, Ru¯der Boškovi´c Institute, Bijeni˘cka cesta 54, 10000 Zagreb, Croatia
* Contributed equally
# Corresponding authors: ahassana@ictp.it, gsk20@cam.ac.uk

Abstract

Disentangling the origin of the optical activity of non-aromatic proteins is challenging due to their size and thus their high computational requisites. Here we show, in a much smaller model system, that the single amino acid glutamine undergoes a chemical transformation leading to an unreported glutamine-like structure which has a similar broad absorption spectrum reported previously for non-aromatic proteins. We further show computationally that the optical activity of the glutamine-like structure is directly coupled to short-hydrogen bonds, but also displays charge and vibrational fluctuations, the latter of which are also present in less optically active structures such as in L-glutamine. Since experimentally the glutamine-like structure is the brightest structure, we conclude that short-hydrogen bonds are the ones responsible for the large Stokes shift observed in optically active non-aromatic proteins.

Introduction

Short peptides void of any aromatic residues have been shown to display an intrinsic fluorescence in the visible range (*1*, *2*). This has primarily been observed in fibrillar protein structures linked to neurodegenerative diseases, such as Alzheimer's, Parkinson's and Huntington's diseases (*3–6*). Furthermore, optical properties of double amino acid based nanowires have also been reported, existing either of two non-aromatic or two aromatic amino acids (*2*, *7*, *8*). We have shown that the fluorescence of non-aromatic short crystal structures forming part of the amyloid-beta protein is enhanced by the presence of short hydrogen bonds (SHB) between the termini of prefibrillar structures (*5*). The SHB permits proton transfer leading to a double-well ground state potential which we have proposed prevents a conical intersection in the excited state (*5*). It has been suggested that one of the prerequisites for this fluorescence observed in either amyloid structures or short peptide nanowires is related to hydrogen bonding or aromatic interlocks which, for the latter, decreases the bandgaps down to the semiconductive regions (*9*).

Despite our previous suggestion that proton delocalisation is strongly coupled to this intrinsic fluorescence, its direct role and more importantly, the role of other vibrational modes on putative fluorescing states, has not been elucidated. We have thus searched for a model system, such as a single amino acid-based structure, that displays similar optical properties to amyloid fibrils and is permissive to more sophisticated computational approaches. We have been inspired by the small peptide nanostructures that have been pioneered by the Gazit laboratory (*9*) and by the fact that there are several neurodegenerative diseases that have been connected with an increased level of glutamines produced as part of a protein, as for example Huntingtin in Huntington's disease which renders the protein more aggregation prone (*10*). It has been known that the amide group in L-glutamine (L-glu) is highly labile and thus can rapidly hydrolyse. We show here that the single amino acid L-glu upon heating in water can form a nanostructural material with optical properties similar to the ones observed in other amyloid fibrils such as in fibrils of amyloid-beta, alpha-synuclein or tau (*4*, *11*, *12*).

Using X-ray diffraction (XRD), we show that L-glu dissolved in water and upon heating

becomes cyclysed forming a previously unreported structure which resembles L-

pyroglutamine (which has been reported to be a component of amyloid-beta in the brain

(*13*)), but involves a low-barrier hydrogen bonded anionic dimer with an ammonium

counterion. We have termed the new structure, i.e. L-pyroglutamine complexed with an

ammonium ion, L-pyro-amm. L-pyro-amm has a microcrystalline plate morphology as shown

by scanning electron microscopy (SEM). The newly formed solid was further characterised

using terahertz time-domain spectroscopy (THz-TDS), which provides information on the

low-frequency modes in the crystal that control the proton transfer. Additionally, the

experiments were interpreted using ground and excited state electronic structure

calculations and molecular dynamics simulations. Ultimately, the combination of static

structural information, atomic vibrational dynamics, and optical properties enable the

origins of fluorescence in this particular structure to be elucidated, shedding light on the

complementary processes in more complex systems.

Methods

Experimental

Sample preparation of L-glutamine

L-glutamine (L-glu) (#G3126, #G8540, Sigma-Aldrich, Gillingham, UK) and L-pyroglutamine

(L-pyro) (#83160, Sigma-Aldrich) were dissolved in 18.2Ω MilliQ H_2O at a concentration of

0.3 M or 1 M. Aliquots were placed in a 65°C oven, since heating up proteins to 65°C

increases the formation of amyloid structures as reported previously (*14*). Each aliquot was

rotated to dissolve the powder once a day. Samples were either analysed in liquid form or

dried on a glass or quartz cover slip (#043210.KG, Alfa Aesar, Lancashire, UK) either at room

temperature (RT) or on a heat block set to 50°C.

Emission and excitation wavelength scans

Emission and excitation spectra were taken on a Hitachi F-4500 FL spectrophotometer

(Hitachi High-Technologies Corporation, Tokyo, Japan) at RT in a quartz cuvette. For

measurements, the excitation slit resolution was 5 nm or 10 nm and the emission slit

resolution was 20 nm. The PMT voltage was set at 950 V and the scan speed set at 240

nm/min. The excitation scan was measured between 250 – 400 nm and the emission filter

set to the emission maxima of the sample stated in the figure legend, with a slit resolution

of 20 nm. The emission scan was measured between 380 - 560 nm and the excitation filter

set to the excitation maxima of the sample stated in the figure legend, using a slit resolution

of 5 nm. Four measurements were taken for each sample which was repeated at least three

times and the background (air or H_2O) was subtracted from the average.

Absorption measurements

Absorption measurements were taken on a UV-Vis-NIR Spectrophotometer, UV-3600 Plus

(Shimadzu, Kyoto, Japan) and Cary 6000i (Agilent, Santa Clara, USA). 1 M or 0.3 M L-glu, L-

pyro or L-pyro-amm solutions were measured in 10 mm QX cuvettes (Hellma Analytics,

Müllheim, Germany) or dried on quartz coverslips. Measurements were taken between

wavelengths 200 – 800 nm using 1 nm steps at a slow scan speed and a 1 nm resolution. The

light source change wavelength was set at 393 nm and the grating change wavelength set at

750 nm. Samples were measured at least three times and the experiments repeated at least

three times, measurements were then averaged and H_2O or coverslip only control was

subtracted.

SEM (scanning electron microscopy)

SEM was performed using a FEI Magellan 400 HR-SEM at an acceleration voltage of 2 kV. L-

pyro-amm samples were lyophilised by freezing in liquid nitrogen and freeze drying in a

LyoQuest 85 (Telstar, Terrassa, Spain) and imaged on a glass coverslip.

X-ray diffraction

L-pyro-amm was dried on a glass coverslip in a 50°C oven and then at RT until crystals

formed. Single crystal X-ray diffraction (SCXRD) measurements were performed at 180 K

with a Bruker D8-QUEST PHOTON-100 diffractometer, which utilised a Cu Kα radiation (λ =

1.54 Å), and an APEX-II CCD. Absorption corrections were made using SDABS, and data

integration and reduction were performed with SAINT+. All non-hydrogen atoms were

refined isotropically and anisotropically, followed by inclusion of the hydrogen atoms

(determined using the excess electron density) and refinement isotropically.

Terahertz Time-Domain Spectroscopy

All THz-TDS spectra were acquired using a commercial Terapulse 4000 spectrometer (TeraView Ltd, Cambridge, UK). Samples were prepared for THz-TDS measurements by diluting the solid air dried L-pyro-amm with polyethylene (~ 10% w/w concentration) by gentle mixing using an agate mortar, followed by pressing into 2 mm thick, 13 mm diameter pellets using a hydraulic press. All THz-TDS spectra shown are a result of division of sample and blank datasets, with the blank dataset represented the THz-TDS response of a pellet of pure polyethylene.

Theoretical

DFT-THz Calculations

Calculations were performed using both the CRYSTAL17 (*15*) and Quantum Espresso (*16*) software packages. Geometry optimisations and vibrational analyses performed with the CRYSTAL17 code utilised the atom-centred split-valence triple-zeta 6-311g(2d,2p) basis set for all atom types. Based on a previous study related to ionic molecular crystals (*17*), the range-corrected WB97-X (*18*) functional was used. The vibrational analysis was performed within harmonic approximation, and infrared intensities were determined using the Berry Phase method. Energy convergence criteria were set to $\Delta E < 10^{-8}$ and 10^{-11} hartree for the geometry and frequency calculations, respectively.

Periodic TD-DFT Excited State Calculations

Simulations were performed using the fully periodic Quantum Espresso software package. The Becke-Lee-Yang-Parr (B3LYP) hybrid density functional was used with an energy cutoff of 40 Ry. The calculations of the excited state were performed within the framework of TDDFT using the Liouville-Lanczos formalism implemented in the freely available Quantum-Espresso package (*19*). In this approach, the optical spectra are obtained directly over the wide spectral range without taking into account the numerically complex calculations of the single exited states. We used plane wave basis set and the electron-ion interactions were taken into account via norm conserving Martins-Troullier pseudopotentials (*20*). To determine the ground state wave function, we used the gamma point of the Brillouin zone. All the periodic calculations employed the computationally demanding B3LYP (*21*) hybrid

functional, the kinetic energy cutoff of 40 Ry was used for the wave functions. The intrinsic

band width for the spectra was set to 0.003 Ry (~0.0408 eV).

Periodic Structure Geometry Optimisation

The structures obtained from the experiments were first geometrically optimized at $0°K$

using the Broyden-Fletcher-Goldfarb-Shanno (BFGS) minimisation algorithm implemented in

CP2K (*22, 23*) package. A convergence criterion for the wave function optimisation was used

as $5x10^{-7}$ a.u. Applying the method of the Gaussian and plane wave, the wave function was

expended in the Gaussian double-zeta valence polarised (DZVP) basis set. The cutoff for the

electronic density was set to 300 Ry. We used the gradient correction to the local density

approximation and the core electrons were treated via Goedecker-Teter-Hutter

pseudopotentials (*24*). In all the calculations, we used the Becke-Lee-Yang-Parr (BLYP) (*25*)

functional with the D3(0) Grimme (*26*) dispersion corrections for the van der Waals

interactions.

Ab Initio Molecular Dynamics Simulations

Ab initio Molecular Dynamics simulations (AIMD) were performed using Quickstep

algorithm implemented in CP2K. In these calculations, the propagation of the nuclei was

taken into account within the framework of the Born-Oppenheimer approximation. The

simulations were performed in the NVT ensemble and the temperature was controlled

during the simulations by using the velocity-rescaling thermostat (*27*). We used the time

step of 0.5 femtosecond to update the nuclear coordinates and velocities while the total

length of the simulations for each system is 50 picoseconds.

Excited State Cluster Calculations

A set of excited state calculations were performed on glutamine clusters in order to

understand the role of the environment on the optical properties. Specifically, the optical

properties of L-pyro-amm were investigated using various isolated cluster models with the

Gaussian09 software package. The clusters were extracted directly from the crystal

structure and used in various combinations (dimers, trimers, tetramers) to perform time-

dependent DFT (TD-DFT) calculations. A split-valence triple-zeta 6-311g(2d,2p) basis set was

used for all atom types together with the hybrid B3LYP functional. Some benchmark

simulations, comparing the optical properties obtained from the periodic calculations using
B3LYP to range corrected hybrid functionals like CAM-B3LYP, were also performed with
these clusters.

We also performed a series of excited state optimisations on various model systems built
from L-pyro-amm in order to examine the nature of the geometrical distortions that occur
on the lowest electronic excited state. These calculations were also performed with the
Gaussian09 software package. All clusters were surrounded with a continuum dielectric
constant of 80, representing pure H_2O. The 6-311G(2d,2p) basis set was used for all atoms
together with the range corrected hybrid functional CAM-B3LYP (28). The clusters were first
optimised in the ground state after which they were optimised on the first electronic
excited state.

Results and Discussion

It has long been known that poly-glutamine can form amyloid-like fibrillar structures *in vitro*.
The more glutamine residues in the poly-glutamine polymer, the faster the aggregation
propensity of the polypeptide chain. This led us to investigate whether L-glu on its own
under conditions which normally promote fibril formation, such as an increase in
temperature (14), was able to form structures with similar optical properties, as recently
observed for amyloid fibrils (5, 29, 30).

We first investigated the structure of L-pyro-amm, which formed after incubation of L-glu
for 8 days at 65°C, using SEM and observed crystal structures shown in Fig.1a. However, in
order to investigate whether L-glu had indeed changed its crystal structure arrangement we
performed XRD analysis of the resulting material. In Fig.1b we show the crystal structure of
the heated L-glu structure, which we termed L-pyro-amm, and the published crystal
structures of L-glu and L-pyroglutamine (L-pyro) in Fig. 1c and d. Note, the L-pyro structure
was analysed as it displayed structural similarities to the newly formed L-pyro-amm. Figures
were obtained from geometry optimisations of the nuclear positions of the atoms using
experimental densities. L-pyro-amm consists of 8 pyroglutamine groups and 4 ammonium
ions (144 atoms) complexed within the crystal (see Fig. 1b). In contrast, as shown in Fig. 1c,

L-glu consists of 4 glutamine molecules (80 atoms) in the unit cell which form hydrogen

bonds involving the termini and side chain. Furthermore, as shown in Fig. 1d, L-pyro consists

of 12 pyroglutamine molecules (192 atoms) in the unit cell forming hydrogen bonds

involving the NH and COOH groups.

L-pyro-amm has a rather unique hydrogen bond network structure since four of the

pyroglutamine molecules are deprotonated and hence have a nominal negative charge,

while the other four molecules are neutral. One of the important implications of this

difference is that L-pyro-amm contains a very strong hydrogen bond. The red circled region

in Fig. 1a corresponds to a short hydrogen bond (SHB) with a length of 2.45 Å, while those in

L-glu and L-pyro range between ~2.55-2.85 Å.

The structural change was further confirmed using THz-TDS measurements, as this

technique is strongly dependent on the bulk packing arrangement (as well as on the internal

covalent structure) of the molecules (*31*). The THz-TDS spectrum of the resulting solid, as

well as the solid-state DFT predicted spectrum based on the single crystal XRD (SCXRD)-

determined structure, is shown in Fig. 1e (full spectral assignment available in Suppl. Fig. 1).

The agreement between the experimental and theoretical spectra further supports that full

conversion of the sample occurs and thus enables additional investigations into the

structural and electronic properties of the material.

Figure 1. L-glu forms L-pyro-amm upon heating.

 (a) SEM image of crystals of L-pyro-amm dried. (b) XRD analysis of heated L-glu sample show the newly formed structure, L-pyro-amm. Geometry optimisations show that 8 pyroglutamine groups and four ammonium ions (144 atoms) are complexed in the crystal and a SHB of 2.45 Å (within red dashed lines) is present near the ammonium ion (linked by bonds highlighted by blue dotted lines) (white-hydrogen, red-oxygen, blue-nitrogen, grey-carbon). (c) Clusters of L-glutamine and (d) L-pyroglutamine. (e) Experimental (blue line) and theoretical (grey line) THz-TDS of the L-pyro-amm sample are in agreement and confirm the presence of the new L-pyro-amm structure.

We first investigated whether there were any differences in the optical properties associated with the three crystal structures. Comparing the absorption of L-glu, L-pyro and L-pyro-amm in water, we show that only L-pyro-amm has a significantly red-shifted absorption which lies in the 275-320 nm range, whereas both L-glu and L-pyro primarily absorb in the deep UV (<250 nm) (see Fig. 2a).

We next compared the experimental absorption spectra of L-glu, L-pyro and L-pyro-amm with the ones obtained from time dependent density functional theory (TDDFT). We highlight here, that the small size of the systems permitted us to determine the spectra

using a hybrid functional, thereby not only advancing the quality of our theoretical
predictions from previous studies (*5, 29, 30*) but also coupling the optical properties directly
to different vibrational modes.

Fig. 2b illustrates the absorption spectra obtained for the TDDFT calculations on the 3
periodic systems in the ground state (i.e. at 0 K). Panel b.i) shows the relative oscillator
strength as a function of the frequency while panel b.ii) illustrates the second derivative of
the oscillator strength permitting the positions of the maxima in the spectra to be more
easily identified. The spectra reveal some striking differences between the different
systems. Interestingly, we observe that L-pyro is essentially dark throughout the frequency
range up to ~ 6eV. On the other hand, L-pyro-amm shows the presence of more structure in
the spectrum. Specifically, it is the only system for which the spectrum features a low
energy excitation at 226 nm (5.5 eV) and subsequently other peaks slightly above 220 nm
(5.625eV) and 216 nm (5.75eV). While L-glu exhibits a peak at 222 nm (5.58eV), it is dark up
to 206 nm (~6eV). These calculations were performed with the B3LYP functional.
Importantly, we have found that upon using more accurate functionals such as CAM-B3LYP
(*28*) and WB97-X (*18*) L-pyro-amm remains the most optically active.

We have previously shown that thermal fluctuations and in particular nuclear vibrations,
such as proton transfer, have a large impact on the absorption spectra of peptide structures
compared to absorption spectra at 0 K (*5, 32–34*). In Fig. 2c we show that, compared to the
0 K spectra in Fig. 2b, thermal fluctuations cause a large red shift to around 3.4 eV (365 nm)
for L-pyro-amm, close to what is observed experimentally. These spectra were computed
averaging over 25 frames sampled from the molecular dynamics simulations. Interestingly,
no such effect is observed for L-glu which remains weakly absorbing up to more than 5 eV
(247 nm) as seen at 0 K.

Similar to our previous studies on the intrinsic fluorescence of amyloid-beta fibrils,
absorption is significantly increased in structures containing SHB, as neither L-glu, nor L-pyro
display a significantly red shifted absorption. In order to understand better the physical
origin of the low energy excitation at 226 nm (~5.5 eV) in L-pyro-amm, we computed the

electron response density at this frequency. This is illustrated in Fig. 2d, where we observe

that most of the electron response involves regions around the pyroglutamine rings as well

as regions near the SHB (see dashed circle in Fig. 2d). The optical response thus involves a

collective charge reorganisation involving several parts of the molecular crystal.

We next investigated whether the above structures also display fluorescence excitation and

emission properties as has been observed for amyloid-like structures reported previously (*5,*

29, 30). Fig. 2e shows the excitation scan from 250-400 nm (solid lines) with the emission

set at 430 nm of L-glu in water at day 0 to 8 after incubation at 65°C. Interestingly, we

observe an excitation peak at around 360 nm which is similar to what we have measured

previously for amyloid proteins (*5*). The corresponding emission scan (dashed lines) with the

excitation set at 360 nm and emission from 380-560 nm showed an emission peak around

430 nm, again lying in the same visible range as for amyloid fibrils. When the L-pyro-amm

solution was dried the excitation and emission peaks were slightly blue shifted (Suppl. Fig.

2a) which may be due to a change in the molecular environment in the dried state.

Importantly, we do not see any fluorescence in L-glu (without heating, i.e. at day 0 Fig. 2e.).

To determine the importance of the ammonium ion experimentally, L-pyro has been

incubated in water and heated at 65°C for 8 days, and only a very weak fluorescence has

been detected (Suppl. Fig. 2b).

Figure 2. Optical properties of L-pyro-amm are distinct from L-glu and L-pyro

(a) Absorption spectra of 0.3 M L-glu (black), L-pyro (blue) and L-pyro-amm (red) (L-glu incubated for 8 days at 65°C) in water taken between 200 – 500 nm shows primarily features of L-pyro-amm. (b) Absorption spectra of L-glu, L-pyro and L-pyro-amm obtained from periodic density functional theory calculations with the B3LYP functional. L-pyro-amm features the lowest lying excited states which are characterised by the largest oscillator strengths. (c) Absorption spectra for L-glu and L-pyro-amm obtained from periodic simulations at room temperature. The spectra were computed by averaging over 25 frames randomly sampled from the *ab initio* molecular dynamics simulations. (d) The excited state electron density computed for L-pyro-amm from the optimised structure computed at the first peak (arrow in panel b). The lowest excited state density shows a response from various parts of the crystal structure including the pyroglutamic acid ring and the SHB region (see

dashed circle). The orange and green surfaces correspond to regions involving a decrease

and increase in electron density respectively, shown at an iso-value of $1x10^{-5}$. (e) 1 M L-glu

was incubated at 65°C and the excitation and emission spectra were measured over time.

Excitation spectra were measured between 250-400 nm with emission set at 420 nm and

emission spectra were measured between 380-560 nm with the excitation set at 360 nm.

It is not possible to directly pinpoint the exact mechanisms and origins of the difference in

the optical properties between the three systems from the ground state calculations at 0 K.

As alluded to earlier, one of the factors that distinguishes L-pyro-amm from the other

systems is the presence of the SHB (highlighted by red circles in Fig. 1a) and the presence of

the ammonium ion. In order to characterise the behaviour of the SHB, we conducted *ab*

initio molecular dynamics simulations of the three systems at 300 K and examined the

proton transfer coordinates defined as the difference in distance between the proton (H)

and the two oxygen atoms (O1 and O2) that sandwich it and is commonly referred to as the

proton transfer coordinate (d_{O1-H}-d_{O2-H}) as shown in Fig. 4a for different types of hydrogen

bonds in the crystals. It is clear that the SHB in L-pyro-amm is characterised by a double-well

potential. The barrier associated with this proton transfer is on the order of thermal energy,

indicating that zero-point energy (ZPE) would make the proton transfer barrierless (*35*). An

examination of similar proton transfer coordinates for hydrogen bonds in L-glu and L-pyro

show that they are characterised by only single-well potentials.

The nature of the optical properties is sensitive to the environment in which the glutamine

molecules reside. It has previously been reported that charged amino acids already display

an absorption in the range of 250-350 nm that is significantly red shifted (*36, 37*). The

origins of the low energy absorption were attributed to charge transfer excitations. The

simulations of these systems were performed in the gas phase, rather than considering the

protein environment such as shown for L-pyro-amm in Fig 2d. In comparison to the results

presented in Fig. 2d, data presented in Fig. 3, show that the origins of the electronic

transitions equally arise from a charge transfer (CT) between the highest occupied

molecular orbital (HOMO) on the anionic dimer, and the lowest unoccupied molecular

orbital (LUMO) centred on the ammonium cation when performed in the gas phase.

Interestingly, the correct transition energy is only predicted when the ammonium cation is

spatially near the centre of the dimer, which corresponds to the delocalisation of the

negative charge and the SHB. Two generalised geometries, with the ammonium cation near

the SHB (as seen in Fig. 3a) and away from the SHB (Fig. 3b), with the corresponding HOMO

and LUMO orbitals are shown. The results predict a transition of 303.50 nm for dimer a. and

668.66 nm for dimer b., with dimer a. most closely resembling the chemical environment

present within the crystalline material.

The results show that charge transfer is capable to lead to absorption in the far UV when

investigated in the gas phase, i.e. neglecting the direct protein environment. However,

including the protein environment in the molecular crystal (Fig. 2d) results instead in the

excitation being a charge reorganisation involving several different molecular groups of the

crystal. Indeed, by shuffling the protons along the SHBs in the ground-state, we observe an

electronic response involving the entire structural units of L-pyro-amm including both the

hydrogen bonded regions and the pyroglutamic acid rings when the protein environment is

accounted for (Suppl. Fig.3b i-iv).

Figure 3. Comparison of the HOMO and LUMO orbitals on two L-Pyro-Amm models.

L-pyro-Amm structures are presented where the ammonium cation is located directly near

the hydrogen bond (a), and where the ammonium cation is located away from the hydrogen

bond (b). While both models predicted charge-transfer HOMO-LUMO states, only in the

case of (a) is the transition predicted to be in the vicinity of the experimentally observed

peak, 303.50 nm compared to 668.66 nm for (b).

Up to this point, we have shown that the vibrations of protons along SHBs are an important
part of the structural fluctuations in the ground state structure of L-pyro-amm. These
calculations however, do not say anything about the nuclear relaxation upon photo-
excitation which is most relevant for fluorescence. In a final set of theoretical studies, to
ascertain the role of the proton transfer in the ground state, as well as to elucidate the
presence of other types of vibrational modes, we performed a series of geometry
optimisations on the first excited state of a cluster carved out from L-glu, L-pyro and L-pyro-
amm crystals and surrounded by a continuum dielectric constant of 80.

Figure 4. Collective vibrations affect the emission in L-pyro-amm

a) Shows the proton transfer free energy profiles along several hydrogen bonds in L-glu, L-pyro and L-pyro-amm obtained from ground-state ab initio molecular dynamics simulations. Note, the SHB in L-pyro-amm is characterised by a double-well potential. b) Scatter plot of the oscillator strengths versus the emission energy (defined as the difference between the first excited state and the ground-state) during the excited optimisations for L-glu, L-pyro and L-pyro-amm. c) Scatter plot of the oscillator strengths versus the distance of the carbonyl bond (C=O) that extends during the excited state optimisation in L-glu, L-pyro and L-pyro-amm. d)-f) Snapshots of three different systems of L-pyro-amm that were optimised on the excited state. In clusters shown in d) and e) the initial position of the proton along the SHB has been changed, and in f) the ammonium ion has been placed closer to the SHB region. The bottom panel of d), e) and f) show zoomed in plots of the main regions of the cluster (circled in the top panels) that undergo significant changes. g)-i) Ground and excited state energies are plotted as a function of the excited state optimisation for the three systems shown in panels d-f). The curves on the excited state are colour-coded with the oscillator strengths. S_0 and S_1 refer to the ground and excited state energies, respectively. The right-side of the y-axis corresponds to C=O bond lengths (dashed black lines) for g) and h). In panel i), where there is a proton transfer on the excited state, the dashed blue lines correspond to the O-H distance along the SHB on the same distance scale.

Fig. 4b shows a scatter plot of the difference between the first excited state and ground-state energies and the corresponding oscillator strengths. The scatter plots were obtained over the course of the excited state optimisation. We observe that the L-pyro-amm system is characterised by the largest oscillator strengths compared to L-glu or L-pyro. One of the major structural changes that occurs upon excitation is an increase in the C=O bond length similar to observations in previous studies (*38–40*). Fig. 4c shows the oscillator strengths as a function of the C=O bond, that increases in length by about 0.1 Angstroms.

Although all three glutamine-related structures display similar trends on the excited state, what makes L-pyro-amm unique is the presence of the SHB. Fig. 4b and c show that L-pyro-amm displays the largest oscillator strength peaking at approximately 3.5eV consistent with our experimental findings. Although L-glu also has a peak at around 3.5 eV it is much weaker

than the one of L-pyro-amm. We thus decided to focus on a series of excited state

optimisations for various clusters of L-pyro-amm. In particular, we focused on the proton

transfer in the ground state as seen in the double-well potential and on the proximity of the

ammonium ion to the short hydrogen bond region, all shown to play an important role in

the optical properties of L-pyro-amm so far. We thus constructed three clusters with

different initial conditions, which are shown in the top panels of Fig. 4d-f. The clusters

shown in Fig. 4d and 4e differ in the initial positions of the protons along the SHB. As seen in

Fig. 4a, the ground-state simulations of L-pyro-amm present a double-well potential in the

finite temperature simulations. We thus moved the proton along one of the SHBs,

constrained the geometry in the ground-state which yielded the cluster shown in Fig. 4e. In

Fig. 4f, the ammonium ion is placed closer to the SHB region.

The circled regions in Fig. 4d-f highlight the main regions, where the nuclear degrees of

freedom respond in the excited state. The bottom panels show a zoomed in image of the

changes that occur upon relaxation, for which the solid and transparent structures

correspond to geometries obtained before and after the relaxation on the excited state,

respectively. The bottom panel of Fig. 4d illustrates that the change involves the C=O stretch

and deplanarisation of the peptide dihedral angle of the ring. The corresponding evolution

of the first excited state and ground state energies as a function of optimisation is shown in

Fig. 4g. The relaxation on the excited state narrows the energy difference between the

ground and excited state to approximately 3.5eV which is consistent with the experiments.

The top panels of Fig. 4e and f involve two other initial conditions for which the proton

vibrations and the proximity of the ammonium ion to the SHB are altered. We see in Fig. 4h

and i that the magnitude of the oscillator strength of L-pyro-amm in the excited state is

sensitive to these new initial conditions. Transferring the protons on the ground-state (see

dotted black rectangles in Fig. 4d-f), significantly changes the oscillator strengths as seen in

the comparison between Fig. 4g-i, across the entire relaxation process in the first excited

state while still maintaining the C=O stretch and deplanarisation as seen in the bottom

panel of Fig. 3e. In the case of Fig. 4f, besides the reorganisation involving the C=O stretch,

we also observe a proton transfer along the SHB. This is seen in Fig. 4i, which shows the O-H

stretch (dashed-blue curve) along the SHB during the optimisation which increases from

1.26 to 1.32 Angstroms. The zoomed in plot in the bottom panel of Fig. 4f shows this subtle

change in the proton transfer coordinate. Thus, the evolution of the excited state in L-pyro-

amm into the region displaying visible fluorescence appears to involve a collective

reorganisation of various vibrational modes forming the hydrogen bond network with the

SHB playing a particular role, as it leads to the brightest structure, also confirmed

experimentally. These findings are also consistent with some recent theoretical work

showing the importance of proton transfer in self-assembled peptides made of aromatic

phenyl-alanine amino acids (*41*). However, the latter structure features an aromatic amino

acid, whereas we observe similar optical properties in non-aromatic structures.

Conclusions

The experimental and theoretical findings presented here, elucidate a rather complex

molecular mechanism associated with the non-aromatic intrinsic fluorescence in protein-

like structures. In the case of L-glutamine, a chemical reaction creates a newly formed

structure involving a cyclised pyroglutamic acid ring. This new structure features absorption

in the UV and emission in the visible range very similar to the chemically distinct amyloid

fibrils (*4, 5, 11, 12, 41–43*).

The structural chromophore responsible for the optical properties in this new protein-

related structure arises from a hydrogen bond network characterised by specific vibrational

fluctuations on the excited state. The presence of strong hydrogen bonds along which

proton transfer occurs and secondly, specific ionic interactions in close proximity, such as

involving the ammonium ion, affect the optical properties. Although the fluorescence

observed in these systems is much weaker than those compared to conventional

fluorophores, the physical and chemical properties of the hydrogen bond network reported

here maybe a generic feature across many other peptide structures.

In summary, we show here that there are several vibrational modes in biological structures

that contribute to fluorescence properties as shown by excited state simulations and

confirmed experimentally. We further show that the protein environment strongly

contributes to the optical activity of the protein like structure and that the SHB significantly

enhances the fluorescence signal detected, potentially by inhibiting a conical intersection in
the excited state as discussed previously (*5*). Interestingly, SHBs have recently been
observed in different biological systems which have long been associated with either
intrinsic fluorescence, such as NADP/NAD (*44*), FAD/FMN (*45*), the light-sensing
chromophore in photoactive yellow protein (*46*), or in the active site of many enzymes, such
as hydrolases and oxidoreductases (*47, 48*), many of which consist of a highly complex H-
bond structure similar to amyloid proteins. Our findings may thus further the design of
novel optically active biomaterial for applications in optical sensing or the design of novel
biocompatible catalysts.

Author Information

Corresponding Authors

Ali A. Hassanali, ahassana@ictp.it

Gabriele S. Kaminski Schierle, gsk20@cam.ac.uk

Author Contributions

*A.D.S and M.N.Q contributed equally. A.D.S, prepared samples for all experimental data.
P.J.W. performed SEM experiments. A.D.B and M.T.R performed XRD measurements and
analysed data. M.T.R. performed THz experiments and DFT-THz calculations and analysed
data. A.D.S. performed excitation and emission measurements and analysed data. A.D.S and
S.T.J. performed absorption measurements and analysed data. M.T.R and E.M.K performed
TD-DFT cluster calculations. M.N.Q, E.P, L.G, R.G and A.H performed AIMD, Periodic TD-DFT
Excited State Calculations and Periodic Structure Geometry Optimisation calculations. A.D.S,
M.N.Q, M.T.R, S.T.J, L.G, J.A.Z, D.C, A.H and G.S.K.S contributed to manuscript writing. All
authors have given approval to the final version of the manuscript.

Notes

The authors declare no competing financial interest.

Acknowledgments

G.S.K.S. acknowledges funding from the Wellcome Trust, the UK Medical Research Council
(MRC), Alzheimer Research UK (ARUK), and Infinitus China Ltd. A.D.S. acknowledge

Alzheimer Research UK the British Biophysical Society (BSS) for travel grants. P.J.W

acknowledges EPSRC funding (EP/L016087/1).

535 **Supplementary data**

Mode Number	Frequency (THz)	Intensity (km mol⁻¹-1)	Assignment
1	0.955	0.22	Antisymmetric translation in b
2	1.215	0.71	Coupled translation in a and rotation about bc
3	1.325	1.52	external rotation about b
4	1.450	0.36	Asymmetric translation in c
5	1.554	4.37	External rotation about a
6	1.676	1.76	Translation in a and external asymmetric rotation about H-bond coordinate
7	1.805	2.28	External asymmetric rotation of dimer pairs
8	1.929	2.11	External symmetric rotation of dimer pairs
9	1.950	1.28	Asymmetric rotation about h-bond coordinate, ammonium translational motion
10	2.113	3.53	Translation and rotation (breathing) around ammonium cation
11	2.208	2.5	Symmetric external rotation of entire formula units
12	2.404	2.37	In phase external rotation perpendicular to h-bond coordinate
13	2.502	0.45	Out of phase external rotataion perpendicular to h-bond coordinate
14	2.564	0.73	External out of phase rotation of h-bonded chains about b
15	2.565	8.91	Out of phase external rotation of individual pyroglutamic molecules with translation of ammonium cation
16	2.654	18.09	External vibration coupled with torsion of the COOH group
17	2.679	0.29	External rotation and torsion of pyroglutamic ring
18	2.776	4.94	External rotation and torsion of ring and carboxyl group
19	2.988	2.68	External rotation and translation of ammonium

Supplementary Figure 1. Full spectral assignment of THz data.

The top chart shows the contribution to each IR-active mode including external translations

and hindered rotations, and internal vibrational motions (i.e. torsions), while the bottom

table lists the detailed assignment for each mode.

Supplementary Figure 2. L-pyro-amm has blue shifted fluorescence when dried and

displays higher fluorescence intensity than L-pyro.

(a)The excitation peak of L-pyro-amm when dried (solid dark red line) is blue shifted with a

peak maximum ~ 340 nm compared to L-pyro-amm in H_2O (solid red line) which has a peak

maximum ~ 360 nm. The emission peak of L-pyro-amm when dried (dashed dark red line) is

also blue shifted, with a peak maximum ~ 420 nm, compared to L-pyro-amm in H_2O (dashed

red line) with a peak maximum ~ 430 nm. (b) 1 M L-glu and 1 M L-pyro (blue) were

incubated in H_2O for 8 days at 65°C. After 9 days, the L-glu had converted into the L-pyro-

amm structure (red). L-pyro-amm has a clear excitation peak maximum at ~360 nm and

emission peak maximum at ~ 430 nm, while L-pyro (blue), although not completely dark, has

no clear excitation or emission peak.

Supplementary Figure 3. The optical properties of L-pyro-amm are sensitive to the environment and involves the electronic response of the entire structure.

(a)(i) L-pyro-amm cluster used to examine the sensitivity of the optical response on different parts of the cluster upon moving different protons. (ii) Charge differences between the ground and excited state are computed using restrained electrostatic potential atomic partial charges (RESP) for the cluster shown in i). Note, the electronic response involves all the atoms of the cluster. (iii) The two protons in the rectangle regions are displaced to be in the centre of the hydrogen bond and the charge differences are then computed. As illustrated, the proton displacement leads to a larger change in magnitude of the charges. (iv) The charge differences are computed for another nuclear configuration for which the protons are kept fixed but the O—O distance is increased from 2.45 to 3.2 Angstroms. The charge differences obtained here are quite similar to the original condition shown in (ii).

References:

1.   A. Shukla, S. Mukherjee, S. Sharma, V. Agrawal, K. V. R. Kishan, P. Guptasarma, A
novel UV laser-induced visible blue radiation from protein crystals and aggregates:
Scattering artifacts or fluorescence transitions of peptide electrons delocalized
through hydrogen bonding? *Arch. Biochem. Biophys.* **428**, 144–153 (2004).

2.   G. Rosenman, N. Amdursky, M. Molotskii, D. Aronov, L. Adler-Abramovich, E. Gazit,
Blue luminescence based on quantum confinement at peptide nanotubes. *Nano Lett.*
9, 3111–3115 (2009).

3.   F. T. Chan, D. Pinotsi, G. S. K. Schierle, C. F. Kaminski, in *Bio-nanoimaging: Protein*
Misfolding and Aggregation (Elsevier, 2013;
http://linkinghub.elsevier.com/retrieve/pii/B9780123944313000134), pp. 147–155.

4.   F. T. S. Chan, G. S. Kaminski Schierle, J. R. Kumita, C. W. Bertoncini, C. M. Dobson, C. F.
Kaminski, C. V Robinson, C. M. Dobson, J. Beard, P. Das, K. Jansen, M. DeLucia, W.-L.
Lin, G. Dolios, R. Wang, C. B. Eckman, D. W. Dickson, M. Hutton, J. Hardy, T. Golde,
Protein amyloids develop an intrinsic fluorescence signature during aggregation.
Analyst. **138**, 2156 (2013).

5.   D. Pinotsi, L. Grisanti, P. Mahou, R. Gebauer, C. F. Kaminski, A. Hassanali, G. S.
Kaminski Schierle, Proton Transfer and Structure-Specific Fluorescence in Hydrogen
Bond-Rich Protein Structures. *J. Am. Chem. Soc.* **138**, 3046–3057 (2016).

6.   J. Pansieri, V. Josserand, S. J. Lee, A. Rongier, D. Imbert, M. M. Sallanon, E. Kövari, T.
G. Dane, C. Vendrely, O. Chaix-Pluchery, M. Guidetti, J. Vollaire, A. Fertin, Y. Usson, P.
Rannou, J. L. Coll, C. Marquette, V. Forge, Ultraviolet–visible–near-infrared optical
properties of amyloid fibrils shed light on amyloidogenesis. *Nat. Photonics.* **13**, 473–
479 (2019).

7.   L. Adler-Abramovich, E. Gazit, Correction: The physical properties of supramolecular
peptide assemblies: From building block association to technological applications.
Chem. Soc. Rev. **43**, 7236–7236 (2014).

8.   L. Adler-Abramovich, M. Reches, V. L. Sedman, S. Allen, S. J. B. Tendler, E. Gazit,
Thermal and chemical stability of diphenylalanine peptide nanotubes: Implications for
nanotechnological applications. *Langmuir.* **22**, 1313–1320 (2006).

9.   K. Tao, P. Makam, R. Aizen, E. Gazit, Self-assembling peptide semiconductors. *Science*
(80-.). **358** (2017), p. eaam9756.

10.   J. M. Andresen, J. Gayán, L. Djoussé, S. Roberts, D. Brocklebank, S. S. Cherny, L. R.

Cardon, J. F. Gusella, M. E. Macdonald, R. H. Myers, D. E. Housman, N. S. Wexler, J.

Lorimer, J. Porter, F. Gomez, C. Moskowitz, K. P. Gerstenhaber, E. Shackell, K. Marder,

G. Penchaszadeh, S. A. Roberts, A. Brickman, D. Brocklebank, J. Gray, S. R. Dlouhy, S.

Wiktorski, M. E. Hodes, P. M. Conneally, J. B. Penney, J. F. Gusella, J. H. Cha, M.

Irizarry, D. Rosas, S. Hersch, Z. Hollingsworth, A. B. Young, D. E. Housman, M. M. de

Young, E. Bonilla, T. Stillings, A. Negrette, S. R. Snodgrass, M. D. Martinez-Jaurrieta,

M. A. Ramos-Arroyoh, J. Bickham, J. S. Ramos, F. Marshall, I. Shoulson, G. J. Rey, A.

Feigin, N. Arnheim, A. Acevedo-Cruz, L. Acosta, J. Alvir, K. Fischbeck, L. M. Thompson,

A. Young, L. Dure, C. J. O'Brien, J. Paulsen, S. P. Moran, D. Krch, P. Hogarth, D. S.

Higgins, B. Landwehrmeyer, M. R. Hayden, E. W. Almqvist, R. R. Brinkman, O.

Suchowersky, A. Durr, C. C. Dodé, F. Squitieri, P. J. Morrison, M. Nance, C. A. Ross, R.

L. Margolis, A. Rosenblatt, G. T. Estrella, D. M. Cabrero, R. J. A. Trent, E. McCusker, A.

Novelletto, M. Frontali, J. S. Paulsen, R. Jones, A. Zanko, T. Ashizawa, A. Lazzarini, J. L.

Li, V. C. Wheeler, A. L. Russ, G. Xu, J. S. Mysore, T. Gillis, M. Hakky, L. A. Cupples, M.

Saint-Hilaire, S. M. Hersch, The relationship between CAG repeat length and age of

onset differs for Huntington's disease patients with juvenile onset or adult onset.

Ann. Hum. Genet. **71**, 295–301 (2007).

11.   D. Pinotsi, A. K. Buell, C. M. Dobson, G. S. Kaminski Schierle, C. F. Kaminski, A Label-

Free, Quantitative Assay of Amyloid Fibril Growth Based on Intrinsic Fluorescence.

ChemBioChem. **14**, 846–850 (2013).

12.   T. N. Tikhonova, N. R. Rovnyagina, A. Y. Zherebker, N. N. Sluchanko, A. A. Rubekina, A.

S. Orekhov, E. N. Nikolaev, V. V. Fadeev, V. N. Uversky, E. A. Shirshin, Dissection of the

deep-blue autofluorescence changes accompanying amyloid fibrillation. *Arch.*

Biochem. Biophys. **651**, 13–20 (2018).

13.   H. Mori, K. Takio, M. Ogawara, D. J. Selkoe, Mass spectrometry of purified amyloid β

protein in Alzheimer's disease. *J. Biol. Chem.* **267**, 17082–17086 (1992).

14.   M. Fändrich, C. M. Dobson, The behaviour of polyamino acids reveals an inverse side

chain effect in amyloid structure formation. *EMBO J.* **21**, 5682–90 (2002).

15.   R. Dovesi, A. Erba, R. Orlando, C. M. Zicovich-Wilson, B. Civalleri, L. Maschio, M. Rérat,

S. Casassa, J. Baima, S. Salustro, B. Kirtman, Quantum-mechanical condensed matter

simulations with CRYSTAL. *Wiley Interdiscip. Rev. Comput. Mol. Sci.* **8**, e1360 (2018).

16.   P. Giannozzi, O. Andreussi, T. Brumme, O. Bunau, M. Buongiorno Nardelli, M.

Calandra, R. Car, C. Cavazzoni, D. Ceresoli, M. Cococcioni, N. Colonna, I. Carnimeo, A.

Dal Corso, S. De Gironcoli, P. Delugas, R. A. Distasio, A. Ferretti, A. Floris, G. Fratesi, G.

Fugallo, R. Gebauer, U. Gerstmann, F. Giustino, T. Gorni, J. Jia, M. Kawamura, H. Y. Ko,

A. Kokalj, E. Kücükbenli, M. Lazzeri, M. Marsili, N. Marzari, F. Mauri, N. L. Nguyen, H.

V. Nguyen, A. Otero-De-La-Roza, L. Paulatto, S. Poncé, D. Rocca, R. Sabatini, B. Santra,

M. Schlipf, A. P. Seitsonen, A. Smogunov, I. Timrov, T. Thonhauser, P. Umari, N. Vast,

X. Wu, S. Baroni, Advanced capabilities for materials modelling with Quantum

ESPRESSO. *J. Phys. Condens. Matter*. **29** (2017), doi:10.1088/1361-648X/aa8f79.

17.   M. T. Ruggiero, J. Gooch, J. Zubieta, T. M. Korter, Evaluation of Range-Corrected

Density Functionals for the Simulation of Pyridinium-Containing Molecular Crystals. *J.*

Phys. Chem. A. **120**, 939–947 (2016).

18.   J. Da Chai, M. Head-Gordon, Long-range corrected hybrid density functionals with

damped atom-atom dispersion corrections. *Phys. Chem. Chem. Phys.* **10**, 6615–6620

(2008).

19.   D. Rocca, R. Gebauer, Y. Saad, S. Baroni, Turbo charging time-dependent density-

functional theory with Lanczos chains. *J. Chem. Phys.* **128** (2008),

doi:10.1063/1.2899649.

20.   N. Troullier, J. L. Martins, Efficient pseudopotentials for plane-wave calculations.

Phys. Rev. B. **43**, 1993–2006 (1991).

21.   A. D. Becke, Density-functional thermochemistry. III. The role of exact exchange. *J.*

Chem. Phys. **98**, 5648–5652 (1993).

22.   J. D. Head, M. C. Zerner, A Broyden-Fletcher-Goldfarb-Shanno optimization procedure

for molecular geometries. *Chem. Phys. Lett.* **122**, 264–270 (1985).

23.   J. VandeVondele, M. Krack, F. Mohamed, M. Parrinello, T. Chassaing, J. Hutter,

Quickstep: Fast and accurate density functional calculations using a mixed Gaussian

and plane waves approach. *Comput. Phys. Commun.* **167**, 103–128 (2005).

24.   S. Goedecker, M. Teter, Separable dual-space Gaussian pseudopotentials. *Phys. Rev.*

B - Condens. Matter Mater. Phys. **54**, 1703–1710 (1996).

25.   A. D. Becke, Density-functional exchange-energy approximation with correct

asymptotic behavior. *Phys. Rev. A*. **38**, 3098–3100 (1988).

26.   S. Grimme, J. Antony, S. Ehrlich, H. Krieg, A consistent and accurate *ab initio*

parametrization of density functional dispersion correction (DFT-D) for the 94

elements H-Pu. *J. Chem. Phys.* **132**, 154104 (2010).

27.   G. Bussi, D. Donadio, M. Parrinello, Canonical sampling through velocity rescaling. *J.*

Chem. Phys. **126** (2007), doi:10.1063/1.2408420.

28.   T. Yanai, D. P. Tew, N. C. Handy, A new hybrid exchange-correlation functional using

the Coulomb-attenuating method (CAM-B3LYP). *Chem. Phys. Lett.* **393**, 51–57 (2004).

29.   K. Jong, Y. Taghipour Azar, L. Grisanti, A. D. Stephens, S. T. E. Jones, D. Credgington, G.

S. Kaminski Schierle, A. A. Hassanali, Low Energy Optical Excitations as an Indicator of

Structural Changes Initiated at the Termini of Amyloid Protein. *Phys. Chem. Chem.*

Phys. (2019), doi:10.1039/c9cp04648h.

30.   L. Grisanti, D. Pinotsi, R. Gebauer, G. S. Kaminski Schierle, A. A. Hassanali, A

computational study on how structure influences the optical properties in model

crystal structures of amyloid fibrils. *Phys. Chem. Chem. Phys.* **19**, 4030–4040 (2017).

31.   E. P. J. Parrott, J. A. Zeitler, Terahertz time-domain and low-frequency Raman

spectroscopy of organic materials. *Appl. Spectrosc.* **69**, 1–25 (2015).

32.   Y. K. Law, A. A. Hassanali, Role of Quantum Vibrations on the Structural, Electronic,

and Optical Properties of 9-Methylguanine. *J. Phys. Chem. A.* **119**, 10816–10827

(2015).

33.   S. Sappati, A. Hassanali, R. Gebauer, P. Ghosh, Nuclear quantum effects in a

HIV/cancer inhibitor: The case of ellipticine. *J. Chem. Phys.* **145** (2016),

doi:10.1063/1.4968046.

34.   Y. K. Law, A. A. Hassanali, The importance of nuclear quantum effects in spectral line

broadening of optical spectra and electrostatic properties in aromatic chromophores.

J. Chem. Phys. **148** (2018), doi:10.1063/1.5005056.

35.   D. Marx, Erratum: Proton transfer 200 years after von grotthuss: Insights from Ab

initio simulations (ChemPhysChem (2006) 7 (1848-1810)). *ChemPhysChem.* **8**, 209–

210 (2007).

36.   S. Prasad, I. Mandal, S. Singh, A. Paul, B. Mandal, R. Venkatramani, R. Swaminathan,

Near UV-Visible electronic absorption originating from charged amino acids in a

monomeric protein. *Chem. Sci.* **8**, 5416–5433 (2017).

37.   I. Mandal, S. Paul, R. Venkatramani, Optical backbone-sidechain charge transfer

transitions in proteins sensitive to secondary structure and modifications. *Faraday*

698    *Discuss.* **207**, 115–135 (2018).

699 38. M. Mališ, Y. Loquais, E. Gloaguen, H. S. Biswal, F. Piuzzi, B. Tardivel, V. Brenner, M.

700    Broquier, C. Jouvet, M. Mons, N. D. S. Došlić, I. Ljubić, Unraveling the mechanisms of

701    nonradiative deactivation in model peptides following photoexcitation of a

702    phenylalanine residue. *J. Am. Chem. Soc.* **134**, 20340–20351 (2012).

703 39. B. P. Fingerhut, K. E. Dorfman, S. Mukamel, Monitoring nonadiabatic dynamics of the

704    RNA base uracil by UV pump-IR probe spectroscopy. *J. Phys. Chem. Lett.* **4**, 1933–

705    1942 (2013).

706 40. M. Sapunar, T. Ayari, N. Došlić, Comparative study of the photodynamics of

707    malonaldehyde and acetylacetone. *Chem. Phys.* **515**, 622–627 (2018).

708 41. S. K. Joseph, N. Kuritz, E. Yahel, N. Lapshina, G. Rosenman, A. Natan, Proton-Transfer-

709    Induced Fluorescence in Self-Assembled Short Peptides. *J. Phys. Chem. A.* **123**, 1758–

710    1765 (2019).

711 42. A. Handelman, N. Kuritz, A. Natan, G. Rosenman, Reconstructive Phase Transition in

712    Ultrashort Peptide Nanostructures and Induced Visible Photoluminescence.

713    *Langmuir.* **32** (2016), pp. 2847–2862.

714 43. L. L. Del Mercato, P. P. Pompa, G. Maruccio, A. Della Torre, S. Sabella, A. M.

715    Tamburro, R. Cingolani, R. Rinaldi, Charge transport and intrinsic fluorescence in

716    amyloid-like fibrils. *Proc Natl Acad Sci U S A.* **104**, 18019–18024 (2007).

717 44. A. Stavrinides, E. C. Tatsis, L. Caputi, E. Foureau, C. E. M. Stevenson, D. M. Lawson, V.

718    Courdavault, S. E. O'Connor, Structural investigation of heteroyohimbine alkaloid

719    synthesis reveals active site elements that control stereoselectivity. *Nat. Commun.* **7**

720    (2016), doi:10.1038/ncomms12116.

721 45. F. Forneris, D. P. H. M. Heuts, M. Delvecchio, S. Rovida, M. W. Fraaije, A. Mattevi,

722    Structural analysis of the catalytic mechanism and stereoselectivity in Streptomyces

723    coelicolor alditol oxidase. *Biochemistry.* **47**, 978–985 (2008).

724 46. S. Yamaguchi, H. Kamikubo, K. Kurihara, R. Kuroki, N. Niimura, N. Shimizu, Y.

725    Yamazaki, M. Kataoka, Low-barrier hydrogen bond in photoactive yellow protein.

726    *Proc. Natl. Acad. Sci. U. S. A.* **106**, 440–444 (2009).

727 47. W. W. Cleland, P. A. Frey, J. A. Gerlt, The low barrier hydrogen bond in enzymatic

728    catalysis. *J. Biol. Chem.* **273** (1998), pp. 25529–25532.

729 48. S. Zhou, L. Wang, Unraveling the structural and chemical features of biological short

730   hydrogen bonds. *Chem. Sci.* **10**, 7734–7745 (2019).
731