1   **Short hydrogen bonds enhance non-aromatic protein-related fluorescence**

2Amberley D. Stephens,*[1] Muhammad Nawaz Qaisrani,*[2,3,4] Michael T. Ruggiero,[5,1] Gonzalo

3Diaz Miron,[6] Uriel N. Morzan,[2] Mariano C. González Lebrero,[6] Saul T.E. Jones,[7] Emiliano

4Poli,[2] Andrew D. Bond,[8] Philippa J. Woodhams,[1] Elyse M. Kleist,[9] Luca Grisanti,[10] Ralph

5Gebauer,[2] J. Axel Zeitler,[1] Dan Credgington,[7] Ali Hassanali,[2#] Gabriele S. Kaminski

6Schierle[1#]

9[1] Chemical Engineering and Biotechnology, University of Cambridge, Cambridge CB3 0AS, UK
10[2] The Abdus Salam International Centre for Theoretical Physics, Strada Costiera 11, 34151 Trieste,
11Italy
12[3] SISSA International School for Advanced Studies, Via Bonomea 265, 34136 Trieste, Italy
13[4] Institut für Physik, Johannes Gutenberg Universität, Staudingerweg 7, 55128-Mainz, Germany
14[5] Department of Chemistry, University of Vermont, 82 University Place, Burlington, VT, 05405 USA
15[6] Departamento de Química Inorgánica, Analítica y Química Física, INQUIMAE, Facultad de
16Ciencias Exactas y Naturales, Universidad de Buenos Aires, Primer Piso Pabellon 2, C1428EHA,
17Ciudad Universitaria, Argentina.
18[7] Cavendish Laboratory, University of Cambridge, JJ Thomson Avenue, Cambridge, CB3 0HE, U.K
19[8] Department of Chemistry, University of Cambridge, Lensfield Road, Cambridge CB2 1EW, United
20Kingdom
21[9] Department of Chemistry, Syracuse University, Syracuse, New York 13244, United States
22[10] Division of Theoretical Physics, Ru̅der Boškovíc Institute, Bijenǐcka cesta 54, 10000 Zagreb,
23Croatia
24* Contributed equally
25[#] Corresponding authors: ahassana@ictp.it, gsk20@cam.ac.uk
26

28**Abstract**

29Fluorescence in biological systems is usually associated with the presence of aromatic

30groups. Here, we show that specific hydrogen bonding networks can significantly affect

31fluorescence employing a combined experimental and computational approach. In particular,

32we reveal that the single amino acid L-glutamine, by undergoing a chemical transformation

33leading to the formation of a short hydrogen bond, displays optical properties that are

34significantly enhanced compared to L-glutamine itself. *Ab initio* molecular dynamics

35simulations highlight that these short hydrogen bonds prevent the appearance of a conical

36intersection between the excited and the ground states and thereby significantly decrease non-

37radiative transition probabilities. Our findings open the doors for the design of new

38photoactive materials with biophotonic applications.

Significance statement

44Intrinsic fluorescence of non-aromatic amino acids is a puzzling phenomenon with an 45enormous potential for technological and biomedical applications. The physical origins of 46this effect, however, remain elusive. Herein, we demonstrate how specific hydrogen bond 47networks can modulate fluorescence. We highlight the key role played by short hydrogen 48bonds in the networks on the ensuing fluorescence and we provide a detailed molecular 49mechanism to explain this unusual non-aromatic optical properties. Our findings should 50benefit the design of novel optically active biomaterials for applications in biosensing and 51imaging.

Introduction

55Short peptides void of any aromatic residues have been shown to display an intrinsic 56fluorescence in the visible range (*1*, *2*). This has primarily been observed in fibrillar protein 57structures linked to neurodegenerative diseases, such as Alzheimer's, Parkinson's and 58Huntington's diseases (*3–6*). Furthermore, optical properties of double amino acid based 59nanowires have also been reported, consisting either of two non-aromatic or two aromatic 60amino acids (*2*, *7*, *8*). We have previously suggested that the fluorescence of non-aromatic 61short crystal structures forming part of the amyloid-beta protein is enhanced by proton 62delocalisation (*5*). We have hypothesised that one of the prerequisites for this fluorescence 63observed in either amyloid structures or short peptide nanowires is related to hydrogen 64bonding or aromatic interlocks which, for the latter, decreases the bandgaps down to the 65semiconductive regions (*9*).

67Despite our previous suggestion that proton delocalisation is strongly coupled to this intrinsic 68fluorescence, its direct role on putative fluorescing states has not been elucidated. We have 69thus searched for a model system, such as a single amino acid-based structure, that displays 70similar optical properties to amyloid fibrils and is permissive to more sophisticated 71computational approaches. We have been inspired by the small peptide nanostructures that 72have been pioneered by the Gazit laboratory (*9*) and by the fact that there are several 73neurodegenerative diseases that have been connected with an increased level of glutamines 74produced as part of a protein, as for example Huntingtin in Huntington's disease which 75renders the protein more aggregation prone (*10*). It has been known that the amide group in

76L-glutamine (L-glu) is highly labile and thus can rapidly hydrolyse. We show here that the
77single amino acid L-glu upon heating in water can form a nanostructural material with optical
78properties similar to the ones observed in other amyloid fibrils such as in fibrils of amyloid-
79beta, alpha-synuclein or tau (*4*, *11*, *12*).

81Using X-ray diffraction (XRD), we show that L-glu dissolved in water and upon heating
82becomes cyclysed forming a previously unreported structure which resembles L-
83pyroglutamine and which has been reported to be a component of amyloid-beta in the brain
84(*13*), but involves a low-barrier hydrogen bonded anionic dimer with an ammonium
85counterion. We have termed the new structure, i.e. L-pyroglutamine complexed with an
86ammonium ion, L-pyro-amm. L-pyro-amm has a microcrystalline plate morphology as shown
87by scanning electron microscopy (SEM). The newly formed solid has a unique hydrogen
88bond network formed by very strong hydrogen bonds (SHB) with a length of approximately
892.45 Å which is confirmed by using terahertz time-domain spectroscopy (THz-TDS).

91By employing a combination of electronic structure calculations and *ab initio* molecular
92dynamics in the ground and excited state we provide an interpretation of the reported
93experiments. We illustrate the important contribution of different vibrational distortions on
94the optical properties. Furthermore, our simulations identify the origins of the non-aromatic
95fluorescence in L-pyro-amm, demonstrating the key role played by the SHBs, which prevent
96the appearance of a conical intersection, significantly reducing the chances of non-radiative
97relaxation towards the ground state and hence increasing the optical properties.

99**Methods and Materials**

101*Experimental*

102*Sample preparation of L-glutamine*

103L-glutamine (L-glu) (#G3126, #G8540, Sigma-Aldrich, Gillingham, UK) and L-
104pyroglutamine (L-pyro) (#83160, Sigma-Aldrich) were dissolved in 18.2Ω MilliQ H_2O at a
105concentration of 0.3 M or 1 M. Aliquots were placed in a 65°C oven, since heating up
106proteins to 65°C increases the formation of amyloid structures as reported previously (*14*).
107Each aliquot was rotated to dissolve the powder once a day. Samples were either analysed in
108liquid form or dried on a glass or quartz cover slip (#043210.KG, Alfa Aesar, Lancashire,

109UK) either at room temperature (RT) or on a heat block set to 50°C. L-pyro-amm formed
110translucent crystals when dried.

112*Emission and excitation wavelength scans*

113Emission and excitation spectra were taken on a Hitachi F-4500 FL spectrophotometer
114(Hitachi High-Technologies Corporation, Tokyo, Japan) at room temperature (RT) in a quartz
115cuvette. For measurements, the excitation slit resolution was 5 nm or 10 nm and the emission
116slit resolution was 20 nm. The PMT voltage was set at 950 V and the scan speed set at 240
117nm/min. The excitation scan was measured between 250 – 400 nm and the emission filter set
118to the emission maxima of the sample stated in the figure legend, with a slit resolution of 20
119nm. The emission scan was measured between 380 - 560 nm and the excitation filter set to
120the excitation maxima of the sample stated in the figure legend, using a slit resolution of 5
121nm. Four measurements were taken for each sample which were repeated at least three times
122and the background (air or H_2O) was subtracted from the average.

124*Absorption measurements*

125Absorption measurements were taken on a UV-Vis-NIR Spectrophotometer, UV-3600 Plus
126(Shimadzu, Kyoto, Japan) and Cary 6000i (Agilent, Santa Clara, USA). 1 M or 0.3 M L-glu,
127L-pyro or L-pyro-amm solutions were measured in 10 mm QX cuvettes (Hellma Analytics,
128Müllheim, Germany) or dried on quartz coverslips. Measurements were taken between
129wavelengths 200 – 800 nm using 1 nm steps at a slow scan speed and a 1 nm resolution. The
130light source change wavelength was set at 393 nm and the grating change wavelength set at
131750 nm. Both direct and integrating sphere measurements were taken and showed little
132difference in results, direct measurements are shown in the manuscript. Samples were
133measured at least three times and the experiments repeated at least three times, measurements
134were then averaged and H_2O or coverslip only control was subtracted.

136*SEM (scanning electron microscopy)*

137SEM was performed using a FEI Magellan 400 HR-SEM at an acceleration voltage of 2 kV.
138L-pyro-amm samples were lyophilised by freezing in liquid nitrogen and freeze drying in a
139LyoQuest 85 (Telstar, Terrassa, Spain) and imaged on a glass coverslip.

141*X-ray diffraction*

142L-pyro-amm was dried on a glass coverslip in a 50°C oven and then at RT until crystals

143formed. Single crystal X-ray diffraction (SCXRD) measurements were performed at 180 °K

144with a Bruker D8-QUEST PHOTON-100 diffractometer, which utilised a Cu Kα radiation (λ

145= 1.54 Å), and an APEX-II CCD. Absorption corrections were made using SDABS, and data

146integration and reduction were performed with SAINT+. All non-hydrogen atoms were

147refined isotropically and anisotropically, followed by inclusion of the hydrogen atoms

148(determined using the excess electron density) and refinement isotropically. The structure is

149deposited in The Cambridge Crystallographic Data Centre, CCDC No. 1981551.

151*Terahertz Time-Domain Spectroscopy*

152All THz-TDS spectra were acquired using a commercial Terapulse 4000 spectrometer

153(TeraView Ltd, Cambridge, UK). Samples were prepared for THz-TDS measurements by

154diluting the solid air dried L-pyro-amm with polyethylene (~ 10% w/w concentration) by

155gentle mixing using an agate mortar, followed by pressing into 2 mm thick, 13 mm diameter

156pellets using a hydraulic press. All THz-TDS spectra shown are a result of division of sample

157and blank datasets, with the blank dataset represented the THz-TDS response of a pellet of

158pure polyethylene.

160*Theoretical*

161*Density functional theory (DFT)-THz Calculations*

162Calculations were performed using both the CRYSTAL17 (*15*) and Quantum Espresso (*16*)

163software packages. Geometry optimisations and vibrational analyses performed with the

164CRYSTAL17 code utilised the atom-centred split-valence triple-zeta 6-311g(2d,2p) basis set

165for all atom types. Based on a previous study related to ionic molecular crystals (*17*), the

166range-corrected WB97-X (*18*) functional was used. The vibrational analysis was performed

167within harmonic approximation, and infrared intensities were determined using the Berry

168Phase method (*19*). Energy convergence criteria were set to $\Delta E < 10^{-8}$ and 10^{-11} hartree for the

169geometry and frequency calculations, respectively.

171*Periodic time dependent (TD)-Density Functional Theory (DFT) Excited State Calculations*

172Simulations were performed using the fully periodic Quantum Espresso software package.

173The Becke-Lee-Yang-Parr (B3LYP) hybrid density functional was used with an energy

174cutoff of 40 Ry. The calculations of the excited state were performed within the framework of

175TDDFT using the Liouville-Lanczos formalism implemented in the freely available

176Quantum-Espresso package (*20*). In this approach, the optical spectra are obtained directly
177over the wide spectral range without taking into account the numerically complex
178calculations of the single exited states. We used plane wave basis set and the electron-ion
179interactions were taken into account via norm conserving Martins-Troullier pseudopotentials
180(*21*). To determine the ground state wave function, we used the gamma point of the Brillouin
181zone. All the periodic calculations employed the computationally demanding B3LYP (*22*)
182hybrid functional, the kinetic energy cutoff of 40 Ry was used for the wave functions. The
183intrinsic band width for the spectra was set to 0.003 Ry (~0.0408 eV).

185*Periodic Structure Geometry Optimisation*

186The structures obtained from the experiments were first geometrically optimized at 0°K using
187the Broyden-Fletcher-Goldfarb-Shanno (BFGS) minimisation algorithm implemented in
188CP2K (*23*, *24*) package. A convergence criterion for the wave function optimisation was used
189as 5×10^{-7} a.u.. Applying the method of the Gaussian and plane wave, the wave function was
190expended in the Gaussian double-zeta valence polarised (DZVP) basis set. The cutoff for the
191electronic density was set to 300 Ry. We used the gradient correction to the local density
192approximation and the core electrons were treated via Goedecker-Teter-Hutter
193pseudopotentials (*25*). In all the calculations, we used the Becke-Lee-Yang-Parr (BLYP) (*26*)
194functional with the D3(0) Grimme (*27*) dispersion corrections for the van der Waals
195interactions.

197*Ab Initio Molecular Dynamics Simulations*

198Ab initio Molecular Dynamics simulations (AIMD) were performed using Quickstep
199algorithm implemented in CP2K. In these calculations, the propagation of the nuclei was
200taken into account within the framework of the Born-Oppenheimer approximation. The
201simulations were performed in the NVT ensemble and the temperature was controlled during
202the simulations by using the velocity-rescaling thermostat (*28*). We used the time step of 0.5
203femtosecond to update the nuclear coordinates and velocities while the total length of the
204simulations for each system is 50 picoseconds.

206*Excited State Cluster Calculations*

207A set of excited state calculations were performed on glutamine clusters in order to
208understand the role of the environment on the optical properties. Specifically, the optical

209properties of L-pyro-amm were investigated using various isolated cluster models with the
210Gaussian09 software package. The clusters were extracted directly from the crystal structure
211and used in various combinations (dimers, trimers, tetramers) to perform TD-DFT
212calculations. A split-valence triple-zeta 6-311g(2d,2p) basis set was used for all atom types
213together with the hybrid B3LYP functional. Some benchmark simulations, comparing the
214optical properties obtained from the periodic calculations using B3LYP to range corrected
215hybrid functionals like CAM-B3LYP, were also performed with these clusters.

217We also performed a series of excited state optimisations on various model systems built
218from L-pyro-amm in order to examine the nature of the geometrical distortions that occur on
219the lowest electronic excited state. These calculations were also performed with the
220Gaussian09 software package. All clusters were surrounded with a continuum dielectric
221constant of 80, representing pure H_2O. The 6-311G(2d,2p) basis set was used for all atoms
222together with the range corrected hybrid functional CAM-B3LYP (*29*). The clusters were
223first optimised in the ground state after which they were optimised on the first electronic
224excited state.

226*Non-adiabatic Decay Probabilities Using Excited State Molecular Dynamics*

227Excited state AIMD was employed, as implemented in the LIO quantum-chemical package
228(https://github.com/MALBECC/lio) (*30–33*), to analyse the influence of three key factors
229determining the optical properties of L-pyro-amm: (i) the formation of a SHB, (ii) the
230presence of ammonium, and (iii) the combination of ring deplanarisation and carbonyl
231stretching. Our model system for this study was the L-pyro-amm dimer with a single SHB. In
232order to assess the influence of the ammonium ion on the optical properties of L-pyro-amm,
233AIMD was performed both on the L-pyro-amm with and without the ammonium ion.
234Analogously, in order to shed light on the role of the SHB in the L-pyro-amm fluorescence,
235in addition to performing simulations of L-pyro at the natural SHB distance of 2.5 Å, we
236performed three replicas constraining the HB distance to 3.0 Å, 3.5 Å and 4.5 Å, respectively.
237

238The initial structures for this analysis were extracted from the ground state AIMD described
239above. Subsequently, 3 ps ground state AIMD at 300K was performed to equilibrate the
240system, followed by 1ps of excited state AIMD. The non-radiative decay probability (NRP)
241was computed every time step using the TDDFT-based Trajectory Surface Hopping
242algorithm without permitting decays to the ground state (*34–36*). The non-adiabatic coupling

243elements between *S0* and *S1* (Equation 1) can be estimated analytically employing the
244method introduced by Tapavicza, *et al.,* (*37*).

245
$$\sigma_{S0,S1} = \langle \Psi_{S0}(r, R(t)) | \nabla_R | \Psi_{S1}(r, R(t)) \rangle \qquad (1)$$

246In this context, the non-radiative decay probability (NRP) from *S1* to *S0* (Equation 2) is
247expressed as:

248
$$NRP(t) = -2 \int_{t}^{t+\Delta t} dt' \frac{Re[c_{S1} c_{S0}^* \sigma_{S0,S1}(t')]}{c_{S0} c_{S0}^*}, \qquad (2)$$

249where C coefficients satisfy (Equation 3)

250
$$i\hbar \frac{\partial C_i(t)}{\partial t} = C_i(t)\omega_i - i\hbar \sum_{j=0}^{N_{states}} C_j \sigma_{ij}. \qquad (3)$$

251NRP(t) is the probability for the system to relax from *S1* to *S0* without fluorescent emission at
252time *t*. Hence, the accumulated non-radiative decay probability (Equation 4) accounts for the
253total relaxation probability from the excitation instant (*t=0*) up to time *t*.

254
$$ANRP(t) = \int_{0}^{t} dt' NRP(t') \qquad (4)$$

256As mentioned above, this methodology was applied to five different initial conditions of the
257dimer at SHB, 3.0, 3.5, and 4.5 Å of hydrogen distance and for the dimer-amm at SHB
258distance. In order to avoid introducing large structural deformations, the distance constrains
259in the HB were introduced in combination with position restrains on the nitrogen and the C_β
260of the pyroglutamic ring.

262The excited state AIMDs were performed employing the density functional theory (DFT)
263using the hybrid functional PBE0 level (*38*), with a 6-31G basis set and a time step of 0.5 fs.
264The computation of excitation energies, excited-state gradients, and non-adiabatic coupling
265vectors, the Linear Response TDDFT method was used within the Tamm-Dancoff
266approximation (*37*, *39*).

268**Results and Discussion**

270It has long been known that poly-glutamine can form amyloid-like fibrillar structures *in vitro*.
271The more glutamine residues in the poly-glutamine polymer, the faster the aggregation
272propensity of the polypeptide chain. This led us to investigate whether L-glu on its own,
273under conditions which normally promote fibril formation, such as an increase in temperature

274(*14*), was able to form structures with similar optical properties, as recently observed for
275amyloid fibrils (*5*, *40*, *41*).

277We first investigated the structure of L-pyro-amm, which formed after incubation of L-glu for
2788 days at 65°C, using SEM and observed crystal structures shown in Fig. 1a. However, in
279order to investigate whether L-glu had indeed changed its crystal structure arrangement we
280performed XRD analysis of the resulting material. In Fig. 1b we show the crystal structure of
281the heated L-glu structure, which we termed L-pyro-amm, and the published crystal
282structures of L-glu and L-pyroglutamine (L-pyro) in Suppl. Fig. 1. Note, the L-pyro structure
283was analysed as it displayed structural similarities to the newly formed L-pyro-amm. Figures
284were obtained from geometry optimisations of the nuclear positions of the atoms using
285experimental densities. L-pyro-amm consists of 8 pyroglutamine groups and 4 ammonium
286ions (144 atoms) complexed within the crystal (see Fig. 1b). In contrast, as shown in Suppl.
287Fig. 1a, L-glu consists of 4 glutamine molecules (80 atoms) in the unit cell which form
288hydrogen bonds involving the termini and side chain. Furthermore, as shown in Suppl. Fig.
2891b, L-pyro consists of 12 pyroglutamine molecules (192 atoms) in the unit cell forming
290hydrogen bonds involving the NH and COOH groups.

292L-pyro-amm has a rather unique hydrogen bond network structure since four of the
293pyroglutamine molecules are deprotonated and hence have a nominal negative charge, while
294the other four molecules are neutral. One of the important implications of this difference is
295that L-pyro-amm contains a very strong hydrogen bond. The red circled regions in Fig. 1b
296correspond to a short hydrogen bonds (SHBs) with a length of 2.45 Å, while those in L-glu
297and L-pyro range between ~2.55-2.85 Å, Suppl. Fig. 1.

299The structural change was further confirmed using THz-TDS measurements, as this technique
300is strongly dependent on the bulk packing arrangement as well as on the internal covalent
301structure of the molecules (*42*). The THz-TDS spectrum of the resulting solid, as well as the
302solid-state DFT predicted spectrum based on the single crystal XRD (SCXRD)-determined
303structure, is shown in Fig. 1c (full spectral assignment available in Suppl. Fig. 2). The
304agreement between the experimental and theoretical spectra further supports that full
305conversion of the sample occurs and thus enables additional investigations into the structural
306and electronic properties of the material. The agreement is also indicative of the ability of the

307theoretical model to not only reproduce the experimental structure, but also the weak forces
308found in solid structures.

310**Figure 1. L-glu forms L-pyro-amm upon heating.**

311 (a) SEM image of crystals of L-pyro-amm dried. (b) XRD analysis of heated L-glu sample
312show the newly formed structure, L-pyro-amm. Geometry optimisations show that 8
313pyroglutamine groups and four ammonium ions (144 atoms) are complexed in the crystal and
314a SHB of 2.45 Å (within red dashed lines) is present near the ammonium ion (white-
315hydrogen, red-oxygen, blue-nitrogen, grey-carbon). (c) Experimental (blue line) and
316theoretical (grey line) THz-TDS of the L-pyro-amm sample are in agreement and confirm the
317presence of the new L-pyro-amm structure.

319We first investigated whether there were any differences in the optical properties associated
320with the three crystal structures. Comparing the absorption of L-glu, L-pyro and L-pyro-amm
321in water, we show that only L-pyro-amm has a significantly red-shifted absorption which lies
322in the 275-320 nm range, whereas both L-glu and L-pyro primarily absorb in the deep UV
323(<250 nm) (see Fig. 2a).

325We next compared the experimental absorption spectra of L-glu, L-pyro and L-pyro-amm
326with the ones obtained from time dependent density functional theory (TDDFT). We
327highlight here, that the small size of the systems permitted us to determine the spectra using a
328hybrid functional, thereby not only advancing the quality of our theoretical predictions from
329previous studies (*5*, *40*, *41*) but also coupling the optical properties directly to different
330vibrational modes.

332Fig. 2b illustrates the absorption spectra obtained for the TDDFT calculations on the 3
333periodic systems in the ground state (i.e. at 0 °K). Panel b.i) shows the relative oscillator
334strength as a function of the frequency while panel b.ii) illustrates the second derivative of the
335oscillator strength permitting the positions of the maxima in the spectra to be more easily
336identified. The spectra reveal some striking differences between the different systems.
337Interestingly, we observe that L-pyro is essentially dark throughout the frequency range up to
338~ 6eV. On the other hand, L-pyro-amm shows the presence of more structure in the spectrum.
339Specifically, it is the only system for which the spectrum features a low energy excitation at
340226 nm (5.5 eV) and subsequently other peaks slightly above 220 nm (5.625eV) and 216 nm
341(5.75eV). While L-glu exhibits a peak at 222 nm (5.58eV), it is dark up to 206 nm (~6eV).
342

343We have previously shown that thermal fluctuations and in particular nuclear vibrations, such
344as proton transfer, have a large impact on the absorption spectra of peptide structures as
345compared to the 0 °K behavior (*5, 43–45*). In Fig. 2c we show that, compared to the 0 °K
346spectra in Fig. 2b, thermal fluctuations cause a large red shift to around 3.4 eV (365 nm) for
347L-pyro-amm, close to what is observed experimentally. These spectra were computed
348averaging over 25 frames sampled from the molecular dynamics simulations. Interestingly,
349no such effect is observed for L-glu which remains weakly absorbing up to more than 5 eV
350(247 nm) as seen at 0 °K.
351

352In order to understand better the physical origin of the low energy excitation at 226 nm (~5.5
353eV) in L-pyro-amm, we computed the electron response density at this frequency. This is
354illustrated in Fig. 2d, where we observe that most of the electron response is localised in
355regions around the pyroglutamine rings as well as regions near the SHB (see dashed circle in
356Fig. 2d). The optical response thus entails a charge reorganisation involving several parts of
357the molecular crystal. However, since L-pyro contains the same pyroglutamine rings as L-
358pyro-amm but absorbs more in the UV-spectrum, we conclude that the structural changes in
359the crystal in the presence of the SHB, which are the main features that distinguish L-pyro
360from L-pyro-amm, are responsible for the large Stokes shift observed in L-pyro
361experimentally and computationally.
362

363We next investigated whether the above structures also display fluorescence excitation and
364emission properties as has been observed for amyloid-like structures reported previously (*5,*

40, 41). Fig. 2e shows the excitation scan from 250-400 nm (solid lines) with the emission set
at 430 nm of L-glu in water at day 0 to 8 after incubation at 65°C. We observe an excitation
peak at around 360 nm which is similar to what we have measured previously for amyloid
proteins (5). The corresponding emission scan (dashed lines) with the excitation set at 360
nm and emission from 380-560 nm showed an emission peak around 430 nm, again lying in
the same visible range as for amyloid fibrils. When the L-pyro-amm solution was dried the
excitation and emission peaks were slightly blue shifted (Suppl. Fig. 3a) which may be due to
a change in the molecular environment in the dried state. Importantly, we do not see any
fluorescence in L-glu (without heating, i.e. at day 0 Fig. 2e.). To determine the importance of
the ammonium ion experimentally, L-pyro was incubated in water and heated at 65°C for 8
days, and only a very weak fluorescence has been detected (Suppl. Fig. 3b).

376
377 **Figure 2. Optical properties of L-pyro-amm are distinct from L-glu and L-pyro.**

378(a) Absorption spectra of 0.3 M L-glu (black), L-pyro (blue) and L-pyro-amm (red) (L-glu
379incubated for 8 days at 65°C) in water taken between 200 – 500 nm shows primarily features
380of L-pyro-amm. (b) Absorption spectra of L-glu, L-pyro and L-pyro-amm obtained from
381periodic density functional theory calculations with the B3LYP functional. L-pyro-amm
382features the lowest lying excited states which are characterised by the largest oscillator
383strengths. (c) Absorption spectra for L-glu and L-pyro-amm obtained from periodic
384simulations at room temperature. The spectra were computed by averaging over 25 frames
385randomly sampled from the *ab initio* molecular dynamics simulations. (d) The excited state
386electron density computed for L-pyro-amm from the optimised structure computed at the first
387peak (arrow in panel b). The lowest excited state density shows a response from various parts
388of the crystal structure including the pyroglutamic acid ring and the SHB region (see dashed
389circle). The orange and green surfaces correspond to regions involving a decrease and
390increase in electron density respectively, shown at an iso-value of 1×10^{-5}. (e) 1 M L-glu was
391incubated at 65°C and the excitation and emission spectra were measured over time.
392Excitation spectra were measured between 250-400 nm with emission set at 420 nm and
393emission spectra were measured between 380-560 nm with the excitation set at 360 nm.
394

395As alluded to earlier, one of the factors that distinguishes L-pyro-amm from the other systems
396is the presence of the SHB (highlighted by red circles in Fig. 1b) and the presence of the
397ammonium ion. In order to characterise the behaviour of the SHB, we conducted *ab initio*
398molecular dynamics simulations of the three systems at 300 °K and examined the proton
399transfer coordinates defined as the difference in distance between the proton (H) and the two
400oxygen atoms (O1 and O2) that sandwich it and is commonly referred to as the proton
401transfer coordinate ($d_{O1-H} - d_{O2-H}$) as shown in Fig. 3 for different types of hydrogen bonds in
402the crystals. It is clear that the SHB in L-pyro-amm is characterised by a double-well
403potential. The barrier associated with this proton transfer is on the order of thermal energy,
404indicating that zero-point energy (ZPE) would make the proton transfer barrierless (*46*). An
405examination of similar proton transfer coordinates for hydrogen bonds in L-glu and L-pyro
406show that they are characterised by only single-well potentials.

Figure 3: Free energy profiles along the proton transfer coordinate show only L-pyro-
amm displays a double well potential. Free energy profiles along the proton transfer
coordinate are displayed for L-glu (black), L-pyro (blue) and L-pyro-amm (red) at room
temperature. The L-glu and L-pyro display single well potentials, while the L-pyro-amm
system is the only one the exhibits a double well potential implying that there is proton
transfer from one side to the other.

The nature of the optical properties is sensitive to the environment in which the glutamine
molecules reside. It has previously been reported that charged amino acids already display an
absorption in the range of 250-350 nm that is significantly red shifted (*47*, *48*). The origins of
the low energy absorption were attributed to charge transfer excitations. The simulations of
these systems were performed in the gas phase, rather than considering the protein
environment such as shown for L-pyro-amm in Fig 2d. In comparison to the results presented
in Fig. 2d, data presented in Fig. 4, show that the origins of the electronic transitions equally
arise from a charge transfer (CT) between the highest occupied molecular orbital (HOMO) on
the anionic dimer, and the lowest unoccupied molecular orbital (LUMO) centred on the
ammonium cation when performed in the gas phase. Interestingly, the correct transition
energy is only predicted when the ammonium cation is spatially near the centre of the dimer,
which corresponds to the delocalisation of the negative charge and the SHB. Two generalised
geometries, with the ammonium cation near the SHB (as seen in Fig. 4a) and away from the
SHB (Fig. 4b), with the corresponding HOMO and LUMO orbitals are shown. The results
predict a transition of 304 nm for dimer (a) and 669 nm for dimer (b), with dimer (a) most
closely resembling the chemical environment present within the crystalline material.

433The results show that charge transfer is capable to lead to absorption in the near UV when 434investigated in the gas phase, i.e. neglecting the direct protein environment. However, 435including the protein environment in the molecular crystal (Fig. 2d) results instead in the 436excitation being a charge reorganisation involving several different molecular groups of the 437crystal. Indeed, by shuffling the protons along the SHBs in the ground-state, we observe an 438electronic response involving the entire structural units of L-pyro-amm including both the 439hydrogen bonded regions and the pyroglutamic acid rings when the protein environment is 440accounted for (Suppl. Fig. 4).

442**Figure 4. Comparison of the HOMO and LUMO orbitals on two L-pyro-amm models.**
443L-pyro-amm structures are presented where the ammonium cation is located directly near the 444hydrogen bond (a), and where the ammonium cation is located away from the hydrogen bond 445(b). While both models predicted charge-transfer HOMO-LUMO states, only in the case of 446(a) is the transition predicted to be in the vicinity of the experimentally observed peak, 304 447nm compared to 669 nm for (b).

449Up to this point, we have shown that proton transfer along SHBs is an important part of the 450structural fluctuations in the ground state structure of L-pyro-amm. In order to characterise 451the nuclear relaxation upon photoexcitation, we conducted both excited state optimisations as 452well as excited state molecular dynamics simulations. We first show the results from the 453excited state optimisations obtained from clusters carved out from the different glutamine 454crystals and surrounded by a continuum dielectric constant of 80. Fig. 5a shows a scatter plot 455of the difference between the first excited state and ground-state energies as well as the 456corresponding oscillator strengths for the three systems, L-glu, L-pyro and L-pyro-amm. The 457scatter plots were obtained over the course of the excited state optimisation. We observe that

458the L-pyro-amm system is characterised by the largest oscillator strengths peaking at
459approximately 3.5eV (354 nm), which is consistent with our experimental findings. Although
460L-glu and L-pyro also have a peak at around 3.5 eV it is much weaker than the one of L-pyro-
461amm. We thus decided to focus on a series of excited state optimisations for various clusters
462of L-pyro-amm.

464Fig. 5b shows the evolution of the first excited state energy for one of the L-pyro-amm
465clusters (Fig. 5c) across the optimisation steps. We find that the excited state energy drops by
4660.5 eV combined with an enhancement of the oscillator strength while the ground state
467energy, shown in solid-black, rises by about 1.5eV. In this case, the closing of the energy gap
468to approximately 3eV is associated with an increase of the carbonyl oxygen bond as observed
469in previous studies (48–50), as well as a de-planarisation of the pyroglutamine ring (see Fig.
4705c). Similar features are also observed in other clusters including one with the ammonium ion
471closer to the short hydrogen bond (Suppl. Fig. 5).

474**Figure 5. L-pyro-amm optical properties as seen through excited state optimisations.**
475(a) Scatter plot of the oscillator strengths versus the emission energy (defined as the
476difference between the first excited state and the ground-state) during the excited
477optimisations of L-glu (black), L-pyro (blue) and L-pyro-amm (red). (b) Ground and excited
478state energies are plotted as a function of the excited state optimisation of the system shown
479in c. The curves on the excited state are colour-coded with the oscillator strengths. S_0 and S_1
480refer to the ground and excited state energies, respectively. (c) A snapshot of the optimised
481excited state cluster also showing the lengthening of the carbonyl bond and the
482deplanarisation of the ring.

484To further explore the preceding mechanisms, we next turn to deploying non-adiabatic decay
485estimations using excited state ab initio molecular dynamics simulations, as implemented in
486the LIO quantum-chemical package (https://github.com/MALBECC/lio) (*30–33*). This
487approach enables investigating the S0←S1 de-excitation probability providing a clear
488interpretation of the ensuing optical properties of L-pyro-amm (see Methods for details). The
489nature of the S0←S1 transition is mainly a $\nu''=0 \leftarrow \nu'=0$ (where ν' and ν'' are the
490vibrational quantum numbers in the electronic ground and excited state respectively) with a
491FC factor between ground vibrational levels of ~0.82 (see Suppl. Fig. 6), therefore the excited
492state simulation was initiated by a vertical excitation to the S1 state. We performed two sets
493of excited state simulations: (i) varying the strength of the SHB to study its influence on the
494fluorescence of L-pyro dimer (Fig.6 a-b), and (ii) comparing L-pyro with L-pyro-amm
495dimers, to assess the role of ammonium ion on the transient excited state dynamics (Fig. 6c).
496Each trajectory was propagated for 1 ps from which the non-radiative decay probability
497(NRP) was determined by the fewest-switches trajectory surface hopping approach (*34–36*).
498Fig. 6d shows the time evolution of the accumulated NRP for various hydrogen bond lengths.
499The accumulated NRP (ANRP) represents the total probability for $S_0 \leftarrow S_1$ non-radiative
500relaxation. The larger value of the NRP implies that the system has a higher likelihood of
501decaying non-radiatively to the ground-state. On the other hand, a lower NRP would involve
502a longer excited-state lifetime increasing the fluorescence probability. Fig. 6d demonstrates
503unequivocally that the presence of the short hydrogen bond significantly reduces the chances
504of non-radiative relaxation towards the ground state. Examining the energy gap between the
505excited and ground state for this system shows that it occurs approximately ~3 eV (413 nm)
506(see Suppl. Fig. 7a) consistent with fluorescence in the blue/green visible regime. The
507presence of the ammonium ion further enhances, albeit by a subtle amount, the probability of
508trapping the system in the excited state.

Figure 6. Excited state AIMD highlight how the presence of a SHB and the ammonium ion is enhancing fluorescence. Excited state AIMD performed for the (a) L-pyro dimer, (b) the L-pyro dimer constraining the SHB distance, and (c) the L-pyro-amm. (d) Accumulated non-radiative $S_0 \leftarrow S_1$ decay probability (ANRP) for L-pyro dimer (dimer-SHB) and the constrained SHB L-pyro (dimer-) where the SHB distance was fixed at 3.0 Å, 3.5 Å and 4.5 Å and for the L-pyro-amm dimer (dimer-amm-SHB). (e.) Accumulated non-radiative decay probability for L-pyro (red discontinuous line) and L-pyro-amm (black dashed line). (f) Carboxyl C=O distance histogram for the L-pyro-amm dimer at the SHB distance (red curve) or constraining the SHB bond to 4.5 Å (blue curve). (g) Ring dihedral angle histogram for the L-pyro-amm dimer at the SHB distance (red curve) or constraining the SHB bond to 4.5 Å (blue curve).

The excited state optimisations shown in Fig. 5, illustrate the important role played by the vibrational distortions in the excited state that on the ensuing optical property. In agreement with this, Figs. 6f and g show the distribution of the C=O bond length and the ring de-planarisation (computed as the sum of its internal dihedral angles) for the dimer systems with

528the SHB and weaker (longer) hydrogen bonds in the left and right panel, respectively. 529Interestingly, we see that the trajectories that undergo non-radiative decay are characterised 530by both, a significant increase in the carbonyl stretch (C=O bond length) by about a tenth of 531an Angstrom, and a large de-planarisation of the dihedral angle. The combined distortion of 532these vibrational coordinates leads to non-radiative decay as observed in the dimer system 533with longer hydrogen bonds. In the system with the short and strong hydrogen bonds, these 534modes are hampered which essentially prevents the system from easily accessing conical 535intersections. This increases the likelihood of observing fluorescence in the L-pyro-amm 536structure characterised by short-hydrogen-bonds. Therefore, both our experiments and 537simulations demonstrate the crucial role played by the SHB and in determining the 538fluorescence of L-pyro-amm.

540**Conclusions**

542The experimental and theoretical findings presented here, elucidate a rather complex 543molecular mechanism associated with the non-aromatic intrinsic fluorescence in protein-like 544structures. In the case of L-glutamine, a chemical reaction creates a newly formed structure 545involving a cyclised pyroglutamic acid ring. This new structure features absorption in the UV 546and emission in the visible range very similar to the chemically distinct amyloid fibrils (*4*, *5*, 547*11*, *12*, *49–51*). The structural chromophore responsible for the optical properties in this new 548protein-related structure arises from a hydrogen bond network associated with structures 549involving short hydrogen bonds. Indeed, we have shown previously that similar to L-pyro-550amm, the crystallised structure of the hydrophobic core of amyloid beta, 2Y3J, L-pyro-amm 551contains a SHB (*52*) leading to a double well potential in the ground state. The presence of 552strong hydrogen bonds along which proton transfer occurs and secondly, specific ionic 553interactions in close proximity, such as involving the ammonium ion of L-pyro-amm, affect 554the optical properties. Although the fluorescence observed in these systems is much weaker 555compared to conventional fluorophores, the physical and chemical properties of the hydrogen 556bond networks reported here maybe a generic feature across many other peptide structures.

558Our non-adiabatic molecular dynamics simulations demonstrate that the presence of short 559hydrogen bonds with specific environmental conditions hinder vibrational deformations that 560can access conical intersections, as was previously speculated (*5*). The role of carbonyl 561groups is also consistent with a recent experimental study showing their importance for non-

562aromatic fluorescence (*53*). SHBs have recently been observed in different biological systems 563which have long been associated with either intrinsic fluorescence, such as NADP/NAD (*54*), 564FAD/FMN (*55*), the light-sensing chromophore in photoactive yellow protein (*56*), or in the 565active site of many enzymes, such as hydrolases and oxidoreductases (*57*, *58*), many of which 566consist of a highly complex H-bond structures similar to amyloid proteins. Furthermore, it 567has been recently reported that 1 in every 16 hydrogen bonds in over 1600 proteins are 568characterised by short hydrogen bonds (*58*). Thus, the mechanisms we espouse here may be 569more general. Our findings offer the possibility of designing novel biomaterial for 570applications in optical sensing or the design of novel biocompatible catalysts.

572**Supporting Information**

573Supporting data include; Supplementary Figure 1. Clusters of L-glu and L-pyro. 574Supplementary Figure 2. Full spectral assignment of THz data. Supplementary Figure 3. L-575pyro-amm has blue shifted fluorescence when dried and displays higher fluorescence 576intensity than L-pyro. Supplementary Figure 4. The optical properties of L-pyro-amm are 577sensitive to the environment and involves the electronic response of the entire structure. 578Supplementary Figure 5. L-pyro-amm undergoes vibrational distortions upon excitation as 579seen through excited state optimisations. Supplementary Figure 6. Potential energy surface 580along the SHB proton transfer coordinate in the ground (S0, panels A and D) and the excited 581(S1, panels B and C) states. Supplementary Figure 7. The energy gap between excited and 582ground state of L-pyro-amm is consistent with the experiments and eventually deactivates via 583a conical intersection back to the ground state. SCXRD data are available at the Cambridge 584Crystallographic Data Centre, CCDC No. 1981551. Raw data are available at the University 585of Cambridge Repository https://doi.org/10.17863/CAM.57945.

587**Author Information**

588**Corresponding Authors**

589Ali A. Hassanali, ahassana@ictp.it

590Gabriele S. Kaminski Schierle, gsk20@cam.ac.uk

592**Author Contributions**

593G.S.K.S and A.H conceptualised the manuscript. *A.D.S and M.N.Q contributed equally. 594A.D.S prepared samples for all experimental data. P.J.W. performed SEM experiments. 595A.D.B and M.T.R performed XRD measurements. M.T.R. performed THz experiments and

596DFT-THz calculations. A.D.S. performed excitation and emission measurements. A.D.S and
597S.T.J. performed absorption measurements. M.T.R and E.M.K performed TD-DFT cluster
598calculations. M.N.Q, E.P, L.G, R.G and A.H performed AIMD, Periodic TD-DFT Excited
599State Calculations and Periodic Structure Geometry Optimisation calculations. G.D.M,
600U.N.M, and M.C.GL performed excited state molecular dynamics. A.D.S, M.N.Q, M.T.R
601S.T.J, L.G, G.D.M, U.N.M, M.C.G.L, J.A.Z, D.C, A.H and G.S.K.S contributed to
602manuscript writing. All authors have given approval to the final version of the manuscript.

604**Notes**

605The authors declare no competing financial interest.

607**Acknowledgments**

608We thank Bluebell Drummond, Talia Shmool and Chetan Poudel for work that is not in the
609final manuscript.

611**Funding Sources**

612G.S.KS. acknowledges funding from the Wellcome Trust (065807/Z/01/Z) (203249/Z/16/Z),
613the UK Medical Research Council (MRC) (MR/K02292X/1), Alzheimer Research UK
614(ARUK) (ARUK-PG013-14), Michael J Fox Foundation (16238) and Infinitus China Ltd.
615M.T.R and J.A.Z acknowledge EPSRC funding (EP/N022769/1). P.J.W acknowledges
616EPSRC funding (EP/L016087/1). A.D.S. acknowledges Alzheimer Research UK the British
617Biophysical Society (BSS) for travel grants.

619**Abbreviations**

620SEM, scanning electron microscopy; XRD, X-ray diffraction; THz-TDS, terahertz time
621domain spectroscopy; DFT, density functional theory; TD-DFT, time dependent DFT; SHB,
622short hydrogen bond; CT, charge transfer; L-glu, L-glutamine; L-pyro, L-pyroglutamine; L-
623pyro-amm, L-pyroglutamine complexed with an ammonium ion.

625**References:**

6261.   A. Shukla, S. Mukherjee, S. Sharma, V. Agrawal, K. V. R. Kishan, P. Guptasarma, A
novel UV laser-induced visible blue radiation from protein crystals and aggregates:
Scattering artifacts or fluorescence transitions of peptide electrons delocalized through
hydrogen bonding? *Arch. Biochem. Biophys.* **428**, 144–153 (2004).

30. G. Rosenman, N. Amdursky, M. Molotskii, D. Aronov, L. Adler-Abramovich, E.
Gazit, Blue luminescence based on quantum confinement at peptide nanotubes. *Nano*
Lett. **9**, 3111–3115 (2009).

33. F. T. Chan, D. Pinotsi, G. S. K. Schierle, C. F. Kaminski, in *Bio-nanoimaging: Protein*
Misfolding and Aggregation (Elsevier, 2013;
http://linkinghub.elsevier.com/retrieve/pii/B9780123944313000134), pp. 147–155.

34. F. T. S. Chan, G. S. Kaminski Schierle, J. R. Kumita, C. W. Bertoncini, C. M. Dobson,
C. F. Kaminski, C. V Robinson, C. M. Dobson, J. Beard, P. Das, K. Jansen, M.
DeLucia, W.-L. Lin, G. Dolios, R. Wang, C. B. Eckman, D. W. Dickson, M. Hutton, J.
Hardy, T. Golde, Protein amyloids develop an intrinsic fluorescence signature during
aggregation. *Analyst.* **138**, 2156 (2013).

35. D. Pinotsi, L. Grisanti, P. Mahou, R. Gebauer, C. F. Kaminski, A. Hassanali, G. S.
Kaminski Schierle, Proton Transfer and Structure-Specific Fluorescence in Hydrogen
Bond-Rich Protein Structures. *J. Am. Chem. Soc.* **138**, 3046–3057 (2016).

36. J. Pansieri, V. Josserand, S. J. Lee, A. Rongier, D. Imbert, M. M. Sallanon, E. Kövari,
T. G. Dane, C. Vendrely, O. Chaix-Pluchery, M. Guidetti, J. Vollaire, A. Fertin, Y.
Usson, P. Rannou, J. L. Coll, C. Marquette, V. Forge, Ultraviolet–visible–near-infrared
optical properties of amyloid fibrils shed light on amyloidogenesis. *Nat. Photonics.* **13**,
473–479 (2019).

37. L. Adler-Abramovich, E. Gazit, The physical properties of supramolecular peptide
assemblies: From building block association to technological applications. *Chem. Soc.*
Rev. **43**, 7236–7236 (2014).

38. L. Adler-Abramovich, M. Reches, V. L. Sedman, S. Allen, S. J. B. Tendler, E. Gazit,
Thermal and chemical stability of diphenylalanine peptide nanotubes: Implications for
nanotechnological applications. *Langmuir.* **22**, 1313–1320 (2006).

39. K. Tao, P. Makam, R. Aizen, E. Gazit, Self-assembling peptide semiconductors.
Science (80-.). **358** (2017), p. eaam9756.

40. J. M. Andresen, J. Gayán, L. Djoussé, S. Roberts, D. Brocklebank, S. S. Cherny, L. R.
Cardon, J. F. Gusella, M. E. Macdonald, R. H. Myers, D. E. Housman, N. S. Wexler, J.
Lorimer, J. Porter, F. Gomez, C. Moskowitz, K. P. Gerstenhaber, E. Shackell, K.
Marder, G. Penchaszadeh, S. A. Roberts, A. Brickman, D. Brocklebank, J. Gray, S. R.
Dlouhy, S. Wiktorski, M. E. Hodes, P. M. Conneally, J. B. Penney, J. F. Gusella, J. H.
Cha, M. Irizarry, D. Rosas, S. Hersch, Z. Hollingsworth, A. B. Young, D. E. Housman,
M. M. de Young, E. Bonilla, T. Stillings, A. Negrette, S. R. Snodgrass, M. D.

Martinez-Jaurrieta, M. A. Ramos-Arroyoh, J. Bickham, J. S. Ramos, F. Marshall, I. Shoulson, G. J. Rey, A. Feigin, N. Arnheim, A. Acevedo-Cruz, L. Acosta, J. Alvir, K. Fischbeck, L. M. Thompson, A. Young, L. Dure, C. J. O'Brien, J. Paulsen, S. P. Moran, D. Krch, P. Hogarth, D. S. Higgins, B. Landwehrmeyer, M. R. Hayden, E. W. Almqvist, R. R. Brinkman, O. Suchowersky, A. Durr, C. C. Dodé, F. Squitieri, P. J. Morrison, M. Nance, C. A. Ross, R. L. Margolis, A. Rosenblatt, G. T. Estrella, D. M. Cabrero, R. J. A. Trent, E. McCusker, A. Novelletto, M. Frontali, J. S. Paulsen, R. Jones, A. Zanko, T. Ashizawa, A. Lazzarini, J. L. Li, V. C. Wheeler, A. L. Russ, G. Xu, J. S. Mysore, T. Gillis, M. Hakky, L. A. Cupples, M. Saint-Hilaire, S. M. Hersch, The relationship between CAG repeat length and age of onset differs for Huntington's disease patients with juvenile onset or adult onset. *Ann. Hum. Genet.* **71**, 295–301 (2007).

11. D. Pinotsi, A. K. Buell, C. M. Dobson, G. S. Kaminski Schierle, C. F. Kaminski, A Label-Free, Quantitative Assay of Amyloid Fibril Growth Based on Intrinsic Fluorescence. *ChemBioChem.* **14**, 846–850 (2013).

12. T. N. Tikhonova, N. R. Rovnyagina, A. Y. Zherebker, N. N. Sluchanko, A. A. Rubekina, A. S. Orekhov, E. N. Nikolaev, V. V. Fadeev, V. N. Uversky, E. A. Shirshin, Dissection of the deep-blue autofluorescence changes accompanying amyloid fibrillation. *Arch. Biochem. Biophys.* **651**, 13–20 (2018).

13. H. Mori, K. Takio, M. Ogawara, D. J. Selkoe, Mass spectrometry of purified amyloid β protein in Alzheimer's disease. *J. Biol. Chem.* **267**, 17082–17086 (1992).

14. M. Fändrich, C. M. Dobson, The behaviour of polyamino acids reveals an inverse side chain effect in amyloid structure formation. *EMBO J.* **21**, 5682–90 (2002).

15. R. Dovesi, A. Erba, R. Orlando, C. M. Zicovich-Wilson, B. Civalleri, L. Maschio, M. Rérat, S. Casassa, J. Baima, S. Salustro, B. Kirtman, Quantum-mechanical condensed matter simulations with CRYSTAL. *Wiley Interdiscip. Rev. Comput. Mol. Sci.* **8**, e1360 (2018).

16. P. Giannozzi, O. Andreussi, T. Brumme, O. Bunau, M. Buongiorno Nardelli, M. Calandra, R. Car, C. Cavazzoni, D. Ceresoli, M. Cococcioni, N. Colonna, I. Carnimeo, A. Dal Corso, S. De Gironcoli, P. Delugas, R. A. Distasio, A. Ferretti, A. Floris, G. Fratesi, G. Fugallo, R. Gebauer, U. Gerstmann, F. Giustino, T. Gorni, J. Jia, M. Kawamura, H. Y. Ko, A. Kokalj, E. Kücükbenli, M. Lazzeri, M. Marsili, N. Marzari, F. Mauri, N. L. Nguyen, H. V. Nguyen, A. Otero-De-La-Roza, L. Paulatto, S. Poncé, D. Rocca, R. Sabatini, B. Santra, M. Schlipf, A. P. Seitsonen, A. Smogunov, I.

698   Timrov, T. Thonhauser, P. Umari, N. Vast, X. Wu, S. Baroni, Advanced capabilities

699   for materials modelling with Quantum ESPRESSO. *J. Phys. Condens. Matter.* **29**

700   (2017), doi:10.1088/1361-648X/aa8f79.

70117.   M. T. Ruggiero, J. Gooch, J. Zubieta, T. M. Korter, Evaluation of Range-Corrected

702   Density Functionals for the Simulation of Pyridinium-Containing Molecular Crystals.

703   *J. Phys. Chem. A*. **120**, 939–947 (2016).

70418.   J. Da Chai, M. Head-Gordon, Long-range corrected hybrid density functionals with

705   damped atom-atom dispersion corrections. *Phys. Chem. Chem. Phys.* **10**, 6615–6620

706   (2008).

70719.   Y. Noel, C. M. Zicovich-Wilson, B. Civalleri, P. D'Arco, R. Dovesi, Polarization

708   properties of ZnO and BeO: An ab initio study through the Berry phase and Wannier

709   functions approaches. *Phys. Rev. B - Condens. Matter Mater. Phys.* **65**, 1–9 (2002).

71020.   D. Rocca, R. Gebauer, Y. Saad, S. Baroni, Turbo charging time-dependent density-

711   functional theory with Lanczos chains. *J. Chem. Phys.* **128** (2008),

712   doi:10.1063/1.2899649.

71321.   N. Troullier, J. L. Martins, Efficient pseudopotentials for plane-wave calculations.

714   *Phys. Rev. B*. **43**, 1993–2006 (1991).

71522.   A. D. Becke, Density-functional thermochemistry. III. The role of exact exchange. *J.*

716   *Chem. Phys.* **98**, 5648–5652 (1993).

71723.   J. D. Head, M. C. Zerner, A Broyden-Fletcher-Goldfarb-Shanno optimization

718   procedure for molecular geometries. *Chem. Phys. Lett.* **122**, 264–270 (1985).

71924.   J. VandeVondele, M. Krack, F. Mohamed, M. Parrinello, T. Chassaing, J. Hutter,

720   Quickstep: Fast and accurate density functional calculations using a mixed Gaussian

721   and plane waves approach. *Comput. Phys. Commun.* **167**, 103–128 (2005).

72225.   S. Goedecker, M. Teter, Separable dual-space Gaussian pseudopotentials. *Phys. Rev. B*

723   *- Condens. Matter Mater. Phys.* **54**, 1703–1710 (1996).

72426.   A. D. Becke, Density-functional exchange-energy approximation with correct

725   asymptotic behavior. *Phys. Rev. A*. **38**, 3098–3100 (1988).

72627.   S. Grimme, J. Antony, S. Ehrlich, H. Krieg, A consistent and accurate *ab initio*

727   parametrization of density functional dispersion correction (DFT-D) for the 94

728   elements H-Pu. *J. Chem. Phys.* **132**, 154104 (2010).

72928.   G. Bussi, D. Donadio, M. Parrinello, Canonical sampling through velocity rescaling. *J.*

730   *Chem. Phys.* **126** (2007), doi:10.1063/1.2408420.

73129.   T. Yanai, D. P. Tew, N. C. Handy, A new hybrid exchange-correlation functional

732   using the Coulomb-attenuating method (CAM-B3LYP). *Chem. Phys. Lett.* **393**, 51–57

733   (2004).

73430.   J. P. Marcolongo, A. Zeida, J. A. Semelak, N. O. Foglia, U. N. Morzan, D. A. Estrin,

735   M. C. González Lebrero, D. A. Scherlis, Chemical Reactivity and Spectroscopy

736   Explored From QM/MM Molecular Dynamics Simulations Using the LIO Code.

737   *Front. Chem.* **6**, 70 (2018).

73831.   U. N. Morzan, F. F. Ramírez, M. B. Oviedo, C. G. Sánchez, D. A. Scherlis, M. C. G.

739   Lebrero, Electron dynamics in complex environments with real-time time dependent

740   density functional theory in a QM-MM framework. *J. Chem. Phys.* **140**, 164105

741   (2014).

74232.   M. A. Nitsche, M. Ferreria, E. E. Mocskos, M. C. G. Lebrero, GPU accelerated

743   implementation of density functional theory for hybrid QM/MM simulations. *J. Chem.*

744   *Theory Comput.* **10**, 959–967 (2014).

74533.   F. Ramírez, · Gonzalo, D. Mirón, M. C. González Lebrero, D. A. Scherlis, QM-MM

746   Ehrenfest dynamics from first principles: photodissociation of diazirine in aqueous

747   solution. *Theor. Chem. Acc.* **137**, 124 (2018).

74834.   E. Tapavicza, I. Tavernelli, U. Rothlisberger, Trajectory surface hopping within linear

749   response time-dependent density-functional theory. *Phys. Rev. Lett.* **98**, 023001

750   (2007).

75135.   J. C. Tully, Molecular dynamics with electronic transitions. *J. Chem. Phys.* **93**, 1061–

752   1071 (1990).

75336.   M. Barbatti, Nonadiabatic dynamics with trajectory surface hopping method. *Wiley*

754   *Interdiscip. Rev. Comput. Mol. Sci.* **1** (2011), pp. 620–633.

75537.   E. Tapavicza, G. D. Bellchambers, J. C. Vincent, F. Furche, Ab initio non-adiabatic

756   molecular dynamics. *Phys. Chem. Chem. Phys.* **15** (2013), pp. 18336–18348.

75738.   C. Adamo, V. Barone, Toward reliable density functional methods without adjustable

758   parameters: The PBE0 model. *J. Chem. Phys.* **110**, 6158–6170 (1999).

75939.   S. Hirata, M. Head-Gordon, Time-dependent density functional theory within the

760   Tamm-Dancoff approximation. *Chem. Phys. Lett.* **314**, 291–299 (1999).

76140.   K. Jong, Y. Taghipour Azar, L. Grisanti, A. D. Stephens, S. T. E. Jones, D.

762   Credgington, G. S. Kaminski Schierle, A. A. Hassanali, Low Energy Optical

763   Excitations as an Indicator of Structural Changes Initiated at the Termini of Amyloid

764   Protein. *Phys. Chem. Chem. Phys.* (2019), doi:10.1039/c9cp04648h.

76541.   L. Grisanti, D. Pinotsi, R. Gebauer, G. S. Kaminski Schierle, A. A. Hassanali, A

computational study on how structure influences the optical properties in model crystal structures of amyloid fibrils. *Phys. Chem. Chem. Phys.* **19**, 4030–4040 (2017).

42. E. P. J. Parrott, J. A. Zeitler, Terahertz time-domain and low-frequency Raman spectroscopy of organic materials. *Appl. Spectrosc.* **69**, 1–25 (2015).

43. Y. K. Law, A. A. Hassanali, Role of Quantum Vibrations on the Structural, Electronic, and Optical Properties of 9-Methylguanine. *J. Phys. Chem. A.* **119**, 10816–10827 (2015).

44. S. Sappati, A. Hassanali, R. Gebauer, P. Ghosh, Nuclear quantum effects in a HIV/cancer inhibitor: The case of ellipticine. *J. Chem. Phys.* **145** (2016), doi:10.1063/1.4968046.

45. Y. K. Law, A. A. Hassanali, The importance of nuclear quantum effects in spectral line broadening of optical spectra and electrostatic properties in aromatic chromophores. *J. Chem. Phys.* **148** (2018), doi:10.1063/1.5005056.

46. D. Marx, Erratum: Proton transfer 200 years after von grotthuss: Insights from Ab initio simulations (ChemPhysChem (2006) 7 (1848-1810)). *ChemPhysChem.* **8**, 209–210 (2007).

47. S. Prasad, I. Mandal, S. Singh, A. Paul, B. Mandal, R. Venkatramani, R. Swaminathan, Near UV-Visible electronic absorption originating from charged amino acids in a monomeric protein. *Chem. Sci.* **8**, 5416–5433 (2017).

48. I. Mandal, S. Paul, R. Venkatramani, Optical backbone-sidechain charge transfer transitions in proteins sensitive to secondary structure and modifications. *Faraday Discuss.* **207**, 115–135 (2018).

49. A. Handelman, N. Kuritz, A. Natan, G. Rosenman, Reconstructive Phase Transition in Ultrashort Peptide Nanostructures and Induced Visible Photoluminescence. *Langmuir.* **32** (2016), pp. 2847–2862.

50. S. K. Joseph, N. Kuritz, E. Yahel, N. Lapshina, G. Rosenman, A. Natan, Proton-Transfer-Induced Fluorescence in Self-Assembled Short Peptides. *J. Phys. Chem. A.* **123**, 1758–1765 (2019).

51. L. L. Del Mercato, P. P. Pompa, G. Maruccio, A. Della Torre, S. Sabella, A. M. Tamburro, R. Cingolani, R. Rinaldi, Charge transport and intrinsic fluorescence in amyloid-like fibrils. *Proc Natl Acad Sci U S A.* **104**, 18019–18024 (2007).

52. K. Jong, L. Grisanti, A. Hassanali, Hydrogen Bond Networks and Hydrophobic Effects in the Amyloid β30-35Chain in Water: A Molecular Dynamics Study. *J. Chem. Inf. Model.* **57**, 1548–1562 (2017).

53. C. Niyangoda, T. Miti, L. Breydo, V. Uversky, M. Muschol, Carbonyl-based blue
autofluorescence of proteins and amino acids. *PLoS One*. **12**, e0176983 (2017).

54. A. Stavrinides, E. C. Tatsis, L. Caputi, E. Foureau, C. E. M. Stevenson, D. M. Lawson,
V. Courdavault, S. E. O'Connor, Structural investigation of heteroyohimbine alkaloid
synthesis reveals active site elements that control stereoselectivity. *Nat. Commun.* **7**
(2016), doi:10.1038/ncomms12116.

55. F. Forneris, D. P. H. M. Heuts, M. Delvecchio, S. Rovida, M. W. Fraaije, A. Mattevi,
Structural analysis of the catalytic mechanism and stereoselectivity in Streptomyces
coelicolor alditol oxidase. *Biochemistry*. **47**, 978–985 (2008).

56. S. Yamaguchi, H. Kamikubo, K. Kurihara, R. Kuroki, N. Niimura, N. Shimizu, Y.
Yamazaki, M. Kataoka, Low-barrier hydrogen bond in photoactive yellow protein.
Proc. Natl. Acad. Sci. U. S. A. **106**, 440–444 (2009).

57. W. W. Cleland, P. A. Frey, J. A. Gerlt, The low barrier hydrogen bond in enzymatic
catalysis. *J. Biol. Chem.* **273** (1998), pp. 25529–25532.

58. S. Zhou, L. Wang, Unraveling the structural and chemical features of biological short
hydrogen bonds. *Chem. Sci.* **10**, 7734–7745 (2019).

834
835
836Supplementary Information for
837
838Short hydrogen bonds enhance non-aromatic protein-related fluorescence
839

840Amberley D. Stephens, Muhammad Nawaz Qaisrani, Michael T. Ruggiero, Gonzalo Diaz Miron, Uriel
841N. Morzan, Mariano C. González Lebrero, Saul T.E. Jones, Emiliano Poli, Andrew D. Bond, Philippa
842J. Woodhams, Elyse M. Kleist, Luca Grisanti, Ralph Gebauer, J. Axel Zeitler, Dan Credgington, Ali
843Hassanali, Gabriele S. Kaminski Schierle
844
845Ali Hassanali, Gabriele S. Kaminski Schierle
846Email: ahassana@ictp.it, gsk20@cam.ac.uk

847
848

867**Supplementary Figure 1**. **Clusters of L-glu and L-pyro**

868Structural representations of (a) L-glutamine (L-glu) and (b) L-pyroglutamine (L-pyro)

869clusters from the published crystal structure.

Mode Number	Frequency (THz)	Intensity (km mol⁻¹)	Assignment
1	0.955	0.22	Antisymmetric translation in b
2	1.215	0.71	Coupled translation in a and rotation about bc
3	1.325	1.52	external rotation about b
4	1.450	0.36	Asymmetric translation in c
5	1.554	4.37	External rotation about a
6	1.676	1.76	Translation in a and external asymmetric rotation about H-bond coordinate
7	1.805	2.28	External asymmetric rotation of dimer pairs
8	1.929	2.11	External symmetric rotation of dimer pairs
9	1.950	1.28	Asymmetric rotation about h-bond coordinate, ammonium translational motion
10	2.113	3.53	Translation and rotation (breathing) around ammonium cation
11	2.208	2.5	Symmetric external rotation of entire formula units
12	2.404	2.37	In phase external rotation perpendicular to h-bond coordinate
13	2.502	0.45	Out of phase external rotation perpendicular to h-bond coordinate
14	2.564	0.73	External out of phase rotation of h-bonded chains about b
15	2.565	8.91	Out of phase external rotation of individual pyroglutamic molecules with translation of ammonium cation
16	2.654	18.09	External vibration coupled with torsion of the COOH group
17	2.679	0.29	External rotation and torsion of pyroglutamic ring
18	2.776	4.94	External rotation and torsion of ring and carboxyl group
19	2.988	2.68	External rotation and translation of ammonium

894 **Supplementary Figure 2. Full spectral assignment of THz data**

895 The top chart shows the contribution to each IR-active mode including external translations
896 and hindered rotations, and internal vibrational motions (i.e. torsions), while the bottom table
897 lists the detailed assignment for each mode.

910**Supplementary Figure 3. L-pyro-amm has blue shifted fluorescence when dried and**
911**displays higher fluorescence intensity than L-pyro**

912(a)The excitation peak of L-pyro-amm when dried (solid dark red line) is blue shifted with a
913peak maximum ~ 340 nm compared to L-pyro-amm in H_2O (solid red line) which has a peak
914maximum ~ 360 nm. The emission peak of L-pyro-amm when dried (dashed dark red line) is
915also blue shifted, with a peak maximum ~ 420 nm, compared to L-pyro-amm in H_2O (dashed
916red line) with a peak maximum ~ 430 nm. (b) 1 M L-glu and 1 M L-pyro (blue) were
917incubated in H_2O for 8 days at 65°C. After 9 days, the L-glu had converted into the L-pyro-
918amm structure (red). L-pyro-amm has a clear excitation peak maximum at ~360 nm and
919emission peak maximum at ~ 430 nm, while L-pyro (blue), although not completely dark, has
920no clear excitation or emission peak.

Supplementary Figure 4. The optical properties of L-pyro-amm are sensitive to the environment and involves the electronic response of the entire structure

(a)(i) L-pyro-amm cluster used to examine the sensitivity of the optical response on different parts of the cluster upon moving different protons. (ii) Charge differences between the ground and excited state are computed using restrained electrostatic potential atomic partial charges (RESP) for the cluster shown in i). Note, the electronic response involves all the atoms of the cluster. (iii) The two protons in the rectangle regions are displaced to be in the centre of the hydrogen bond and the charge differences are then computed. As illustrated, the proton displacement leads to a larger change in magnitude of the charges. (iv) The charge differences are computed for another nuclear configuration for which the protons are kept fixed but the O—O distance is increased from 2.45 to 3.2 Angstroms. The charge differences obtained here are quite similar to the original condition shown in (ii).

Supplementary Figure 5. L-pyro-amm undergoes vibrational distortions upon excitation as seen through excited state optimisations

(a-b) correspond to two additional clusters to the one reported in the main text that were built from the L-pyro-mm crystal structure. c) and (d) correspond to the evolution of the excited state energy and ground-state energy over the course of the optimisation. Also shown, are the changes in the C-O bond lengths (dashed black) which increases on the excited state surface. In (d) we show the O-H bond stretch where a proton transfer is observed on the excited state.

952Supplementary Figure 6. The S0 and S1 nuclear wavefunctions associated to the SHB
953proton transfer coordinate reveal a vibronic transition dominated by a 0←0 character.
954The calculated potential energy surfaces (black continuous lines) are prolongated with a
955quadratic fitting (gray dashed lines). Panels C and D illustrate the ground-state vibrational
956wavefunction in the S0 and the S1 electronic states (and   respectively). The resulting
957Franck-Condon factor between   and   is 0.82, showing that the nature of the S1 ← S0
958transition is mainly a $\nu''=0 \leftarrow \nu'=0$ (where ν' and ν'' are the vibrational quantum numbers
959in the electronic ground and excited state respectively).

966**Supplementary Figure 7. The energy gap between excited and ground state of L-pyro-**
967**amm is consistent with the experiments and eventually deactivates via a conical**
968**intersection back to the ground state**

969(a) Shows the energy of the excited and ground state obtained from the excited state
970molecular dynamics simulations where non-adiabatic decay probabilities are estimated. We
971see clearly that the energy difference between the excited and ground state is approximately 3
972eV (413 nm) consistent with experimental data. (b) Shows the same excited state and ground
973state energy for another simulation where the system de-activates very quickly within 10s of
974femtoseconds through a conical intersection where the ground and excited state energies
975intersect.

976