Proteogenomic single cell analysis of skeletal muscle myocytes

Katherine M. Fomchenko[1,4], Rohan X. Verma[1,4], Suraj Kannan[2], Brian L. Lin[2], Xiaoping Yang[1],

Tim O. Nieuwenhuis[1], Arun H. Patil[1], Karen Fox-Talbot[1], Matthew N. McCall[3], Chulan Kwon[2],

David A. Kass[2], Avi Z. Rosenberg[1], Marc K. Halushka[1]*

1 Department of Pathology, Johns Hopkins University School of Medicine, Baltimore, MD, USA
2 Division of Cardiology, Department of Medicine, Johns Hopkins University School of
Medicine, Baltimore, MD, USA
3 Department of Biostatistics and Computational Biology, University of Rochester Medical
Center, Rochester, NY, USA
4 These authors contributed equally to this project
13

Email Addresses:
Kfomche1@jhmi.edu
Rverma6@jhmi.edu
Skannan4@jhmi.edu
Blin29@jhmi.edu
Xyang15@jhmi.edu
Tnieuwe1@jhmi.edu
Ahanuma2@jhmi.edu
ktalbot@jhmi.edu
matthew_mccall@urmc.rochester.edu
ckwon13@jhmi.edu
dkass@jhmi.edu
arosen34@jhmi.edu
mhalush1@jhmi.edu
28

* Correspondence and address for reprints to:
Marc K. Halushka, M.D., Ph.D.
Johns Hopkins University School of Medicine
Ross Bldg. Rm 632B
720 Rutland Avenue
Baltimore, MD 21205
410-614-8138 (ph)
410-502-5862 (fax)
mhalush1@jhmi.edu
38

Abstract

Skeletal muscle myocytes have evolved into slow and fast-twitch types. These types are functionally distinct as a result of differential gene and protein expression. However, an understanding of the complexity of gene and protein variation between myofibers is unknown. We performed deep, whole cell, single cell RNA-seq on intact and fragments of skeletal myocytes from the mouse flexor digitorum brevis muscle. We compared the genomic expression data of 171 of these cells with two human proteomic datasets. The first was a spatial proteomics survey of mosaic patterns of protein expression utilizing the Human Protein Atlas (HPA) and the HPASubC tool. The second was a mass-spectrometry (MS) derived proteomic dataset of single human muscle fibers. Immunohistochemistry and RNA-ISH were used to understand variable expression. scRNA-seq identified three distinct clusters of myocytes (a slow/fast 2A cluster and two fast 2X clusters). Utilizing 1,605 mosaic patterned proteins from visual proteomics, and 596 differentially expressed proteins by MS methods, we explore this fast 2X division. Only 36 genes/proteins had variable expression across all three studies, of which nine are newly described as variable between fast/slow twitch myofibers. An additional 414 genes/proteins were identified as variable by two methods. Immunohistochemistry and RNA-ISH generally validated variable expression across methods presumably due to species-related differences. In this first integrated proteogenomic analysis of mature skeletal muscle myocytes we confirm the main fiber types and greatly expand the known repertoire of twitch-type specific genes/proteins. We also demonstrate the importance of integrating genomic and proteomic datasets.

Introduction

Skeletal muscle is a voluntary, striated muscle found throughout the body with contraction regulated by nerve impulses through the neuromuscular junction (NMJ). Skeletal muscles consist of different fiber types delineated by the isoform of the myosin heavy chain they express, metabolic function, and other properties (1). In humans, slow fibers (type 1) and some fast fibers (type 2A) exhibit oxidative metabolic properties, while fast type 2X fibers exhibit glycolytic metabolic properties (2). Mice have an additional type 2B fast fiber. These fiber types are variable across different muscles of the body reflecting different functional needs (2, 3).

Multiple proteins and protein classes vary across fiber types (1, 4). These include isoforms of the myosin heavy and light chains, calcium ATPase pumps, troponin T, and tropomyosin proteins, as well as metabolic proteins, such as pyruvate kinase, GAP dehydrogenase, and succinate dehydrogenase. Beyond these classes, there have been few efforts to catalog the entirety of fast/slow twitch expression differences by proteomics or genomics.

Among proteins, the deepest effort, to date, has been the single fiber proteomics work of the Mann laboratory (5, 6). In separate studies of mouse and human single fiber skeletal muscles, 1,723 and 3,585 proteins were reported, respectively, many of which were variably expressed among slow and fast twitch fibers. The most comprehensive gene expression study was performed in mice using DNA microarrays across ten type 1 and ten type 2B fibers (7). Single cell RNA-sequencing (scRNA-seq) also has been performed in skeletal muscle and muscle cultures. However, the large size of skeletal myocytes has precluded them from these datasets, which are instead predominately satellite cells, and other supporting cell types (8-15). A recent publication used SMART-Seq to evaluate three fast twitch mouse fibers (16). The totality of

these studies strongly suggests there are numerous expression differences between skeletal

muscle fiber types and a need for new approaches to capture this diversity.

The Kwon laboratory recently developed a large cell sorting method to isolate mature

cardiac myocytes (17). We ascertained if this method could be used to isolate the even larger

skeletal muscle myocytes for scRNA-seq. Our goal was to combine this genetic data with single

cell spatial proteomic data from the Human Protein Atlas (HPA) and an established mass

spectrometry human skeletal muscle proteomic dataset for a unique proteogenomic

characterization of skeletal muscle expression mosaicism.

Results:

scRNA-seq and identification of fast/slow twitch fiber types

We performed single cell RNA-seq using the established mcSCRB-seq protocol (18, 19).

We recovered data for 763 cells and sequenced to a median depth of 108,110 reads per cell. As

we were unsure of where the ideal skeletal myocytes might arise from our flow-sorting method,

they were taken from two different gates set on extinction (EXT) always "high" and time of

flight (TOF) being both high or low (Supplementary Fig. 1). Additional cells were collected

from a pseudo-biopsy approach with fragmented skeletal myocytes (see methods). Preliminary

analyses, however, indicated a distinct cluster of cells with a high percentage of mitochondrial

reads or otherwise low abundance reads. Notably, almost all of our pseudo-biopsy myocyte

fragments and many TOF-low cells fell into this category. These quality control metrics likely

indicated poor quality or sheared cells with loss of RNA. Thus, we excluded these cells and

narrowed our analysis to the best 171 cells remaining with a median read count of 239,252 per

cell.

An average of 12,098 transcripts were identified in these cells and all had the expression

patterns of mature skeletal myocytes, highly expressing a myosin heavy chain isoform. Because

of the narrow focus of this work to delineate cell subtypes and expression variability of just

skeletal muscle myocytes, this isolation strategy linked to deep sequencing, proved to be

advantageous.

We performed PCA of the data, corrected the data for the top 20 PCAs and utilized the

top 3,000 variable genes (by +/- standard deviation) to cluster these cell types (Fig. 1a). Three

Supplementary Figure 1. Mouse skeletal muscle myocyte preparation. A) Flow cytometry showing three gated areas representing EXT-high/TOF-high, EXT-high/TOF-low and EXT-low populations of flexor digitorum brevis myocytes. B) Representative images of Gate 1 EXT-high/TOF-high. C) Representative images of Gate 2 EXT-high/TOF-low. D) Representative images of Gate 3 EXT-low. E) Representative image of pseudo-biopsy isolated myocyte fragments. Gates 1 and 2 were used for library preparation. White size bar is 400 µm.

Fig. 1. a) UMAP graph of 171 skeletal muscle cells based on variable gene expression determined by scRNA-seq. **b)** UMAP graph based on mouse orthologous expression of HPASubC variable proteins. **c)** UMAP graph based on mouse orthologous expression of MS variable proteins **d-f)** Heat maps of major genes expression differences between the different fiber types based on the different datasets. **g)** Major myosin heavy chain distributions across the 171 cells as a percentage of each heavy chain. The colored areas are the assignments of each cell based on the scRNA-seq-based data **h)** Violin plots of 7 genes that varied between the two fast 2X$_a$ groups in the HPASubC-based data set. **i)** Assignment of each skeletal myocyte to a fiber type across the three methods. Strong agreement existed for the slow/fast 2A cells by any method of analysis

groups were observed in a UMAP dimensionality reduction plot. The first cluster, containing 69

cells (40% of all cells) had elevated expression of *Myh1* and *Myh8* clearly identifying this group

as containing fast 2X type cells and denoted as fast 2X$_a$ (Fig. 1d). A second cluster (N=53 cells)

had slightly more variable *Myh1* and *Myh8* differential expression, but by overall *Myh* gene

expression, Fig. 1g, also appeared to be a fast 2X cell type (denoted fast 2X$_b$). Of note, *Myh4*, a

myosin heavy chain associated with fiber type 2B, was elevated in a single cell in this group

(Fig. 1g) (3).

A third cluster (3) containing 49 cells (29% of the total) was defined by high expression

of *Tnnt1* and *Myh2.* A deeper analysis of this group showed that 12 cells had high to modestly

elevated *Myh7* expression (a slow-twitch marker), indicating this cluster was a combination of

slow-twitch cells and fast 2A fibers (Fig. 1g). The flexor digitorum brevis is a fast twitch

muscle, thus the overall distribution of significantly more fast (159) to slow fibers (12) is

consistent with expected.

Interestingly, the expression patterns of the main fast/slow differentiating *Myh* genes was

not as dichotomous as noted in protein based fiber type data (5). Here there were many more

cells with intermediate levels and coexpression of *Myh1* and *Myh2* suggesting higher gene

plasticity and more cell hybrids (Fig. 1g**)** (3).

HPA-based mosaic protein discovery

To complement variable gene expression data, we generated mosaic protein data by

performing an analysis of the IHC-based HPA dataset of skeletal muscle images using the

HPASubC suite of tools (20). The HPASubC tool, obtains a selected organs' images from the

Human Protein Atlas (HPA) and allows rapid and agnostic interrogation of images for staining

patterns of interest. This approach established a protein-based list of mosaically-expressed

proteins. Out of 50,351 images reviewed for 10,301 unique proteins, 2,164 proteins had possible

mosaic expression in skeletal muscle. Based on the aggregate image scores assigned to each

protein, they were subsetted into categories of "real" mosaicism (374 proteins), "likely"

mosaicism (1,231 proteins), and "unknown" probability of mosaicism (559 proteins)

(Supplementary Data 1, Supplementary Fig. 2). For analysis purposes, we focused on the 1,605

proteins that were in the "real" or "likely" categories to reduce the incidence of false positive

staining.

This method identified the well-known fiber type specific proteins such as MYH1,

MYH2, MY4, MHY6, MHY7, and MYH8 that were categorized as both "real" or "likely" based

on staining patterns (Supplementary Data 1). It also identified numerous uncharacterized or

poorly characterized proteins, such as the zinc finger proteins ZNF213, ZNF282, ZNF343,

Supplementary Figure 2. Scoring schema for HPASubC-based skeletal muscle mosaicism. A score of 1 indicated an "unknown" mosaicism based on subtle differences in stain intensity, or inconsistent patterns. A score of 2, "likely," was a clear distinction of staining by myofiber but the staining was not robust. A score of 3 "real" identified clear and robust staining differences by muscle cell. The score was primarily about the pattern and secondarily about the intensity of the staining difference.

ZNF350 and ZNF367 all of which had "real" patterns of mosaicism. A limitation of this spatial,

IHC-based approach is that each protein image is independent of other proteins. Thus, one

cannot identify co-expression patterns to assign proteins to certain fiber types.

We therefore investigated how these 1,605 proteins might inform on fiber type of

skeletal muscle cells by using this list to subset the orthologous mouse gene data from the

scRNA-seq experiment. Using just these orthologous mouse genes, we regenerated the UMAP

plot that identified four clusters (Fig. 1b). It essentially recapitulated the fast and slow fibers

types noted from the exclusive scRNA-seq data, despite being based on a different set of genes.

Uniquely, it subsetted the fast $2X_a$ cluster into two groups, one denoted by high expression of

Myom2 and *Gdf11* and the other denoted by high *Ucp1* and *Adamts4* (Fig. 1e). A t-test of gene

expression comparing genes from just these two subsets of the fast $2X_a$ cluster identified

multiple genes variably expressed between them (Fig. 1h). Although the cell clustering was

generally similar between mouse scRNA-seq gene data and HPASubC data with regard to

slow/fast 2A vs fast 2X, it was unclear which method was more representative. Therefore, we

obtained a public MS dataset as a third method to classify slow and fast twitch fibers.

Fast/slow twitch variation by MS-based proteomics

The human skeletal muscle fiber MS data in Murgia et al. is based on 152 fibers from

eight donors (5). This dataset had 596 proteins with >2.3 fold variation between type 1 and type

Supplementary Figure 3. UMAP of MS-based protein data by cell type. **A)** Seurat identified four cell clusters. **B)** UMAP was coloured based on cell assignments of Murgia et al. The slow type cells are generally Seurat cluster 2. Fast 2A cells are generally in Seurat cluster 3, although they are also detected in clusters 1 and 3. Fast 2X clusters are predominately in Seurat cluster 1.

2A fibers. We analyzed the full LFQ dataset of protein expression and constructed a UMAP plot

that showed four clusters (Supplementary Fig. 3). One cluster was composed primarily of slow

type 1 fibers and was adjacent to a second cluster with a small mixture of slow and other cell

types. Two other clusters were primarily a collection of fast 2X and fast 2A cell types. Similar to

the HPASubC approach above, we subsetted the orthologous mouse genes to these 596 proteins

to explore cell fiber type assignment.

As seen in the UMAP plot, five groups were identified (Fig. 1c). Similar to the other two

datasets (scRNA-seq and HPASubC), a slow/fast 2A fiber type was denoted by elevated

expression of several genes including *Tnnt1* and *Myl2* (Fig. 1f). One fast 2X fiber group ($2X_b$)

was identified by high expression of *Myh1* and *Myh8*. The second fast 2X fiber group was then

subdivided into three groups based on alternative elevated expression of genes that include

Gdf11 and *Ucp1* (group 3), *S100A8* (group 4) and *Adamts4* and *Mpz* (group 5). Unlike the

protein expression level based UMAP, slow fibers and fast 2A fibers were not distinct.

(Supplementary Fig. 3). This difference may be a result of the higher percentage of slow fibers in

the MS dataset.

Cross comparisons of the three approaches yield similar cell types.

We identified the cluster assignment of each skeletal muscle cell based on the scRNA-

seq, HPASubC, and MS approaches. We then plotted this information to demonstrate the extent

to which there was fluidity in assignment by fiber type (Fig. 1i). All but one cell (48/49)

assigned to the slow/fast 2A cluster based on scRNA-seq data remained in that cluster using

other methods of clustering (HPASubC and MS). An additional 7-8 cells from the fast 2X groups

became assigned to the slow/fast 2A cluster using the other methods of cell assignment. Cells

moved interchangeably between the fast $2X_a$ and fast $2X_b$ clusters depending on the method used

to cluster. We used this information to try and understand what distinguished fast $2X_a$ and fast

$2X_b$ clusters.

The $2X_a$ and fast $2X_b$ clusters differ by axonal genes.

To understand if the two fast 2X clusters represent unique cell types, cell states, or some

technical division, we performed a differential expression to determine what genes drove their

differences. Of 5,260 genes compared, 557 genes were differentially expressed (t. test; adj. p.

value <0.01). A Gene Ontology (GO) analysis on the 557 genes identified an enrichment of the

cellular component "neuronal synapse," suggesting variability at the NMJ. A further review of

the top significant genes showed that >20 genes appear to have neuronal origins (*Cdh4, Cdkl5,*

Cntn4, Dscam, Gabbr2, Kirrel3, Lingo2, Lrp1, L1cam, Nrcam, Ntn1, Ntrk3, Ptprt, Ptpro, Robo2,

Sdk1, Sema5a, Sema6d, Shank2, Sox5, Tnr, and *Wwox*). Of these, NRTK3, LRP1, and ROBO2

were identified as mosaic in skeletal muscle cells by HPASubC. Additionally, in HPA images,

seven orthologous proteins of these "neuronal" genes showed moderate staining, but each of

these had a TPM <1 (from GTEx expression data). Only LRP1 was identified in the orthologous

MS-dataset. This variability made us wonder how frequently the same genes/proteins were noted

to be mosaic by each of the three methods.

There is limited overlap of shared expression information

We compared the 3,000 most variable genes, the 1,605 HPASubC proteins, and the 596

MS proteins for shared patterns of mosaicism (Fig. 2). Only 23 genes/proteins were mosaic by

all three approaches using the Seurat analysis method (Fig. 2a). An additional 300

genes/proteins were shared across two methods, with the most overlap identified between the

two datasets with the most genes/proteins. Thus, we reasoned the abundance of genes/proteins

by method was a major driver of overlap leading us to focus on the 3,052 transcripts shared by

all three approaches, regardless of their mosaic/variable status. This resulted in 157

genes/proteins shared across any two methods with the most overlap between the two protein

datasets (77 proteins) (Fig. 2c).

As so few genes were shared with the protein sets, we wondered if the computational

approaches of the Seurat method limited the discovery of the correct variable genes. Therefore,

Fig. 2. Venn diagrams comparing the three methods and two analysis types with the full datasets (top) and the limited datasets (below). **a)** A Seurat-based overlap including all mosaic genes/proteins. **b)** A simpleSingleCell-based overlap including all mosaic genes/proteins. **c & d)** Seurat and simpleSingleCell-based methods limited to the 3,051 genes/proteins shared across the three studies.

we tried a second analysis approach, simpleSingleCell, to identify variable genes (21). By this

method, there was an increase (N=36) in overlap of genes/proteins being identified by all three

methods and more genes/proteins being identified by two methods (414) (Fig. 2b). Interestingly,

comparisons limited to the shared gene/protein list resulted in the highest overlap between the

MS- and gene-based datasets (Fig. 2d). A third method of using differential expression on the

scRNA-seq data to compare the subset of 12 slow-twitch cells to all fast twitch (2X and 2A) or

just fast 2X cells gave equivalent data to the simpleSingleCell approach.

Shared, abundant transcripts by cell type

We then wondered about the extent to which highly abundant proteins/genes were driving

our ability to detect mosaic proteins/genes. By normalized read counts of the scRNA-seq data,

we determined the 50 most abundant transcripts by the average of each cell type in the three

clusters determined by Seurat (Supplementary Data 2). Not surprisingly, the overall most

abundant transcripts were *Ttn*, *Acta1* and *mt-Rnr2*. Of the 23 mosaic genes/proteins found by all

three methods (using Seurat analysis), only *Myh1* and *Tnnt1* were on the list. Adding the mosaic

genes from the simpleSingleCell analysis, seven additional genes (*Mylpf*, *Tnnt3*, *Tmp1*, *Tnni2*,

Eno3, *Atp2a1* and *Pfkm*) were noted. This overall indicates that most abundant genes (≥41/50)

are not consistently mosaic in skeletal myocytes.

Species dichotomy in protein expression patterns

The generally low amount of overlap across the methods was unexpected. We wondered

if this discrepancy particularly between the gene and protein data was the result of species

differences in twitch type expression. To address this, we investigated staining patterns for three

proteins. Two (DCAF11, ENO3) were selected as they had clear mosaic staining by human

HPASubC images and no gene variation by Seurat analysis of the scRNA-seq. PVALB was

selected for showing variation by the mouse scRNA-seq data, but no variation by HPASubC.

DCAF11 was robustly mosaic in human but non-mosaic in mouse. ENO3 was mosaic in

both and PVALB was weakly mosaic in human but robustly mosaic in the mouse tissue (Fig. 3).

This data suggested that discrepancies may relate to differences in mosaic protein expression

between species (DCAF11) and possible technical causes (PVALB). Because ENO3 was mosaic

in the mouse skeletal muscle, but not mosaic by Seurat gene expression analysis, we explored if

a posttranscriptional form of regulation was occurring.

Fig. 3. Representative IHC and RNA-ISH of discrepant proteins and genes. a) HPA images (top row) are mosaic for DCAF11 and ENO3 and negative for PVALB staining. Follow up staining validated the DCAF11 and ENO3 staining while suggesting a subtle mosaicism of PVALB in humans. In mice, ENO3 and PVALB are clearly mosaic, while DCAF11 is not. **b)** RNA-ISH demonstrates co-expression of *CYB5R1* and *ENO3* in a mosaic pattern. **c)** Only *Eno3* was observed (in a mosaic pattern) in mouse muscle by RNA-ISH.

RNA-ISH indicates variable mosaicism

We performed RNA-ISH in both mouse and human skeletal muscles for *Eno3*, *Srsf11* and

Cyb5r1. All of their protein products were mosaic by HPASubC and MS protein expression and

had high or reasonably abundant gene expression (6552.8, 19.3, 201.5 pTPM respectively,

HPA). None of these genes were variably mosaic in the mouse gene data. We found mosaic co-

expression of all three genes in human skeletal muscle (Fig. 3). Whereas *ENO3* and *CYB5R1*

RNA was diffusely present across human skeletal myocytes, *SRSF11* was localized to sub-cell

membrane areas. In mouse muscle, *Eno3* was variably expressed, but neither *Cyb5r1* or *Srsf11*

were identified, although their levels of expression (~1,000x lower than *Eno3* in mouse) may be

too low to be seen by this method.

Many highly-supported variably expressed proteins were not previously identified

259   Thirty-six gene/proteins were variably expressed based on the simpleSingleCell,

HPASubC and MS based analyses (Fig. 2d, Table 1). Of these, based on an extensive literature

search, nine functionally diverse proteins are uniquely reported here as mosaic. Of the full 36,

22 were present in fast twitch myocytes and 14 in slow twitch myocytes based on the MS data.

In addition to these 36, another 414 genes/proteins were identified by two complementary

methods (Fig. 2). This includes well-known type specific proteins TNNC1 and TNNI1 (present

in the HPASubC and simpleSingleCell datasets, but not variably expressed in the MS dataset).

266   Finally, another 4,217 genes/proteins were variably expressed by one method. Of this

267   group, 1,211 were detected by HPASubC and 270 of these proteins were scored as "real" with

268   clear patterns of mosaicism (Supplementary Fig. 4).

Gene Name	Gene Symbol	Muscle Type- MS based	HPASubC Confidence	Seurat Norm. Counts	Mosaic Status
ArfGAP With Coiled-Coil, Ankyrin Repeat And PH Domains 2	Acap2	Fast	Likely	1.79	Unknown
Adenylate Kinase 1	Ak1	Fast	Likely	5.75	Known
Aldolase, Fructose-Bisphosphate A	Aldoa	Slow	Likely	7.98	Known
ATPase Sarcoplasmic/Endoplasmic Reticulum Ca2+ Transporting 1	Atp2a1	Fast	Real	6.68	Known
ATPase Sarcoplasmic/Endoplasmic Reticulum Ca2+ Transporting 2	Atp2a2	Slow	Real	2.08	Known
Calsequestrin 2	Casq2	Slow	Likely	1.61	Known

CD36 Molecule	Cd36	Slow	Likely	2.71	Known
Creatine Kinase, Mitochondrial 2	Ckmt2	Slow	Likely	4.54	Known
DnaJ Heat Shock Protein Family (Hsp40) Member C3	Dnajc3	Fast	Likely	2.2	Unknown
Enolase 3	Eno3	Fast	Real	6.97	Known
ELKS/RAB6-interacting/CAST Family Member 1	Erc1	Fast	Likely	3.14	Unknown
Glyceraldehyde-3-Phosphate Dehydrogenase	Gapdh	Fast	Likely	5.91	Known
Glyoxalase Domain Containing 4	Glod4	Slow	Likely	1.79	Unknown
Glycerol-3-Phosphate Dehydrogenase 1	Gpd1	Fast	Likely	3.22	Known
Kinesin Family Member 5B	Kif5b	Fast	Likely	3.83	Unknown
Lactate Dehydrogenase A	Ldha	Fast	Real	5.81	Known
Lactate Dehydrogenase B	Ldhb	Slow	Real	4.5	Known
Myosin Binding Protein C, Fast Type	Mybpc2	Fast	Real	5.2	Known
Myosin Heavy Chain 1	Myh1	Fast	Real	7.91	Known
Myosin Heavy Chain 6	Myh6	Slow	Likely	1.79	Known
Myosin Heavy Chain 8	Myh8	Fast	Likely	5.41	Known
Myosin Light Chain, Phosphorylatable, Fast Skeletal Muscle	Mylpf	Fast	Real	3.47	Known
Myosin Light Chain 3	Myl3	Slow	Real	1.95	Known
Myozenin 2	Myoz2	Slow	Real	8.06	Known
PDZ And LIM Domain 1	Pdlim1	Slow	Likely	3.43	Known
Peroxisomal Biogenesis Factor 19	Pex19	Fast	Likely	2.41	Unknown
Phosphofructokinase, Muscle	Pfkm	Fast	Likely	2.2	Known
Phosphoglycerate Kinase 1	Pgk1	Fast	Likely	3.67	Known
Ribosomal Protein S15a	Rps15a	Slow	Likely	4.94	Unknown
Thymosin Beta 4 X-Linked	Tmsb4x	Fast	Likely	3.58	Unknown
Troponin C2, Fast Skeletal Type	Tnnc2	Fast	Real	1.1	Known
Troponin I2, Fast Skeletal Type	Tnni2	Fast	Likely	5.19	Known
Troponin T1, Slow Skeletal Type	Tnnt1	Slow	Real	7.47	Known
Troponin T3, Fast Skeletal Type	Tnnt3	Fast	Real	6.31	Known
Topomyosin 1	Tpm1	Fast	Likely	7.97	Known
UDP-Glucose 6-Dehydrogenase	Ugdh	Slow	Likely	8.01	Unknown

Table 1. 37 Genes/Proteins identified as mosaic by all three methods based on simpleSingleCell analysis.

270 **Discussion**

Supplementary Figure 4. Nine representative images of 270 proteins scored as real mosaicism using HPASubC, but not identified by other methods. All images from HPA.

We describe the first proteogenomic analysis of skeletal muscle single fiber types using

combined scRNA-seq, spatial proteomics, and MS proteomics. Because delineations of skeletal

muscle fiber types are known and this project was exclusive to this one cell type, our study is a

useful model system to evaluate combining and synthesizing gene and protein data into a

coherent description of a cell. Also, by utilizing a deep sequencing approach and fewer cells, we

were not limited to just classifying a cell, but rather had sufficient data to delve into full gene

expression. Our data identifies common themes across the methods, but also significant

differences and complexities in gene/protein assignments.

Regardless of the method and genes/proteins used to cluster, we found general agreement

on the major types of skeletal muscle myocytes. We identified a small group of slow twitch cells

that clustered with fast 2A cells. These groups were consistently clustered away from two

clusters of fast 2X cells. The differences between these two fast 2X groups, described herein as

$2X_a$ and $2X_b$, are open to interpretation. The simplest explanation is that some axonal material

remained variably adherent to skeletal muscle cells through the NMJ, and these 20+ genes

resulted in the separation observed by UMAP (Fig. 1a). This would imply a technical cause of

the two fast 2X cell subtypes as a result of myocyte isolation. Adherent cell fragments are likely

to be a global issue for some cell type isolation, although it would not impact nuclear scRNA-seq

studies. A more interesting explanation is variable neuronal transfer of mRNAs across the NMJ

into the skeletal muscles via extracellular vesicles (22, 23). This would imply a real state-

difference in these cells, notable only by the deep sequencing strategy employed. Regardless, of

which is accurate, this division is unlikely to indicate true separate fast 2X subtypes. In fact, the

cross-referenced proteomic data was useful in demonstrating the arbitrary nature of this

delineation (Fig. 1i).

The extent of overlap of mosaic genes/proteins across the methods was surprisingly low.

Only 36 genes/proteins were cross-validated across all three approaches using the

simpleSingleCell method (Table 1). This list included well-known, fiber-type specific proteins

such as MYH1 and MYH6 and newly described mosaic proteins like DNAJC3 and GLOD4. The

lack of agreement across methods has made it difficult to confidently state how many

proteins/genes are variable by twitch pattern and further demonstrates the challenge of relying on

a single method. If a gene or protein is mosaic by two methods, this number climbs to 450. If all

mosaic genes and proteins are included, this increased to >4,500 genes/proteins. Over 1,600

proteins appear to be mosaic by the HPASubC method alone (Supplementary Data 1 and

Supplementary Fig. 4).

The reason for the variability in mosaic genes/proteins is certainly multifactorial. One

potential major difference is the comparison across two separate species (mouse and human). As

we noted with the DCAF11 IHC, this protein was mosaic in human muscle but did not appear to

be mosaic in mouse. Secondly, some genes have markedly variable expression levels between

the two species. While *CYB5R1* and *SRSF11* are robustly expressed in human muscle at 201.5

and 19.3 pTPM (in HPA), they were only 17.7 and 1.9 FPKM in our mouse scRNA-seq. It is

also possible that post-transcriptional regulation leads to more extreme expression variation in

proteins than genes. As described above, extreme expression dichotomy in *Myh* genes was less

than in similar MYH protein data (Fig. 1d) (5).

Our study represents the first use of LP-FACS to isolate single myofibers for scRNA-seq.

As skeletal myocytes are often long, stretching across the length of a muscle, isolation

techniques (particularly from human samples) may rely on the use of biopsies or otherwise

fragmented myocytes. To test the effect of myocyte fragmentation on scRNA-seq data quality,

we used a liberal gating strategy of our dissociated myocytes (including both EXT-high/TOF-

low and EXT-high/TOF-high populations) as well as directly sequencing fragmented myocytes

generated through a pseudo-biopsy approach. Disappointingly, we found that a large portion of

our sequenced myocytes were of poor quality, including those from our pseudo-biopsy approach.

By contrast, the highest quality data likely came from fully intact myocytes, in particular the

EXT-high/TOF-high population. Because this population is almost completely enriched for

intact myocytes, we believe that future experiments using LP-FACS to isolate skeletal myocytes

should focus solely on the EXT-high/TOF-high population. We are confident that this will allow

for a much higher percentage of good quality scRNA-seq libraries, akin to what we have

observed previously with LP-FACS isolation of cardiac myocytes (17). These results also mean

that more work must be done to identify better isolation methods for human skeletal muscle.

Current methods of human skeletal muscle biopsying from the quadriceps only obtains muscle

fragments and thus more creative methods to obtain full length fibers or non-damaged fibers

must be considered.

Technical factors also impact our ability to detect mosaicism on all platforms. Discovery

mass spectrometry is challenged to identify low abundance proteins. Having low input from

single fibers was further limiting and reduced the ability to computationally distinguish

expression differences in low abundance proteins. Most fibers had between 500-700 proteins

identified. As we have stated repeatedly, IHC in the HPA is subject to false positive staining

from shared epitopes (20, 24-26). It also incurs false negative staining for failed antibodies or

antibodies with staining parameters designed for other tissues. Further, some genes/proteins

observed in the other datasets were missing from the HPA data. The gene data was also limited

in the number of total cells analyzed (171) and the rarity of slow twitch cells from this muscle.

Cross-cell contamination, may have also stunted the differences between cell types (27).

In conclusion, we have created the first proteogenomic analysis of gene/protein

mosaicism in skeletal muscle. We replicated the known fiber types of slow, fast 2A, and fast 2X,

as well as greatly expanded our understanding of genes and with variable expression across these

cell types.

Methods

Isolation and Sequencing of Adult Skeletal Myocytes

Experiments were performed using C57BL/6J mice greater than 3 months of age. To isolate

skeletal myocytes, we performed collagenase-based digestion of the flexor digitorum brevis

(FDB), a short muscle of the hind feet, as per previously established protocols (28). We tested

two separate approaches to isolating myocytes. In the first approach, we dissected the FDB from

tendon to tendon prior to digestion, enabling isolation of fully intact myocytes. In the second

approach, we cut small portions of the FDB muscle using scissors. We reasoned that the latter

approach would broadly mimic skeletal muscle biopsy as might be done, for example, from a

human patient sample. The FDB was transferred to a dish containing DMEM with 1%

penicillin/streptomycin, 1% fetal bovine serum, and 2mg/mL Collagenase Type II

(Worthington). Muscle was digested for 1.5 hours in a 37C cell incubator with 5% CO_2.

Subsequently, the muscle was transferred to a dish containing media without collagenase, and

gently triturated to release single myocytes. Large undigested chunks and tendons were removed

with tweezers prior to single cell isolation.

We subsequently isolated single myocytes through large particle fluorescent-activated cell

sorting (LP-FACS), using a flow channel size of 500 µm. The COPAS SELECT Flow Pilot

Platform (Union Biometrica) was employed. Using time-of-flight (TOF, measuring axial length)

and optical extinction (EXT, measuring optical density) parameters, we found that skeletal

myocytes separated into three populations – an EXT-low population, EXT-high/TOF-low

population, and EXT-high/TOF-high population (Supplementary Fig. 1A). The EXT-high/TOF-

high population was comprised almost entirely of intact myofibers with lengths > 400 µm,

suggesting successful sorting of large myocytes (Supplementary Fig. 1B). Interestingly, the

EXT-high/TOF-low population was composed of what appeared to be rod-shaped fragments that

maintained sarcomeric proteins, albeit disrupted (Supplementary Fig. 1C). The EXT-low

population was comprised mostly of debris and dead cells, as previously observed with cardiac

myocytes (Supplementary Fig. 1D). The EXT-high/TOF-low population qualitatively resembled

our pseudo-biopsy isolated myocyte fragments (Supplementary Fig. 1E), which also shared

similar TOF and EXT parameters (not shown). To our knowledge, this is the first FACS-based

single cell RNA-seq study of skeletal myocytes; thus, we adopted a broad gating strategy for

isolation of single cells. We sorted 700 EXT-high myocytes (comprised of both TOF-high and

TOF-low populations) as well as 100 myocyte fragments isolated through the pseudo-biopsy

method.

These sorted cells were placed individually into 96-well plates. Capture plate wells contained 5

µl of capture solution (1:500 Phusion High-Fidelity Reaction Buffer, New England Biolabs;

1:250 RnaseOUT Ribonuclease Inhibitor, Invitrogen). Single cell libraries were then prepared

using the previously described mcSCRB-seq protocol (18, 19). Briefly, cells were subjected to

proteinase K treatment followed by RNA desiccation to reduce the reaction volume. RNA was

subsequently reverse transcribed using a custom template-switching primer as well as a barcoded

adapter primer. The customized mcSCRB-seq barcode primers contain a unique 6 base pair cell-

specific barcode as well as a 10 base pair unique molecular identifier (UMI). Transcribed

products were pooled and concentrated, with unincorporated barcode primers subsequently

digested using Exonuclease I treatment. cDNA was PCR-amplified using Terra PCR Direct

Polymerase (Takara Bio). Final libraries were prepared using 1ng of cDNA per library with the

Nextera XT kit (Illumina) using a custom P5 primer as previously described.

scRNA-seq sequencing and analysis

Pooled libraries were sequenced on two high-output lanes of the Illumina NextSeq500 with a 16

base pair barcode read, 8 base pair i7 index read, and a 66 base pair cDNA read design. To

analyze sequencing data, reads were mapped and counted using zUMIs 2.2.3 with default

396 settings and barcodes provided as a list (29). zUMIs utilizes STAR (2.5.4b) (30) to map reads to

397 an input reference genome and featureCounts through Rsubread (1.28.1) to tabulate counts and

398 UMI tables (30, 31). Reads were mapped to the mm10 version of the mouse genome. We used

399 GRCm38 from Ensembl concatenated with ERCC spike-in references for the reference genome

400 and gene annotations. Dimensionality reduction and cluster analysis were performed with Seurat

401 (2.3.4) (32).

402 **Seurat and simpleSingleCell**

403 Analysis was performed using the Seurat R toolkit V3.1.1 for this dataset (32). Initial filtering

404 removed lower quality cells (read count <5000 RNAs detected or >20% mitochondrial genes)

405 before sctransform normalization (33). A standard Seurat workflow was initially used for data

406 analysis. This workflow identifies a subset of genes with high cell-to-cell variation within the

407 scRNA-seq data. This subset is subsequently used as input to principal component analysis as

408 well as downstream nonlinear dimensionality reduction methods such as Uniform Manifold

409 Approximation and Projection (UMAP). Additionally, Seurat also allows for use of custom gene

410 lists as input to downstream analysis. This allowed us to use two custom gene lists, specifically

411 those derived from orthologous genes to mosaic proteins in the visual (HPASubC) dataset (20) or

412 the differentially expressed proteins in the MS proteomic dataset (5). Thus each of our three gene

413 lists, one produced by Seurat's workflow, another visual proteomic-based gene list, and a final

414 mass spectrometry-based gene list defining known muscle cell types, were used one at a time to

415 subset our initial data set and generate principal components for downstream analysis.

416 After determining clustering via these three approaches, UMAPs were generated alongside with

417 heat maps representing the top genes in clusters as determined by each gene set used for PCA.

418 Overlapping genes between the HPAsubC data, MS data, and significant genes determined by

Seurat were also examined for overlaps. Gene expression for *Trdh, Lama4, Ryr1, Dpp10,*

Pde4dip, Sugct, and *Myh1* was plotted across two fast $2X_a$ clusters based on the HPASubC data.

Simple Single Cell and Scran

Simple single cell 1.8.0 workflow was followed using scran 1.12.1 for normalization of raw

counts and fitting a mean-dependent trend to the gene-specific variances in single-cell RNA-seq

data (21). In line with this, we decomposed the gene-specific variance into biological and

technical components and selected the top 3000 genes for comparisons.

RNA-ISH

Mouse and human skeletal muscles were obtained at necropsy (>3 month old) and rapid autopsy

(66 year old male), the latter under an IRB-approved protocol. Tissues were immediately fixed in

formalin and paraffin-embedded blocks were created, from which 5 micron slides were made.

Custom probes for RNA *in situ* hybridization (RISH) were obtained from RNAscope (ACDBio).

These probes were designed to detect human and mouse forms of the following genes: *ENO3*

(GenBank accession nm_001976.5), *CYB5R1* (nm_016243.3), *SRSF11* (nm_004768.5), *Eno3*

(nm_007933.3), *Cyb5r1* (nm_028057.3), and *Srsf11* **(**nm_001093753.2). Each probe set targeted

all validated NCBI refseq transcript variants of the gene.

The Multiplex Fluorescent Reagent Kit v2 (ACDBio) was used following the manufacturer's

instructions. Briefly, FFPE tissue slides were baked for one hour at 60°C. The slides were

subsequently deparaffinized with xylene, rinsed with 100% ethanol and air-dried. After

application of hydrogen peroxide and washing, slides were treated with target retrieval reagent in

a steamer (>99°C) for 20 minutes. Then, the tissue was permeabilized using a protease.

Hybridization of the probes to the targeted mRNAs was performed by incubation in a 40°C oven

for 2 hours. After washes, the slides were processed for the standard signal amplification and

application of fluorescent dye (Opal dye 520, 570 and 620, AKOYA Biosciences) steps. Finally,

the slides were counterstained with DIPA, mounted with Prolong Gold Antifade Mounting

solution (Invitrogen) and stored in a 4°C room. The fluorescent images were obtained in the

Johns Hopkins Microscope Core Facility using a Zeiss LSM700 Laser scanning confocal

microscope.

Immunohistochemistry

The same tissues described above were used for standard immunohistochemistry. Antibodies

were obtained for WDR23/DCAF11 (bs-8388R, Bioss Antibodies), PVALB (A2781, Abclonal),

and ENO3 (ARP48203_T100, Aviva Systems Biology) that were reported to cross react to both

human and mouse. Immunohistochemistry was performed as described previously (25, 34).

HPA and HPASubC

The HPA is a comprehensive repository of IHC stained tissue microarrays for numerous

tissues, including skeletal muscle (35, 36). The HPASubC tool can rapidly and agnostically

interrogate images of the HPA to characterize specific staining patterns in organs (20, 24, 26).

HPASubC v1.2.4 was used to download 50,351 skeletal muscle tissue microarray images

covering 10,301 unique proteins from the HPA website (v18). The images were individually

reviewed using HPASubC by K.M.F to evaluate the presence of a mosaic pattern of protein

expression based on IHC staining. The classification of mosaicism was based on a pre-study

training set of 300 images from HPA reviewed collaboratively (K.M.F and M.K.H). Mosaicism

was defined as a dispersed pattern of differential staining in which a significant number of non-

adjacent muscle fibers had a higher staining intensity than the surrounding fibers, preferably

persisting across the entire microarray. All positive selections made by the trainee were reviewed

and rescored, as needed, by a board-certified pathologist (M.K.H.).

After an initial fast review of the images, a re-review to score the images was performed.

A three-tiered classification system was used indicating increasing certainty of mosaicism: 0

indicated the absence of mosaic staining; 1 indicated unknown mosaic staining; 2 indicated

likely mosaic staining; 3 indicated real mosaic staining. Scoring evaluation was based on the

quality of the mosaic pattern, including stain intensity differential between fibers, the presence of

"blush"/incomplete staining within cells, and the consistency and completeness of the fiber

staining pattern throughout the sample. HPASubC was used on an Apple MacBook Pro running

macOS Sierra v10.12.6 with 8 GB RAM and 3.1 GHz CPU and a Dell Precision Tower 3620

running Windows 10 with 16 GB RMA and a 3.7 GHz CPU.

Conversion of gene and protein symbols

To identify orthologs across human and mouse genes/proteins we had to synchronize

across gene/protein names and across the species. We used the David Gene ID Conversion Tool

(https://david.ncifcrf.gov/conversion.jsp), BioMart at Ensembl

(http://useast.ensembl.org/biomart/martview/e8a4fba4cb5c0be7a30841471b55674d), UniProt

Retreive/ID mapping (https://www.uniprot.org/uploadlists/) and direct searches at both UniProt

and GeneCards (https://www.genecards.org/), to cross integrate the human protein symbols,

mouse gene symbols, human gene symbols and ENSG IDs (37-39).

Gene Ontology (GO) Validation

GO was performed on the 557 most variable genes between two fast 2X clusters (2X$_a$ and 2X$_b$) using the Gene Ontology resource (http://geneontology.org/) and selecting for cellular component.

Mass Spectrometry (MS) Data Set

We utilized the Murgia et al. human skeletal muscle fiber MS-based proteomic dataset (5). This contained information from 3,585 proteins across 152 fibers from 8 donors (5). The ratio of expression of proteins between Type 1 and Type 2A cells were determined using Table S6 of Murgia et al. Five hundred and ninety-six proteins with >2.3 fold differences between cell types were selected. Label-free quantification (LFQ) data, from Supplemental Table S4, for the 154 human single muscle fiber proteomics was obtained. The log2 transformed LFQ data was converted to raw values and only proteins expressed across all fiber types (n=94) were considered for plotting UMAP as described (5). Functions of the R-package Seurat (Version 3.1.1) were executed sequentially to derive a UMAP along with its dependency library "uwot (Version 0.1.4)" in R (Version 3.6.1) (40, 41). A Seurat object of the data matrix was created using 'CreateSeuratObject' with default parameters. This data was normalized using the 'NormalizeData' function and outlier proteins were identified using the 'FindVariableFeatures.' Proteins across the fiber types were scaled and centered to create a PCA object using 'ScaleData' and 'RunPCA' respectively. Further, k-nearest neighbors and shared nearest neighbor for each fiber type were generated on the Seurat object using 'FindNeighbors' and 'FindClusters'to plot UMAP using 'RunUMAP'. All of these functions were executed using default parameters. The

clustering obtained with UMAP was overlaid with the classification of muscle fiber types based

on Murgia et al. using ggplot2 (Version 3.2.1).

Data availability

Mouse skeletal muscle sequencing was deposited at the Sequence Read Archive (SRA –

SRP241908) and the Gene Expression Omnibus (GSE143636).

Code availability

All analysis scripts are available at GitHub

(https://github.com/mhalushka/Skeletal_muscle_mosaicism).

Acknowledgements:

The authors thank Efrain Ribeiro for his helpful comments on the project. M.K.H. was supported

by grants 1R01HL137811, R01GM130564, and P30CA006973 from the National Institutes of

Health and 17GRNT33670405 from the American Heart Association. T.O.N. was supported by

grant R01GM130564. M.N.M. was supported by R01HL137811 and the University of Rochester

CTSA award number UL1TR002001. A.Z.R was supported by R01GM130564. C.K. and S.K.

were supported by NIH R01HD086026, TEDCO 2019-MSCRFD-5044, and the JHU Discovery

Award. S.K. was supported by fellowship 20PRE35200028 from the American Heart

Association.

Contributions K.M.F helped conceive the project and generated proteomic data. S.K. and

B.L.L. generated the skeletal muscle sequencing library. X.Y and K.F-T. performed IHC and

RISH. R.X.V., S.K., T.O.N, A.H.P. and M.N.M. performed analysis. C.K. and D.A.K. oversaw

the library preparation. A.Z.R helped develop the project. M.K.H. conceived the project,

performed analyses and wrote the manuscript. All authors contributed toward revisions of the

manuscript.

Conflicts of interest

The authors declare no conflicts of interest.

References

1.   Okumura N, Hashida-Okumura A, Kita K, Matsubae M, Matsubara T, Takao T, et al.
Proteomic analysis of slow- and fast-twitch skeletal muscles. Proteomics. 2005;5(11):2896-906.
2.   Gonzalez-Freire M, Semba RD, Ubaida-Mohien C, Fabbri E, Scalzo P, Hojlund K, et al.
The Human Skeletal Muscle Proteome Project: a reappraisal of the current literature. J Cachexia
Sarcopenia Muscle. 2017;8(1):5-18.
3.   Schiaffino S, Reggiani C. Fiber types in mammalian skeletal muscles. Physiological
reviews. 2011;91(4):1447-531.
4.   Drexler HC, Ruhs A, Konzer A, Mendler L, Bruckskotten M, Looso M, et al. On
marathons and Sprints: an integrated quantitative proteomics and transcriptomics analysis of
differences between slow and fast muscle fibers. Molecular & cellular proteomics : MCP.
2012;11(6):M111 010801.
5.   Murgia M, Toniolo L, Nagaraj N, Ciciliot S, Vindigni V, Schiaffino S, et al. Single
Muscle Fiber Proteomics Reveals Fiber-Type-Specific Features of Human Muscle Aging. Cell
reports. 2017;19(11):2396-409.
6.   Murgia M, Nagaraj N, Deshmukh AS, Zeiler M, Cancellara P, Moretti I, et al. Single
muscle fiber proteomics reveals unexpected mitochondrial specialization. EMBO Rep.
2015;16(3):387-95.
7.   Chemello F, Bean C, Cancellara P, Laveder P, Reggiani C, Lanfranchi G. Microgenomic
analysis in skeletal muscle: expression signatures of individual fast and slow myofibers. PLoS
One. 2011;6(2):e16807.
8.   Giordani L, He GJ, Negroni E, Sakai H, Law JYC, Siu MM, et al. High-Dimensional
Single-Cell Cartography Reveals Novel Skeletal Muscle-Resident Cell Populations. Molecular
cell. 2019;74(3):609-21 e6.
9.   Porpiglia E, Samusik N, Ho ATV, Cosgrove BD, Mai T, Davis KL, et al. High-resolution
myogenic lineage mapping by single-cell mass cytometry. Nature cell biology. 2017;19(5):558-
67.
10.   Dell'Orso S, Juan AH, Ko KD, Naz F, Perovanovic J, Gutierrez-Cruz G, et al. Single cell
analysis of adult mouse skeletal muscle stem cells in homeostatic and regenerative conditions.
Development. 2019;146(12).
11.   Cho DS, Doles JD. Single cell transcriptome analysis of muscle satellite cells reveals
widespread transcriptional heterogeneity. Gene. 2017;636:54-63.
12.   Cornelison DD, Wold BJ. Single-cell analysis of regulatory gene expression in quiescent
and activated mouse skeletal muscle satellite cells. Developmental biology. 1997;191(2):270-83.
13.   Cacchiarelli D, Qiu X, Srivatsan S, Manfredi A, Ziller M, Overbey E, et al. Aligning
Single-Cell Developmental and Reprogramming Trajectories Identifies Molecular Determinants
of Myogenic Reprogramming Outcome. Cell Syst. 2018;7(3):258-68 e3.

14.   Trapnell C, Cacchiarelli D, Grimsby J, Pokharel P, Li S, Morse M, et al. The dynamics
and regulators of cell fate decisions are revealed by pseudotemporal ordering of single cells.
Nature biotechnology. 2014;32(4):381-6.
15.   Rubenstein AB, Smith GR, Raue U, Begue G, Minchev K, Ruf-Zamojski F, et al. Single-
cell transcriptional profiles in human skeletal muscle. Scientific reports. 2020;10(1):229.
16.   Blackburn DM, Lazure F, Corchado AH, Perkins TJ, Najafabadi HS, Soleimani VD.
High-Resolution Genome-Wide Expression Analysis of Single Myofibers Using SMART-Seq. J
Biol Chem. 2019.
17.   Kannan S, Miyamoto M, Lin BL, Zhu R, Murphy S, Kass DA, et al. Large Particle
Fluorescence-Activated Cell Sorting Enables High-Quality Single-Cell RNA Sequencing and
Functional Analysis of Adult Cardiomyocytes. Circ Res. 2019;125(5):567-9.
18.   Soumillon M, Cacchiarelli D, Semrau S, van Oudenaarden A, Mikkelsen TS.
Characterization of directed differentiation by high-throughput single-cell RNA-Seq. BioRxiv.
2014.
19.   Ziegenhain C, Vieth B, Parekh S, Reinius B, Guillaumet-Adkins A, Smets M, et al.
Comparative Analysis of Single-Cell RNA Sequencing Methods. Molecular cell.
2017;65(4):631-43 e4.
20.   Cornish TC, Chakravarti A, Kapoor A, Halushka MK. HPASubC: A suite of tools for
user subclassification of human protein atlas tissue images. Journal of pathology informatics.
2015;6:36.
21.   Lun AT, McCarthy DJ, Marioni JC. A step-by-step workflow for low-level analysis of
single-cell RNA-seq data with Bioconductor. F1000Research. 2016;5:2122.
22.   Ashley J, Cordy B, Lucia D, Fradkin LG, Budnik V, Thomson T. Retrovirus-like Gag
Protein Arc1 Binds RNA and Traffics across Synaptic Boutons. Cell. 2018;172(1-2):262-74 e11.
23.   Korkut C, Ataman B, Ramachandran P, Ashley J, Barria R, Gherbesi N, et al. Trans-
synaptic transmission of vesicular Wnt signals through Evi/Wntless. Cell. 2009;139(2):393-404.
24.   Anene DF, Rosenberg AZ, Kleiner DE, Cornish TC, Halushka MK. Utilization of
HPASubC for the identification of sinusoid-specific proteins in the liver. Journal of proteome
research. 2016;15(5):1623-9.
25.   Wang TY, Lee D, Fox-Talbot K, Arking DE, Chakravarti A, Halushka MK.
Cardiomyocytes have mosaic patterns of protein expression. Cardiovasc Pathol. 2018;34:50-7.
26.   Cheah JX, Nieuwenhuis TO, Halushka MK. An expanded proteome of cardiac t-tubules.
Cardiovasc Pathol. 2019;42:15-20.
27.   Nieuwenhuis TO, Yang S, Verma RX, Pillalamarri V, Arking DE, Rosenberg AZ, et al.
Basal Contamination of Sequencing: Lessons from the GTEx dataset. BioRxiv. 2019.
28.   Shefer G, Yablonka-Reuveni Z. Isolation and culture of skeletal muscle myofibers as a
means to analyze satellite cells. Methods Mol Biol. 2005;290:281-304.
29.   Parekh S, Ziegenhain C, Vieth B, Enard W, Hellmann I. zUMIs - A fast and flexible
pipeline to process RNA sequencing data with UMIs. Gigascience. 2018;7(6).
30.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast
universal RNA-seq aligner. Bioinformatics. 2013;29(1):15-21.
31.   Liao Y, Smyth GK, Shi W. The Subread aligner: fast, accurate and scalable read mapping
by seed-and-vote. Nucleic Acids Res. 2013;41(10):e108.
32.   Butler A, Hoffman P, Smibert P, Papalexi E, Satija R. Integrating single-cell
transcriptomic data across different conditions, technologies, and species. Nature biotechnology.
2018;36(5):411-20.

33.   Hafemeister C, Satija R. Normalization and variance stabilization of single-cell RNA-seq
data using regularized negative binomial regression. BioRxiv. 2019.
34.   Wang TY, Arking DE, Maleszewski JJ, Fox-Talbot K, Nieuwenhuis TO, Santhanam L, et
al. Human cardiac myosin light chain 4 (MYL4) mosaic expression patterns vary by sex.
Scientific reports. 2019;9(1):12681.
35.   Uhlen M, Fagerberg L, Hallstrom BM, Lindskog C, Oksvold P, Mardinoglu A, et al.
Proteomics. Tissue-based map of the human proteome. Science. 2015;347(6220):1260419.
36.   Uhlen M, Oksvold P, Fagerberg L, Lundberg E, Jonasson K, Forsberg M, et al. Towards
a knowledge-based Human Protein Atlas. Nature biotechnology. 2010;28(12):1248-50.
37.   Yates A, Akanni W, Amode MR, Barrell D, Billis K, Carvalho-Silva D, et al. Ensembl
2016. Nucleic Acids Res. 2016;44(D1):D710-6.
38.   The UniProt C. UniProt: the universal protein knowledgebase. Nucleic Acids Res.
2017;45(D1):D158-D69.
39.   Fishilevich S, Zimmerman S, Kohn A, Iny Stein T, Olender T, Kolker E, et al. Genic
insights from integrated human proteomics in GeneCards. Database : the journal of biological
databases and curation. 2016;2016.
40.   Stuart T, Butler A, Hoffman P, Hafemeister C, Papalexi E, Mauck WM, 3rd, et al.
Comprehensive Integration of Single-Cell Data. Cell. 2019;177(7):1888-902 e21.
41.   Becht E, McInnes L, Healy J, Dutertre CA, Kwok IWH, Ng LG, et al. Dimensionality
reduction for visualizing single-cell data using UMAP. Nature biotechnology. 2018.
639