Title: Hmx3a does not require its homeodomain for its essential functions in zebrafish spinal cord, ear and lateral line development.

Authors:

Samantha J. England[*], Gustavo A. Cerda[†], Angelica Kowalchuk[*], Taylor Sorice[*], Ginny Grieb[*] and Katharine E. Lewis[*§]

[*]Department of Biology, Syracuse University, 107 College Place, Syracuse, NY 13244, USA

[†]Department of Physiology, Development and Neuroscience, University of Cambridge, Downing Street, Cambridge CB2 3DY, UK.

[§]Corresponding author. Department of Biology, Syracuse University, 107 College Place, Syracuse, NY 13244, USA. Telephone: +1 315 443 5902; Email: kelewi02@syr.edu

Running Title: Hmx2 and Hmx3a functions in zebrafish development

Key Words: Hmx2, morpholino, dI2, interneuron, genetic compensation

Abstract

Transcription factors that contain a homeodomain DNA-binding domain have crucial functions in most aspects of cellular function and embryonic development in both animals and plants. Hmx proteins are a sub-family of NK homeodomain-containing proteins that have fundamental roles in development of sensory structures such as the eye and the ear. However, Hmx functions in spinal cord development have not been analyzed. Here we show that zebrafish (*Danio rerio*) *hmx2* and *hmx3a* are co-expressed in spinal dI2 and V1 interneurons, whereas *hmx3b*, *hmx1* and *hmx4* are not expressed in spinal cord. Using mutational analyses, we demonstrate that, in addition to its previously reported role in ear development, Hmx3a is required for correct specification of a subset of spinal interneuron neurotransmitter phenotypes, as well as correct lateral line progression and survival to adulthood. Surprisingly, despite similar expression patterns of *hmx2* and *hmx3a* during embryonic development, zebrafish *hmx2* mutants are viable and have no obviously abnormal phenotypes in sensory structures or neurons that require Hmx3a. In addition, embryos homozygous for deletions of both *hmx2* and *hmx3a* have identical phenotypes to severe *hmx3a* single mutants. However, mutating *hmx2* in hypomorphic *hmx3a* mutants that usually develop normally results in abnormal ear and lateral line phenotypes. This suggests that while Hmx2 cannot compensate for loss of Hmx3a, it does function in these developmental processes, although to a much lesser extent than Hmx3a. More surprisingly, our mutational analyses strongly suggest that Hmx3a does not require its homeodomain DNA-binding domain for its roles in viability or embryonic development.

Introduction

Homeobox-containing genes and the Homeodomain-containing transcription factors that they encode, have crucial functions in most aspects of cellular function and embryonic development in both animals and plants (Burglin and Affolter 2016). They were also some of the first examples discovered of invertebrate developmental genes that are highly conserved in vertebrates (Carrasco *et al.* 1984; Gehring 1985). One important subclass of homeodomain proteins are NK proteins. *NK* genes are evolutionarily ancient and are part of the ANTP megacluster, which also includes *Hox* and *ParaHox* genes. NK proteins have fundamental roles in the development of mesoderm, endoderm, the nervous system and the heart in all bilaterian animals examined so far (Wotton *et al.* 2010; Holland 2013; Treffkorn *et al.* 2018) and they are also found in sponges, one of the most basal animals still alive, and potentially the sister group to all other animals (Larroux *et al.* 2007; Fortunato *et al.* 2014; Pisani *et al.* 2015; Simion *et al.* 2017).

Hmx proteins (H6 Family Homeodomain proteins, previously called Nk5 or Nkx5 proteins, see Table 1) are a key sub-family of NK proteins. In vertebrates there are usually three or four different *Hmx* genes as *Hmx4* is only found in some species (Wotton *et al.* 2010). Interestingly, *Hmx2* and *Hmx3* are usually located adjacent to each other on the same chromosome and this is also the case for *Hmx1* and *Hmx4*, suggesting that both pairs of genes arose from tandem duplication events rather than the two rounds of whole genome duplication that occurred at the base of the vertebrates (Wotton *et al.* 2010). In teleosts, there are occasionally extra duplicates of one or more of these genes as the result of the additional genome duplication in this lineage, although interestingly, the retained genes are not consistent between different teleost species (Wotton *et al.* 2010). In zebrafish there are two *hmx3* genes, *hmx3a* and *hmx3b,* but only one *hmx1*, *hmx2* and *hmx4* gene.

Previous research has shown that Hmx2 and Hmx3 have crucial functions in ear development in mouse and our recent work shows that this is also the case for Hmx3a in zebrafish (Wang *et al.* 1998; Wang *et*

al. 2001; Wang *et al.* 2004; Wang and Lufkin 2005; Hartwell *et al.* 2019). In mouse, both Hmx2 and Hmx3 mutants have ear defects and these are more severe in double mutants (Wang *et al.* 2001; Wang *et al.* 2004). *Hmx2* and *Hmx3* are also required for correct specification of the mouse hypothalamus (Wang *et al.* 2004) and morpholino knock-down experiments have suggested that they are required for correct lateral line development in zebrafish (Feng and Xu 2010). *Hmx2* and *Hmx3* are also expressed in two distinct domains in mouse spinal cord but the spinal cord functions of these genes are unknown (Bober *et al.* 1994; Wang *et al.* 2000; Wang *et al.* 2004).

Here we show that zebrafish *hmx2* and *hmx3a* are co-expressed in spinal dI2 and V1 interneurons, whereas *hmx3b*, *hmx1* and *hmx4* are not expressed in spinal cord. Using mutational analyses, we demonstrate that, in addition to its role in ear development, Hmx3a is required for correct specification of a subset of spinal cord interneuron neurotransmitter phenotypes as well as lateral line progression and viability (survival to adulthood). Our data suggest that in the absence of Hmx3a function, a subset of dI2 spinal interneurons switch their neurotransmitter phenotype from glutamatergic (excitatory) to GABAergic (inhibitory). This is important because currently very little is known about how dI2 spinal interneuron neurotransmitter phenotypes are specified, or indeed how spinal cord excitatory neurotransmitter phenotypes in general are specified, and if neurons do not acquire the correct neurotransmitter phenotypes, they cannot function appropriately in spinal cord circuitry.

Surprisingly, despite the fact that *hmx2* and *hmx3a* have similar expression patterns during embryonic development and both genes are required for correct ear development in mouse, we have not uncovered any requirement for Hmx2, by itself, in viability, ear development, lateral line progression or specification of spinal cord interneuron neurotransmitter phenotypes in zebrafish. Zebrafish *hmx2* mutants are viable and have no obviously abnormal phenotypes in these sensory structures and neurons that require Hmx3a, even when almost all of the *hmx2* locus is deleted. (In our most severe mutant allele,

hmx2^{SU39}, only 84 nucleotides of 5' and 60 nucleotides of 3' coding sequence remain). In addition, zebrafish embryos homozygous for deletions of both *hmx2* and *hmx3a* have identical phenotypes to severe *hmx3a* single mutants. However, mutating *hmx2* in hypomorphic *hmx3a* mutants, that usually develop normally, results in abnormal ear and lateral line progression phenotypes, suggesting that while Hmx2 cannot compensate for loss of Hmx3a, it does function in these developmental processes, although to a much lesser extent than Hmx3a. Our results also suggest that Hmx3a does not require its homeodomain for its roles in viability or embryonic development. This is very surprising, as homeodomain proteins usually function by binding DNA through their homeodomain and regulating gene expression. In contrast, our mutational analyses suggest that Hmx3a only requires its N terminal-domain for its vital functions in viability and sensory organ and spinal cord interneuron development.

Materials and Methods

Ethics statement

All zebrafish experiments in this research were carried out in accordance with the recommendations and approval of either the UK Home Office or the Syracuse University IACUC committee.

Zebrafish husbandry and fish lines

Zebrafish (*Danio rerio*) were maintained on a 14-h light/10-h dark cycle at 28.5°C. Embryos were obtained from natural paired and/or grouped spawnings of wild-type (WT; AB, TL or AB/TL hybrid) fish, heterozygous or homozygous *hmx2*, *hmx3a* or *hmx2/3a* mutants (created as part of this study and reported here, see Fig. 4), *Tg(evx1:EGFP)^{SU1}* or *Tg(evx1:EGFP)^{SU2}* transgenic fish (Juárez-Morales *et al.* 2016)*, Tg(UAS:mRFP)* transgenic fish (Balciuniene *et al.* 2013), heterozygous *mindbomb1 (mib1^{ta52b})* mutants (Jiang *et al.* 1996) or heterozygous or homozygous *hmx3a^{sa23054}* mutants (Kettleborough *et al.* 2013).

CRISPR mutagenesis and screening

The *hmx3a^{SU3}* allele was described previously (Hartwell *et al.* 2019). With the exception of *hmx3a^{sa23054}* (generated in the Zebrafish Mutation Project and obtained from ZIRC), we created all of the other *hmx2, hmx3a* and *hmx2/3a* double deletion mutants described in this paper using CRISPR mutagenesis. For all alleles, other than the *hmx2^{MENTHU}* allele, we designed and synthesized single guide RNA (sgRNA) and *Cas9* mRNA as in (Hartwell *et al.* 2019). For the *hmx2^{MENTHU}* allele, we designed the crRNA using the Microhomology-mediated End joining kNockout Target Heuristic Utility (MENTHU) tool (version 2.1.2), in the Gene Sculpt Suite (Ata *et al.* 2018; Mann *et al.* 2019). The MENTHU allele crRNA design was verified with CHOPCHOP (version 3.0.0) (Montague *et al.* 2014; Labun *et al.* 2016; Labun *et al.*

2019) and the CRISPR-Cas9 gRNA design checker tool (Integrated DNA Technologies). The *hmx2^{MENTHU}* crRNA was purchased together with a universal 67mer tracrRNA (1072533) and Alt-R S.p. Cas9 Nuclease V3 (1081058) from integrated DNA Technologies. See Table 2 for gRNA sequences. *hmx2^{SU35}, hmx2^{SU36}, hmx2^{SU37}* and *hmx3a^{SU42}* alleles were all generated with a single sgRNA. *hmx2^{MENTHU}* was generated with a single crRNA. *hmx2^{SU38}, hmx2^{SU39}, hmx2;hmx3a^{SU44}* and *hmx2;hmx3a^{SU45}* alleles are all deletions, generated by combinatorial use of two sgRNAs. The *hmx2^{SU38}*, sgRNAs flank the homeodomain. For *hmx2^{SU39}* we used the same 3' sgRNA and a more 5' sgRNA. To make *hmx2;hmx3a* double deletion alleles we designed sgRNAs that flanked the two genes, which are adjacent on chromosome 17. In all cases except the *hmx2^{MENTHU}* mutant, we injected 2-4 nl of a mixture of 200 ng/µl of each sgRNA + 600 ng/µl *nls-ZCas9-nls* mRNA into the single-cell of a one-cell stage AB WT embryo. To create the *hmx2^{MENTHU}* mutant allele, we injected 1 nl of a 5 µM crRNA:tracrRNA:Cas9 ribonucleoprotein complex into the single-cell of a very early one-cell stage embryo from an incross of heterozygous *hmx3a^{SU42}* fish. The 5 µM crRNA:tracrRNA:Cas9 ribonucleoprotein complex was synthesized as described in (Hoshijima *et al.* 2019). *hmx3a^{SU43}* is a *hsp70:Gal4* knock-in allele. We co-injected a donor template containing *Gal4*, under the control of a minimal *hsp70* promoter, (Mbait-hsp70:Gal4, a kind gift of Dr Shin-ichi Higashijima (Kimura *et al.* 2014)) with two sgRNA molecules: one specifically targeting *hmx3a* (see Table 2), and one, (*Mbait* sgRNA (GGCTGCTGCGGTTCCAGAGG)), specifically linearizing the donor template *in vivo*, into the single-cell of one-cell stage embryos from an incross of heterozygous *Tg(UAS:mRFP)* fish. For these experiments, embryos were injected with 2-4 nl of a mixture of 130 ng/µl of each sgRNA + 180 ng/µl *nlz-ZCas9-nls* mRNA + 66 ng/µl *pMbait-hsp70:Gal4* donor DNA. We screened injected embryos for RFP fluorescence in patterns consistent with *hmx3a* expression (i.e. ear, lateral line primordium and/or spinal cord) from 1 day post fertilization (dpf) onwards and raised injected embryos displaying appropriate expression patterns to adulthood. We then assessed germline transmission by outcrossing to

heterozygous or homozygous *Tg(UAS:mRFP)* fish. Gal4 expression in *hmx3a*[SU43] recapitulates *hmx3a* spinal expression but is not expressed in the ear or lateral line primordium (data not shown).

We identified founder fish for **hmx2**[SU35]**, hmx2**[SU36]**, hmx2**[SU37] and **hmx3a**[SU42] alleles using high resolution melt analysis (HRMA), and the supermix and amplification programs described in (Hartwell *et al.* 2019). For the PCRs described below, we used Phusion High-Fidelity DNA Polymerase (M0530L, NEB) unless otherwise stated. HRMA primers and PCR primers for sequencing are provided in Table 2.

We used the following PCR conditions to identify **hmx2**[SU38] founder fish: 98.0ºC for 30 seconds, 35 cycles of: 98.0ºC for 10 seconds, 67.0ºC for 30 seconds, 72.0ºC for 40 seconds, followed by a final extension at 72.0ºC for 5 minutes. We distinguished the mutant allele by gel electrophoresis on a 1% TAE agarose gel (110V for 30 minutes). The WT allele generated a 1098 bp product, compared with a 671 bp mutant allele product.

We used nested PCR to identify **hmx2**[SU39] founder fish, with the following conditions: Nested PCR 1: 98.0ºC for 30 seconds, 35 cycles of: 98.0ºC for 10 seconds, 69.0ºC for 20 seconds, 72.0ºC for 75 seconds, followed by a final extension at 72.0ºC for 5 minutes. The mutant allele was distinguished by gel electrophoresis on a 1% TAE agarose gel (110V for 30 minutes). The WT allele generated a 2012 bp product (which may or may not be detected on the gel), compared with a 576 bp mutant allele product. We then diluted the nested 1 PCR product 1:5 in sterile distilled water and performed the Nested PCR 2 reaction using the following conditions: 98.0ºC for 30 seconds, 35 cycles of: 98.0ºC for 10 seconds, 66.0ºC for 20 seconds, 72.0ºC for 60 seconds, followed by a final extension at 72.0ºC for 5 minutes. The mutant allele was distinguished by gel electrophoresis on a 1% TAE agarose gel (110V for 30 minutes). The WT allele generated a 1758 bp product compared with a 322 bp mutant allele product.

We identified **hmx2**^{MENTHU} F0 embryos by PCR, followed by sequencing with the FW primer that generated the amplicon (Table 2). The PCR was performed on DNA extracted from individual embryos using the following conditions: 98.0°C for 30 seconds, 35 cycles of: 98.0°C for 10 seconds, 64.0°C for 20 seconds, 72.0°C for 15 seconds, followed by a final extension at 72.0°C for 5 minutes. We assayed that the PCR was successful by gel electrophoresis on a 2.5% TBE agarose gel (100V for 40 minutes). The PCR generates a 155 bp product. The PCR product was purified using EZ-10 Spin Column PCR Products Purification Kit (BS664, Bio Basic) and eluted in 30 µl sterile water prior to sequencing.

We used either assessment of germline transmission, as described above, or PCR to identify **hmx3a**^{SU43} founder fish. PCR conditions were: 98.0°C for 30 seconds, 35 cycles of: 98.0°C for 10 seconds, 69.0°C for 20 seconds, 72.0°C for 60 seconds, followed by a final extension at 72.0°C for 5 minutes. The mutant allele was distinguished by gel electrophoresis on a 1% TAE agarose gel (110V for 30 minutes). A 1471 bp PCR product was only generated by fish heterozygous for the allele. It was not produced from WT animals since the reverse primer only recognizes the inserted donor DNA sequence.

We identified **hmx2;hmx3a**^{SU44} and **hmx2;hmx3a**^{SU45} founder fish by nested PCR, using the following conditions: <u>Nested PCR 1:</u> 98.0°C for 30 seconds, 35 cycles of: 98.0°C for 10 seconds, 67.0°C for 20 seconds, 72.0°C for 30 seconds, followed by a final extension at 72.0°C for 5 minutes. The mutant allele was distinguished by gel electrophoresis on a 1% TAE agarose gel (110V for 30 minutes). The WT product was too large to be generated by these PCR conditions, so only heterozygous animals are detected by the presence of a 514 bp product on the gel. We then diluted the nested 1 PCR 1:5 in sterile distilled water and performed the <u>Nested PCR 2</u> reaction using the following conditions: 98.0°C for 30 seconds, 35 cycles of: 98.0°C for 10 seconds, 66.0°C for 20 seconds, 72.0°C for 30 seconds, followed by a final extension at 72.0°C for 5 minutes. The mutant allele was distinguished by gel electrophoresis on

a 1% TAE agarose gel (110V for 30 minutes). Again, the WT product was too large to be generated by these PCR conditions, so only heterozygous animals were detected by the presence of a 445 bp product.

Once stable lines were established, we identified *hmx2^{SU35}* fish by PCR, followed by sequencing (the mutation introduces a 1 bp insertion that cannot be resolved by restriction digestion, and we cannot distinguish heterozygotes from homozygotes using HRMA). We performed this PCR using Taq DNA Polymerase (M0320S, NEB) and the following conditions: 95.0°C for 30 seconds, 35 cycles of: 95.0°C for 20 seconds, 52.0°C for 30 seconds, 68.0°C for 45 seconds, followed by a final extension at 68.0°C for 5 minutes. We used HRMA and the conditions described above to identify *hmx2^{SU36}* stable mutants. Homozygous mutants segregate from heterozygous animals by the scale of their deflection in the HRMA plot. We identified *hmx2^{SU37}* mutants by performing the PCR used to sequence *hmx2^{SU35}* stable mutants (see above). When we analyzed the products on a 1% TAE gel (110V for 30 minutes), the WT allele generated a 580 bp product, compared with a 528 bp mutant product. We identified *hmx2^{SU38}* stable mutants using the same PCR conditions initially used to identify founders (see above). We used the same nested PCR conditions to identify *hmx2^{SU39}* mutants. However, the WT product was not always visible on the gel. Therefore, we also performed a WT amplicon PCR identical to that described above for identifying stable *hmx2^{SU35}* fish, as this genomic region is only present in WT and heterozygous animals.

We identified stable *hmx3a^{SU42}* mutants by PCR, using Taq DNA Polymerase (M0320S, NEB) and the following conditions: 94.0°C for 2 minutes, 35 cycles of: 94.0°C for 30 seconds, 64.9°C for 30 seconds, 72.0°C for 30 seconds, followed by a final extension at 72.0°C for 2 minutes. Whilst the mutant PCR product (321 bp) could sometimes be distinguished from the WT product (331 bp) by running on a 2% TBE gel (100V for 55 minutes), the mutation also deletes a BanI restriction site. Following digestion with BanI (R0118S, NEB), the products were run on a 2% TBE gel (100V for 40 minutes). The WT amplicon digested to completion, producing 120 bp + 211 bp bands, whereas the mutant product did not

cut. We identified stable **hmx3a^{SU3}** mutants by running the same PCR used to identify *hmx3a^{SU42}* mutants. The insertion in *hmx3a^{SU3}* was easily visualized on a 2% TBE gel. The WT product was 331 bp, compared to a mutant product of 400 bp. Since the PCR used to detect **hmx3a^{SU43}** mutants was specific to the inserted donor DNA, and the WT amplicon in *hmx2;hmx3a^{SU44}* and *hmx2;hmx3a^{SU45}* mutants was too large to detect by standard PCR, for these alleles we also performed a WT amplicon PCR to distinguish WTs from hets. The WT amplicon PCR was identical to that performed prior to BanI digestion on *hmx3a^{SU42}* mutants (see above). For **hmx3a^{SU43}**, the WT amplicon PCR results were compared to the PCR results (identical PCR to that first used to identify founders, see above), and for **hmx2;hmx3a^{SU44}** and **hmx2;hmx3a^{SU45}**, the WT amplicon PCR results were compared to the Nested 2 PCR results (identical Nested 2 PCR to that first used to identify founders – see above).

In all cases, stable F1 heterozygous fish were confirmed by sequencing. See Fig. 4 for details of how individual mutant alleles differ from one another. To further confirm the mutant sequences of *hmx2^{SU39}* and *hmx3a^{SU42}*, we extracted total RNA from embryos produced by incrosses of homozygous viable adults using TRIzol Reagent (15596018, ThermoFisher Scientific) and the RNEasy Mini Kit (74104, Qiagen). Total RNA was converted to cDNA using the iScript cDNA synthesis kit (1708891, Bio-Rad). We performed transcript-specific PCRs using the following primers and conditions: *hmx2*-FW: TGAACTGTTATGAGACGAGAATGAA and *hmx2*-RV: GTGTATTTTGTACGTCTTAGTGTGTGT (PCR: 98.0°C for 30 seconds; 35 cycles of: 98.0°C for 10 seconds, 64.2°C for 20 seconds; 72.0°C for 30 seconds, followed by final extension at 72.0°C for 5 minutes) or *hmx3a*-FW: AACCGCGTTTAAGTTCCCATTG and *hmx3a*-RV: GTGCGAGTAGTAAACCGGATGAG (PCR: 98.0°C for 30 seconds; 35 cycles of: 98.0°C for 10 seconds, 71.0°C for 20 seconds; 72.0°C for 30 seconds, followed by final extension at 72.0°C for 5 minutes). We then confirmed these homozygous mutant transcript sequences by sequencing.

Morpholino injections

For single knockdown (SKD) experiments, 3.5 nl of a mixture containing either 2 ng/nl of a translation-blocking *hmx2* morpholino (5' TTCCGCTGTCCTCCGAATTATTCAT) or 2 ng/nl of a translation-blocking *hmx3a* morpholino (5' ACGTATCCTGTGTTGTTTCGGGCAT) plus 5 ng/nl of a control zebrafish p53 morpholino (5' GCGCCATTGCTTTGCAAGAATTG) was injected into the single-cell of a one-cell stage WT embryo (all morpholinos obtained from Gene Tools). For double knockdown (DKD) experiments, 3.5 nl of a mixture containing 2 ng/nl of both *hmx* morpholinos plus 5 ng/nl of the control zebrafish p53 morpholino was injected. DKD but not SKD embryos exhibit delayed development from somitogenesis stages onwards when compared to uninjected controls. To circumvent this, they were incubated at 32°C from 9 hours post fertilization (h) onwards. This ensured that control and injected embryos reached the desired developmental stage of 27 h at approximately the same time. The lateral line primordium does not migrate in DKD animals, so this could not be used to stage injected embryos. Instead, these embryos were visually inspected and fixed when they displayed the same head-trunk angle, head size and eye size as prim-staged uninjected control embryos (Kimmel 1995). Migration of the lateral line primordium is unaffected in SKD embryos, so these were prim-staged prior to fixing for experiments (Kimmel 1995). Morpholino injections always produce a spectrum of phenotypes, since it is hard to ensure that every cell receives the same dose. Therefore, prior to fixing at 27 h, we removed any embryos with severely abnormal morphology (stunted length and/or severely developmentally delayed, likely caused by receiving too much morpholino). Embryos injected with *hmx2/3a* morpholinos (SKD and DKD) display a slight curled-tail-down morphology. Embryos that lacked this morphology (and may therefore not have received any or sufficient morpholino) were also removed before fixing.

For the mRNA rescue experiments, we co-injected both morpholinos (at the same volume and dose described above), together with a total dose of up to 500 pg of morpholino resistant (MOR) *hmx3a* mRNA. This mRNA had 7 nucleotides altered in the *hmx3* morpholino recognition sequence. Each

change was in the third nucleotide of a codon. This codon wobble was used so that the same amino acid was encoded in each case, but the mRNA would not be recognized by the morpholino. The protein encoded by the injected RNA is therefore the same as endogenous Hmx3a.

WT *hmx3a*:   ATG CCC GAA ACA ACA CAG GAT ACG (Met, Pro, Glu, Thr, Thr, Gln, Asp, Thr)

MOR-hmx3a: ATG CCG GAG ACT ACT CAA GAC ACC (Met, Pro, Glu, Thr, Thr, Gln, Asp, Thr)

To assess whether genetic compensation occurs in either $hmx2^{SU39}$ or $hmx3a^{SU42}$ mutants, which lack obvious phenotypes, or $hmx3a^{SU3}$ mutants, which have milder spinal cord phenotypes than *hmx2/3a* DKD embryos, we injected the same dose of either *hmx2 + p53* MOs ($hmx2^{SU39}$) or *hmx3a + p53* MOs ($hmx3a^{SU42}$, $hmx3a^{SU3}$) as described above, into the single-cell of one-cell stage embryos generated from incrosses of heterozygous $hmx2^{SU39}$, $hmx3a^{SU42}$ or $hmx3a^{SU3}$ parents respectively. If genetic compensation is occurring, the upregulated compensating gene(s) will not be knocked-down by the *hmx* morpholino and the phenotype of homozygous mutants should be unchanged. In contrast, WT and heterozygous animals, which contain at least one WT copy of the respective *hmx* gene will be susceptible to the *hmx* morpholino and should exhibit stronger, morphant-like phenotypes. For these experiments, while we removed any embryos with severely abnormal morphology, we did not remove embryos that lacked the curled-tail-down morphology, incase these were morpholino-resistant mutant embryos. After fixing, we performed an *in situ* hybridization for the glutamatergic marker, *slc17a6a/b*. We visually inspected the embryos on a dissecting microscope and categorized them as either the stronger, morphant-like phenotype (large reduction in the number of *slc17a6a/b*-expressing cells) or a more subtle phenotype (WT-like in the case of $hmx2^{SU39}$ and $hmx3a^{SU42}$ or a smaller reduction in the number of *slc17a6a/b*-expressing cells in the case of $hmx3a^{SU3}$ embryos). Embryos within each class were then genotyped as described in the CRISPR mutagenesis section above.

Genotyping

We isolated DNA for genotyping from both anesthetized adult fish and fixed embryos via fin biopsy or head dissections, respectively. For assaying ear phenotypes, we dissected tail tips instead. We genotyped the *hmx* CRISPR mutants as described above. For *mib1^{ta52b}* and *hmx3a^{sa23054}* mutants, we used KASP assays designed by LGC Genomics LLC. KASP assays use allele-specific PCR primers, which differentially bind fluorescent dyes that we quantified with a BioRad CFX96 real-time PCR machine to distinguish genotypes. The proprietary primers used were: *mib_ta52b* and *hmx3a_sa23054*. Heads or tail tips of fixed embryos were dissected in 70% glycerol/30% distilled water with insect pins. Embryo trunks were stored in 70% glycerol/30% distilled water at 4°C for later analysis. For all experiments except Phalloidin staining experiments, DNA was extracted via the HotSHOT method (Truett *et al.* 2000) using 10 µl of 50 mM NaOH and 1 µl of 1M Tris-HCl (pH-7.4). For Phalloidin staining experiments, the tail up until the end of the yolk extension was dissected in 70% glycerol/30% distilled water as described above and transferred to PBST. The PBST was then replaced with 50 µl of DNA extraction buffer (10 mM Tris, pH 8.0, 10 mM EDTA, 200 mM NaCl, 0.5% SDS, 200 µg/ml Proteinase K (Proteinase K, recombinant, PCR Grade, 3115879001, Sigma Aldrich), before incubating for 3 hours in a 55°C water bath. The samples were vortexed periodically to ensure thorough digestion of the tissue. Subsequently, the Proteinase K was inactivated by heating the samples for 10 minutes at 100°C, prior to centrifuging for 20 minutes at 14,000 rpm at room temperature to pellet debris. The supernatant was transferred to sterile microcentrifuge tubes before adding 20 µg UltraPure Glycogen (10814010, ThermoFisher Scientific) and 2 volumes of ice-cold RNase-free ethanol. Samples were precipitated at -20°C overnight. Genomic DNA was recovered by centrifugation at +4°C, followed by washing with 70% RNase-free ethanol and further centrifugation at +4°C. After carefully removing the ethanolic supernatant, the pellets were air dried for 5-10 minutes at room temperature before resuspending in 15 µl of sterile distilled water.

<u>*in situ* hybridization and immunohistochemistry</u>

We fixed embryos in 4% paraformaldehyde / phosphate-buffered saline (PBS) and performed single and double *in situ* hybridizations and immunohistochemistry plus *in situ* hybridization double labeling experiments as previously described (Concordet *et al.* 1996; Batista *et al.* 2008). Sources of *in situ* hybridization probes are provided in Table 1. To amplify *in situ* hybridization probe templates for *hmx1* and *hmx3b*, we created zebrafish cDNA from 27 h WT zebrafish embryos. We extracted total RNA by homogenizing 50–100 mg of embryos in 1 mL of TRIzol reagent (Ambion, 15596-026). We confirmed RNA integrity (2:1 ratio of 28S:18S rRNA bands) and quality (A260/A280 ratio of ~2.0) using agarose gel electrophoresis and spectrophotometry respectively. We synthesized cDNA using Bio-Rad iScript Reverse Transcription Supermix kit (Bio-Rad, 170-8891). We amplified *hmx1* sequence from the cDNA using Phusion High-Fidelity DNA Polymerase (M0530L, NEB), primers *hmx1*-FW: CTGGTATATTTGCTCAAGACATGC and *hmx1*-RV: GCTTCTGCTGAACACAGTTCG and PCR conditions: 98.0ºC for 10 seconds, 30 cycles of: 98.0ºC for 6 seconds; 57.0ºC for 30 seconds and 72.0ºC for 30 seconds, followed by a final extension for 45 seconds at 72.0ºC. The PCR product was assessed on a 1% TAE gel, before purifying with QIAquick PCR Purification Kit (28104, Qiagen). We used Taq DNA Polymerase (M0320S, NEB) to add 3'A overhangs prior to TOPO TA-cloning (K4600-01, Invitrogen). We then performed colony PCR using the same PCR primers and conditions used to amplify the *hmx1* sequence from cDNA. We extracted plasmid DNA from positive colonies using QIAprep Spin Miniprep Kit (27104, Qiagen) and then verified the sequence using standard SP6 and T7 primers for Sanger sequencing. To make the anti-sense RNA riboprobe, we linearized DNA with HindIII-HF (R3104S, NEB) and transcribed with T7 RNA Polymerase (10881767001, Roche). We used a PCR-based DNA template to make the *hmx3b* ISH probe. The reverse primer contains the T3 RNA Polymerase minimal promoter sequence (underlined). We used primers *hmx3b*-FW: GTGTGCCCGTCATCTACCAC and

hmx3b-RV: <u>AATTAACCCTCACTAAAGGGA</u>TGAAGATGATGAAGATGCGCAAC, 27 h WT cDNA, Phusion High-Fidelity DNA Polymerase (M0530L, NEB) and PCR conditions: 94.0°C for 3 minutes, 35 cycles of: 94.0°C for 30 seconds, 56.5°C for 30 seconds and 72.0°C for 1.5 minutes, followed by a final extension step of 72.0°C for 10 minutes. We purified the template through phenol:chloroform:isoamyl alcohol extraction and precipitation with 0.2 M NaCl and ice-cold ethanol prior to *in situ* probe synthesis using 1 μg purified PCR product, T3 RNA Polymerase (11031171001, Roche) and DIG RNA Labeling Mix (11277073910, Roche).

Embryos older than 24 h were usually incubated in 0.003% 1- phenyl-2-thiourea (PTU) to prevent pigment formation. For some experiments we added 5% of dextran sulfate to the hybridization buffer (* in Table 3). Dextran sulfate can increase specific staining in *in situ* hybridization experiments as it facilitates molecular crowding (Ku *et al.* 2004; Lauter *et al.* 2011).

In cases where we did not detect expression of a particular gene in the spinal cord, we checked for low levels of expression by exposing embryos to prolonged staining. In some cases, this produced higher background (diffuse, non-specific staining), especially in the hindbrain, where ventricles can sometimes trap anti-sense riboprobes.

To determine neurotransmitter phenotypes we used probes for genes that encode proteins that transport or synthesize specific neurotransmitters as these are some of the most specific molecular markers of these cell fates (Higashijima *et al.* 2004b; Higashijima *et al.* 2004c and references therein). A mixture of probes to *slc17a6a* and *slc17a6b* (previously called *vglut*), which encode glutamate transporters, was used to label glutamatergic neurons (Higashijima *et al.* 2004b; Higashijima *et al.* 2004c). GABAergic neurons were labeled using probes to *gad1b* (probes previously called *gad67a* and *gad67b*) (Higashijima

et al. 2004b; Higashijima *et al.* 2004c). The *gad1b* gene encodes for a glutamic acid decarboxylase, which is necessary for the synthesis of GABA from glutamate. A mixture of probes (*glyt2a* and *glyt2b*) for *slc6a5* (previously called *glyt2*) was used to label glycinergic cells (Higashijima *et al.* 2004b; Higashijima *et al.* 2004c). *slc6a5* encodes for a glycine transporter necessary for glycine reuptake and transport across the plasma membrane.

The antibodies that we used for fluorescent *in situ* hybridization were mouse anti-Dig (200-002-156, Jackson ImmunoResearch, 1:5000) and rabbit anti-Flu (A889, Invitrogen, 1:2500). These were detected using secondary antibodies: goat anti-rabbit-HRP (G-21234, ThermoFisher Scientific, 1:750) and goat anti-mouse-HRP (G-21040, ThermoFisher Scientific, 1:750) and Tyramide SuperBoost Kits B40922 and B40915 (ThermoFisher Scientific).

For double fluorescent *in situ* hybridization and immunohistochemistry, after detection of the *in situ* hybridization reaction using Tyramide SuperBoost Kit B40915 (with HRP, Goat anti-mouse IgG and Alexa Fluor 594 Tyramide), embryos were washed 8 × 15min in PBST (PBS with 0.1% Tween-20) and incubated in Image-iT FX Signal Enhancer (ThermoFisher Scientific, I36933) for 30 mins at room temperature. Immunohistochemistry was performed using chicken polyclonal anti-GFP primary antibody (Ab13970, Abcam, 1:500) and a Goat anti-chicken IgY (H+L), Alexa Fluor 488 secondary antibody (A-11039, ThermoFisher Scientific, 1:1000).

Phalloidin staining

4 d old embryos generated from incrosses of heterozygous *hmx2*[SU39] or *hmx2;hmx3a*[SU44] parents were fixed and processed for phalloidin staining as described in (Hartwell *et al.* 2019). Stained embryos were

stored in DABCO (2% w/v 1,4-Diazabicyclo[2.2.2]octane (D27802, Sigma Aldrich) in 80% glycerol in sterile distilled water).

qPCR analyses

We collected embryos from incrosses of AB wild-type parents and flash-froze them at 16-cell, 6 h, 14 h, 27 h and 48 h stages. We collected 40-50 embryos per biological replicate per developmental stage and performed duplicate biological replicates. We isolated total RNA by homogenizing each sample in 1 mL of TRIzol reagent (Ambion, 15596-026). Following chloroform extraction, we added 20 µg UltraPure Glycogen (10814010, ThermoFisher Scientific) to the aqueous phase followed by one volume of RNase-free ethanol. We performed RNA purification and genomic DNA removal using the Monarch Total RNA Miniprep Kit (T2010S, NEB) following manufacturer's instructions for purifying TRIzol-extracted samples. RNA concentration was measured using Nanodrop 2000 (ND2000, ThermoFisher Scientific), prior to synthesizing cDNA using the Bio-Rad iScript Reverse Transcription Supermix kit (Bio-Rad, 170-8891). We also included controls lacking Reverse-Transcriptase to assay for the presence of genomic DNA contamination. qPCR was performed in triplicate for each sample using iTaq Universal SYBR Green Supermix (1725121, Bio-Rad) and a BioRad CFX96 real-time PCR machine. The following qPCR primers were used:

hmx2-qPCR-FW: CCCATTTCAAGTTTCACGATCCAGTC,

hmx2-qPCR-RV: TGCTCCTCTTTGTAATCCGGTAG,

hmx3a-qPCR-FW: TTGATGGCAGCTTCTCCCTTTC,

hmx3a-qPCR-RV: ACTCTTCTTCCAGTCGTCTATGC,

mob4-qPCR-FW: CACCCGTTTCGTGATGAAGTACAA,

mob4-qPCR-RV: GTTAAGCAGGATTTACAATGGAG.

The *hmx2* and *hmx3a* primers were generated in this study. The *mob4* primers were generated by Hu and colleagues (Hu *et al.* 2016). They demonstrated that *mob4* is a more effective reference gene than *actb2*

across a broad range of zebrafish developmental stages, including early stages where only maternal RNAs should be present (Hu *et al.* 2016). To generate amplification data the program used was: 95.0ºC for 30 seconds, 40 cycles of: 95.0ºC for 5 seconds, 63.3ºC (*hmx2*)/64.5ºC (*hmx3a*)/60.0ºC (*mob4*) for 30 seconds, with imaging after each cycle. To assay amplification specificity and exclude false positives from primer dimers we then generated melt data using: 65.0ºC for 30 seconds, 40 cycles of: 65.0ºC-95.0ºC, + 0.5ºC/second increment, with each increment held for 5 seconds prior to imaging, 95.0ºC for 15 seconds.

Screening lateral line and otolith phenotypes

We examined whether any of the *hmx* mutants generated in this study had lateral line and/or fused otolith phenotypes, as reported for *hmx2;hmx3a* double-knockdown embryos (Feng and Xu 2010). To assay live lateral line phenotypes, we anaesthetized embryos from incrosses of heterozygous mutant fish in 0.016% tricaine (A5040, Sigma Aldrich) in embryo medium (EM, 5 mM NaCl, 0.17 mM KCl, 0.33 mM $CaCl_2$ $2H_2O$, 0.33 mM $MgSO_4 \cdot 7H_2O$, 0.017% w/v (0.7mM) HEPES pH 7.8 and 0.00004% methylene blue in autoclaved reverse osmosis water) and mounted them on coverslip bridges (2 x 22 mm square glass coverslips (16004-094, VWR) glued together on either side of a 24 x 60 mm glass cover slip (12460S, ThermoFisher Scientific), overlaid with a third 22 mm square glass coverslip). Using a Zeiss Axio Imager M1 compound microscope, we located the tip of the lateral line primordium and counted the somite number adjacent to this position. We also used this method routinely to determine the developmental stage of embryos prior to fixing for *in situ* hybridization. To assay lateral line phenotypes in fixed embryos, we performed *in situ* hybridizations for *hmx3a* or *krt15* (both of which label the migrating primordium and neuromasts) and then determined the lateral line position as in live embryos. To examine live otolith phenotypes, embryos were raised until 3 dpf, before anaesthetizing (as for assessing live lateral line phenotypes) and examining the spatial location of otoliths in both ears. WT

embryos have two otoliths in each ear: one smaller, anterior (utricular) otolith, and one larger, posterior (saccular) otolith. These are separate from each other and spatially distinct. We classified otoliths as fused if only one large, amalgamated otolith was visible in a mid-ventral position within the otic vesicle (see Fig. 5U).

Imaging

Embryos were mounted in 70% glycerol:30% distilled water and Differential Interference Contrast (DIC) pictures were taken using an AxioCam MRc5 camera mounted on a Zeiss Axio Imager M1 compound microscope. Fluorescent images were taken on a Zeiss LSM 710 confocal microscope. Images were processed using Adobe Photoshop software (Adobe, Inc) and Image J software (Abramoff et al., 2004). In some cases, different focal planes were merged to show labeled cells at different medial lateral positions in the spinal cord. All images were processed for brightness-contrast and color balance using Adobe Photoshop software (Adobe, Inc.). Images of control and mutant embryos from the same experiment were processed identically. Figures were assembled using Adobe Photoshop and Adobe Illustrator (Adobe, Inc.).

Cell counts and statistics

In all cases, cell counts are for both sides of a five-somite length of spinal cord adjacent to somites 6-10. Embryos were mounted laterally with the somite boundaries on each side of the embryo exactly aligned and the apex of the somite over the middle of the notochord. This ensures that the spinal cord is straight along its dorsal-ventral axis and that cells in the same dorsal/ventral position on opposite sides of the spinal cord will be directly above and below each other. Embryos from mutant crosses were counted blind to genotype. Labeled cells in embryos analyzed by DIC were counted while examining embryos on a Zeiss Axio Imager M1 compound microscope. We identified somites 6–10 in each embryo and counted the number of labeled cells in that stretch of the spinal cord. We adjusted the focal plane as we

examined the embryo to count cells at all medial/lateral positions (both sides of the spinal cord; also see Batista *et al.* 2008; Batista and Lewis 2008; England *et al.* 2011; Hilinski *et al.* 2016; Juárez-Morales *et al.* 2016).

In some cases, cell count data were pooled from different experiments. Prior to pooling, all pairwise combinations of data sets were tested to determine if there were any statistically significant differences between them as described below. Data were only pooled if none of the pairwise comparisons were statistically significantly different from each other. In addition, as *in situ* hybridization staining can vary slightly between experiments, we only compared different mutant results when the counts from their corresponding WT sibling embryos were not statistically significantly different from each other.

To determine whether differences in values are statistically significant, data were first analyzed for normality using the Shapiro-Wilk test. Data sets with non-normal distributions were subsequently analyzed using the Wilcoxon-Mann-Whitney test (also called the Mann Whitney U test). For data sets with normal distributions, the F-test for equal variances was performed, prior to conducting either a type 2 (for equal variances) or type 3 (for non-equal variances) student's *t*-test. P values generated by Wilcoxon-Mann-Whitney, type 2 student's *t*-test and type 3 student's *t*-test are indicated by [^], [+] and [§] respectively. *n*-values for each experiment are provided in the figure legends. To assess whether mutant phenotypes occurred at Mendelian frequencies, we performed Chi-squared tests. To test whether a small number of embryos with abnormal phenotypes was statistically significantly different from zero we performed a binomial distribution test, using the cumulative distribution function, the number of embryos without mutant phenotypes, the total number of embryos examined (n) and a probability argument of n-1/n. P values >0.05 support the null hypothesis that the number of embryos with abnormal phenotypes is not statistically significantly different from zero.

<u>Microarray expression profiling experiments</u>

These experiments are described in detail in (Cerda *et al.* 2009). P values were corrected for multiple testing (Benjamini and Hochberg 1995; Gentleman *et al.* 2004; Tarraga *et al.* 2008). These data have been deposited in the NCBI Gene Expression Omnibus with accession number GSE145916.

<u>Data and Reagent Availability</u>

Plasmids and zebrafish strains are available upon request. One supplemental figure is available at FigShare. This contains an alignment of mouse and zebrafish Hmx2 and Hmx3a protein sequences. Microarray data have been previously deposited in the NCBI Gene Expression Omnibus with accession number GSE145916.

Results

hmx2 and *hmx3a* are the only *hmx* genes expressed in spinal cord.

While the expression and functions of zebrafish *hmx* genes have been analyzed during the development of sensory structures such as the eye and the ear, the expression of *hmx1*, *hmx2*, *hmx3a* and *hmx4* in the developing spinal cord has not been investigated and no expression data has previously been reported for *hmx3b,* which only appeared in more recent versions of the zebrafish genome sequence (Zv9 and above). Therefore, to determine which of the *hmx* genes are expressed in the spinal cord we performed *in situ* hybridizations for *hmx1*, *hmx2*, *hmx3a*, *hmx3b* and *hmx4* at different developmental stages (Fig. 1). At all of these stages, we observed no spinal cord expression of *hmx1*, *hmx3b* or *hmx4* (Fig. 1). However, consistent with previous reports, both *hmx1* and *hmx4* were expressed in the developing eye, ear and anterior lateral line neuromasts (Fig. 1; French *et al.* 2007; Feng and Xu 2010; Gongal *et al.* 2011; Boisset and Schorderet 2012; Marcelli *et al.* 2014). In contrast, the only expression of *hmx3b* that we observed was weak hindbrain expression at later stages of development (36-48 h; Fig. 1N', S' & X').

In contrast, *hmx2* and *hmx3a* are expressed in the spinal cord at all of the stages that we examined (Fig. 1). The spinal cord expression patterns of these two genes are very similar, with the exception that initially, *hmx2* expression appears to be weaker than *hmx3a* and it does not extend as far caudally (Fig. 1B & C). Consistent with previous reports, both of these genes are also expressed in the lateral line and developing ear (Fig. 1; Adamska *et al.* 2000; Feng and Xu 2010; Hartwell *et al.* 2019) as well as distinct regions of the brain (data not shown).

As *hmx3a* is expressed in the spinal cord, and teleost duplicate genes often have similar expression patterns, we wanted to further test whether there was any spinal cord expression of *hmx3b*. Therefore, we performed *in situ* hybridization on *mindbomb1 (mib1^{ta52b})* mutants at 24 h. *mib1* encodes an E3-

ubiquitin protein ligase required for efficient Notch signaling. Consequently, Notch signaling is lost in *mib1^{ta52b}* mutants and this causes most spinal progenitor cells to precociously differentiate into early-forming classes of spinal neurons at the expense of later-forming classes of neurons and glia (Jiang *et al.* 1996; Schier *et al.* 1996; Itoh *et al.* 2003; Park and Appel 2003; Batista *et al.* 2008). As a result, weak expression in spinal neurons is often expanded and stronger and hence easier to observe in *mib1^{ta52b}* mutants at 24 h (Batista *et al.* 2008; England *et al.* 2017). However, even in *mib1^{ta52b}* mutants we detected no expression of *hmx3b* (Fig. 1D').

<u>*hmx2* and *hmx3a* are expressed in V1 and dI2 interneurons in the spinal cord.</u>

To identify the spinal cord neurons that express *hmx2* and *hmx3a*, we performed several different double-labeling experiments. Double *in situ* hybridization with *hmx2* and *hmx3a* confirmed that these genes are co-expressed in the exact same cells in the spinal cord (Fig. 2C). Approximately half of these *hmx2* and *hmx3a (hmx2/3a)* co-expressing spinal cells also co-express *slc32a1*, which is only expressed by inhibitory (glycinergic and GABAergic) interneurons, and approximately half co-express *slc17a6a/b,* which are only expressed by excitatory (glutamatergic) interneurons (Fig. 2D & F; see materials and methods for a more detailed description of probes used to determine neurotransmitter phenotypes). In addition, the inhibitory *hmx2/3a*-expressing cells are generally more ventral than the excitatory double-labelled cells. Our previous expression-profiling of V1 interneurons, suggested that these cells might be the ventral inhibitory neurons that express *hmx3a* (Fig. 2B, for a description of these experiments see (Cerda *et al.* 2009)). Results from our lab and others, have established that V1 interneurons are the only spinal cord cells that express *engrailed1b* (*en1b*) (Higashijima *et al.* 2004a; Batista and Lewis 2008). Therefore, to confirm that V1 interneurons also express *hmx3a* we performed double *in situ* hybridizations for *hmx3a* and *en1b*. These experiments showed that all of the *en1b*-expressing spinal cells co-express *hmx3a,* and that approximately half of the *hmx2/3a*-expressing spinal cells co-express

en1b (Fig. 2E). Taken together, these data clearly identify the inhibitory *hmx2/3a*-expressing spinal cells as V1 interneurons.

As mentioned above, the glutamatergic *hmx2/3a*-expressing cells are generally located more dorsal to the inhibitory *hmx2/3a*-expressing cells. Therefore, these excitatory cells could be V0v, dI5, dI3, dI2 or dI1 interneurons (Fig. 2A; Cheng *et al.* 2005; Grossmann *et al.* 2010; Satou *et al.* 2012; Talpalar *et al.* 2013). The zebrafish embryonic spinal cord is relatively small. For example, at 27h, the dorsal-ventral axis is only about 10 cells high. As a result, the different neuronal populations are often intermingled, rather than clearly separated as they are in amniotes (e.g. see Batista *et al.* 2008; England *et al.* 2011). In addition, studies in amniotes suggest that many dorsal neurons migrate dorsally or ventrally soon after they are born (e.g. see Gross *et al.* 2002; Müller *et al.* 2002). Taken together, this means that it is hard to accurately identify dorsal spinal cord cell types by position alone. Therefore, to identify the excitatory *hmx2/3a*-expressing neurons we performed double labeling experiments with markers of different dorsal excitatory cell types. We found that all *hmx2/3a*-expressing spinal cells co-express *lhx1a* and *lhx5* (Fig. 2B & G). Lhx1a and Lhx5 are predominantly expressed by inhibitory spinal cord interneurons, but they are also expressed by dI2 interneurons, which are excitatory (Gowan *et al.* 2001; Moran-Rivard *et al.* 2001; Gross *et al.* 2002; Muller *et al.* 2002; Cheng *et al.* 2004; Wine-Lee *et al.* 2004; Muller *et al.* 2005; Alaynick *et al.* 2011; Satou *et al.* 2012). The only other excitatory neurons that might express these two *lhx* genes are V0v interneurons. Recent single-cell RNA-Seq (scRNA-Seq) data from mouse spinal cord identified possible V0 cells that expressed *lhx1a* and *lhx5*, although, given the relatively small size of this recovered population, it is not clear whether these cells were excitatory V0v and/or inhibitory V0$_D$ cells (Delile *et al.* 2019). V0v cells are the only spinal cord cells that express *evx1* and *evx2* (Juárez-Morales *et al.* 2016 and references therein). Therefore, we tested whether there was any co-expression of *hmx3a* and *evx1/2* using double *in situ* hybridization as well as immunohistochemistry for GFP and *in situ* hybridization for *hmx3a* in *Tg(evx1:EGFP)*[SU1] and *Tg(evx1:EGFP)*[SU2] embryos. However, we did

not observe any co-expression in any of these experiments (Fig. 2H and data not shown). Therefore, we are confident that the excitatory *hmx2/3a*-expressing spinal cells are dI2 interneurons. Consistent with this, recent mouse scRNA-Seq spinal cord data suggests that mouse *Hmx2* and *Hmx3* are also expressed in V1 and dI2 spinal cord interneurons (Delile *et al.* 2019).

Knock-down experiments suggest that Hmx2 and Hmx3a may be redundantly required for correct
specification of a subset of spinal interneuron glutamatergic phenotypes.

As an initial step to try and identify the function(s) of Hmx2 and Hmx3a in spinal cord development we performed morpholino knock-down experiments. As previous analyses suggested that these genes have redundant roles in ear and lateral line development (Feng and Xu 2010), we designed and injected morpholinos for both of these genes (see materials and methods). In embryos co-injected with the two morpholinos (*hmx2;hmx3a* double knock-down (DKD) animals), we observed stalled lateral line progression and fused otoliths in the ears. Normally there is an anterior (utricular) and a posterior (saccular) otolith in each ear, but in DKD embryos there was just one fused otolith in a medio-ventral region of each ear (data not shown). When we analyzed spinal cord phenotypes, we detected no change in the number of *hmx3a-* or *en1b*-expressing cells in DKD embryos, suggesting that dI2 and V1 interneurons still form in normal numbers (Fig. 3A-F, Table 3). However, when we examined markers of neurotransmitter phenotypes, we observed a reduction in the number of spinal excitatory (glutamatergic) cells and a corresponding increase in inhibitory cells. As *hmx2* and *hmx3a* are only expressed by dI2 neurons (which are glutamatergic) and V1 neurons (which are inhibitory) in the spinal cord, this suggested that at least some dI2 interneurons had switched their neurotransmitter phenotype from glutamatergic to inhibitory (Fig. 3G-L, Table 3). To confirm the specificity of these knock-down results, we tested whether co-injecting a morpholino-resistant *hmx3a* mRNA with the *hmx2* and *hmx3a* morpholinos could rescue the spinal cord phenotype of DKD embryos. Indeed, we were able to obtain a partial rescue in the number of glutamatergic neurons (Fig. 3M-O and Table 3). Injections of higher

amounts of mRNA led to embryo death, probably because of the toxic effects of injecting considerable amounts of both mRNA and morpholinos into the embryos during early development.

Mutational analyses suggest that Hmx2 is not, by itself, required, and that Hmx3a does not require its DNA-binding homeodomain, for viability, correct migration of lateral line primordium, or correct development of ear otoliths or a subset of spinal cord interneurons.

To further and more robustly test the hypothesis that *hmx2* and *hmx3a* are required for the correct specification of a subset of spinal cord interneuron neurotransmitter phenotypes we created CRISPR mutants in each of these genes, targeting a region upstream of the homeobox (see materials and methods; Fig. 4). We also obtained a *hmx3a*[sa23054] allele from the Sanger zebrafish mutation project (Kettleborough *et al.* 2013) that introduces a stop codon upstream of the homeobox.

Our analyses of homozygous mutant embryos demonstrate that *hmx3a*[SU3] and *hmx3a*[SU43] mutants have fused otoliths, stalled lateral line progression and are homozygous lethal (Fig. 4, Fig. 5O, U, Table 4 and data not shown, also see (Hartwell *et al.* 2019) for a detailed description of the *hmx3a*[SU3] ear phenotype). When we examined the spinal cords of these mutants, we observed a statistically significant reduction in the number of glutamatergic cells, but in both cases the reduction was smaller than we had previously observed for morpholino-injected DKD embryos (Fig. 5D & I, Table 3). There was also an increase in the number of inhibitory spinal cord interneurons, although, again, the increase was less than in the DKD morpholino-injected embryos (Fig. 6C, H, M, Table 3). However, similar to the DKD embryos, there was no change in the number of spinal *hmx3a*- or *en1b*-expressing cells in *hmx3a*[SU3] mutants, suggesting that dI2 and V1 interneurons are forming in normal numbers and not dying or changing into different classes of interneurons (Fig. 6A, B, F, G, K, L, Table 3).

In contrast, embryos homozygous for $hmx3a^{SU42}$ do not have fused otoliths or stalled lateral line progression (Fig. 4, Fig. 5M, Table 4) and unlike $hmx3a^{SU3}$ mutants, they have normal expression of $hmx3a$ in the anterior otic epithelium and adjacent anterior neuroblasts (Fig. 5S and cf Fig. 5R & U'). $hmx3a^{SU42}$ mutants also have normal numbers of spinal cord glutamatergic neurons (Fig. 5B & G, Table 3) and are homozygous viable (Fig. 4, Table 4). Interestingly, embryos homozygous for $hmx3a^{sa23054}$ have variable, incompletely penetrant otolith fusion phenotypes that range from no fusion, through incomplete fusion (Fig. 5T), to complete fusion, despite the fact that any protein encoded by this allele should retain more WT sequence than that encoded by $hmx3a^{SU42}$ (Fig. 4). However, $hmx3a^{sa23054}$ mutants have normal lateral line progression, no reduction in the number of spinal cord glutamatergic neurons and they are also viable (Fig. 4, Fig. 5C, H & N, Tables 3 & 4).

Surprisingly, we also found that all four of the different $hmx2$ alleles we created in these experiments ($hmx2^{SU35}$, $hmx2^{SU36}$, $hmx2^{SU37}$, $hmx2^{SU38}$; Fig. 4) are homozygous viable and have no obvious defects in otolith development. We also examined lateral line progression and the number of spinal cord glutamatergic neurons in $hmx2^{SU37}$ and $hmx2^{SU38}$ homozygous mutants and did not detect any change compared to WT embryos (Fig. 4, Tables 3 & 4 and data not shown).

To test whether our $hmx2$ mutants had no obvious phenotypes because they had retained some Hmx2 function, we created a large deletion allele, $hmx2^{SU39}$, that deletes most of the $hmx2$ genomic sequence. Only 84 nucleotides of 5' and 60 nucleotides of the most 3' coding sequence remain (Fig. 4). However, $hmx2^{SU39}$ homozygous mutants also lack fused otoliths and are homozygous viable. They also have normal expression of $hmx3a$ and $pax5$ in the anterior otic epithelium, normal expression of $hmx3a$ in the adjacent anterior neuroblasts, the normal complement of three distinct cristae and two distinct maculae in the ear, and normal lateral line progression (Fig. 4, Fig. 5Q, W, AA, AE & AI, Tables 3 & 4). In

addition, there is no change in the number of glutamatergic cells in the spinal cords of these mutants, when compared to WT sibling embryos (Fig. 5F & K, Table 3).

While this was surprising, we hypothesized that the lack of obvious phenotypes could be because Hmx3a was compensating for loss of Hmx2. To test this hypothesis, we created double mutant embryos. As *hmx2* and *hmx3a* are adjacent on chromosome 17, it is not possible to create double mutants by breeding single mutants. Both mutations have to exist on the same chromosome. Therefore, we created two different deletion alleles, *hmx2;hmx3a^{SU44}* and *hmx2;hmx3a^{SU45}*, that lack the entire *hmx3a* coding sequence and all but the last 66 nucleotides of *hmx2* coding sequence (Fig. 4 and materials and methods; the alleles differ only in the amount of remaining sequence upstream of the *hmx3a* locus). However, in embryos homozygous for these large deletion alleles, the reduction in the number of excitatory spinal cord interneurons and the increase in the number of inhibitory spinal neurons is equivalent to that in *hmx3a^{SU3}* and *hmx3a^{SU43}* single mutants (Fig. 5E & J, Fig. 6D, I & N and Table 3). There are also no obvious differences in the otolith fusion or lateral line progression phenotypes between these single mutants and embryos homozygous for the double deletion alleles (Fig. 5P & V, and compare to Fig. 5 O and U). To further interrogate whether there might be subtle differences in ear phenotypes between *hmx3a^{SU3}* and double deletion mutants, we examined the expression of *pax5* in the anterior otic epithelium and the presence and integrity of the cristae and maculae of the ear using phalloidin staining. Both the reduction in *pax5* expression in the anterior otic epithelium (cf Fig.5Z & Y), the fusion/juxtaposition of the maculae within a more ventro-medial position in the ear (cf Fig.5AH & AG) and the size and number of cristae in the ear (Fig.5AB-AD) are equivalent in *hmx2;hmx3a^{SU44}* double deletion and *hmx3a^{SU3}* single mutants.

To determine whether the increase in the number of spinal inhibitory interneurons reflects an increase in glycinergic or GABAergic neurons, we examined the expression of genes expressed exclusively by cells

with these inhibitory neurotransmitter phenotypes (*slc6a5* for glycinergic and *gad1b* for GABAergic, see materials and methods). While we found no statistically significant difference in the number of spinal cord glycinergic cells in *hmx3a^{SU3}* mutants or *hmx2;hmx3a^{SU44}* deletion mutants, there was a statistically significant increase in the number of GABAergic cells in *hmx3a^{SU3}* and *hmx2;hmx3a^{SU44}* mutants compared to WT sibling embryos (Fig. 6E, J, O, P, Q, V, W, AB, AC, Table 3).

As *hmx2* spinal expression is initially weaker than *hmx3a* expression (Fig.1) we also analyzed *hmx2^{SU39}* single mutants at 48 h, to determine if there was a spinal cord neurotransmitter phenotype at this later stage of development. We examined the number of GABAergic cells, as these are easier than glutamatergic cells to count at this stage. However, there was no change in the number of GABAergic spinal cells in *hmx2^{SU39}* single mutants compared to WT embryos (Fig. 6 T, Z, AF, Table 3). We also tested whether the spinal phenotype of embryos homozygous for the double deletion alleles was more severe than *hmx3a^{SU3}* single mutants at 48 h. However, while the number of GABAergic cells was increased in both *hmx3a^{SU3}* single mutants and *hmx2;hmx3a^{SU44}* double deletion mutants, there was no statistically significant difference between these two phenotypes (Fig. 6R, S, X, Y, AD, AE, Table 3).

Trans-heterozygous crosses suggest that *hmx3a^{SU42}* is a hypomorphic allele.

hmx3a^{SU3}, *hmx3a^{SU42}* and *hmx3a^{SU43}* mutant alleles all introduce a frameshift within 4 nucleotides of each other (nucleotides 319, 320, and 323 of the coding sequence respectively). Assuming that all three of these alleles are translated into truncated proteins, *hmx3a^{SU3}* would retain 106 WT amino acids whereas the other two alleles would retain 107 WT amino acids (Fig. 4). However, despite the similarity of these mutant alleles, embryos homozygous for *hmx3a^{SU3}* and *hmx3a^{SU43}* have fused otoliths, stalled lateral line progression, altered spinal interneuron neurotransmitter phenotypes and do not survive to adulthood whereas embryos homozygous for *hmx3a^{SU42}* are viable and lack all of these phenotypes. In addition, *hmx3a^{sa23054}* mutants have variable otolith fusion phenotypes, despite retaining more WT

sequence than $hmx3a^{SU3}$, $hmx3a^{SU42}$ or $hmx3a^{SU43}$ alleles, and embryos homozygous for $hmx2^{SU39}$, which almost completely deletes the $hmx2$ coding sequence, have no obviously abnormal phenotypes. Given the surprising nature of these results, we decided to further investigate these alleles by creating trans-heterozygous animals. To do this we performed different pair-wise crosses between fish heterozygous for $hmx2^{SU39}$, $hmx3a^{SU3}$, $hmx3a^{SU42}$, $hmx3a^{SU43}$, $hmx3a^{sa23054}$ and $hmx2;hmx3a^{SU44}$ and analyzed ear and lateral line development in the resulting embryos.

Interestingly, when we crossed fish heterozygous for $hmx2;hmx3a^{SU44}$ with fish heterozygous for $hmx2^{SU39}$ the resulting embryos had normal otolith development and lateral line progression (Table 5). Given that approximately a quarter of these embryos should lack almost all of the coding sequence for both alleles of $hmx2$ ($hmx2;hmx3a^{SU44}$ lacks all but the last 66 bp and $hmx2^{SU39}$ lacks all but the first 84 and last 60 bp of $hmx2a$ coding sequence, see Fig. 4) as well as all of the coding sequence for one allele of $hmx3a$, this demonstrates that one WT allele of $hmx3a$ is sufficient for normal otolith development and lateral line progression.

As expected, as both alleles produce homozygous mutant phenotypes, when we crossed fish heterozygous for $hmx3a^{SU3}$ with fish heterozygous for $hmx3a^{SU43}$ we observed mendelian ratios of embryos with fused otoliths and stalled lateral line progression (Table 5). Similarly, we obtained mendelian ratios of embryos with fused otoliths and stalled lateral line progression when we crossed fish heterozygous for $hmx3a^{SU3}$ with fish heterozygous for $hmx2;hmx3a^{SU44}$, or fish heterozygous for $hmx3a^{SU43}$ with fish heterozygous for $hmx2;hmx3a^{SU44}$.

More interestingly, when we crossed fish heterozygous for $hmx3a^{sa23054}$ with fish heterozygous for either $hmx2;hmx3a^{SU44}$ or $hmx3a^{SU3}$, we also obtained mendelian ratios of embryos with fully penetrant otolith fusion phenotypes and stalled lateral line progression, as opposed to the variable otolith fusion

phenotypes and normal lateral line progression that occurs in $hmx3a^{sa23054}$ homozygous mutants (cf Table 4 & Table 5). Some of the embryos with stalled lateral line progression had the strong phenotype that we observe in $hmx3a^{SU3}$ and $hmx2;hmx3a^{SU44}$ homozygous mutants, and the rest had a slightly weaker phenotype (Table 5, similar to Fig. 6AA). This suggests that while two $hmx3a^{sa23054}$ alleles provide sufficient Hmx3a activity for normal lateral line progression and, in some cases, normal otolith development, this is not the case for either the combination of one $hmx3a^{sa23054}$ and one $hmx3a^{SU3}$ allele or the combination of one $hmx3a^{sa23054}$ allele over a $hmx3a$ deletion.

Even more surprisingly, when we crossed fish heterozygous for $hmx3a^{SU42}$ with fish heterozygous for either $hmx2;hmx3a^{SU44}$ or $hmx3a^{SU3}$ we also obtained embryos with fused otoliths and stalled lateral line progression, although most of the embryos had the slightly weaker lateral line phenotype mentioned above (Fig. 6 AA & AG, Table 5). For the combination of $hmx3a^{SU42}$ and $hmx2;hmx3a^{SU44}$, these phenotypes occurred in mendelian ratios. However, for the combination of $hmx3a^{SU3}$ and $hmx3a^{SU42}$, while we observed a mendelian ratio of embryos with stalled lateral line progression, only 17% of embryos had abnormal otolith phenotypes and in most of these cases the otoliths were either adjacent but not fused in both ears or there was an abnormal otolith phenotype in only one ear (Table 5). When we genotyped a subset of these embryos, we found that all of the embryos with abnormal otolith phenotypes (n=18, 6 embryos each with either fused otoliths in both ears, adjacent otoliths in both ears, or a fused or adjacent otolith phenotype in only one ear) were $hmx3a^{SU3/+};hmx3a^{SU42/+}$ trans-hets. Interestingly, when we genotyped embryos with WT otolith phenotypes (two normal otoliths per ear, n=173), 5.20% were actually $hmx3a^{SU3/+};hmx3a^{SU42/+}$ trans-hets, 29.48% were only heterozygous for $hmx3a^{SU3}$, 31.79% were only heterozygous for $hmx3a^{SU42}$ and 33.53% were homozygous wild type. Taken together, these data suggest that even though $hmx3a^{SU42}$ homozygous mutants have no obvious abnormal phenotypes, $hmx3a^{SU42}$ is a hypomorphic allele: while two alleles of $hmx3a^{SU42}$ provide sufficient Hmx3a activity for normal otolith development and lateral line progression, one $hmx3a^{SU42}$

allele combined with either one $hmx2;hmx3a^{SU44}$ or one $hmx3a^{SU3}$ allele does not. In contrast, one WT allele of $hmx3a$ is sufficient for normal development.

Loss of Hmx2 function can enhance hypomorphic $hmx3a$ phenotypes.

Our comparisons of $hmx3a^{SU3}$, $hmx3a^{SU43}$, $hmx2;hmx3a^{SU44}$ and $hmx2;hmx3a^{SU45}$ mutant phenotypes (Figs 5 & 6 and Table 3) suggested that $hmx2$ does not act redundantly with $hmx3a$ in zebrafish otolith development, lateral line progression or specification of correct spinal interneuron neurotransmitter phenotypes as complete removal of both genes does not result in more severe phenotypes than loss of just Hmx3a function. However, if loss of Hmx3a function already produces maximal mutant phenotypes, we wouldn't detect stronger phenotypes in embryos homozygous for the double deletion alleles. Therefore, a more sensitive way to test if Hmx2 functions in these developmental processes, would be to remove Hmx2 function in embryos homozygous for a "weaker" hypomorphic $hmx3a$ mutant allele such as $hmx3a^{SU42}$. Unfortunately, we cannot mate fish with the $hmx2^{SU39}$ and $hmx3a^{SU42}$ mutant alleles to make double mutants, as $hmx2$ and $hmx3a$ are adjacent on the same chromosome, so each mutant allele is tightly linked to a WT allele for the other gene. Therefore, we decided to knock-down Hmx2 function in $hmx3a^{SU43}$ mutants using CRISPR-mediated mutagenesis.

We injected CRISPR reagents to mutate $hmx2$ into embryos from an incross of fish that were heterozygous for $hmx3a^{SU42}$. We used the Microhomology-mediated End joining kNockout Target Heuristic Utility (MENTHU) tool to identify a sgRNA target site that should predominantly result in the same 5 bp deletion frame-shift allele (Fig. 4, Fig. 7C), being generated through microhomology-mediated end joining (MMEJ; Ata et $al.$ 2018; Mann et $al.$ 2019). We also used a two-part crRNA + tracrRNA system + Cas9 protein ribonucleoprotein complex for the injections, as this can produce a high efficiency of biallelic mutations and F0 phenotypes (DiNapoli et $al.$ 2019; Hoshijima et $al.$ 2019; Lewis Lab unpublished data). When we did this, we found that at ~3.5 dpf, 28.25% (n=807) of $hmx2$ CRISPR-

injected embryos had an abnormal otolith phenotype (21.31% had fused otoliths in both ears and 6.94% had fused or adjacent otoliths in only one ear; Fig.7B, cf to uninjected control, Fig.7A). In comparison, only 0.6% (n=670) of uninjected embryos and 2% (n=347) of embryos injected with a CRISPR crRNA ribonucleoprotein complex that we have used successfully to make mutations in an unrelated gene, had abnormal otolith phenotypes. These control experiments were performed at the same time as the *hmx2* CRISPR injections, using embryos obtained from the same heterozygous *hmx3a^{SU42}* parent fish.

We examined 40 of the *hmx2* CRISPR-injected embryos at ~30 h for lateral line progression phenotypes and then let these embryos develop to ~3.5 dpf, so we could correlate lateral line and otolith phenotypes. 25% of the embryos had strong or medium stalled lateral line progression phenotypes and all of these embryos also developed fused otoliths in both ears (Table 6A). A few additional embryos had a weaker lateral line progression defect (migration of the primordium was only delayed by 2-3 somites compared to stage-matched injected siblings) and two of these also developed fused otoliths in both ears. When we genotyped these embryos for *hmx3a^{SU42}*, we found that all of the embryos with fused otoliths were homozygous for *hmx3a^{SU42}* (Table 6B).

We also genotyped 72 additional *hmx2* CRISPR-injected embryos, just over half of which had otolith defects. The vast majority of the embryos with otolith phenotypes were homozygous for *hmx3a^{SU42}* (Fig,7B, Table 6C, one embryo with fused otoliths in both ears and one embryo with an otolith defect in one ear only were heterozygous). In contrast, all except one of the 35 embryos that did not have obvious defects in otolith development were heterozygous for *hmx3a^{SU42}* or WT (Table 6C).

Taken together, these results suggest that we obtained a high efficiency of *hmx2* mutations in our injected embryos and that CRISPR-mediated knock-down of Hmx2 causes *hmx3a^{SU42}* mutants to have defects in otolith development and lateral line progression. To test this, we sequenced the *hmx2* allele from 23 of

the embryos that we had genotyped for $hmx3a^{SU42}$ that had different otolith phenotypes (Table 6D). We found that all of these embryos had a substantial frequency of $hmx2$ nonsense alleles. As predicted by the MENTHU algorithm, the mutated alleles all contained a 5 bp deletion, although some also had additional mismatches in the three bases immediately prior to the deletion and the location of the deletion differed by 1 bp in a few cases. In all cases, we estimate that at least 60% of the amplified $hmx2$ sequences were mutant (Table 6D; Fig.7C, D). In one of the WT embryos that lacked a phenotype, approximately 90% of the amplified $hmx2$ sequences were mutant, suggesting that, consistent with the lack of abnormal phenotypes in $hmx2^{SU39}$ mutants, CRISPR mutagenesis of $hmx2$ is not sufficient for abnormal otolith development (Fig.7C, D).

$hmx2$ and $hmx3a$ are not expressed maternally.

One possible explanation for why the spinal cord phenotype is less severe in $hmx2/3a$ deletion mutants than in morpholino-injected DKD embryos would be if $hmx2$ and/or $hmx3a$ are maternally expressed, as in this case the morpholinos might knock-down both maternal and zygotic function whereas the mutants would only remove zygotic function. In addition, maternal expression of $hmx2$ might explain the lack of any obvious abnormal phenotypes in $hmx2$ single mutants. To test this, we performed *in situ* hybridization for $hmx2$ and $hmx3a$ at the 16-cell stage. However, we did not detect any maternal expression of $hmx2$, $hmx3a$ or any of the other hmx genes (Fig. 8I-M). We also performed qRT-PCR for $hmx2$ and $hmx3a$ on whole embryos at different developmental stages. We did not observe expression of either gene at either the 16-cell stage or at 6 h, suggesting that neither $hmx2$ nor $hmx3a$ are maternally expressed (Fig. 8H). At 14 h, shortly after when both of these genes start to be expressed in the ear and spinal cord, we observed low levels of expression and, for both genes, as expected, this became more abundant at 27 h and 48 h. (Fig. 8H). Finally, we also generated embryos from adults that were homozygous mutant for $hmx2^{SU38}$, $hmx2^{SU39}$ and $hmx3a^{sa23054}$. However, even though half of the embryos

in each of these crosses should have been maternal zygotic mutants, we did not observe any embryos with fused otoliths (Fig. 4, Table 4).

hmx3b, hmx1 and *hmx4* expression is not upregulated in *hmx2;hmx3a*[SU44] deletion mutants.

Even though *hmx3b*, *hmx1* and *hmx4* are not normally expressed in the spinal cord (Fig.1), it was theoretically possible that they are upregulated in response to the absence, or reduced levels, of either Hmx2 and/or Hmx3a function, in which case they could partially substitute for the loss of Hmx2 and/or Hmx3a. To test this, we performed *in situ* hybridization for these genes in *hmx3a*[SU3] and *hmx2;hmx3a*[SU44] mutants at 27 h. In both cases, we did not observe any spinal cord expression of these genes in either genotyped mutants or their sibling embryos, although, as observed previously in WT embryos (Fig. 1) *hmx1* and *hmx4* were expressed in the eye, ear and anterior lateral line neuromasts in both mutants and WT sibling embryos (Fig. 8E-G and data not shown).

hmx2[SU39] mutants do not lack a spinal cord phenotype because of genetic compensation.

Recent reports have demonstrated that genetic compensation (upregulation of other genes that can compensate for loss of the mutated gene) can result in loss-of-function mutants having a less severe phenotype than embryos injected with a morpholino against the same gene (Rossi *et al.* 2015; El-Brolosy and Stainier 2017; Zhu *et al.* 2017; Sztal *et al.* 2018; El-Brolosy *et al.* 2019; Peng 2019). If this is the case, then morpholino knock-down should have less effect on mutant embryos than on WT sibling embryos, because the morpholino will not affect upregulated compensating genes (e.g. see Rossi *et al.* 2015; Sztal *et al.* 2018). Therefore, to test whether the lack of a spinal cord phenotype in *hmx2*[SU39] deletion mutants is due to genetic compensation, we injected the *hmx2* morpholino into embryos from a cross of fish heterozygous for *hmx2*[SU39] and performed *in situ* hybridization for *slc17a6a/b* to label glutamatergic spinal cord interneurons. We predicted that if there was genetic compensation in *hmx2*[SU39] mutants, morpholino-injected WT sibling embryos should have a reduced number of spinal cord

glutamatergic interneurons, whereas *hmx2^SU39^* homozygous mutant embryos should have normal numbers of these cells. However, if there is no genetic compensation, we would expect a similar frequency of homozygous mutant and WT embryos to have spinal cord phenotypes and those phenotypes to be roughly equivalent in severity. In contrast, if morpholino-injected homozygous mutant embryos have more severe phenotypes than WT siblings, this might suggest that *hmx2^SU39^* is a hypomorphic allele. However, this seems highly unlikely given that the *hmx2* gene is almost completely deleted in this allele (Fig. 4).

We initially examined the injected embryos down a stereomicroscope and divided them into two groups: those that had an obvious reduction in glutamatergic cells and others that either lacked a phenotype or had a more subtle phenotype. When we genotyped these embryos, we found both homozygous mutant and WT embryos at frequencies that were not statistically significantly different from Mendelian ratios in each group (Table 7). In addition, when we compared the average number of glutamatergic cells in morpholino-injected WT and mutant embryos, there was no statistically significant difference between them, <u>regardless</u> of whether we compared all of the morpholino-injected embryos of each genotype or just compared embryos within the same phenotypic group (Table 7). This suggests that the lack of an abnormal spinal cord phenotype in *hmx2^SU39^* homozygous mutant embryos is not due to genetic compensation and that the differences that we observed between the two groups of injected embryos instead probably reflect exposure to different levels of morpholino (see materials and methods).

<u>*hmx3a^SU42^* mutant alleles do not lack a spinal cord phenotype because of genetic compensation.</u>

We also tested whether the spinal cord phenotype of *hmx3a^SU42^* mutants is less severe than embryos injected with a *hmx3a* morpholino because of genetic compensation. As above, we injected the *hmx3a* morpholino into embryos from a cross of fish heterozygous for *hmx3a^SU42^* and performed *in situ* hybridization for *slc17a6a/b*. When we examined the injected embryos down a stereomicroscope, we

were able to separate them into one group that had an obvious reduction in spinal glutamatergic cells and another group that either lacked, or had a more subtle, phenotype. However, when we genotyped the embryos in these two groups, we found similar numbers of homozygous mutant and WT embryos in both groups and the frequencies of different genotypes in each group were not statistically significantly different from Mendelian ratios (Table 8). In addition, when we compared the average number of glutamatergic cells in morpholino-injected WT and mutant embryos, there was no statistically significant difference between them, <u>regardless</u> of whether we compared all of the morpholino-injected embryos of each genotype or just compared embryos within the same phenotypic group (Table 8). These data suggest that the lack of an abnormal spinal cord phenotype in *hmx3a*^{SU42} homozygous mutant embryos is not due to genetic compensation and that the differences that we observed between the two groups of injected embryos just reflect exposure to different levels of morpholino (see materials and methods). Consistent with this, we also do not observe any nonsense-mediated decay (NMD) of *hmx3a* mRNA in *hmx3a*^{SU42} mutants (Fig. 8A; NMD has been suggested to play a key role in at least some instances of genetic compensation; El-Brolosy *et al.* 2019).

hmx3a^{SU3} mutants also do not have genetic compensation.

We also tested whether *hmx3a*^{SU3} mutants have genetic compensation. As above, we injected the *hmx3a* morpholino into embryos from a cross of fish heterozygous for *hmx3a*^{SU3} and performed *in situ* hybridization for *slc17a6a/b*. When we examined these embryos down a stereomicroscope, some of the embryos appeared to have a severe reduction in glutamatergic cells that resembled the morpholino knock-down phenotype, whereas in the other embryos any reduction was more subtle. However, when we genotyped these embryos, we again found similar numbers of homozygous mutant and WT embryos in both groups and the frequencies of the different genotypes in each group were not statistically significantly different from Mendelian ratios (Table 9). In addition, when we compared the average number of glutamatergic cells in morpholino-injected WT and mutant embryos, there was no statistically

significant difference between them, <u>regardless</u> of whether we compared all of the morpholino-injected embryos, or just compared the injected embryos within a particular phenotypic group (Table 9). For the "weaker" phenotypic group, the difference between WT and mutant embryos approached statistical significance. However, this is probably because some of the WT embryos in this category had almost no reduction in the number of glutamatergic cells, whereas all of the mutants in this category had at least their normal mutant phenotypes. These results suggest that the differences that we observed between the two groups of injected-embryos probably just reflect exposure to different levels of morpholino (see materials and methods), and that embryos in the "weaker" phenotypic group did not receive sufficient morpholino to effectively knock-down *hmx3a* mRNA or cause the more severe morpholino phenotype. Taken together, these data suggest that the spinal cord phenotype in *hmx3a*^{SU3} homozygous mutant embryos is not less severe than the morpholino knock-down phenotype because of genetic compensation. Consistent with this, we also do not detect any NMD of *hmx3a* mRNA in *hmx3a*^{SU3} mutants (Fig. 8C).

Discussion

Hmx3a is required for correct neurotransmitter phenotypes of a subset of spinal cord interneurons.

In this paper, we identify for the first time, a requirement for Hmx3a in spinal cord interneuron development. We demonstrate that *hmx2* and *hmx3a* are expressed by V1 and dI2 interneurons, which is consistent with very recent scRNA-Seq data from mouse spinal cord (Delile *et al.* 2019). Of these cell types, only dI2 interneurons are glutamatergic. Therefore, the most likely explanation for the reduction in the number of glutamatergic spinal cord cells in *hmx3a* mutants is that dI2 interneurons are losing their glutamatergic phenotypes. Given that we also detect a corresponding increase in GABAergic spinal cord cells, but the number of V1 cells (indicated by *en1b* expression) does not change, it is likely that the dI2 interneurons that are losing their glutamatergic phenotypes are becoming GABAergic instead. Unfortunately, the respective *in situ* hybridization probes are not strong enough to formally confirm with double-labeling experiments that dI2 interneurons switch their neurotransmitter phenotype from glutamatergic to GABAergic. However, unless Hmx3a is acting in a cell non-autonomous manner, which we think is unlikely as this protein has a nuclear localization sequence and no obvious signal peptide, this is the most likely explanation of our data.

Hmx3a is required for progression of the posterior lateral line primordium.

Feng and Xu previously reported that the number of posterior lateral line primordium neuromasts was either severely reduced or completely lost at 3 dpf in *hmx2/3a* DKD animals (Feng & Xu, 2010). Intriguingly, the few neuromasts that sometimes persisted were located very rostrally in the embryo, close to the earliest forming somites. Our analyses demonstrate that at 27 h, when the posterior lateral line primordium has migrated to somite 10 in WT embryos, in *hmx3a*^{SU3}, *hmx3a*^{SU43}, *hmx2;hmx3a*^{SU44} and *hmx2;hmx3a*^{SU45} mutants the primordium is stalled adjacent to somites 1-4. This suggests that the previously reported loss of neuromasts at 3 dpf is probably caused by the posterior

lateral line primordium failing to migrate and deposit neuromasts. Feng & Xu also described reduced cell proliferation (at 15 h) and reduced *hmx3a* expression (at 24 h) in the posterior lateral line primordium of *hmx2/3a* DKD animals. Whilst we cannot rule out the possibility that the lateral line primordium fails to migrate because it has not formed correctly, we observe persistent expression of both *hmx3a* and *krt15* in the stalled primordium of our *hmx3a^SU3^, hmx3a^SU43^, hmx2;hmx3a^SU44^* and *hmx2;hmx3a^SU45^* mutants (data not shown), suggesting that some other mechanism, possibly chemosensory, might underlie the stalled migration.

Hmx3a does not require its homeodomain for its functions in viability and otolith, lateral line and spinal cord interneuron development.

Our results also suggest that Hmx3a does not require its homeodomain for either its role in viability or its essential functions in otolith development, lateral line progression or correct specification of a subset of spinal cord neurotransmitter phenotypes. This is surprising because most homeodomain proteins act as transcription factors and use their homeodomain to bind DNA and regulate gene expression. Instead, our data suggest that there must be at least one other, as yet undiscovered, crucial functional domain in the N-terminal region of Hmx3a, that is required for its functions in embryo development and viability, as embryos homozygous for *hmx3a^SU42^* are viable and have no obvious abnormal phenotypes and embryos homozygous for *hmx3a^sa23054^* are also viable, have normal lateral line progression and spinal cord interneuron neurotransmitter phenotypes and produce viable progeny. It is highly unlikely that the lack of obvious abnormal phenotypes in these two different mutants is due to an alternative translation start site creating a truncated Hmx3a protein that contains the homeodomain, as the only downstream methionine in *hmx3a* is more than a third of the way through the homeodomain. While we cannot rule out the possibility that Hmx3a may bind to other DNA-binding proteins and function in transcriptional complexes, unless the N-terminal of Hmx3a contains a novel DNA-binding domain, our results also suggest that Hmx3a is not acting as a classic transcription factor (defined in the strict sense as a protein

that binds DNA and regulates transcription) during its functions in viability and otolith, lateral line and spinal cord interneuron development. Nevertheless, as the homeodomain in highly conserved, suggesting that it is still under evolutionary pressure to be maintained, it is possible that Hmx3a has additional functions that do require this domain, either in adult fish or in aspects of development that we did not assay. However, if this is the case, it is still striking that these functions are not required for such fundamental processes as viability or reproduction.

There are a few other examples of homeodomain proteins that can function in some contexts without their homeodomain. For example, protein interaction and over-expression experiments suggest that Lbx2 does not require its homeodomain to enhance Wnt signaling during gastrulation in zebrafish embryos (Lu *et al.* 2014). Instead it sequesters TLE/Groucho, preventing this protein from binding to TCF7L1 and reducing TLE/TCF co-repressor activity. In addition, *homothorax* (*hth*) does not require its homeobox for its functions during Drosophila head development and proximo-distal patterning of the appendages, although the homeodomain is required for antennal development (Noro *et al.* 2006). *hth* has 16 exons and three alternative splice forms. Only one of these isoforms contains the homeobox, but all three contain a protein interaction domain, called the HM domain, that binds to, and can induce the nuclear localization of, Extradenticle (Noro *et al.* 2006). However, in contrast to *hth*, zebrafish *hmx3a* has only two exons and one splice form. While relatively rare, there also examples of transcription factors from other families that only need to bind DNA for some of their functions. For example, Scl/Tal1 has both DNA-binding dependent and DNA-binding independent functions in hematopoietic and vascular development (Porcher *et al.* 1999; Ravet *et al.* 2004).

Zebrafish Hmx2 is not, by itself, required for viability or correct development of otoliths, lateral line or spinal cord neurotransmitter phenotypes.

The experiments described in this paper also show that *hmx2* single mutants, including *hmx2^{SU39}* mutants that lack almost all *hmx2* coding sequence, are viable and have no obvious otolith, lateral line or spinal cord interneuron neurotransmitter mutant phenotypes, and that *hmx2;hmx3a^{SU44}* and *hmx2;hmx3a^{SU45}* deletion mutants do not have more severe phenotypes than *hmx3a^{SU3}* or *hmx3a^{SU43}* single mutants. These results are surprising because zebrafish *hmx2* and *hmx3a* have very similar expression domains during embryonic development (although the spinal cord expression of *hmx3a* does briefly precede that of *hmx2*, Fig. 1B & C), these overlapping expression domains are highly conserved in different vertebrates and studies in other animals suggest that Hmx2 and Hmx3 often act redundantly during development (Wang *et al.* 2004; Wang and Lufkin 2005; Wotton *et al.* 2010). Most notably, previous analyses demonstrated that mouse *Hmx2* mutants had defects in ear development, although interestingly these were more severe in about 70% of homozygous mutants than in the other 30%, showing that there was some variability in the requirement for Hmx2 (Wang *et al.* 2001). In addition, mouse *Hmx2;Hmx3* double mutants had more severe ear phenotypes than either single mutant, as well as defects in hypothalamus and pituitary development that were not found in either single mutant and most of the double mutants died around the 5th day after birth, whereas the single mutants were viable (Wang *et al.* 2004). When considered in combination, these data suggest that mouse Hmx2 has important functions in ear and brain development, although some of these are redundant with Hmx3. In contrast, the only experiment where we detected any function for zebrafish Hmx2, was when we introduced *hmx2* mutations into hypomorphic *hmx3a^{SU42}* mutants. Taken together, our data suggest that while zebrafish Hmx2 can function in otolith and lateral line development, it may only impact the development of these structures in embryos with significantly reduced Hmx3a function (less activity than one functional *hmx3a* allele, as embryos trans-heterozygous for *hmx2;hmx3a^{SU44}* and *hmx2^{SU39}* develop normally) but more Hmx3a activity than in *hmx3a^{SU3}* or *hmx3a^{SU43}* mutants.

Our experiments also suggest that the lack of any obvious abnormal phenotypes in $hmx2^{SU39}$ single mutants is not due to genetic compensation or maternal expression of *hmx* genes. One possible explanation for why zebrafish Hmx2 might have a diminished role in development compared to mouse Hmx2 or zebrafish Hmx3a could be if zebrafish Hmx2 has evolved to be less conserved with mouse Hmx2 and Hmx3 than zebrafish Hmx3a. However, a comparison of all four proteins only reveals six residues that are shared between mouse Hmx2 and Hmx3 and zebrafish Hmx3a but not zebrafish Hmx2, and four of these residues are upstream of the $hmx3a^{SU3}$ mutation (Supp. Data Fig. 1). Our prior research identified Hmx2 and Hmx3 in all of the different vertebrates that we analyzed, including five different teleost species, and our phylogenetic analyses of these proteins did not suggest that Hmx2 has evolved any faster in zebrafish than in other species, or that Hmx2 has evolved faster than Hmx3 (Wotton *et al.* 2010). Taken together, these observations suggest that there is still considerable evolutionary pressure to maintain zebrafish Hmx2, which in turn suggests that it should have an important role(s) in zebrafish survival and/or reproduction. Therefore, it is surprising that we did not detect more severe consequences from loss of Hmx2. It is possible that Hmx2 has important functions later in development and/or in aspects of development that we did not assay. However, if this is the case, these functions are not required for viability or reproduction as even $hmx2^{SU39}$ homozygous mutants survive to adulthood and produce viable progeny. It is also possible that *hmx2* has important function(s) in adult fish, as our assays would not have detected this. It would be interesting to investigate these possibilities in future studies.

Very similar *hmx3a* mutant alleles have different homozygous mutant phenotypes.

It is currently unclear why $hmx3a^{SU42}$ retains more WT function than $hmx3a^{SU43}$ when both should encode proteins with only 107 WT amino acids. We are confident that this is not due to alternative splicing or exon skipping as we have sequenced cDNA made from extracted total RNA from homozygous $hmx3a^{SU42}$ individuals (see materials and methods). We also do not observe any NMD of *hmx3a* mRNA for any of our *hmx3a* single mutant alleles (Fig. 8A-D; the double deletion mutants lack

all *hmx3a* coding sequence, so there is no mRNA to assess), so the difference in allelic strength is not due to some of the mutant mRNAs being degraded. However, it is possible that different mutant alleles result in different amounts of truncated protein due to differences in translation efficiency or protein stability. Unfortunately, we were not able to test this as there are currently no antibodies that uniquely recognize the N-terminal region of Hmx3a that is conserved in all four of our *hmx3a* single mutant proteins. In addition, we were unable to detect any Hmx3a peptides in a SWATH analysis (data not shown; Hmx3a has also not been detected in other SWATH analyses Blattmann *et al.* 2019; Lin *et al.* 2019), presumably because, like many transcription factors, it is expressed in either two few cells and/or at too low a level.

It is also possible that the overall length of the mutant protein is important for retaining function and that the additional abnormal amino acids after the frameshift but before the premature stop codon in *hmx3a^{SU42}* help this allele retain more WT function. A longer protein sequence might facilitate a required protein conformation and/or binding with other proteins or molecules. If this is the case, then it could also explain why Hmx3a^{SU42} protein (which is predicted to contain 107 WT + 42 abnormal amino acids, Fig.4) appears to retain more WT function than Hmx3a^{sa23054} protein (which should contain only 118 WT amino acids; Fig.4). Currently, there are no known binding partners of Hmx3a. However, if future analyses identify any it would be interesting to test if they can bind to Hmx3a^{SU42} and Hmx3a^{SU3}.

Another related possibility is that the different stretches of abnormal amino acids after the frameshift in *hmx3a^{SU3}*, *hmx3a^{SU42}* and *hmx3a^{SU43}* might introduce sequences that influence protein stability, degradation or function. Using a variety of online analysis tools, we did not detect any sumoylation, or PEST sequences in any of the predicted mutant protein sequences and the only ubiquitination motifs that we identified are located in the WT sequence present in all four mutant proteins (Rice *et al.* 2000; Sarachu and Colet 2005; Brameier *et al.* 2007; Radivojac *et al.* 2010; Zhao *et al.* 2014). However, the

eukaryotic linear motif prediction tool identified a monopartite variant of a classic, basically-charged nuclear localization signal in Hmx3a^{SU42} that is not present in any of the other predicted mutant protein sequences (Via *et al.* 2009; Gould *et al.* 2010; Kumar *et al.* 2020), although this domain was not detected using default parameters with cNLS Mapper or NucPred (Brameier *et al.* 2007; Kosugi *et al.* 2009). This is potentially very interesting as WT Hmx3a has a nuclear localization signal located between amino acids 167-177, overlapping the start of the homeodomain at amino acid 171, which is downstream of the mutations in all of these alleles.

In addition, online tools that identify disordered versus ordered protein structure suggest that both the WT amino acids in Hmx3a^{sa23054} that are not found in the other predicted Hmx3a mutant proteins and the non-WT amino acids in the predicted protein product of *hmx3a^{SU42}* may provide longer stretches of disordered sequence than are present at the end of the predicted protein products of *hmx3a^{SU3}* or *hmx3a^{SU43}* (Linding *et al.* 2003; Dosztanyi *et al.* 2005; Ishida and Kinoshita 2007; Dosztanyi 2018; Meszaros *et al.* 2018; Erdos and Dosztanyi 2020). These findings raise the intriguing possibility that *hmx3a^{SU42}* might retain Hmx3a function because it can still localize to the nucleus and/or that *hmx3a^{sa23054}* and *hmx3a^{SU42}* might retain some WT activity because of the disordered sequences at the end of their predicted proteins. As disordered protein regions can switch between disordered and ordered states in the presence of a binding partner, it is tempting to speculate that these disordered stretches at the C-termini of Hmx3a^{SU42} and Hmx3a^{sa23054} proteins may still be able to bind proteins essential for Hmx3a function that the other alleles cannot (Meszaros *et al.* 2018). Investigation of these possibilities is outside the scope of the current study but would be interesting to address in future work.

hmx2 and *hmx3a* morpholino injections produce more severe spinal interneuron phenotypes than *hmx2* and *hmx3a* mutants.

Our original morpholino data suggested that all dI2 interneurons might be switching their neurotransmitter phenotypes as the increase in the number of spinal inhibitory cells and the reduction in the number of excitatory spinal cells in DKD embryos were both roughly equal to the number of dI2 interneurons (glutamatergic *hmx3a*-expressing cells). However, even in our *hmx2;hmx3a* deletion mutants, the number of cells changing their neurotransmitter phenotypes is lower than this. The differences between morpholino-injected embryos and mutant embryos can be seen most clearly in our experiment to test whether genetic compensation occurs in *hmx3a^{SU3}* homozygous mutants. These data directly compare uninjected mutants with morpholino-injected mutants and WT siblings from the same experiment. While there were a range of morpholino-injected phenotypes, overall the *hmx3a* morpholino-injected WT and *hmx3a^{SU3}* mutant embryos had a more severe reduction of glutamatergic cells than uninjected mutants (Table 9). While it is possible that *hmx3a^{SU3}* mutants may be slightly hypomorphic, their spinal cord phenotype is the same as *hmx2;hmx3a^{SU44}* mutants, in which the *hmx3a* coding sequence is completely deleted. Therefore, the more severe phenotypes in some of the *hmx3a^{SU3}* mutants injected with *hmx3a* morpholino cannot be explained by the morpholino removing any residual Hmx3a function. This experiment also suggests that the less severe phenotype in uninjected *hmx3a^{SU3}* mutants is not caused by genetic compensation. Consistent with this, we have also shown that this less severe mutant phenotype is not due to other *hmx* genes being upregulated in these mutants or maternal expression of *hmx3a* or any other *hmx* genes. Taken together, these data suggest that the *hmx3a* morpholino may have non-specific effects on neurotransmitter phenotypes in the spinal cord, although see the discussion below. Similarly, as *hmx2* morpholino-injected single knock-down embryos have reduced numbers of spinal cord glutamatergic cells, while *hmx2^{SU39}* mutants do not, and our experiments suggest that this is also not due to genetic compensation, *hmx2* morpholinos may also have non-specific effects on neurotransmitter phenotypes in the spinal cord. Consistent with the idea that these morpholinos have non-specific effects, there is a body curvature phenotype in the morpholino-injected embryos that is not present in the mutants. However, our data also suggest that the *hmx3a* morpholino has specific

effects, as we were able to partially rescue the DKD embryo phenotype by co-injecting a morpholino-resistant *hmx3a* mRNA. Interestingly, we rescued about 19 glutamatergic cells in these experiments, which is roughly equivalent to the number of cells that lose their glutamatergic phenotypes and/or gain inhibitory phenotypes in *hmx3a*^{SU3} and *hmx2;hmx3a*^{SU44} mutants at the same stage, suggesting that we may have rescued all of the specific effects of the *hmx3a* morpholino.

One puzzling thing, however, about this interpretation of the data is that it is not clear why non-specific effects from either of these morpholinos should exacerbate the specific effect, namely a switch in neurotransmitter phenotype, that we see in *hmx3a* mutants. For example, it is very surprising, given our mutant analyses, that injections of the *hmx2* morpholino cause a reduction in the number of spinal cord glutamatergic interneurons. This is not due to cross-reactivity with *hmx3a*, as the two morpholino sequences are completely different from each other and there is no homology between the *hmx2* morpholino and *hmx3a* upstream or coding sequence. When we BLAST the *hmx2* morpholino sequence against the zebrafish genome the only homology is with *hmx2* (25/25 residues) and with intronic sequence for a gene *si:dkey-73p2.3* on chromosome 3 (18/25 residues), which is predicted to encode a protein with GTP-binding activity. Similarly, when we blast the *hmx3a* morpholino sequence against the zebrafish genome the only homology is with *hmx3a* (25/25 residues). It is also surprising that non-specific effects of the *hmx3a* morpholino would, just by chance, exacerbate the real loss of Hmx3a phenotype, causing additional spinal cord interneurons to lose their glutamatergic phenotypes and instead become inhibitory. It is unlikely that this more severe phenotype is due to cell death or a delay in embryo development as there was no change in the number of *hmx3a*- or *en1b*-expressing spinal cord cells in DKD morpholino-injected embryos and there was an increase in the number of inhibitory spinal cord interneurons equivalent to the reduction in glutamatergic neurons. This suggests that if the more severe morpholino-injection phenotypes are due to non-specific effects of the morpholinos, these non-

specific effects produce an identical phenotype to the specific knock-down effect, namely a switch in neurotransmitter phenotype.

One intriguing alternative possibility that could explain the apparent specificity of the additional phenotype in the morpholino-injected embryos could be that Hmx3a is acting as a fate guarantor, to make the normal neurotransmitter phenotype more robust, as has previously been described for some transcription factors (Topalidou *et al.* 2011; Zheng *et al.* 2015; Zheng and Chalfie 2016). In these cases, mutating the gene usually caused a partially penetrant phenotype in ideal conditions but a more severe phenotype in stressed conditions. In this case, the injection of morpholinos could be such a stressed condition, and this could account for the more severe phenotype. Future analyses could investigate this possibility by testing if other stressed conditions increase the severity of $hmx3a^{SU3}$ single mutant or *hmx2;hmx3a* deletion mutant phenotypes. Even if this is not the case, our results suggest that something other than just non-specific effects from the morpholinos may be occurring. Therefore, we felt that it was crucial to report this morpholino injection data, as an intriguing and hopefully thought-provoking contribution to the continuing discussion about the pros and cons of using morpholinos to investigate gene function.

<u>*hmx3b* has very limited expression during early zebrafish embryogenesis.</u>

The data in this paper also provide the first characterization of zebrafish *hmx3b* expression. Surprisingly, given the expression of all of the other four *hmx* genes during zebrafish embryonic development, the only expression of *hmx3b* that we have been able to detect, is very weak expression in the hindbrain from 36-48 h (Fig. 1 N', S' & X'). When we performed our earlier analyses of vertebrate NK genes, we did not find *hmx3b* in either the Zv7 or Zv8 versions of the zebrafish genome (Wotton *et al.* 2010). It did not appear until Zv9. Interestingly, despite *Hmx2* and *Hmx3* being closely linked in all vertebrates examined so far, only one *hmx2* gene has been found in the zebrafish genome and zebrafish *hmx3b* is

not located within any of the previously described duplicated NK homeobox clusters, including the teleost duplications: it is located on chromosome 12, separate from any other *nk* genes (Wotton *et al.* 2010). However, *hmx3b* is flanked by two genes, *bub3* and *acadsb*, that flank *Hmx2* and *Hmx3* in the human, mouse, chicken and xenopus genomes, suggesting that part of the NK cluster previously identified on chromosome 1 may have translocated to chromosome 12. This is maybe not surprising as zebrafish *hmx2* and *hmx3a* have also translocated from the rest of their NK cluster on chromosome 13 to chromosome 17 and zebrafish *hmx1* and *hmx4* have translocated from the rest of their cluster on chromosome 14 to chromosome 1 (Wotton *et al.* 2010). It is possible that this translocation and the loss of surrounding sequences may explain the different expression pattern of *hmx3b* compared to *hmx3a*. For example, in previous analyses we identified three highly conserved non-coding regions in the vicinity of *hmx2* and *hmx3a* (two upstream of *hmx3a* and one in between *hmx3a* and *hmx2*) that are conserved in mammals, frog and teleosts, (Wotton *et al.* 2010) but none of these regions are present near *hmx3b* (data not shown).

In conclusion, in this paper we provide the first description of zebrafish *hmx3b* expression. Our results also identify the spinal cord cells that express *hmx2* and *hmx3a* (dI2 and V1 interneurons) and uncover novel functions for Hmx3a in correct specification of a subset of spinal cord neurotransmitter phenotypes and in lateral line progression. Our data suggest that while Hmx3a is required for viability, correct otolith development, lateral line progression and specification of a subset of spinal neurotransmitter phenotypes, Hmx2 is not, by itself, required for any of these developmental processes, although it can act partially redundantly with Hmx3a in situations where Hmx3a function is significantly reduced, but not completely eliminated. Finally, our results also suggest that Hmx3a does not require its homeodomain for its roles in viability or embryonic development. Taken together, these findings significantly enhance our understanding of spinal cord, ear and lateral line development and suggest that, intriguingly, more homeobox proteins may not require their homeobox domain for many of their essential functions.

Acknowledgments

We thank ZFIN for providing information on nomenclature and other essential zebrafish resources and Henry Putz, Jessica Bouchard, Paul Campbell, Annika Swanson and several SU undergraduate fish husbandry workers for help with maintaining zebrafish lines. We also thank Professor Jason Fridley and Associate Professor Qiu Wang for help with statistical analyses and William Haws and Arshi Mustafa for comments on previous drafts of the manuscript. This research was primarily funded by NINDS R01:NS077947 with some support from New York State Spinal Cord Injury Fund Contract # C32253GG and NSF IOS 1755354.

Author Contributions

SJE created all the *hmx* mutants described in this study except *hmx3a*^{sa23054}, performed most of the morpholino and mutant experiments, all of the statistical analyses and prepared the figures and some of the tables; GAC performed most of the double-labeling experiments and some of the morpholino analyses; AK, TS and GG performed some of the genotyping and *in situ* hybridization experiments and AK also performed some of the double-labeling experiments; KEL conceptualized and directed the study, acquired the financial support for the project, contributed to data analysis, made some of the tables and wrote the paper. All authors read and commented on drafts of the paper and approved the final version.

References

Adamska, M., S. Leger, M. Brand, T. Hadrys, T. Braun *et al.*, 2000 Inner ear and lateral line expression of a zebrafish Nkx5-1 gene and its downregulation in the ears of FGF8 mutant, ace. Mech Dev 97**:** 161-165.

Alaynick, W. A., T. M. Jessell and S. L. Pfaff, 2011 SnapShot: spinal cord development. Cell 146**:** 178-178 e171.

Ata, H., T. L. Ekstrom, G. Martinez-Galvez, C. M. Mann, A. V. Dvornikov *et al.*, 2018 Robust activation of microhomology-mediated end joining for precision gene editing applications. PLoS Genet 14**:** e1007652.

Balciuniene, J., D. Nagelberg, K. T. Walsh, D. Camerota, D. Georlette *et al.*, 2013 Efficient disruption of Zebrafish genes using a Gal4-containing gene trap. BMC Genomics 14**:** 619.

Batista, M. F., J. Jacobstein and K. E. Lewis, 2008 Zebrafish V2 cells develop into excitatory CiD and Notch signalling dependent inhibitory VeLD interneurons. Dev Biol 322**:** 263-275.

Batista, M. F., and K. E. Lewis, 2008 Pax2/8 act redundantly to specify glycinergic and GABAergic fates of multiple spinal interneurons. Dev Biol 323**:** 88-97.

Benjamini, Y., and Y. Hochberg, 1995 Controlling the false discovery rate: a practical and powerful approach to multiple testing. Journal of the Royal Statistical Society Series B 57**:** 289–300.

Blattmann, P., V. Stutz, G. Lizzo, J. Richard, P. Gut *et al.*, 2019 Generation of a zebrafish SWATH-MS spectral library to quantify 10,000 proteins. Sci Data 6**:** 190011.

Bober, E., C. Baum, T. Braun and H. H. Arnold, 1994 A novel NK-related mouse homeobox gene: expression in central and peripheral nervous structures during embryonic development. Dev Biol 162**:** 288-303.

Boisset, G., and D. F. Schorderet, 2012 Zebrafish hmx1 promotes retinogenesis. Exp Eye Res 105**:** 34-42.

Brameier, M., A. Krings and R. M. MacCallum, 2007 NucPred--predicting nuclear localization of proteins. Bioinformatics 23**:** 1159-1160.

Burglin, T. R., and M. Affolter, 2016 Homeodomain proteins: an update. Chromosoma 125**:** 497-521.

Carrasco, A. E., W. McGinnis, W. J. Gehring and E. M. De Robertis, 1984 Cloning of an X. laevis gene expressed during early embryogenesis coding for a peptide region homologous to Drosophila homeotic genes. Cell 37**:** 409-414.

Cerda, G. A., M. Hargrave and K. E. Lewis, 2009 RNA profiling of FAC-sorted neurons from the developing zebrafish spinal cord. Dev Dyn 238**:** 150-161.

Cheng, L., A. Arata, R. Mizuguchi, Y. Qian, A. Karunaratne *et al.*, 2004 Tlx3 and Tlx1 are post-mitotic selector genes determining glutamatergic over GABAergic cell fates. Nat Neurosci 7**:** 510-517.

Cheng, L., O. A. Samad, Y. Xu, R. Mizuguchi, P. Luo *et al.*, 2005 Lbx1 and Tlx3 are opposing switches in determining GABAergic versus glutamatergic transmitter phenotypes. Nat Neurosci 8**:** 1510-1515.

Concordet, J. P., K. E. Lewis, J. W. Moore, L. V. Goodrich, R. L. Johnson *et al.*, 1996 Spatial regulation of a zebrafish patched homologue reflects the roles of sonic hedgehog and protein kinase A in neural tube and somite patterning. Development 122**:** 2835-2846.

Delile, J., T. Rayon, M. Melchionda, A. Edwards, J. Briscoe *et al.*, 2019 Single cell transcriptomics reveals spatial and temporal dynamics of gene expression in the developing mouse spinal cord. Development 146.

DiNapoli, S., R. Martinez-McFaline, K. Gribbin, C. Balgobin, I. Nelson *et al.*, 2019 A total synthetic approach to CRISPR/Cas9 genome editing and homology directed repair. bioRxiv.

Dosztanyi, Z., 2018 Prediction of protein disorder based on IUPred. Protein Sci 27**:** 331-340.

Dosztanyi, Z., V. Csizmok, P. Tompa and I. Simon, 2005 IUPred: web server for the prediction of intrinsically unstructured regions of proteins based on estimated energy content. Bioinformatics 21**:** 3433-3434.

El-Brolosy, M. A., Z. Kontarakis, A. Rossi, C. Kuenne, S. Gunther *et al.*, 2019 Genetic compensation triggered by mutant mRNA degradation. Nature 568**:** 193-197.

El-Brolosy, M. A., and D. Y. R. Stainier, 2017 Genetic compensation: A phenomenon in search of mechanisms. PLoS Genet 13**:** e1006780.

England, S., M. F. Batista, J. K. Mich, J. K. Chen and K. E. Lewis, 2011 Roles of Hedgehog pathway components and retinoic acid signalling in specifying zebrafish ventral spinal cord neurons. Development 138**:** 5121-5134.

England, S. J., P. C. Campbell, S. Banerjee, A. J. Swanson and K. E. Lewis, 2017 Identification and Expression Analysis of the Complete Family of Zebrafish pkd Genes. Front Cell Dev Biol 5**:** 5.

Erdos, G., and Z. Dosztanyi, 2020 Analyzing Protein Disorder with IUPred2A. Curr Protoc Bioinformatics 70**:** e99.

Feng, Y., and Q. Xu, 2010 Pivotal role of hmx2 and hmx3 in zebrafish inner ear and lateral line development. Dev Biol 339**:** 507-518.

Fortunato, S. A., M. Adamski, O. M. Ramos, S. Leininger, J. Liu *et al.*, 2014 Calcisponges have a ParaHox gene and dynamic expression of dispersed NK homeobox genes. Nature 514**:** 620-623.

French, C. R., T. Erickson, D. Callander, K. M. Berry, R. Koss *et al.*, 2007 Pbx homeodomain proteins pattern both the zebrafish retina and tectum. BMC Dev Biol 7**:** 85.

Gehring, W. J., 1985 The homeo box: a key to the understanding of development? Cell 40**:** 3-5.

Gentleman, R. C., V. J. Carey, D. M. Bates, B. Bolstad, M. Dettling *et al.*, 2004 Bioconductor: open software development for computational biology and bioinformatics. Genome Biol 5**:** R80.

Gongal, P. A., L. D. March, V. L. Holly, L. M. Pillay, K. M. Berry-Wynne *et al.*, 2011 Hmx4 regulates Sonic hedgehog signaling through control of retinoic acid synthesis during forebrain patterning. Dev Biol 355**:** 55-64.

Gould, C. M., F. Diella, A. Via, P. Puntervoll, C. Gemund *et al.*, 2010 ELM: the status of the 2010 eukaryotic linear motif resource. Nucleic Acids Res 38**:** D167-180.

Gowan, K., A. W. Helms, T. L. Hunsaker, T. Collisson, P. J. Ebert *et al.*, 2001 Crossinhibitory activities of Ngn1 and Math1 allow specification of distinct dorsal interneurons. Neuron 31**:** 219-232.

Gross, M. K., M. Dottori and M. Goulding, 2002 Lbx1 specifies somatosensory association interneurons in the dorsal spinal cord. Neuron 34**:** 535-549.

Grossmann, K. S., A. Giraudin, O. Britz, J. Zhang and M. Goulding, 2010 Genetic Dissection of Rhythmic Motor Networks in Mice. Progress in Brain Research 187**:** 19-37.

Hartwell, R. D., S. J. England, N. A. M. Monk, N. J. van Hateren, S. Baxendale *et al.*, 2019 Anteroposterior patterning of the zebrafish ear through Fgf- and Hh-dependent regulation of hmx3a expression. PLoS Genet 15**:** e1008051.

Higashijima, S., M. A. Masino, G. Mandel and J. R. Fetcho, 2004a Engrailed-1 expression marks a primitive class of inhibitory spinal interneuron. J Neurosci 24**:** 5827-5839.

Higashijima, S., M. Schaefer and J. R. Fetcho, 2004b Neurotransmitter properties of spinal interneurons in embryonic and larval zebrafish. J Comp Neurol 480**:** 19-37.

Higashijima, S. I., G. Mandel and J. R. Fetcho, 2004c Distribution of prospective glutamatergic, glycinergic, and gabaergic neurons in embryonic and larval zebrafish. Journal of Comparative Neurology 480**:** 1-8.

Higashijima, S. I., M. Schaefer and J. R. Fetcho, 2004d Neurotransmitter properties of spinal interneurons in embryonic and larval zebrafish. Journal of Comparative Neurology 480**:** 19-37.

Hilinski, W. C., J. R. Bostrom, S. J. England, J. L. Juárez-Morales, S. de Jager *et al.*, 2016 Lmx1b is required for the glutamatergic fates of a subset of spinal cord neurons. Neural Dev 11**:** 16.

Holland, P. W., 2013 Evolution of homeobox genes. Wiley Interdiscip Rev Dev Biol 2**:** 31-45.

Hoshijima, K., M. J. Jurynec, D. Klatt Shaw, A. M. Jacobi, M. A. Behlke *et al.*, 2019 Highly Efficient CRISPR-Cas9-Based Methods for Generating Deletion Mutations and F0 Embryos that Lack Gene Function in Zebrafish. Dev Cell 51**:** 645-657 e644.

Hu, Y., S. Xie and J. Yao, 2016 Identification of Novel Reference Genes Suitable for qRT-PCR Normalization with Respect to the Zebrafish Developmental Stage. PLoS One 11**:** e0149277.

Ishida, T., and K. Kinoshita, 2007 PrDOS: prediction of disordered protein regions from amino acid sequence. Nucleic Acids Res 35**:** W460-464.

Itoh, M., C. Kim, G. Palardy, T. Oda, Y. Jiang *et al.*, 2003 Mind bomb is a ubiquitin ligase that is essential for efficient activation of Notch signaling by Delta. Dev Cell. 4**:** 67-82.

Jiang, Y. J., M. Brand, C. P. Heisenberg, D. Beuchle, M. Furutani-Seiki *et al.*, 1996 Mutations affecting neurogenesis and brain morphology in the zebrafish, *Danio rerio*. Development 123**:** 205-216.

Juárez-Morales, J. L., C. J. Schulte, S. A. Pezoa, G. K. Vallejo, W. C. Hilinski *et al.*, 2016 Evx1 and Evx2 specify excitatory neurotransmitter fates and suppress inhibitory fates through a Pax2-independent mechanism. Neural Dev 11**:** 5.

Kettleborough, R. N. W., E. M. Busch-nentwich, S. A. Harvey, C. M. Dooley, E. D. Bruijn *et al.*, 2013 A systematic genome-wide analysis of zebrafish protein-coding gene function. 496**:** 494-497.

Kimmel, C. B., 1995 Stages of Embryonic Development of the Zebrafish pp.

Kimura, Y., Y. Hisano, A. Kawahara and S. Higashijima, 2014 Efficient generation of knock-in transgenic zebrafish carrying reporter/driver genes by CRISPR/Cas9-mediated genome engineering. Sci Rep 4**:** 6545.

Kimura, Y., Y. Okamura and S. Higashijima, 2006 alx, a zebrafish homolog of Chx10, marks ipsilateral descending excitatory interneurons that participate in the regulation of spinal locomotor circuits. J Neurosci 26**:** 5684-5697.

Kosugi, S., M. Hasebe, M. Tomita and H. Yanagawa, 2009 Systematic identification of cell cycle-dependent yeast nucleocytoplasmic shuttling proteins by prediction of composite motifs. Proc Natl Acad Sci U S A 106**:** 10171-10176.

Ku, W. C., W. K. Lau, Y. T. Tseng, C. M. Tzeng and S. K. Chiu, 2004 Dextran sulfate provides a quantitative and quick microarray hybridization reaction. Biochem Biophys Res Commun 315**:** 30-37.

Kumar, M., M. Gouw, S. Michael, H. Samano-Sanchez, R. Pancsa *et al.*, 2020 ELM-the eukaryotic linear motif resource in 2020. Nucleic Acids Res 48**:** D296-D306.

Labun, K., T. G. Montague, J. A. Gagnon, S. B. Thyme and E. Valen, 2016 CHOPCHOP v2: a web tool for the next generation of CRISPR genome engineering. Nucleic Acids Res 44**:** W272-276.

Labun, K., T. G. Montague, M. Krause, Y. N. Torres Cleuren, H. Tjeldnes *et al.*, 2019 CHOPCHOP v3: expanding the CRISPR web toolbox beyond genome editing. Nucleic Acids Res 47**:** W171-W174.

Larroux, C., B. Fahey, S. M. Degnan, M. Adamski, D. S. Rokhsar *et al.*, 2007 The NK homeobox gene cluster predates the origin of Hox genes. Curr Biol 17**:** 706-710.

Lauter, G., I. Soll and G. Hauptmann, 2011 Two-color fluorescent in situ hybridization in the embryonic zebrafish brain using differential detection systems. BMC Dev Biol 11**:** 43.

Lin, Q., L. W. L. Low, A. Lau, E. W. L. Chua, Y. Matsuoka *et al.*, 2019 Tracking genome-editing and associated molecular perturbations by SWATH mass spectrometry. Sci Rep 9**:** 15240.

Linding, R., L. J. Jensen, F. Diella, P. Bork, T. J. Gibson *et al.*, 2003 Protein disorder prediction: implications for structural proteomics. Structure 11**:** 1453-1459.

Lu, F. I., Y. H. Sun, C. Y. Wei, C. Thisse and B. Thisse, 2014 Tissue-specific derepression of TCF/LEF controls the activity of the Wnt/beta-catenin pathway. Nat Commun 5**:** 5368.

Madeira, F., Y. M. Park, J. Lee, N. Buso, T. Gur *et al.*, 2019 The EMBL-EBI search and sequence analysis tools APIs in 2019. Nucleic Acids Res 47**:** W636-W641.

Mann, C. M., G. Martinez-Galvez, J. M. Welker, W. A. Wierson, H. Ata *et al.*, 2019 The Gene Sculpt Suite: a set of tools for genome editing. Nucleic Acids Res 47**:** W175-W182.

Marcelli, F., G. Boisset and D. F. Schorderet, 2014 A dimerized HMX1 inhibits EPHA6/epha4b in mouse and zebrafish retinas. PLoS One 9**:** e100096.

Meszaros, B., G. Erdos and Z. Dosztanyi, 2018 IUPred2A: context-dependent prediction of protein disorder as a function of redox state and protein binding. Nucleic Acids Res 46**:** W329-W337.

Montague, T. G., J. M. Cruz, J. A. Gagnon, G. M. Church and E. Valen, 2014 CHOPCHOP: a CRISPR/Cas9 and TALEN web tool for genome editing. Nucleic Acids Res 42**:** W401-407.

Moran-Rivard, L., T. Kagawa, H. Saueressig, M. K. Gross, J. Burrill *et al.*, 2001 Evx1 is a postmitotic determinant of V0 interneuron identity in the spinal cord. Neuron 29**:** 385-399.

Muller, T., K. Anlag, H. Wildner, S. Britsch, M. Treier *et al.*, 2005 The bHLH factor Olig3 coordinates the specification of dorsal neurons in the spinal cord. Genes Dev 19**:** 733-743.

Muller, T., H. Brohmann, A. Pierani, P. A. Heppenstall, G. R. Lewin *et al.*, 2002 The homeodomain factor lbx1 distinguishes two major programs of neuronal differentiation in the dorsal spinal cord. Neuron 34**:** 551-562.

Müller, T., H. Brohmann, A. Pierani, P. a. Heppenstall, G. R. Lewin *et al.*, 2002 The homeodomain factor Lbx1 distinguishes two major programs of neuronal differentiation in the dorsal spinal cord. Neuron 34**:** 551-562.

Noro, B., J. Culi, D. J. McKay, W. Zhang and R. S. Mann, 2006 Distinct functions of homeodomain-containing and homeodomain-less isoforms encoded by homothorax. Genes Dev 20**:** 1636-1650.

Park, H. C., and B. Appel, 2003 Delta-Notch signaling regulates oligodendrocyte specification. Development 130**:** 3747-3755.

Peng, J., 2019 Gene redundancy and gene compensation: An updated view. J Genet Genomics 46**:** 329-333.

Pfeffer, P. L., T. Gerster, K. Lun, M. Brand and M. Busslinger, 1998 Characterization of three novel members of the zebrafish *Pax2/5/8* family: dependency of *Pax5* and *Pax8* expression on the *Pax2.1* (*noi*) function. Development 125**:** 3063-3074.

Pisani, D., W. Pett, M. Dohrmann, R. Feuda, O. Rota-Stabelli *et al.*, 2015 Genomic data do not support comb jellies as the sister group to all other animals. Proc Natl Acad Sci U S A 112**:** 15402-15407.

Porcher, C., E. C. Liao, Y. Fujiwara, L. I. Zon and S. H. Orkin, 1999 Specification of hematopoietic and vascular development by the bHLH transcription factor SCL without direct DNA binding. Development 126**:** 4603-4615.

Radivojac, P., V. Vacic, C. Haynes, R. R. Cocklin, A. Mohan *et al.*, 2010 Identification, analysis, and prediction of protein ubiquitination sites. Proteins 78**:** 365-380.

Ravet, E., D. Reynaud, M. Titeux, B. Izac, S. Fichelson *et al.*, 2004 Characterization of DNA-binding-dependent and -independent functions of SCL/TAL1 during human erythropoiesis. Blood 103**:** 3326-3335.

Rice, P., I. Longden and A. Bleasby, 2000 EMBOSS: the European Molecular Biology Open Software Suite. Trends Genet 16**:** 276-277.

Rossi, A., Z. Kontarakis, C. Gerri, H. Nolte, S. Holper *et al.*, 2015 Genetic compensation induced by deleterious mutations but not gene knockdowns. Nature 524**:** 230-233.

Sarachu, M., and M. Colet, 2005 wEMBOSS: a web interface for EMBOSS. Bioinformatics 21**:** 540-541.

Satou, C., Y. Kimura and S. Higashijima, 2012 Generation of multiple classes of V0 neurons in zebrafish spinal cord: progenitor heterogeneity and temporal control of neuronal diversity. J Neurosci 32**:** 1771-1783.

Schier, A. F., S. C. F. Neuhauss, M. Harvey, J. Malicki, L. Solnica-Krezel *et al.*, 1996 Mutations affecting the development of the embryonic zebrafish brain. Development 123**:** 165-178.

Simion, P., H. Philippe, D. Baurain, M. Jager, D. J. Richter *et al.*, 2017 A Large and Consistent Phylogenomic Dataset Supports Sponges as the Sister Group to All Other Animals. Curr Biol 27**:** 958-967.

Sordino, P., D. Duboule and T. Kondo, 1996 Zebrafish *Hoxa* and *Evx-2* genes: cloning, developmental expression and implications for the functional evolution of posterior *Hox* genes. Mech. Dev. 59**:** 165 - 175.

Sztal, T. E., E. A. McKaige, C. Williams, A. A. Ruparelia and R. J. Bryson-Richardson, 2018 Genetic compensation triggered by actin mutation prevents the muscle damage caused by loss of actin protein. PLoS Genet 14**:** e1007212.

Talpalar, A. E., J. Bouvier, L. Borgius, G. Fortin, A. Pierani *et al.*, 2013 Dual-mode operation of neuronal networks involved in left-right alternation. Nature 500**:** 85-88.

Tarraga, J., I. Medina, J. Carbonell, J. Huerta-Cepas, P. Minguez *et al.*, 2008 GEPAS, a web-based tool for microarray data analysis and interpretation. Nucleic Acids Res 36**:** W308-314.

Thaeron, C., F. Avaron, D. Casane, V. Borday, B. Thisse *et al.*, 2000 Zebrafish *evx1* is dynamically expressed during embryogenesis in subsets of interneurones, posterior gut and urogenital system. Mechanisms of Development 99**:** 167-172.

Topalidou, I., A. van Oudenaarden and M. Chalfie, 2011 Caenorhabditis elegans aristaless/Arx gene alr-1 restricts variable gene expression. Proc Natl Acad Sci U S A 108**:** 4063-4068.

Toyama, R., and I. Dawid, 1997 *lim6*, a novel LIM homeobox gene in the zebrafish: comparison of its expression pattern with *lim1*. Dev. Dyn. 209**:** 406-417.

Treffkorn, S., L. Kahnke, L. Hering and G. Mayer, 2018 Expression of NK cluster genes in the onychophoran Euperipatoides rowelli: implications for the evolution of NK family genes in nephrozoans. Evodevo 9**:** 17.

Truett, G. E., P. Heeger, R. L. Mynatt, A. A. Truett, J. A. Walker *et al.*, 2000 Preparation of PCR-quality mouse genomic DNA with hot sodium hydroxide and tris (HotSHOT). Biotechniques 29**:** 52, 54.

Valentin, G., P. Haas and D. Gilmour, 2007 The chemokine SDF1a coordinates tissue migration through the spatially restricted activation of Cxcr7 and Cxcr4b. Curr Biol 17**:** 1026-1031.

Via, A., C. M. Gould, C. Gemund, T. J. Gibson and M. Helmer-Citterich, 2009 A structure filter for the Eukaryotic Linear Motif Resource. BMC Bioinformatics 10**:** 351.

Wang, W., E. K. Chan, S. Baron, T. Van de Water and T. Lufkin, 2001 Hmx2 homeobox gene control of murine vestibular morphogenesis. Development 128**:** 5017-5029.

Wang, W., J. F. Grimmer, T. R. Van De Water and T. Lufkin, 2004 Hmx2 and Hmx3 homeobox genes direct development of the murine inner ear and hypothalamus and can be functionally replaced by Drosophila Hmx. Dev Cell 7**:** 439-453.

Wang, W., P. Lo, M. Frasch and T. Lufkin, 2000 Hmx: an evolutionary conserved homeobox gene family expressed in the developing nervous system in mice and Drosophila. Mech Dev 99**:** 123-137.

Wang, W., and T. Lufkin, 2005 Hmx homeobox gene function in inner ear and nervous system cell-type specification and development. Exp Cell Res 306**:** 373-379.

Wang, W., T. Van De Water and T. Lufkin, 1998 Inner ear and maternal reproductive defects in mice lacking the Hmx3 homeobox gene. Development 125**:** 621-634.

Wine-Lee, L., K. J. Ahn, R. D. Richardson, Y. Mishina, K. M. Lyons *et al.*, 2004 Signaling through BMP type 1 receptors is required for development of interneuron cell types in the dorsal spinal cord. Development 131**:** 5393-5403.

Wotton, K. R., F. K. Weierud, J. L. Juarez-Morales, L. E. Alvares, S. Dietrich *et al.*, 2010 Conservation of gene linkage in dispersed vertebrate NK homeobox clusters. Dev Genes Evol 219**:** 481-496.

Zhao, Q., Y. Xie, Y. Zheng, S. Jiang, W. Liu *et al.*, 2014 GPS-SUMO: a tool for the prediction of sumoylation sites and SUMO-interaction motifs. Nucleic Acids Res 42**:** W325-330.

Zheng, C., and M. Chalfie, 2016 Securing Neuronal Cell Fate in C. elegans. Curr Top Dev Biol 116**:** 167-180.

Zheng, C., F. Q. Jin and M. Chalfie, 2015 Hox Proteins Act as Transcriptional Guarantors to Ensure Terminal Differentiation. Cell Rep 13**:** 1343-1352.

Zhu, P., Z. Ma, L. Guo, W. Zhang, Q. Zhang *et al.*, 2017 Short body length phenotype is compensated by the upregulation of nidogen family members in a deleterious nid1a mutation of zebrafish. J Genet Genomics 44**:** 553-556.

Tables and Table Legends

Table 1.

Gene name	Previous names	ZFIN ID	Reference for probe
hmx1	*nkx5.3,nkx5-3, im:7155045, zgc:172164*	ZDB-GENE-080204-54	This publication – see materials and methods.
hmx2	*nkx5.2*	ZDB-GENE-080506-2	(Feng and Xu 2010)
hmx3a	*hmx3, Nkx5-1, nkx5.1, zgc:109845*	ZDB-GENE-001020-1	(Feng and Xu 2010)
hmx3b	*si:zfos-911d5.2*	ZDB-GENE-130603-102	This publication – see materials and methods.
hmx4	*hmx3b, zfSOHo, wu:fc21b03, zgc:154104*	ZDB-GENE-060825-142	(Feng and Xu, 2010)
slc32a1	*viaat, zgc:158324*	ZDB-GENE-061201-1	(Kimura *et al.* 2006)
en1b	*eng1b, eng4, engrailed 4, wu:fc84e11, zgc:113499*	ZDB-GENE-980526-6	Gift of Drs. Kikuci & Westerfield, University of Oregon.
slc17a6a	*slc17a6l, VGLUT2.2, vglut2b, wu:fj12c01*	ZDB-GENE-050105-4	(Higashijima *et al.* 2004c; Higashijima *et al.* 2004d)
slc17a6b	*slc17a6, blu, blumenkohl, si:packt73.2; si:packt90.2, si:xx-packt73.2, VGLUT2.1, vglut2a*	ZDB-GENE-030616-554	(Higashijima *et al.* 2004c; Higashijima *et al.* 2004d)
lhx1a	*lim1, lim1a, wu:fe25d04, wu:fj36b06*	ZDB-GENE-980526-347	(Toyama and Dawid 1997)
evx1	*cb416*	ZDB-GENE-980526-364	(Thaeron *et al.* 2000)
evx2	None	ZDB-GENE-980526-215	(Sordino *et al.* 1996)
slc6a5	*glyt2, si:dkey-7n10.12*	ZDB-GENE-050105-2	(Higashijima *et al.* 2004c; Higashijima *et al.* 2004d)
gad1b	*gad1, GAD67, gad67*	ZDB-GENE-030909-3	(Higashijima *et al.* 2004c; Higashijima *et al.* 2004d)
krt15	*wu:fa91c06, wu:fa95d04, zgc:55334*	ZDB-GENE-040426-2931	(Valentin *et al.* 2007)
pax5	None	ZDB-GENE-001030-2	(Pfeffer *et al.* 1998)

Table 1. Gene names, previous names, ZFIN identifiers and references for *in situ* hybridization probes. Column 1 lists genes used in this study. Previous names, where known, are provided in column 2. Column 3 contains the unique ZFIN identification number for each gene. Column 4 provides the reference for the *in situ* hybridization RNA riboprobe used in our experiments.

Table 2.

SU Allele	Gene(s)	sgRNA(s) Sequence(s)	F0 Screening Method(s)	F0 Screening Primer Sequences	Stable Mutant Screening Method(s)	Stable Mutant Screening Primer Sequences
35	*hmx2*	GGTATGGATGGTGTCAAATG	HRMA	FW: TGTTTGTGTGTCGCATTGGTTG RV: TGTCTTCAGAAACGGGAGACAT	Sequencing	FW: CAACAGTCGATCAAACAGTCAGAG RV: GTGGACTCGAGTTGATAAACCTG
36	*hmx2*	GGTATGGATGGTGTCAAATG	HRMA	FW: TGTTTGTGTGTCGCATTGGTTG RV: TGTCTTCAGAAACGGGAGACAT	HRMA	FW: TGTTTGTGTGTCGCATTGGTTG RV: TGTCTTCAGAAACGGGAGACAT
37	*hmx2*	GGTATGGATGGTGTCAAATG	HRMA	FW: TGTTTGTGTGTCGCATTGGTTG RV: TGTCTTCAGAAACGGGAGACAT	PCR	FW: CAACAGTCGATCAAACAGTCAGAG RV: GTGGACTCGAGTTGATAAACCTG
38	*hmx2*	5': GGTATGGATGGTGTCAAATG 3': GGTTCCAGGATAATACAGTG	PCR	FW: TCAACAGTCGATCAAACAGTCAG RV: GAACCGTGTTGCTTTTGTGTTTTAC	PCR	FW: TCAACAGTCGATCAAACAGTCAG RV: GAACCGTGTTGCTTTTGTGTTTTAC
39	*hmx2*	5': GGACCGGACTCCGTCGTTTG 3': GGTTCCAGGATAATACAGTG	Nested PCR	Nested 1 FW: GCAAATGAGGGCACGTTTCAC RV: GGACATATACAGAACCGTGTTGC Nested 2 FW: TGAATAATTCGGAGGACAGCGGAAG RV: CGTAGACTCGTTTAATTGCTGC	Nested PCR (for deletion) and WT amplicon*	Nested 1 FW: GCAAATGAGGGCACGTTTCAC RV: GGACATATACAGAACCGTGTTGC Nested 2 FW: TGAATAATTCGGAGGACAGCGGAAG RV: CGTAGACTCGTTTAATTGCTGC WT Amplicon FW: CAACAGTCGATCAAACAGTCAGAG RV: GTGGACTCGAGTTGATAAACCTG
MENTHU	*hmx2*	TCGGAGGATGACTGCAGCGC	T7 Endonuclease I	FW: GTCTATTCTGGGCACCTCAAAC RV: GTACACCAGGCTCAGAGCAGTA	Sequencing	FW: GTCTATTCTGGGCACCTCAAAC RV: GTACACCAGGCTCAGAGCAGTA
42	*hmx3a*	GGAATCTCGGTCGGTGCCTG	HRMA	FW: CGAATGCTAATTTGGCCTCTATTACT RV: TTTTGTTGTCGTCTTCATCGTCC	PCR + restriction digest	FW: TGGCAAAGTGACACGACCAG RV: GAGAACACCGTGCGAGTTTTC Restriction enzyme: BanI
3	*hmx3a*	GGAATCTCGGTCGGTGCCTG	HRMA	FW: CGAATGCTAATTTGGCCTCTATTACT RV: TTTTGTTGTCGTCTTCATCGTCC	PCR	FW: TGGCAAAGTGACACGACCAG RV: GAGAACACCGTGCGAGTTTTC
43	*hmx3a*	GGAATCTCGGTCGGTGCCTG	Fluorescence and/or PCR	FW: GAGAACACCGTGCGAGTTTTC RV: AGTTTCCTGGCAGCATATCCAG	Fluorescence and/or PCR and WT amplicon*	PCR FW: GAGAACACCGTGCGAGTTTTC RV: AGTTTCCTGGCAGCATATCCAG WT Amplicon FW: TGGCAAAGTGACACGACCAG RV: GAGAACACCGTGCGAGTTTTC

SU Allele	Gene(s)	sgRNA(s) Sequence(s)	F0 Screening		Stable Mutant Screening	
			Method(s)	Primer Sequences	Method(s)	Primer Sequences
44	hmx2; hmx3a	5': GGGGCACGTACAATAGCAAA 3': GGTTCCAGGATAATACAGTG	Nested PCR	Nested 1: FW: GTTAGACTTTGCAATGCTCGTG RV: GGACATATACAGAACCGTGTTGC Nested 2: FW: CCTTTTTGCTATTGCCTTCATC RV: CGTAGACTCGTTTAATTGCTGC	Nested PCR (for deletion) and WT amplicon*	Nested 1: FW: GTTAGACTTTGCAATGCTCGTG RV: GGACATATACAGAACCGTGTTGC Nested 2: FW: CCTTTTTGCTATTGCCTTCATC RV: CGTAGACTCGTTTAATTGCTGC WT Amplicon FW: TGGCAAAGTGACACGACCAG RV: GAGAACACCGTGCGAGTTTTC
45	hmx2; hmx3a	5': GGGGCACGTACAATAGCAAA 3': GGTTCCAGGATAATACAGTG	Nested PCR	Nested 1: FW: GTTAGACTTTGCAATGCTCGTG RV: GGACATATACAGAACCGTGTTGC Nested 2: FW: CCTTTTTGCTATTGCCTTCATC RV: CGTAGACTCGTTTAATTGCTGC	Nested PCR (for deletion) and WT amplicon*	Nested 1: FW: GTTAGACTTTGCAATGCTCGTG RV: GGACATATACAGAACCGTGTTGC Nested 2: FW: CCTTTTTGCTATTGCCTTCATC RV: CGTAGACTCGTTTAATTGCTGC WT Amplicon FW: TGGCAAAGTGACACGACCAG RV: GAGAACACCGTGCGAGTTTTC

Table 2. sgRNA and primer sequences used for *hmx2*, *hmx3a* and *hmx2;hmx3a* CRISPR mutagenesis. Column 1 lists SU allele designation. Column 2 provides the name of the targeted gene or genes. Column 3 shows the sgRNA sequences used. Columns 4 and 5 indicate the method and primer sequences used to screen founder fish (HRMA = high resolution melt analysis). Columns 6 and 7 indicate the method and primer sequences used to screen stable mutant fish. *In the case of large deletion alleles (where the WT PCR product would be too large to detect by standard PCR conditions), a second PCR (a WT amplicon PCR) is performed. The genomic region amplified by this PCR is only present in WT and heterozygous animals. For experimental conditions, see materials and methods.

Table 3.

Fig.	Comparison	Gene	Difference between two means	P value
3C	Uninjected control (73.1 ± 0.9) vs *hmx2;hmx3a* DKD (71.8 ± 1.3)	*hmx3a*	1↓	0.328^
3F	Uninjected control (47.8 ± 0.4) vs *hmx2;hmx3a* DKD (47.2 ± 1.0)	*en1b*	1↓	0.580[+]
3I	Uninjected control (105.4 ± 1.5) vs *hmx2* SKD (86.8 ± 1.6)	*slc17a6a/b*	19↓	**<0.001[+]**
3I	Uninjected control (105.4 ± 1.5) vs *hmx3a* SKD (82.8 ± 1.8)	*slc17a6a/b*	23↓	**<0.001[+]**
3I	Uninjected control (105.4 ± 1.5) vs *hmx2;hmx3a* DKD (70.3 ± 1.3)	*slc17a6a/b*	35↓	**<0.001[+]**
3I	*hmx2* SKD (86.8 ± 1.6) vs *hmx2;hmx3a* DKD (70.3 ± 1.3)	*slc17a6a/b*	17↓	**<0.001[+]**
3I	*hmx3a* SKD (82.8 ± 1.8) vs *hmx2;hmx3a* DKD (70.3 ± 1.3)	*slc17a6a/b*	13↓	**<0.001[+]**
3I	*hmx2* SKD (86.8 ± 1.6) vs *hmx3a* SKD (82.8 ± 1.8)	*slc17a6a/b*	4↓	0.138[+]
3L	Uninjected control (138.6 ± 2.2) vs *hmx2;hmx3a* DKD (167.5 ± 2.9)	*slc32a1*	29↑	**<0.001[+]**
3O	Uninjected control (105.4 ± 1.5) vs *hmx2;hmx3a* DKD + *MOR-hmx3a* mRNA (89.2 ± 2.0)	*slc17a6a/b*	16↓	**<0.001[+]**
3O	*hmx2;hmx3a* DKD (70.3 ± 1.3) vs *hmx2;hmx3a* DKD + *MOR-hmx3a* mRNA (89.2 ± 2.0)	*slc17a6a/b*	19↑	**<0.001[+]**
NS	WT (105.5 ± 2.2) vs *hmx2^{SU37}* (103.3 ± 0.7)	*slc17a6a/b*	2↓	0.582^
NS	WT (102.8 ± 0.9) vs *hmx2^{SU38}* (103.6 ± 1.8)	*slc17a6a/b*	1↓	0.700[+]
5K	WT (96.4 ± 1.3) vs *hmx2^{SU39}* (94.7 ± 1.1)	*slc17a6a/b*	2↓	0.351[+]
5G	WT (106.0 ± 0.9) vs *hmx3a^{SU42}* (108.0 ± 3.1)	*slc17a6a/b*	2↑	0.673^
5H	WT (102.2 ± 1.1) vs *hmx3a^{sa23054}* (101.0 ± 1.7)	*slc17a6a/b*	1↓	0.567[+]
NS	WT (97.0 ± 1.2) vs *hmx3a^{SU43}* (85.2 ± 1.2)	*slc17a6a/b*	12↓	**0.019^**
5I	WT (99.8 ± 1.9) vs *hmx3a^{SU3}* (86.4 ± 0.8)	*slc17a6a/b*	13↓	**<0.001[+]**
5J	WT (102.2 ± 1.9) vs *hmx2;hmx3a^{SU44}* (89.0 ± 1.9)	*slc17a6a/b*	13↓	**0.001[+]**
NS	WT (102.0 ± 1.7) vs *hmx2;hmx3a^{SU45}* (87.8 ± 2.2)	*slc17a6a/b*	14↓	**<0.001[+]**
5I&J	*hmx3a^{SU3}* (86.4 ± 0.8) vs *hmx2;hmx3a^{SU44}* (89.0 ± 1.9)	*slc17a6a/b*	3↑	0.243[+]
NS	*hmx3a^{SU3}* (86.4 ± 0.8) vs *hmx2;hmx3a^{SU45}* (87.8 ± 2.2)	*slc17a6a/b*	1↑	0.559§
NS	*hmx3a^{SU43}* (85.2 ± 1.2) vs *hmx2;hmx3a^{SU44}* (89.0 ± 1.9)	*slc17a6a/b*	4↑	0.126[+]
NS	*hmx3a^{SU43}* (85.2 ± 1.2) vs *hmx2;hmx3a^{SU45}* (87.8 ± 2.2)	*slc17a6a/b*	3↑	0.309§
6K	WT (69.6 ± 0.7) vs *hmx3a^{SU3}* (70.2 ± 0.9)	*hmx3a*	1↑	0.613[+]
6L	WT (47.6 ± 2.0) vs *hmx3a^{SU3}* (45.0 ± 1.7)	*en1b*	3↓	0.356[+]
6M	WT (154.4 ± 2.9) vs *hmx3a^{SU3}* (167.8 ± 2.3)	*slc32a1*	13↑	**0.004^**
6N	WT (158.0 ± 2.0) vs *hmx2;hmx3a^{SU44}* (169.2 ± 1.7)	*slc32a1*	11↑	**0.002[+]**
NS	WT (158.0 ± 1.3) vs *hmx3a^{SU43}* (167.6 ± 2.7)	*slc32a1*	10↑	**0.012[+]**
6O	WT (130.4 ± 1.4) vs *hmx3a^{SU3}* (130.0 + 3.0)	*slc6a5*	0	0.907[+]
NS	WT (111.8 ± 3.9) vs *hmx2;hmx3a^{SU44}* (118.3 ± 3.0)	*slc6a5*	7↑	0.234[+]
6AB	WT (79.0 ± 2.1) vs *hmx3a^{SU3}* (90.4 ± 2.4)	*gad1b*	11↑	**0.022^**
6AC	WT (77.5± 2.5) vs *hmx2;hmx3a^{SU44}* (87.4 ± 2.3)	*gad1b**	10↑	**0.014[+]**
6AB&AC	*hmx3a^{SU3}* (90.4 ± 2.4) vs *hmx2;hmx3a^{SU44}* (87.4 ± 2.3)	*gad1b**	3↓	0.382[+]
6AD	WT (151.4 ± 2.7) vs *hmx3a^{SU3}* (175.7 ± 3.5)	*gad1b* (48h)*	24↑	**<0.001[+]**
6AE	WT (143.5 ± 6.7) vs *hmx2;hmx3a^{SU44}* (170.2 ± 8.5)	*gad1b* (48h)*	27↑	**0.024[+]**
6AF	WT (152.4 ± 4.8) vs *hmx2^{SU39}* (153.6 ± 2.5)	*gad1b* (48h)*	1↑	0.608^

Fig.	Comparison	Gene	Difference between two means	P value
6AD&AE	*hmx3a*^{SU3} (175.7 ± 3.5) vs *hmx2;hmx3a*^{SU44} (170.2 ± 8.5)	*gad1b* (48h)*	5↓	0.562§

Table 3. Statistical comparisons of numbers of cells expressing particular genes.

Statistical comparisons between uninjected WT control or WT sibling embryos and knockdown or mutant embryos. First column indicates the figure panel that contains the relevant bar chart for the comparison. NS = not shown in a figure. Second column states the genotypes being compared. Single mutant and double deletion embryos have to be obtained from different parents because *hmx2* and *hmx3a* are adjacent to each other on chromosome 17. Therefore, these were always analyzed as separate experiments and single mutants were only compared to double deletion mutants when there was no statistically significant difference between the WT sibling cell counts in the two experiments. Numbers within parentheses indicate mean numbers of cells ± S.E.M. In all cases, numbers are an average of at least 4 embryos and cells were counted in all dorsal-ventral spinal cord rows. All of the experiments were conducted on 27 h embryos except those indicated with (48 h) next to the gene name, which used 48 h embryos. 27 h embryos fixed on different days varied slightly in stage from prim stage 9 to 12. This explains the small differences in numbers of cells labelled with a particular probe in WT embryos in different experiments. Column three lists the gene that the cell counts and statistical comparison refer to. Asterisks indicate experiments performed with dextran sulfate (see materials and methods). The fourth column indicates the difference between the two mean values for the embryos being compared. All values are rounded to the nearest whole number. ↑ = increase, ↓ = decrease. Last column shows the P value for the comparison, rounded to three decimal places. Statistically significant (P<0.05) values are indicated in bold. Statistical test used is indicated by superscript symbol: Wilcoxon-Mann-Whitney test ($^\wedge$), type 2 Student's t-test ($^+$) or type 3 Student's t-test (§). For a discussion of why particular tests were used, see materials and methods.

Table 4.

Mutant allele	% Otolith fusion phenotype	P value for otolith fusion phenotype	% Survival	P value for survival	% Stalled lateral line progression	P value for stalled lateral line progression
SU35	0% (n=100; z)	N/A	18.03% (n=61)	0.234	0% (n=28)	N/A
SU36	0% (n=100; z)	N/A	27.27% (n=66)	0.566	0% (n=26)	N/A
SU37	0% (n=112; z)	N/A	20.19% (n=104)	0.258	0% (n=48)	N/A
SU38	0% (n=56; z) 0% (n=77; mz)	N/A	33.30% (n=54)	0.214	0% (n=104)	N/A
SU39	0% (n=49; z) 0% (n=228; mz)	N/A	30.26% (n=76)	0.289	0.69% (n=145)	**<0.001**
sa23054	12.35% (n=332; z) 27.41% (n=135; mz)	**<0.001** 0.552	20.78% (n=77)	0.428	0% (n=124)	N/A
SU42	0% (n=121; z)	N/A	20.00% (n=65)	0.388	0% (n=104)	N/A
SU3	24.85% (n=2370; z)	0.850	NONE	N/A	20.36% (n=334)	0.935
SU43	23.33% (n=463, z)	0.391	NONE	N/A	22.94% (n=109)	0.995
SU44	25.41% (n=1334; z)	0.752	NONE	N/A	25.23% (n=329)	1.000
SU45	29.52% (n=227; z)	0.126	NONE	N/A	19.44% (n=108)	0.879

Table 4. Statistical analyses of whether the frequencies of homozygous mutants that survive to adulthood or have abnormal otolith development or lateral line progression phenotypes are Mendelian.

Column one indicates the mutant allele. These are listed in the same order as in Fig. 4. Columns two, four and five show the frequency of embryos with otolith fusion phenotypes from incrosses of heterozygous (z) or homozygous (mz) parents, the frequency of viable adult homozygous mutants and the frequency of embryos with stalled lateral line progression respectively. n = total number of animals examined. Columns three, five and seven list the P value, rounded up to 3 decimal places, for a Chi-squared test of the hypothesis that the frequency of embryos with fused otoliths, adult homozygous viability or stalled lateral line progression respectively is Mendelian (25% for crosses of heterozygous parents (z) and 100% for crosses of homozygous parents (mz)). Statistically significant values are indicated in bold. N/A = not applicable (cases where no homozygous mutants survived, had otolith fusions or stalled lateral line phenotypes).

Table 5.

Allele combination	% Ear phenotypes	# Embryos analyzed	P value	% Severe phenotype	% Moderate phenotype	% Weak phenotype	% Stalled lateral line progression	# Embryos analyzed	P value	% Strong phenotype	% Weaker phenotype
SU39 x SU44	0.7%	268	**<0.001**	0.0%	0.0%	0.7%	0.0%	38	**<0.001**	0.0%	0.0%
SU3 x SU43	22.6%	623	0.165	22.5%	0.0%	0.1%	32.4%	37	0.250	32.4%	0.0%
SU3 x SU44	26.8%	276	0.487	26.8%	0.0%	0.0%	19.4%	36	0.441	19.4%	0.0%
SU43 x SU44	28.3%	279	0.214	28.0%	0.0%	0.3%	31.8%	44	0.296	22.7%	9.1%
sa23054 x SU44	22.4%	303	0.289	22.4%	0.0%	0.0%	26.3%	38	0.703	18.4%	7.9%
sa23054 x SU3	26.6%	139	0.696	26.6%	0.0%	0.0%	17.6%	34	0.244	0.0%	17.6%
SU42 x SU44	22.1%	208	0.337	19.7%	1.4%	1.0%	18.9%	37	0.703	0.0%	18.9%
SU3 x SU42	17.1%	1794	**<0.001**	4.7%	6.0%	6.4%	23.3%	192	0.614	2.1%	21.2%

Table 5: Trans-heterozygous mutant analyses identify an allelic series of *hmx3a* mutants.

Parents heterozygous for different mutant alleles were mated and otolith and lateral line progression phenotypes were assayed in their progeny. Otoliths were assayed at 3 dpf by visual inspection down a stereomicroscope. Lateral line progression was assayed by *in situ* hybridization for *krt15* at 27 h. Column one indicates the allele combination tested. Column two shows the percentage of embryos with otolith phenotypes, column three indicates the total number of embryos analyzed. In contrast to incrosses of *hmx3a^{SU3}* or *hmx3a^{SU43}*, where mutant embryos have fused otoliths in both ears ("severe" phenotype), in some trans-heterozygous crosses we observed two additional types of ear phenotypes. Phenotypes were classified as "moderate" if otoliths were adjacent but not fused in both ears, and as "weak" if there was an otolith fusion or adjacent otolith phenotype in only one ear. The percentage of embryos with each of these phenotypes is provided in columns five (severe), six (moderate) and seven (weak). Column eight indicates the percentage of embryos with stalled lateral line progression and column nine shows the total number of embryos analyzed. In contrast to incrosses of *hmx3a^{SU3}* or *hmx3a^{SU43}*, where mutant embryos lack any migration of the lateral line primordium along the trunk ("strong" phenotype), in some trans-heterozygous crosses, we observed embryos where the lateral line primordium had migrated slightly more caudally ("weaker"

phenotype). Column 11 shows the percentage of embryos with the strong phenotype, column 12 shows the percentage of embryos with the slightly weaker phenotype. Chi-squared tests were performed to test if the frequency of embryos with otolith fusion or lateral line phenotypes was Mendelian and the P values for these tests are provided in columns 4 or 10 respectively. Statistically significant values are indicated in bold. We also performed a binomial distribution test, using the cumulative distribution function, to test whether the number of embryos from the SU39 x SU44 cross that had fused otoliths was statistically significantly different from zero. P=0.264 for this test.

Table 6A-D.

Table 6A. Lateral line and otolith phenotypes of 40 embryos from *hmx3a^SU42/+* parents injected with *hmx2* CRISPR reagents

Lateral line phenotype	Fused otoliths both ears	Normal otoliths both ears	Total	
Strong	6	0	6	
Medium	4	0	4	
Weak	2	5	7	
Normal	1	22	23	
Total	13	27	40	

Table 6B. *hmx3a^SU42* genotypes of the 40 embryos included in Table 6A

Lateral line /Otolith phenotype	*hmx3a^SU42* homozgous	*hmx3a^SU42* heterozygous	*hmx3a^SU42* WT	Total
Strong/Fused	6	0	0	6
Medium/Fused	4	0	0	4
Weak/Fused	2	0	0	2
Normal/Fused	1	0	0	1
Weak/Normal	0	4	1	5
Normal/Normal	0	16	6	22
Total	13	20	7	40

Table 6C. Genotyping embryos with abnormal and normal otolith phenotypes

Otolith Phenotype	*hmx3a^SU42* homozygous	*hmx3a^SU42* heterozygous	*hmx3a^SU42* WT	Total
Fused both ears	32	1	0	33
Fused or adjacent one ear	3	1	0	4
Normal both ears	1	26	8	35
Total	36	28	8	72

Table 6D. *hmx3a^SU42* genotypes and otolith phenotypes of embryos genotyped for *hmx2*

Otolith phenotype	*hmx3a^SU42* homozygous	*hmx3a^SU42* heterozygous	*hmx3a^SU42* WT	% of *hmx2* mutant sequences
Fused both ears	9	1	0	60-90%
Fused one ear	1	1	0	60-75%
Normal both ears	0	4	7	60-90%

Table 6. Lateral line primordium and otolith phenotypes in embryos from incrosses of *hmx3a^SU42/+* parents, injected with *hmx2^MENTHU* CRISPR reagents

Table 6A. Embryos from an incross of $hmx3a^{SU42/+}$ parents were injected with $hmx2^{MENTHU}$ CRISPR reagents at the one-cell stage and assayed for lateral line primordium and otolith phenotypes at 30 h and 72 h respectively. Rows 2-5 show stalled lateral line primordium migration phenotypes: normal = primordium in expected position (over somite 15 at 30h); weak = primordium migration stalled by 2-3 somites; medium = primordium moderately stalled (over somite 6-7 at 30 h); strong = primordium not detected. Columns 2-3 show otolith phenotypes. Column 4 = total number of embryos with each lateral line phenotype. Row 6 = total number of embryos with each otolith phenotype.

Table 6B. $hmx3a^{SU42}$ genotypes of the 40 injected embryos screened for lateral line primordium and otolith phenotypes in Table 6A. Rows 2-7 show combinations of lateral line primordium phenotype (normal, weak, medium, strong, as in Table 6A) and otolith phenotype (fused in both ears, two normal otoliths in both ears, as in Table 6A). Columns 2-4 show $hmx3a^{SU42}$ genotypes. Column 5 = total number of embryos with each combination of lateral line primordium and otolith fusion phenotypes. Row 7 = total number of embryos with each $hmx3a^{SU42}$ genotype.

Table 6C. $hmx3a^{SU42}$ genotypes of embryos from a second incross of $hmx3a^{SU42/+}$ parents injected with $hmx2^{MENTHU}$ CRISPR reagents at the one-cell stage and assayed for otolith fusion phenotypes at 72 h. Rows 2-4 show otolith phenotypes. Columns 2-4 show $hmx3a^{SU42}$ genotypes. Column 5 = total number of embryos with each otolith phenotype. Row 5 = total number of embryos with each $hmx3a^{SU42}$ genotype.

Table 6D. A subset of 23 embryos screened in Table 6C were also sequenced to assess whether they contained any $hmx2^{MENTHU}$ mutant sequences. Rows 2-4 indicate otolith phenotypes. Columns 2-4 indicate $hmx3a^{SU42}$ genotypes. Column 5 = approximate percentage of $hmx2^{MENTHU}$ mutant sequences

detected in the PCR amplicons for embryos with each otolith phenotype. In all cases, the $hmx2^{MENTHU}$

mutant sequences represented at least 60% of all $hmx2$ sequences in each PCR amplicon.

Table 7.

Genotype / Embryo class	WT	Het	*hmx2*[SU39] mutants	N value	P value
Uninjected control	33.3%	57.6%	9.1%	33	0.11
WT-like	44.4%	22.2%	33.3%	9	0.12
Morphant-like	32.3%	41.9%	25.8%	31	0.68

Genotype / Embryo class	WT	*hmx2*[SU39] mutants	P value
Uninjected control	106.36 ± 1.75	109.67 ± 0.67	$0.21^{\wedge}$
WT-like	104.25 ± 5.30	98.33 ± 4.06	$0.86^{\wedge}$
Morphant-like	78.10 ± 2.50	71.25 ± 3.25	0.11^{+}
All injected	85.57 ± 3.96	78.64 ± 4.56	0.26^{+}

Table 7. The lack of spinal cord phenotypes in *hmx2*[SU39] mutants is not due to genetic compensation

Spinal cord phenotypes were assessed by *slc17a6a/b in situ* hybridization. *hmx2* morpholino-injected embryos from an incross of *hmx2*[SU39/+] parents were visually inspected on a stereomicroscope and categorized as resembling either a "WT-like" or "morphant-like" spinal cord phenotype. Embryos were genotyped to identify homozygous WTs and mutants. The spinal cord phenotypes of these embryos were then analyzed on a compound microscope while blind to genotype. Values in columns 2 and 3 of lower table indicate the mean number of labelled cells in the spinal cord region adjacent to somites 6-10 $\pm$ S.E.M. P value in upper table is from Chi-squared test. P value in bottom table is either from a Wilcoxon-Mann-Whitney test ($^{\wedge}$, performed when data was not normally distributed) or from a type 2 Student's t-test ($^{+}$, performed when data was normally distributed and variances were equal) for the comparison of homozygous WT embryos to homozygous *hmx2*[SU39] mutants for a particular classification (values on same row). All P values are rounded up to 2 decimal places. See materials and methods for more information on statistical tests.

Table 8.

Genotype Embryo class	WT	Het	$hmx3a^{SU42}$ mutants	N value	P value
Uninjected control	22.5%	52.5%	25.0%	40	0.93
WT-like	31.25%	31.25%	37.5%	16	0.30
Morphant-like	20.5%	56.4%	23.1%	39	0.61

Genotype Embryo class	WT		$hmx3a^{SU42}$ mutants	P value
Uninjected control	117.56 ± 2.26		116.10 ± 1.38	0.58[+]
WT-like	116.80 ± 2.89		114.67 ± 2.69	0.60[+]
Morphant-like	86.38 ± 2.15		88.22 ± 5.07	0.74[§]
All injected	98.08 ± 4.58		98.80 ± 4.67	0.91[+]

Table 8. The lack of spinal cord phenotypes in $hmx3a^{SU42}$ mutants is not due to genetic compensation

Spinal cord phenotypes were assessed by *slc17a6a/b in situ* hybridization. *hmx3a* morpholino-injected embryos from an incross of $hmx3a^{SU42/+}$ parents were visually inspected on a stereomicroscope and categorized as resembling either a "WT-like" or "morphant-like" spinal cord phenotype. Embryos were genotyped to identify homozygous WTs and mutants. The number of glutamatergic cells in the spinal cord region adjacent to somites 6-10 was then counted in each embryo, using a compound microscope, while blind to genotype. Values in columns 2 and 3 of the lower table indicate the mean number of labelled cells ± S.E.M. The P value in the upper table is from a Chi-squared test to determine if the frequencies of different genotypes were Mendelian. The P value in the bottom table is either from a type 2 ([+], performed when data was normally distributed and variances were equal), or type 3 Student's t-test ([§], performed when data was normally distributed and variances were unequal) for the comparison of homozygous WT embryos to homozygous $hmx3a^{SU42}$ mutants for a particular classification (values on same row). All P values are rounded up to 2 decimal places. See materials and methods for more information on statistical tests.

Table 9.

Genotype / Embryo class	WT	Het	*hmx3a*^{SU3} mutants	N value	P value
Uninjected control	20.4%	54.6%	25%	44	0.76
Weaker phenotype	19.5%	56.1%	24.4%	41	0.74
Morphant-like	21.7%	55.4%	22.9%	83	0.54

Genotype / Embryo class	WT	*hmx3a*^{SU3} mutants	P value
Uninjected control	106 ± 2.44	94.5 ± 2.62	**0.01**[+]
Weaker phenotype	102.75 ± 2.27	96.8 ± 1.77	0.05[+]
Morphant-like	70.22 ± 3.76	74.0 ± 2.75	0.72[^]
All injected	80.23 ± 4.01	81.86 ± 2.78	0.74[+]

Table 9. The incomplete penetrance of the spinal cord phenotype in *hmx3a*^{SU3} mutants is not due to genetic compensation

Spinal cord phenotypes were assessed by *slc17a6a/b in situ* hybridization. *hmx3a* morpholino-injected embryos from an incross of *hmx3a*$^{SU3/+}$ parents were visually inspected on a stereomicroscope and categorized as resembling either a "weaker" or more severe, "morphant-like" spinal cord phenotype. Embryos were genotyped to identify homozygous WTs and mutants. The number of glutamatergic cells in the spinal cord region adjacent to somites 6-10 was then counted in each embryo, using a compound microscope, while blind to genotype. Values in columns 2 and 3 of the lower table indicate the mean number of labelled cells ± S.E.M. The P value in the upper table is from a Chi-squared test to determine if the frequencies of different genotypes were Mendelian. The P value in the bottom table is either from a Wilcoxon-Mann-Whitney test (^, performed when data was not normally distributed) or from a type 2 Student's t-test (+, performed when data was normally distributed and variances were equal) for the comparison of homozygous WT embryos to homozygous *hmx3a*^{SU3} mutants for a particular classification

(values on same row). All P values are rounded up to 2 decimal places. See materials and methods for more information on statistical tests. Statistically significant values are indicated in bold.

<u>Figure Legends</u>

Figure 1. <u>Expression of *hmx* genes in WT zebrafish embryos</u>.

Lateral views of *hmx* expression in spinal cord (**A-Y**), hindbrain (**N', S', X'**), eye and ear (**A', E', F', J',** **K', O', P', T', U', Y'**) and lateral line primordium and neuromasts (**B-C, G-H, L-M, Q-R**) at 19 h (**A-E, A', E'**), 24 h (**D'**), 27 h (**F-J, F', J'**), 36 h (**K-O, K'-O'**), 42 h (**P-T, P', S'**) and 48 h (**U-Y, U', X',** **Y'**). Rostral, left; Dorsal, up. (**B-C, G-H, L-M, Q-R, V-W**) *hmx2* and *hmx3a* are expressed in spinal cord, lateral line primordium (white asterisks), neuromasts (black asterisks), and anterior ear (data not shown) at all stages examined, although *hmx2* spinal cord expression initially appears weaker than *hmx3a* and does not extend as far caudally. (**A, A', E, E', F, F', J, J', K, K', O, O', P, P', T, T', U, U',** **Y, Y'**) *hmx1* and *hmx4* are not expressed in spinal cord at any of these stages but are expressed in the eye (black arrowheads) and posterior-ventral ear and adjacent ganglion of the anterior lateral line (white arrowheads). (**D, I, N, N', S, S', X, X'**) *hmx3b* is not expressed in WT spinal cord at any of these stages. The only expression we observed was in the hindbrain between 36 and 48 h (black arrows). (**D'**) *hmx3b* is also not expressed in *mib1^(ta52b)* mutants. Inset shows spinal cord but there is also no expression in any other tissue. (**F', I, N', O', P', S, S', T, U', X, X', Y'**). The background (diffuse, non-specific staining) in these pictures is higher because we exposed the embryos to prolonged staining to ensure that there was no weak spinal cord expression. Especially in the brain, this can lead to background staining as the large ventricles of the hindbrain trap anti-sense riboprobes. Scale Bar = 50 μm (**A-Y**), 120 μm (**A', E',** **F', J', K', N', O', P', S', T', U', X', Y'**).

Figure 2. <u>*hmx2* and *hmx3a* are expressed in V1 and dI2 interneurons.</u>

(**A**) Schematic showing neurotransmitter and post-mitotic transcription factor phenotypes of spinal cord interneurons found in the dorsal-ventral spinal cord region shown in panels C-H. It is currently unclear whether V0v interneurons express Lhx1/5 (*, see discussion in the results). (**B**) Heatmap analysis of

gene expression profiling of V1 interneurons. A three-class ANOVA analysis of differential expression was performed on different FAC-sorted populations of cells. Class 1: All post-mitotic spinal neurons. Class 2: V1 interneurons. Class 3: All trunk cells. Each column is a different biological replicate. Rows show relative expression levels for a single transcription factor gene as normalized data transformed to a mean of 0, with standard deviation of +1 (highly expressed) or -1 (weakly/not expressed) sigma units. Adjusted P values corrected for multiple testing are <0.000001 for all genes shown. For more information on these experiments see (Cerda *et al.* 2009). Expression profiles are included for positive control genes (green), *en1b* and *pax2a,* that are expressed by V1 cells and negative control genes (purple), *gata3* and *vsx1,* that are expressed by other spinal cord interneurons but not V1 cells. Data for *lhx1a, lhx5* and *hmx3a* shows that they are co-expressed in V1 interneurons. (**C-H**) Lateral views of spinal cord at 27 h. Rostral, left; Dorsal, up. *hmx2* and *hmx3a* are co-expressed (**C**) in V1 and dI2 interneurons. (**D-H**) V1 interneurons are inhibitory (*slc32a1*-expressing) (**D**) and express *en1b* (**E**). dI2 interneurons are glutamatergic (excitatory, express *slc17a6a/b*) (**F**) and express *lhx1a* (**G**) but not *evx1* (**H**). Asterisks = double-labeled and white crosses = single-labeled *hmx3a*-expressing cells (green). Expression of other genes is red. Scale bar = 20 μm.

Figure 3. *hmx2/3a* double knockdown (DKD) embryos have fewer excitatory (glutamatergic) and more inhibitory spinal cord interneurons.

(**A-B, D-E, G-H, J-K, M-N**) Lateral views of spinal cord at 27 h. Rostral, left; Dorsal, up. (**C, F, I, L, O**) Mean number of cells expressing specific genes in a precisely-defined spinal cord region adjacent to somites 6-10 at 27 h. All counts are an average of at least 5 embryos. Error bars indicate S.E.M. Statistically significant ($p < 0.001$) comparisons are indicated with brackets and three asterisks. All data were first analyzed for normality using the Shapiro-Wilk test. Data set in **C** is non-normally distributed and was, therefore, analyzed with the Wilcoxon-Mann-Whitney test. Data sets in **F, I, L, O** are normally distributed and so the F test for equal variances was performed. Each pairwise combination of data sets

has equal variances and so a type 2 (for equal variances) student's t-test was performed. Mean numbers of cells and P-values are provided in Table 3. (**A-F**) There is no change in the number of *hmx3a-* or *en1b*-expressing spinal cells in *hmx2/3a* DKD compared to uninjected control embryos, suggesting that V1 and dI2 interneurons do not die or transfate/change into different cell types. (**G-I**) The number of *slc17a6a/b*-expressing (excitatory) spinal cells is reduced in both double and single knockdown (SKD) embryos, with the reduction being more severe in DKD embryos. (**J-L**) Concomitantly, there is a statistically significant increase in the number of *slc32a1*-expressing (inhibitory) cells in DKD embryos. (**M-O**) Injection of a morpholino-resistant *hmx3a* mRNA partially rescues the number of spinal excitatory cells in DKD embryos. Scale bar = 30 μm.

Figure 4. Summary of *hmx2/3a* mutant alleles analyzed and their phenotypes when homozygous.

Left-hand side: schematics of 11 mutant alleles and one double mutant analyzed. Top row = genomic locus; lower rows indicate predicted protein products. There is a 6454 bp gap between *hmx2* and *hmx3a*. Right-hand side: Column 1: allele number. Column 2: indicates whether embryos with fused otoliths were observed in incrosses of heterozygous (z) or homozygous (mz) parents. (Also see Table 4 and Fig. 5). Column 3 indicates whether a reduction in the number of spinal excitatory cells was observed at 27 h in homozygous mutants, as assayed by *in situ* hybridization for *slc17a6a/b* (Fig. 5 and data not shown). Column 4: indicates whether viable adult homozygous mutants were recovered. In all cases where adult homozygous mutants were identified, the numbers of these fish are not statistically significantly different (P> 0.214) from expected mendelian ratios, as assayed by a Chi-squared test. P values are provided in Table 4. Column 5: indicates whether embryos with stalled lateral line progression phenotypes were observed in incrosses of heterozygous parents. All *hmx2* stable mutant alleles recovered to date are homozygous viable and homozygous mutants do not have a reduction in the number of glutamatergic spinal cord interneurons or otolith or stalled lateral line progression phenotypes. This is the case, even for embryos from incrosses of fish homozygous mutant for the most severely deleted *hmx2* alleles

(*hmx2^{SU38}* and *hmx2^{SU39}*). *hmx3a^{sa23054}* mutants are also homozygous viable. They have variable, incompletely penetrant, otolith fusion phenotypes. Strikingly, only 27.41% of embryos from an incross of homozygous mutant parents have otolith fusion phenotypes (Table 4). *hmx3a^{sa23054}* mutants do not have a reduction in the number of glutamatergic spinal cord interneurons or stalled lateral line progression phenotypes. *hmx3a^{SU42}* mutants are homozygous viable and do not have any obvious abnormal phenotypes, even though this allele should encode a protein with the same number of WT amino acids as *hmx3a^{SU43}* and only one more WT amino acid than *hmx3a^{SU3}*. *hmx2;hmx3a^{SU44}* and *hmx2;hmx3a^{SU45}* differ only in the amount of upstream sequence that is deleted and have identical phenotypes.

Figure 5. Only some *hmx2/3a* alleles have mutant phenotypes.

(**A-F, L-AI, L', R' U'**) Lateral views. Rostral, left; Dorsal, up. (**A-F**) Expression of *slc17a6a/b* in spinal cord at 27 h. (**G-K**) Mean number of cells expressing *slc17a6a/b* in a precisely-defined spinal cord region adjacent to somites 6-10 at 27 h. All counts are an average of at least 4 embryos. Error bars indicate S.E.M. Statistically significant ($p < 0.001$) comparisons are indicated with brackets and three asterisks. All data were first analyzed for normality using the Shapiro-Wilk test. Data in **G** is not normally distributed and so a Wilcoxon-Mann-Whitney test was performed. Data sets in **H-K** are normally distributed and so the F test for equal variances was performed, followed by a type 2 student's t-test (for equal variances). P-values are provided in Table 3. (**L-Q, L'**) Lateral line primordium phenotypes examined either by *hmx3a* expression (**L', M-N, Q**) or live (**L, O-P**) at 27 h. (**R-AI, R', U'**) Ear phenotypes examined either by *hmx3a* expression at 27 h (**R-S, U' & W**), live at 4 dpf (**R', T-V**), *pax5* expression at 24 h (**X-AA**) or phalloidin staining at 4 dpf (**AB-AI**). (**A-K**). There is no change in the number of spinal excitatory neurons in *hmx3a^{SU42}* (**B, G**) and *hmx3a^{sa23054}* (**C, H**) and *hmx2^{SU39}* (**F, K**) mutant embryos compared to WT (**A**). In contrast, there is a statistically significant reduction in *hmx3a^{SU3}* (**D, I**) and *hmx2;hmx3a^{SU44}* (**E, J**) mutant embryos. (**L-Q, L'**). At 27 h, the tip of the lateral

line primordium (black dotted line, **L**) has reached somite 10 (S10 = somite 10) in WT embryos (**L, L'**). This rate of migration is unchanged in *hmx3a^{SU42}*, *hmx3a^{sa23054}* and *hmx2^{SU39}* mutant embryos (**M-N, Q**). (**O-P**) In contrast, the lateral line primordium fails to migrate in *hmx3a^{SU3}* (**O**) and *hmx2;hmx3a^{SU44}* (**P**) mutant embryos. Instead, it is stalled adjacent to somites 1-4 (S4 = somite 4). (**R-S, U' & W**) *hmx3a* expression in the ear (inside white dotted lines) and in presumptive neuroblasts anterior to the ear (white arrowheads) is unchanged in *hmx3a^{SU42}* (**S**) and *hmx2^{SU39}* (**W**) mutants, compared to WT embryos (**R**), but is severely reduced in both the presumptive neuroblasts anterior to the ear and the anterior ear (black arrowhead) in *hmx3a^{SU3}* mutants (**U'**). (**T**) *hmx3a^{sa23054}* mutants show incompletely penetrant, variable otolith fusion phenotypes, ranging from no fusion (like WT ear in **R'**), through incomplete fusion (**T**), to complete fusion, like that observed at full penetrance in *hmx3a^{SU3}* (**U**) and *hmx2;hmx3a^{SU44}* (**V**) mutant embryos. (**X-AA**) The expression of *pax5* in the anterior ear (inside black dotted lines) is unchanged in *hmx2^{SU39}* (**AA**) mutants, compared to WT embryos (**X**), but is severely reduced (blue arrowhead) in both *hmx3a^{SU3}* (**Y**) and *hmx2;hmx3a^{SU44}* (**Z**) mutants. (**AB-AI**) The three cristae of the ear (white asterisks) form normally in WT (**AB**), *hmx3a^{SU3}* (**AC**), *hmx2;hmxsa^{SU44}* (**AD**) and *hmx2^{SU39}* (**AE**) mutants. In contrast, the spatially distinct anterior (utricular, red arrow) and posterior (saccular, white arrow) maculae are unchanged in *hmx2^{SU39}* (**AI**) mutants, compared to WT embryos (**AF**), but are fused and are located in a more medio-ventral position (white cross) in *hmx3a^{SU3}* (**AG**) and *hmx2;hmx3a^{SU44}* (**AH**) mutants. However, there are no obvious differences between *hmx3a^{SU3}* and *hmx2;hmx3a^{SU44}* mutants. Scale bar = 50 μm (**A-F**), 30 μm (**L-W**), 20 μm (**X-AA**), 60 μm (**AB-AI**). Panels **X-Y, AB-AC** and **AF-AG** are reproduced from (Hartwell *et al.* 2019) as per the Creative Commons Attribution (CC BY) license at *PLoS Genetics*.

Figure 6. Analysis of *hmx3a* single and *hmx2;hmx3a* deletion mutants.

(**A-J, P-AA & AG**) Lateral views of *hmx3a* (**A, F**), *en1b* (**B, G**) *slc32a1* (**C, D, H, I**) *slc6a5* (**E, J**), *gad1b* (**P-T, V-Z**) or *krt15* (**U, AA, AG**) expression in spinal cord (**A-J, P-T, V-Z**) or lateral line

primordium (**U, AA, AG**) at 27 h (**A-J, P, Q, U-W, AA, AG**) or 48 h (**R-T, X-Z**). Rostral, left; Dorsal, up. (**K-O, AB-AF**). Mean number of cells expressing specific genes in a precisely-defined spinal cord region adjacent to somites 6-10 at 27 h (**K-O, AB & AC**) or 48 h (**AD-AF**). All counts are an average of at least 5 embryos. Error bars indicate S.E.M. Statistically significant ($p < 0.05$) comparisons are indicated with brackets and asterisks. $p < 0.05$ = *, $p < 0.01$ = **, $p < 0.001$ = ***. White bars indicate WT data and black bars the appropriate mutant data as indicated in key under panel AG. All data were first analyzed for normality using the Shapiro-Wilk test. Data sets in M, AB and AF are non-normally distributed and were analyzed with the Wilcoxon-Mann-Whitney test. Data sets in K, L, N, O, AC, AD, AE are normally distributed and so the F test for equal variances was performed. All of these had equal variances, so a type 2 student's t-test was performed. P-values are provided in Table 3. (**A, B, F, G, K, L**) As in DKD embryos (Fig. 3), dI2 and V1 interneurons do not die, nor do dI2 interneurons transfate/change into V1 interneurons in *hmx3a*^{SU3} mutant embryos, since the numbers of *hmx3a*- (**A, F & K**) and *en1b*-expressing cells (**B, G & L**) do not change compared to WT embryos. There is a statistically significant increase in the number of inhibitory, *slc32a1*-expressing cells in *hmx3a*^{SU3} mutants (**C, H, M**) and *hmx2;hmx3a*^{SU44} mutants (**D, I, N**) compared to WT embryos. (**E, J, O**) However, at 27 h, the number of *slc6a5*-expressing cells is unchanged between WT and *hmx3a*^{SU3} mutants, whereas there is an increase in the number of GABAergic (*gad1b*-positive) in *hmx3a*^{SU3} (**P, V, AB**) and *hmx2;hmx3a*^{SU44} mutants (**Q, W, AC**), suggesting that the additional inhibitory cells in the mutant embryos are GABAergic and not glycinergic. (**R, S, X, Y, AD, AE**) There is an equivalent increase in GABAergic (*gad1b*-positive) cells at 48 h in *hmx3a*^{SU3} and *hmx2;hmx3a*^{SU44} mutant embryos. However, there is no change in the number of GABAergic (*gad1b*-positive) cells at 48 h in *hmx2*^{SU39} mutants, compared to WT embryos (**T, Z, AF**). (**U, AA, AG**). *hmx3a*$^{SU42/+}$*;hmx3a*$^{SU3/+}$ trans-het embryos have two different lateral line primordium progression phenotypes at 27 h. Scale bar = 50 μm.

Figure 7. Phenotypic and genotypic analysis of embryos from an incross of *hmx3a*$^{SU42/+}$ parents injected with *hmx2*^{MENTHU} CRISPR reagents.

(**A-B**) Lateral views of ear phenotypes at 4 dpf in live uninjected (**A**) and *hmx2*^{MENTHU} CRISPR-injected (**B**) *hmx3a*^{SU42} homozygous mutants. Rostral, left; Dorsal, top. The uninjected *hmx3a*^{SU42} homozygous mutant embryo has two normal otoliths in each ear (**A**). In contrast, the *hmx2*^{MENTHU} CRISPR-injected *hmx3a*^{SU42} homozygous mutant embryo has fused otoliths in both ears. (**C-D**) Wild type (top row) and *hmx2*^{MENTHU} (bottom row) genomic sequences on top. Each colored box represents a specific nucleotide in the *hmx2* coding sequence: A = green, C = blue, G = black, T = red. The *hmx2*^{MENTHU} mutant sequence contains a 5 bp deletion (CGCAG (red line)) which introduces a premature stop codon (red dashed box) 14-16 bases after the deletion. Sequencing traces (below) from individual embryos from an incross of *hmx3a*$^{SU42/+}$ parents injected with *hmx2*^{MENTHU} CRISPR reagents. In the injected embryos, *hmx2*^{MENTHU} mutant sequences (with the 5 bp deletion) comprise approximately 60% (**C**) to 90% (**D**) of all amplified sequences at this locus. Scale bar = 50 μm (A& B).

Figure 8. Expression of *hmx* genes in mutant zebrafish embryos and before the mid-blastula transition.

(**A-G, I-M**) Lateral views of expression in the spinal cord (**A-G**) at 27 h or whole embryos at 1.5 h (16-cells, **I-M**). (**A-G**) Rostral, left; Dorsal, up. (**I-M**) Animal pole, up. (**A-G**) *hmx3a* mRNA does not exhibit nonsense-mediated decay (NMD) in *hmx3a*^{SU42} (**A**), *hmx3a*^{sa23054} (**B**), *hmx3a*^{SU3} (**C**) or *hmx3a*^{SU43} (**D**) mutant embryos. *hmx1* (**E**), *hmx3b* (**F**) and *hmx4* (**G**) are not expressed in the spinal cord of *hmx2;hmx3a*^{SU44} deletion mutants. However, *hmx1* and *hmx4,* were still expressed in the head as shown in Fig. 1 (data not shown), confirming that the *in situ* hybridization experiment had worked. We never detect expression of *hmx3b* in WT embryos at 27 h (see Fig. 1). None of the *hmx* genes are maternally expressed at 1.5 h, as assessed by *in situ* hybridization (**I-M**), and, in the case of *hmx2* and *hmx3a*, qRT-PCR on whole embryos (**H**). No maternal expression of *hmx2* and *hmx3a* was detected and zygotic expression was not observed via qRT-PCR until 14 h (**H**). Scale Bar = 50 μm (**A-G**), 280 μm (**I-M**).

Supplementary Data Figure 1. Alignment of mouse and zebrafish Hmx2 and Hmx3(a) protein sequences.

Alignment of full-length mouse and zebrafish Hmx2 and Hmx3(a) protein sequences. Alignment performed using Clustal Omega version 1.2.4 and default parameters (Madeira *et al.* 2019). Blue text = homeobox domain. Green underlined text = last WT amino acid of Hmx3a^{SU3} protein. Gray shaded residues are conserved between all four Hmx proteins. Yellow shaded residues are conserved between both mouse Hmx proteins and zebrafish Hmx3a protein, but not zebrafish Hmx2 protein. : = amino acids with highly similar properties; . = amino acids with weakly similar properties. Other than the start methionine, upstream of the homeobox domain only 8 other amino acids are conserved between all four Hmx proteins. In contrast, 26 amino acids are conserved between all four Hmx proteins downstream of the homeobox domain. As expected, the homeobox domain is highly conserved (91%, 52/57 amino acids) across all four Hmx proteins.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

A — *hmx3a^{SU42/SU42}* Uninjected Control

B — *hmx3a^{SU42/SU42}; hmx2^{MENTHU/+}*

C

Wild-type *hmx2* sequence

G A C T G C A G C G C A G G A G A G G A C T C G

Mutant *hmx2* sequence (5 bp deletion)

G A C T G C A G G A G A G G A C T C G G G T G A

G A C T G C A G   G A G A G G A C T C G G G   T G A C

Site of 5 bp deletion

Premature stop codon

Wild-type sequence   ~60% Mutant sequences

D

Wild-type *hmx2* sequence

G A C T G C A G C G C A G G A G A G G A C T C G

Mutant *hmx2* sequence (5 bp deletion)

G A C T G C A G G A G A G G A C T C G G G T G A

G A C T G C A G   G A G A G G A C T C G G G   T G A C

Site of 5 bp deletion

Premature stop codon

Wild-type sequence   ~90% Mutant sequences

Figure 8

Supplementary Data Figure 1

```
Mouse Hmx3       MPEPGPDASGTASAPPPQPPPQPPAPKESPFSIRNLLNGDHHRPPP------KPQPPPRT 54
Zebrafish Hmx3a  MPE--------------TTQDTCASAKDSPFFIKNLLNSDSKPSKP------KPILAP-- 38
Mouse Hmx2       MGSK----------EDVGKGCPAAGGVSSFTIQSILGGGPSEAP---REPAGWPARKRS 46
Zebrafish Hmx2   MNNS----------EDSGSKCSP-APISSFTIQSILGTSNDGVRSAGKDSPKSQPRKRT 48
                 *  .              * * *::.:*. .

Mouse Hmx3       LFAPASAAAAAAAAAAAAAAKGALEGAAGFALSQVGDLAFPRFEIPAQRFALPAHYLERSP 114
Zebrafish Hmx3a  ----T-------------KAGL--DGSFSLSQVGEINFPRFELPTQRFALPA-YLERAS 77
Mouse Hmx2       LS-------------------------------------------------VSSEEEEPEE 58
Zebrafish Hmx2   LS-------------------------------------------------VSSED-DCSA 59
                                                                    :  :    .

Mouse Hmx3       AWWYPYTLTP--AGGHLPRPEASEKA------LLRDSSPASGTDRDSPEPLLKADPDHKE 166
Zebrafish Hmx3a  AWWYPYTLSA--S-AHLHRTEAAQ--------KARDSSPTTGTDRDSPELVLKSDPDAKD 126
Mouse Hmx2       GWKAPACFCPDPHGPKEPSPKHHTPIPFPCLGTPKGSGGAGPAASERTPFLSPSHPDFKE 118
Zebrafish Hmx2   GEDSGDCYCSEPGVPESCNPH--QPLN-FCLGATKGLLPVQDGIDRRPHLTPSILPDYKE 116
                 .          .        .            ::    .            ** *:

Mouse Hmx3       --LDSKSPDEIILEESDSEEGKKEGEAVPGAAGTTVGATTATPGSEDWKAGAESPEKKPA 224
Zebrafish Hmx3a  DEDDNKSGDEIVLEESDTEDGKKEG------------------GIDDWKKSDDGAD-KKP 167
Mouse Hmx2       EK--------------------E--------RLLPAGSPSPGPER--PRDGGAERQTG 146
Zebrafish Hmx2   EQ--------------------G--------RACSQMS--PVSED--RQRDGPDKQNN 142
                                                :        . : :

Mouse Hmx3       CRKKKTRTVFSRSQVFQLESTFDMKRYLSSSERAGLAASLHLTETQVKIWFQNRRNKWKR 284
Zebrafish Hmx3a  CRKKKTRTVFSRSQVFQLESTFDMKRYLSSSERAGLAASLHLTETQVKIWFQNRRNKWKR 227
Mouse Hmx2       AAKKKTRTVFSRSQVYQLESTFDMKRYLSSSERACLASSLQLTETQVKTWFQNRRNKWKR 206
Zebrafish Hmx2   SAKKKTRTVFSRSQVYQLESTFDMKRYLSSSERACLASSLQLTETQVKTWFQNRRNKWKR 202
                 . ****************:****************** **:**:******* *********

Mouse Hmx3       QLAAELEAANLSHAAAQRIVRVPILYHENSAAEGAAAAAGAPVPVSQPLLTFPHPVYYSH 344
Zebrafish Hmx3a  QLAAELEAANLSHAAAQRIVRVPILYHENSASESTNT--AGNVPVSQPLLTFPHPVYYSH 285
Mouse Hmx2       QLSAELEAANMAHASAQTLVGMPLVFRDSSLLR--------VPVP-RSLAFPAPLYYPS 256
Zebrafish Hmx2   QLSAELEAANMAHASAQTLVGMPLVFRENSLLR--------VPVP-RSIAFPTPLYYPG 252
                 **:*******::**:** :* :*::::.*  .       *** ::** *:**

Mouse Hmx3       PVVSSVPLLRPV----- 356
Zebrafish Hmx3a  PIVTSVPLLRPV----- 297
Mouse Hmx2       SNLSALPLYNLYNKLDY 273
Zebrafish Hmx2   SNLPALPLYNLYNKIEY 269
                 :  ::**  .
```

Homeodomain   Last wild-type amino acid in Zebrafish Hmx3a^{SU3} protein

* Conserved in all four Hmx proteins   Conserved in all but Zebrafish Hmx2 protein

: Residues with strongly similar properties

. Residues with weakly similar properties