1   **Evolution and maintenance of microbe-mediated protection**

2   **under occasional pathogen infection**

4   **Running title: Evolution of microbe-mediated protection**

6   Kloock, A.[1*], Bonsall, M.B.[1], King, K.C.[1]

[1] Department of Zoology, University of Oxford, 11a Mansfield Road, Oxford OX1 3SZ

* Corresponding author (anke.kloock@zoo.ox.ac.uk)

Authorship Contributions

KCK and AK conceived the project. KCK, MBB, and AK designed the project. AK performed

the evolution experiment and all subsequent assays. AK and MBB performed the statistical

analysis. AK, KCK and MBB wrote the manuscript and agreed on the final version.

Acknowledgments

We would like to thank the King group for help and support during the performance of the

evolution experiment, especially Charlotte Rafaluk-Mohr and Maria Ordovas-Montanes. AK was

supported by a fellowship from the "Studienstiftung des Deutschen Volkes". KCK is grateful for

a Leverhulme Trust project grant (RPG-2015-165) and ERC Starting Grant (COEVOPRO

802242).

Conflict of Interest Statement

The authors declare no conflict of interest.

Abstract

Every host is colonized by a variety of microbes, some of which can protect their hosts from

pathogen infection. However, pathogen presence naturally varies over time in nature, such as in

the case of seasonal epidemics. We experimentally coevolved populations of *Caenorhabditis*

elegans worm hosts with bacteria possessing protective traits (*Enterococcus faecalis*), in

treatments varying the infection frequency with pathogenic *Staphylococcus aureus* every host

generation, alternating host generations, every fifth host generation or never. We additionally

investigated the effect of initial pathogen presence at the formation of the defensive symbiosis.

Our results show that enhanced microbe-mediated protection evolved during host-protective

microbe coevolution when faced with rare infections by a pathogen. Initial pathogen presence had

no effect on the evolutionary outcome of microbe-mediated protection. We also found that

protection was only effective at preventing mortality during the time of pathogen infection.

Overall, our results suggest that resident microbes can be a form of transgenerational immunity

against rare pathogen infection.

Keywords: (1) Experimental evolution, (2) Defensive Symbiosis, (3) Heterogeneity, (4) Host-

pathogen interactions, (5) Protection,

Introduction

In nature, all plants and animals are colonised by microbes (Barriere, 2006; Ley et al., 2006; Vántus et al., 2014). The composition of these microbial communities is highly diverse and includes harmful, neutral, and beneficial microbial species (Ley et al., 2006), including those that can be important players in host defence against parasites, a phenomenon referred to as 'defensive mutualism' (King, 2019; May and Nelson, 2014). Recognised for over a century, defensive mutualism has been observed in plants (Mendes et al., 2011) and in a range of animals (Dillon et al., 2000; Dong et al., 2009; Jaenike et al., 2010; Koch and Schmid-Hempel, 2011), including humans (Kamada et al., 2013; Ley et al., 2006; Maynard et al., 2012) wherein microbes can supplement host immune systems (Abt and Artis, 2013; Hooper et al., 2012; McFall-Ngai et al., 2013).

The net benefits of defensive mutualism are dependent upon the presence of pathogens (Clay et al., 2005; King and Bonsall, 2017; Lively et al., 2005). Whilst hosts can benefit from microbe-mediated protection, defensive symbionts can be less beneficial to the host in the absence of enemies, due to metabolic and physiological costs (King, 2019). For example, in the interaction of aphids and the bacterium *Hamiltonella defensa*, the host tissue is harmed by defensive toxins that protect against infection from parasitoids (Vorburger and Gouskov, 2011). In some cases, possessing protective microbes might be more beneficial to the host than investing in its own immune system (Martinez et al., 2016). From the perspective of the symbiont, it is most useful to its host under high pathogen prevalence, and thus can persist in the host population (Palmer et al., 2008). Nevertheless, a stable symbiotic interaction is hypothesized to be evolved and maintained (Kwiatkowski and Vorburger, 2012) only when the host benefit of carrying defensive symbionts

outweighs any costs. The interactions of obligate and defensive symbionts and hosts can be stable

for millions of years (Moran et al., 2005).

Not all environments are constantly pathogen rich which might shift the balance of costs and

benefits during defensive mutualisms, particularly during coevolutionary interactions (King and

Bonsall, 2017). Pathogen prevalence can be spatially (King et al., 2009) or temporally variable,

the latter in the case of seasonal epidemics (e.g., flu peaks each winter in the northern hemisphere

(Finkelman, 2007) or rabies in North American skunks which peaks in Autumn (Gremillion-Smith

and Woolf, 1988)). Different environmental factors can influence disease transmission such as an

increase in malaria risk in warmer regions after rainfall (Altizer et al., 2006), or an increase in

contact rate and thus higher flu infection rate during the winter months (London and Yorke, 1973).

The impact of other temporally heterogeneous factors on the strength and direction of selection on

species interactions have been explored (oxygen concentration (Dey et al., 2016), resource

availability (Friman et al., 2011; Friman and Laakso, 2011; Hiltunen et al., 2012), environmental

productivity (Harrison et al., 2013)). Whether the varied presence of pathogens can similarly alter

selection for symbiotic interactions has been explored theoretically (Fenton et al., 2011), but

remains to be empirically tested.

Here, we examined the impact of temporal variation in pathogen infection on the evolution of

microbe-mediated protection. We used *Caenorhabditis elegans* as a worm host and allowed it to

be colonised by a bacterium (*Enterococcus faecalis*) that protects against infection by

Staphylococcus aureus (King et al., 2016). *Enterococcus faecalis* has been shown to be protective

across animal microbiomes (Kommineni et al., 2015; Martín-Vivaldi et al., 2010). It has been

previously shown that *E. faecalis* can evolve to provide enhanced protection when residing in

C. elegans hosts during constant pathogen infection (King et al., 2016; Rafaluk-Mohr et al., 2018).

From this, we predict that variation in pathogen infection might limit the evolution of microbe-

mediated protection. In the present study, we experimentally co-passaged *C. elegans* with

protective *E. faecalis* and infected the host with evolutionary static pathogenic *S. aureus* at

different intervals of host evolution. We also examined whether pathogen presence at the initial

formation of the coevolving interaction is crucial to the evolution of protection. We show that

enhanced microbe-mediated protection emerged out of novel coevolutionary host-microbe

interactions and during pathogen infection, regardless of its temporal variability or the time point

of first infection. Enhanced protection was only effective during pathogen infection. If hosts

survived infection, they could recover and had the same longevity and reproductive output across

treatments. These results thus suggest that even occasional pathogen infection can select for

defensive mutualism, revealing the potential for this phenomenon to be widespread in nature.

Materials and Methods

Worm host and bacteria system

As a bacteriovore, *Caenorhabditis elegans* interacts constantly with a variety of bacteria either by feeding or hosting them (Cabreiro and Gems, 2013; Garsin et al., 2001; Schulenburg and Ewbank, 2004). Consequently, *C. elegans* is an established model for studying innate immunity (Gravato-Nobre and Hodgkin, 2005), as it can be infected with its natural (Jansson, 1994; Schulenburg and Ewbank, 2004) as well as opportunistic pathogens (Garsin et al., 2001; Tan et al., 1999). Most pathogens are taken up orally by the worm (Marsh and May, 2012), and some can proliferate and colonize the worm gut (King et al., 2016; Rafaluk-Mohr et al., 2018).

Naturally, *C. elegans* is a self-fertilising hermaphrodite (Brenner, 1974), but in this experiment obligate outcrossing worm populations (line EEVD00) with males and females (hermaphrodites that carry the *fog-2(q71)* mutation) were used (Theologidis et al., 2014). This lineage was generated by Henrique Teotonio (ENS Paris) and encompasses the genetic diversity of 16 natural worm isolates (Theologidis et al., 2014). Worms were kept on Nematode Growth Medium (NGM), inoculated with *Salmonella*, hereafter referred to as food. Worms were infected with the pathogenic *S. aureus* (MSSA476) (Holden et al., 2004), which is virulent and kills worm hosts by lysing the intestinal cells lining the gut wall (Sifri et al., 2003). Worms were exposed to *E. faecalis* (OG1RF) (Garsin et al., 2001), which was isolated from the human digestive system, but has been previously shown to colonize and proliferate in the host gut (Ford et al., 2017; King et al., 2016; Rafaluk-Mohr et al., 2018), where it provides protection.

Experimental evolution - Design

Six single clones of *E. faecalis* (one for each of the six replicate populations) and a single population of *C. elegans* were the ancestors (hereafter referred to as the Ancestor) for all evolving populations. To account for potential differences in virulence, a stock of four clones of *S. aureus* was used for pathogen infections. Both *C. elegans* and colonising *E. faecalis* were allowed to evolve in presence of each other, while *S. aureus* was kept evolutionarily static. Infection with *S. aureus* was varied over host evolutionary time (indicated by purple in Table 1) to represent temporal heterogeneity in pathogen infection, including a range from always to every 2nd generation, every 5th generation, and never (Table 1). Moreover, we included differences in whether pathogens were present at the initial formation of the symbiotic interaction or later (2.1. vs. 2.2., and 5.1. vs. 5.2. in Table 1). Controls for lab adaptation were maintained for the host (No Protective-Microbe control, NPM in Table 1) and *E. faecalis* (No Host Control, NHC in Table 1).

Table 1: Experimental procedure for the evolution experiment. Columns indicate the number of experimental host generations (1- 20), while rows show the eight treatments. Host generations were infected with *S. aureus* (purple) or given food (green), while constantly coevolving with *E. faecalis*. Two controls for lab effects on host evolution (dark brown, No Protective Microbe, NPM) and *E. faecalis* evolution (light brown, No Host Control, NHC) were also included, where the NPM treatment was only ever exposed to food alone. Each evolutionary treatment consisted of six independent evolutionary replicates.

Experimental evolution - Culturing and passaging methods

At the start of each generation, worms were bleached as described previously and left in M9 buffer

overnight for larvae to hatch (Stiernagle, 2006). Simultaneously, *E. faecalis* clones were cultured

overnight in Todd-Hewitt Broth (THB) in 600µl at 30ºC, while food was cultured overnight in LB

broth. Subsequently, 9cm NGM plates were inoculated with 300µl of each overnight culture. Plates

with freshly inoculated bacteria were dried at room temperature before approximately 1000 L1

worms were added to each NGM plate. After these plates dried at room temperature, they were

transferred to a 20ºC incubator and left for 48h. Simultaneously, a liquid culture of *S. aureus* was

grown in THB from frozen stock, while a liquid culture of food was grown in LB, and both were

incubated under shaking conditions at 30ºC. The following day, 100µl of each overnight culture

were spread on 9cm plates, *S. aureus* on Tryptone Soy Broth agar (TSB) plates and food on NGM

plates and incubated at 30ºC overnight. To transfer worms to the pathogen or food plates,

nematodes were washed off the *E. faecalis* plates with M9 buffer and washed three times over

small-pore filters to remove all externally attached bacteria, as previously described (Jansen et al.,

2015; Papkou et al., 2019; Rafaluk-Mohr et al., 2018). Worms were infected with either *S. aureus*

or exposed to food (Table 1) and left at 25ºC for 24h. After this time, worms were then washed off

the plates with M9 buffer once more to plate them on NGM plates seeded with food for laying

eggs. Roughly 10% of these worms were crushed and plated on *E. faecalis* selective medium (TSB

+ 100mg/ml Rifampicin). The remaining worms were left on food plates for 48h to allow for egg

laying.

To passage *E. faecalis*, roughly 100 *E. faecalis* colonies were picked and grown up shaking

overnight in 600µl THB at 30ºC, while worms were bleached and left to hatch overnight. This

cycle was repeated for 20 experimental host generations.

All passaged worms and *E. faecalis* samples were cryopreserved at -80 ºC. A proportion of the

offspring of surviving worms were frozen in 40% DMSO, and 100µl of *E. faecalis* liquid culture

was mixed with 100µl of glycerol before cryopreservation.

Host survival and fecundity assays

All assays were conducted at the end of the evolution experiment on archived samples. Plates were

randomized and fully encoded during each experiment to ensure the experimenter was blind to

different treatments whilst collecting data.

Basic procedures were adopted from the experimental evolution, but with the following alterations

to keep the assays feasible with higher accuracy when scoring dead and alive worms: 400 L1

worms were exposed to 200µl of food and *E. faecalis* on 6cm NGM plates, while 60µl *S. aureus*

overnight culture was used to inoculate 6cm TSB plates.

To assess microbe-mediated protection of different combinations of worms and *E. faecalis*, 400

L1s were exposed to 50:50 mixtures of *E. faecalis* and food for 48h. Worms were then washed off

these plates as described above and infected with *S. aureus* for 24h at 25°C. Survival in form of

counting dead and alive worms was then scored.

To assess any long-term fitness consequences after protective microbe exposure and pathogen

infection, long-term survival and fecundity were measured. Worms were exposed as described for

the survival assays. Subsequently, five females and five males were picked onto 3cm food seeded

NGM plates at 25°C and then transferred to new plates every 36h to avoid any confusion between

offspring produced and original adults. At each time point, survival was scored. To measure

fecundity, the number of worm eggs on the plates at 120h since bleaching were counted.

Statistical Analysis

Statistical analyses were carried out with RStudio (Version 1.1.463 for Mac), graphs created with

the ggplot2 package (Version 2.1.0) and edited with Inkscape (Version 0.91). All host survival and

fecundity data were analysed with nested binomial mixed effects models (R package lme4),

followed by a Tukey multiple-comparison tests (R package multcomp). Life-span data were

analysed with Kaplan Meier Log Rank test with FDR correction for multiple testing.

Results

Figure 1: **Host survival showing protective effects of *E. faecalis*.** Early exposure of worms to *E. faecalis* (both ancestors) provides some degree of protection from the infection of *S. aureus*. 24h host survival levels reveal a benefit to *E. faecalis* colonisation independent of pathogen presence or absence. Circles indicate those treatment being exposed to *E. faecalis* and food in the earlier stage (L1-L4), while squares indicate food alone treatment in the earlier stage (L1-L4). Filled symbols indicate those treatments being exposed to food in the later stage, while open symbols indicate those treatments being exposed to the pathogen *S. aureus* in the later stage. Each symbol indicates the mean ± S.E of five replicates. Axis scales were chosen to be the same across all plots.

Before the start of the evolution experiment, the starting conditions were tested. Confirming previous results, *E. faecalis* showed some spontaneous host-protective potential against *S. aureus*. Worms raised on *E. faecalis* and food survived better than those raised on food alone, independent of food or pathogen present at the later stage (General Linear Model, X^2=10.205, df=1, p=0.001; Figure 1). Worms infected with *S. aureus* in later life survived worse than those being exposed to

190   food (General Linear Model, X^2=119.643, df=1, p<0.001; Figure 1). These results demonstrate the

191   beneficial and protective effects for the host after exposure to the protective microbe *E. faecalis*.

Figure 2: **Host survival for coevolving sympatric and allopatric pairs of worms and *E. faecalis***. Microbe-mediated protection was assessed for (A) sympatric pairs of coevolved worms and *E. faecalis,* (B) allopatric pairs of evolved worms and ancestral *E. faecalis*, and (C) allopatric pairs of ancestral worms and evolved *E. faecalis*. Bigger symbols represent mean ± S.E. and consists of six biological replicates and four technical replicates. Smaller symbols indicate the data distribution. Circles indicate sympatric pairs of coevolved *E. faecalis* and worms, squares indicate ancestral pairs of *E. faecalis* and worms and triangles indicate allopatric pairs of *E. faecalis* and worms. Letters indicate results of a GLMM, followed by a Tukey Post-hoc Test. The same letter indicates no significant difference. Axis scales were chosen to be the same across all plots.

Infection with *S. aureus* over evolutionary time in the experiment led to the substantial

enhancement of microbe-mediated protection, with the evolutionary background of the sympatric

pair of host and *E. faecalis* having a significant impact on host survival (Mixed Effects Model,

X^2=42.479, df=4, p<0.001; Figure 2A). Higher microbe-mediated protection in comparison to the

Ancestor occurred in all evolutionary histories involving pathogen presence across the temporal

heterogeneity treatments in our evolution experiment (Always, 2.1. and 5.1.). However, this did

not occur in the pathogen absence (Never) treatment. Host evolutionary history alone had a

significant effect on host survival (Mixed Effects Model, X^2=35.779, df=5, p<0.001; Figure 2B),

but did not reveal the same pattern as for sympatric pairs. No effect of bacteria evolutionary history

alone on infected host survival was observed (Mixed Effects Model, X^2=3.2511, df=5, p=0.6613;

Figure 2C). Taken together, enhanced microbe-mediated protection evolved only as a product of

coevolution and pathogen presence for sympatric pairs; this occurred regardless of the temporal

heterogeneity.

Figure 3: **Host survival in evolutionary treatments differing in initial pathogen exposure time points**. The time point of initial infection varied for infection to the pathogen every two generations (2.1. and 2.2) or every five generations (5.1. or 5.2.) but does not influence the outcome. Closed symbols indicate initial pathogen presence (Host generation 1), open symbols indicate later pathogen presence (Generation 2 for 2.1. and 2.2. and Generation 5 for 5.1. and 5.2.). Bigger symbols represent mean ± S.E and consists of six biological replicates and four technical replicates of the sympatric pairs. Smaller symbols indicate the data distribution. Letters indicate results of a GLMM, followed by a Tukey Post-hoc Test. The same letter indicates no significant difference. Axis scales were chosen to be the same across all plots.

As an additional form of pathogen heterogeneity, the impact of the timing of initial pathogen

infection on the evolution of microbe-mediated protection was investigated. An effect of different

initial pathogen infection time points on host survival following pathogen infection was observed

(Mixed Effects Model: X^2=7.945, df=3, p=0.04716 Figure 3), although a Tukey Post-Hoc test

revealed no significant differences (Table A1).

Figure 4: **Long-term survival and fecundity of *E. faecalis*-colonised hosts that survived pathogen infection**. (A)
Long-term host survival was measured. Survival curves for sympatric pairs of worms and *E. faecalis* are shown as
Kaplan-Meier estimates. Worms were exposed to *E. faecalis* and food (green), then to *S. aureus* (purple), and long-
term survival was monitored on food (orange). The dotted line indicates the time point at which fecundity was
measured. (B) Number of eggs/Female across sympatric pairs of coevolved worms and *E. faecalis*. Bigger symbols
represent mean ± S.E. and consists of six biological replicates and four technical replicates. Smaller symbols indicate
the data distribution. Circles indicate sympatric pairs of coevolved *E. faecalis* and worms, squares indicate ancestral
pairs of *E. faecalis* and worms. Letters indicate results of a GLMM, followed by a Tukey Post-hoc Test. The same
letter indicates no significant difference.

Furthermore, we investigated the long-term consequences to hosts colonised by *E. faecalis* after

24h of pathogen infection. No significant differences were observed in the long-term survival post-

infection of worm hosts colonised by their sympatric *E. faecalis* across treatments (Kaplan Meier

Log Rank Test, FDR corrected, all comparisons p>0.05, Figure 4A). In addition, we did not find

significant differences in fecundity among sympatric host-*E. faecalis* pairs (Mixed Effects Model,

X^2=3.9418, df=4, p=0.4278, Figure 4B).

Discussion

It has been shown that hosts receive the greatest benefits from protective microbes under constant pathogen infection. We hypothesized that variation in pathogen presence over time would limit the evolution of microbe-mediated protection due to the reduced benefits to the host and bacterial symbiont. In our study, enhanced pathogen defence emerged out of host-symbiont coevolutionary interactions only when pathogens were present, independent of the interval or initial presence of the pathogen. Notably, the ultimate strength of microbe-mediated protection that evolved was not impacted by the number of host generations between pathogen infections, the proportion of generations infected, or the presence of the pathogen at the first host-microbe interaction. These results suggest that resident microbes can be a form of transgenerational immunity against rare pathogen infections.

We found that microbe-mediated protection is maintained even in the prolonged absence of pathogen, but that pathogen presence is necessary for microbe-mediated protection to evolve, as previously hypothesized (Clay et al., 2005; King and Bonsall, 2017; Lively et al., 2005). This result is unlike previous work showing that the scale of heterogeneity in abiotic conditions can affect the strength of selection for traits in some symbiotic interactions (Harrison et al., 2013). This discrepancy is potentially due to costs in our symbiotic system being ameliorated (at least in terms of host survival) in well-provisioned hosts, as hosts are provided with food alongside *E. faecalis* and are thus rescued from starvation (also see (Dasgupta et al., 2019)). Although protective symbionts can incur costs (e.g., Vorburger & Gouskov, 2011) for their hosts, with potential for impacts on coevolutionary interactions (King and Bonsall, 2017), it is possible that potential costs of bacterial colonisation might be only detectable when hosts are stressed (Lively, 2006) or that

the costs were not strong enough for us to detect(Little et al., 2002). Different measures of cost

remain to be explored (e.g. lifespan in the complete absence of a protective microbe and a

pathogen). Higher protection also does not always come with higher costs, as found in the black

bean aphid-*Hamiltonella defensa* interaction (Cayetano et al., 2015). Thus, protective traits in an

organism's commensal microbiota could be selected for under pathogen infection and easily

maintained in subsequent uninfected generations.

Microbe-mediated protection was strongest between sympatric pairs when pathogens were present

over evolutionary time, consistent with previous findings (Rafaluk-Mohr et al., 2018). In our study,

protection emerged during coevolution after only 20 host generations, and not due to the

independent evolution of either interacting species, but due to the coevolution of both species

(King and Bonsall, 2017). The time-scale of these interactions is short compared to the longer

shared evolutionary histories shared by other defensive mutualisms (Jousselin et al., 2003; Quek

et al., 2004; Shoemaker et al., 2002). Nevertheless, our findings reveal the potential for microbe-

mediated protection to become enhanced during the formation of a coevolving host-microbiota

relationship.

In conclusion, our results show that enhanced protection in host-microbe interactions can rapidly

evolve and be maintained even under infrequent pathogen infection, suggesting that resident

microbes can be a form of stable, transgenerational immunity. The protective benefit of an

organism's microbiota might remain undetected for several host generations until pathogens re-

emerge. Future research on the failure of pathogens transmit within host populations should

consider the contribution of the protective microbiota to prevent disease spread.

References

Abt, M.C., Artis, D., 2013. The dynamic influence of commensal bacteria on the immune response to pathogens. Curr. Opin. Microbiol. 16, 4–9. https://doi.org/10.1016/j.mib.2012.12.002

Altizer, S., Dobson, A., Hosseini, P., Hudson, P., Pascual, M., 2006. Seasonality and the dynamics of infectious diseases 18.

Barriere, A., 2006. Isolation of C. elegans and related nematodes. WormBook. https://doi.org/10.1895/wormbook.1.115.1

Brenner, S., 1974. The Genetics of Caenorhabditis Elegans. Genetics 77, 71–94.

Cabreiro, F., Gems, D., 2013. Worms need microbes too: microbiota, health and aging in *Caenorhabditis elegans*: The *C. elegans* -microbe holobiont. EMBO Mol. Med. 5, 1300–1310. https://doi.org/10.1002/emmm.201100972

Cayetano, L., Rothacher, L., Simon, J.-C., Vorburger, C., 2015. Cheaper is not always worse: strongly protective isolates of a defensive symbiont are less costly to the aphid host. Proc. R. Soc. B Biol. Sci. 282, 20142333–20142333. https://doi.org/10.1098/rspb.2014.2333

Clay, K., Holah, J., Rudgers, J.A., 2005. Herbivores cause a rapid increase in hereditary symbiosis and alter plant community composition. Proc. Natl. Acad. Sci. 102, 12465–12470. https://doi.org/10.1073/pnas.0503059102

Dasgupta, M., Shashikanth, M., Bojanala, N., Gupta, A., Javed, S., Singh, V., 2019. Nuclear hormone receptor NHR-49 shapes immuno-metabolic response of Caenorhabditis elegans to Enterococcus faecalis infection. (preprint). Immunology. https://doi.org/10.1101/549907

Dey, S., Proulx, S.R., Teotónio, H., 2016. Adaptation to Temporally Fluctuating Environments by the Evolution of Maternal Effects. PLOS Biol. 29.

Dillon, R.J., Vennard, C.T., Charnley, A.K., 2000. Exploitation of gut bacteria in the locust. Nature 403, 851–851. https://doi.org/10.1038/35002669

Dong, Y., Manfredini, F., Dimopoulos, G., 2009. Implication of the Mosquito Midgut Microbiota in the Defense against Malaria Parasites. PLoS Pathog. 5, e1000423. https://doi.org/10.1371/journal.ppat.1000423

Fenton, A., Johnson, K.N., Brownlie, J.C., Hurst, G.D.D., 2011. Solving the Wolbachia Paradox: Modeling the Tripartite Interaction between Host, Wolbachia, and a Natural Enemy. Am. Nat. 178, 333–342. https://doi.org/10.1086/661247

Finkelman, B.S., 2007. Global Patterns in Seasonal Activity of Influenza A/H3N2, A/H1N1, and B from 1997 to 2005: Viral Coexistence and Latitudinal Gradients. PLoS ONE 10.

Ford, S.A., Williams, D., Paterson, S., King, K.C., 2017. Co-evolutionary dynamics between a defensive microbe and a pathogen driven by fluctuating selection. Mol. Ecol. 26, 1778–1789. https://doi.org/10.1111/mec.13906

Friman, V.-P., Laakso, J., 2011. Pulsed-Resource Dynamics Constrain the Evolution of Predator-Prey Interactions. Am. Nat. 177, 334–345. https://doi.org/10.1086/658364

Friman, V.-P., Laakso, J., Koivu-Orava, M., Hiltunen, T., 2011. Pulsed-resource dynamics increase the asymmetry of antagonistic coevolution between a predatory protist and a prey bacterium: Pulsed-resource dynamics and coevolution. J. Evol. Biol. 24, 2563–2573. https://doi.org/10.1111/j.1420-9101.2011.02379.x

Garsin, D.A., Sifri, C.D., Mylonakis, E., Qin, X., Singh, K. V., Murray, B.E., Calderwood, S.B., Ausubel, F.M., 2001. A simple model host for identifying Gram-positive virulence factors. PNAS 98, 10892–10897.

Gravato-Nobre, M.J., Hodgkin, J., 2005. Caenorhabditis elegans as a model for innate immunity
to pathogens. Cell. Microbiol. 7, 741–751. https://doi.org/10.1111/j.1462-
5822.2005.00523.x
Gremillion-Smith, C., Woolf, A., 1988. EPIZOOTIOLOGY OF SKUNK RABIES IN NORTH
AMERICA. J. Wildl. Dis. 24, 620–626. https://doi.org/10.7589/0090-3558-24.4.620
Harrison, E., Laine, A.-L., Hietala, M., Brockhurst, M.A., 2013. Rapidly fluctuating environments
constrain coevolutionary arms races by impeding selective sweeps. Proc. R. Soc. B Biol.
Sci. 280, 20130937–20130937. https://doi.org/10.1098/rspb.2013.0937
Hiltunen, T., Friman, V.-P., Kaitala, V., Mappes, J., Laakso, J., 2012. Predation and resource
fluctuations drive eco-evolutionary dynamics of a bacterial community. Acta Oecologica
38, 77–83. https://doi.org/10.1016/j.actao.2011.09.010
Holden, M.T.G., Feil, E.J., Lindsay, J.A., Peacock, S.J., Day, N.P.J., Enright, M.C., Foster, T.J.,
Moore, C.E., Hurst, L., Atkin, R., Barron, A., Bason, N., Bentley, S.D., Chillingworth, C.,
Chillingworth, T., Churcher, C., Clark, L., Corton, C., Cronin, A., Doggett, J., Dowd, L.,
Feltwell, T., Hance, Z., Harris, B., Hauser, H., Holroyd, S., Jagels, K., James, K.D.,
Lennard, N., Line, A., Mayes, R., Moule, S., Mungall, K., Ormond, D., Quail, M.A.,
Rabbinowitsch, E., Rutherford, K., Sanders, M., Sharp, S., Simmonds, M., Stevens, K.,
Whitehead, S., Barrell, B.G., Spratt, B.G., Parkhill, J., 2004. Complete genomes of two
clinical Staphylococcus aureus strains: Evidence for the rapid evolution of virulence and
drug resistance. Proc. Natl. Acad. Sci. 101, 9786–9791.
https://doi.org/10.1073/pnas.0402521101
Hooper, L.V., Littman, D.R., Macpherson, A.J., 2012. Interactions Between the Microbiota and
the Immune System. Science 336, 1268–1273. https://doi.org/10.1126/science.1223490
Jaenike, J., Unckless, R., Cockburn, S.N., Boelio, L.M., Perlman, S.J., 2010. Spread of a
Drosophila Defensive Symbiont. Science 329, 5.
Jansen, G., Crummenerl, L.L., Gilbert, F., Mohr, T., Pfefferkorn, R., Thänert, R., Rosenstiel, P.,
Schulenburg, H., 2015. Evolutionary Transition from Pathogenicity to Commensalism:
Global Regulator Mutations Mediate Fitness Gains through Virulence Attenuation. Mol.
Biol. Evol. 32, 2883–2896. https://doi.org/10.1093/molbev/msv160
Jansson, H.-B., 1994. Adhesion of Conidia of Drechmeria coniospora to Caenorhabditis elegans
Wild Type and Mutants. J. Nematol. 26, 6.
Jousselin, E., Rasplus, J.-Y., Kjellberg, F., 2003. Convergence and Coevolution in a mutualism:
Evidence from a molecular Phylogeny of Ficus. Evolution 57, 1255.
https://doi.org/10.1554/02-445
Kamada, N., Seo, S.-U., Chen, G.Y., Núñez, G., 2013. Role of the gut microbiota in immunity and
inflammatory disease. Nat. Rev. Immunol. 13, 321–335. https://doi.org/10.1038/nri3430
King, K.C., 2019. Defensive symbionts. Curr. Biol. 29, R78–R80.
https://doi.org/10.1016/j.cub.2018.11.028
King, K.C., Bonsall, M.B., 2017. The evolutionary and coevolutionary consequences of defensive
microbes for host-parasite interactions. BMC Evol. Biol. 17, 190.
https://doi.org/10.1186/s12862-017-1030-z
King, K.C., Brockhurst, M.A., Vasieva, O., Paterson, S., Betts, A., Ford, S.A., Frost, C.L.,
Horsburgh, M.J., Haldenby, S., Hurst, G.D., 2016. Rapid evolution of microbe-mediated
protection against pathogens in a worm host. ISME J. 10, 1915–1924.
https://doi.org/10.1038/ismej.2015.259

King, K.C., Delph, L.F., Jokela, J., Lively, C.M., 2009. The Geographic Mosaic of Sex and the Red Queen. Curr. Biol. 19, 1438–1441. https://doi.org/10.1016/j.cub.2009.06.062

Koch, H., Schmid-Hempel, P., 2011. Socially transmitted gut microbiota protect bumble bees against an intestinal parasite. Proc. Natl. Acad. Sci. 108, 19288–19292. https://doi.org/10.1073/pnas.1110474108

Kommineni, S., Bretl, D.J., Lam, V., Chakraborty, R., Hayward, M., Simpson, P., Cao, Y., Bousounis, P., Kristich, C.J., Salzman, N.H., 2015. Bacteriocin production augments niche competition by enterococci in the mammalian gastrointestinal tract. Nature 526, 719–722. https://doi.org/10.1038/nature15524

Kwiatkowski, M., Vorburger, C., 2012. Modeling the Ecology of Symbiont-Mediated Protection against Parasites. Am. Nat. 179, 595–605. https://doi.org/10.1086/665003

Ley, R.E., Peterson, D.A., Gordon, J.I., 2006. Ecological and Evolutionary Forces Shaping Microbial Diversity in the Human Intestine. Cell 124, 837–848. https://doi.org/10.1016/j.cell.2006.02.017

Little, T.J., Carius, H.-J., Sakwinska, O., Ebert, D., 2002. Competitiveness and life-history characteristics of Daphnia with respect to susceptibility to a bacterial pathogen: Cost of resistance in Daphnia. J. Evol. Biol. 15, 796–802. https://doi.org/10.1046/j.1420-9101.2002.00436.x

Lively, C.M., 2006. The ecology of virulence: Ecology of virulence. Ecol. Lett. 9, 1089–1095. https://doi.org/10.1111/j.1461-0248.2006.00969.x

Lively, C.M., Clay, K., Wade, M.J., Fuqua, C., 2005. Competitive coexistence of vertically and horizontally transmitted parasites. Evol. Ecol. Res.

London, W.P., Yorke, J.A., 1973. Recurrent Outbreaks of Measles, Chickenpox, and Mumps. Am. J. Epidemiol. 98.

Marsh, E.K., May, R.C., 2012. Caenorhabditis elegans, a Model Organism for Investigating Immunity. Appl. Environ. Microbiol. 78, 2075–2081. https://doi.org/10.1128/AEM.07486-11

Martinez, J., Cogni, R., Cao, C., Smith, S., Illingworth, C.J.R., Jiggins, F.M., 2016. Addicted? Reduced host resistance in populations with defensive symbionts. Proc. R. Soc. B Biol. Sci. 283, 20160778. https://doi.org/10.1098/rspb.2016.0778

Martín-Vivaldi, M., Peña, A., Peralta-Sánchez, J.M., Sánchez, L., Ananou, S., Ruiz-Rodríguez, M., Soler, J.J., 2010. Antimicrobial chemicals in hoopoe preen secretions are produced by symbiotic bacteria. Proc. R. Soc. B Biol. Sci. 277, 123–130. https://doi.org/10.1098/rspb.2009.1377

May, G., Nelson, P., 2014. Defensive mutualisms: do microbial interactions within hosts drive the evolution of defensive traits? Funct. Ecol. 28, 356–363. https://doi.org/10.1111/1365-2435.12166

Maynard, C.L., Elson, C.O., Hatton, R.D., Weaver, C.T., 2012. Reciprocal interactions of the intestinal microbiota and immune system. Nature 489, 231–241. https://doi.org/10.1038/nature11551

McFall-Ngai, M., Hadfield, M.G., Bosch, T.C.G., Carey, H.V., Domazet-Lošo, T., Douglas, A.E., Dubilier, N., Eberl, G., Fukami, T., Gilbert, S.F., Hentschel, U., King, N., Kjelleberg, S., Knoll, A.H., Kremer, N., Mazmanian, S.K., Metcalf, J.L., Nealson, K., Pierce, N.E., Rawls, J.F., Reid, A., Ruby, E.G., Rumpho, M., Sanders, J.G., Tautz, D., Wernegreen, J.J., 2013. Animals in a bacterial world, a new imperative for the life sciences. Proc. Natl. Acad. Sci. 110, 3229–3236. https://doi.org/10.1073/pnas.1218525110

Mendes, R., Kruijt, M., de Bruijn, I., Dekkers, E., van der Voort, M., Schneider, J.H.M., Piceno,
Y.M., DeSantis, T.Z., Andersen, G.L., Bakker, P.A.H.M., Raaijmakers, J.M., 2011.
Deciphering the Rhizosphere Microbiome for Disease-Suppressive Bacteria. Science 332,
1097–1100. https://doi.org/10.1126/science.1203980
Moran, N.A., Tran, P., Gerardo, N.M., 2005. Symbiosis and Insect Diversification: an Ancient
Symbiont of Sap-Feeding Insects from the Bacterial Phylum Bacteroidetes. Appl. Environ.
Microbiol. 71, 8802–8810. https://doi.org/10.1128/AEM.71.12.8802-8810.2005
Palmer, T.M., Stanton, M.L., Young, T.P., Goheen, J.R., Pringle, R.M., Karban, R., 2008.
Breakdown of an Ant-Plant Mutualism Follows the Loss of Large Herbivores from an
African Savanna. Science 319, 192–195. https://doi.org/10.1126/science.1151579
Papkou, A., Guzella, T., Yang, W., Koepper, S., Pees, B., Schalkowski, R., Barg, M.-C.,
Rosenstiel, P.C., Teotónio, H., Schulenburg, H., 2019. The genomic basis of Red Queen
dynamics during rapid reciprocal host–pathogen coevolution. Proc. Natl. Acad. Sci. 116,
923–928. https://doi.org/10.1073/pnas.1810402116
Quek, S.-P., Davies, S.J., Itino, T., Pierce, N.E., 2004. Codiversification in an Ant-Plant
Mutualism: Stem Texture and the Evolution of Host Use in Crematogaster (Formicidae:
Myrmicinae) inhabitants of Macaranga (Euphorbiaceae). Evolution 58, 554.
https://doi.org/10.1554/03-361
Rafaluk-Mohr, C., Ashby, B., Dahan, D.A., King, K.C., 2018. Mutual fitness benefits arise during
coevolution in a nematode-defensive microbe model. Evol. Lett. 2, 246–256.
https://doi.org/10.1002/evl3.58
Schulenburg, H., Ewbank, J.J., 2004. Diversity and specificity in the interaction between
Caenorhabditis elegans and the pathogen Serratia marcescens. BMC Evol. Biol. 4, 49.
https://doi.org/10.1186/1471-2148-4-49
Shoemaker, D.D., Machado, C.A., Molbo, D., Werren, J.H., Windsor, D.M., Herre, E.A., 2002.
The distribution of Wolbachia in fig wasps: correlations with host phylogeny, ecology and
population structure. Proc. R. Soc. Lond. B Biol. Sci. 269, 2257–2267.
https://doi.org/10.1098/rspb.2002.2100
Sifri, C.D., Begun, J., Ausubel, F.M., Calderwood, S.B., 2003. Caenorhabditis elegans as a Model
Host for Staphylococcus aureus Pathogenesis. Infect. Immun. 71, 2208–2217.
https://doi.org/10.1128/IAI.71.4.2208-2217.2003
Stiernagle, T., 2006. Maintenance of C. elegans. WormBook.
https://doi.org/10.1895/wormbook.1.101.1
Tan, M.-W., Mahajan-Miklos, S., Ausubel, F.M., 1999. Killing of Caenorhabditis elegans by
Pseudomonas aeruginosa used to model mammalian bacterial pathogenesis. Proc. Natl.
Acad. Sci. 96, 715–720. https://doi.org/10.1073/pnas.96.2.715
Theologidis, I., Chelo, I.M., Goy, C., Teotónio, H., 2014. Reproductive assurance drives
transitions to self-fertilization in experimental Caenorhabditis elegans 21.
Vántus, V.B., Kovács, M., Zsolnai, A., 2014. The rabbit caecal microbiota: development,
composition and its role in the prevention of digestive diseases - a review on recent
literature in the light of molecular genetic methods. Acta Agrar. Kaposváriensis 18, 55–65.
Vorburger, C., Gouskov, A., 2011. Only helpful when required: a longevity cost of harbouring
defensive symbionts: Defensive symbionts reduce host longevity. J. Evol. Biol. 24, 1611–
1617. https://doi.org/10.1111/j.1420-9101.2011.02292.x
474
475

Appendix

Experimental Procedure

Figure A1: The experimental procedure of the evolution experiment in detail.

The procedure of the evolution experiment can be seen in Figure S1 and will be the following:

1. On the first day of each generation, worms were bleached(Stiernagle, 2006). During this step worms were removed from the plates by washing with M9 buffer. Afterwards a 50:50 mixture of 10% NaClO and 5M NaOH was used to bleach all the remaining bacteria in the solution and to release eggs from bleached adult worms. Only eggs survive this step and were then left in the M9 buffer over-night. All bleached eggs hatched overnight, but do not develop any further and thus lead to increased synchronisation for each worm population. Every bleach solution was plated out on a TSA plate to control for carried over contaminations. 100 colonies of *E. faecalis* were picked from the plates that were grown on *E. faecalis* selective medium for 48 hours at 30°C. These colonies were picked into 600μl of THB and then grown up over-night. Simultaneously, a single colony of *Salmonella* food was picked and added to 25ml of LB broth to be grown under shaking conditions over night.
2. On the second day of each generation, the overnight shaken and hatched worms were exposed to 600μl of a 50:50 mixture of *E. faecalis* and food. At this step the population size was adjusted to only contain 1000 individuals. Worms remained on these plates with *E. faecalis* for 48 hours.

3. On the third day of each generation, TSA plates to which worms were going to be exposed on day
4 were inoculated with 100µl of either *S. aureus* or food and were incubated at 30°C overnight.
4. On the fourth day of each generation, worms were washed off the plates seeded with
E. faecalis by filter tip washing. For this purpose, worms were washed off the plates with twice
1.5ml of M9 buffer +1% Triton X100, as previously described(Jansen et al., 2015; Papkou et
al., 2019; Rafaluk-Mohr et al., 2018).This worm and bacteria suspension was spun down for 1
minute, 1 ml of the supernatant was discarded and the rest of the pellet was pipetted on the top of
a filter of a filter tip and spun down for 3 min. Worms were left on the top of the filter were washed
with 400µl of M9 three times, before being re-suspended in 100µl of M9 to bring onto plate. During
this method, most of the externally attached bacteria are washed off the worms to ensure that worm
survival can be attributed to gut colonization of *E. faecalis* and not external attachment of the
protective microbe. *Enterococcus faecalis* remains in the worm's intestine and will establish a
protective effect. After worms were transferred to the plates containing either *S. aureus* or food, all
plates were moved to 25°C.
5. On the fifth day of each generation, worms were washed off the plates again by filter tip washing
(as described for day 4 and previously (Jansen et al., 2015; Papkou et al., 2019; Rafaluk-Mohr
et al., 2018)). Worms were left on plates seeded with food at 20°C for 48 hours to lay eggs. The
amount of transferred bacteria (either *S. aureus* or food can be neglected to have any influence on
the further development of the worms. These plates were then used for bleaching on day 1 of the
following generation. 10% of the worm mixture was separated and used to isolate *E. faecalis*. For
this purpose, the suspension of worms was crushed and then plated on TSA plates with Rifampicin,
as the *E. faecalis* strain carries a Rifampicin resistance. The plated gut content was allowed to grow
at 30°C for 48 hours.

Statistical results:

Table A1: All statistical results summarized, including the statistical test, the specifics associated

with each test, the relevant degrees of freedom and p-values.

Figure	Results
1	Binomial GLM | Comparison	X^2	Degrees of Freedom	p-value | |---|---|---|---| | L1-L4 stage | 10.205 | 1 | 0.001401 | | L4 – Adult | 119.643 | 1 | <2.2 x10^{-16} | | Interaction | 0.053 | 1 | 0.871 |
2A	Binomial GLM Evolution effect: X^2: 42.479, Df=4, p=1.327x10^{-8} Tukey Post-hoc comparisons: | Comparison | Estimate Std. | Error | z-value | p-value |

	Always – Ancestor	-0.5490	0.1354	-4.055	<0.001
	2.1. - Ancestor	-0.5902	0.1348	-4.378	<0.001
	5.1. – Ancestor	-0.8224	0.1448	-5.680	<0.001
	Never – Ancestor	-0.3174	0.1290	-2.460	0.0985

2B Binomial GLM

Evolution effect: X^2: 35.779, Df=5, p=1.051×10^{-6}

Tukey Post-hoc comparisons:

Comparison	Estimate Std.	Error	z-value	p-value
Always – Ancestor	-0.92335	0.19399	-4.760	<0.001
2.1. - Ancestor	-0.84208	0.18860	-4.465	<0.001
5.1. – Ancestor	-0.49494	0.18527	-2.671	0.0791
Never – Ancestor	-0.78893	0.19395	-4.068	<0.001
NPM – Ancestor	-0.52794	0.18847	-2.801	0.0560

2C Binomial GLM

Evolution effect: X^2: 3.2511, Df=5, p= 0.6613

3 Binomial GLM

Evolution effect: X^2: 7.945, Df=3, p= 0.04716

Comparison	Estimate Std.	Error	z value	p-value
2.2. - 2.1.	0.05039	0.10064	0.501	0.9576
5.1. - 2.1.	-0.26298	0.12387	-2.123	0.1414
5.2. - 2.1.	-0.07281	0.12634	-0.576	0.9373
5.1. - 2.2.	-0.31337	0.12399	-2.527	0.0538
5.2. - 2.2.	-0.12320	0.12646	-0.974	0.7585
5.2. - 5.1.	0.19017	0.09467	2.009	0.1795

4A Kaplan Meier Estimates

Comparison	p-value	FDR corrected p-value
Always – Ancestor	0.16086355	0.3217271
2.1. - Ancestor	0.71526638	0.7152664
5.1. – Ancestor	0.04131051	0.1652421
Never – Ancestor	0.04131051	0.7152664

4B	Binomial GLM
	Evolution effect: X^2: 3.8418, Df=4, p= 0.4278