Fine-scale genomic analyses of admixed individuals reveal

unrecognized genetic ancestry components in Argentina

Native American, African and European genetic ancestries in Argentina

Pierre Luisi[1,*], Angelina García[1,2,3], Juan Manuel Berros[4], Josefina M. B. Motti[5], Darío
Demarchi[1,2,3], Emma Alfaro[6,7], Eliana Aquilano[8], Carina Argüelles[9,10], Sergio
Avena[11,12], Graciela Bailliet[8], Julieta Beltramo[8,13], Claudio M. Bravi[8], Mariela Cuello[8],
Cristina Dejean[11,12], José Edgardo Dipierri[7], Laura S. Jurado Medina[8], José Luis
Lanata[14], Marina Muzzio[8], María Laura Parolin[15], Maia Pauro[1,2,3], Paula B. Paz
Sepúlveda[8], Daniela Rodríguez Golpe[8], María Rita Santos[8], Marisol Schwab[8], Natalia
Silvero[8], Jeremias Zubrzycki[16], Virginia Ramallo[17], Hernán Dopazo[18,19,*]
11
12
13 **1.** Universidad Nacional de Córdoba, Facultad de Filosofía y Humanidades,
14 Departamento de Antropología, Argentina
15
16 **2.** Instituto de Antropología de Córdoba (IDACOR), Consejo Nacional de
17 Investigaciones Científicas y Técnicas – Universidad Nacional de Córdoba, Córdoba,
18 Argentina
19
20 **3.** Universidad Nacional de Córdoba, Facultad de Filosofía y Humanidades, Museo de
21 Antropología, Córdoba, Argentina.
22
23 **4.** Consejo Nacional de Investigaciones Científicas y Técnicas – Laboratorio de
24 Análisis de Datos, Biocódices S.A, Buenos Aires, Argentina
25
26 **5.** Núcleo de Estudios Interdisciplinarios de Poblaciones Humanas de Patagonia
27 Austral (NEIPHA), Consejo Nacional de Investigaciones Científicas y Técnicas –
28 Universidad Nacional del Centro de la Provincia de Buenos Aires, Quequén,
29 Argentina.
30
31 **6.** Instituto de Ecorregiones Andinas (INECOA), Consejo Nacional de Investigaciones
32 Científicas y Técnicas – Universidad Nacional de Jujuy, Argentina
33
34 **7.** Instituto de Biología de la Altura, Universidad Nacional de Jujuy, Argentina
35
36 **8.** Instituto Multidisciplinario de Biología Celular (IMBICE), Consejo Nacional de
37 Investigaciones Científicas y Técnicas – Comisión de Investigaciones Científicas –
38 Universidad Nacional de La Plata, Argentina
39
40 **9.** Grupo de Investigación en Genética Aplicada (GIGA), Facultad de Ciencias Exactas,
41 Químicas y Naturales, Departamento de Genética, Instituto de Biología Subtropical

(IBS) - Nodo Posadas, Universidad Nacional de Misiones (UNaM) – Consejo Nacional de Investigaciones Científicas y Técnicas, Posadas, Argentina.

10. Cátedra de Biología Molecular. Carrera de Medicina. Facultad de Ciencias de la Salud. Universidad Católica de las Misiones (UCAMI), Posadas, Argentina.

11. Instituto de Ciencias Antropológicas (ICA), Facultad de Filosofía y Letras, Universidad de Buenos Aires, Argentina

12. Centro de Estudios Biomédicos, Biotecnológicos, Ambientales y Diagnóstico (CEBBAD), Universidad Maimónides, Buenos Aires, Argentina,

13. Laboratorio de Análisis Comparativo de ADN, Corte Suprema de Justicia de la Provincia de Buenos Aires, La Plata, Argentina

14. Instituto de Investigaciones en Diversidad Cultural y Procesos de Cambio (IIDyPCa), Consejo Nacional de Investigaciones Científicas y Técnicas – Universidad Nacional de Río Negro, San Carlos de Bariloche, Argentina

15. Instituto de Diversidad y Evolución Austral (IDEAus), Consejo Nacional de Investigaciones Científicas y Técnicas – Centro Nacional Patagónico, Puerto Madryn, Argentina

16. Laboratorio de Genómica. Biocódices S.A., Buenos Aires, Argentina

17. Instituto Patagónico de Ciencias Sociales y Humanas (IPCSH) – Consejo Nacional de Investigaciones Científicas y Técnicas - Centro Nacional Patagónico, Puerto Madryn, Argentina

18. Departamento de Ecología, Genética y Evolución, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Argentina

19. Consejo Nacional de Investigaciones Científicas y Técnicas – Biocodices S.A., Buenos Aires, Argentina

* Corresponding Authors:
Emails: pierrespc@gmail.com (PL), dopazoh@gmail.com (HD)

Abstract

Similarly to other populations across the Americas, Argentinean populations trace back their genetic ancestry into African, European and Native American ancestors, reflecting a complex demographic history with multiple migration and admixture events in pre- and post-colonial times. However, little is known about the sub-continental origins of these three main ancestries. We present new high-throughput genotyping data for 87 admixed individuals across Argentina. This data was combined to previously published data for admixed individuals in the region and then compared to different reference panels specifically built to perform population structure analyses at a sub-continental level. Concerning the Native American ancestry, we could identify four Native American components segregating in modern Argentinean populations. Three of them are also found in modern South American populations and are specifically represented in Central Andes, Central Chile/Patagonia, and Subtropical and Tropical Forests geographic areas. The fourth component might be specific to the Central Western region of Argentina, and it is not well represented in any genomic data from the literature. As for the European and African ancestries, we confirmed previous results about origins from Southern Europe, Western and Central Western Africa, and we provide evidences for the presence of Northern European and Eastern African ancestries.

Author Summary

The human genetic diversity in Argentina reflects demographic processes during which the European colonists invaded a territory where Native American populations were settled. During the colonial period, the slave trade also prompted many African people to move to Argentina. Little is known about the origins of these three continental ancestry components in Argentinean populations nowadays. Genotyping data for 87 admixed individuals throughout Argentina was generated and data from the literature was re-analyzed to shed light on this question. We confirmed that most of the European genetic ancestry comes from the South, although several individuals are related to Northern Europeans. We confirmed that the African origins in Argentina mainly trace back from Western and Central/Western regions, and we document some proportion of Eastern African origins poorly described before. As for the Native American ancestry, we identified that it can be divided into four main components that correspond to Central Chile/Patagonia, Subtropical and Tropical Forests, Central Andes and Central Western region of Argentina. In order to understand the specificity of the genetic diversity in Argentina, more effort is required to generate specific massive genomic knowledge at the local level.

Introduction

The first systematic investigation of human genetic variation in Argentina focused on a limited number of markers either uniparental (mtDNA, Y-STRs, Y-SNP; [1–10]

or autosomal (Short Tandem Repeats, Ancestry Informative Markers, Alu sequences,

indels, and blood groups [11–16]). Studies based on autosomal markers identified

an important inter-individual heterogeneity for the African, Native American and

European genetic ancestry proportions [11–16]. Accordingly, most of the studies

based on uniparental markers showed large differences in the genetic composition

of Argentinean populations, accounting for the different demographic histories

within the country [1,2,17–20,3–9,16]. Although the idea of a 'white' country with

most of inhabitant's descendants from European immigrants has now been rejected

by these studies, the Argentine founding myth of a white and European nation

remains perceptible today [21].

There is a wealth of information about the European ancestors, both from familial

stories and from historical records. On the contrary, little is known about the African

and Native American populations that contributed to the admixture events between

the continental components. The great wave of European immigration occurred

between the mid-19th to mid-20th centuries. Although immigrants came from all

over the continent, historical records attest that immigration waves from Southern

Europe (mainly Italy and Spain) were predominant [22].

As for the African genetic origins in Argentina, historical records about the arrival of

African slaves to to Río de la Plata show that Luanda, in current Angola territory in

Central Western region of Africa, was the main departure harbor, followed by ports

located in the Gulf of Guinean and on the coast of the present-day Senegalese,

Gambian and Sierra Leone territories. The Indian Ocean coasts, in the current

territory of Mozambique, were also an important departure location [23–25].

Although the departure harbor is a poor proxy to infer the actual slaves' origins

[25,26], the African uniparental lineages observed in Argentina are consistent with

the historical records [27,28].

As for the Native American component, it is difficult to study its origin focusing on

present-day communities since their organization has changed drastically after the

arrival of the first conquerors in the 16th century [21]. During the period of

conquest and colonization, wars, diseases and forced labor decimated the Native

populations. The system of colonial exploitation also often meant the relocation of

individuals, families, and communities [29]. Then, the expansion of the nation-state

by the late 19th Century can be described as a territorial annexation process and

subjugation of the indigenous peoples perpetrated by the Argentinean national

armed forces between 1876 and 1917 [30].

Due to the specificity of the Argentinean demographic history, a remaining challenge

is to unravel which populations from each continent contributed to the genetic pool

in nowadays Argentinean populations leveraging genotype data for hundreds of

thousands of autosomal markers from the whole genome. Recently, two articles

presented high-throughput genotyping data for modern Argentine individuals, and

provided the first insights to decipher which populations from each continent

contributed to the genetic pool in nowadays Argentinean populations [31,32]. In

both studies, it was found that the European ancestry in Argentina is mainly

explained by Italian and Iberian ancestry components. Homburger *et al.* observed a

strong gradient in Native American ancestry of South American Latinos between

Andean and other South American Native American populations [31]. In addition,

Muzzio *et al.* found that African ancestry is explained by a Central Western (Bantu-

influenced) component and a Western component.

Despite these progresses, the efforts to understand the demographic history that

shaped genetic diversity in Argentina have inevitably been scarce, particularly for

African and Native American components. Studies of a wider region, namely South

America can provide important insights. A previous study showed that Western and

Central Western African ancestries are common across the Americas, particularly in

Northern latitudes, while the influence of South/Eastern African ancestry is greater

in South America [33]. In another study in Brazil, two African ancestries have been

observed: a Western African one and another associated with Central East African

and Bantu populations, the latter being more present in the Southeastern and

Southern regions [34]. Ancient DNA analyses of the peopling of the Americas

suggest that Native American populations of South America descend from two

streams from Northern America: one mainly present in the Andes and another one

present elsewhere [35]. These streams have replaced the first people settled in the

region, whose ancestry was related to the Clovis culture [35]. Genetic continuity for

the Native American component appears to have prevailed in the region ever since

these replacements [35–37]. However, little is known about the legacy in modern

populations of the different ancestries associated to the several waves of population

arrivals in South America. Modern Native American populations in the Southern

Cone of South America seems to be divided between a component that includes

populations in Tropical and Subtropical Forests and another component that

includes Andean, Central Chile and Patagonian populations [38]. A recent genomic

study of ancient and modern populations from Central Chile and Western Patagonia

further identified that they are differentiated from the Andean and Subtropical and

Tropical Forests populations [37]. In another genomic study of modern samples (in

which the Southern Cone is only represented by the Gran Chaco region), it has been

found that all non-Andean South American populations are likely to share a common

lineage, while they are unlikely to share with the Andeans any common ancestor

from Central America [39], supporting the hypothesis of many back migrations to

Central America from non-Andean South American populations [39]. It has also

been described that the genetic interactions between the peopling routes on both

sides of the Andes were limited [39,40].

Here, we carry out a fine-scale population genomic study to get insight into the

genetic structure and the complex origins of the African, European and Native

American ancestries in Argentina populations. Using Affymetrix, Axiom-Lat1 array

technology, we genotyped 87 admixed individuals nationwide for more than

800,000 SNPs covering coding and non-coding regions of the genome. Additionally,

a dataset with ~500 individuals from modern and ancient populations throughout

South America was constructed. We also pulled together genotype data for Sub-

Saharan African and European populations from the literature. The new data

generated in the present study, compared to those data sets, allows to broaden the

knowledge of the sub-continental origins of the three main genetic ancestries of

Argentinean populations.

Results and Discussion

Studied Populations

High-throughput genotyping data was generated for 87 admixed individuals

throughout Argentina (**Fig 1** and **S1 Table**). For clarity in the visualization of the

results, we used provinces and regions to classify the admixed individuals analyzed

as shown in **Fig 1**. This classification has not been used for any statistical analyses.

The generated data was compared to different data sets to understand the origins of

the genetic diversity in the country. These data sets are called **DS<n>** and are

described in **S2 Table**.

Fig 1. Sample locations from the present study.
The samples are divided according to regions for visualization only. N: The number
of samples that passed genotyping Quality Controls.

Ancestry in a worldwide context

In order to characterize the genetic diversity observed in Argentina within a

worldwide genetic context, Principal Component Analysis (PCA [41]; **S1 Fig**) was

applied to the dataset **DS1,** which contains genotype data for the samples from

Argentina generated in the present study, as well as other admixed and Native

American individuals from South America [31,37,38,42] and individuals from

Europe and Africa from the 1000 Genomes Project (1KGP; [42]). The PCA shows that

Argentinean individuals have different proportions of European, Native American

and, much less represented, African ancestry (**S1A Fig**). This pattern, which has

already been documented in other admixed populations from South America

[31,42], including in Argentina [13–16,32] supports the heterogeneous genetic

origins throughout the country. In addition, PC4 discriminates between Southern

and Northern European individuals (**S1B Fig**) while PC6 separates Luhya

population, a Bantu-speaking population in Kenya, from Western African

populations (**S1C Fig**). The Argentinean samples do not exhibit any gradient along

these two PCs. But, PC3 and PC5 discriminate three different main Native American

ancestries that are also represented in the Argentinean samples.

We ran unsupervised clustering models with Admixture software [43] on **DS1**. We

used from 3 to 12 putative ancestral populations. Comparing the cross-validation

scores obtained for each run, we estimated that the genotype data analyzed was

best explained with a model with K=8 ancestral populations (**S2A Fig**). At K=3 (**S2B**

Fig and **S3 Table**), the algorithm allows estimating the proportions of European,

Native American and African ancestry. For K=4 to K=7, (**S2C-F Figs**), the model

detects sub-continental ancestries. At $K=8$ (**Fig 2; S3 Table**), the European ancestry

is divided into Northern and Southern components (dark and light blue,

respectively), while African ancestry is composed of Westernmost African (green),

Gulf of Guinea (light green) and Bantu-influenced (dark green) components.

Moreover, three components are observed for Native American ancestry: one in

Central Chile/Patagonia populations (in orange; hereafter referred to as **CCP**),

another in Central Andes populations (in yellow; hereafter referred to as **CAN**) and a

third one in populations from Tropical and Subtropical Forests (in pink; here after

referred to as **STF**).

| **Fig 2. Admixture Analyses in a Worldwide Context.**
| Admixture for *K=8.* . **1st row**: 1000 Genomes Samples from Europe, Africa and South
| America [42]. **2nd row**: Modern Native American samples grouped following [38].
| **3rd row**: Chilean, Peruvian and Argentinean admixed samples from [31]. **4th row:**
| Argentinean admixed samples from the present study. Samples are grouped
| according to regions (**Fig 1**); CYA: Cuyo Region in Argentina; NEA: Northeastern
| Region, NWA: Northwestern Region; PPA: Pampean Region; PTA: Patagonia Region.
| CLM: Colombians from Medellin; PEL: Peruvians from Lima.

We confirmed that the estimates of the continental ancestry proportion obtained

from a model with $K=3$ and $K=8$ are highly consistent (**S3 Fig**). Moreover, the

eigenvectors from PCA and the ancestry proportion estimates with Admixture are

well correlated (**S4 Fig**).

From the Admixture results for $K=8$ (**Fig 2**), we observed that the European

ancestry for Argentinean samples, is divided in Southern and Northern components,

the former being the most abundant. The low proportion of African ancestry in

Argentinean samples makes difficult to interpret its sub-continental origins from

analyses within a global context. Surprisingly, all three components of Native

American ancestry are present in most Argentinean samples (**Fig 2**). They exhibit

mid proportions of different Native ancestries suggesting either the result of a

mixture between these three ancestry components or an underrepresentation of

Native American in the reference panels currently available. Such mixture pattern is

not observed in other South American countries. Indeed, the Native American

ancestry for Peruvian, Chilean and Colombian admixed samples is mainly

represented by CAN, CCP and STF, respectively. This is consistent with the

geographical area where the admixed individuals have been sampled, and the

genetic ancestry of the indigenous communities from each country.

Sub-continental ancestry components in Argentina

To decipher the sub-continental origins of the European, African and Native

American ancestries, we first estimated local ancestry patterns in phased **DS2** and

DS3 (**DS2p** and **DS3p**) separately. Across phased chromosomes, we assigned

whether a genomic region was of Native American, European or African ancestry

using RFMix v2 [44]. We applied principal component and unsupervised clustering

analyses on the masked data (**S5 Fig**) compiled with reference data sets describing

the genetic diversity within each continent (**S2 Table**).

The genetic legacy of European migration in Argentina

We used **DS4**, a combination of the masked genotype data for admixed individuals.

with a set of European individuals carefully selected to be representative of the

genetic diversity in their sampling area [45,46].

From the PCA, we observed that most Argentinean individuals cluster with Iberians

and Italians (**Fig 3**), as previously described [31,32]. However, some individuals

cluster with Central and Northern Europeans.

Fig 3. European Ancestry Specific Principal Component Analysis.
(A) Multidimensional scaling scatterplot (MDS) from Euclidian distances calculated from weighted 5-first European-specific Principal Components performed using the European reference samples, and admixed samples masked for African ancestry. Individuals from the European reference panel are colored according to main European geographic regions as shown in **(B)** while South American individuals are represented as shown in the legend. Elbow method to choose the best number of PCs to compute MDS is shown in **S6 Fig**.

We estimated that the genotype data analyzed was best explained with an

admixture model with *K*=2 (**S7A Fig)**. Although it is rather difficult to assign a clear

label to each of these two ancestral populations, it seems that the algorithm

discriminates between Central/Northern (dark blue) and Southern Europe (light

blue) components (**S7B Fig**). For *K*=3, we still observe a Central/Northern European

component (dark blue), while the South can be divided into a component

particularly represented in Iberian individuals (turquoise) and another component

more frequent in Southeastern Europeans and Italians (light blue). The Iberian

component is highly represented in Southern American samples (**S7C Fig**), most

likely reflecting the legacy of first Iberian migrations in South America during

colonial period [11]. Samples from Argentina and Chile exhibit higher proportions of

Southeastern/Italian and Northern European ancestries than Colombians, as well as

lower Iberian ancestry proportions (**S8 Fig**). We observed no significant difference

in the proportions of any European ancestry between Argentinean and Chilean

samples (**S8 Fig**). However, both PCA and Admixture shows that the individuals

with most Southeastern/Italian ancestry are from Argentina, This is consistent with

a previous study [31], and it can be explained by the many arrivals from Italy during

the great wave of European immigration in the 19th and 20th centuries [22].

Moreover, Argentinean individuals with higher proportion of Central/Northern

European ancestry are from Misiones province (NEA), consistent with the historical

record of settlement of Polish, German, Danish and Swedish colonies promoted by

governmental or private enterprises in the province [47].

Different African ancestry components in Argentina

To investigate the sub-continental components that explains African ancestry in Argentina, we used **DS5** which combines masked genotype data from admixed individuals with a published data set of African individuals [42,48–50] (**S9A Fig**). The African reference populations used here can be divided into five main groups: Bantu-influenced, Hunter-Gatherers, Western African, San and Eastern African populations [48].

From the PCA applied to this data, we observed that African ancestry in Argentina has small genetic affinity with San and Hunter-Gatherers populations. Indeed, according to PC1 and PC2, Argentinean individuals are located within a group of African individuals belonging to Bantu-influenced, Western African and Eastern African populations, and clearly distinct from San and Hunter-Gatherer populations (**S9B Fig**). Moreover, on PC3, most Argentinean are closer to Western African and Bantu-influenced populations, while several Argentinean individuals, particularly from Northern Argentina, are located within a gradient towards Eastern African populations (**S9D Fig**). According to PC4, most of the Argentinean individuals are closer to the Bantu-influenced populations while others individuals are closer to Western Africa populations, and several other Argentinean individuals are found in a cline between these two groups (**S9C Fig**). These patterns were confirmed by applying Admixture algorithm. The Cross-Validation procedure points to a best fit of the data with K=5 (**S10A Fig**), a model for which we observe that he Bantu-influenced and Western Africa components are the most represented in Argentinean individuals. Moreover, this analysis also showed that some individuals exhibit

smaller, yet important, proportions of Eastern African ancestry, particularly in

Northern Argentina (**Fig 4**). Although, the important missing genotype rate in

masked data for admixed individuals could bias PCA and Admixture results, the

results obtained by both methods are highly consistent for admixed individuals

(**S11 Fig**).

Fig 4. African Ancestry-Specific Admixture Analysis.
Admixture for *K=5*. **1st row**: 1,685 reference samples with >99% of African ancestry.
Populations are grouped following [48]. **2nd and 3rd rows**: Peruvian and
Argentinean admixed samples from [31] and from the present study. Samples are
grouped according to region. CYA: Cuyo Region; NEA: Northeastern Region, NWA:
Northwestern Region; PPA: Pampean Region; PTA: Patagonia Region. Genotype data
for admixed samples were masked for African ancestry.

The legacy of Western Africa on the African genetic diversity in the Americas has

been preeminent [51–53], along with an impact of Bantu-influenced populations

from Central Western Africa, particularly in Brazil and the Caribbean [51,52]. These

two African ancestries have also been previously documented in Argentina from

studies of autosomal [32] and maternal markers [27,28]. Maternal lineages specific

to populations from Central Western Africa –particularly from Angola- are the most

common African lineages in Argentina according to studies in the Central region

[27] and in four urban centers (Puerto Madryn, Rosario, Resistencia and Salta) [28].

These results are concordant with the predominance of the Bantu-influenced origin

identified in the present study. In addition, the presence of Southeastern Africa

maternal lineages in Argentina [27,28] is consistent with African ancestry of this

origin identified in previous studies in other South America countries [33,34], and

with the Eastern African ancestry identified here.

An unrecognized genetic Native American component in Argentina

We compared masked genotype data for admixed individuals with masked genotype

data of modern Native American reference populations from South America [37,38]

and a set of ancient DNA data from the region [35–37,54,55] (**DS6**; **Fig 5A**). The PCA

(**Fig 5B**) and Admixture analyses (best model with *K=3*) confirmed the three main

South Native American ancestry components described by de la Fuente *et al.* (2018),

and identified in our analyses at the global level.

Fig 5. Native American ancestry-specific principal component analysis.
(A) Localization map of the studied samples. Color and point coding matches sample
groups: ancient populations are grouped following [35], modern Native American
populations are grouped following [38] while Argentinean admixed sample
locations are grouped according to Regions (**Fig 1**). For clarity, some geographic
coordinates have been slightly changed. **(B)** Multidimensional scaling scatterplot
from Euclidian distance calculated from weighted Native American ancestry specific
Principal Components. Elbow method to choose the best number of PCs to compute
MDS is shown in **S12 Fig.**

Similar results were also observed when applying Admixture algorithm (best model

with *K=3*; **S13A Fig**). The proportion estimates for the three main Native American

ancestries observed for the South America Native American individuals and for the

ancient samples (**Fig 6**) highly correlate with estimates from unmasked data (**S14**

Fig), and are consistent with CAN, CCP and STF labels that we attributed. In

Argentina, the three Native American ancestries are observed in almost all the

admixed individuals studied. In Northeastern Argentina (NEA), STF ancestry is the

most frequent. In Northwestern Argentina (NWA), CAN ancestry is not clearly

dominating since STF is also observed in important proportions. In the South, CCP is

observed in greater proportions.

Fig 6. Native American Ancestry-Specific Admixture Analysis.
Admixture for *K=3.* . **1st row**: Ancient samples grouped following [35]. **2nd row**:
Modern Native American samples grouped following [38]. **3rd row**: Chilean,
Peruvian and Argentinean admixed samples from [31]. **4th row:** Argentinean
admixed samples from the present study; sample locations are grouped according to
Regions (**Fig 1**); CYA: Cuyo Region; NEA: Northeastern Region, NWA: Northwestern
Region; PPA: Pampean Region; PTA: Patagonia Region. Genotype data for all modern
samples were masked for Native American ancestry..

Although migration events tend to reduce the genetic distance among these

components, we still identified correlations between geographic coordinates and

the proportions of each Native American ancestry component. Indeed, CAN is found

in higher frequencies further North (**S15A Fig**) and West (**S15A Fig**), STF is higher

further North (**S15B Fig**) and East (**S15E Fig**), while the proportion of CCP is higher

further South (**S15C Fig**).

Many individuals from the Cuyo and Pampean regions of Argentina (San Juan and

Córdoba provinces as well as South of Buenos Aires province) exhibit intermediate

position in PCA (**Fig 5**) and mid proportion estimates with Admixture (**Fig 6**). This

pattern can be interpreted as the result of a mixture between different ancestries

(scenario 1) or relative limited shared history with any of them (scenario 2). In the

intent to discriminate between both scenarios, we increased the number of

ancestral populations to 4 and 5. Model with *K=4* (**S13C Fig**) does not help to

address this question, and model with *K*=5 informs that scenario 2 is the most likely

for Middle Holocene samples from the Southern Cone (**S13D Fig**). Models including

more ancestral populations (K>5) did not allow robust ancestry proportion

estimates most likely due to the masked data leveraged here. Many regions of

Argentina are underrepresented in the reference panel because of the scarcity of

Native American communities in most of the Argentinean territory. Altogether, PCA

and admixture analyses are not sufficient to unequivocally contrast whether some

Native American genetic diversity specific to Argentina is not represented in the

reference panel.

In order to circumvent these limitations, we performed an objective quantitative

approach based on *K*-means clustering (**S16 Fig**). We assigned the 452 modern

South American individuals (the 53 ancient samples included in **DS6** were not

analyzed here) to a Native American ancestry cluster. We assigned 163, 161 and 70

individuals to the clusters representing CAN, STF and CCP, respectively, and 32

individuals were assigned to a fourth cluster (**S4 Table**). The remaining 26

individuals were removed for further analyses because their group assignation was

not consistent across the three clustering approaches. We acknowledge that these

groups are culturally, ethnically and linguistically heterogeneous. However, we

argue that analyzing such groups built from genetic similarities may provide

interesting insights into evolutionary mechanisms that shaped the Native American

ancestry in South America.

Most individuals from Calingasta, (located in the Northwest Monte and Thistle of the

Prepuna ecoregion; San Juan Province) and from Santiago de Chile were assigned to

the fourth group. The genealogical record for the Calingasta individuals attests to a

local origin of their direct ancestors up to two generations ago, and they have

mtDNA sub-haplogroups predominant in the Cuyo region (**S1 Table**; [56,57]). On

the other hand, all the Huilliche and Pehuenche individuals from Central Chile [37]

have been consistently assigned to CCP. Altogether, we decided to refer to this

fourth group hereafter as Central Western Argentina hereafter (CWA).

Relationship among the four identified Native American
451
components
452
Significant positive f_3 scores to test for Treeness were obtained for the six possible

pairwise comparisons among the four Native American clusters identified (**Fig 7A**

and **S5A Table**). In addition to confirming the differentiation among the three

components described in our reference panel (CCP, CAN and STF), this analysis

showed that the fourth group (CWA) most likely represents a Native American

component never described in any previous study based on autosomal genetic

markers. In fact, the genetic differentiation (measured with F_{ST} index) between CWA

and any other cluster is not lower than for other comparisons (**Fig 7B**). The lowest

F_{ST} was observed between STF and CAN, probably due to the fact that STF

encompasses the Northern Andes region (**Fig 7B**). Moreover, the distribution of 1 –

$f_3(YRI; Ind1, Ind2)$ between pairs of individuals from different groups (**S17 Fig**) is an

additional argument to discard a scenario of mixture (mentioned before as scenario

1).

Furthermore, f_4 analyses showed that (i) CAN has no particular genetic affinity with

any component relative to the others; (ii) STF is closer to CAN as compared to CCP

and CWA; and (iii) CWA and CCP exhibit higher genetic affinity between them than

with CAN or STF (**Fig 7C**; **S5B Table**). However, a neighbor-joining analysis [58]

from distances of the form 1/$f_3(YRI; X, Y)$ suggests that CAN is more closely related

to CCP and CWA than to STF (**Fig 7D**; **S5C Table**).

Fig 7. Relationship among the four Native American groups identified.
(A) f_3(*Target; S1, S2*) to test for treeness. **(B)** F_{ST} between pairs of Native American
groups. **(C)** f_4*(YRI, Target; S1, S2)* to test whether Target shares more ancestry with
S1 or S2. Since f_4 is symmetrical when switching *S1* and *S2,* only positive
comparisons are shown. **(D)** Neighbor-joining tree estimated from distance matrix
of the form 1/ f_3(YRI; X, Y). The tree was estimated using ancient sample from
Upward Sun River site in Beringia (USR1) as outgroup. CAN: Central Andes; STF:
Subtropical and Tropical Forests; CCP: Central Chile / Patagonia; CWA: Central
Western Argentina; YRI: Yoruba from 1KGP.
Vertical segments are the +/- 3 standard errors intervals in A and B.

Based on these evidences, we argue that CWA   may represent a Native American

ancestry that diverged from CCP and established on the Eastern side of the Andes in

the Cuyo region.

The existence of a specific differentiated component in the Central Western and

Central regions of Argentina has been previously suggested from maternal lineages

analyses accounting for the genetic relationship between these two regions [10] and

the presence of specific clades underrepresented elsewhere [10,27,59]. These

studies support the hypothesis of a common origin and/or important gene flow [57],

and the authors referred to a meta-population with great temporal depth and

differentiated from other regions in Argentina. Moreover, the ethnographic

description of the populations that were settled at the moment of contact with the

Spanish colonies accounts to a potential relationship between Huarpes in the Cuyo

region and Comechingones in present-day Córdoba province ([60], cited by [61]).

Altogether, these facts support the Central Western Argentina (**CWA**) label that we

attributed to this fourth component.

Complementary historical facts may explain the representation of this component in

Santiago de Chile and its absence in Huilliche and Pehuenche populations. First, in

pre-Columbian times, it seems to have existed an ethnic boundary around the 36° S

Latitude, since Central Andean civilizations could not expand further South. [62,63].

Later, the Spanish could neither settle those territories. Second, the Spanish

colonists that established in Santiago de Chile organized massive deportation of

Huarpes individuals from Cuyo region to palliate the lack of indigenous workforce

[64]. For example, in Santiago de Chile, in 1614, 37% of the indigenous people that

lived in the suburbs were Huarpes according to the chronicler Vázquez de Espinosa

[65].

The present study, in which we analyzed individuals that do not belong to any

indigenous community, made possible the identification of a Native American

component not previously reported from autosomal markers.

Genetic affinity of the four Native American ancestry components with ancient populations

We evaluated the genetic affinity of the four Native American ancestry components

to ancient samples from the literature [35–37,54,55,66,67]. Graphical summaries of

pairwise f_3-based distances are presented in **S18 Fig**.

When comparing the genetic affinity of a given component with the different ancient

groups using either the f_3-*outgroup* or the f_4 statistics (**S19F Fig** and **S5D,E Tables**),

we identified that CAN tends to exhibit greater genetic affinity with ancient Andean

populations than with other ancient groups (**S19A,E Figs**). Strikingly, the genetic

affinity of this component with both Late Andes and Early Andean ancient groups

would point to a genetic continuity across the whole temporal transect that the

archaeological samples provide. This would suppose that the replacement of an

early population arrival by a later stream of gene flow in Central Andes identified

previously [35] does not fully explain the current-day genetic diversity for CAN. We

observed higher genetic affinity with ancient Southern Cone groups for both CCP

and CWA (**S19C,D,G,H Figs**). As for STF, we observed intermediate genetic affinity

with ancient samples from both the Andes and the Southern Cone (**S19B,F Figs**).

The fact that there is no ancient sample group exhibiting outstanding genetic affinity

with STF points to the underrepresentation of this component in ancient samples.

First, the geographical range covered by ancient samples that could represent this

component is restricted to Brazil, while STF is a heterogeneous group that includes

relatively isolated populations [39] from the Gran Chaco, the Amazonas and

Northern Andes. Moreover, the most recent samples that could represent STF are

aged ~6700BP, and gene flow with other components since then may have

contributed to dissolve the genetic affinity of STF with ancient samples in Brazil

analyzed here.

Then, we evaluated the relationship of the time depth of the ancient samples from

either the Andes or the Southern Cone, with their genetic affinity to the modern

components of Native American ancestry (**Fig 8**). We observed a statistically

significant relationship between the age of the ancient Southern Cone samples and

their genetic affinity with CCP and CWA. This means that the older the ancient

sample from the Southern Cone, the lower the shared drift with CCP and CWA. On

the other hand, no statistically significant relationship was identified for STF and

CAN (P = 0.523 and P=0.596, respectively; **Fig 8A**). These patterns could be due to a

relationship between geography and the age of the ancient samples because the

most recent samples are concentrated in the Southern tip of the subcontinent (**Fig**

5A). Moreover, the number of SNPs with genotype data tends to decrease with the

age of the ancient samples due to DNA damage, and thus inducing a potential bias

towards significant positive correlations. To simultaneity correct both these two

putative confounding effects, we repeated the analyses but using a correction for the

ancient sample age (the residuals of the linear regression between the age of the

ancient samples and their geographic coordinates) and a correction for genetic

affinity estimates (the residuals of the linear regression between f_3 and the number

of SNPs to estimate it). This correction intensified the relationship described for CCP

and CWA (**Fig 8C**). It also allowed to actually identifying significant relationships for

STF and CAN. On the other hand, CAN is the only modern Native American

component that exhibits a significant relationship between its genetic affinity with

ancient Andean samples and their age (**Fig 8B**). This pattern holds after correction

for geography (**Fig 8D**). Repeating the same analyses using f_4 statistics, we reached

the same conclusions (**S20 Fig**).

Fig 8. Changes across time of genetic affinity of the four Native American with
ancient samples.
Each point represents a f_3-Outgroup score of the form $f_3(YRI; X, Ancient)$ vs the age of
ancient samples, where *X* is one of the four identified modern Native American
components, and *Ancient* is an ancient group. *X* is represented by the color of the
square while the symbol inside the square represents *Ancient*. The legend for
plotted symbols is shown in Fig 5**. (A)** f_3 vs age of ancient samples from Southern
Cone. **(B)** f_3 vs age of ancient samples from the Andes. **(C)** f_3 vs age of ancient
samples from Southern Cone considering correction for both f_3 and age. **(D)** f_3 vs age
of ancient samples from the Andes considering correction for both f_3 and age
correction. Linear regression slopes and the associated *P*-values are shown. CAN:
Central Andes; STF: Subtropical and Tropical Forests; CCP: Central Chile /
Patagonia; CWA: Central Western Argentina; YRI: Yoruba from 1KGP

Early divergence among the four Native American components in Argentina

Using another setting of the f_4 statistics (**S5F Table**; **S21 Fig**), we observed that all Southern Cone ancient samples – except Los Rieles ~10900 BP – are more similar to CCP than to CWA (**S21F Fig**), pointing to divergence time between those two components older than ~7700 BP. CWA is not closer to any Brazilian and Andean ancient sample as compared to CCP, reinforcing the idea that CWA is not a mix of CCP, CAN and STF related ancestries. Moreover, we observed that all the ancient groups of the Southern Cone have greater genetic affinity with CCP than with CAN and STF (**S21B,D Figs**). Ancient groups of the Southern Cone (with the exception of Los Rieles ~10900 BP) tend to exhibit higher genetic affinity with CWA than with STF (**S21C Fig**). In addition, the comparisons including STF and CWA also point to a closer genetic relation between ancient groups from the Southern Cone with CWA, although statistical significance is reached only for Late Holocene Patagonia samples.

All the ancient samples from the Andes, independently of their previous assignment to the Early or Late population stream [35], are genetically more similar to CAN than to any of the three other components (**S21A,D,E Figs**). These results support further the hypothesis that the ancestry associated with an early Andean population arrival remains present to some extent in modern Central Andean populations and has not been totally replaced by a later stream of gene flow. Moreover, the archaeological samples in Brazil have higher genetic affinity with CAN than with CWA and CCP (not statistically significant for the latter comparisons). This suggests that divergence of

STF with CCP and CWA could have occurred before the divergence of STF with CAN

(also observed in **S18A Fig**).

Altogether, the trends depicted by these analyzes are expected under a model of

early divergence among the four components. Our results also support the

hypothesis of genetic continuity over long period of time for the different

components after their split.

In order to get insights into the past genetic influence among the four components

since their divergence, we applied a last f_4-statistics analysis (**S5G Table**; **S22 Fig**).

We observed that the influence of CAN appears to have been more pronounced into

STF component than into CWA and CCP components (**S22B,G,I Figs**). The genetic

affinity of CAN with STF, CCP and CWA is reduced as compared to both ancient Early

and Late Andes samples (**S22A,H,J Figs**), implying that that the genetic legacy of

Andean ancient populations into CAN remains important nowadays. The mutual

influence of CCP and CWA components seems to have been important in remote

times (**S22K,L Figs**). However, the available ancient samples provide much more

precise information on the influence of CCP into CWA component than the other

way around. Indeed, CWA does not exhibit higher genetic affinity to any ancient

group relative to CCP (**S22K Fig**). The archaeological record for which genetic data

has been generated misrepresents CWA since its early divergence with CCP, as well

as the common ancestors specific to these two components.

Conclusions

European ancestry in modern Argentinean populations is mostly explained by
Southern European origins, although we identified several individuals with
Northern European ancestry. African ancestry in Argentina traces its origins from
three main components. Although the Bantu-influenced and Western components
are clearly the most represented, we also found that an Eastern origin explains some
of the African ancestry in Argentina and represents up to ~30% of the African
ancestry for some Argentinean individuals. Our work shows that studying more
admixed individuals, with a particular focus on extending the geographical coverage
of the Argentinean territory, would help to identify the genetic legacy from
secondary migration streams, and thus to get a better representation of the complex
origins of African and European ancestries in the country.

Concerning Native American ancestry, we concluded that Argentinian populations
share, in varying proportions, from three distinct components previously described:
Central Andes, Central Chile/Patagonia and Subtropical and Tropical Forests.
Moreover, we present evidence supporting the hypothesis that the Native American
ancestry in the Central Western region of Argentina may derive from a fourth
component that diverged early from the other Native American components, and
maintained a tight link with the Central Chile/Patagonia component. This relation is
not explained by a putative contribution of admixed individuals from Santiago de
Chile (**S23 Fig**). Having identified this component from admixed individuals
demonstrates that focusing only on indigenous communities is insufficient, at least

in Argentina, to fully characterize the Native American genetic diversity and

decipher the pre-Columbian history of Native Americans. Indeed, most indigenous

communities have been culturally annihilated and invisibilized [68,69] to the point

that several Argentinean regions were considered "Indian free" in the mid-20[st]

century [70]. However, the cultural incorporation did not necessarily imply a

biological extinction. Although studies based on samples from indigenous

communities [37–39] provide decisive information to understand the evolutionary

history of Native American ancestry, alternative strategies must be considered to fill

this gap in the effort to more fully describe the Native American ancestry (e.g. see

[71]). Studying admixed individuals can be complex, and leveraging a pure

statistical approach, we grouped individuals from rather culturally, ethnically,

linguistically, and genetically heterogeneous groups to represent the four Native

American components discussed here. Yet, the present study provides useful

insights into the routes followed by the main population arrivals in the Southern

Cone (**S24 Fig**).

Further efforts are needed to better characterize the Native American ancestry

component identified in the Central Western region of Argentina. Particularly, we

encourage future studies to confirm the tentative geographical label that we suggest

here, and to estimate its influence in the region. Besides these specific questions,

many other general questions remain to be answered to better understand the pre-

Columbian population dynamics in the Southern Cone such as the time and place of

the splits among the components described here, and the extent of genetic

exchanges among them. More genotype data for ancient samples, modern

indigenous communities and admixed individuals, particularly in Central,

Northeastern and Patagonia regions of Argentina, would help to decipher these

issues.

The genomic characterization of populations is an unavoidable practice for many

issues ranging from the understanding of our biological heritage, the rational use of

biobanks, the definition of an adequate reference genome, the estimation of

polygenic risk scores, the study and treatment of simple and complex diseases, and

the design of a national program of genomic medicine in our country. This study is a

joint effort of Argentinean institutions funded by the national scientific system, and

represents the first milestone of the *Consorcio Poblar*, a national consortium for

creating a public reference biobank to support biomedical genomic research in

Argentina [72]. Genomic knowledge of local populations should be a priority for

developing countries to achieve an unbiased representation of diversity in public

databases and the scientific development in periphery countries.

Material and Methods

Studied Samples

We genotyped 94 individuals with the Axiom LAT1 array (Affymetrix) from 24

localities and 17 provinces across Argentina (**Fig 1**). These samples were selected

among 240 collected by different population genetics groups (Consorcio PoblAR)

during past sampling campaigns with a biological anthropology focus. According to

the available information (e.g. interviews, genealogical information, etc.), each

PoblAR research group selected for this study some samples, maximizing the odds

that they come from individuals with greater Native American ancestry. For

example, surnames were used as a proxy to achieve this objective and the

permanence of ancestors in national territory has been another variable that was

taken into account. The analyzed 94 samples were also selected to ensure an

extended geographical range and were included when they presented sufficient

DNA concentration and Native American maternal lineage. Moreover, among the

males, we prioritized those with Native American paternal lineage.

All saliva and blood samples were collected under written informed consent to

participate to this study. The informed consents for research use were approved by

several Ethics Committees from (i) Hospital Zonal de Puerto Madryn (Resolución

009/2015), (ii) Hospital Zonal de San Carlos de Bariloche (Resolución 1510/2015),

(iii) Hospital Provincial de Pediatría Dr. Fernando Barreyro, (iv) Investigaciones

Biomédicas IMBICE (RENIS CE000023), (v) Provincia de Jujuy, (vi) Hospital Italiano

of Buenos Aires (protócolo 1356/09); and (vii). Centro de Educación Médica e

Investigaciones Clínicas "Norberto Quirno" (Resolución 612/2018). The biological

samples were coded and anonymized, as per the Argentina National Law 25.326 of

Protection of Personal Data.

87 samples and 791,543 autosomal variants passed the standard Affymetrix

genotyping Quality Controls (**S1 Table**).

Most of the genotype data processing was performed using in-house scripts in R

[73] and perl [74], leveraging *plink2* [75], *vcftools v1.13* [76], and *bedTools v2* [77].

We compiled the genotype data for the 87 Argentinean samples with different

genotype data available in the literature. We focused our study on biallelic SNPs

(removing indels and SNPs with more than 2 alleles). Any putative inconsistent

strand had been fixed processing to the relevant flip, filtering out any SNP with

ambiguous genotype (A/T, G/C).

Cryptic relatedness among samples were assessed using King software [78]. To

avoid any 1st degree relationship, we filtered out individuals, minimizing the total

number of removals. No admixed individuals had been removed at this step.

We thus build different dataset arrangements (named **DS\<n\>**) that we analyzed

through this work (**S2 Table**).

714   Argentinean genetic diversity within a worldwide context

To analyze genetic diversity in Argentina within a worldwide context, we built the

Dataset1 (**DS1**). This dataset contains 87 Argentinean samples, 654 African, 503

European and 179 South American samples from 1KGP [42], 54 modern unrelated

Chilean samples [37], and 161 Native-American individuals from South America

[38]. Moreover, **DS1** included genotype data from [31] which consists in 82

individuals from Lima (Peru), 27 from Santiago de Chile, and 161 from Argentinean

urban centers.

We filtered out any variant and individual exhibiting in the compiled data set more

than 2% and 5% of missing genotypes (--geno 0.02 and --mind 0.05 flags in plink

1.9), respectively, as well as Minor Allele Frequency below 1% (--maf 0.01 flag in

plink 1.9). The filtered data was then pruned for Linkage-Disequilibrium (--indep-

pairwise 50 5 0.5 flag in plink 1.9). The combined data set has a total intersection of

59,237 SNPs and 1,908 individuals. With this curated data we performed Principal

Component Analyses [41] (**S1 Fig**) and Admixture [43] (**Fig 2, S2 Fig and S3 Table**).

Local Ancestry

Local ancestry analyses rely on haplotype reconstruction (phasing) and require high

SNP density. Since, admixed individuals of interest were genotyped on different

microarray platforms, we decided to perform two separate local ancestry analyses

on two different data sets (**DS2** and **DS3**). **DS2** consists of 87 Argentinean

individuals from the present study, and 54 Chilean individuals [37], all of them

genotyped with the Axiom LAT1 microarray. **DS3** includes 175 Argentinean, 27

Chilean, and 119 Peruvian genotyped with the Illumina OMNI1 microarray [31].

Both **DS2**, **DS3** were merged with 1KGP data consisting in 503 phased reference

samples for each of the African and European genetic ancestry, and 347 Latin

American individuals.

The 87 Argentinean, and the 54 Chilean samples were phased with shapeIT2

[79,80]. The genetic map, and 5,008 haplotypes panel provided by 1KGP were

downloaded from http://mathgen.stats.ox.ac.uk/impute/1000GP_Phase3/.

Algorithm, and model parameters were used by default, filtering out monomorphic

SNPs, and those with more than 2% of missing genotype. We obtained phased

genotype data for 608,501 autosomal SNPs. This data was then merged with phased

1KGP genotype data for African, American and European samples described before.

Since this data set derived from DS2, we call it **DS2p** (phased DS2).

The 175 Argentinean, 27 Chilean and 119 Peruvian samples from [31] were phased

separately using the same procedure with shapeIT2. After filtering for missing

genotypes and merging with phased 1KGP genotype data for African, American and

European samples we obtained phased data for 694,626 autosomal SNPs. Since this

data set derived from DS3, we call it **DS3p** (phased DS3).

In **DS2** and **DS3** we ran Admixture using K parameter minimizing the Cross-

Validation score. We used individuals with more than 99% of Native American

ancestry as references for local ancestry estimation. For **DS2** we used $K = 7$, and

Native American ancestry was defined as the sum of the two American specific

components observed (**S25 Fig**). According to this criterion, 48 individuals were

assigned as Native American reference. All the other American samples were

defined as Admixed.

For **DS3**, we used $K = 5$, and Native American ancestry was defined as the single

American specific component observed (**S26 Fig**). According to this criterion, 19

individuals were assigned as Native American reference. All the other American

samples were defined as Admixed.

RFMix v2 ([44] downloaded at https://github.com/slowkoni/rfmix on 15th August

2018) was run on **DS2P** and **DS3P** separately using parameter settings similar to

[81]. The reference panels consist of the African and European samples from 1KGP,

as well as Native American individuals identified through Admixture procedure

described before. We used 1 Expectation-Maximization iteration (-e 1), actualizing

the reference panel in this process (--reanalyze-reference). We used CRF spacing

size and random forest window size of 0.2 cM (-c 0.2 and -r 0.2). We use a node size

of 5 (-n 5). We set the number of generations since admixture to 11 (-G 11)

considering the estimates from [31]. The forward-backward output was then

interpreted to assign allele ancestry to the one exhibiting major posterior

probability, conditioning that it was greater than 0.9. Otherwise, the allele ancestry

was assigned to Unknown (UKN). As a sanity check, the global ancestry proportions

estimated through this RFMix analysis were compared with those obtained with

Admixture software. The global ancestry proportion estimates obtained by both

procedures matched very well: spearman's correlation greater than 0.9 in American

samples for any of the 3 continental ancestries (**S27 Fig**). Moreover, the ancestry

ditypes assigned in admixed individuals from 1KGP (included in both DS2p and

DS3p) were highly consistent between both independent masking procedures (**S28**

Fig).

Ancestry Specific Population Structure

784   In order to analyze the ancestry-specific population structure we masked the data,

785   i.e. for each individual, we assigned missing genotype for any position for which at

786   least one of the two alleles was not assigned to the relevant ancestry. In other

787   words, to study ancestry A, we kept for each individual, regions exhibiting ancestry

788   A on both haplotypes (ditypes) as illustrated in **S5 Fig**

European ancestry specific population structure

To study European ancestry specific population structure, we analyzed together

masked data for this ancestry for Colombian individuals from 1KG and individuals

from **DS2P** and **DS3P** excluding individuals from Chilean Native American

communities [37] . This data was merged with a set of reference individuals with

European ancestry [46], which is a subset of the POPRES dataset [45]. We call this

data set as **DS4**. We removed individuals with less than 30% SNPs with the ancestry

ditypes (--mind 0.7 with *plink* 1.9). We also removed SNPs with more than 50% of

missing genotypes (--geno 0.5 with *plink* 1.9). Thus, **DS4** contains 132 modern

Argentinean individuals (29 from the present study and 103 from [31]), 17

individuals from Santiago de Chile, 4 from Lima and 74 from Colombia [42]. **DS4**

encompasses 29,347 SNPs of which 27,634 remained after LD-pruning (--indep-

pairwise 50 5 0.5 flag in *plink2*).

Smartpca from Eigensoft v7.2.0 was run on **DS4** [41] with the *lsqproject* option ON*.

We report the PCA results summarized into a 2-dimensional space by applying

Multidimensional Scaling on weighted Euclidian distance matrix for the first N PCs.

We weighted each PC by the proportion of variance it explains. We selected the N

most informative PCs according to the Elbow method on the proportion of explained

variance. Admixture [43] was run with K ranging from 2 to 10 with cross-validation

procedure.

African ancestry specific population structure

To study African ancestry specific population structure, we analyzed together

masked data for this ancestry for individuals from **DS2P** and **DS3P**. This data was

merged with a compilation of reference individuals with African ancestry from

[42,48–50]. We removed African individuals with less than 99% of African ancestry

when comparing them to the 2504 individuals from 1KGP (Admixture with K=7

minimizing cross-validation score). We thus reduced the African reference to 1685

individuals. We call as **DS5** the data set containing both the masked data for

admixed South American individuals and African reference individuals.

We removed SNPs with more than 10% of missing genotypes (--geno 0.1 with *plink*

1.9), and individuals with less than 5% of the ancestry ditypes (--mind 0.95 with

plink 1.9). Thus, **DS5** contains, 26 modern Argentinean individuals (all from the

present study), and 12 individuals from Lima (*9*). **DS5** consisted in 137,136 SNPs, of

which 128,086 remained after LD-pruning (--indep-pairwise 50 5 0.5 flag in *plink2*).

PCA and Admixture were performed as for European ancestry specific population

structure analyses (described before).

Native American ancestry specific population structure

To study Native American ancestry specific population structure, we analyzed

together masked data for this ancestry for individuals from **DS2P** and **DS3P**. This

data was merged with pseudo-haploid data for ancient samples within South and

Central America [35–37,54,55], as well as with masked data for Native American

individuals from [38]. The pseudo-haploid data for ancient samples was

downloaded from the Reich Laboratory webpage

(https://reich.hms.harvard.edu/downloadable-genotypes-present-day-and-ancient-

dna-data-compiled-published-papers) the 15th of May 2019. For each sample, we

used annotations (geographic coordinates and approximate date) from the metafile

provided at the same url.

We call this data set **DS6.** We removed individuals with less than 30% SNPs with the

ancestry ditypes (--mind 0.7 with *plink* 1.9). We also removed SNPs with less than

50% individuals with the ancestry ditypes (--geno 0.5 with *plink* 1.9). **DS6** contains

146 modern Argentinean individuals (74 from the present study and 72 from [31]),

22 individuals from Santiago and 77 from Lima, along with 207 Native South

American individuals from [37,38] and 53 ancient samples. **DS6** encompasses

47,003 SNPs, of which 39,423 remained after LD-pruning (--indep-pairwise 50 5 0.5

flag in *plink2*).

PCA and Admixture were performed as for European ancestry specific population

structure analyses (described before), with the difference that the *poplistname*

option was set for *smartpca* in order to estimate the PC using only modern samples

and project the ancient samples.

Statistical assignation of modern South American individuals to Native American components

848
849

Given a distance matrix among modern individuals from **DS6**, we performed *K-*

means clustering with the number of clusters ranging from 2 to 20 and selected the

K-means output minimizing the Bayesian Information Criterion (BIC). This

procedure was applied to three different distance matrix: (i) Weighted Euclidean

distance based on the two first PCs from Principal Component Analysis (ii)

Euclidean distance based on the ancestry proportions estimated from Admixture

(K=3) and (iii) $1 - f_3(Ind1,Ind2,Yoruba)$, where *Ind1* and *Ind2* are two individuals. In

the three cases, the *K*-means procedure minimized the BIC when considering 4

clusters. These clusters correspond to the three Native American Component

discussed in this paper (CAN, CCP and STF), along with one laying in-between, which

we finally attributed to Central Western Argentina. However, individual assignment

to one of these four clusters was not totally consistent according to the distance

matrix we used. To obtain a robust assignation, an individual was assigned to a

given cluster when it consistently belonged to it across the three *K*-means

procedures, otherwise it had been removed for following analyses (**S16 Fig**). The

cluster assignation for each individual is given in **S4 Table**.

We computed pairwise F_{ST} among the groups using smartpca [41] with fstonly and

lsqproject set to YES, and all the other settings left at default. Standard errors were

estimated with the block jackknife procedure [41].

F-statistics to infer relationship between the four Native
American components in Argentina and their genetic affinity
with ancient populations

Starting from genotype data for the individuals in **DS6**, we also included genotype

data from Yoruba (YRI) population from [42], Mixe population from [38] and,

pseudo-haploid data for Anzick individual from the Clovis culture [67] and USR1

individual from Upward Sun River in Beringia [66]. The resulting data set is called

DS7. Within **DS7,** we grouped modern individuals according to their assigned Native

American group and we removed those with inconsistent assignation (**S16 Fig** and

S4 Table). Thus, **DS7** contains 426 modern individuals, 55 ancient samples, 108

Yoruba and 17 Mixe individuals. **DS7** encompasses 88,564 SNPs. Note that we did

not apply any SNP filtering overall **DS7** in order to maximize the number of SNPs

included in each group comparison considered.

Using modern individuals from **DS7**, and considering any possible combination of

the four groups identified (STF, CCP, CAN and CWA), we computed the f_3 statistics

[82] in the form of $f_3(Target; S1, S2)$. This allowed to contrast whether *Target, S1* and

S2 could be organized in the form of a phylogenetic tree *(positive f_3)* or whether the

Target group is the result of mixture between *S1* and *S2* groups (negative f_3). We

also computed $f_4(YRI, Target; S1, S2)$ to test whether group *Target* shares more

evolutionary history with group *S1* (negative f_4) or group *S2* (positive f_4). Moreover,

we reconstructed the Neighbour-joining tree from the matrix of distances with

Phylip v3.2 [58] and *USR1* as outgroup. The distances were estimated as $1/f_3$-

outgroup(YRI; X, Y).

We computed f_3-Outgroup and f_4 statistics in order to estimate the genetic affinity of

the four Native American components with ancient populations from South and

Central America. We computed the *f_3-outgroup* and f_4 statistics in the form of *$f_3(YRI$;*

X, Ancient) and *$f_4(YRI, X; USR1, Ancient)$*, where *X* the represent the cluster

containing all individuals assigned to a given Native American ancestry component.

We also computed f_4 of the form $f_4(YRI, Ancient; X, Y)$. This f_4 setting allowed testing

which of *X* or *Y,* each referring to one of the four Native American components.

shares more ancestry with a given *Ancient* group. *X* and *Y*.

We finally computed f_4 of the form $f_4(Ancient, X; Y, YRI)$ to test whether a given

modern Native American component *Y* shares exhibit closer genetic affinity with a

given *Ancient* group (negative f_4) or with another modern Native American

component *X* (positive f_4).

All the results based on *F-statistics* are listed in **S5 Table**. We assessed significance

of a comparison considering 3 standard errors ($|Z| > 3$) from block jackknife

procedure [82].

Geographical maps

All maps have been generated in R using, *maps* package , (https://cran.r-

[project.org/web/packages/maps/index.html](https://cran.r-project.org/web/packages/maps/index.html)). **S24 Fig** includes a raster

downloaded at https://www.naturalearthdata.com/ and integrated in R with *raster*

(https://CRAN.R-project.org/package=raster) and *RStoolbox*

(http://bleutner.github.io/RStoolbox) packages.

Abbreviations

1KGP: 1000 Genomes Project

AMBA: Metropolitan Area of Buenos Aires

BIC: Bayesian Information Criterion

CAN: Central Andes

CCP: Central Chile / Patagonia

CLM: Colombians from Medellin

CWA: Central Western Region in Argentina

CYA: Cuyo Region in Argentina

DS: Data set

LD: Linkage Disequilibrium

NEA: Northeastern Region in Argentina

NWA: Northwestern Region in Argentina

PCA: Principal Component Analysis

PEL: Peruvians from Lima

PPA: Pampean Region in Argentina

PTA: Patagonia Region in Argentina

SNP: Single Nucleotide Polymorphism

STF: Subtropical and Tropical Forests

YRI: Yoruba individuals in Ibadan, Nigeria

Acknowledgments

In memoriam of Raúl Carnese.

We thank Rolando González-José, Andrea Llera and Mariana Berenstein for the

management and promotion of the Consorcio Poblar.

We thank all the participants of the different sampling campaigns across Argentina

from which the newly genotyped samples are derived from.

We thank Ricardo A. Verdugo, Etienne Patin, Luca Pagani, Marta E. Alarcón

Riquelme and María Teruel Artacho who kindly shared genotype data included in

this study.

We thank Laura Fejerman for her proofreading of the manuscript.

946   Disclosure Declaration

PL provides consulting services to myDNAmap S.L.

JMB and JZ are employed by Biocódices S.A; HD is the scientific director of

Biocódices S.A.

Biocodices provided support in the form of part-time employment salaries for

authors HD, JMB and JZ.

Neither Biocódices nor myDNAmap have made any decision in the design of the

study, sampling biological material, the collection, and the analysis of the data.

These companies have also not made any decision on the preparation and

publication of the work. Biocódices provided inputs for DNA's sample extraction and

access to computing cluster resources.

957   Data Availability Statement

958   The data analyzed here comprises both newly generated and previously reported

959   data sets. Access to publicly available data sets should be requested through the

960   distribution channels indicated in each published study. For newly genotyped

961   samples, individual genotype data will be made available at

962   http://biocodices.hopto.org/Affymetrix_Pop_Arg/ at time of publication.

Bibliography

1.   Bravi CM, Sans M, Bailliet G, Martínez-Marignac V, Portas M, Barreto I, et al.

Characterization of mitochondrial DNA and Y-chromosome haplotypes in a

Uruguayan population of African ancestry. Hum Biol. 1997;69: 641–652.

2.   Sala A, Penacino G, Corach D. Comparison of allele frequencies of eight STR

loci from Argentinean Amerindian and European populations. Hum Biol.

1998;70: 937–947.

3.   Goicoechea A., Carnese FR, Dejean C, Avena SA, Weimer TA, Franco MHLP, et

al. Genetic Relationships Between Amerindian Populations of Argentina. Am J

Phys Anthropol. 2001;115: 133–143.

4.   Dejean CB, Crouau-Roy B, Goicoechea AS, Avena SA, Carnese FR. Genetic

variability in Amerindian populations of northern Argentina. Genet Mol Biol.

2004;27: 489–495.

5.   Fejerman L, Carnese F, Goicoechea A, Avena S, Dejean C, Ward R. African

ancestry of the population of Buenos Aires. Am J Phys Anthropol. 2005;128:

164–70.

6.   Wang S, Ray N, Rojas W, Parra M V., Bedoya G, Gallo C, et al. Geographic

patterns of genome admixture in latin American mestizos. PLoS Genet.

2008;4: e1000037.

7.   Bobillo MC, Zimmermann B, Sala A, Huber G, Röck A, Bandelt H-J, et al.

Amerindian mitochondrial DNA haplogroups predominate in the population

of Argentina: towards a first nationwide forensic mitochondrial DNA

sequence database. Int J Legal Med. 2010;124: 263–268.

8.   Pauro M, Garcia A, Bravi CM, Demarchi DA. Distribucion de haplogrupos

mitocondriales aloctonos en poblaciones rurales de Cordoba y San Luis. Rev

Argentina Antropol Biol. 2010;12: 47–55.

9.   Pauro M, García A, Nores R, Demarchi D a. Analysis of uniparental lineages in

two villages of Santiago Del Estero, Argentina, seat of Pueblos de Indios in

colonial times. Hum Biol. 2013;85: 699–720.

10.   García A, Pauro M, Bailliet G, Bravi CM, Demarchi DA. Genetic variation in

populations from central Argentina based on mitochondrial and Y

chromosome DNA evidence. J Hum Genet. 2018.

11.   Martínez Marignac VL, Bertoni B, Parra EJ, Bianchi NO. Characterization of

admixture in an urban sample from Buenos Aires, Argentina, using

uniparentally and biparentally inherited genetic markers. Hum Biol. 2004;76:

543–57.

12.   Seldin MF, Tian C, Shigeta R, Scherbarth HR, Silva G, Belmont JW, et al.

Argentine Population Genetic Structure: Large Variance in Amerindian

Contribution. Am J Phys Anthropol. 2007;132: 455–462.

13.   Corach D, Lao O, Bobillo C, van Der Gaag K, Zuniga S, Vermeulen M, et al.

1003 Inferring continental ancestry of argentineans from autosomal, Y-

1004 chromosomal and mitochondrial DNA. Ann Hum Genet. 2010;74: 65–76.

1005 14. Parolin ML, Toscanini UF, Velázquez CL, Berardi GL, Holley A, Tamburrini C, et

1006 al. Genetic admixture patterns in Argentinian Patagonia. PLoS One. 2019;14:

1007 e0214830.

1008 15. Avena S, Via M, Ziv E, P??rez-Stable EJ, Gignoux CR, Dejean C, et al.

1009 Heterogeneity in genetic admixture across different regions of argentina. PLoS

1010 One. 2012;7: e34695.

1011 16. García A, Dermarchi DA, Tovo-Rodrigues L, Pauro M, Callegari-Jacques SM,

1012 Salzano FM, et al. High interpopulation homogeneity in Central Argentina as

1013 assessed by ancestry informative markers (AIMs). Genet Mol Biol. 2015;38:

1014 324–331.

1015 17. Salas A, Richards M, Lareu M-V, Scozzari R, Coppa A, Torroni A, et al. The

1016 African diaspora: mitochondrial DNA and the Atlantic slave trade. Am J Hum

1017 Genet. 2004;74: 454–465.

1018 18. García A, Pauro M, Nores R, Bravi CM, Demarchi DA. Phylogeography of

1019 mitochondrial haplogroup D1: An early spread of subhaplogroup D1j from

1020 Central Argentina. Am J Phys Anthropol. 2012;149: 583–590.

1021 19. Corach D, Figueira Risso L, Marino M, Penacino G, Sala A. Routine Y-STR

1022 typing in forensic casework. Forensic Sci Int. 2001;118: 131–135.

1023 20. Salas A, Jaime JC, Álvarez-Iglesias V, Carracedo Á. Gender bias in the

1024 multiethnic genetic composition of central Argentina. J Hum Genet. 2008;53:

1025   662–674.

1026   21.   Martínez Sarasola C. Nuestros paisanos los indios. 9th ed. Buenos Aires: Del

1027   Nuevo Extremo; 2011.

1028   22.   Devoto F. Historia de la inmigración en la Argentina. 1a ed. Buenos Aires:

1029   Editorial Sudamericana; 2003.

1030   23.   Studer E. La trata de negros en el Río de la Plata durante el siglo XVIII. Buenos

1031   Aires: Instituto de Historia de Argentina "Doctor Emilio Ravignani"; 1958.

1032   24.   Rodríguez Molas R. Los sometidos de la conquista. Centro Edi. Buenos Aires;

1033   1985.

1034   25.   Borucki A. The Slave Trade to the Río de la Plata, 1777–1812: Trans-Imperial

1035   Networks and Atlantic Warfare. Colon latin Am Rev. 2011;20: 81–107.

1036   26.   Guzmán F. Africanos en la Argentina. Una reflexión desprevenida. Andes.

1037   2006;17.

1038   27.   Pauro M. Análisis molecular de linajes uniparentales en poblaciones humanas

1039   del centro de Argentina. Universidad de Córdoba. 2015.

1040   28.   Di Fabio Rocca F. La presencia subsahariana en el acervo génico de

1041   poblaciones cosmopolitas de la Argentina. Universidad de Buenos Aires. 2016.

1042   29.   Mandrini R. La Argentina aborigen. De los primeros pobladores a 1910. Siglo

1043   XXi. Buenos Aires; 2008.

1044   30.   Lenton DI, Delrio WM, Pérez PMV, Papazian AER, Nagy MA, Musante M.

1045   Argentina's Constituent Genocide: Challenging the Hegemonic National

Narrative and Laying the Foundation for Reparations to Indigenous Peoples.

Armen Rev. 2012;53: 63–84.

31.   Homburger JR, Moreno-Estrada A, Gignoux CR, Nelson D, Sanchez E, Ortiz-

Tello P, et al. Genomic Insights into the Ancestry and Demographic History of

South America. PLOS Genet. 2015;11: e1005602.

32.   Muzzio M, Motti JMB, Sepulveda PBP, Yee M, Santos R, Ramallo V, et al.

Population structure in Argentina. PLoS One. 2018;13: e0196325.

33.   Gouveia MH, Borda V, Leal TP, Moreira RG, Bergen AW, Kehdy FSG, et al.

Origins, admixture dynamics and homogenization of the African gene pool in

the Americas. Mol Biol Evol. 2020;Epub ahead: pii: msaa033.

34.   Kehdy FSG, Gouveia MH, Machado M, Magalhães WCS, Horimoto AR. Origin

and dynamics of admixture in Brazilians and its effect on the pattern of

deleterious mutations. Proc Natl Acad Sci U S A. 2015;112: 8696–8701.

35.   Posth C, Nakatsuka N, Lazaridis I, Fehren-schmitz L, Krause J, Reich D, et al.

Reconstructing the Deep Population History of Central and South America

Article Reconstructing the Deep Population History of Central and South

America. Cell. 2018;175: 1185–1197.

36.   Moreno-Mayar JV, Vinner L, Damgaard PDB, Fuente C De, Margaryan A,

Orbegozo MI, et al. Early human dispersals within the Americas. Science.

2018;362: pii: eaav2621.

37.   de la Fuente C, Ávila-arcos MC, Galimany J, Carpenter ML, Homburger JR.

Genomic insights into the origin and diversification of late maritime hunter-

gatherers from the Chilean Patagonia. Proc Natl Acad Sci U S A. 2018;115:

E4006–E4012.

38.   Reich D, Patterson N, Campbell D, Tandon A, Mazieres S, Ray N, et al.

Reconstructing Native American population history. Nature. 2012;488: 370–4.

39.   Gnecchi-ruscone GA, Sarno S, Fanti S De, Gianvincenzo L, Giuliani C, Boattini A,

et al. Dissecting the Pre-Columbian Genomic Ancestry of Native Americans

along the Andes – Amazonia Divide. Mol Biol Evol. 2019;36: 1254–1269.

40.   Gomez-Carballa A, Pardo-Seco, Brandini S, Achilli A, Perego UA, Coble MD, et

al. The peopling of South America and the trans-Andean gene flow of the first

settlers. Genome Res. 2018 Jun;28(6):767-779.

41.   Patterson N, Price AL, Reich D. Population structure and eigenanalysis. PLoS

Genet. 2006;2: 2074–2093.

42.   The 1000 Genomes Project Consortium. A global reference for human genetic

variation. Nature. 2015;526: 68–74.

43.   Alexander DH, Novembre J, Lange K. Fast model-based estimation of ancestry

in unrelated individuals. Genome Res. 2009;19: 1655–64.

44.   Maples BK, Gravel S, Kenny EE, Bustamante CD. RFMix: A discriminative

modeling approach for rapid and robust local-ancestry inference. Am J Hum

Genet. 2013;93: 278–288.

45.   Nelson MR, Bryc K, King KS, Indap A, Boyko AR, Novembre J, et al. The

Population Reference Sample, POPRES: A Resource for Population, Disease,

and Pharmacological Genetics Research. Am J Hum Genet. 2008;83: 347–358.

46.   Novembre J, Johnson T, Bryc K, Kutalik Z, Boyko AR, Auton A, et al. Genes

mirror geography within Europe. Nature. 2008;456: 98–101.

47.   Gallero MC, Krautstofl EM. Proceso de poblamiento y migraciones en la

Provincia de Misiones , Argentina (1881-1970). Avá Rev Antropol. 2010;16:

1881–1970.

48.   Patin E, Lopez M, Grollemund R, Verdu P, Harmant C, Quach H, et al.

Dispersals and genetic adaptation of Bantu-speaking populations in Africa and

North America. Science. 2017;546: 543–546.

49.   Pagani L, Kivisild T, Tarekegn A, Ekong R, Plaster C, Romero IG, et al.

Ethiopian Genetic Diversity Reveals Linguistic Stratification and Complex

Influences on the Ethiopian Gene Pool. Am J Hum Genet. 2012;91: 83–96.

50.   Schlebusch CM, Skoglund P, Sjödin P, Gattepaille LM, Hernandez D, Jay F, et al.

Genomic Variation in Seven Khoe-San Groups Reveals Adaptation and

Complex African History. Science. 2012;338: 374–9.

51.   Moreno-Estrada A, Gravel S, Zakharia F, McCauley JL, Byrnes JK, Gignoux CR,

et al. Reconstructing the Population Genetic History of the Caribbean. PLoS

Genet. 2013;9.

52.   Fortes-lima C, Gessain A, Ruiz-linares A, Restrepo BN, Rojas W, Migot-nabias

F, et al. Genome-wide Ancestry and Demographic History of African-

Descendant Maroon Communities from French Guiana and Suriname. Am J

Hum Genet. 2017;101: 725–736.

53.   Bryc K, Auton A, Nelson MR, Oksenberg JR, Hauser SL, Williams S, et al.

1112   Genome-wide patterns of population structure and admixture in West

1113   Africans and African Americans. Proc Natl Acad Sci U S A. 2010;107: 786–791.

1114   54.   Raghavan M, Steinrücken M, Harris K, Schiffels S, Rasmussen S, DeGiorgio M,

1115   et al. Genomic evidence for the Pleistocene and recent population history of

1116   Native Americans. Science (80-). 2015;349: aab3884.

1117   55.   Lindo J, Haas R, Hofman C, Apata M, Moraga M, Verdugo RA, et al. The genetic

1118   prehistory of the Andean highlands 7000 years BP though European contact.

1119   Sci Adv. 2018;4: eauu4921.

1120   56.   Motti JMB, Muzzio M, Ramallo V, Kladniew BR, Alfaro EL, Dipierri JE, et al.

1121   Origen Y Distribución Espacial De Linajes Maternos Nativos En El Noroeste Y

1122   Centro Oeste Argentinos. Rev Argentina Antropol Biol. 2013;15: 3–14.

1123   57.   Motti J. Caracterización de Linajes Maternos en la Población Actual del

1124   Noroeste y Centro-Oeste Argentinos. Universidad Nacional de La Plata. 2012.

1125   58.   Felsenstein J. PHYLIP-Phylogeny Inference Package (Ver. 3.2). Cladistics.

1126   1989;5: 164–166.

1127   59.   Motti JMB, Schwab ME, Beltramo J, Jurado-medina LS, Muzzio M, Ramallo V, et

1128   al. Regional differentiation of native American populations from the analysis

1129   of maternal lineages. Intersecc en Antropol. 2017;18: 271–282.

1130   60.   Canals Frau S. Etnología de los Huarpes. Una síntesis. Anales del Instituto de

1131   Etnología Americana. Anales del Instituto de Etnología Americana, Tomo VII.

1132   Mendoza; 1946. pp. 9–147.

61.   Escolar D. Los dones étnicos de la Nación. Identidades huarpe y modos de
producción de soberanía en Argentina. Prometeo E. Buenos Aires; 2007.

62.   Bengoa J. Historia de los antiguos mapuches del sur. Desde antes de la llegada
de los españoles hasta las paces de Quilín. Editorial. Santiago; 2003.

63.   Galdames S. ¿Detuvo la batalla del Maule la expansión inca hacia el sur de
Chile? Cuardernos Hist. 2017;3: 7–25.

64.   Michieli CT. Antigua historia de Cuyo. Ansilta Ed. San Juan; 1994.

65.   Gascón M. Cuyo en el espacio imperial. La fase de configuración: 1580-1680.
Rev Tefros. 2011;9: 1–20.

66.   Moreno-Mayar JV, Potter BA, Vinner L, Steinrücken M, Rasmussen S, Terhorst
J, et al. Terminal Pleistocene Alaskan genome reveals first founding
population of Native Americans. Nature. 2018;553: 203–207.

67.   Rasmussen M, Anzick SL, Waters MR, Skoglund P, Degiorgio M, Jr TWS, et al.
The genome of a Late Pleistocene human from a Clovis burial site in western
Montana. Nature. 2014;506: 225–229.

68.   Michieli CT. La disolución de la categoría Jurídico-Social de "indio" en el siglo
XVIII: El caso de San Juan (Región de Cuyo). Publicacio. Universidad Nacional
de San Juan; 2000.

69.   Katzer L. El mestizaje como dispositivo biopolítico. Biblos. Pueblos indígenas
Interculturalidad, colonialidad, política. Biblos. Buenos Aires; 2009. pp. 59–7.

70.   Briones C. Construcciones de aboriginalidad en Argentina. Société suisse des

Américanistes / Schweizerische Amerikanisten-Gesellschaft. 2004;Bulletin 6:

73–90.

71.   Mas-sandoval A, Arauna LR, Gouveia MH, Barreto ML, Horta BL, Lima-costa

1157   MF, et al. Reconstructed Lost Native American Populations from Eastern

1158   Brazil Are Shaped by Differential Jê/Tupi Ancestry. Genome Biol Evol.

1159   2019;11: 2593–2604.

1160   72.   Dopazo H, Llera AS, Berenstein M, Gonzáles-josé R. Genomas , enfermedades y

1161   medicina de precisión : un Proyecto Nacional. Ciencia, Tecnol y Política.

1162   2019;2: 1–10.

1163   73.   Team RC. R: A language and environment for statistical computing. R

1164   Foundation for Statistical Computing, Vienna, Austria. 2019.

1165   74.   Wall L, Christiansen T, Orwant J. Programming perl. O'Reilly M. 2000.

1166   75.   Chang CC, Chow CC, Tellier LCAM, Vattikuti S, Purcell SM, Lee JJ. Second-

1167   generation PLINK: Rising to the challenge of larger and richer datasets.

1168   Gigascience. 2015;4: 1–16.

1169   76.   Danecek P, Auton A, Abecasis G, Albers CA, Banks E, Depristo MA, et al. The

1170   variant call format and VCFtools. Bioinformatics. 2011;27: 2156–2158.

1171   77.   Quinlan AR, Hall IM. BEDTools : a flexible suite of utilities for comparing

1172   genomic features. Bioinformatics. 2010;26: 841–842.

1173   78.   Manichaikul A, Mychaleckyj JC, Rich SS, Daly K, Sale M, Chen W-M. Robust

1174   relationship inference in genome-wide association studies. Bioinformatics.

2010;26: 2867–73.

79.   Delaneau O, Marchini J, Zagury J-F. A linear complexity phasing method for

thousands of genomes. Nat Methods. 2011;9: 179–181.

80.   Delaneau O, Marchini J, Zagury J. A linear complexity phasing method for

thousands of genomes. Nat Methods. 2012;9: 179–181.

81.   Martin AR, Gignoux CR, Walters RK, Wojcik GL, Neale BM, Gravel S, et al.

Human Demographic History Impacts Genetic Risk Prediction across Diverse

Populations. Am J Hum Genet. 2017;100: 635–649.

82.   Patterson N, Moorjani P, Luo Y, Mallick S, Rohland N, Zhan Y, et al. Ancient

admixture in human history. Genetics. 2012;192: 1065–93.

1185

Supplementary Figure Captions

S1 Fig. Principal component analysis in a worldwide Context.
A: PC2 vs PC1; **B:** PC4 vs PC3; **C:** PC6 vs PC5. The percentage of variance explained by each principal component (PC) is given. Each point represents an individual following the color and point codes given in legend.

S2 Fig. Admixture analysis in a worldwide context.
(A) Cross-Validation score for Admixture runs on the worldwide meta dataset (**DS1**) with K from 3 to 12. **(B-G)** Admixture results with K=3 to K=8. 1KGP: 1000 Genomes Project; CYA: Cuyo Region; NEA: Northeastern Region, NWA: Northwestern Region; PPA: Pampean Region; PTA: Patagonia Region.

S3 Fig. Comparison of different admixture continental ancestry proportion estimates in a worldwide context.
Comparison of the African, European and Native American ancestry proportion estimates obtained with Admixture models with K=3 and K=8 applied to **DS1. (A)** African ancestry proportions for K=3 are as observed in green in **S2B Fig**, while for K=8 they are estimated as the sum of the three greenish colors observed in **Main Fig 2**. **(B)** European ancestry proportions for K=3 are as observed in blue in **S2B Fig**, while for K=8 they are estimated as the sum of the two bluish colors observed in **Main Fig 2**. **(C)** Native American ancestry proportions for K=3 are as observed in orange in **S2B Fig**, while for K=8 they are estimated as the sum of the three reddish colors observed in **Main Fig 2**.

S4 Fig. Correlation between eigenvectors and ancestry proportion estimates from analyses in a worldwide context.
Comparison of ancestry proportion estimates from Admixture model with $K=8$ and the 6 first Principal Components (PCs) in **DS1. (A)** PC1 vs African ancestry proportions (estimated as the sum of the three greenish colors observed in **Main Main Fig 2**). **(B)** PC2 vs European ancestry proportions (estimated as the sum of the two bluish colors observed in **Main Fig 2**). **(C)** PC2 vs Native American ancestry proportions (estimated as the sum of the three redish colors observed in **Main Fig 2**). **(D)** PC6 vs Bantu-influenced ancestry proportions (dark olive green in **Main Fig 2**). **(E)** PC6 vs Western African ancestry proportions (dark green in **Main Fig 2**). **(F)** PC4 vs Southern European ancestry proportions (light blue in **Main Fig 2**). **(G)** PC4 vs Northern European ancestry proportions (dark blue in **Main Fig 2**). **(H)** PC3 vs Cenral Chile / Patagonia ancestry proportions (orange in **Main Fig 2**). **(I)** PC3 vs Central Andes ancestry proportions (yellow in **Main Fig 2**). **(J)** PC5 vs Subtropical and Tropical Forests ancestry proportions (pink in **Main Fig 2**).

S5 Fig. Example of a local ancestry output.
(A) RFMIX output for a given admixed individual. **(B)** Masked genotype showing ditypes of Native American (red), European (blue) and African (green) ancestry.

Gaps are represented in grey and regions with unassigned ancestry (Unknown) are
in black.

S6 Fig. Choice of the number of principal components from European
ancestry-specific principal component analysis.
Elbow method to determine which PC minimizes the angle of the curve from the
chart "Percentage of variance explained *versus* Number of PCs"

S7 Fig. European ancestry specific admixture analysis.
(A) Cross-Validation scores for *K* from 2 to 10. **(B)** Admixture for *K=2*. **(C)**
Admixture for *K=3*. CYA: Cuyo Region; NEA: Northeastern Region, NWA:
Northwestern Region; PPA: Pampean Region; PTA: Patagonia Region.

S8 Fig. Comparision of different European ancestry proportions in South
America.
Comparison of ancestry proportion estimates from European Ancestry Specific
Admixture Analyses (*K=3*) among samples from the Argentina, Chile and Colombia
(A) Southeaster/Italian ancestry (light blue in **S7C Fig**). **(B)** Iberian ancestry
(turquoise in **S7C Fig**). **(C)** Northern Ancestry (dark blue in **S7C Fig**). *P*-value of the
Wilcoxon test for each pairwise comparison is shown.

S9 Fig. African ancestry-specific principal component analysis.
(A) Localization map of the 1685 reference samples with >99% of African ancestry.
(B-C) Principal Components performed using the African reference samples
(represented as in panel **A**), and South American samples masked for African
ancestry.

S10 Fig. African ancestry-specific admixture analysis.
(A) Cross-validation scores *K* from 2 to 10. **(B)** Admixture for *K=2*. **(C)** Admixture
for *K=3*. **(D)** Admixture plots for *K=4*. CYA: Cuyo Region; NEA: Northeastern Region,
NWA: Northwestern Region; PPA: Pampean Region; PTA: Patagonia Region.

S11 Fig. Correlation between eigenvectors and ancestry proportion estimates
from African ancestry specific analyses
Comparison of ancestry proportion estimates from Admixture model with *K=5* and
some Principal Components (PCs) in admixed samples from **DS5. (A)** PC3 vs
Western African ancestry proportions (yellow in **S10 Fig**). **(B)** PC3 vs Bantu-
influenced ancestry proportions (blue in **S10 Fig**). **(C)** PC4 vs Eastern African
ancestry proportions (orange in **S10 Fig**).

S12 Fig. Choice of the number of principal components from Native American
ancestry-specific principal component analysis.
Elbow method to determine which PC minimizes the angle of the curve from the
chart "Percentage of variance explained *versus* Number of PCs"

S13 Fig. Native American ancestry-specific admixture analysis.

(A) Cross-validation scores *K* from 2 to 10. **(B)** Admixture for *K=2*. **(C)** Admixture for *K=4*. **(D)** Admixture plots for *K=5*. CYA: Cuyo Region; NEA: Northeastern Region, NWA: Northwestern Region; PPA: Pampean Region; PTA: Patagonia Region.

S14 Fig. Comparison of Native American ancestry proportion estimates obtained with admixture on unmasked data (*K=8)* and masked data (*K=3*).
Unmasked and masked data refers to **DS1** and **DS6**, respectively. Spearman correlation coefficients and associated *P*-values are shown. CAN: Central Andes; STF: Subtropical and Tropical Forests; CCP: Central Chile / Patagonia.

S15 Fig. Correlation of Native American ancestry proportions and geographic coordinates in Argentina
(A) Central Andes ancestry proportions vs Latitude. **(B)** Central Andes ancestry proportions vs Longitude. **(C)** Subtropical and Tropical Forests ancestry proportions vs Latitude. **(D)** Subtropical and Tropical Forests ancestry proportions vs Longitude. **(E)** Central Chile/Patagonia ancestry proportions vs Latitude. **(F)** Central Chile/Patagonia ancestry proportions vs Longitude. Linear regression slopes and the associated *P*-values are shown.

S16 Fig. Individual assignation to a Native American ancestry cluster.
Consensus cluster assignment of South American individuals based on three *K-means* procedures run with different pairwise distances among individuals. **(Top)** *K-means* results using Ancestry-Specific PCA and Admixture (ASPCA and AS-Admixture), and *f₃* results to compute pairwise distances. Individuals are represented as in **Main Fig 5**. Insets: BIC score for number of clusters set to *K-means* ranging from 2 to 20. In all the three cases, *K*-means BIC was minimized when considering 4 clusters. **(Bottom)** Same as top with point colors corresponding to the assigned cluster.

S17 Fig. Pairwise genetic affinity among individuals assigned to different groups.
Boxplots for *1- f₃(YRI; Ind1, Ind2),* where *Ind1* and *Ind2* are two individuals belonging to Group 1 and Group 2, respectively. The groups are either the fourth Native American components identified or ancient Middle Holocene Southern Cone groups. For clarity, boxplot outliers are not shown. YRI: Yoruba from 1KGP.

S18 Fig. Graphical visualization of pairwise genetic distances among modern and ancient groups in South America
(A) Neighbor-joining tree from distances of the form 1/*f₃(YRI; X, Y).* USR1 from Ancient Beringia was used as *outgroup* **(B)** Multidimensional-scaling from distances of the form 1-*f₃(YRI; X, Y).* Each group is represented as appearing in the leaf of (A). USR1 and Anzick-1 were not considered in (B). YRI: Yoruba from 1KGP

S19 Fig. Genetic affinity of the four Native American components with ancient groups.
(A-D) *f₃(YRI; X; Ancient).* **(*E-H)* *f₄(YRI, X; Ancient Beringia, Ancient).* **(A)** and **(E)**: with Central Andes (CAN) as *X*. **(B)** and **(F)**: With Subtropical and Tropical Forests (STF)

as *X*. **(C)** and **(G)**: With Central Chile / Patagonia (CCP) as *X*. **(D)** and **(H)**: With

Central Western Argentina (CWA) as *X*. YRI: Yoruba from 1KGP; *Ancient Beringia*:

USR1 individual from [66]; *X*: Native American component in Argentina (one plot

per *X). Ancient*: ancient group labeled on the *x*-axis and represented with a

point/color scheme as in **Main Fig 5**. Vertical segments are the +/- 3 standard

errors intervals.

S20 Fig. Changes across time of genetic affinity of the four Native American

components with ancient groups.

Each point represents a f_4 score of the form f_4*(YRI, X; Ancient Beringia, Ancient)* vs

the age of ancient sample, where *X* is one of the four identified Native American

components, and *Ancient* is an ancient group. *X* is represented by the color of the

square while *Ancient* is represented by the point within the square. The point code

of the ancient samples is shown in **Main Fig 5**. *Ancient Beringia*: USR1 individual

from [66]. **(A)** f_4 vs age of ancient samples from Southern Cone. **(B)** f_4 vs age of

ancient samples from Andes. **(C)** f_4 vs age of ancient samples from Southern Cone

considering correction for both f_4 *and* age. **(D)** f_4 vs age of ancient samples from

Andes considering correction for both f_4 and age. Linear regression slopes and the

associated *P*-values are shown. CAN: Central Andes; STF: Subtropical and Tropical

Forests; CCP: Central Chile / Patagonia; CWA: Central Western Argentina; YRI:

Yoruba from 1KGP

S21 Fig. Comparison of genetic affinity of an ancient group to a Native

American component relative to another.

f_4 *(*YRI*, Ancient, X, Y)* where *X* and *Y* are two of the four identified Native American

components (one plot per *X-Y* combination), and *Ancient* is ancient group labeled on

the *x*-axis and represented with a point/color scheme as in **Main Fig 5**. Vertical

segments are the +/- 3 standard errors intervals. Note this setting for f_4 statistics is

symmetrical when switching *X* and *Y.*

S22 Fig. Comparison of genetic affinity of a Native American component to

another relative to an ancient group.

f_4*(Ancient, X; Y, YRI)* where *X* and *Y* are two of the four identified Native American

components (one plot per *X-Y* combination), and *Ancient* is ancient group labeled on

the *x*-axis and represented with a point/color scheme as in **Main Fig 5**. Vertical

segments are the +/- 3 standard errors intervals. CAN: Central Andes; STF:

Subtropical and Tropical Forests; CCP: Central Chile / Patagonia; CWA: Central

Western Argentina; YRI: Yoruba from 1KGP

S23 Fig. Removing admixed Santiago de Chile individuals to compute *F*-

statistics does not affect the results.

Admixed individuals from Santiago de Chile were removed to perform the analyses

presented in this figure. **(A)** f_3*(Target; S1, S2*) to test for treeness; **(B)** f_4*(YRI, Target;*

S1, S2) to test whether Target shares more ancestry with S1 or S2; **(C)** f_3*(YRI; CWA;*

Ancient); **(D)** f_4*(YRI, CWA; Ancient Beringia, Ancient)*; **(E)** f_4*(Ancient, CCP; CWA, YRI)*;

(F) f_4*(Ancient, CWA; CCP YRI)*; **(G)** f_4*(YRI, Ancient; CWA, CCP).* CCP: Central Chile /

Patagonia.

S24 Fig. Schematic routes for the main population arrivals in the Southern
Cone.
Each arrow represents one of the four components discussed throughout the article.
Neither the time and place of the splits among these components nor gene flow
among them have been addressed in this study.

S25 Fig. Admixture analyses in DS2 to define European, African and Native
American reference individuals for local ancestry analyses
(A) Cross-validation scores from K=3 to K=10. **(B)** Admixture for $K=7$

S26 Fig. Admixture analyses in DS3 to define European, African and Native
American reference individuals for local ancestry analyses
(A) Cross-validation scores from K=3 to K=10. **(B)** Admixture for $K=5$

S27 Fig. Comparison of admixture and RFMix ancestry proportion estimates in
DS2p and DS3p.
(A-C) For DS2p: Argentinean samples from the present study with reference panel
that consists in 1KGP individuals from Africa, Europe and America [42] and Chilean
individuals from [37]. Native American, European and African ancestry proportions
estimates with RFMix vs with Admixture with $K=7$. **(D-F) For DS3p:** Argentinean
samples from [31] with reference panel that consists in 1KGP individuals from
Africa, Europe and America [42]. Native American, European and African ancestry
proportions estimates with RFMIx vs with Admixture with $K=5$.

S28 Fig. Consistency of the masking procedure applied to DS2p and DS3p.
We compared the percentage of variants with same ancestry ditypes in DS2p and
DS3p for American admixed individuals from the 1000 Genomes. Project.

Supplementary Table Captions
1370

S1 Table. Sample information.
Sampling location, gender, uniparental lineages, Affymetrix QC metrics, color and
point coding used for plots.

S2 Table. Data sets (DS) analyzed throughout the article.

S3 Table. Ancestry proportion estimates in a worldwide context.
Ancestry proportion estimates from Admixture analyses with $K=3$ and $K=8$ at the
worldwide level. The column names describe the labels attributed to each ancestry
detecting for both Admixture analyses, as well as the hexadecimal code for the color
used to represent it in the corresponding admixture plot. The columns "Point",
"Color" and "cex" list the graphical parameters used to represent each individual in
the different plots throughout the article.

S4 Table. Native American cluster assignation.
Individual Native American cluster assignation is given for each of the three *K*-means procedures and for the consensus call (columns "F3", "PCA", "Admixture" and "Consensus"). The ancestry proportion estimates from Admixture analyses with *K=3* on the masked data for Native American ancestry are also provided. The column names explicit the labels attributed to each ancestry detecting for both Admixture analyses as well as the hexadecimal code for the color. For admixed individuals (from the present study and [31]), the "Population" and "Region" columns list the locality and province, respectively, while for Native American population (from [37,38]) the "Population" and "Region" columns list the ethnic and main ethnic groups, respectively.

S5 Table. *F*-statistics
(A) f_3*(Target; S1, S2)* only for comparisons including Native American components **(B)** f_4*(YRI, Target; S1, S2)* only for comparisons including Native American components **(C)** f_3*(YRI; X, Y)* only for comparisons including Ancient Beringia, Mixe and Native American components **(D)** f_3*(YRI; X, Y)* where *X* and *Y* can be either an ancient group or one of the four Native American components. **(E)** f_4*(YRI, X; Ancient Beringia, Ancient)* only for comparisons including between a Native American components (*X*) and an ancient group (*Ancient*). **(F)** f_4*(YRI, Ancient, X, Y)* only for comparisons including two Native American components (*X, Y*) and an ancient group (*Ancient*). **(G)** f_4*(Ancient, X; Y, YRI)* only for comparisons including two Native American components (*X, Y*) and an ancient group (*Ancient*). YRI: Yoruba from 1KGP

Fig. 1. Sample ocations from the present study.

Fig. 2. Admixture analyses in a worldwide context.

Fig. 3. European ancestry specific principal component analysis.

Fig. 4. African ancestry-specific admixture analysis.

Fig. 5. Native American ancestry-specific principal component analysis.

Fig. 6. Native American ancestry-specific admixture analysis

Fig. 7. Relationship among the four Native American groups identified.

Fig. 8. Changes across time of genetic affinity of the four Native American with ancient samples.