TITLE

Comparing DNA replication programs reveals large timing shifts at centromeres of endocycling cells in maize roots

SHORT TITLE

Replication timing shifts at centromeres of endocycling cells in maize roots

Authors:

Emily E. Wear[1][*], Jawon Song[2], Gregory J. Zynda[2], Leigh Mickelson-Young[1], Chantal LeBlanc[3][#a], Tae-Jin Lee[1][#b], David O. Deppong[1], George C. Allen[4], Robert A. Martienssen[3], Matthew W. Vaughn[2], Linda Hanley-Bowdoin[1], William F. Thompson[1]

Affiliations:
[1] Department of Plant and Microbial Biology, North Carolina State University, Raleigh, North Carolina, United States of America

[2] Texas Advanced Computing Center, University of Texas, Austin, Texas, United States of America

[3] Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, United States of America

[4] Department of Horticultural Science, North Carolina State University, Raleigh, North Carolina, United States of America

[#a] Current Address: Department of Molecular, Cellular and Developmental Biology, Yale University, New Haven, Connecticut, United States of America

[#b] Current Address: Syngenta Crop Protection, Research Triangle Park, North Carolina, United States of America

* Corresponding author
E-mail: emily_wear@ncsu.edu

ABSTRACT

Plant cells undergo two types of cell cycles – the mitotic cycle in which DNA replication is coupled to mitosis, and the endocycle in which DNA replication occurs in the absence of cell division. To investigate DNA replication programs in these two types of cell cycles, we pulse labeled intact root tips of maize (*Zea mays*) with 5-ethynyl-2'-deoxyuridine (EdU) and used flow sorting of nuclei to examine DNA replication timing (RT) during the transition from a mitotic cycle to an endocycle. Here, we compare sequence-based RT profiles and found that most regions of the maize genome replicate at the same time during S phase in mitotic and endocycling cells, despite the need to replicate twice as much DNA in the endocycle. However, regions collectively corresponding to 2% of the genome displayed significant changes in timing between the two types of cell cycles. The majority of these regions are small, with a median size of 135 kb, and shift to a later RT in the endocycle. However, we found larger regions that shifted RT in centromeres of seven of the ten maize chromosomes. These regions covered the majority of the previously defined functional centromere in each case, which are ~1–2 Mb in size in the reference genome. They replicate mainly during mid S phase in mitotic cells, but primarily in late S phase of the endocycle. Strikingly, the immediately adjacent pericentromere sequences are primarily late replicating in both cell cycles. Analysis of CENH3 enrichment levels in nuclei of different ploidies suggested that there is only a partial replacement of CENH3 nucleosomes after endocycle replication is complete. The shift to later replication of centromeres and reduced CENH3 enrichment after endocycle replication is consistent with the hypothesis that centromeres are being inactivated as their function is no longer needed.

AUTHOR SUMMARY

In traditional cell division, or mitosis, a cell's genetic material is duplicated and then split

between two daughter cells. In contrast, in some specialized cell types, the DNA is duplicated a

second time without an intervening division step, resulting in cells that carry twice as much DNA

– a phenomenon called an endocycle, which is common during plant development. At each step,

DNA replication follows an ordered program, in which highly compacted DNA is unraveled and

replicated in sections at different times during the synthesis (S) phase. In plants, it is unclear

whether traditional and endocycle programs are the same. Using root tips of maize, we found a

small portion of the genome whose replication in the endocycle is shifted in time, usually to later

in S phase. Some of these regions are scattered around the genome, and mostly coincide with

active genes. However, the most prominent shifts occur in centromeres. This location is

noteworthy because centromeres orchestrate the process of separating duplicated chromosomes

into daughter cells, a function that is not needed in the endocycle. Our observation that

centromeres replicate later in the endocycle suggests there is an important link between the time

of replication and the function of centromeres.

INTRODUCTION

Developmentally programmed DNA replication without nuclear breakdown, chromosome condensation or cell division, a phenomenon known as endoreduplication or endocycling, occurs in a wide variety of plants and animals [1-3]. In plants, endoreduplication is a systemic feature [4] and often an important step in the development of tissues and organs such as fruit, endosperm, leaf epidermal cells, and trichomes [5]. Initiation of endocycling is frequently associated with a transition from cell proliferation to cell differentiation and expansion [6]. In plant roots, cells at the tip divide actively by normal mitosis, while endocycling cells become frequent further from the tip, in a zone associated with differentiation and increases in cell size [7, 8].

We developed a system to analyze DNA replication in *Zea mays* (maize) roots [8, 9], with similar approaches being applied in our work with *Arabidopsis* cell suspensions [10]. In this system, newly replicated DNA is labeled *in vivo* with the thymidine analog, 5-ethynyl-2'-deoxyuridine (EdU), and labeled nuclei are separated by flow cytometry into populations representing different stages of S phase. Cytological analysis showed that spatiotemporal features of maize DNA replication are significantly different from those of animal cells [11]. We then characterized the replication timing (RT) program in mitotic cells of the apical 1-mm root segment [12], using a modified replication timing by sequencing protocol (Repli-seq) [13, 14]. In mitotic cells, we found evidence for a gradient of early replicating, open chromatin that transitions gradually into less open and less transcriptionally active chromatin replicating in mid S phase. We also confirmed previous cytological observations showing that heavily compacted classical heterochromatin, including knobs and pericentromeres, replicate primarily in late S phase [11, 15]. While these relationships between RT and chromatin packaging are generally

similar to those found in other systems, we did not find evidence for megabase-scale replication

domains that have been characterized in mammalian cells (reviewed in [16] and references

therein).

Although replication in the first 1-mm of the root is mostly mitotic, with DNA contents

of labeled nuclei ranging from 2C to 4C, flow cytometry profiles of nuclei derived from root

tissue between 1 and 3 mm from the tip also included a substantial population of labeled nuclei

from endocycling cells, with DNA contents between 4C and 8C. Cytological analysis showed

that the spatiotemporal patterns of replication in endocycling nuclei are very similar to those in

mitotic nuclei [11]. However, it remained to be determined whether the entire genome is

uniformly replicated during the endocycle, and whether the temporal program is altered when

replication occurs without an intervening mitosis.

Both under-replication and over-replication (amplification) have been observed in

multiple animal systems, notably including *Drosophila* (reviewed in [17]). In addition to the

well-known amplification of chorion genes and under-replication of heterochromatin, under-

replication also occurs in a number of euchromatic regions, with a degree of tissue specificity

suggesting a possible role in differentiation [18-20].

Even though endopolyploidy is common in plants, there are very few reports of over- or

under-replication of specific sequences. Some orchids exhibit a phenomenon in which only a

fraction of the genome is endoreplicated [21, 22], but in most cases, endopolyploid cells have

DNA contents that are multiples of the 2C value. Both highly repetitive heterochromatic regions

and highly expressed genes are extensively endoreduplicated in maize endosperm nuclei, as

would be expected for uniform replication of the entire genome [23]. More definitively, whole

genome sequencing in *Arabidopsis* showed that leaf nuclear DNA is evenly endoreduplicated in

wild-type plants, although the same series of experiments clearly demonstrated selective over-

replication in *atxr5* and *atxr6* mutants [24].

In addition, there is as yet no information as to whether changes in RT programs are

associated with endoreduplication or differentiation in plant systems. That such changes might

occur in association with differentiation is supported by reports of extensive changes in RT

between animal cell cultures representing different embryonic or differentiated cell types (e.g.

[13, 25-27]).

To address these questions in the maize root tip system, we carried out a detailed

comparison of RT dynamics in mitotic and endocycling cells. To isolate endocycling nuclei, we

focused on a root segment 1–3 mm from the apex where there is a higher proportion of

endocycling cells and used flow cytometry to separate nuclei of higher ploidy. We found very

little evidence for changes in copy number that would be associated with over- or under-

replication, and the RT profiles for the vast majority of the genome are very similar. However,

we found significant changes in timing for a number of loci that together correspond to 2% of the

genome. Most notably, we found major changes in the RT of centromeres, which replicate

mainly during mid S phase in mitotic cells, but primarily in late S phase of the endocycle.

RESULTS

Separating endocycling from mitotic nuclei

As reported previously and described in Methods, we used a 20-min pulse of the thymidine

analog, EdU, to label newly replicated DNA in intact maize roots. This was followed by

formaldehyde fixation and isolation of nuclei from defined segments of root tips (Fig 1A).

Incorporated EdU was conjugated with Alexa Fluor 488 (AF-488) by "click" chemistry [28]. The

nuclei were then stained with DAPI and fractionated by two-color fluorescence activated flow

sorting to generate populations at different stages of the mitotic cell cycle or the endocycle [8, 9].

Fig 1B and 1C show flow cytometry profiles obtained for root segments 0–1 mm and 1–3 mm

from the tip, respectively. Fluorescent signals from nuclei that incorporated EdU during S phase

of a normal mitosis form an "arc" between 2C and 4C DNA contents, while nuclei labeled

during the endocycle S phase form a similar arc between 4C and 8C. As seen in Fig 1C, the

endocycle arc is more prominent in nuclei preparations from 1–3 mm root segments. To analyze

endocycle RT, which we will describe in detail below, we separated labeled nuclei representing

early, mid, and late S-phase fractions using the sorting gates shown in Fig 1C, adjusting the

endocycle early gate to avoid contamination with mitotic nuclei in late S phase. Reanalysis of the

sorted nuclei confirmed that there was good separation between the nuclei populations from the

adjusted early sorting gate and the mid sorting gate (S1 Fig). The flexibility of the EdU labeling

and flow sorting system also allowed us to collect unlabeled nuclei, representing non S-phase

cells with 2C, 4C and 8C DNA contents. These nuclei were used to characterize selected histone

marks following mitotic or endocycle replication and to investigate the copy number of

individual loci across the genome.

Fig 1. Global comparison of mitotic cycle and endocycle replication timing programs.

(**A**) Schematic of a maize root showing the meristem zone (0–1 mm region) and transition zone

(1–3 mm region) used for replication timing experiments. (**B and C**) Flow cytograms of nuclei

isolated from the 0–1 mm root segments (**B**) and 1–3 mm root segments (**C**). Dots are pseudo-

colored by density and black rectangles represent the sorting gates used to collect the pre-

replicative 2C reference sample and early (E), mid (M) and late (L) S-phase fractions from either

the mitotic cycle or endocycle. (**D**) Global scale view of replication timing (RT) for chromosome

10, comparing mitotic and endocycling profiles in early, mid and late S phase. Uniquely

mapping reads were aggregated in 3-kb windows, normalized for sequencing depth, divided by

the normalized 2C reference read counts, and Haar wavelet smoothed (see Methods). The global

RT profiles for mitotic and endocycling cells are very similar to each other for all ten

chromosomes. The schematic of chromosome 10 at the bottom shows the location of the

centromere (black oval) and the 10 Mb region that is expanded in panel **E** (red rectangle). (**E**)

Expanded view of a 10 Mb region on chromosome 10 with overlaid mitotic and endocycle RT

profiles. Unmappable or multi-mapping regions ("blacklist") are indicated as tick marks in the

bottom track. This example illustrates the similarity between the mitotic and endocycle RT

profiles that is observed throughout most of the genome. Scale for all panels: 0–5 normalized

signal ratio.

Evidence for complete genome replication during the endocycle

Given the well documented examples of over- and under-replication during the endocycle in

animal systems, we investigated whether there are local copy number differences in the maize

genome after endocycle replication. To do this, we used the non S-phase 2C, 4C, and 8C nuclei

populations described above, and carried out whole genome paired-end sequencing. To gain a

better representation of the copy number of repeat regions in the genome, reads that could not be

uniquely mapped to a single location were included, but we retained only the primary alignment

location for each read pair. These data were examined for regions in which normalized read

frequencies in 5-kb windows differed between 8C and 4C or 4C and 2C nuclei, using procedures

described by Yarosh et al. ([29]; S1 Text). We found about 5% of the 5-kb windows had ratio

values that fell outside of two standard deviations of the mean ratio for 8C and 4C or 4C and 2C

(1.0 ± 0.2 S. D. for both; S2A and B Fig). However, these windows all either occurred as

singleton 5-kb windows scattered around the genome (S2C Fig) or coincided with regions that

had very low read mapping in the 2C sample, indicating they are likely the spurious result of

making a ratio between windows with very few reads in both samples. As such, there is very

little evidence of meaningful over- or under-replication of genomic regions in nuclei with

different ploidy levels.

To further investigate whether there is complete replication of high-copy repeats that are

not well represented in the genome assembly, we used BLAST software to query all reads, not

just those that can be mapped to the genome, to determine the percentage of reads corresponding

to each of several consensus sequences for high-copy repeats (S1 Text). Analyzed sequences

included the knob repeats *knob180* and *TR-1* [30, 31], 5S and 45S rDNA repeats [32], and

centromere-associated *CentC* satellite repeats [33]. We also queried consensus sequences for

centromere retrotransposons of maize (*CRM*) families 1–4 [34-37]. In all cases, we found the

percentages to be similar in the 2C, 4C and 8C samples (S2D and E Fig), further suggesting that

there is little or no over- or under-replication.

Replication timing analysis

As described above, we sorted endocycling nuclei from the S-phase populations in Fig 1C, and

extracted and sheared the DNA in each fraction. EdU-containing DNA fragments were

immunoprecipitated with an antibody to AF-488, resulting in sequence populations representing

DNA replicating during early, middle, or late S phase of the endocycle. We also prepared DNA

from the unlabeled 2C nuclei pool to provide a reference dataset representing pre-replicative

nuclei. DNA from three biological replicates of each sample was sequenced to generate paired-

end reads.

To compare the RT programs in endocycling and mitotic nuclei, we mapped our previous

Repli-seq data for mitotic nuclei [12] and our new data for endocycling nuclei to the new maize

B73 RefGen_v4 genome, which includes improved assemblies of centromeres and more

complete annotations of transposable elements (TEs) [38, 39]. Uniquely mapped read depth

varied between ~3 and 11× genome coverage per S-phase sample, so all samples were randomly

downsampled to ~3× coverage to ensure comparable results (see Methods and S1 Spreadsheet).

We used the *Repliscan* analysis pipeline [14] to generate profiles of replication activity in

early, mid and late fractions of each S phase. These profiles were generated by aggregating the

Repli-seq read densities for each S-phase sample in 3-kb static windows, scaling the reads to 1×

genome coverage, and then dividing by the scaled read counts from the unlabeled 2C reference

data and smoothing by Haar wavelet transform (see Methods and [14]). Normalizing with the 2C

reference corrected for differences in sequencing efficiencies and collapsed repeats that caused

"spikes" in the data (illustrated for late replication in the endocycle in S3 Fig), producing an

estimate of replication intensity or "signal" in each 3-kb window. We also excluded 3-kb

windows with extremely low read coverage in the 2C reference sample (see Methods) from all

analyses ("blacklist" windows, indicated by black tick marks in Fig 1E).

Fig 1D shows that the global RT patterns are remarkably similar in endocycling and

mitotic nuclei, and overlays of the corresponding profiles show mostly minor differences (Fig

1E). Pearson's correlation coefficient values between corresponding S-phase fractions from the

mitotic and endocycle data are very high (r values of 0.91, 0.89 and 0.96 for early, mid and late,

respectively). These values are similar to those found between individual biological replicates

within each sample (S4 Fig).

Identifying regions of altered timing

Despite the global similarity of the RT programs of mitotic and endocycling cells, there are

regions scattered around the maize genome that show a shift in RT. To identify timing

differences, we first calculated the difference in normalized replication signal between the

mitotic and endocycle data at each genomic location for the early, mid and late profiles

separately (S1 Table; S5 Fig). We then constrained our analysis by focusing only on regions

where there was an equal and opposite timing difference in at least one other S-phase fraction

(for example, regions in which a decrease in early replication signal in endocycling cells was

associated with a corresponding increase in mid and/or late S-phase signal at the same location).

We allowed a gap distance of 6 kb when searching for regions with timing differences to account

for small blacklist regions that break up larger regions of change. We found that 11% of the

genome showed a difference in timing of at least 10% of the total difference range for a given

profile (difference in replication signal ≥ 0.4; S1 Table), with an opposite timing difference at the

same threshold criterion at the identical location in another S phase profile. Many of these

regions are small, with the lower 50% of regions ranging in size from 3 kb to the median size of

33 kb (S2 Table), and it is not clear if such small alterations are biologically relevant.

To identify more robust differences, designated Regions of Altered Timing (RATs), we

identified regions in which the difference in replication signal was ≥ 25% of the total difference

range for a given profile (difference in replication signal ≥ 1.0; S1 Table), and which also met

the criterion of having an opposite difference in at least one other profile. To highlight larger and

contiguous regions of change, we included ≥ 10% regions that were adjacent to the original ≥

25% regions. However, RATs had to have at least one core region where the timing change was

at least 25% (S2 Table) to be included in our analysis. Representative ≥ 25% and ≥ 10% regions

are indicated by different shades of red and blue bars in Fig 2 (additional examples are in S6

Fig). Finally, we examined the profiles for the RATs in individual biological replicates to verify

there was good agreement between the replicates (Figs 2B and S6). By selecting only the most

robust RATs we excluded other regions where timing changes are less dramatic – for example

those indicated by dashed boxes in Fig 2. In such regions, the timing difference did not meet our

criteria of a ≥ 25% difference in signal (box 2 in Fig 2A) and/or there is not an equal and

opposite ("compensated") timing difference (box 3 in Fig 2A).

Fig 2. Identifying regions of altered timing.

(**A**) An example region (5 Mb) on chromosome 10 containing two robust Regions of Altered

Timing (RATs), indicated by boxes outlined with solid lines. The RAT in box 1 (red) shifts from

Earlier-to-Later, and the RAT in box 4 (blue) shifts from Later-to-Earlier. Dashed boxes denote

regions with some level of RT difference in which the magnitude of the difference did not meet

our ≥ 25% criterion (box 2), or in which the change in one S-phase fraction was not compensated

by an opposite change in at least one other S-phase fraction (box 3). Annotated genes (purple)

and unmappable or multi-mapping regions ("blacklist", black) are indicated as tick marks in the

bottom tracks. (**B**) The same chromosome region as in (**A**) with the individual biological

replicate profiles overlaid to demonstrate that RATs are not caused by local regions of technical

variation between replicates. Scale for panels **A** and **B**: 0–5 normalized signal ratio. (**C**) Boxplots

representing the distribution of RAT sizes in the three categories: Later-to-Earlier, Earlier-to-

Later, and a subset of Earlier-to-Later RATs found in functional centromeres (CEN) [38].

Boxplot whiskers represent 1.5 x interquartile range (IQR). The axis is broken to show two

values that are much higher than the others and correspond to large RATs in CEN 9 and CEN 10.

However, it is important to note that the sizes of CEN RATs are underestimated, because

centromeres contain variable numbers and sizes of blacklist regions, which break up what would

probably be long continuous RATs (see Fig 3).

Robust RATs fall into two categories, those where the strongest replication signal occurs

later in the mitotic cycle than it does in the endocycle ("Later-to-Earlier" shift), and those in

which the strongest signal occurs earlier in the mitotic cycle than in the endocycle ("Earlier-to-

Later" shift). In addition, we separately characterized a subset of the Earlier-to-Later RATs that

are located in functional centromeres ("Earlier-to-Later-CEN") using centromere (CEN)

coordinates from [38]. Our stringent criteria identified RATs comprising only about 2% of the

maize genome (Table 1), with the vast majority (1.7% of the genome) in the Earlier-to-Later

category. Non-CEN Later-to-Earlier and Earlier-to-Later RATs have similar size distributions,

with median sizes of 141 and 135 kb, respectively (Fig 2C and Table 1). All of the CEN RATs

fall into the Earlier-to-Later category and have a median size of 132 kb, similar to the non-CEN

RATs. It is important to note, however, that the sizes of CEN RATs are likely underestimated

because of numerous blacklist regions within the centromeres that break what are likely

continuous RATs into several smaller parts in our analysis. Even though maize centromeres are

remarkably well sequenced [38], they still contain some gaps and regions where reads cannot be

uniquely mapped in the current B73 RefGen_v4 genome assembly, as indicated by the black tick

marks in the bottom tracks of Fig 3A–3D.

304 **Table 1.**

RAT category	Count	Median size (kb)	Coverage (kb)	% of genome	RATs with gene (%)	RATs with expressed gene (%)
Later-to-Earlier	41	141	6,291	0.3	92.7	82.9
Earlier-to-Later	192	135	26,907	1.3	96.4 *	91.1 *
Earlier-to-Later-CEN	41	132	7,668	0.4	43.9	22.0

305 **Table 1. RAT summary table.**

306 A summary of the region count, median size, total genome coverage, and percentage of the entire

307 genome represented in each RAT category. The number of RATs that overlap genes or expressed

308 genes is also presented. Asterisks denote one RAT category in which the indicated percent

309 overlap was greater than expected by chance (permutation P value ≤ 0.001), estimated by

310 permutation analysis (see Methods and S7 Fig.).

311

312 **Fig 3. Large RATs correspond to functional centromeres.** Our analysis found large RATs,

313 sometimes broken by blacklist regions (black tick marks at the bottom of each panel) at each of

314 the seven "complex" maize centromeres. The remaining three "simple" centromeres (on

315 chromosomes 1, 6, and 7) showed various levels of timing differences that did not meet the

316 criteria for calling RATs in our initial analysis. (**A–D**) Each 5-Mb region shown contains early

317 (E), mid (M) and late (L) RT profiles with mitotic and endocycle data overlaid (scale: 0–5

318 normalized signal ratio). The difference in late replication signal profiles (endocycle minus

319 mitotic; labeled "L dRT") for windows where the difference was compensated by an equal and

320 opposite difference in the early and/or mid profiles is also shown. Late differences compensated

321 at the $\geq 10\%$ threshold (light red), and those compensated at the $\geq 25\%$ threshold (dark red) are

322 shown, but only regions that contained at least one $\geq 25\%$ shift were classified as robust RATs in

our initial analysis. Two examples of simple centromeres, CEN 1 (**A**) and CEN 6 (**B**), and two

examples of complex centromeres, CEN 9 (**C**) and CEN 10 (**D**) are presented. The black

arrowheads in panels **A–D** denote example regions with a peak of early replication signal within

or adjacent to the centromere (for other examples, see S12 Fig). Colored boxes below the RT

profiles denote Earlier-to-Later RATs (red) and the functional centromere (black; [38]).

Chromosome 9 contains two called CEN regions labeled 9a and 9b. The colored tick marks (see

legend for colors) correspond to elements of centromeric retrotransposons of maize (*CRM*)

families 1–4 [39], gene annotations in RefGen_v4 [38] and the locations of mappable *CentC*

satellite repeats [40]. Blacklist regions are indicated by black tick marks in the lowest track. (**E**

and F) Timing differences (endocycle - mitotic) between late profiles for each centromere (**E**)

and corresponding pericentromere (**F**; ± 1 Mb) were calculated in 100-kb static windows. In

panel **F**, asterisks indicate difference values from windows where an Earlier-to-Later-CEN RAT

extends past the called CEN boundary [38] into the pericentromere; open circles indicate

windows that contain a non-CEN Earlier-to-Later RAT that met our compensation criteria.

Timing differences between early and mid profiles are shown in S13 Fig.

Non-centromeric RATs

We analyzed the non-CEN RATs for the content of genes and TEs, as well as the presence of

histone modifications and functional annotations related to the genes within RATs. To assess

whether the percentage of RATs containing genes differed from random expectation, we

randomly shuffled coordinates corresponding to the non-CEN Later-to-Earlier and Earlier-to-

Later RATs around the genome 1000 times and calculated the percentage of regions that overlap

genes in each set. We found that 93% and 96% of Later-to-Earlier and Earlier-to-Later RATs,

respectively, contain at least one annotated gene and usually contain a small cluster of genes

(Tables 1 and S3). Using root-tip RNA-seq data that are not specific to mitotic or endocycle

cells, we found that although only 50% of the 682 genes found in non-CEN RATs are expressed

at a meaningful level (FPKM ≥ 1; S3 Table), 83% and 91% of Later-to-Earlier and Earlier-to-

Later RATs, respectively, contain at least one expressed gene (Table 1). The observed percent

overlap of Earlier-to-Later RATs with genes and expressed genes are both significantly greater

than expected by random chance (permutation P value ≤ 0.001; S7B and D Fig). Differences

from random expectation were less obvious for Later-to-Earlier RATs, although the percent

overlap of expressed genes is on the edge of significance (permutation P value = 0.035; S7C

Fig).

We were unable to directly compare expression of genes in RATs in mitotic and

endocycling cells because we could not obtain RNA of sufficient quality to sequence from fixed,

sorted nuclei. Instead, we assessed a selection of gene-associated histone post-translational

modifications in sorted non S-phase 2C, 4C and 8C nuclei. In our previous work in maize root

mitotic cells, we showed that trimethylation of H3 lysine 4 (H3K4me3) and acetylation of H3

lysine 56 (H3K56ac) modifications tend to colocalize on active genes and are associated with

earlier replicating regions, while trimethylation of H3 lysine 27 (H3K27me3) tends to be on

repressed genes regardless of their RT [12]. For each ploidy level, we quantified the percentage

of genes within RATs that have each mark, as well as the fold enrichment relative to input for

called peaks within genes. There are very few differences between ploidy levels in the number of

genes bearing these marks (S8D Fig), but there are some minor shifts in the peak enrichment in

8C nuclei compared to 2C (S8A–C Fig). The clearest shift is a decrease in H3K4me3 enrichment

found on expressed genes in Earlier-to-Later RATs (S8B Fig), which suggests these genes may

have decreased expression in endocycling cells.

We also performed a gene ontology (GO) analysis for the genes found in non-CEN RATs

to ask if there are functional annotations enriched in genes that shift replication timing. For this

analysis, we focused on the genes that we identified as expressed in the root tip (S2 Spreadsheet).

We found 44 significantly enriched GO terms for genes within Earlier-to-Later RATs, including

biological process and molecular function terms related to gene expression, DNA/RNA

metabolism, and the cell cycle (S9 Fig). A wide variety of significant cellular component GO

terms were also found, which may relate to various differentiation processes occurring in

endocycling cells. There are no significant GO terms for genes within Later-to-Earlier RATs,

though the presence of only 52 expressed genes in this RAT category made it difficult to fully

assess significance. Taken together, these analyses of transcription-related histone modifications

and functional annotations suggest a role for gene expression changes in the Earlier-to-Later

RATs. Given that these regions are shifting to a later RT in the endocycle, a decrease in gene

expression would be expected [12]. Clearly, however, more work will be needed to confirm this

hypothesis.

The general organization of the maize genome is genes clustered in "islands" interspersed

with blocks of transposable elements [41-43]. We used a similar permutation strategy as for the

genes to estimate the significance of any differences in percent coverage of each TE superfamily

in non-CEN RATs as compared to random expectation, estimated from 1000 randomly shuffled

sets. The TE annotations were from the recent RefGen_v4 TEv2 disjoined annotation, where

every bp is assigned to a single TE [39]. We found the coverage of the RLG/Gypsy superfamily

in Earlier-to-Later RATs is significantly less than random expectation (permutation P value $\leq$

0.001; S4 Table). There are other, less significant, positive and negative associations with TE

superfamilies in non-CEN RATs, including RLC/Copia, DTT/Tc1-Mariner, DTM/Mutator and

DHH/Helitron (S4 Table). We also found that the percent AT content in RATs is similar to that

of the genome as a whole, with median values of 55% and 56% for Later-to-Earlier and Earlier-

to-Later RATs, respectively, and a median value of 55% for the whole genome (S10 Fig).

Centromeric RATs

Functional centromeres are defined by their content of nucleosomes containing the centromere-

specific histone variant known as CENH3 in plants and CENP-A in animals. CENH3/CENP-A

makes up only a small percentage of the total H3 population in centromeres, but plays an

important role in recruiting kinetochore proteins [44-46]. Maize is unusual among higher

eukaryotes in that a majority of centromeric reads can be uniquely mapped [47]. In our

replication timing data, for example, we found that on average 45% of all reads that map to

centromeres could be uniquely mapped to a single location (S11 Fig). Only these uniquely

mapping reads were used for further analysis. In addition, most of the maize centromere

assemblies are relatively intact, and functional centromeres have been located by mapping ChIP-

seq reads for CENH3 [38]. When combined with our replication timing data, these features of

the maize system create a unique opportunity to assess RT programs for centromeres.

Our analysis found large, robust RATs across seven of the ten centromeres (Figs 3C, 3D

and S12), with replication occurring mainly in mid S in mitotic cells, but changing to primarily

late S in endocycling cells. It is also noteworthy that though replication occurs mainly in mid S

in mitotic cells, there are some distinct peaks of early replication inside or directly adjacent to the

called centromere (indicated by black arrowheads in Fig 3 and S12) in all but one of the maize

centromeres. These early peaks remain in the endocycle, though in some cases there is a

reduction in early signal with a concomitant increase in mid signal at the same location. The

seven centromeres that contain robust RATs (CEN 2, 3, 4, 5, 8, 9 and 10) were previously

classified as "complex" because they contain a mixture of retrotransposons with some

centromere satellite repeat arrays (*CentC*; [40, 47]). In the RefGen_v4 genome assembly, CEN 9

has two called CENH3-binding regions [38], which we refer to as CEN 9a and 9b (Fig 3C; black

bars). Interestingly, we only found a robust RAT in the larger CEN 9a, with the smaller CEN 9b

showing almost no timing shift.

The remaining three centromeres (CEN 1, 6, and 7) were previously characterized as

"simple" because they mainly contain large arrays of the *CentC* repeat [40, 47]. In our analysis,

the simple centromeres showed, at most, small timing shifts that did not meet our criteria for a

robust RAT (Figs 3A, 3B and S12). However, *CentC* repeats are not well represented in the

reference genome assembly, so our ability to analyze replication of the complete simple

centromeres is limited. Portions of CEN 7 that are present in the assembly replicate mainly in

mid S phase in both mitotic and endocycling cells (S12 Fig), while sequences in the assemblies

for CEN 1 and CEN 6 are mostly late replicating in both types of cells, with some minor timing

changes across small regions (Fig 3A and 3B).

The robust RATs on the seven complex centromeres correspond quite closely to the

boundaries of the functional centromeres defined from CENH3 ChIP-seq data [38]. The

cumulative coverage of RATs in each complex centromere ranges from 405–1518 kb (S5 Table).

However, because each centromere includes blacklist regions that vary in size and number,

automated analysis did not identify the true sizes of the RATs. To avoid this problem, we have

chosen to focus the following analyses on the entire functional centromere instead of on

computationally identified RATs.

For the entire CENH3-binding region of each chromosome (excluding blacklist regions),

we calculated the difference in early, mid and late replication signal (endocycle minus mitotic)

from RT profiles by averaging across 100-kb static windows. For comparison, we also calculated

the replication signal differences in pericentromeres, which were arbitrarily defined as the ± 1

Mb flanking the CENH3 region. We inspected all RT differences in the centromeres and

pericentromeres by not requiring that the RT differences be compensated by an opposite shift in

the other S-phase fractions. Early and mid replication signals across the complex centromeres

decrease and late replication signals increase in endocycling cells, reflecting a large shift toward

late replication. The RT difference values for the late profile in centromeres and pericentromeres

are shown in Fig 3E and 3F, respectively, while the difference values for early and mid profiles

are shown in S13 Fig. Interestingly, the timing difference tapers off towards the edges of the

functional centromere (see profiles in Figs 3C, 3D and S12), and there is striking congruity in the

replication signals for mitotic and endocycling cells in the immediately adjacent pericentromere

regions (Fig 3A–D). The few timing shifts in pericentromeric regions are smaller in size and

much less dramatic than those in the centromere proper (Fig 3F). Moreover, very few (8%) of

pericentromeric windows with timing shifts are compensated by an equal and opposite shift in

the other S-phase profiles (S6 Table), suggesting many of these uncompensated differences may

result from technical variation rather than from meaningful biological differences. In contrast,

nearly all (85%) of the centromeric windows have compensated RT shifts.

Genomic elements and features in centromeres

Maize centromeres contain varying amounts of tandemly arrayed *CentC* repeats (single repeats

of 156 bp in length; [33]) as well as several *CRM* retrotransposon families interspersed with

elements from a few other retrotransposon families [36, 43, 48, 49]. *CentC* repeats and *CRM*

elements are also present in the adjacent pericentromeres where there is no CENH3 binding [43,

48]. In RefGen_v4, there are also fifty annotated genes within centromeres. We asked if all of

these sequence elements in centromeres behave similarly in the mitotic to endocycle transition,

or if certain elements show larger timing shifts than others. We also asked if all three types of

sequence elements show similar RT changes in centromeres versus pericentromeres. Given that

the RT signal values were aggregated in 3-kb windows, we only included elements that covered

at least half a window (1.5 kb) in our analysis. Fig 4 summarizes data on these questions for the

complex centromeres, while data for the simple centromeres are shown in S14 Fig. Similar

results were found when all elements were included (S14 Fig).

Fig 4. Comparing replication times for genomic features in complex centromeres and

corresponding pericentromeres. (**A–D**) Boxplots comparing replication signals during mitotic

and endocycle S phases for centromeres, pericentromeres (± 1 Mb), and genomic features within

them. The panels show the distributions of replication signals in early (E), mid (M), and late (L)

S for all 3-kb windows (**A**), annotated genes (**B**), mapped *CentC* repeats (**C**), and *CRM1/2*

elements (**D**) in centromeres and pericentromeres. For panels **A** and **C**, colored violin plots are

overlaid, while for panels **B** and **D**, individual data points are shown. Only elements that covered

at least 50% of a 3-kb window were included in each analysis, though results were similar when

all elements were included (S14 Fig). The number of windows or elements included in each

analysis is indicated above each graph. Boxplots for all elements in simple centromeres, as well

as for the individual *CRM1* and *CRM2* families are in S14 Fig.

The results for the two dominant *CRM* families, *CRM1* and *CRM2*, are similar (S14 Fig),

so these families were grouped together in Fig 4C. When present in centromeres, all three major

classes of elements – genes, *CRM1/2*, and *CentC* repeats – clearly replicate later during the

endocycle than in the mitotic cycle (Fig 4). In contrast, genes and *CRM* elements in the

pericentromere show little or no timing shifts. A full analysis of the replication times of *CentC*

repeats in pericentromeres is hampered by the limited representation of this repeat class in the

genome assembly (Fig 4D and S14E).

Chromatin features in centromeres

We also examined activating (H3K56ac and H3K4me3) and repressive (H3K27me3) histone

post-translational modifications to look for epigenetic changes in centromeres after endocycle

replication. It was previously reported that some H3K4me3 and H3K27me3 peaks of enrichment

occur in the centromere, mainly associated with genes [50]. We asked whether genes that have

these modifications continue to have them after mitotic and endocycle replication, and found

very few changes in the number of genes with these modifications at each ploidy level (S15 Fig).

There was also very little change in the fold enrichment of these histone marks in centromere

genes when comparing 2C, 4C and 8C nuclei.

We also investigated the levels of dimethylation of histone H3 lysine 9 (H3K9me2)

enrichment in each centromere. Previous work indicated there is a depletion of H3K9me2 in

centromeres relative to adjacent pericentromeres [51, 52], which we observed as well (S16 Fig).

Traditional peak calling tools are not effective for H3K9me2 because of its even distribution

across the maize genome. Instead, we estimated the fold enrichment by calculating the percent of

total H3K9me2 ChIP reads in a given centromere region (using coordinates from [38]) and

dividing by the percent of total input reads corresponding to that centromere in three biological

replicates). We found a similar H3K9me2 average fold enrichment for all centromeres and for

2C, 4C and 8C nuclei, although values for 4C and 8C nuclei were consistently slightly higher

than those for 2C nuclei (S16A Fig). CENH3 nucleosomes lack the lysine 9 residue found in

canonical histone H3 [53], so H3K9me2 enrichment must occur in the interspersed H3

nucleosomes.

Centromeric histone H3 in mitotic and endocycling centromeres

Unlike the canonical histone H3, CENH3 is not replaced in a replication dependent manner in

higher eukaryotes, resulting in a dilution of CENH3 relative to centromeric DNA during S phase

[54, 55]. New CENH3 is incorporated into nucleosomes after the completion of S phase, but the

timing of its integration into centromeric chromatin differs for plants, flies and humans

(reviewed in [56]). In the plants tested thus far, deposition of CENH3 has been reported to occur

between late G2 and metaphase [57-60].

Because mitosis does not occur in the endocycle and centromere function is presumably

not required, we speculated that CENH3 might remain at low levels following DNA replication

in endocycling cells. This hypothesis is supported by cytological studies of *Arabidopsis*

endopolyploid nuclei showing the CENH3 signal does not increase in parallel with the total

DNA content or the signal for 180-bp centromeric repeats [58, 59]. To test this hypothesis with

maize centromeres, we used a maize anti-CENH3 antibody [48] for ChIP-seq analysis of CENH3

binding in sorted non S-phase 2C, 4C, and 8C populations of nuclei. It is important to note that

the 4C nuclei come from a mixture of cells, some of which will return to the mitotic cycle and

others that will continue on to the endocycle (at least 13% of nuclei in the 1–3 mm region). We

asked whether the location or level of CENH3 enrichment changed after DNA replication in the

mitotic cycle or the endocycle. For visualization of CENH3 localization, ChIP-seq read counts

from three biological replicates for each ploidy level were aggregated in 3-kb windows and

normalized to the level of a uniform 1× genome coverage, so that corresponding windows in the

different ploidy level profiles were comparable. The normalized read count in each 3-kb window

was then divided by the corresponding normalized read count for the corresponding ploidy input

DNA to calculate a fold enrichment relative to DNA content value for CENH3 binding

sequences in that window. The spatial distribution of CENH3 enrichment across the centromeres

remained the same in 2C, 4C, and 8C cells. This is illustrated for CEN 9 and CEN 10 in Fig 5A

and 5B, and data for the rest of the centromeres are shown in S17 Fig. There are also a few small

spikes of CENH3 enrichment outside the called centromere (e.g. seen in Fig 5 and S17, but also

occasionally further out on the arms). These spikes also remain in the same location between 2C,

4C and 8C cells, some of which could be related to misassembly of the reference genome.

However, if real, these ectopic CENH3 peaks are less numerous and more persistent in G2 than

those recently observed in HeLa cells [61].

Fig 5. CENH3 localization and enrichment in mitotic and endocycling centromeres. We

profiled CENH3 binding by ChIP-seq in flow sorted, non S-phase nuclei with 2C (before mitotic

replication), 4C (after mitotic replication) and 8C (after endocycle replication) DNA contents. (**A**

and B) CENH3 localization patterns for 2C, 4C and 8C nuclei in CEN 9a and 9b (**A**) and CEN

10 (**B**). Scale in both panels is 0–120 fold CENH3 enrichment relative to input. Colored boxes

below the CENH3 profiles denote the previously identified functional centromere (black; [38]),

and Earlier-to-Later-CEN RATs (red). Tick marks in the bottom two tracks indicate blacklist

regions (black) and mapped *CentC* repeats (teal). (**C**) We used the ChIP-seq datasets from 2C,

4C and 8C nuclei to estimate the CENH3 average fold enrichment relative to DNA content for

complex centromeres by calculating the percent of total CENH3 reads found in a given

centromere (using coordinates from [38] and dividing by the percent of total input reads

corresponding to that centromere. Black dots represent the individual values from biological

replicates. Data for simple centromeres are shown in S17B Fig.

To compare total CENH3 content of entire centromeres at different ploidy levels, we

calculated the percent of total CENH3 reads found in a given centromere and made a ratio to the

percent of total reads from the corresponding input DNA in that centromere separately for each

biological replicate, as described above for H3K9me2. The CENH3 average fold enrichment

relative to total DNA content is similar for 2C and 4C nuclei in each of the complex centromeres

(Fig 5C), with an average 4C/2C enrichment ratio of 1.1 (S7 Table). However, CENH3

enrichment decreases with the increase in ploidy from 4C to 8C (Fig 5C), with an average 8C to

4C enrichment ratio of only 0.7 (S7 Table). Average CENH3 enrichment values for simple

centromeres were lower and slightly more variable, likely because of assembly issues. In both

cases, however, the ratio of CENH3 enrichment in 8C cells to that in 4C cells is clearly higher

than 0.5, which would be expected if there was no incorporation of new CENH3 after endocycle

replication, but smaller than the 1.0 ratio expected if there was full replacement (S7 Table). It is

worth noting that these data refer to post-replication 8C nuclei, which exited S phase prior to the

time of analysis, and that post-replication 4C nuclei show no dilution of CENH3 relative to DNA

content. Thus, our data are consistent with a model in which the CENH3 to DNA ratio is reduced

as DNA replicates during the endocycle S phase, and only partially restored after completion of

S phase.

DISCUSSION

The maize root tip includes a naturally occurring developmental gradient, with cells in the

meristem region (ca 0–1 mm) primarily undergoing mitotic cell cycles, while a subpopulation of

cells in the transition zone (ca 1–3 mm) enters a developmentally programmed endocycle prior to

further differentiation [8, 62]. Even though endocycling is very common in plants and plays

essential roles in differentiation and the development of specialized tissues, cell size increases,

and stress responses [2, 5, 63, 64], replication timing (RT) programs have not yet been

characterized for alternative cell cycles, such as the endocycle.

We generated whole genome Repli-seq data for root cell nuclei undergoing DNA

replication in either the mitotic cycle or the endocycle, making use of *in vivo* EdU labeling of

intact root tips and two-color fluorescence activated nuclei sorting. By doing so, we avoided

potential artefacts caused by cell synchronization [65] and chromosome aberrations often found

in plant and animal cell cultures (e.g. [66-68]). We present replication activity profiles for early,

mid and late replication separately, instead of collapsing the data into an early:late ratio as many

studies do. The rationale for this approach is that, for roughly one third of the maize genome, we

previously found heterogeneity in mitotic RT – e.g. regions of the genome in which root tip cells

exhibit significant replication activity in both early and mid S, or both mid and late S [12]. An

additional advantage to presenting the replication profiles separately is the ability to assess

whether there are concomitant or "compensated" changes in a region at multiple stages of S

phase. This compensation criterion helped us separate RT shifts that could be subject to technical

error, such as alterations in flow sorting gates, from shifts that are more likely to represent

meaningful changes in the population preference to replicate a replicon or cluster of replicons at

a particular time in S phase.

The current study sought to investigate whether the mitotic RT program is maintained in

the first round of the endocycle in maize root cells, despite the need to replicate twice as much

DNA and the initiation of various root cell differentiation pathways. Extending our previous

cytological observation that spatiotemporal patterns of replication are similar in mitotic and

endocycling cells [11], we found that RT programs at the sequence level are strikingly similar as

well. Pearson's correlation coefficient values comparing data from the two types of cell cycles

were similar to those for biological replicates within each type. The high level of reproducibility

is particularly noteworthy in the case of the early replication profiles, given that the flow sorting

gate for early replicating nuclei in the endocycle had to be adjusted to minimize contamination

from late replicating mitotic nuclei (Fig 1C). This overall conservation of RT programs suggests

that the process of re-establishing the RT program must be similar for the two types of cell

cycles in maize roots. In animal systems, re-establishment of the RT program has been shown to

occur in G1 of each cell cycle at a "timing decision point"[69], however the details of this

process have not been studied in plants.

Most plants fully replicate their genome during endocycles [70], although there are a few

exceptions (e.g. various orchid species; [21, 22]). We found very little evidence for over- or

under-replication occurring in endocycling maize root cells, unlike the distinctive over- and

under-replication found in *Drosophila* endocycles (reviewed in [17] and references therein). Our

result is consistent with earlier cytological reports that whole chromosomes, as well as repetitive

knobs and centromeres, are completely replicated in the highly endopolyploid maize endosperm

[23].

In contrast to the global maintenance of RT, we observed a small fraction of the maize

genome that exhibits some difference in RT between the two types of cell cycles. Approximately

11% of the genome showed compensated differences at a stringency level of $\geq$ 10% difference in

replication signal (see Methods). However, with the notable exception of centromeric regions,

which are discussed in more detail below, we chose to characterize only the most robust Regions

of Altered Timing (RATs), defined by the criteria of containing a core region with compensated

differences at a stringency level of $\geq$ 25% difference in replication signal. These robust non-

centromeric RATs comprise only 1.6% of the genome, and the size range of individual regions

(39–387 kb, median 138 kb) is consistent with our previous observation that regions of

coordinate replication in maize are ~50–300 kb in size [12]. This may include from one to a few

replicons, based on previous estimates of replicon size in monocot plants [71].

The first 1 mm of the maize root contains the meristem and precursors for at least ten

different cell types. Only some of these cell types enter the endocycle prior to cell elongation

[62]. If there are differences in the RT programs of different cell types, some or all of the non-

centromeric RATs may be associated with shifts in the relative contribution of different cell

types to the two samples of nuclei, rather than to endocycling *per se*. Research in metazoans has

revealed ~8-20% of their genomes can shift RT between cell types [13, 25, 26, 72-74]. In

mammals, these shifts generally involve large regions or "domains" in the megabase size range

(reviewed in [16]). These RT domains are much larger than the non-centromeric RATs in maize,

even though the maize genome is similar in size to the human and mouse genomes. However, in

the much smaller *Drosophila* genome, regions that show timing shifts between cell types are

more similar in size to the maize non-centromeric RATs [72, 74].

The vast majority of the non-centromeric RATs involved RT shifts from Earlier-to-Later,

with a significant enrichment for not only genes, but genes expressed in the root tip. This result

suggests the possibility that RT shifts may be related to shifts in gene expression. Unfortunately,

we have been unable to follow transcriptional changes in endocycling nuclei directly, as we have

as yet been unable to isolate RNA of sufficient quality to characterize transcripts from fixed

nuclei. However, our analysis of activating and repressive histone modifications uncovered only

minor changes in the enrichment and location of these marks within RAT genes after endocycle

replication. The lack of notable changes in the proportion of RAT genes bearing H3K56ac and

H3K4me3 modifications after the endocycle suggests that these histone marks are permissive to

changes in RT. Nonetheless, the direction of the change in H3K4me3 enrichment on genes in

Earlier-to-Later RATs after endocycle replication (S8B Fig) is consistent with the hypothesis that

a shift to later RT may accompany a decrease in gene expression. Many studies have identified a

correlation between RT and transcriptional activity (reviewed in [16]), but there are also multiple

examples of these processes being uncoupled (e.g. [27, 75]).

In the case of centromeres, it is easy to imagine that the large shifts to later replication are

related specifically to endocycling, because endocycling cells no longer require functional

centromeres. Though often broken by unmappable and multi-mapping ("blacklist") regions in the

genome assembly, when combined, centromeric RATs are much larger in size than the non-

centromeric RATs and cover the majority of each of the seven complex centromeres (S5 Table).

These seven centromeres, which are well assembled in the maize B73 RefGen_v4 genome,

contain satellite repeats interspersed with retrotransposons [38, 47], enabling almost 50% of our

sequencing reads that map to these centromeres to be uniquely positioned. In most species, in

which centromeres contain large numbers of tandemly arrayed satellite repeats, it is difficult to

map centromeric sequence reads to unique positions and, thus, to fully assess centromeric RT

patterns [76]. Though yeast centromeres replicate in early S phase [77-80], most higher

eukaryotes replicate centromeres asynchronously through mid to late S phase [54, 81-86]. Many

of the reports in higher eukaryotes are based on cytological observations, membrane

hybridization, or PCR data with limited resolution. Even a recent genomic analysis of

centromeric RT in human cell lines was significantly limited by the quality of the human

centromere assemblies, and could only uniquely map ~15% of centromeric reads [76].

Centromere replication in plant species, assessed mostly by cytological methods, has variously

been reported to occur in early, mid or late S [87-90], though it is often unclear if the analysis

was of sufficient resolution to distinguish the RT of centromeres from that of adjacent

pericentromeres. In contrast, we have provided a high-resolution analysis of the distribution of

replication times across maize centromeres, and compared RT of centromeres to adjacent

pericentromeres. These analyses revealed several features shared by the RT programs of the

seven complex maize centromeres. For example, in mitotic cells there are a few distinct peaks of

early replication (e.g. arrowheads in Figs 3 and S12), interspersed with mainly mid replication

activity that transitions to late replication at the edges of the functional centromere. In the

endocycle, entire centromeres – including regions with early and mid replication activity and the

genes, retroelements and *CentC* repeats within them – undergo a shift to later replication. As a

result, the RT of the complex centromeres in the endocycle becomes much more similar to that

of the immediately adjacent pericentromeric regions, which replicate primarily in late S phase in

both mitotic and endocycling cells.

The presence of distinct peaks of early replication in or adjacent to functional

centromeres (arrowheads in Fig 3 and S12) is noteworthy because they signify a population

preference for initiation in early S phase at these loci. This observation is of particular interest

because yeast centromeres contain a replication origin that is the first to initiate on its respective

chromosome and plays a role in centromere specification [80]. In maize, there is no evidence that

these early regions in centromeres are the first to replicate on the entire chromosome, but they

are earlier replicating than their surroundings. Origin mapping experiments (e.g. [91, 92]) would

be required to distinguish whether these early regions contain single or small clusters of origins,

and the location of any other origins in centromeres that may fire in mid or late S phase.

Unlike complex centromeres, the three simple centromeres of maize show less drastic

timing changes, that occur over smaller regions. These simple centromeres are not as well

assembled as the complex centromeres [40, 47], and we cannot assess RT for the possibly large

portions of these centromeres not present in the genome assembly. One potential interpretation of

our results is that the simple centromeres have distinct RT programs that show less timing shift

in the endocycle, possibly related to their different sequence composition. Alternatively, the

missing portions of the simple centromere assemblies could be replicating more like the complex

centromeres. Because simple centromeres are known to primarily contain large *CentC* arrays [40,

47], the second hypothesis is supported by our analysis of mapped *CentC* satellite repeats in all

centromeres, which showed that, as a group, these repeats consistently shift RT from mid to late.

Another piece of evidence comes from our analysis of complex centromeres, which showed that

the magnitude of the RT change tapers off toward the outer edges of the functional centromere.

One can speculate that the simple centromere assemblies are comprised mostly of the sequences

at the edges of the actual centromere, which would still be anchored to nonrepetitive regions in

the genome assembly. As in complex centromeres, these edge sequences might have a smaller

RT shift than internal sequences. Future cytological experiments, using a combination of flow

sorted EdU-labeled nuclei and techniques for identifying maize chromosomes [93, 94] could

help address questions related to the RT of simple centromeres.

713   The centromere-specific histone variant, CENH3 (also called CENP-A in animal

systems) plays an important role in recruiting kinetochore proteins [44-46]. In metazoans, it has

been shown that CENP-A is distributed among sister centromeres during replication, but the full

complement of new molecules is not redeposited until later [55, 95]. However, there are

differences in the timing of deposition of CENH3/CENP-A among eukaryotes. Deposition

occurs from S phase to G2 in yeasts, while in plants and protozoans it occurs from late G2 to

metaphase, and in metazoans it occurs mostly during G1 (with the exception of some *Drosophila*

cell types in metaphase to G1; reviewed in [46, 56, 60]). These interesting differences between

phylogenetic groups in the timing of CENH3/CENP-A deposition suggest there may also be

differences in the mechanisms and regulation of deposition that need to be explored further [59].

In our analysis of CENH3 enrichment relative to DNA content in maize root cells, the population

of 4C nuclei appear to have a full complement of CENH3, which would be consistent with the

previous results for plant species. This result suggests a model in which the sub-population of 4C

cells entering the endocycle also carry a full complement of CENH3. If that model is correct, our

data for 8C nuclei imply that CENH3 is only partially replaced after DNA replication in the

endocycle. Because the population of 8C nuclei we analyzed likely represents a mixture of cells

that recently exited endocycle S phase and others that exited some time ago we cannot determine

whether CENH3 will be fully restored in all cells at a later time. However, it is clear that the

ratio of CENH3 to DNA is not immediately restored, and the lower ratio is widely distributed

across all ten centromeres.

It is unlikely that endocycling cells will ever re-enter the mitotic cycle [1, 96, 97], and it

is not clear why endocycling cells would maintain or redeposit CENH3 nucleosomes at all unless

CENH3 has roles outside of mitotic cell division. A recent study in *Drosophila* midgut cells

found that CENP-A is required even in post-mitotic and differentiated cells, and proposed that

the loading of CENP-A in endocycling cells is essential for maintaining chromosome cohesion

[98]. This possibility has not yet been tested.

Centromeres are considered to be epigenetically specified, as there are no unique

sequences in the functional centromere that are not also found in the adjacent pericentromere

(e.g. reviewed in [44, 99, 100]). With this in mind, we tested whether changes in enrichment

levels of CENH3 nucleosomes, or several modifications to canonical H3 nucleosomes, could

explain the large shift to later replication of centromeres in endocycling cells. These studies only

uncovered very small changes in activating and repressive histone H3 modifications in

centromeres after endocycle replication. The magnitude of the change in CENH3, while

somewhat larger, was not on the scale of the change in RT. It is possible that the decrease in

dosage of CENH3 proteins has an effect on the recruitment of replication proteins, as has been

proposed in the yeast *Candida albicans* [80]. If replication proteins were not recruited as

efficiently, this could contribute to a delay in replication time of the centromere. It is also

possible that more significant changes might be found in epigenetic marks that we did not

investigate, for example changes in DNA methylation patterns or other histone post-translational

modifications. A variety of modifications to CENP-A nucleosomes have been identified,

(reviewed in [101]), but very little is known about CENH3 modifications in plants [102, 103],

highlighting an area for future research. Experiments in human cells identified cell cycle related

interchanges of acetylation, monomethylation and ubiquitination at the lysine 124 residue of

CENP-A [104, 105]. Mutations of this residue led to replication defects and alterations to

centromeric RT [105]. Another interesting question is whether changes in chromatin

conformation or 3D positioning in the nucleus are associated with the large shift in centromeric

RT. In mammals, RT is considered a functional readout of large-scale chromatin structure [16,

27, 73], and regions that shift RT have been shown to also change 3D localization [106].

Additionally, a study in mouse showed that when late replicating pericentric heterochromatin

was experimentally repositioned to the nuclear periphery, a location where mid replicating

chromatin is usually found in that system, the RT of those regions was advanced [107].

Investigating the interplay of chromatin environment, gene transcription and DNA

replication in plant systems, particularly in important crop species, has proven difficult in the

past. Numerous reasons for these difficulties exist, for example, plants have cell walls and are

rich in nucleases, actively dividing cells are sequestered in tiny meristematic regions, and many

genomes have a high content of retrotransposons and other repeats. As a result, understanding of

such critical areas has lagged behind that in yeast and animal systems. However, with recent

progress in assembling genomic resources and anticipated advances in the ability to isolate

individual cell types [108], perform sophisticated analyses of genome conformation [109, 110]

and follow individual chromosome regions using elegant cytological paints [94], the maize root

tip system is poised to contribute to rapid progress in these and many other important areas of

plant genome biology.

METHODS

Plant material

Seeds of *Zea mays* inbred line B73 (GRIN NPGS PI 550473) were germinated on damp paper towels and grown for three days. Seedling roots were labeled by immersion in sterile water containing 25 μM EdU (Life Technologies) for 20 min, using growth and experimental conditions described previously [8, 9, 12]. Biological replicate material was grown independently and harvested on different days. For the endocycle Repli-seq experiment, after rinsing roots well with sterile water, the 1–3 mm segments (Fig 1A) were excised from primary and seminal roots. The root segments were fixed, washed and snap-frozen as described previously [9].

Flow cytometry and sorting of root nuclei

Details of the flow sorting for Repli-seq analysis were described previously [9, 12]. Briefly, nuclei were isolated from the fixed root segments, and the incorporated EdU was conjugated to AF-488 using a Click-iT® EdU Alexa Fluor 488 Imaging Kit (Life Technologies). The nuclei were then resuspended in cell lysis buffer (CLB) [9] containing 2 μg/mL DAPI and 40 μg/mL Ribonuclease A and filtered through a CellTrics® 20-μm nylon mesh filter (Partec) just before flow sorting on an InFlux™ flow cytometer (BD Biosciences) equipped with UV (355 nm) and blue (488 nm) lasers. Nuclei prepared from the 1–3 mm root segments were sorted to collect populations of EdU/AF-488-labeled nuclei with DNA contents in three defined sub-stage gates between 4C and 8C, corresponding to early, mid and late S phase of the endocycle. The early endocycle gate was shifted slightly to the right to exclude mitotic nuclei in late S phase (Fig 1C). For each biological replicate, between 50,000 and 200,000 nuclei were sorted from each fraction

of the endocycle S phase. A small sample of nuclei from each gate was sorted into CLB buffer

containing DAPI and reanalyzed to determine the sort purity (S1 Fig). Sorting and reanalysis

details for the mitotic nuclei are described in [12].

For ChIP-seq experiments, roots were labeled with EdU, and nuclei were isolated from

0–3 mm (H3K27me3 and H3K4me3) or 0–5 mm (H3K56ac) root segments and conjugated to

AF-488 as described above. The 2C, 4C and 8C unlabeled, non S-phase populations of nuclei

were sorted into 2× extraction buffer 2 (EB2) [111] using the same sorting conditions as in Wear

et al. [12]. After sorting, the 2× EB2 was diluted to 1× with 1× STE. All flow cytometry data

were analyzed using FlowJo v10.0.6 (TreeStar, Inc.) as described in Wear et al. [12].

DNA and chromatin immunoprecipitations

For endocycle Repli-seq samples, reversal of formaldehyde cross links, nuclear DNA

purification and isolation, DNA shearing, EdU/AF-488 DNA immunoprecipitation with an anti-

Alexa Fluor 488 antibody (Molecular Probes, #A-11094, lot 895897), and DNA fragment

purification were performed as described in Wear et al. [12].

ChIP procedures were performed as in Wear et al. [12] except the chromatin was sheared

using a Covaris S220 ultrasonicator to an average fragment size of 200 bp using a peak incident

power of 140 W, 10% duty cycle, and 200 cycles per burst for 6 min. Three percent of the

chromatin volume was set aside to use as the input control for each of the 2C, 4C and 8C

samples and frozen at -70°C until the formaldehyde cross link reversal step. The antibodies used

for ChIP were as follows: *Zea mays* anti-CENH3 antibody at a 1:250 dilution (gift from R.K.

Dawe) [48], anti-H3K9me2 antibody at a 1:25 dilution (Cell Signaling Technologies; 9753, lot

4), anti-H3K56ac antibody at a 1:200 dilution (Millipore; 07-677, lot DAM1462569), anti-

H3K4me3 antibody at a 1:300 dilution (Millipore; 07-473, lot DAM1779237) and anti-

H3K27me3 antibody at a 1:300 dilution (Millipore; 07-449, lot 2,275,589). See S18 Fig for

antibody validation experiments for anti-H3K9me2 and anti-CENH3.

Library construction and sequencing

For Repli-seq and ChIP-seq samples, the final purified DNA was used to construct paired-end

libraries as described [12]. After adapter ligation, all samples underwent 17 cycles of PCR. For

each Repli-seq or ChIP-seq experiment, individual samples from three biological replicates

collected on different days were barcoded, pooled and sequenced on either the Illumina HiSeq

2000 or NextSeq platforms. However, in the case of the Repli-seq mitotic late-S samples and

CENH3 ChIP 4C samples, one biological replicate failed during library generation or

sequencing, resulting in data from only two biological replicates. Repli-seq and ChIP-seq read

mapping statistics are shown in S1 Spreadsheet.

Replication timing data analysis

Trimming and quality control of 100-bp paired-end Repli-seq reads were carried out as described

previously [12], and reads were aligned to the maize B73 RefGen_v4 reference genome [38]

(Ensembl Plants release 33; ftp://ftp.ensemblgenomes.org/pub/plants/release-33/gff3/zea_mays/)

using BWA-MEM v0.7.12 with default parameters [112]. Redundant reads resulting from PCR

amplification were removed from each of the alignment files using Picard

(http://broadinstitute.github.io/picard/) and SAMtools [113]. Properly paired, uniquely mapping

reads (MAPQ score > 10) were retained with SAMtools [113] for downstream analysis. The

resulting mitotic Repli-seq data were more than 3× the sequencing coverage of the endocycle

Repli-seq data (S1 Spreadsheet). Repli-seq results are robust at various sequencing depths [14],

but to ensure that the mitotic and endocycle data were comparable, the reads were downsampled

by a uniform random process using a custom python script incorporating the BEDTools suite

[114] to a total of 65.7 million reads per sample, representing almost 3× genome coverage for

each S-phase fraction (S1 Spreadsheet). We preferred this to normalization so that any possible

sampling bias due to sequencing depth would be similar in all samples.

Repli-seq data were analyzed using *Repliscan* [14]. Individual biological replicates of

Repli-seq data were independently analyzed, and after finding good correlation between

replicates (Pearson correlation coefficients from 0.80–0.99; S4 Fig) the replicates were

aggregated by sum and normalized to 1× genome coverage using the reads per genomic content

(RPGC) method. The following changes from the *Repliscan* default parameters described in [12]

were used. Read densities were aggregated in 3-kb windows across the genome (parameter -*w*

3000). Additionally, we customized the cutoff for reducing type one errors which excluded

genomic windows with extremely low coverage in the 2C reference sample. To identify these

low read mapping windows, which we labeled "blacklist", *Repliscan* log-transformed the read

counts from the pre-replicative 2C reference sample and windows with read counts in the lower

2.5% tail of a fitted normal distribution were excluded from all samples (parameter --*pcut* 2.5-

100). The upper 2.5% tail containing extremely high coverage windows or "spikes" was not

removed at this step, because we found that these data spikes were adequately normalized in the

subsequent step of dividing each 3-kb window in the S-phase samples by the 2C reference data –

which also normalized for sequencing biases and collapsed repeats (S3 Fig). The data were then

Haar wavelet smoothed [14] to produce the final profiles for early, mid and late S-phase

replication signals in the mitotic cycle and endocycle. Processed data files, formatted for the

Integrative Genomics Viewer (IGV) [115], are available for download from CyVerse (formerly

the iPlant Collaborative; [116]) via the information in S1 Spreadsheet.

Identifying regions of altered replication timing

The difference between normalized signal profiles of mitotic and endocycle Repli-seq data for

early, mid, and late S was calculated in 3-kb windows, and the maximum negative and positive

differences were then calculated for each chromosome and averaged. Regions showing a timing

difference of ≥ 25% (difference in replication signal ≥ 1.0) or ≥ 10% (difference in replication

signal ≥ 0.4) of the total range of differences in each profile were identified (S1 Table; S5 Fig)

using the data filter tool in SAS JMP Pro v14 (SAS Institute Inc.). Windows were kept in the

analysis only if their timing differences were "compensated" by opposite timing difference(s) of

≥ 25% or ≥ 10%, respectively, in one or both of the other two S-phase fractions. For example, a

decrease in early replication signal in endocycling cells must be compensated by an increase in

mid and/or late S-phase signal in the same cell population. Adjacent 3-kb windows with timing

differences that met either the ≥ 10% or ≥ 25% threshold were merged, keeping the two files

separate, using mergeBED in the BEDTools suite, and allowing a 6 kb gap distance (parameter -

d 6000) [114]. This initial step resulted in many very small regions being identified (S2 Table).

As a second step, if ≥ 10% regions were immediately adjacent to ≥ 25% regions, they were

merged together using mergeBED to highlight larger regions of contiguous change (S2 Table).

Only regions that contained at least one ≥ 25% region were kept for further analysis, and termed

regions of alternate timing (RATs). By requiring a ≥ 25% RT change core region to be included,

all of the stand-alone, extremely small regions (< 24 kb) were effectively filtered out, without the

requirement of an arbitrary size filter. RATs were categorized into three groups: 1) later in

mitotic to earlier in endocycle (Later-to-Earlier), 2) earlier in mitotic to later in endocycle

(Earlier-to-Later) and 3) a subset of the Earlier-to-Later RATS that were located in the

previously identified functional centromeres (Earlier-to-Later-CEN) (coordinates from [38]).

There were no Later-to-Earlier-CEN RATs. For a list of RAT regions, including genomic

coordinates and genes within them, see S2 and S3 Spreadsheets.

ChIP-seq data analysis

ChIP-seq reads for H3K27me3, H3K4me3, H3K56ac (100-bp paired-end reads), H3K9me2 and

CENH3 (150-bp paired-end reads) were trimmed, mapped to maize B73 RefGen_v4.33, and

filtered to retain only properly-paired, uniquely-mapped reads (MAPQ score > 10) as described

above for Repli-seq reads. The 2C ChIP and input data for H3K27me3, H3K4me3, H3K56ac is

from [12], while the 4C and 8C ChIP data was generated for this study, see S1 Spreadsheet. For

details on peak calling and analysis for H3K27me3, H3K4me3, H3K56ac, see S1 Text.

For visualization of CENH3 localization in 2C, 4C and 8C nuclei, read counts for

individual biological replicates of CENH3 or input samples were scaled to 1× genome coverage

using the reads per genomic content (RPGC) method. Biological replicate data had good

agreement (Pearson's correlation coefficient values between biological replicates of 0.97-0.99;

S1 Spreadsheet), and were merged and scaled again to 1× coverage so the samples would be

comparable. CENH3 scaled read counts in each 3-kb window were divided by the scaled read

counts from the input sample for the corresponding ploidy level, resulting in CENH3 fold

enrichment values relative to input.

To compare CENH3 enrichment relative to DNA content in 2C, 4C and 8C cells over

entire centromeres, we calculated the percent of total CENH3 reads found in a given centromere

(using coordinates from [38]), divided by the percent of total input reads corresponding to that

centromere. This was done separately for individual biological replicates; we then calculated the

mean fold enrichment estimates. H3K9me2 fold enrichment over entire centromeres and

pericentromeres was calculated in the same way.

Genomic features

The maize filtered gene set Zm00001d.2 annotation from B73 RefGen_v4 [38] was downloaded

from Ensembl Plants (ftp://ftp.ensemblgenomes.org/pub/plants/release-33/gff3/zea_mays/). The

updated B73 Refgen_v4 TEv2 disjoined annotation [39] was downloaded from

http://mcstitzer.github.io/maize_TEs. Coordinates for mapped *CentC* satellite repeat regions are

described in Gent et al. [40]. The percent AT content was calculated in 3-kb static windows

across the genome.

Analysis of features in RATs and random permutation analysis

We tested the association of various genomic features with the non-CEN RAT categories by

determining the overlap of a particular feature with each RAT type. The coordinates for genomic

features (genes, expressed genes, TE superfamilies) were intersected with RAT coordinate

intervals using intersectBED (parameters *-wa -wb*) in the BEDtools suite [114]. The percent of

RATs containing a feature or the percent coverage of genes and TE superfamilies were computed

and compared to values for the genome as a whole. The number of genes per RAT was also

determined using intersectBED (parameter *-u*).

For comparison, the coordinates for the non-CEN Earlier-to-Later and Later-to-Earlier

RAT sets were randomly shuffled around the genome, excluding functional centromeres, using

BEDTools shuffle [114]. These random sets preserved the number of regions and region size of

the original RAT sets, and are labeled "EtoL shuffle1" and "LtoE shuffle1" for the Earlier-to-

Later and Later-to-Earlier RATs, respectively. When there appeared to be differences in the

observed overlap values with genomic features between non-CEN RATs and their corresponding

random shuffle sets, a permutation or feature randomization test, as described in [12] was used to

assess the statistical significance of the observed value. To do so, the coordinates for the non-

CEN RAT sets were randomly shuffled around the genome 1000 times, as described above.

Analysis of features in centromeres and pericentromeres

For comparison to CEN regions (coordinates from [38]), pericentromeres were arbitrarily

defined as the ± 1 Mb flanking each CEN. In the case of chromosome 9, the pericentromere

included the ± 1 Mb flanking both CEN 9a and 9b. Replication timing signal values in CENs and

pericentromeres were intersected with genes, *CRM1* and *CRM2* families and mapped *CentC*

regions using intersectBED (parameters -*wa* -*wb*) in the BEDtools suite [114]. Only elements

that covered at least half of a 3-kb window of Repli-seq data were included in Fig 4, while

elements with any amount of overlap were included in S14 Fig. Additionally, if a single gene or

CRM element spanned more than one of the 3-kb windows, the replication signals were averaged

using mergeBED (parameter -*o* mean) to compute a single value for the entire gene or element.

ACKNOWLEDGEMENTS

We thank the following people for helpful discussions of this work: Ashley Brooks, Emily

Wheeler and Hank Bass. We are grateful to Kelly Dawe and Jonathan Gent for sharing the maize

CENH3 antibody, mapped *CentC* data and advice. We thank Patrick Mulvaney and James

Harrison Priode for help with harvesting plant material.

REFERENCES

1.   Edgar BA, Zielke N, Gutierrez C. Endocycles: a recurrent evolutionary innovation for post-mitotic cell growth. Nat Rev Mol Cell Biol. 2014;15(3):197-210. Epub 2014/02/22. doi: 10.1038/nrm3756. PubMed PMID: 24556841.

2.   Lee HO, Davidson JM, Duronio RJ. Endoreplication: polyploidy with purpose. Genes Dev. 2009;23(21):2461-77. Epub 2009/11/04. doi: 10.1101/gad.1829209. PubMed PMID: 19884253; PubMed Central PMCID: PMCPMC2779750.

3.   Fox DT, Duronio RJ. Endoreplication and polyploidy: insights into development and disease. Development. 2013;140(1):3-12. Epub 2012/12/12. doi: 10.1242/dev.080531. PubMed PMID: 23222436; PubMed Central PMCID: PMCPMC3513989.

4.   Galbraith DW, Harkins KR, Knapp S. Systemic endopolyploidy in *Arabidopsis thaliana*. Plant Physiol. 1991;96(3):985-9. Epub 1991/07/01. doi: 10.1104/pp.96.3.985. PubMed PMID: 16668285; PubMed Central PMCID: PMCPMC1080875.

5.   Joubes J, Chevalier C. Endoreduplication in higher plants. Plant Mol Biol. 2000;43(5-6):735-45. Epub 2000/11/23. doi: 10.1023/a:1006446417196. PubMed PMID: 11089873.

6.   Breuer C, Ishida T, Sugimoto K. Developmental control of endocycles and cell growth in plants. Curr Opin Plant Biol. 2010;13(6):654-60. Epub 2010/11/26. doi: 10.1016/j.pbi.2010.10.006. PubMed PMID: 21094078.

7.   Hayashi K, Hasegawa J, Matsunaga S. The boundary of the meristematic and elongation zones in roots: endoreduplication precedes rapid cell expansion. Sci Rep. 2013;3:2723. Epub 2013/10/15. doi: 10.1038/srep02723. PubMed PMID: 24121463; PubMed Central PMCID: PMCPMC3796303.

8.   Bass HW, Wear EE, Lee TJ, Hoffman GG, Gumber HK, Allen GC, et al. A maize root tip system to study DNA replication programmes in somatic and endocycling nuclei during plant development. J Exp Bot. 2014;65(10):2747-56. Epub 2014/01/23. doi: 10.1093/jxb/ert470. PubMed PMID: 24449386.

9.   Wear EE, Concia L, Brooks AM, Markham EA, Lee T-J, Allen GC, et al. Isolation of plant nuclei at defined cell cycle stages using EdU labeling and flow cytometry. Plant Cell Division: Methods and Protocols. 2016:69-86.

10.   Concia L, Brooks AM, Wheeler E, Zynda GJ, Wear EE, LeBlanc C, et al. Genome-wide analysis of the *Arabidopsis* replication timing program. Plant Physiol. 2018;176(3):2166-85. Epub 2018/01/06. doi: 10.1104/pp.17.01537. PubMed PMID: 29301956; PubMed Central PMCID: PMCPMC5841712.

11.   Bass HW, Hoffman GG, Lee TJ, Wear EE, Joseph SR, Allen GC, et al. Defining multiple, distinct, and shared spatiotemporal patterns of DNA replication and endoreduplication from 3D image analysis of developing maize (*Zea mays* L.) root tip nuclei. Plant Mol Biol. 2015;89(4-5):339-51. Epub 2015/09/24. doi: 10.1007/s11103-015-0364-4. PubMed PMID: 26394866; PubMed Central PMCID: PMCPMC4631726.

12.   Wear EE, Song J, Zynda GJ, LeBlanc C, Lee TJ, Mickelson-Young L, et al. Genomic analysis of the DNA replication timing program during mitotic S phase in maize (*Zea mays*) root tips. Plant Cell. 2017;29(9):2126-49. Epub 2017/08/27. doi: 10.1105/tpc.17.00037. PubMed PMID: 28842533; PubMed Central PMCID: PMCPMC5635974.

13.   Hansen RS, Thomas S, Sandstrom R, Canfield TK, Thurman RE, Weaver M, et al. Sequencing newly replicated DNA reveals widespread plasticity in human replication timing. Proc Natl Acad Sci U S A. 2010;107(1):139-44. Epub 2009/12/08. doi:

10.1073/pnas.0912402107. PubMed PMID: 19966280; PubMed Central PMCID:
PMCPMC2806781.
14.   Zynda GJ, Song J, Concia L, Wear EE, Hanley-Bowdoin L, Thompson WF, et al.
Repliscan: a tool for classifying replication timing regions. BMC Bioinformatics.
2017;18(1):362. Epub 2017/08/09. doi: 10.1186/s12859-017-1774-x. PubMed PMID: 28784090;
PubMed Central PMCID: PMCPMC5547489.
15.   Pryor A, Faulkner K, Rhoades MM, Peacock WJ. Asynchronous replication of
heterochromatin in maize. Proc Natl Acad Sci U S A. 1980;77(11):6705-9. Epub 1980/11/01.
doi: 10.1073/pnas.77.11.6705. PubMed PMID: 16592919; PubMed Central PMCID:
PMCPMC350357.
16.   Marchal C, Sima J, Gilbert DM. Control of DNA replication timing in the 3D genome.
Nat Rev Mol Cell Biol. 2019;20(12):721-37. Epub 2019/09/04. doi: 10.1038/s41580-019-0162-
y. PubMed PMID: 31477886.
17.   Hua BL, Orr-Weaver TL. DNA replication control during *Drosophila* development:
insights into the onset of S phase, replication initiation, and fork progression. Genetics.
2017;207(1):29-47. Epub 2017/09/07. doi: 10.1534/genetics.115.186627. PubMed PMID:
28874453; PubMed Central PMCID: PMCPMC5586379.
18.   Hammond MP, Laird CD. Chromosome structure and DNA replication in nurse and
follicle cells of *Drosophila melanogaster*. Chromosoma. 1985;91(3-4):267-78. Epub 1985/01/01.
doi: 10.1007/bf00328222. PubMed PMID: 3920017.
19.   Hammond MP, Laird CD. Control of DNA replication and spatial distribution of defined
DNA sequences in salivary gland cells of *Drosophila melanogaster*. Chromosoma. 1985;91(3-
4):279-86. Epub 1985/01/01. doi: 10.1007/bf00328223. PubMed PMID: 3920018.
20.   Nordman J, Li S, Eng T, Macalpine D, Orr-Weaver TL. Developmental control of the
DNA replication and transcription programs. Genome Res. 2011;21(2):175-81. Epub
2010/12/24. doi: 10.1101/gr.114611.110. PubMed PMID: 21177957; PubMed Central PMCID:
PMCPMC3032921.
21.   Hribova E, Holusova K, Travnicek P, Petrovska B, Ponert J, Simkova H, et al. The
enigma of progressively partial endoreplication: new insights provided by flow cytometry and
next-generation sequencing. Genome Biol Evol. 2016;8(6):1996-2005. doi:
10.1093/gbe/evw141. PubMed PMID: WOS:000386368200011.
22.   Travnicek P, Certner M, Ponert J, Chumova Z, Jersakova J, Suda J. Diversity in genome
size and GC content shows adaptive potential in orchids and is closely linked to partial
endoreplication, plant life-history traits and climatic conditions. New Phytol. 2019. doi:
10.1111/nph.15996. PubMed PMID: WOS:000489166400001.
23.   Bauer MJ, Birchler JA. Organization of endoreduplicated chromosomes in the endosperm
of *Zea mays* L. Chromosoma. 2006;115(5):383-94. Epub 2006/06/03. doi: 10.1007/s00412-006-
0068-2. PubMed PMID: 16741707.
24.   Jacob Y, Stroud H, Leblanc C, Feng S, Zhuo L, Caro E, et al. Regulation of
heterochromatic DNA replication by histone H3 lysine 27 methyltransferases. Nature.
2010;466(7309):987-91. doi: 10.1038/nature09290. PubMed PMID: 20631708; PubMed Central
PMCID: PMCPMC2964344.
25.   Hiratani I, Ryba T, Itoh M, Yokochi T, Schwaiger M, Chang CW, et al. Global
reorganization of replication domains during embryonic stem cell differentiation. PLoS Biol.
2008;6(10):e245. doi: 10.1371/journal.pbio.0060245. PubMed PMID: 18842067; PubMed
Central PMCID: PMCPMC2561079.

26.   Hiratani I, Ryba T, Itoh M, Rathjen J, Kulik M, Papp B, et al. Genome-wide dynamics of
replication timing revealed by in vitro models of mouse embryogenesis. Genome Res.
2010;20(2):155-69. doi: 10.1101/gr.099796.109. PubMed PMID: 19952138; PubMed Central
PMCID: PMCPMC2813472.
27.   Rivera-Mulia JC, Buckley Q, Sasaki T, Zimmerman J, Didier RA, Nazor K, et al.
Dynamic changes in replication timing and gene expression during lineage specification of
human pluripotent stem cells. Genome Res. 2015;25(8):1091-103. Epub 2015/06/10. doi:
10.1101/gr.187989.114. PubMed PMID: 26055160; PubMed Central PMCID:
PMCPMC4509994.
28.   Salic A, Mitchison TJ. A chemical method for fast and sensitive detection of DNA
synthesis *in vivo*. Proc Natl Acad Sci USA. 2008;105(7):2415-20. doi:
10.1073/pnas.0712168105. PubMed PMID: 18272492; PubMed Central PMCID:
PMCPMC2268151.
29.   Yarosh W, Spradling AC. Incomplete replication generates somatic DNA alterations
within *Drosophila* polytene salivary gland cells. Genes Dev. 2014;28(16):1840-55. Epub
2014/08/17. doi: 10.1101/gad.245811.114. PubMed PMID: 25128500; PubMed Central PMCID:
PMCPMC4197960.
30.   Peacock WJ, Dennis ES, Rhoades MM, Pryor AJ. Highly repeated DNA sequence
limited to knob heterochromatin in maize. Proc Natl Acad Sci USA. 1981;78(7):4490-4. PubMed
PMID: 16593063; PubMed Central PMCID: PMCPMC319817.
31.   Ananiev EV, Phillips RL, Rines HW. A knob-associated tandem repeat in maize capable
of forming fold-back DNA segments: are chromosome knobs megatransposons? Proc Natl Acad
Sci USA. 1998;95(18):10785-90. PubMed PMID: 9724782; PubMed Central PMCID:
PMCPMC27973.
32.   Rivin CJ, Cullis CA, Walbot V. Evaluating quantitative variation in the genome of *Zea*
mays. Genetics. 1986;113(4):1009-19. PubMed PMID: 3744025; PubMed Central PMCID:
PMCPMC1202908.
33.   Ananiev EV, Phillips RL, Rines HW. Chromosome-specific molecular organization of
maize (*Zea mays* L.) centromeric regions. Proc Natl Acad Sci USA. 1998;95(22):13073-8. doi:
DOI 10.1073/pnas.95.22.13073. PubMed PMID: 9789043; PubMed Central PMCID:
PMCPMC23713.
34.   Presting GG, Malysheva L, Fuchs J, Schubert I. A TY3/GYPSY retrotransposon-like
sequence localizes to the centromeric regions of cereal chromosomes. The Plant Journal.
1998;16(6):721-8.
35.   Miller JT, Dong FG, Jackson SA, Song J, Jiang JM. Retrotransposon-related DNA
sequences in the centromeres of grass chromosomes. Genetics. 1998;150(4):1615-23. PubMed
PMID: WOS:000077401300025.
36.   Baucom RS, Estill JC, Chaparro C, Upshaw N, Jogi A, Deragon JM, et al. Exceptional
diversity, non-random distribution, and rapid evolution of retroelements in the B73 maize
genome. PLoS Genet. 2009;5(11):e1000732. doi: 10.1371/journal.pgen.1000732. PubMed
PMID: 19936065; PubMed Central PMCID: PMCPMC2774510.
37.   Gent JI, Schneider KL, Topp CN, Rodriguez C, Presting GG, Dawe RK. Distinct
influences of tandem repeats and retrotransposons on CENH3 nucleosome positioning. Epigenet
Chromatin. 2011;4. doi: Artn 3
10.1186/1756-8935-4-3. PubMed PMID: WOS:000288223700001.

38.   Jiao Y, Peluso P, Shi J, Liang T, Stitzer MC, Wang B, et al. Improved maize reference
genome with single-molecule technologies. Nature. 2017;546(7659):524-7. doi:
10.1038/nature22971. PubMed PMID: 28605751.
39.   Anderson SN, Stitzer MC, Brohammer AB, Zhou P, Noshay JM, O'Connor CH, et al.
Transposable elements contribute to dynamic genome content in maize. Plant Journal. 2019. doi:
10.1111/tpj.14489. PubMed PMID: WOS:000486761400001.
40.   Gent JI, Wang N, Dawe RK. Stable centromere positioning in diverse sequence contexts
of complex and satellite centromeres of maize and wild relatives. Genome Biology. 2017;18. doi:
ARTN 121
10.1186/s13059-017-1249-4. PubMed PMID: WOS:000403802400001.
41.   SanMiguel P, Tikhonov A, Jin YK, Motchoulskaia N, Zakharov D, Melake-Berhan A, et
al. Nested retrotransposons in the intergenic regions of the maize genome. Science.
1996;274(5288):765-8. doi: DOI 10.1126/science.274.5288.765. PubMed PMID: 8864112.
42.   Liu R, Vitte C, Ma J, Mahama AA, Dhliwayo T, Lee M, et al. A GeneTrek analysis of
the maize genome. Proc Natl Acad Sci USA. 2007;104(28):11844-9. doi:
10.1073/pnas.0704258104. PubMed PMID: 17615239; PubMed Central PMCID:
PMCPMC1913904.
43.   Schnable PS, Ware D, Fulton RS, Stein JC, Wei F, Pasternak S, et al. The B73 maize
genome: complexity, diversity, and dynamics. Science. 2009;326(5956):1112-5. doi:
10.1126/science.1178534. PubMed PMID: 19965430.
44.   Comai L, Maheshwari S, Marimuthu MPA. Plant centromeres. Current Opinion in Plant
Biology. 2017;36(Supplement C):158-67. doi: 10.1016/j.pbi.2017.03.003.
45.   Fukagawa T, Earnshaw WC. The centromere: chromatin foundation for the kinetochore
machinery. Developmental Cell. 2014;30(5):497-509. doi: 10.1016/j.devcel.2014.08.016.
PubMed PMID: WOS:000341296100005.
46.   McKinley KL, Cheeseman IM. The molecular basis for centromere identity and function.
Nat Rev Mol Cell Biol. 2016;17(1):16-29. Epub 2015/11/26. doi: 10.1038/nrm.2015.5. PubMed
PMID: 26601620.
47.   Gent JI, Nannas NJ, Liu Y, Su H, Zhao H, Gao Z, et al. Genomics of maize centromeres.
The Maize Genome. Compendium of Plant Genomes2018. p. 59-80.
48.   Zhong CX, Marshall JB, Topp C, Mroczek R, Kato A, Nagaki K, et al. Centromeric
retroelements and satellites interact with maize kinetochore protein CENH3. Plant Cell.
2002;14(11):2825-36. Epub 2002/11/06. doi: 10.1105/tpc.006106. PubMed PMID: 12417704;
PubMed Central PMCID: PMCPMC152730.
49.   Presting GG. Centromeric retrotransposons and centromere function. Current Opinion in
Genetics & Development. 2018;49:79-84. doi: 10.1016/j.gde.2018.03.004. PubMed PMID:
WOS:000433211500012.
50.   Zhao H, Zhu X, Wang K, Gent JI, Zhang W, Dawe RK, et al. Gene expression and
chromatin modifications associated with maize centromeres. G3 (Bethesda). 2015;6(1):183-92.
Epub 2015/11/14. doi: 10.1534/g3.115.022764. PubMed PMID: 26564952; PubMed Central
PMCID: PMCPMC4704717.
51.   Zhang W, Lee HR, Koo DH, Jiang J. Epigenetic modification of centromeric chromatin:
hypomethylation of DNA sequences in the CENH3-associated chromatin in *Arabidopsis thaliana*
and maize. Plant Cell. 2008;20(1):25-34. Epub 2008/02/02. doi: 10.1105/tpc.107.057083.
PubMed PMID: 18239133; PubMed Central PMCID: PMCPMC2254920.

52. Gent JI, Madzima TF, Bader R, Kent MR, Zhang X, Stam M, et al. Accessible DNA and
relative depletion of H3K9me2 at maize loci undergoing RNA-directed DNA methylation. Plant
Cell. 2014;26(12):4903-17. Epub 2014/12/04. doi: 10.1105/tpc.114.130427. PubMed PMID:
25465407; PubMed Central PMCID: PMCPMC4311197.
53. Bailey AO, Panchenko T, Sathyan KM, Petkowski JJ, Pai PJ, Bai DL, et al.
Posttranslational modification of CENP-A influences the conformation of centromeric
chromatin. Proc Natl Acad Sci U S A. 2013;110(29):11827-32. Epub 2013/07/03. doi:
10.1073/pnas.1300325110. PubMed PMID: 23818633; PubMed Central PMCID:
PMCPMC3718089.
54. Shelby RD, Monier K, Sullivan KF. Chromatin assembly at kinetochores is uncoupled
from DNA replication. J Cell Biol. 2000;151(5):1113-8. Epub 2000/11/22. doi:
10.1083/jcb.151.5.1113. PubMed PMID: 11086012; PubMed Central PMCID:
PMCPMC2174364.
55. Jansen LE, Black BE, Foltz DR, Cleveland DW. Propagation of centromeric chromatin
requires exit from mitosis. J Cell Biol. 2007;176(6):795-805. Epub 2007/03/07. doi:
10.1083/jcb.200701066. PubMed PMID: 17339380; PubMed Central PMCID:
PMCPMC2064054.
56. Boyarchuk E, Montes de Oca R, Almouzni G. Cell cycle dynamics of histone variants at
the centromere, a model for chromosomal landmarks. Curr Opin Cell Biol. 2011;23(3):266-76.
Epub 2011/04/08. doi: 10.1016/j.ceb.2011.03.006. PubMed PMID: 21470840.
57. Nagaki K, Kashihara K, Murata M. Visualization of diffuse centromeres with
centromere-specific histone H3 in the holocentric plant *Luzula nivea*. Plant Cell.
2005;17(7):1886-93. doi: 10.1105/tpc.105.032961. PubMed PMID: WOS:000230165400004.
58. Lermontova I, Schubert V, Fuchs J, Klatte S, Macas J, Schubert I. Loading of
Arabidopsis centromeric histone CENH3 occurs mainly during G2 and requires the presence of
the histone fold domain. Plant Cell. 2006;18(10):2443-51. doi: 10.1105/tpc.106.043174. PubMed
PMID: WOS:000241818300004.
59. Lermontova I, Fuchs J, Schubert V, Schubert I. Loading time of the centromeric histone
H3 variant differs between plants and animals. Chromosoma. 2007;116(6):507-10. doi:
10.1007/s00412-007-0122-8. PubMed PMID: WOS:000250880100002.
60. Schubert V, Lermontova I, Schubert I. Loading of the centromeric histone H3 variant
during meiosis–how does it differ from mitosis? Chromosoma. 2014;123(5):491-7.
61. Nechemia-Arbely Y, Miga KH, Shoshani O, Aslanian A, McMahon MA, Lee AY, et al.
DNA replication acts as an error correction mechanism to maintain centromere identity by
restricting CENP-A to centromeres. Nat Cell Biol. 2019;21(6):743-+. doi: 10.1038/s41556-019-
0331-4. PubMed PMID: WOS:000470080100011.
62. Baluska F. Nuclear size, DNA content, and chromatin condensation are different in
individual tissues of the maize root apex. Protoplasma. 1990;158(1-2):45-52. doi: Doi
10.1007/Bf01323273. PubMed PMID: WOS:A1990EK62800007.
63. Sugimoto-Shirasu K, Roberts K. "Big it up": endoreduplication and cell-size control in
plants. Current Opinion in Plant Biology. 2003;6(6):544-53. doi: 10.1016/j.pbi.2003.09.009.
64. Bhosale R, Boudolf V, Cuevas F, Lu R, Eekhout T, Hu Z, et al. A spatiotemporal DNA
endoploidy map of the *Arabidopsis* root reveals roles for the endocycle in root development and
stress adaptation. The Plant Cell. 2018:tpc. 00983.2017.

65.   Cooper S. Rethinking synchronization of mammalian cells for cell cycle analysis.
Cellular and Molecular Life Sciences. 2003;60(6):1099-106. Epub 2003/07/16. doi:
10.1007/s00018-003-2253-2. PubMed PMID: 12861378.
66.   Phillips RL, Kaeppler SM, Olhoft P. Genetic instability of plant tissue cultures:
breakdown of normal controls. Proc Natl Acad Sci USA. 1994;91(12):5222-6. PubMed PMID:
8202472; PubMed Central PMCID: PMCPMC43966.
67.   Mayshar Y, Ben-David U, Lavon N, Biancotti JC, Yakir B, Clark AT, et al. Identification
and classification of chromosomal aberrations in human induced pluripotent stem cells. Cell
Stem Cell. 2010;7(4):521-31. doi: 10.1016/j.stem.2010.07.017. PubMed PMID: 20887957.
68.   Laurent LC, Ulitsky I, Slavin I, Tran H, Schork A, Morey R, et al. Dynamic changes in
the copy number of pluripotency and cell proliferation genes in human ESCs and iPSCs during
reprogramming and time in culture. Cell Stem Cell. 2011;8(1):106-18. doi:
10.1016/j.stem.2010.12.003. PubMed PMID: 21211785; PubMed Central PMCID:
PMCPMC3043464.
69.   Dimitrova DS, Gilbert DM. The spatial position and replication timing of chromosomal
domains are both established in early G1 phase. Molecular Cell. 1999;4(6):983-93. PubMed
PMID: ISI:000084485900010.
70.   Frawley LE, Orr-Weaver TL. Polyploidy. Curr Biol. 2015;25(9):R353-R8. doi: DOI
10.1016/j.cub.2015.03.037. PubMed PMID: WOS:000353999000005.
71.   Van't Hof J. DNA replication in plants: Cold spring harbor laboratory press; 1996.
72.   Schwaiger M, Stadler MB, Bell O, Kohler H, Oakeley EJ, Schubeler D. Chromatin state
marks cell-type- and gender-specific replication of the *Drosophila* genome. Genes Dev.
2009;23(5):589-601. doi: 10.1101/gad.511809. PubMed PMID: 19270159; PubMed Central
PMCID: PMCPMC2658520.
73.   Ryba T, Hiratani I, Lu J, Itoh M, Kulik M, Zhang J, et al. Evolutionarily conserved
replication timing profiles predict long-range chromatin interactions and distinguish closely
related cell types. Genome Res. 2010;20(6):761-70. doi: 10.1101/gr.099655.109. PubMed
PMID: 20430782; PubMed Central PMCID: PMCPMC2877573.
74.   Lubelsky Y, Prinz JA, DeNapoli L, Li YL, Belsky JA, MacAlpine DM. DNA replication
and transcription programs respond to the same chromatin cues. Genome Research.
2014;24(7):1102-14. doi: 10.1101/gr.160010.113. PubMed PMID: WOS:000338185000005.
75.   Armstrong RL, Penke TJR, Strahl BD, Matera AG, McKay DJ, MacAlpine DM, et al.
Chromatin conformation and transcriptional activity are permissive regulators of DNA
replication initiation in *Drosophila*. Genome Research. 2018;28(11):1688-700. doi:
10.1101/gr.239913.118. PubMed PMID: WOS:000448950400009.
76.   Massey DJ, Kim D, Brooks KE, Smolka MB, Koren A. Next-generation sequencing
enables spatiotemporal resolution of human centromere replication timing. Genes-Basel.
2019;10(4):269.
77.   McCarroll RM, Fangman WL. Time of replication of yeast centromeres and telomeres.
Cell. 1988;54(4):505-13.
78.   Raghuraman MK, Winzeler EA, Collingwood D, Hunt S, Wodicka L, Conway A, et al.
Replication dynamics of the yeast genome. Science. 2001;294(5540):115-21. doi:
10.1126/science.294.5540.115. PubMed PMID: WOS:000171448800041.
79.   Kim SM, Dubey DD, Huberman JA. Early-replicating heterochromatin. Genes Dev.
2003;17(3):330-5. doi: 10.1101/gad.1046203. PubMed PMID: 12569122; PubMed Central
PMCID: PMCPMC195982.

80.	Koren A, Tsai HJ, Tirosh I, Burrack LS, Barkai N, Berman J. Epigenetically-inherited
centromere and neocentromere DNA replicates earliest in S-phase. Plos Genetics. 2010;6(8). doi:
ARTN e1001068
10.1371/journal.pgen.1001068. PubMed PMID: WOS:000281383800023.
81.	Tenhagen KG, Gilbert DM, Willard HF, Cohen SN. Replication timing of DNA-
sequences associated with human centromeres and telomeres. Molecular and Cellular Biology.
1990;10(12):6348-55. PubMed PMID: ISI:A1990EJ60200026.
82.	O'Keefe R, Henderson S, Spector DL. Dynamic organization of DNA replication in
mammalian cell nuclei: spatially and temporally defined replication of chromosome-specific
satelite DNA sequences. Journal of Cell Biology. 1992;116.
83.	Hultdin M, Gronlund E, Norrback KF, Just T, Taneja K, Roos G. Replication timing of
human telomeric DNA and other repetitive sequences analyzed by fluorescence in situ
hybridization and flow cytometry. Exp Cell Res. 2001;271(2):223-9. doi:
10.1006/excr.2001.5391. PubMed PMID: ISI:000172749500003.
84.	Sullivan B, Karpen G. Centromere identity in *Drosophila* is not determined in vivo by
replication timing. Journal of Cell Biology. 2001;154(4):683-90. doi: DOI
10.1083/jcb.200103001. PubMed PMID: WOS:000170683900002.
85.	Ouspenski II, Van Hooser AA, Brinkley BR. Relevance of histone acetylation and
replication timing for deposition of centromeric histone CENP-A. Exp Cell Res.
2003;285(2):175-88. doi: 10.1016/S0014-4827(03)00011-9. PubMed PMID:
WOS:000182507500002.
86.	Shang WH, Hori T, Martins NM, Toyoda A, Misu S, Monma N, et al. Chromosome
engineering allows the efficient isolation of vertebrate neocentromeres. Dev Cell.
2013;24(6):635-48. Epub 2013/03/19. doi: 10.1016/j.devcel.2013.02.009. PubMed PMID:
23499358; PubMed Central PMCID: PMCPMC3925796.
87.	Fuchs J, Strehl S, Brandes A, Schweizer D, Schubert I. Molecular--cytogenetic
characterization of the *Vicia faba* genome--heterochromatin differentiation, replication patterns
and sequence localization. Chromosome Res. 1998;6(3):219-30.
88.	Schubert I. Late-replicating satellites: something for all centromeres? Trends in Genetics.
1998;14(10):385-6.
89.	Jasencakova Z, Meister A, Schubert I. Chromatin organization and its relation to
replication and histone acetylation during the cell cycle in barley. Chromosoma. 2001;110(2):83-
92. doi: 10.1007/s004120100132. PubMed PMID: 11453558.
90.	Samaniego R, de la Torre C, de la Espina SMD. Dynamics of replication foci and nuclear
matrix during S phase in *Allium cepa* L. cells. Planta. 2002;215(2):195-204. doi:
10.1007/s00425-002-0741-5. PubMed PMID: WOS:000176778800005.
91.	Macheret M, Halazonetis TD. Intragenic origins due to short G1 phases underlie
oncogene-induced DNA replication stress. Nature. 2018.
92.	Smith OK, Kim R, Fu HQ, Martin MM, Lin CM, Utani K, et al. Distinct epigenetic
features of differentiation-regulated replication origins. Epigenet Chromatin. 2016;9. doi: ARTN
18
10.1186/s13072-016-0067-3. PubMed PMID: WOS:000376861500001.
93.	Albert PS, Gao Z, Danilova TV, Birchler JA. Diversity of chromosomal karyotypes in
maize and its relatives. Cytogenet Genome Res. 2010;129(1-3):6-16. doi: 10.1159/000314342.
PubMed PMID: 20551613.

94.   Albert PS, Zhang T, Semrau K, Rouillard J-M, Kao Y-H, Wang C-JR, et al. Whole-
chromosome paints in maize reveal rearrangements, nuclear domains, and chromosomal
relationships. Proceedings of the National Academy of Sciences. 2019:201813957.
95.   Hemmerich P, Weidtkamp-Peters S, Hoischen C, Schmiedeberg L, Erliandri I, Diekmann
S. Dynamics of inner kinetochore assembly and maintenance in living cells. Journal of Cell
Biology. 2008;180(6):1101-14. doi: DOI 10.1083/jcb.200710052. PubMed PMID:
WOS:000254295400007.
96.   Dembinsky D, Woll K, Saleem M, Liu Y, Fu Y, Borsuk LA, et al. Transcriptomic and
proteomic analyses of pericycle cells of the maize primary root. Plant Physiology.
2007;145(3):575-88. doi: 10.1104/pp.107.106203. PubMed PMID: WOS:000250712800001.
97.   Orr-Weaver TL. When bigger is better: the role of polyploidy in organogenesis. Trends in
Genetics. 2015;31(6):307-15. doi: 10.1016/j.tig.2015.03.011. PubMed PMID:
WOS:000356636400005.
98.   del Arco AG, Edgar BA, Erhardt S. In vivo analysis of centromeric proteins reveals a
stem cell-specific asymmetry and an essential role in differentiated, non-proliferating cells. Cell
Rep. 2018;22(8):1982-93.
99.   Birchler JA, Han FP. Maize centromeres: structure, function, epigenetics. Annu Rev
Genet. 2009;43:287-303. doi: Doi 10.1146/Annurev-Genet-102108-134834. PubMed PMID:
ISI:000273580300013.
100.   Henikoff S, Furuyama T, editors. Epigenetic inheritance of centromeres. Cold Spring
Harb Sym; 2010: Cold Spring Harbor Laboratory Press.
101.   Muller S, Almouzni G. Chromatin dynamics during the cell cycle at centromeres. Nature
Reviews Genetics. 2017;18(3):192-208. doi: 10.1038/nrg.2016.157. PubMed PMID:
WOS:000394428300010.
102.   Zhang XL, Li XX, Marshall JB, Zhong CX, Dawe RK. Phosphoserines on maize
CENTROMERIC HISTONE H3 and histone H3 demarcate the centromere and pericentromere
during chromosome segregation. Plant Cell. 2005;17(2):572-83. doi: 10.1105/tpc.104.028522.
PubMed PMID: WOS:000227043800019.
103.   Demidov D, Heckmann S, Weiss O, Rutten T, Tomastikova ED, Kuhlmann M, et al.
Deregulated phosphorylation of CENH3 at Ser65 affects the development of floral meristems in
Arabidopsis thaliana. Frontiers in Plant Science. 2019;10. doi: ARTN 928
10.3389/fpls.2019.00928. PubMed PMID: WOS:000477060000001.
104.   Niikura Y, Kitagawa R, Ogi H, Abdulle R, Pagala V, Kitagawa K. CENP-A K124
ubiquitylation is required for CENP-A deposition at the centromere. Dev Cell. 2015;32(5):589-
603. Epub 2015/03/03. doi: 10.1016/j.devcel.2015.01.024. PubMed PMID: 25727006; PubMed
Central PMCID: PMCPMC4374629.
105.   Bui M, Pitman M, Nuccio A, Roque S, Donlin-Asp PG, Nita-Lazar A, et al. Internal
modifications in the CENP-A nucleosome modulate centromeric dynamics. Epigenet Chromatin.
2017;10. doi: ARTN 17
10.1186/s13072-017-0124-6. PubMed PMID: WOS:000398515300001.
106.   Takebayashi S, Dileep V, Ryba T, Dennis JH, Gilbert DM. Chromatin-interaction
compartment switch at developmentally regulated chromosomal domains reveals an unusual
principle of chromatin folding. Proc Natl Acad Sci U S A. 2012;109(31):12574-9. Epub
2012/07/19. doi: 10.1073/pnas.1207185109. PubMed PMID: 22807480; PubMed Central
PMCID: PMCPMC3411983.

107.   Heinz KS, Casas-Delucchi CS, Torok T, Cmarko D, Rapp A, Raska I, et al. Peripheral re-
localization of constitutive heterochromatin advances its replication timing and impairs
maintenance of silencing marks. Nucleic Acids Res. 2018;46(12):6112-28. Epub 2018/05/12.
doi: 10.1093/nar/gky368. PubMed PMID: 29750270; PubMed Central PMCID:
PMCPMC6158597.
108.   Krishnakumar V, Choi Y, Beck E, Wu Q, Luo A, Sylvester A, et al. A maize database
resource that captures tissue-specific and subcellular-localized gene expression, via fluorescent
tags and confocal imaging (Maize Cell Genomics Database). Plant Cell Physiol. 2015;56(1):e12.
Epub 2014/11/30. doi: 10.1093/pcp/pcu178. PubMed PMID: 25432973.
109.   Dong P, Tu X, Chu PY, Lu P, Zhu N, Grierson D, et al. 3D chromatin architecture of
large plant genomes determined by local A/B compartments. Mol Plant. 2017;10(12):1497-509.
Epub 2017/11/28. doi: 10.1016/j.molp.2017.11.005. PubMed PMID: 29175436.
110.   Sotelo-Silveira M, Chavez Montes RA, Sotelo-Silveira JR, Marsch-Martinez N, de Folter
S. Entering the next dimension: plant genomes in 3D. Trends Plant Sci. 2018;23(7):598-612.
Epub 2018/04/29. doi: 10.1016/j.tplants.2018.03.014. PubMed PMID: 29703667.
111.   Gendrel AV, Lippman Z, Martienssen R, Colot V. Profiling histone modification patterns
in plants using genomic tiling microarrays. Nat Methods. 2005;2(3):213-8. Epub 2005/09/17.
doi: 10.1038/nmeth0305-213. PubMed PMID: 16163802.
112.   Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
arXiv preprint arXiv:13033997. 2013.
113.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence
Alignment/Map format and SAMtools. Bioinformatics. 2009;25(16):2078-9. doi:
10.1093/bioinformatics/btp352. PubMed PMID: 19505943; PubMed Central PMCID:
PMCPMC2723002.
114.   Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic
features. Bioinformatics. 2010;26(6):841-2. doi: 10.1093/bioinformatics/btq033. PubMed PMID:
20110278; PubMed Central PMCID: PMCPMC2832824.
115.   Robinson JT, Thorvaldsdottir H, Winckler W, Guttman M, Lander ES, Getz G, et al.
Integrative genomics viewer. Nat Biotechnol. 2011;29(1):24-6. doi: 10.1038/nbt.1754. PubMed
PMID: 21221095; PubMed Central PMCID: PMCPMC3346182.
116.   Merchant N, Lyons E, Goff S, Vaughn M, Ware D, Micklos D, et al. The iPlant
collaborative: cyberinfrastructure for enabling data to discovery for the life sciences. PLoS Biol.
2016;14(1):e1002342. doi: 10.1371/journal.pbio.1002342. PubMed PMID: 26752627; PubMed
Central PMCID: PMCPMC4709069.

1355

 SUPPORTING INFORMATION

 S1 Text. Supplemental Methods.

 S1 Fig. (related to Fig 1) Assessment of purity of flow sorted endocycling nuclei. Maize root

 tip nuclei were isolated from the 1–3 mm root region and sorted on a BD InFlux flow sorter. A

 small sample from each of the three S-phase sort gates was re-analyzed to determine the purity of

 the sorted nuclei. Histograms of relative DNA content (DAPI fluorescence) from re-analyzed

 sorted nuclei are overlaid for early (E), mid (M), and late (L) S-phase gates from the endocycle

 arc to show the separation between sorted samples. Similar separation was found for sorted early,

 mid and late nuclei from the mitotic cycle (see Supplemental Fig. 1 in [12]). The histogram of

 relative DNA content for the entire unsorted nuclei population (black line) is shown for

 reference.

 S2 Fig. (related to Fig 1) Genomic copy number analysis. Whole genome sequence data from

 sorted non S-phase 2C, 4C and 8C nuclei were used to assess copy number per DNA content

 across the genome. To better represent the copy number of repeat regions, the primary alignment

 location for each read pair – even those that map to multiple locations – were included in the

 analysis. (**A and B**) Histograms of the normalized read frequency ratios, calculated in 5-kb static

 windows, for 2C/4C (**A**) and 8C/4C (**B**) nuclei. The black dashed lines indicate the overall mean

 and the red dashed lines indicate ± 2 S. D. from the mean. (**C**) The 8C/4C read frequency ratios

 plotted as a function of genomic location, which shows that the values outside ± 2 S. D. all occur

 as singleton 5-kb windows. (**D and E**) We used consensus sequences for 45S rDNA and

 knob180 (**D**), and for 5S rDNA, *TR-1*, *CentC* and *CRM1–4* families (**E**) to individually query all

1389   of the trimmed whole genome sequence reads using BLAST software and a non-stringent E

1390   value to allow for variants of each repeat (S1 Text). The mean percentage of total reads that align

1391   to each repeat type was calculated for three biological replicates of 2C, 4C and 8C data. Black

1392   dots represent the individual biological replicate values. The apparent slight under-replication of

1393   several elements (e.g., *knob180* and *CRM2*) is not statistically significant.

1394

S3 Fig. (related to Figs 1 and 3) Example of Repli-seq data processing with *Repliscan*. An

example region from CEN 10 is shown to illustrate that the pre-replicative 2C reference data

effectively normalizes spikes of signal in the S-phase data. (**A and B**) Read densities were

calculated in 3-kb windows for the 2C reference (**A**) and each S-phase sample (endocycle late

profile shown; **B**). After excluding blacklist regions (e.g. unmappable and multi-mapping

regions), reads were scaled for overall sequence depth in each sample. (**C**) Scaled reads in each

S-phase sample were normalized by making a ratio to 2C reference scaled reads in each 3-kb

window. (**D**) Replication signal profiles were smoothed using a Haar wavelet transform to

remove noise without altering peak boundaries.

1404

S4 Fig. (related to Fig 1) Pearson's correlation coefficient values between individual

biological replicates of mitotic and endocycle Repli-seq data. (**A and B**) Biological replicates

(BR) of early (E), mid (M) and late (L) Repli-seq data for the mitotic cycle (Mit; panel **A**) and

endocycle (En; panel **B**) was analyzed independently using *Repliscan* [14]. The agreement

between biological replicates was assessed by calculating Pearson's correlation coefficients. (**C**)

The Pearson's correlation coefficients for E, M, L data between mitotic cycle and endocycle.

S5 Fig. (related to Fig 2) Boxplots of differences in early, mid and late replication signal

profiles for each chromosome. Differences in replication (dRT) signal were calculated by

subtracting the mitotic signal from the endocycle signal for early (E), mid (M) and late (L) S-

phase fractions in each 3-kb window across the genome. The distributions of dRT signal values

are represented as violin plots for each chromosome. Median values are indicated by colored

squares and 1.5 x IQR of the distribution is indicated by colored whisker lines. Dashed lines

indicate the thresholds used in subsequent steps for identifying RATs ($\geq 10\%$ and $\geq 25\%$ of the

total difference range; S1 Table).

1419

S6 Fig. (related to Fig 2) Additional examples of non-CEN RATs. (A–F) Example regions on

chromosomes 1 (**A**), 3 (**B**), 4 (**C**), 5 (**D**), 6 (**E**) and 7 (**F**) that include RATs. See main text Fig 2

legend for description. Dashed boxes denote regions with some level of RT difference in which

the magnitude of the difference did not meet our $\geq 25\%$ criterion (boxes labeled "a" in panels **A**,

B, **C** and **F**), or in which the change in one S-phase fraction was not compensated by an opposite

change in at least one other S-phase fraction (boxes labeled "b" in panels **C** and **D**).

1426

1427 **S7 Fig. (related to Fig 2 and Table 1) Permutation analysis of the percentage overlap of**

1428 **non-CEN RATs and genes. (A–D)** The percentage of RATs that overlap genes (**A and B**) or

1429 expressed genes (**C and D**) was calculated for non-CEN RATS and 1000 randomly shuffled sets

1430 (see Methods). The observed percentage for RATs (red line) and the frequency distribution of

1431 the random sets (green) are plotted.

1432

S8 Fig. (related to Fig 2) Activating and repressive histone marks in non-CEN RATs.

To assess whether changes in selected histone modifications related to gene transcription and

chromatin accessibility occur in RATs, ChIP-seq data was generated for H3K56ac and

H3K4me3 (active transcription and early replication) and H3K27me (repressive transcription and

facultative heterochromatin) from sorted non S-phase 2C, 4C and 8C nuclei. (**A–C**) The

distributions of fold enrichment values for H3K56ac (**A**), H3K4me3 (**B**) and H3K27me3 (**C**)

peaks in expressed and non-expressed genes (see S1 Text) in 2C, 4C and 8C nuclei are plotted as

boxplots for Later-to-Earlier and Earlier-to-Later RATs and their corresponding randomly

shuffled sets (see Methods). Asterisks indicate statistically significant differences by the non-

parametric Steel-Dwass-Critchlow-Fligner test at the following P value levels: ***, $P < 0.0001$;

**, $P < 0.001$; *, $P < 0.01$. The increase in the fold enrichment of H3K56ac for expressed genes

in Earlier-to-Later RATs (panel **A**) may be associated with increases in peak enrichment we

observed near the 3' end of some genes. (**D**) The count and percentage of expressed and non-

expressed genes with each histone modification shown in the boxplots in panels **A–C**. The

8C/2C ratio of genes with each mark is also shown to demonstrate there is very little change in

the number of genes with each mark. The total number of expressed and non-expressed genes in

each RAT or random category are shown at the bottom for reference.

1450

S9 Fig. (related to Fig 2) Gene ontology analysis of genes in non-CEN RATs. Using the

Plant GO slim ontology subset, we identified 44 significant GO terms in the biological process

(P), molecular function (F), and cellular component (C) GO categories that were enriched in

expressed genes (S1 Text; S3 Spreadsheet) in Earlier-to-Later RATs. Genes in the corresponding

randomly shuffled set shared a few of the significantly enriched cellular component terms as

genes in Earlier-to-Later RATs, suggesting that these terms may be related to common

components of the root, and not RATs specifically. The total number of expressed genes in each

input gene list was as follows: Later-to-Earlier RATs, 52; LtoE shuffle1 random regions, 68;

Earlier-to-Later RATs, 292; EtoL shuffle1 random regions, 275.

1460

1461   **S10 Fig. (related to Fig 2) AT content composition in non-CEN RATs. (A)** The distributions

1462   of percent AT content, calculated in 3-kb static windows, for Later-to-Earlier and Earlier-to-

1463   Later non-CEN RATs and the corresponding random shuffle sets are plotted as boxplots. Values

1464   outside the boxplot whiskers (1.5 x IQR) are represented as grey dots. The dashed line indicates

1465   the genome wide median value.

1466

1467   **S11 Fig. (related to Fig 3) Uniquely mapping Repli-seq reads in centromeres.** The average

1468   percentage of centromeric reads that map to unique locations is shown for each replication

1469   timing sample. Black dots represent the individual values for biological replicates.

1470

1471   **S12 Fig. (related to Fig 3) Replication signal profiles and RATs in complex and simple**

1472   **centromeres.** 5-Mb regions are shown for complex CENs 2, 3, 4, 5, and 8 and simple CEN 7.

1473   See main text Fig 3 legend for description.

1474

S13 Fig. (related to Fig 3) Timing differences in centromeres and pericentromeres. Timing

differences (endocycle minus mitotic) between early (**A and D**), mid (**B and E**) and late (**C and**

F) profiles for each centromere and corresponding pericentromere (± 1 Mb) were calculated in

100-kb static windows. In panels **D**, **E**, and **F** asterisks indicate difference values from windows

where an Earlier-to-Later-CEN RAT extends past the called CEN boundary [38] into the

pericentromere; open circles indicate windows that contain a non-CEN Earlier-to-Later RAT that

met our compensation criteria.

1482

1483 **S14 Fig. (related to Fig 4) Replication times for all genomic features in complex and simple**

1484 **centromeres and corresponding pericentromeres.** All elements within centromeres and

1485 pericentromeres are included, not just those that cover at least half of a 3-kb window, as in Fig 4.

1486 See main text Fig 4 legend for description.

1487

S15 Fig. (related to Fig 4) Activating and repressive histone mark peaks of enrichment in

centromeres. ChIP-seq data were generated for H3K56ac, H3K4me3 (active transcription) and

H3K27me (repressive transcription) from 2C, 4C and 8C nuclei. (**A–C**) The fold enrichment

values for peaks in expressed and non-expressed genes for H3K56ac (**A**), H3K4me3 (**B**) and

H3K27me3 (**C**) in 2C, 4C and 8C nuclei. Red lines indicate the median value. (**D**) The number

of expressed and non-expressed genes with each mark in 2C, 4C and 8C nuclei.

1494

S16 Fig. (related to Fig 5) H3K9me2 fold enrichment relative to DNA content in complex

and simple centromeres. We used the ChIP-seq datasets from 2C, 4C and 8C nuclei to estimate

the H3K9me2 average fold enrichment relative to DNA content by calculating the percent of

total H3K9me2 reads found in a given centromere (**A and B**) using coordinates from [38] or

pericentromere (**C and D**) and dividing by the percent of total input reads corresponding to that

centromere or pericentromere. Black dots represent the individual values from biological

replicates.

S17 Fig. (related to Fig 5) CENH3 localization and enrichment in mitotic and endocycling

centromeres. (**A**) CENH3 localization patterns for 2C, 4C and 8C nuclei for CEN 1–CEN 8. (**B**)

CENH3 average fold enrichment relative to DNA content for complex and simple centromeres.

See main text Fig 5 legend for CEN 9 and CEN 10 localization patterns and description.

1506

S18 Fig. (related to Fig 5) ChIP-qPCR antibody validations for anti-CENH3 and anti-

H3K9me2 antibodies. The percentage of input (%IP) was calculated for various antibody

dilutions and primer sets for the *Zea mays* anti-CENH3 antibody (**A**) and anti-H3K9me2

antibody (**B**). Black dots in panel **A** represent the individual values from two biological

replicates. Positive control primer sets (*CRM2* and Copia retrotransposons) and negative control

primer sets (18S rDNA and Actin1 UTR) were used. The no antibody control (NoAB) values are

too small to see on the graph. See S1 Text for Supplemental Methods.

1514

S1 Table. (related to Fig 2) Replication timing signal differences and thresholds. The

difference in replication timing signal between mitotic and endocycle profiles (endocycle minus

mitotic) was calculated for each 3-kb window across the genome. The maximum negative

difference value, which indicates a higher signal in the mitotic cycle, and the maximum positive

difference value, which indicates a higher signal in the endocycle, are shown for early and late

profiles. The average total difference range between these two values was used to calculate

percentage thresholds for identifying RATs (see S2 Table and main text).

1522

S2 Table. (related to Fig 2) Summary statistics of preliminary RAT calling steps. The

thresholds from S1 Table ($\geq 10\%$ or $\geq 25\%$) were used to identify regions with RT difference in

early or late S phase that were compensated by difference(s) with an opposite sign in one or both

of the other two S-phase fractions (early + mid or mid + late) with greater than or equal to the

same magnitude. The count, minimum, maximum and median region size, and the total coverage

of the B73 RefGen_v4 genome are shown. Final robust RATs included at least one core region

with a $\geq 25\%$ RT difference, but immediately adjacent regions of $\geq 10\%$ differences were

merged together with the $\geq 25\%$ regions to identify larger regions of contiguous change.

1531

1532   **S3 Table. (related to Fig 2) Gene summary in non-CEN RATs.** The percent of RATs that

1533   contain genes, the total number of genes and expressed genes and the mean gene count per RAT

1534   are shown.

1535

S4 Table. (related to Fig 3) Permutation analysis results for gene and TE coverage in non-

CEN RATs. The permutation P values derived from calculating percent coverage in 1000

random permutations of each RAT set (e.g. see S7 Fig). All permutation P values shown are

associated with a test for whether the observed percent coverage value is greater than expected

by chance, unless marked "NEG" which indicates the P value is associated with a test for

whether the observed percent coverage value is less than expected by chance.

1542

S5 Table. (related to Fig 3) Cumulative RAT coverage in centromeres.

The cumulative coverage and number of RATs called in each centromere are shown. For

reference, the previously determined centromere sizes are shown [38], as well as the sizes after

unmappable regions are subtracted out. There are also some unmappable regions of unknown

size missing from the genome assembly [38], which we cannot account for here.

S6 Table. (related to Fig 4) Compensated timing shifts in complex centromeres and

corresponding pericentromeres. We calculated the total number of 3-kb windows in complex

centromeres and pericentromeres (± 1 Mb), as well as the number of windows that show timing

shifts that are compensated (threshold ≥ 10%) by equal and opposite shifts in the other two S-

phase fractions.

1553

S7 Table. (related to Fig 5) CENH3 average fold enrichment relative to DNA content in

centromeres. CENH3 fold enrichment relative to DNA content and the ratio of enrichments

between 4C and 2C and 8C and 4C are shown for each centromere. Fold enrichment values are

the mean ± S. D. of three biological replicates for 2C and 8C and two biological replicates of 4C.

See main text Fig 5 legend for further description. Two sets of theoretical ratio values are also

presented. The first set, labeled "proportional redeposition", corresponds to the hypothesis that

CENH3 is diluted relative to total DNA during replication, and is then redeposited to a level

proportional to the DNA content during the subsequent gap phase. The second set, labeled "no

redeposition", corresponds to an alternate hypothesis that CENH3 is diluted relative to total

DNA during replication, and is not redeposited in the subsequent gap phase.

1564

1565 **S1 Spreadsheet. (related to Figs 1–5) Mapping statistics and data availability for all**

1566 **included datasets.**

1567

1568 **S2 Spreadsheet. (related to Figs 2 and 3) RAT regions list.**

1569

1570 **S3 Spreadsheet. (related to Figs 2 and 3) Genes found in RATs.**

Figure

Figure

Figure

Figure

Figure