Adaptive divergence generates distinct plastic responses in two closely related *Senecio* species

Greg M. Walter[1,2]*, James Clark[1,3], Antonia Cristaudo[4], Delia Terranova[4], Bruno Nevado[3,5], Stefania

Catara[4], Momchil Paunov[6], Violeta Velikova[7], Dmitry Filatov[3], Salvatore Cozzolino[8], Simon J. Hiscock[3] and

Jon R. Bridle[1,9]

[1]University of Bristol, School of Biological Sciences, Bristol BS8 1TQ, UK

[2]Current address: Monash University, School of Biological Sciences, Melbourne 3800, Australia

[3]University of Oxford, Department of Plant Sciences, Oxford, OX1 3RB, UK

[4]University of Catania, Department of Biological, Geological and Environmental Sciences, Catania 95128,

Italy

[4]Current address: Universidade de Lisboa, Centre for Ecology, Evolution and Environmental Changes,

Lisboa 1749-016, Portugal

[6]Sofia University St. Kliment Ohridski, Faculty of Biology, Sofia 1164, Bulgaria

[7]Bulgarian Academy of Sciences, Institute of Plant Physiology and Genetics, Sofia 1113, Bulgaria

[8]University of Naples Federico II, Department of Biology, Naples 80126, Italy

[9]Current address: University College London, Department of Genetics, Evolution and Environment, London

WC1E 6BT, UK

* Corresponding Author: Greg M. Walter

18   Email: greg.walter@monash.edu

Abstract

The evolution of plastic responses to external cues allows species to track the environmental variation they regularly experience. However, it remains unclear how plasticity evolves during adaptation. To test whether distinct patterns of plasticity is associated with recent adaptive divergence, we quantified plasticity for two closely related but ecologically divergent Sicilian daisy species (*Senecio,* Asteraceae). We sampled c.40 genotypes of each species from natural populations on and around Mt Etna and then reciprocally transplanted multiple clones of each genotype into four field sites along an elevational gradient representing each species' native range, the edge of their range, and conditions outside their native range. At each elevation we quantified survival and measured leaf traits that included investment (specific leaf area), morphology, chlorophyll fluorescence, pigment content and gene expression. As evidence of adaptive divergence, both species performed better at their native site and better than the species from the other habitat. Traits and differentially expressed genes that changed with elevation in one species often showed little change in the other species, or changed in the opposite direction. Adaptive divergence is therefore associated with the evolution of distinct plastic responses to environmental variation, despite these two species sharing a recent common ancestor.

Keywords: adaptation, adaptive divergence, differential gene expression, environmental sensitivity, evolutionary history, phenotypic plasticity

Introduction

The resilience of many natural populations to novel or changing environments relies on their ability to adjust their phenotype to track local conditions (Chevin et al. 2010). This phenotypic plasticity generates different phenotypes from the same genotype depending on the environment to which it is exposed (Via et al. 1995; Ghalambor et al. 2007; Charmantier et al. 2008). The ability for plasticity to help a species track environmental variation is shaped by selection within environments routinely experienced by that given species (Ghalambor et al. 2007). Plasticity is therefore only likely to buffer populations by maintaining fitness when exposed to familiar environmental regimes (Bradshaw 1965; Schlichting 1986; Baythavong and Stanton 2010), which will likely affect how populations respond to novel environmental conditions (Chevin and Hoffmann 2017). Characterising plasticity for closely related, but ecologically divergent species can test to what extent adaptive divergence is also associated with divergence in phenotypic plasticity.

The effect of adaptive divergence on plasticity will depend on how selection interacts with plasticity (de Jong 2005; Radersma et al. 2020). In particular, phylogenetic relatedness (Pigliucci et al. 1999; Kellermann et al. 2018), ecology (Kulkarni et al. 2011) and the predictability of the environment (Oostra et al. 2018) can determine the nature and amount of variation in plastic responses among given taxa, as well as the range of environments within which plastic responses are adaptive. Although contrasting environments are expected to select for different forms and different magnitudes of plasticity (Schlichting 1986; Donohue et al. 2001), it is not known to what extent a shared common ancestry will constrain such divergence in plasticity. Few studies have assessed whether adaptation to contrasting environments causes not only adaptive divergence, but also divergence in plastic responses when exposed to the same environmental variation.

Assaying variation in gene expression across environments can reveal key aspects of the genomic basis of plasticity. As allelic (sequence changes in regulatory genes) or epiallelic (e.g. DNA methylation, chromatin remodelling, post-transcriptional modifications) variation underlying trait plasticity becomes fixed during adaptation to a particular environment, differences in plasticity are likely to arise among populations

occupying contrasting environments (Gibson and Wagner 2000; Shaw et al. 2014). Highly predictable

environments will reduce genetic variation in plasticity through stabilising selection acting either on the

genetic regulators (e.g. transcription factors) or long-term epiallelic changes, such as transgenerational DNA

methylation (Colicchio et al. 2015; Oostra et al. 2018). If such stabilising selection varies among

environments, divergence in phenotypic plasticity will occur. Determining the gene expression profiles for

closely related but ecologically divergent species can therefore reveal how plasticity evolves during adaptive

divergence.

In this study, we first use a common garden experiment to identify physiological differences between two

closely related species of *Senecio* that inhabit contrasting elevations on Mt. Etna, Sicily. We then quantify

variation in survival and phenotypic plasticity across four field transplant sites that lie within and beyond the

elevational range of either species. *Senecio chrysanthemifolius* (**Fig. 1a**) is a short-lived perennial

(individuals generally live for less than two years) with highly dissected leaves that occupies disturbed

habitats (e.g., vineyards, abandoned land and roadsides) in the foothills of Mt. Etna c. 400-1,000m a.s.l

(above sea level) and at similar elevations throughout Sicily. By contrast, *S. aethnensis* (**Fig. 1b**) is a longer-

lived perennial with entire, glaucous leaves and is endemic to lava flows c. 2,000-2,600m a.s.l on Mt. Etna

that are covered by snow each winter. These two species are proposed to have diverged recently, possibly

c.150,000 years before present, around the same time that the uplift of Mt. Etna created the high elevation

environment in which *S. aethnensis* is found (Chapman et al. 2013; Osborne et al. 2013). Their recent shared

ancestry is reflected by very low genetic divergence across the genome, despite large differences in habitat,

phenotype and life history (Chapman et al. 2016). In particular, rapid divergence in leaf shape and dissection

is key feature of speciation in this system (Brennan et al. 2009), as well as in other *Senecio* species (Walter et

al. 2018).

Fig. 1 (a) *Senecio chrysanthemifolius* occupies disturbed habitats below c.1,000m a.s.l, and has thin, dissected leaves. **(a)** *Senecio aethnensis* inhabits lava flows and has thicker, smooth-margined leaves with a thick waxy cuticle. **(c)** Map of sampling locations (*S. chrysanthemifolius*: green circles; *S. aethnensis*: blue circles) and transplant sites (yellow squares). Sampling locations for *S. aethnensis* and the transplant sites are labelled by their elevation. Inset map shows the location of the study system within Europe.

We predicted that: (1) The occupation of contrasting habitats by these species would be reflected by differences in physiology in a common garden, suggesting functional trait divergence; (2) As evidence of adaptive divergence, species would perform better at their native elevations than beyond their existing range, and better than the other species within their native habitat; (3) Despite their recent common ancestry, and reflecting their adaptive divergence into contrasting habitats, we predicted that the two species would show different patterns of plasticity at both the phenotypic and gene expression level. To test these hypotheses, we sampled 79 genotypes of *S. chrysanthemifolius* and *S. aethnensis* from natural populations and conducted two reciprocal transplant experiments in 2017 and 2019. The relative ease by which both species can be propagated from cuttings means we are able to control for genetic variation by assaying identical genotypes at multiple sites as clones, allowing biological replicates in analyses of both phenotypic traits and gene expression profiles. Both experiments involved transplanting multiple cuttings of each genotype to four

transplant sites across an elevational range that included the home range of each species, and two

intermediate elevations. Genotypes that were propagated for use in the 2017 experiment were sampled

directly from natural populations as cuttings, while genotypes propagated for the 2019 experiment were

raised in a common garden from seeds collected from individuals growing in natural populations. In each

transplant experiment, we quantified survival, growth, and leaf traits including leaf shape, gene expression,

photosynthetic activity, pigment content and leaf investment.

Materials and methods

Sampling natural populations

We sampled achenes (i.e. seeds), and took cuttings from naturally growing individuals of both species after

plants started flowering. Sampling was conducted in May-June 2017 for *S. chrysanthemifolius* and July 2017

for *S. aethnensis* because *S. aethnensis* grows more slowly and flowers later than *S. chrysanthemifolius*,

given its higher elevation habitat. For *S. chrysanthemifolius*, we sampled from 88 individuals from five sites

between 500-800 m.a.s.l (**Fig 1c**, **Table S1**). For *S. aethnensis*, we sampled from 87 individuals at four

different elevations (2,600m, 2,500m, 2,400m and 2,300m.a.s.l) on both the North and South slopes of Mt.

Etna (**Fig 1c**, **Table S1**). Although this species occurs at 2,000m, we avoided sampling below 2,300m to

avoid the risk of sampling hybrids associated with a stable hybrid zone present at 1,500-1,700m (Brennan et

al. 2009). To minimise the risk of sampling close relatives, most plants sampled were more than 10m apart.

We then used the seeds and cuttings sampled from the individuals in the natural populations for three

separate experiments. First, to test whether the two species have evolved differences in physiology under

common garden conditions, we germinated seeds and grew plants in the laboratory to measure a suite of

physiological traits. Second, to test for differences in plasticity, in 2017 we propagated the cuttings from the

field genotypes and reciprocally transplanted them at four elevations on Mt. Etna. Third, to verify the 2017

experiment, we conducted a second reciprocal transplant of cuttings in 2019, but this time we germinated

field-collected seeds and grew plants under common garden conditions, from which we propagated cuttings

and conducted the transplant.

Physiological differences between species under common garden conditions

To assess differences in physiology between the species, we grew plants from seeds in a growth cabinet (see

Methods S1). From eight maternal families of *S. chrysanthemifolius* we grew 34 individuals, and from ten

maternal families of *S. aethnensis* we grew 41 individuals. Seeds were scarified mechanically and placed in

petri dishes containing moist filter paper. Seedlings were transplanted into 70mm square pots with standard

potting mix and grown for two months before physiological measurements were taken. We used a Dualex+®

instrument (ForceA, France) to measure the leaf content of chlorophyll, anthocyanin and flavonol pigments.

Using an LCpro gas analyser (ADC BioScientific, UK), we measured photosynthetic gas exchange. Intrinsic

water use efficiency (iWUE) was calculated as the ratio between photosynthesis and stomatal conductance.

Chlorophyll fluorescence was measured using an IMAGING-PAM M-series chlorophyll fluorometer (Heinz

Walz GmbH, Germany). Using output from the fluorometer, we quantified two mechanisms of physiological

light defence of leaves (**Methods S1**): the unregulated [Y(NO)] and regulated [Y(NPQ)] dissipation of heat.

2017 field transplant experiment

In the glasshouse (Giarre, Italy), cuttings from all individuals sampled from natural populations (hereafter,

genotypes) were cut into 5cm stem segments, each possessing 2-3 leaf nodes. Each smaller cutting was then

dipped in rooting plant growth regulator for softwood cuttings (Germon® Bew., Der. NAA 0.5%, L. Gobbi,

Italy) and placed in a compressed mix of coconut coir and perlite (1:1) in one cell of an 84-cell tray. All

cuttings from each genotype were kept together in one half of a tray and tray positions were randomised

regularly. Trays were kept moist and checked regularly for cuttings that successfully produced roots

extending out of the bottom of tray. For each genotype, rooted cuttings were randomised into experimental

blocks and transplanted at four field sites. From the initial genotypes sampled, we transplanted 37 *S.*

chrysanthemifolius genotypes and 42 *S. aethnensis* genotypes.

We transplanted multiple cuttings of each genotype into three experimental blocks at each transplant site.

Cuttings were transplanted into grids of 20×7 plants, with the position of cuttings randomised with respect to

genotype, and separated from each other by 40cm (*S. chrysanthemifolius* block n=109; site n=327; total

N=1,308; *S. aethnensis* block n=130; site n=390; total N=1,560). Depending on the number of cuttings that

successfully produced roots, we transplanted 6-15 cuttings per genotype at each transplant site (see **Table**

S1). The two species were transplanted at different times (*S. chrysanthemifolius* in June-July 2017; *S.*

aethnensis in August 2017) because seasonal constraints meant that sampling from natural populations of *S.*

aethnensis was only possible a month after *S. chrysanthemifolius* had been sampled. Following each

transplant, cuttings were watered daily for three weeks to encourage establishment. To prevent death during

high temperatures in July-August (consistently >35ºC), we watered cuttings daily during this period, which

allowed us to assess the phenotypic responses of genotypes to what were still stressful conditions. We

recorded mortality approximately every two weeks and measured the phenotypic traits of all plants at a single

time point when both species showed substantial post-transplant growth (November 2017).

In 2019, we repeated the 2017 experiment, but transplanted cuttings of genotypes that were grown from

field-collected seed in the glasshouse at the same time for both species, and in the same experimental blocks.

This meant that the 2017 experiment reflects differences among adult genotypes sampled directly from

natural populations and transplanted at different points in the season. By contrast, the 2019 experiment

represents cuttings propagated from genotypes raised from seed in a common garden (glasshouse).

2019 field transplant experiment

We germinated five seeds collected from multiple genotypes (*S. aethnensis n=25*; *S. chrysanthemifolius*

n=21) that were sampled from the natural populations using the protocol described above. After one week,

one seedling from each maternal genotype was chosen at random and placed in 14cm pots containing

standard potting media and left to grow under supplemental lighting (25W LED tubes; TSA Technology,

Italy). When plants reached c.30cm high, we removed all branches and propagated cuttings for the transplant

using the same protocol as the 2017 experiment. Plants were left to re-grow, and then we took a second round

of cuttings to increase replication in the transplant experiment.

We prepared the field sites in the same manner as in 2017 and transplanted multiple cuttings of each

genotype into four experimental blocks at each elevation. Cuttings were transplanted into grids of 20×7 plants

and randomised with respect to species and genotype (i.e. both species were planted together). We

transplanted the first round of cuttings 16th April 2019 (3 cuttings/genotype/block; $n=139$ plants/block;

N=1,668 plants), and the second round of cuttings 24th May (2 cuttings/genotype/block; $n=92$ plants/block;

N=368 plants). Overall, we transplanted 2,036 plants. Initial watering was daily for three weeks, after which

it was reduced to once per week to maintain a more natural watering regime than the 2017 experiment.

Field transplant sites

Field transplant sites were at four elevations (500m, 1,000m, 1,500m and 2,000m a.s.l) along a transect on

the south-eastern side of Mt. Etna (**Fig. 1c**). The 500m site was located in a garden among fruit trees, the

1,000m site on an abandoned vineyard among *Quercus ilex*, the 1,500m site among an apple and pear

orchard, and the 2,000m site surrounded by pine trees on a lava flow from 1983. Both the native elevations

(500m for *S. chrysanthemifolius* and 2,000m for *S. aethnensis*) were located less than 1km from natural

populations. Plants of both species were often observed at intermediate elevations, but were never observed

within the native range of the other species. There is an elevational transition in soil type from a silty sand at

elevations between 500m and 1,500m, to volcanic sand at 2,000m. At each transplant site four data loggers

(Tinytag Plus, Gemini Data Loggers, UK) recorded temperature hourly. We took three soil samples at each

transplant site, which were analysed for 21 variables that included nutrients, salts and ions (Nucleo Chimico

Mediterraneo laboratories, Catania, Italy). To analyse the soil data, we used Multi-Dimensional Scaling

(MDS) to calculate the scaled distance between the soil samples taken at all transplant sites.

Characterising leaf morphology and pigment content

To characterise leaf morphology, we sampled and pressed young but fully expanded leaves from each cutting

once they showed extensive growth (at least three months after transplantation). For the 2017 experiment, we

sampled 3-5 leaves per cutting in November, and for the 2019 experiment we sampled two leaves in

September. Leaves were weighed, and then scanned and morphology quantified using the program Lamina

(Bylesjo et al. 2008), which generates estimates of leaf area, perimeter and the number of indentations (leaf

serrations and lobes). To estimate the density of indentations along the leaf margin, we standardised the

number of indentations by the perimeter. To estimate leaf complexity we calculated perimeter^2/area, where

lower numbers indicate less complex (i.e. more entire) leaves. As measures of leaf investment we included

leaf area, and calculated Specific Leaf Area (SLA $= \frac{leaf\ area}{leaf\ weight}$), where greater values represent thinner

leaves. For the 2019 field experiment, we used a Dualex Scientific+ instrument (Force-A, France) to quantify

the concentration of leaf chlorophyll and flavonol pigments. Flavonols are secondary metabolites that help to

reduce oxidative stress caused by abiotic (e.g. temperature or light) and biotic (e.g. herbivory) stressors

(Mierziak et al. 2014).

Quantifying chlorophyll fluorescence (2017 field experiment)

To quantify photosynthetic capacity across elevation for both species, we measured chlorophyll *a*

fluorescence, which estimates the efficiency of the photosynthetic response to intense light. We selected five

genotypes at random from each species and measured chlorophyll fluorescence on four cuttings per genotype

at each elevation (site *n*=40 plants, total N=160). We took measurements at two transplant sites each day,

completing all four sites within one week in October 2017. For each cutting we measured four leaves, and to

temporally replicate measurements we measured the same cuttings at each site on a second day. To take

measurements, we put leaf clips on four leaves of each plant and dark-adapted the plants for 30 minutes by

covering them with large black plastic containers. We then took fluorescence induction curve measurements

for 2 seconds at 3,500μmol s^{-1}m^{-2} photosynthetic photon flux density from each leaf (clip) using a Handy

PEA instrument (Hansatech Instruments Ltd., UK). We calculated PI$_{total}$, the total performance of

photosystem I and II (**Methods S1**).

Statistical analyses of survival and univariate plasticity

We first tested for significant differences in survival across elevation using mixed effects cox proportional

hazards models from the *coxme* v2.2-14 (Therneau 2012) package in R v3.6.1 (R Core Team 2019). Survival

at the end of the experiment was used as the response variable (censored at the last point of data collection),

and we included transplant elevation and species as fixed effects, and genotype and experimental block as

random effects (see equation 1).

We then quantified plasticity as the change in leaf morphology, pigment content and chlorophyll

fluorescence across elevation (all leaf traits were first averaged for each clone) using the linear mixed model

within the package *lme4* v1.1-23 (Bates et al. 2015)

$$y_{ijklm} = T_i + S_j + T_i \times S_j + T_i \times G_{k(j)} + B_{l(i)} + e_{m(ijkl)} \, . \tag{1}$$

Changes in the response variable across transplant elevation were modelled by the *j*th species (S_j) in the *i*th

transplant elevation (T_i) and their interaction ($T_i \times S_j$), which were all included as fixed effects. Random

effects included the interaction between transplant elevation and genotype $T_i \times G_{k(j)}$, which accounted for

differences among genotypes at each elevation, and experimental block within each environment ($B_{l(i)}$). The

residual error variance was captured by $e_{m(ijkl)}$. Separate implementations of equation 1 were used for each

(normally distributed) univariate response variable of interest (y_{ijklm}). For each implementation of equation

1, we tested the significance of the interaction between transplant site and species using likelihood ratio tests.

To correct for multiple comparisons, we adjusted the P-value by multiplying by the number of tests, and

keeping α=0.05. To test whether differences in morphology between transplant sites were significant for each

species, we used *emmeans* v1.4 (Lenth 2019) to conduct pairwise t-tests adjusted for multiple comparisons.

Estimating multivariate plasticity

To quantify multivariate plasticity across elevation, we mean standardised each morphological trait (divided

by its mean). We then used a multivariate analysis of variance (MANOVA) to test for significant differences

in multivariate mean phenotype across elevation and between species. We analysed each experiment

separately by including the morphological traits as a multivariate response variable. We included species,

transplant elevation and their interaction as main effects. We included the interaction between transplant site

and genotype as the error term, which is the appropriate denominator to calculate F-ratios for the main effects

because it tests whether differences among species and transplant elevations are significantly greater than

differences among genotypes. To visualise multivariate differences, we used the output of the MANOVA to

calculate the D-matrix (**methods S1**), which represents differences in mean multivariate phenotype across

species and elevation. We also quantified how the two species differed in their direction of plasticity in

response to elevation by calculating the vector of plasticity ($\Delta\bar{x}$) for each species using

$$\Delta\bar{x} = \bar{x}_{native} - \bar{x}_{novel}, \qquad (2)$$

where $\Delta\bar{x}$ is the vector (standardised to unit length) that represents the change in multivariate phenotype from

the native site ($\bar{x}_{native}$) to the novel elevation ($\bar{x}_{novel}$), for each species separately (Noble et al. 2019).

Calculating the angle that separates $\Delta\bar{x}$ for each species quantifies the difference in the direction of plasticity

between species.

Sampling of plant tissue and RNA extraction

To quantify gene expression variation across elevation and between species, we sampled young leaves from

12 genotypes of each species. For each genotype, we collected 3-4 newly emerged leaves (15-20mm in

length) from three cuttings at each transplant site (2 species × 12 genotypes × 3 clones × 4 elevations; total N

= 288) in summer (22nd to 26th July 2019), 86-90 days after the initial transplant, which was after cuttings

showed sufficient growth. All leaves for a cutting were placed in the same Eppendorf tube and stored in

RNAlater at -20ºC. We extracted RNA from each sample using QIAgen RNeasy kits. Library preparation

and 3' QuantSeq RNA sequencing was performed at the Oxford Genomics Centre on an Illumina NextSeq

platform, producing 75bp single-end reads.

Quantifying differential expression across transplant sites, genotypes, and species

A reference transcriptome was assembled for each species (**Methods S1**). Trimmed reads were mapped to

each species' reference transcriptome using *Salmon* v0.13.1 (Patro et al. 2017). Read counts were normalised

by transcript size, library size and filtered based on counts >5 across half of all samples. All estimates were

repeated using *DESeq2* (Love et al. 2014) and *limma/voom* (Law et al. 2014) according to

$$Counts \sim Species + Transplant\ Site + Species{\times}Transplant\ Site + Genotype\ . \qquad (3)$$

In *limma/voom*, the genotype was modelled as a random effect. For comparisons within species, each

treatment was compared with the home transplant site of each species, with differentially expressed genes

determined based on an adjusted p-value <0.01 (Benjamini and Hochberg 1995) and a log fold change >2 for

overexpression or <-2 for underexpression. In *DESeq2*, log fold changes were shrunk using the 'apeglm'

method and then used to rank genes based on high overexpression and underexpression.

Reference transcriptomes were annotated (**Methods S1**), and to identify functional categories for

differentially expressed genes, gene ontology enrichment analyses were performed using *topGO* v2.3.6

(Alexa and Rahnenführer 2019). Enrichment was determined with a Kolmogarov-Smirnoff test using genes

that were significantly differentially expressed (adjusted p values <0.01) between the native elevation and the

furthest transplant site.

Weighted network construction of differentially expressed genes

Weighted Gene Coexpression Network Analysis (WGCNA) identifies correlations of expression among all

genes and then forms modules of coexpressed genes within that network. Consensus modules were

constructed for each species (**Methods S1**). Each module was then summarized using its first principal

component as a module eigengene, which represents the expression profile of the module. We tested for a

correlation between each module eigengene in each species and transplant elevation. Each module was tested

for gene ontology enrichment in *topGO*, using Fisher's exact test.

Results

Physiological differences between species under common garden conditions

Under common garden conditions, both species show a similar ability to regulate heat dissipation from their leaves [Y(NPQ)] (**Fig. 2a**; t(65)=0.0116, adj. P=1.000), and similar values for non-regulated heat dissipation [Y(NO)] (**Fig. 2b**; t(65)=2.351, adj. P=0.1085). This indicates that photochemical energy conversion and mechanisms protecting leaves from heat are similar for both species under common garden conditions. *Senecio chrysanthemifolius* showed evidence of higher intrinsic water use efficiency than *S. aethnensis* (**Fig. 2b**; t(69)=3.875, adj. P=0.0010), suggesting that leaves of *S. chrysanthemifolius* conserve water more effectively. *Senecio aethnensis* showed slightly greater (but non-significant) leaf concentrations of chlorophyll (**Fig. 2c**; t(143.8)=2.085, adj. P=0.1940), and greater concentrations of flavonols (**Fig. 2d**; t(200.9)=4.399, adj. P<0.0001) than *S. chrysanthemifolius*. Greater concentrations of flavonols indicates a better ability for leaves of *S. aethnensis* to defend against high UV light and lower temperatures (Jugran et al. 2016).

Fig. 2 Physiological differences between species grown from seeds under common garden conditions in the laboratory. Filled circles represent *S. chrysanthemifolius* (*S.ch*), while unfilled circles represent *S. aethnensis* (*S.ae*). Gray circles represent individual plants measured and credible intervals represent the 95% confidence intervals of the mean. **(a)** Both species showed similar values for regulated heat dissipation, Y(NPQ). **(b)** *Senecio aethnensis* showed greater values for the unregulated dissipation of heat, Y(NO). **(c)** *Senecio chrysanthemifolius* showed higher intrinsic water use efficiency, while *S. aethnensis* showed higher leaf chlorophyll content **(d)** and a higher flavonol content **(e)**.

Survival of transplanted cuttings reflects adaptive divergence between species

The transplant sites experience contrasting climatic conditions associated with elevation, with extreme heat

(regularly exceeding 40°C) at 500m and 1,000m during summer, and extreme cold (regularly below 0°C) at

1,500m and 2,000m during winter (**Fig. 3a**). Soil profiles roughly separated the four transplant sites in a

linear fashion along the first axis (MDS1), which represented a transition in soil type as a reduction in

nutrients (amount of organic material, total nitrogen, cation exchange capacity and exchangeable ions) at

higher elevations (**Fig. 3b**; **Table S2**). The second axis (MDS2) characterised differences between the

1,000m site and the other sites, associated with greater concentrations of various salts at 1,000m (soluble

nitrates, calcium and magnesium; **Table S2**).

24
Fig. 3 Differences in environment for the four transplant sites at four elevations. **(a)** Average daily maximum
(solid lines) and minimum (dashed lines) for three data loggers at each site, for the duration of the transplant.
Gray shading represents one standard error in estimating the curves. Higher elevations remained below 40ºC
in the summer and dropped well below zero in the winter. **(b)** Differences in soil composition for 21 soil
variables captured by a multidimensional scaling analysis (**Table S2** presents the soil data).

Both experiments showed strong evidence for adaptive divergence between species whereby each species

survived consistently better at their home ('native') than the species from the other elevation extreme (**Fig.**

4a-b). Evidence of adaptive divergence was supported by a significant species×elevation interaction for both

experiments (2017: $\chi^2(3)=30.21$, P<0.00001, and 2019: $\chi^2(3)=251.00$, P<0.00001). *Senecio*

chrysanthemifolius showed significantly greater mortality at higher elevations away from its native 500m site

(**Fig. 4a-b**; **Table S3**). By contrast, *S. aethnensis* showed significantly greater mortality at lower elevations

away from its native 2,000m site (**Fig. 4a-b**; **Table S3**). In contrast to 2017, in 2019, *S. chrysanthemifolius*

suffered c.35% greater mortality at 500m (its native site) than at 1,000m (which is within the native range).

This is likely to be due to cuttings suffering under intense heat without well-developed root systems,

especially given they received less supplemental water compared to 2017. However, mortality in 2019 was

still c.20% greater at the novel elevation (2,000m) than at 500m, indicating better performance at the native

site.

Fig. 4 Differences in survival between species across all transplant elevations, for **(a)** 2017 and **(b)** 2019.
Solid lines represent *S. chrysanthemifolius*, and broken lines *S. aethnensis*. Grey shading represents the 95%
confidence interval around the mean survival of each species. *Senecio chrysanthemifolius* showed the lowest
survival at the highest elevation in both experiments, while *S. aethnensis* showed lower survival at all three
lower elevations.

Species differences in plasticity: Univariate trait changes across elevation

Both species showed phenotypic plasticity as a change in trait mean with elevation for most leaf morphology

traits **(Fig. 5)**, with patterns of plasticity that were consistent across both experiments. However, the degree

to which plastic changed the phenotype across elevation differed between species depended on the trait. Both

species showed similar increases in leaf area at lower elevations (**Fig. 5a-b**; species×elevation in 2017:

$\chi^2(3)=8.94$, adj. P=0.1202, and 2019: $\chi^2(3)=42.08$, adj. P<0.0001). However, *Senecio chrysanthemifolius*

showed a reduction in leaf complexity at higher elevations, contrasting with no change across elevation in *S.*

aethnensis (**Fig. 5c-d**; species×elevation in 2017: $\chi^2(3)=29.09$, adj. P<0.0001, and 2019: $\chi^2(3)=83.87$, adj.

P<0.0001). By contrast, *S. aethnensis* showed a reduction in leaf indentation at higher elevations, while *S.*

chrysanthemifolius showed no significant change in leaf indentation across elevation in 2017, but a

significant increase in leaf indentation at 2,000m in 2019 (**Fig. 5e-f**; species×elevation in 2017: $\chi^2(3)=29.07$,

adj. P<0.0001, and 2019: $\chi^2(3)=66.68$, adj. P<0.0001). Both species also showed a similar increase in SLA

(Specific Leaf Area) at lower elevations, but *S. aethnensis* showed a much greater increase at 500m and

1,000m (**Fig. 5g-h**; species×elevation in 2017: $\chi^2(3)=21.95$, adj. P=0.00027, and 2019: $\chi^2(3)=37.71$, adj.

P<0.0001), suggesting that *S. aethnensis* produced much thinner leaves than *S. chrysanthemifolius* at lower

elevations (and at all elevations in 2019).

In 2017, we measured chlorophyll fluorescence to calculate the total performance index (PI_{total}), which

estimates the capacity of the photosynthetic machinery. Although *S. chrysanthemifolius* showed no change in

PI_{total} across elevation, *S. aethnensis* showed a steady decline, suggesting reduced photosynthetic activity of

S. aethnensis at lower elevations (**Fig. 5i**; species×elevation $\chi^2(3)=24.59$, P<0.0001). In 2019, we measured

the leaf concentration of chlorophyll and flavonols. Compared to *S. aethnensis, S. chrysanthemifolius* showed

higher concentrations of both pigments. From high to low elevation, both species showed increases in

chlorophyll content, but a reduction at 500m (**Fig. 5j**; species×elevation: $\chi^2(3)=64.23$, adj. P<0.0001). The

two species show contrasting patterns of plasticity in flavonol content across elevation (**Fig. 5k**;

species×elevation: $\chi^2(3)=98.57$, adj. P<0.0001): although both species show similar values at 500m, flavonol

content in *S. chrysanthemifolius* decreases at higher elevations, while in *S. aethnensis* it increases at higher

elevations.

Fig. 5 Changes in univariate leaf traits across elevation for both species measured in 2017 (top row) and 2019
(bottom row). Filled circles, solid lines and upper case letters represent *S. chrysanthemifolius*, while unfilled
circles, dashed lines and lowercase letters represent *S. aethnensis*. Credible intervals represent 95%
confidence intervals for the estimate of the mean of each species at each elevation. Letters denote significant
differences between transplant sites calculated using pairwise tests conducted within each species and
adjusted for multiple comparisons (full statistical summaries are located in **Table S3**). Grey points represent
the mean of all cuttings for each genotype. Traits include: **(a-b)** leaf area, **(c-d)** leaf complexity, **(e-f)** number
of indents, **(g-h)** specific leaf area, **(i)** total photosynthetic performance, **(j)** chlorophyll content and **(k)**
flavonol content. Results are consistent across the 2017 and 2019 experiments, and show that the two species
display distinct plastic responses across elevation for leaf complexity, number of indents, specific leaf area,
photosynthetic performance and flavonol content.

Species differences in plasticity: Change in multivariate phenotype across elevation

To estimate multivariate plasticity, we quantified variation in leaf morphology between species and across

elevation by analysing all leaf traits using a multivariate analysis of variance (MANOVA). For both

experiments, we found significant species×elevation interactions (**Fig. 6;** 2017: Wilks' λ=0.375, $F_{3,298}$=29.19,

P<0.0001; 2019: Wilks' λ=1.330, $F_{3,176}$=22.97, P<0.0001). In both experiments, the first axis of **D** (**d_{max}**)

described >75% of the difference in multivariate phenotype, which primarily separated the two species,

generated by a negative correlation between leaf area and complexity (larger simple leaves versus smaller,

more complex leaves). The second axis (**d_2**) described 15-19% of the difference in multivariate

phenotype, which captured differences across elevation as reductions in most traits at higher elevations.

Outside its native range (i.e. at 500m), plasticity moved the multivariate phenotype of *S. aethnensis* away

from, rather than towards, the native phenotype of *S. chrysanthemifolius*. By contrast, *S. chrysanthemifolius*

genotypes outside their native range (i.e. at 2000m) shifted their phenotypes towards that expressed by *S.*

aethnensis at high elevations (**Fig. 6**). The vectors representing elevational plasticity ($\Delta\overline{x}$) differed between

species by 54° and 56° respectively for the 2017 and 2019 experiments (grey lines in **Fig. 6**), which suggests

the two species showed plastic responses to elevation that differ consistently in direction across years.

Species ○ *S. aethnensis*   ● *S. chrysanthemifolius*   Elevation (m.a.s.l) ● 500 ● 1,000 ● 1,500 ● 2,000

Fig. 6 Multivariate analysis (MANOVA used to calculate the D-matrix) of leaf morphology across elevation
for both species measured in **(a)** 2017 and **(b)** 2019. For both experiments, d_{max} represents a trade-off
between leaf size and complexity, and separates both species. By contrast, d_2 represents similar elevational
changes in most traits. Small filled circles represent genotypes of *S. chrysanthemifolius*, and unfilled circles
S. aethnensis. Large circles represent the multivariate mean at each elevation. Table inset shows the trait
loadings for both multivariate axes. Inset leaf images represent a genotype of each species with a value close
to the multivariate mean at the elevational extremes (500m and 2,000m). Gray lines represent the vector of
plastic responses ($\Delta\overline{x}$) across elevation for each species. The angle between plasticity vectors show that the
two species differ in the direction of plasticity by c.55°, suggesting distinct patterns of plasticity between
species.

Differential gene expression between species

Comparisons of gene expression between species for the 2019 transplant revealed that many genes were

differentially expressed between species at each transplant site (**Fig. 7a**). In total, the largest number of

differentially expressed genes between the two species was at 1,500m (332 genes) with the fewest at 500m

(279 genes). Despite observing the largest number of differentially expressed genes at 1,500m, functional

enrichment revealed that the two species show the fewest significantly enriched GO terms (9 terms) at

1,500m while the largest number of significantly enriched GO terms was at 2,000m (16 terms). Furthermore,

140 genes were differentially expressed between species at all transplant sites, which represents 42-50% of

the total number of differentially expressed genes at each site (and 29% of all genes differentially expressed

at all elevations). Therefore, almost half the genes that were differentially expressed between species

remained consistent across elevation. Despite this, no GO terms were significantly enriched between species

at all transplant sites (**Fig. 7b**). These results suggest that although some genes are consistently differentially

expressed between the two species, they are not related to known functional categories. This means that the

functional categories of genes showing large differences in gene expression between species at each

elevation are not shared across elevations, indicating that the two species differ in gene expression profiles in

response to elevation.

Differential gene expression across elevation within species

The two methods of estimating differential expression (*DESeq2* vs *limma/voom*) showed similar patterns of

gene expression variation, although *limma/voom* detected fewer strongly differentially expressed genes in

each species (**Fig. S1**). For both species, more genes were differentially expressed as genotypes were moved

further from their native elevations (**Fig. S1**). We identified c.350 loci (out of c. 19,000) per species that

showed differential expression between the extreme elevations. Only a small proportion of genes that were

differentially expressed between the native site and the most novel environment (i.e. the elevational

extremes) were common to both species, with 43% and 23 % respectively of overexpressed and

underexpressed genes showing similar patterns across elevation in both species (**Fig. 7c**). However, the

majority of differentially expressed genes showed a species-specific response to elevation providing evidence

for distinct patterns of plasticity in gene expression for the two species (**Fig. 7c**). This meant that most genes

showed responses to elevation that were distinct to each species: strong overexpression or underexpression in

one species contrasted with relatively unchanged expression across elevation in the other species (**Fig. 8**).

Functional enrichment analyses of differentially expressed genes between the elevational extremes revealed 9

significant GO terms in *S. chrysanthemifolius* and 20 in *S. aethnensis*. Only 2 functional categories of these

genes were shared between species, again suggesting that many genes responded in a species-specific manner

(**Tables S4-S5**).

Network reconstruction generated 13 network modules ranging in size from 185 to 3345 genes (**Table S6**). In

6 modules the module eigengene correlated with elevation in both species and 2 modules (**Table S6**: GN3

and GN8) showed a correlation with elevation unique to *S. aethnensis*. No gene modules that correlated with

elevation were specific to *S. chrysanthemifolius*. The two modules that changed in *S. aethnensis* but not *S.*

chrysanthemifolius were associated with functions related to pathogen detection and responses to light

intensity and UV.

Fig. 7

Contrasting patterns of gene expression plasticity between species. (**a**) Total numbers of differentially
expressed genes (-2 < lfc > 2) between species at each transplant site. Many genes that were differentially
expressed between the species were unique to each elevation. (**b**) Total numbers of significantly enriched
(Fisher's exact test, p < 0.05) GO terms between species at each transplant site. Only found significant
functional categories were found for differentially expressed genes at each elevation, which were rarely
shared across elevations. (**c**) Overlapping overexpressed and underexpressed genes between the home and
furthest transplant site in each species. Many genes that were differentially expressed across elevation were
unique to each species.

Fig. 8. Strong overexpression or underexpression in one species was typically reflected by little to no change in gene expression in the same gene for the other species. Normalised expression profiles across all transplant sites for *S. chrysanthemifolius* (solid lines and circles) compared to *S. aethnensis* (dashed lines and unfilled circles). This includes the genes (represented by different shapes) that most strongly underexpressed in *S. aethnensis* but not in *S. chrysanthemifolius* **(a)**, overexpressed in *S. aethnensis* but not in *S. chrysanthemifolius* **(b)**, and five genes strongly underexpressed in *S. chrysanthemifolius* but not in *S. aethnensis* **(c)** and overexpressed in *S. chrysanthemifolius* but not in *S. aethnensis* **(d)**.

Discussion

Given the two *Senecio* species on Mt. Etna are the result of recent ecological divergence (Chapman et al. 2013), we predicted that: (1) the two species would show differences in physiology when grown in common garden conditions; (2) they would perform better at their native elevations than beyond their existing range, and better than the other species within their native habitat; and (3) given their adaptation to contrasting environments, they would display different patterns of plasticity at the phenotypic and gene expression levels.

Consistent with our first prediction, both species showed distinct behaviour in the common garden

experiment (**Fig. 2**), indicating that adaptation to contrasting habitats has generated substantial differences in

physiology between the two species. Supporting our second prediction, both species performed better at their

native site than the other species. Each species also tended to perform best at their native site compared to

other elevations, with strong reductions in survival observed at the elevation furthest from their native site

(i.e. at the native site of the other species; **Fig. 4**). This provides strong evidence that these two species are

adapted to their natural habitats, which is consistent with another field experiment that demonstrated adaptive

divergence between these two species when transplanted as seedlings rather than as cuttings (Walter et al.

2021). Consistent with our third prediction, we found different patterns of plasticity in phenotype (**Figs. 5-6**)

and gene expression profiles (**Figs. 7-8**) between the two species.

Species differences in gene expression plasticity

The native conditions experienced by any given species can determine how gene expression will change in

response to environmental variation (Akman et al. 2016). We observed both differential expression between

species as well as between-species differences in the plasticity of gene expression along the elevational

gradient. Although differential expression between species was observed at each transplant site, we found

that only 29% (140 genes) of genes that were differentially expressed between species were common to all

transplant sites. This suggests that each species has recently evolved distinct patterns of gene expression

associated with their native habitats.

Within a species, plasticity in gene expression in this experiment is represented by genes that are

differentially expressed between their native elevation and other elevations. We found that only 33% of

genes that were differentially expressed across elevational extremes were shared by the two species. The

differences in plasticity between species are evident in the expression profiles of individual genes, where

genes that showed strong changes in one species were weaker or absent in the other species (**Fig 8**). This

meant that highly divergent patterns of gene expression between the species were rare and expression profiles

tended to reflect differences in the magnitude of the response between species, rather than responses in

opposite directions (**Fig 8**). We also found unique networks associated with the response of *S. aethnensis* to

elevation, suggesting that differential expression in specific gene networks underlies differences in plasticity

observed between the two species, particularly in response to biotic stressors and changes in light conditions

associated with higher elevation. Together, these results confirm that these species show distinct patterns of

gene expression responses to elevation, most often arising from species-specific patterns of expression at

similar loci, but with some loci and gene networks displaying contrasting patterns of expression between the

species.

Adaptive divergence creates differences in plasticity

Our results suggest that adaptive divergence between these species is not only associated with divergence in

mean phenotype and physiology under common garden conditions, but also divergence in patterns of

phenotypic plasticity and in the genetic basis of plasticity. It remains possible that genetic drift during the

formation of these two species could cause the species-specific differences in plasticity that we observed.

However, given substantial divergence in leaf form and physiology under common garden conditions, the

differences in plasticity of these same traits among species, and the higher fitness of each species at their

native versus novel habitats, it seems far more likely that adaptive divergence between *S. aethnensis* and *S.*

chrysanthemifolius is responsible for their distinct plastic responses (Taylor and Aarssen 1988; Emery et al.

1994; Donohue et al. 2001; Ho and Zhang 2018).

Other studies have shown that species from different genera can evolve distinct forms of plasticity

(Schlichting and Levin 1986). One study that compared closely related species of *Phlox* wildflowers found

inconsistent differences that could not be explained by ecology or relatedness (Schlichting and Levin 1984).

Here, by comparing plasticity for two closely related species in field experiments that include their natural

habitats, we show that adaptive divergence is associated with the evolution of different forms of plasticity.

We also show that the magnitude and direction of plasticity in response to elevation is trait dependent, in the

sense that the two species showed differences in the magnitude of plasticity across elevation in some traits

(e.g. leaf complexity and specific leaf area), and a contrasting pattern of plasticity in other traits (e.g.

leaf indents and flavonol content). This suggests that adaptive divergence can change both the magnitude and

direction of plasticity, which is likely to depend on how selection varies its action on different traits (Via

1993).

An ecological explanation for differences in plasticity

As a consequence of adapting to the environmental variation specific to their native elevations, these species

could show different patterns of plasticity because they only maintain plasticity in traits important for

tracking variation within their native environment. For example, as a perennial that grows from rootstock

after winter each year, it is likely that the new leaves *S. aethnensis* produces each year possess different

characteristics that are optimal for the specific environmental conditions of that year, allowing *S. aethnensis*

to track environmental variation across years. In this case, plasticity in leaf traits would be critical to cope

with environmental variation across years at high elevations, and so would also show plasticity when

transplanted to lower elevations, even if such plastic changes were maladaptive. This would mean that how

selection acts on a trait in different environments will likely determine the levels of plasticity that evolves in

populations adapting to each environment.

As divergent selection drives trait divergence between the two species, it will likely also change the level of

plasticity in the traits that are diverging. Species will therefore show reduced plasticity in the genes (i.e. the

genes that show differential expression across elevations) and in the traits that need to be continually

expressed in their native habitat. This would occur if the plastic responses required to initially colonise a high

or low elevation habitat become genetically fixed (via genetic assimilation) because consistently strong

stabilising selection for a single phenotype is favoured (Waddington 1953). In this scenario, plasticity would

only be retained in traits that maintain fitness in response to the environmental variation experienced within

their native habitat. For example, *S. chrysanthemifolius* shows both greater leaf complexity than *S.*

aethnensis, as well as greater plasticity in this trait (**Fig. 5b & 5g**), while *S. aethnensis* shows a greater

number of indents, as well as greater plasticity in this trait (**Fig. 5c & 5h**). At high elevations *Senecio*

aethnensis is likely to experience consistently strong selection for less complex leaves, leading to a loss

of plasticity in this trait, perhaps via genetic assimilation. By contrast, *S. chrysanthemifolius* is likely to

experience selection for more complex leaves, and the greater variety of habitats that this species occupies

could maintain plasticity in this trait. Future work should identify how selection affects the evolution of

plasticity in different traits and environmental regimes (Schmitt et al. 1999; Pratt and Mooney 2013; McLean

et al. 2014), and could reveal how the fixation of alleles during adaptation affects trait plasticity (Schlichting

and Levin 1986; Forsman 2015).

Predicting species responses to novel environments

Understanding how plasticity evolves is important for understanding how species can respond to novel

environmental variation (Bradshaw 1965; Baythavong and Stanton 2010). For example, plants experiencing

strong stabilizing selection for a particular phenotype will become more specialised and evolve reduced

plasticity (Emery et al. 1994; Alpert and Simms 2002; Baythavong 2011). In other studies, high-elevation

populations showed reduced plasticity in flowering time (Schmid et al. 2017) and morphology (Emery et al.

1994) compared to low elevation populations. Similarly, we found that the high elevation species, *S.*

aethnensis, showed a steep decline in survival (**Fig. 4**), which was associated with stronger reductions in leaf

investment (**Fig. 5d & 6i**), lower photosynthetic activity (**Fig. 5e**) and reduced capacity for plasticity in leaf

traits to produce phenotypes more similar to the native phenotype of *S. chrysanthemifolius* at lower

elevations (**Fig. 6**). Therefore, the contrasting patterns of plasticity observed for the two species may be

caused by *S. aethnensis* exhibiting maladaptive plasticity at lower elevations (1,500m and below) as a

consequence of adapting to the high elevation habitat.

Limits to adaptive plasticity could threaten the persistence of populations exposed to novel environments

unless genotypic variation in plasticity (genotype×environment interactions) can allow rapid adaptation

(Lande 2009; Chevin et al. 2010). Understanding whether genetic variation in plasticity will allow natural

populations to adapt to ongoing environmental change requires quantifying plasticity and fitness across

environmental variation for genotypes from natural populations.

References

Akman, M., J. E. Carlson, K. E. Holsinger, and A. M. Latimer. 2016. Transcriptome sequencing reveals
population differentiation in gene expression linked to functional traits and environmental gradients
in the South African shrub *Protea repens*. New Phytol. 210:295-309.

Alexa, A. and J. Rahnenführer. 2019. Gene set enrichment analysis with topGO. R package version 2.38.1.

Alpert, P. and E. L. Simms. 2002. The relative advantages of plasticity and fixity in different environments:
when is it good for a plant to adjust? Evolutionary Ecology 16:285-297.

Bates, D., M. Machler, B. M. Bolker, and S. C. Walker. 2015. Fitting linear mixed-effects models using
lme4. Journal of Statistical Software 67:1-48.

Baythavong, B. S. 2011. Linking the spatial scale of environmental variation and the evolution of phenotypic
plasticity: selection favors adaptive plasticity in fine-grained environments. The American Naturalist
178:75-87.

Baythavong, B. S. and M. L. Stanton. 2010. Characterizing Selection on Phenotypic Plasticity in Response to
Natural Environmental Heterogeneity. Evolution 64:2904-2920.

Benjamini, Y. and Y. Hochberg. 1995. Controlling the False Discovery Rate: A Practical and Powerful
Approach to Multiple Testing. Journal of the Royal Statistical Society. Series B (Methodological)
57:289-300.

Bradshaw, A. D. 1965. Evolutionary significance of phenotypic plasticity in plants. Adv. Genet. 13:115-155.

Brennan, A. C., J. R. Bridle, A. L. Wang, S. J. Hiscock, and R. J. Abbott. 2009. Adaptation and selection in
the *Senecio* (Asteraceae) hybrid zone on Mount Etna, Sicily. New Phytol. 183:702-717.

Bylesjo, M., V. Segura, R. Y. Soolanayakanahally, A. M. Rae, J. Trygg, P. Gustafsson, S. Jansson, and N. R. Street. 2008. LAMINA: a tool for rapid quantification of leaf size and shape parameters. BMC Plant Biol. 8.

Chapman, M. A., S. J. Hiscock, and D. A. Filatov. 2013. Genomic divergence during speciation driven by adaptation to altitude. Mol. Biol. Evol. 30:2553-2567.

Chapman, M. A., S. J. Hiscock, and D. A. Filatov. 2016. The genomic bases of morphological divergence and reproductive isolation driven by ecological speciation in *Senecio* (Asteraceae). J. Evol. Biol. 29:98-113.

Charmantier, A., R. H. McCleery, L. R. Cole, C. Perrins, L. E. B. Kruuk, and B. C. Sheldon. 2008. Adaptive phenotypic plasticity in response to climate change in a wild bird population. Science 320:800-803.

Chevin, L. M. and A. A. Hoffmann. 2017. Evolution of phenotypic plasticity in extreme environments. Philosophical Transactions of the Royal Society of London Series B-Biological Sciences 372:20160138.

Chevin, L. M., R. Lande, and G. M. Mace. 2010. Adaptation, plasticity, and extinction in a changing environment: towards a predictive theory. PLoS Biol. 8:e1000357.

Colicchio, J. M., P. J. Monnahan, J. K. Kelly, and L. C. Hileman. 2015. Gene expression plasticity resulting from parental leaf damage in Mimulus guttatus. New Phytologist 205:894-906.

de Jong, G. 2005. Evolution of phenotypic plasticity: patterns of plasticity and the emergence of ecotypes. New Phytol. 166:101-117.

Donohue, K., E. H. Pyle, D. Messiqua, M. S. Heschel, and J. Schmitt. 2001. Adaptive divergence in plasticity in natural populations of *Impatiens capensis* and its consequences for performance in novel habitats. Evolution 55:692-702.

Emery, R. J. N., C. C. Chinnappa, and J. G. Chmielewski. 1994. Specialization, Plant Strategies, and
Phenotypic Plasticity in Populations of *Stellaria longipes* Along an Elevational Gradient. Int. J. Plant
Sci. 155:203-219.

Forsman, A. 2015. Rethinking phenotypic plasticity and its consequences for individuals, populations and
species. Heredity 115:276-284.

Ghalambor, C. K., J. K. McKay, S. P. Carroll, and D. N. Reznick. 2007. Adaptive versus non-adaptive
phenotypic plasticity and the potential for contemporary adaptation in new environments. Funct. Ecol.
21:394-407.

Gibson, G. and G. Wagner. 2000. Canalization in evolutionary genetics: a stabilizing theory? Bioessays
22:372-380.

Ho, W. C. and J. Z. Zhang. 2018. Evolutionary adaptations to new environments generally reverse plastic
phenotypic changes. Nature Communications 9:1-11.

Jugran, A. K., A. Bahukhandi, P. Dhyani, I. D. Bhatt, R. S. Rawal, and S. K. Nandi. 2016. Impact of altitudes
and habitats on valerenic acid, total phenolics, flavonoids, tannins, and antioxidant activity of
Valeriana jatamansi. Appl. Biochem. Biotechnol. 179:911-926.

Kellermann, V., A. A. Hoffmann, J. Overgaard, V. Loeschcke, and C. M. Sgrò. 2018. Plasticity for
desiccation tolerance across *Drosophila* species is affected by phylogeny and climate in complex
ways. Proceedings of the Royal Society B-Biological Sciences 285.

Kulkarni, S. S., I. Gomez-Mestre, C. L. Moskalik, B. L. Storz, and D. R. Buchholz. 2011. Evolutionary
reduction of developmental plasticity in desert spadefoot toads. J. Evol. Biol. 24:2445-2455.

Lande, R. 2009. Adaptation to an extraordinary environment by evolution of phenotypic plasticity and
genetic assimilation. J. Evol. Biol. 22:1435-1446.

Law, C. W., Y. Chen, W. Shi, and G. K. Smyth. 2014. voom: precision weights unlock linear model analysis tools for RNA-seq read counts. Genome Biology 15:R29.

Lenth, R. V. 2019. emmeans: Estimated Marginal Means, aka Least-Squares Means. *R Package*, https://CRAN.R-project.org/package=emmeans.

Love, M. I., W. Huber, and S. Anders. 2014. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biology 15.

McLean, E. H., S. M. Prober, W. D. Stock, D. A. Steane, B. M. Potts, R. E. Vaillancourt, and M. Byrne. 2014. Plasticity of functional traits varies clinally along a rainfall gradient in *Eucalyptus tricarpa*. Plant, Cell Environ. 37:1440-1451.

Mierziak, J., K. Kostyn, and A. Kulma. 2014. Flavonoids as important molecules of plant interactions with the environment. Molecules 19:16240-16265.

Noble, D. W. A., R. Radersma, and T. Uller. 2019. Plastic responses to novel environments are biased towards phenotype dimensions with high additive genetic variation. Proc. Natl. Acad. Sci. USA 116:13452-13461.

Oostra, V., M. Saastamoinen, B. J. Zwaan, and C. W. Wheat. 2018. Strong phenotypic plasticity limits potential for evolutionary responses to climate change. Nature Communications 9:1-11.

Osborne, O. G., T. E. Batstone, S. J. Hiscock, and D. A. Filatov. 2013. Rapid speciation with gene flow following the formation of Mt. Etna. Genome Biology and Evolution 5:1704-1715.

Patro, R., G. Duggal, M. I. Love, R. A. Irizarry, and C. Kingsford. 2017. Salmon provides fast and bias-aware quantification of transcript expression. Nat. Methods 14:417.

Pigliucci, M., K. Cammell, and J. Schmitt. 1999. Evolution of phenotypic plasticity a comparative approach in the phylogenetic neighbourhood of *Arabidopsis thaliana*. J. Evol. Biol. 12:779-791.

Pratt, J. D. and K. A. Mooney. 2013. Clinal adaptation and adaptive plasticity in *Artemisia californica*: implications for the response of a foundation species to predicted climate change. Global Change Biol. 19:2454-2466.

R Core Team. 2019 R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.

Radersma, R., D. W. A. Noble, and T. Uller. 2020. Plasticity leaves a phenotypic signature during local adaptation. Evolution Letters.

Schlichting, C. D. 1986. The Evolution of Phenotypic Plasticity in Plants. Annu. Rev. Ecol. Syst. 17:667-693.

Schlichting, C. D. and D. A. Levin. 1984. Phenotypic Plasticity of Annual *Phlox*: Tests of Some Hypotheses. Am. J. Bot. 71:252-260.

Schlichting, C. D. and D. A. Levin. 1986. Phenotypic plasticity: an evolving plant character. Biol. J. Linn. Soc. 29:37-47.

Schmid, S. F., J. Stocklin, E. Hamann, and H. Kesselring. 2017. High-elevation plants have reduced plasticity in flowering time in response to warming compared to low-elevation congeners. Basic Appl. Ecol. 21:1-12.

Schmitt, J., S. A. Dudley, and M. Pigliucci. 1999. Manipulative approaches to testing adaptive plasticity: phytochrome-mediated shade-avoidance responses in plants. The American Naturalist 154:S43-S54.

Shaw, J. R., T. H. Hampton, B. L. King, A. Whitehead, F. Galvez, R. H. Gross, N. Keith, E. Notch, D. Jung, S. P. Glaholt, C. Y. Chen, J. K. Colbourne, and B. A. Stanton. 2014. Natural Selection Canalizes Expression Variation of Environmentally Induced Plasticity-Enabling Genes. Mol. Biol. Evol. 31:3002-3015.

Taylor, D. R. and L. W. Aarssen. 1988. An Interpretation of Phenotypic Plasticity in *Agropyron repens*
(Graminae). Am. J. Bot. 75:401-413.

Therneau, T. 2012. Mixed Effects Cox Models. Mayo Clinic, Mayo Foundation for Medical Education and
Research.

Via, S. 1993. Adaptive phenotypic plasticity: target or by-product of selection in a variable environment?
The American Naturalist 142:352-365.

Via, S., R. Gomulkiewicz, G. de Jong, S. M. Scheiner, C. D. Schlichting, and P. H. Van Tienderen. 1995.
Adaptive phenotypic plasticity: consensus and controversy. Trends Ecol. Evol. 10:212-217.

Waddington, C. H. 1953. Genetic Assimilation of an Acquired Character. Evolution 7:118-126.

Walter, G. M., J. D. Aguirre, M. W. Blows, and D. Ortiz-Barrientos. 2018. Evolution of genetic variance
during adaptive radiation. The American Naturalist 191:E108–E128.

Walter, G. M., S. du Plessis, D. Terranova, E. la Spina, M. G. Majorana, G. Pepe, J. Clark, S. Cozzolino, A.
Cristaudo, S. J. Hiscock, and J. R. Bridle. 2021. Adaptive maternal effects in early life history traits
help maintain ecological resilience in novel environments for two contrasting *Senecio* species,
bioRxiv.