1 # Temporal selectivity declines in the aging human auditory cortex

2 Julia Erb[1], Lea-Maria Schmitt[1], Jonas Obleser[1]

3 [1] Department of Psychology, University of Lübeck, Maria-Goeppert-Str. 9A, 23562 Lübeck, Germany

4 *Correspondence: Julia Erb, Maria-Goeppert-Str. 9a, 23562 Lübeck, Germany

5   Phone: +49 451 3101 3631, e-mail: julia.erb@uni-luebeck.de

Abstract

Current models successfully describe the auditory cortical response to natural sounds with a set of

spectro-temporal features. However, these models have hardly been linked to the ill-understood

neurobiological changes that occur in the aging auditory cortex. Modelling the hemodynamic

response to a rich natural sound mixture in N=64 listeners of varying age, we here show that in older

listeners' auditory cortex, the key feature of temporal rate is represented with a markedly broader

tuning. This loss of temporal selectivity is most prominent in primary auditory cortex and planum

temporale, with no such changes in adjacent auditory or other brain areas. Amongst older listeners,

we observe a direct relationship between chronological age and temporal-rate tuning, unconfounded

by auditory acuity or model goodness of fit. In line with senescent neural dedifferentiation more

generally, our results highlight decreased selectivity to temporal information as a hallmark of the

aging auditory cortex.

Keywords: fMRI, spectro-temporal modulations, healthy aging, auditory system, presbycusis,

hearing loss, temporal rate coding

Impact Statement

Age-dependent broadening of tuning to temporal modulation rate in older humans' cortical sound

representations highlights decreased temporal selectivity as an anatomically and functionally specific

hallmark of the aging auditory cortex.

Introduction

Age-related hearing loss is a frequent cause for a decline in speech comprehension, particularly in complex acoustic environments. As the temporal envelope of speech is crucial for speech comprehension (Shannon et al., 1995), the ability to accurately encode temporal features in the auditory system is pivotal for successful perception of speech and conspecific vocalizations (Peelle and Wingfield, 2016). Age-related deficits have been observed in different temporal psychoacoustic tasks (e.g. gap detection; Snell et al., 2002). Those psychoacoustic studies suggest that the precision of processing of temporal cues declines with age (e.g. Gordon-Salant et al., 2006).

While age-related hearing loss has been attributed to a variety of peripheral declines in auditory coding (e.g. Gates and Mills, 2005), fewer studies have looked at more central levels of auditory processing. Yet, declines in temporal processing are hypothesized to be not exclusively peripheral in origin, but more central than at least the mid-brain (Recanzone, 2018). In the aged macaque auditory cortex, neurons become more broadly tuned to temporal modulations and temporal fidelity of cortical responses decreases (Ng and Recanzone, 2018). Topographically, a neural dedifferentiation is suggested to take place in the older brain: In young macaques, the neuronal response to the inter-stimulus interval in primary field A1 and caudolateral belt area (CL) differs, in that A1 neurons have shorter response latencies. In aged animals, however, no difference was observed between A1 and CL such that aged A1 neurons had an equivalent response latency to CL neurons. Further, in aged neurons, a shift in neural coding strategy from less temporal coding towards more rate coding of temporal modulations was observed (Overton and Recanzone, 2016; for a review of both coding strategies see Joris et al., 2004). For those neurons that still adhered to a temporal coding strategy, a decrease in temporal fidelity was observed, although the absolute number of neurons responsive to temporal modulations was unaffected by age.

At higher processing levels, only a number of relatively unspecific changes have been reported in

the older human listener (for review see Peelle and Wingfield, 2016). For example, the frontal and

cingulo-opercular cortical response is elevated during a challenging speech comprehension or gap

detection task (Erb and Obleser, 2013, Vaden et al., 2015, Vaden et al., 2019). This has been interpreted

as compensatory mechanism in response to loss of sensory acuity. Aging has been hypothesized to

lead to a neural dedifferentiation in sensory cortices (e.g., Park et al., 2004). A recent more unspecific

observation is that with age, decoding accuracy of stimulus conditions from auditory cortical fMRI

responses declines (Lalwani et al., 2019; but see discussion for contrasting findings in MEG speech

tracking, e.g., Presacco et al., 2016). This finding in turn has been linked to an imbalance of excitation

and inhibition in older adults, more specifically a reduction of GABA levels in auditory cortex (Lalwani

et al., 2019). Accumulating evidence supports the theory of an age-related loss of inhibition in sensory

cortices (Caspary et al., 2008).

A recent model of cortical processing (Chi et al., 2005) has led to substantial progress in our

understanding of how natural sounds become represented in the auditory cortex. A series of studies

have shown that cortical processing of sounds is optimized to represent the spectro-temporal

modulations which are typically present in conspecific vocalizations such as speech or animal calls

(Santoro et al., 2014, Hullett et al., 2016, Santoro et al., 2017, Erb et al., 2019). However, those studies

have not looked at the pressing case of the aged human cortex. It remains unknown whether — and if

so, which — general response properties are altered in aged auditory cortex.

The present study applies an fMRI encoding and decoding approach to compare auditory cortical

responses to natural sounds in young and older humans. We focus on the age-group comparison at

the level of representation of fundamental acoustic features (spectro-temporal modulations). Results

show that while the large-scale topographic organization of acoustic features is preserved in the older

auditory cortex, age groups differ in tuning to temporal modulations: Human aging thus appears to

be accompanied by an anatomically and functionally specific broadening of temporal-rate tuning in

auditory cortex.

Results

fMRI responses to sounds from a total of N=64 younger and older listeners were modelled using

both model-based encoding (Santoro et al., 2014) and decoding (Santoro et al., 2017; Figure 1). In

brief, sounds were decomposed into acoustic features, that is, the frequency-dependent spectro-

temporal modulation content (Chi et al., 2005). Analyses were restricted to voxels in an anatomic mask

of the auditory cortex. In a fourfold cross-validation procedure, we used ridge regression to derive

single- and multi-voxel MTFs.

Figure 1. Experimental design and analysis pipeline. Participants listened to a story embedded in sound

textures at an SNR of 0 dB while we acquired 3T-fMRI data. For audiograms of the participants see Figure 1 –

figure supplement 1). The sounds were decomposed into their modulation spectrum (Chi et al., 2005; for

modulation spectra of the sounds see Figure 1 – figure supplement 2). In a first *univariate encoding* approach, we

calculated a modulation transfer function (MTF) for each individual voxel. In a second *multivariate decoding*

approach, linear decoders were trained on the response patterns in auditory cortex for each feature of the

modulation representation. Predictions were then tested on a left-out testing data set.

Encoding results

We compared three models of hemodynamic responses to sound. The models respectively

describe the fMRI responses to the (1) speech stream (foreground), (2) textures (background), or (3) the

mixture of both streams (for modulation spectra of the different streams see Figure 1 – figure

supplement 2).

Sound identification accuracies. In a first encoding analysis, we estimated a modulation transfer

function (MTF) for each voxel based on a subset of fMRI data (training) for each model. We then

assessed the ability of these models to accurately predict the fMRI responses to sounds of a new,

independent data set (testing data). We quantified prediction accuracy by means of a sound

identification analysis (see Materials and Methods).

Sound identification accuracy was higher than chance level (0.5) for all models and age groups

(Figure 2). The sound mixture model had the highest sound identification accuracies (Figure 2a) and

thus described most accurately the fMRI responses to sounds. All subsequent encoding and decoding

analyses were based on this mixture model. Young participants had significantly higher identification

accuracies for mixture than old participants (Figure 2a). There was no significant correlation between

chronological age and sound identification accuracy for mixture within the young or older group

(Figure 2b).

Figure 2. Sound identification accuracies for the different encoding models. (a) Accuracies for sound

mixture, texture (background stream) and speech (foreground stream), for young (left, green) and older

participants (right, violet). Bars indicate the mean (± standard error of the mean, SEM) of all participants per age

group. Accuracies are normalized between 0 and 1; 0 denotes that the predicted activity pattern for a given

stimulus was least similar to the measured one among all test stimuli; 1 denotes correct identification; chance

level is 0.5. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.005$, exact permutation test. (b) Identification accuracies for sound

mixture were uncorrelated with chronological age within each age group.

Topographical best-feature maps. Best features maps were obtained from univariate encoding

models by marginalizing the single-voxel MTFs for the dimension of interest. In particular, tonotopic

maps were obtained by averaging over the rate and scale dimension and assigning the frequency with

the maximal response to a given voxel. Tonotopic maps showed the typical mirror-symmetric

frequency gradients along Heschl's gyrus in both young and aged auditory cortex, irrespective of age

(Figure 3). Tonotopic gradients were most obvious in the young group (Figure 3a, upper panel).

Consistent with previous human fMRI data (Moerel et al., 2012), a low frequency region was observed

in the central region of Heschl's gyrus (HG), presumably marking the boundary between the human

homologue of primary auditory fields A1 and R (Figure 3a, red-yellow). This low frequency region was

surrounded antero-medially and posteriorly by high frequency regions (Figure 3a, green-blue). The

antero-medial high frequency areas clustered on the planum polare (PP). The posterior regions

preferring high frequencies covered planum temporale (PT). We did not find significant age-group

differences using two-sample voxel-wise t-tests (corrected for multiple comparisons). In fact,

tonotopic group maps were correlated, indicating that topography for best frequency was preserved

in older adults (Figure 3a).

Figure 3. Tonotopic maps were derived by marginalizing modulation transfer functions (MTFs) for
frequency and assigning the feature with the maximal response to a given voxel. (a) Group maps were obtained
as median across subjects and are shown on FreeSurfer's *fsaverage5* template. Tonotopic maps for young group
(upper panel) and older group (bottom panel) are correlated in both the right and left hemisphere. (b)
Exemplary individual tonotopic maps for left hemisphere (LH) of a young (upper panel) and old participant
(bottom panel) are displayed on individual surfaces. Black outlines indicate Heschl's gyrus. *** $p < 0.005$.

Topographical best-feature maps for rate and scale were more complex. Similarly, we did not find

significant voxel-wise age-group differences using two-sample *t*-tests (corrected for multiple

comparisons). On the contrary, both for temporal and spectral modulations, group maps were

correlated between age groups (Figure 4). For best temporal modulation maps, we observed a medial-

to-lateral gradient for increasing temporal rate (Figure 4a). For best spectral modulation maps, the

locus of highest spectral resolution coincided with the low frequency region on Heschl's gyrus (Figure

3a, Figure 4b).

Figure 4. Group best-feature maps for temporal rate (a) and spectral scale (b). Group maps were obtained as median across subjects and shown on FreeSurfer's *fsaverage5* template. Black outlines indicate Heschl's gyrus. Topographic organization in young (upper panel) and older group (bottom panel) is significantly correlated (Pearson's correlation coefficient) for best temporal rate (a) and best spectral scale (b). Histograms of best feature values are shown for the young group (green) and older group (violet) above the color bars. Note the shift of the distribution in older relative to young adults for temporal rate (a) but not spectral scale (b), indicating that older adults' voxel-wise preferred temporal rates are shifted towards higher modulation rates. *** $p < 0.005$.

Decoding results

Feature reconstruction from auditory cortex. Cortical sensitivity to acoustic features was quantified using model-based decoding. Decoders were trained on a subset of the data (training) for each acoustic feature separately (Figure 1). Reconstruction accuracies were obtained as Pearson's *r* between predicted and actual acoustic features in the testing data. Results were compared across age groups using exact permutation tests.

Our main hypothesis was that cortical sensitivity is highest for slow temporal modulations, based on previous observations of human (but not monkey) auditory cortex being most sensitive to the modulations present speech (Santoro et al., 2017, Erb et al., 2019a). Decoding yielded highest

accuracies at frequencies of 230–580 Hz (mean r = 0.47) and spectral scales of 0.25 cyc/oct (mean r =

0.47), irrespective of age (Figure 5 a,b), indicating that brain responses followed the frequency and

spectral modulation content of the stimuli (Figure 1 – figure supplement 2 a). Conversely, for temporal

modulations, reconstruction accuracies peaked at rates of 4–8 Hz (mean r = 0.4) in both age groups

(Figure 5 a,b). Those peaks were not present in the stimuli (Figure 1 – figure supplement 2 a) and imply

preferred processing of temporal rates in the speech-relevant range (Poeppel and Assaneo, 2020),

corroborating our hypothesis (see also discussion).

Selectivity index. A key observation were age differences in selectivity of tuning to temporal

modulations (as quantified by the selectivity index SI, see materials and methods). Selectivity of

temporal rate coding was higher in young than in older participants (mean SI age difference = 0.01, p

= 0.004 [permutation test], Cohen's d = 0.68, Figure 5d, left). Note that removing the extreme case in

the young group (outliers were defined as exceeding the grand average by ± 2 SD) for rate SI does not

alter results qualitatively: It reduces the mean age-group difference to a still significant SI difference =

0.009, p = 0.026. Removing the two extreme cases in the older group for scale SI increases the mean

age-group difference to a significant SI difference = 0.037, p = 0.024. A lower selectivity index is

consistent with a broadening of temporal tuning in the aged auditory cortex.

Figure 5. Decoding from auditory cortex. (a) Auditory cortical modulation transfer functions (MTFs) for the young and older group quantified by reconstruction accuracies (Pearson's *r* between predicted an actual acoustic features). (b) Mean ± SEM of the MTFs' marginal profiles for rate, scale and frequency for young (green) and older (violet) participants separately. (c) We quantified tuning selectivity by the selectivity index (see Eq. [7]). For illustration purposes, we show the marginal profiles for rate in a young (green) and older individual (violet) with extreme *SI* values. (d) *SI* was compared per acoustic dimension between age groups using an exact permutation test. Black line indicates the median *SI*. Black line indicates the median *SI*. Note that when removing the extreme case in the young group (> 2 SD away from the mean) for rate *SI*, the mean age-group difference remains significant (*p* = 0.026). Removing the two extreme cases in the older group for scale *SI* leads to a significant age-group difference (*p* = 0.024). Note the different scaling of the plots. See Figure 5 – figure supplement 1 for decoding from auditory cortex with ICA (AROMA)-cleaned data.

Careful artefact removal from fMRI data is particularly critical for age comparisons. Therefore, we re-ran the decoding analysis on ICA-cleaned data (using AROMA, Pruim et al., 2015). Irrespective of preprocessing without (Figure 5) or with ICA-cleaning (Figure 5–figure supplement 1), the key results remained unchanged, that is reconstruction accuracies peaking at temporal rates of 4 – 8 Hz as well as selectivity for temporal rates being higher in young than in older participants.

Also, this key result of a broadened tuning to temporal rate was corroborated by an even simpler

measure of selectivity: The variance in the reconstruction accuracy profiles across the six temporal rate

bins (1–32 Hz, 'rate variance') was higher in the young than in the older participants (mean difference

in variance = 3.7 x 10^{-4}, $p < 0.001$ [permutation test], Cohen's $d = 0.93$, Figure 6, a, left panel).

In the older group, the rate variance was significantly correlated with the rate selectivity index and

chronological age but was uncorrelated with hearing loss (Figure 6, b). Note however that the rate

selectivity index (see above) in the older group was not significantly correlated with chronological age

(Spearman's rho = -0.2, $p = 0.289$), nor with hearing loss (Spearman's rho = 0.34, $p = 0.062$).

In a control analysis, we ensured that these results cannot be explained by the overall slightly

worse model fits in the older group as quantified by sound identification accuracy (as evident in

Figure 2): In a multiple regression predicting older participants' variance in rate from chronological

age with covariates PTA and sound identification accuracy, neither PTA ($t(26) = 0.73$, $p = 0.473$,

permutation test) nor sound identification accuracy ($t(26) = 1.02$, $p = 0.312$) were significant, while age

was significant ($t(26) = -2.8$, $p = 0.003$; *adjusted $R^2 = 0.39$*). Age remained significant when excluding

two multivariate outliers identified based on Cook's distance ($t(24) = -2.25$, $p = 0.021$).

Figure 6. (a) Variance of the MTFs' marginal profiles compared between age groups. *P*-values are based on an exact permutation test. (b) Amongst the older group, the variance of the rate profiles ('rate variance') correlates with the *SI* for rate (left panel) and chronological age (middle panel), but not with hearing loss (right panel; Spearman's correlation coefficient). *SI*: selectivity index, PTA: pure tone average.

Feature reconstruction from auditory subregions. To test the differential sensitivity of auditory cortical subregions to acoustic features, we ran the decoding analyses in the following regions of interest (ROIs) derived from the FreeSurfer atlas (Destrieux et al., 2010): Heschl's gyrus and sulcus (HG/HS), planum polare (PP), planum temporale (PT), superior temporal gyrus (STG) and superior temporal sulcus (STS, Figure 7 a – f)). As control regions, we chose primary visual cortex (V1, calcarine sulcus) as another primary sensory area, and two higher-level areas, namely the middle frontal (MFG) and superior parietal gyrus (SPG; Figure 7 a, g – i).

A decrease in the rate selectivity index with age was confirmed for HG/HS (mean *SI* difference = 0.012, *p* = 0.009 [permutation test]) and PT (mean *SI* difference = 0.016, *p* = 0.005 [permutation test]; see Figure 7 b, c). Removing the extreme cases in the young group exceeding the grand average by ±

2 SD reduces the mean *SI* age difference to 0.012 (p = 0.008) in HG/HS, and to 0.012 (p = 0.04) in PT.

The alternative measure of selectivity, variance of the rate profiles, was also significantly higher in

young than older adults in HG/HS (variance difference = 4 x 10^{-4}, p = 0.0004) and PT (variance

difference = 5 x 10^{-4}, p = 0.0003).

Note that reconstruction accuracies in all ROIs but the control regions (V1, SPG, MFG) were

significant (permutation tests; for the z-scored reconstruction accuracies in an exemplary region

HG/HS see Figure 7–figure supplement 1). In contrast, there were no such age differences in temporal

selectivity in PP, STG, STS or the control regions V1, SPG and MFG (Figure 7 d – i).

Qualitatively, this observation was confirmed by a control analysis, where we derived topographic

maps of the rate selectivity index from the encoding models. To this end, we marginalized MTFs for

rate and calculated the *SI* for a given voxel. The topographic distribution confirmed highest voxel-wise

SI for rate in area PT of the younger group (Figure 7–figure supplement 2).

These results indicate that age-related changes in temporal selectivity selectively occur in primary

auditory cortex and the planum temporale, but not surrounding belt areas. For spectral scale, we also

found a decreased *SI* in older adults in HG/HS (*SI* difference = 0.012, p = 0.008) and PT (*SI* difference =

0.016, p = 0.021), but none of the other ROIs (Figure 7–figure supplement 3). For frequency, we did not

find any age differences in selectivity index at all (Figure 7–figure supplement 3).

Figure 7. Decoding of temporal rate from regions of interest (ROIs). (a) ROIs derived from the FreeSurfer atlas (Destrieux et al., 2010) are displayed on FreeSurfer's *fsaverage* template. Mean (± SEM) reconstruction accuracy profiles (left) and selectivity index (right) per age group for temporal rate in (b) Heschl's gyrus and sulcus, (c) planum temporale, (d) planum polare, (e) superior temporal gyrus, (f) superior temporal sulcus and the control regions (g) calcarine sulcus, (h) superior parietal gyrus, (i) middle frontal gyrus. The selectivity index (*SI*) was compared per ROI between age groups using an exact permutation test. When removing young-subject *SI* exceeding the grand average by ± 2 SD, the *SI* age difference remains significant in HG/HS (p = 0.008), and in PT (p = 0.04). Consistently, our alternative measure of selectivity, variance across rate bins, was significantly higher in young than older adults in HG/HS (p = 0.0004) and PT (p = 0.0003). See Figure 7 – figure supplement 1 for the reconstruction accuracies z-scored with respect to the empirical null distribution in an exemplary region HG/HS. See Figure 7–figure supplement 2 for group maps of the rate selectivity index obtained from encoding models. See Figure 7–figure supplement 3 for decoding of spectral scale and frequency from ROIs.

Discussion

The cross-sectional approach used encoding and decoding models of hemodynamic responses, to

compare the spectro-temporal fidelity in auditory cortex of younger and older listeners. We pursued

the long-standing but largely understudied question whether temporal and/or spectral modulations,

which are key acoustic dimensions of behaviorally relevant signals such as speech and music, are

represented in a less differentiated fashion in the aging auditory cortex.

We find the large-scale topographic organization of acoustic features entirely preserved in the

aged auditory cortex, and best feature maps for frequency, scale and rate accordingly were correlated

very highly between age groups.

However, age-related differences are evident in the cortical sensitivity to mainly temporal–

modulation features as quantified by the decoding analysis. Decoding of temporal rate is most

accurate at slow rates of 4–8 Hz irrespective of age, but the tuning to temporal rate is significantly

sharper in young than in older participants. Also, amongst older adults, broadening of rate tuning

correlates with chronological age. Importantly, the age-related rate selectivity difference was driven

by primary auditory (HG) and adjacent area PT, while secondary auditory areas PP, STG, and STS did

not show age differences.

Preserved topographic organization in aging auditory cortex

Using natural stimuli, we revealed topographic maps of acoustic feature preference for frequency,

spectral and temporal modulations. Thus, we show that topographic maps of feature preference can

be observed under approximately natural hearing situations. Tonotopic maps exhibited the well-

established mirror-symmetric high-low-high frequency gradients across Heschl's gyrus and adjacent

areas (Figure 3) consistent with abundant evidence in human (Formisano et al., 2003, Talavage et al.,

2004, Humphries et al., 2010, Woods et al., 2010, Da Costa et al., 2011, Striem-Amit et al., 2011, Langers

and van Dijk, 2012, Moerel et al., 2012, 2014) and non-human primates (Merzenich and Brugge, 1973,

Morel et al., 1993, Kosaki et al., 1997, Rauschecker et al., 1997, Bendor and Wang, 2008, Baumann et al.,

2013, Joly et al., 2014, Baumann et al., 2015). Notably, tonotopic maps in younger versus older adults

were highly correlated, indicating topographical stability of tonotopic organization across age groups.

The well-demonstrated existence of tonotopic maps notwithstanding, a topographic organization

for temporal or spectral modulations in auditory cortex is less clear (Schonwiesner and Zatorre, 2009).

Here, in both young and older adults we observed a medial-to-lateral low-to-high gradient for

temporal modulations across the superior temporal plane (Figure 4a), while for spectral modulations,

the locus with the highest resolution (~1.5 cyc/oct) was observed in lateral Heschl's gyrus (Figure 4b).

A topographic representation for modulation rate has previously been observed in the primate

auditory cortex (Baumann et al., 2015) where fast temporal acoustic information was preferably

encoded in caudal auditory regions (Camalier et al., 2012, Kusmierek and Rauschecker, 2014) and slow

temporal information in rostral areas R and RT .

Whereas earlier human fMRI studies had failed to show a clear topography for modulation rate

(Giraud et al., 2000, Schonwiesner and Zatorre, 2009, Overath et al., 2012, Leaver and Rauschecker,

2016), recent human fMRI (Santoro et al., 2014) and electrocorticography (ECoG) studies (Hullett et al.,

2016) proposed the presence of a posterior-to-anterior high-to-low rate gradient in human auditory

cortex. However, comparison of those studies is hampered by the differences in coverage and ranges

of temporal modulations: While Hullett et al. (2016) examined representation of slow temporal

modulation rates present in speech (1–2 Hz) only along the STG with ECoG, Santoro at al. (2014)

analyzed rates of 1–27 Hz, whereas Baumann et al. (2015) presented rates of up to 512 Hz with more

comprehensive coverage using fMRI.

Functional mapping from natural sounds is feasible using conventional MR field strength in

aging populations

While previous studies had presented single natural sounds (Erb et al., 2019a), or even synthetic sounds such as AM stimuli (Baumann et al., 2015) or dynamic ripples (Schonwiesner and Zatorre, 2009), here, we presented a continuous stream of speech embedded in an acoustically rich background of sound textures. As an important foundation for the main conclusions of this manuscript, we here demonstrate the capacity to derive meaningful tonotopic maps from natural stimulus conditions, at conventional field strengths (3 T) and in special populations (older participants).

Although an inherent caveat of the use of natural stimulus conditions is the difficulty of measuring psychoacoustic performance, this experimental design has the advantage that it closely approximates natural listening conditions (Hamilton and Huth, 2018). As synthetic sounds lack both the behavioral relevance and the statistical structure of natural sounds, they activate auditory cortex differently than natural stimuli (Theunissen et al., 2000, Bitterman et al., 2008).

Methodologically, our encoding approach relies on the assumption that acoustic features that maximally activate single voxels are more accurately encoded. However, higher responses may not necessarily mean better encoding. The second, complementary decoding analysis incorporates the whole range of values rather than only the maximal values.. The reconstruction accuracy explicitly quantifies the amount of information about a set of stimulus features that is available in a region of interest. Accurate reconstruction of acoustic features in test sounds indicates that these features are reproducibly mapped into distinct spatial patterns. As data from individual voxels are jointly modelled, this combined analysis of signals from multiple voxels increases the sensitivity for stimulus information that may be represented in patterns of responses, rather than in individual voxels.

Broadened temporal tuning in aging auditory cortex

Under the assumption that the reconstruction accuracy reflects tuning properties of neuronal populations, the observed decrease in selectivity reveals a broadening of temporal tuning functions with age. This finding provides mechanistic insights into the changes of temporal processing in the aging auditory cortex: The broadened temporal tuning, reminiscent of an age-related decrease in temporal precision in the midbrain (Anderson et al., 2012), may underlie the difficulties older adults experience in speech comprehension in noisy environments (Anderson et al., 2011). While the present study lacks more direct measures of encoding along the auditory pathway, it is important to note that rate selectivity in the older listeners was statistically unrelated to their pure-tone audiograms (this holds true for both presented measures of tuning, the selectivity index and the variance across temporal rates [Figure 6]). Thus, the broadening of temporal tuning is not explicable merely by peripheral hearing loss. Future studies need to carefully assess the consequences of altered cortical temporal tuning on behavioral and psychoacoustic performance (Erb et al., 2019b, Flinker et al., 2019, Holmes and Griffiths, 2019), a measure that was precluded by the use of natural sounds in the current study.

The observation of decreased temporal selectivity may seem at odds with previous accounts of elevated cortical responses to the temporal speech envelope in aging. Evidence for better reconstruction of temporal envelopes from older listeners' MEG signals (Presacco et al., 2016) and larger N1 amplitudes in older adults with widened auditory filters (Herrmann et al., 2016, Henry et al., 2017) suggest an enhanced cortical envelope representation. However, an inflated cortical representation of the speech envelope has been argued to be rather a maladaptive response as compensation of the lack of temporal precision in the midbrain (Anderson et al., 2012).

Neuronal coding of temporal modulations follows two distinct principles, that is rate and temporal coding (Joris et al., 2004). While rate coding constitutes a variation of overall spike rate as a function of

modulation frequency, temporal coding is achieved through phase-locking to the stimulus envelope.

Fast temporal modulations (> 50 Hz) are typically encoded through the rate code, while neurons

tuned to slow temporal rates (< 50 Hz) mostly synchronize to the sounds' modulations (temporal

code, Sachs, 1984, Joris et al., 2004). It remains unclear how hemodynamic responses reflect these two

types of neuronal coding and how those are in turn related to the observed age-related broadening of

tuning functions. Evidence from electrophysiology suggests a change in coding strategy across

neuronal populations in A1 from less temporal coding towards more rate coding of temporal

modulations in aged animals (Ng and Recanzone, 2018).

Interestingly, the observed age-related decrease in selectivity to temporal modulations was

restricted to areas HG/HS and PT, the putative homologues of areas A1 and CL in the macaque (Kaas

and Hackett, 2000). Both areas have been shown to exhibit altered temporal encoding in older

macaques (Ng and Recanzone, 2018). In a direct comparison of the vector strengths (a metric of

periodicity of the neuronal response to a modulated signal) as a function of the inter-stimulus interval,

responses differed in young A1 and CL neurons such that A1 neurons had higher vector strength. This

is consistent with the notion that CL neurons process predominantly spatial (rather than temporal)

information. Therefore, so the theory, CL neurons would not necessarily need to pass on the temporal

coding of A1 neurons. This difference between belt and core areas was abandoned in the older

animals, where aged A1 were similar to both the aged and young CL neurons, indicating that the

temporal fidelity of A1 responses decreases with age.

However, a good temporal fidelity of A1 neurons is thought to be critical for temporal processing.

We speculate that such an age-related decline in temporal processing may be linked to an imbalance

of excitation and inhibition observed in aging (Voytek et al., 2015). Inhibitory (e.g. GABAergic or

glycinergic) neurotransmitters have been shown to increase response synchrony to modulated stimuli

in both the cochlear nucleus and the inferior colliculus (Koch and Grothe, 1998, Backoff et al., 1999),

for review see (Caspary et al., 2008), suggesting that inhibitory neurotransmission is of particular

importance to precise neural timing and, thus, the adequate tracking of temporal sound features.

An inherent challenge to age group comparisons is to distinguish any unspecific decrease of SNR in

the data of older participants (e.g., due to vascular changes that impact the BOLD response (Garrett et

al., 2017) or due to the typical increase in movement artefacts) from specific age-related sensory

processing changes. Most reassuringly, however, a multiple regression model taking into account

overall goodness of the decoding model fit (i.e., sound identification accuracy) did not alter the

observed relationship between age and temporal rate tuning.

Conclusions

The present study aimed to narrow the gap between recent progress in modelling the neural

processing of complex features from natural stimuli on the one hand, and potential senescent change

in these stimulus representations in the aging auditory cortex on the other hand. Although the large-

scale topographic organization of acoustic features appears preserved in the auditory cortex of older

compared to younger listeners, age-related differences in the marginal profiles of multi-voxel MTFs

were evident. Tuning to slow temporal rates which abounds in natural sounds and especially in

speech (Erb et al., 2019) was markedly sharper in young compared to older participants. Consistent

with previous findings in the macaque, these results suggest that temporal rate selectivity in auditory

cortex declines in normal aging. The specificity of this decline, confined to primary auditory and

adjacent areas and sparing cochleotopic representations, opens a new lead in the ongoing hunt for

tractable neurobiological signatures of older adults' widely observed deficits in speech

comprehension in noisy environments.

Materials and Methods

Participants

We invited a total of $n = 75$ participants for scanning from which we had to exclude one participant due to excessive movement, three participants due to incidental neurological findings, five participants due to their inability to understand speech in noise, one participant due to broken headphones, one participant due to age (43 years, i.e., could not be assigned to young or older group). The remaining $n = 64$ participants were right-handed, young ($n = 33$; aged 18–32, mean 24.7 years18 female) and older ($n = 31$; aged 51–78, mean 63.8 years, 15 female) native German speakers. Simulations (custom matlab code) showed that a two-sample permutation test at a conventional type I error rate of 5 % with a sample size of $N = 27$ per group can detect medium to large effects of Cohen's $d = 0.75$ (as can be expected for the neural fMRI measures under consideration, e.g. Alavash et al., 2019)with a satisfactory power of 80 %. The power of our procedure reduces accordingly if the true effect in the population is smaller but remains over 60 % also for true effects closer to Cohen's $d = 0.5$.

Younger participants had self-reported normal hearing. Older participants' hearing ranged from normal hearing to mild hearing loss. The older participants were part of a cohort which is regularly tested in the lab (e.g., Alavash et al., 2019). They were recruited based on their audiograms that had been acquired within the two years prior to the experiment (for audiograms see Figure 1 – figure supplement 1 a) and were excluded from the study if the pure-tone-average (PTA) of the better ear exceeded 30 dB HL. On average, hearing thresholds are expected to increase by approximately 0.25–1.7 dB HL per year at the age range of 48–79 years (Wiley et al., 2008). Thus, we considered 2-year-old audiograms to be valid. All participants gave informed consent and were financially compensated or received course credit. All procedures were approved by the local ethics committee of the University of Lübeck (ethical approval AZ 16-107).

Stimuli and task

Participants listened to 64 minutes of a freely narrated audiobook (Hertha Müller, "*Die Nacht ist aus Tinte gemacht*") presented against a competing stream of resynthesized natural sounds ("sound textures"; McDermott and Simoncelli, 2011) at 0 dB SNR. Textures were synthesized from the spectro-temporal modulation content of a large set of real-life sounds (n = 192), including speech and vocal samples, music pieces, animal cries, scenes from nature and tool sounds that had been used in previous studies (Moerel et al., 2013, Santoro et al., 2017). Texture synthesis parameters were as follows: Frequency range = 0.02 – 10 kHz, number of frequency bands = 30, sampling rate = 20 kHz, temporal modulation range = 0.5 – 200 Hz, sampling rate = 400 Hz; maximum number of iterations = 60. Texture exemplars of 5 s length were concatenated to form the background stream. The order of exemplars was pseudo-randomized across participants (in four different sound orders). In total, each exemplar was repeated four times.

Participants were asked to listen to the story and answer three four-choice questions on its semantic content after each run (24 questions in total); chance level was thus proportion correct = 0.25. Four-answer choice questions were displayed on a screen and participants responded via a button box in their right hand. Both the young and older group performed on average above chance (see Figure 1 – figure supplement 1b), but young participants performed significantly better than older participants (mean proportion correct difference = 0.1, p = 0.009 (permutation test), Cohen's d = 0.65).

Acquisition of MRI data

We acquired functional and structural MRI at 3T (Siemens Magnetom Skyra) with a 64-channel RF head array coil.

Functional T2*-weighted data were acquired using an echo planar imaging sequence. We collected

eight runs of eight minutes each (8 runs à 519 volumes) using continuous scanning. The acquisition

parameters were as follows: repetition time (TR) = acquisition time (TA) = 947 ms, echo time (TE) = 28

ms, acceleration factor = 4, flip angle = 60°, field of view (FOV) = 200 x 200 mm, 52 slices; voxel size =

2.5 mm isotropic (whole-brain coverage). Field maps for intensity inhomogeneity correction were

acquired after every second run (TR = 610 ms, TE1 = 4.92, TE2 = 7.38, flip angle = 60°, FOV = 200 x 200

mm, 62 slices, voxel size = 2.5 mm isotropic).

Anatomical T1-weighted images were acquired using an MPRAGE sequence (TR = 2400 ms, time to

inversion [TI] = 1000 ms, TE = 3.16 ms, flip angle = 8°, FOV = 256 x 256 mm, number of slices = 176,

voxel size = 1 mm isotropic, GRAPPA acceleration factor = 2). T2-weighted images were collected at

the end of the session (TR = 3200 ms, TE = 449 ms, FOV = 256 x 256 mm, number of slices = 176, voxel

size = 1 mm isotropic, GRAPPA acceleration factor = 2).

Preprocessing of MR data

Results included in this manuscript are based on a preprocessing pipeline of *fMRIPprep* 1.2.4 which

is based on *Nipype* 1.1.6 (Esteban et al., 2019). The following two paragraphs on preprocessing are

based on an automatically generated output of *fMRIPprep*.

Anatomical data preprocessing

The T1-weighted (T1w) image was corrected for intensity non-uniformity (INU)

using N4BiasFieldCorrection (ANTs 2.2.0) and used as T1w-reference throughout the workflow. The

T1w-reference was then skull-stripped using antsBrainExtraction.sh (ANTs 2.2.0), using OASIS as target

template. Brain surfaces were reconstructed using recon-all (FreeSurfer 6.0.1), and the brain mask

estimated previously was refined with a custom variation of the method to reconcile ANTs-derived

and FreeSurfer-derived segmentations of the cortical gray-matter of Mindboggle. Spatial

normalization to the ICBM 152 Nonlinear Asymmetrical template version 2009c was performed

through nonlinear registration with antsRegistration (ANTs 2.2.0), using brain-extracted versions of

both T1w volume and template. Brain tissue segmentation of cerebrospinal fluid (CSF), white-matter

(WM) and gray-matter (GM) was performed on the brain-extracted T1w using fast (FSL 5.0.9).

Functional data preprocessing

For each of the eight BOLD runs, the following preprocessing was performed. First, a reference

volume and its skull-stripped version were generated using a custom methodology of *fMRIPrep*. A

deformation field to correct for susceptibility distortions was estimated based on a field map that was

co-registered to the BOLD reference, using a custom workflow of *fMRIPrep*. Based on the estimated

susceptibility distortion, an unwarped BOLD reference was calculated for a more accurate co-

registration with the anatomical reference. The BOLD reference was then co-registered to the T1w

reference using bbregister (FreeSurfer) which implements boundary-based registration. Co-

registration was configured with nine degrees of freedom to account for distortions remaining in the

BOLD reference. Head-motion parameters with respect to the BOLD reference (transformation

matrices, and six corresponding rotation and translation parameters) are estimated before any

spatiotemporal filtering using mcflirt (FSL 5.0.9). BOLD runs were slice-time corrected

using 3dTshift from AFNI 20160207. The BOLD time-series (including slice-timing correction when

applied) were resampled onto their original, native space by applying a single, composite transform to

correct for head-motion and susceptibility distortions. In a control analysis, ICA-based Automatic

Removal Of Motion Artifacts (AROMA) was used to generate a variant of data that is non-aggressively

denoised (Pruim et al., 2015). All analyses (see below) were performed in native space. Subsequently,

results (best feature maps) were resampled to surfaces on the *fsaverage5* template using mri_vol2surf

(FreeSurfer).

Modulation representation

The modulation content of the stimuli was obtained by filtering the sounds within a biologically

plausible model of auditory processing (Chi et al., 2005). This auditory model consists of an early stage

that models the transformations that acoustic signals undergo from the cochlea to the midbrain; and a

cortical stage that accounts for the processing of the sounds at the level of the auditory cortex. We

derived the spectrogram and its modulation content using the "NSL Tools" package (available at

http://www.isr.umd.edu/Labs/NSL/Software.htm) and customized Matlab code (The MathWorks Inc.,

Matlab 2014b/2018a).

Spectrograms for all sounds were obtained using a bank of 128 overlapping bandpass filters with

equal width (Q_{10dB} = 3), spaced along a logarithmic frequency axis over a range of f = 180–7040 Hz. The

output of the filter bank was band-pass filtered (hair cell stage). A midbrain stage modelled the

enhancement of frequency selectivity as a first-order derivative with respect to the frequency axis,

followed by a half-wave rectification and a short-term temporal integration (time constant τ = 8 ms).

Next, the auditory spectrogram was further analyzed by the cortical stage, where the modulation

content of the auditory spectrogram was computed through a bank of 2-dimensional filters selective

for a combination of spectral and temporal modulations. The filter bank performs a complex wavelet

decomposition of the auditory spectrogram. The magnitude of such decomposition yields a phase-

invariant measure of modulation content. The modulation selective filters have joint selectivity for

spectral and temporal modulations, and are directional, i.e. they respond either to upward or

downward frequency sweeps.

Filters were tuned to spectral modulation frequencies of Ω = [0.3, 0.4, 0.8, 1.3, 2.3, 4] cyc/oct,

temporal modulation frequencies of ω = [1, 2, 4, 8, 16, 32] Hz, and centre frequencies of f = [232, 367,

580, 918, 1452, 2297, 3633, 5746] Hz. Our rationale for this choice of values was to use a

decomposition roughly covering the temporal and spectral modulations present in the acoustic

energy of natural sounds we used (for spectro-temporal modulation content of the sounds see Figure

1 – figure supplement 2). To avoid overfitting, for the decoding analyses in regions of interest (ROIs),

we reduced the number of features such that filters were tuned to spectral modulation frequencies of

Ω = [0.25, 0.5, 1, 2 , 4] cyc/oct, temporal modulation frequencies of ω = [1, 2.4, 5.7, 13.5, 32] Hz, and

centre frequencies of f = [277, 576, 1201, 2502, 5213] Hz.

The filter bank output was computed at each frequency along the tonotopic axis and then

averaged over time. This resulted for the encoding (ROI decoding) analysis in a representation with 6

(5) spectral modulation frequencies $\times$ 6 (5) temporal modulation frequencies $\times$ 8 (5) tonotopic

frequencies = 288 (125) parameters to learn. The time-averaged output of the filter bank was averaged

across the upward and downward filter directions (Santoro et al., 2014). Those processing steps were

applied to all stimuli, resulting in an [N x F] feature matrix S of modulation energy, where N is the

number of sounds, and F is the number of features in the modulation representation.

Prior to acoustic feature extraction, the continuous sounds (~8 min per run) were cut into

snippets of the length of the TR (947 ms) resulting in a total of n = 4152 sounds which were subdivided

into training (n = 3114) and test sounds (n = 1038) for the cross-validation procedure (see below). This

resulted in an equivalent temporal resolution of the feature matrix and the fMRI data.

Note that for the encoding analysis (see below), the number of parameters to estimate is thus

smaller than the number of observations in the training set (n = 3114 training sounds). In the decoding

analysis, the number of fitted features is limited by the number of voxels instead (Equation 5). Each

feature was convolved with the standard double gamma model for the hemodynamic response

function peaking at 4 s.

528 **Encoding and decoding models**

529 We applied two modelling approaches to fMRI data as described in Erb et al. (2019; Figure 1). In a

530 first *univariate encoding* approach, we calculated a modulation transfer function (MTF) for each

531 individual voxel. The MTF characterizes how faithfully the modulation content of the stimulus gets

532 transferred to the voxel. By assigning the feature value with the maximal response to each voxel, we

533 obtained maps of a voxel's best features across the auditory cortex.

534 In a second *multivariate decoding* approach, data from voxels were jointly modelled within a

535 model-based decoding framework. The combined analysis of signals from multiple voxels increases

536 the sensitivity for stimulus information that is represented in patterns of activity, rather than in

537 individual voxels. Further, the accuracy with which those features can be reconstructed provides an

538 explicit measure of the amount of information about sound features available in cortex.

539 **Univariate encoding analysis: Model estimation**

540 Based on the training data only, the fMRI activity Y_i [$N_{train} \times 1$] at voxel i was modeled as a linear

541 transformation of the feature matrix S_{train} [$N_{train} \times F$] plus a noise term n [$N_{train} \times 1$]:

$$Y_i = S_{train}C_i + n \qquad (1)$$

543 where N_{train} is the number of sounds in the training set, and C_i is an [$F \times 1$] vector of model

544 parameters, whose elements c_{ij} quantify the contribution of feature j to the overall response of voxel i.

545 Columns of matrices S_{train} and Y_i were converted to standardized z-scores. Therefore, Equation (1) does

546 not include a constant term. The solution to Equation (1) was computed using kernel ridge regression

547 (Hoerl and Kennard, 1970). The regularization parameter λ was selected independently for each voxel

548 via generalized cross-validation (Golub et al., 1979). The search grid included 25 values between 10^{-6}

549 and 10^6 logarithmically spaced with a grid grain of $10^{0.5}$.

550   To obtain more stable estimates of the voxels' feature profiles, this computation was performed in

551   a fourfold cross-validation procedure within each participant using different subsets of the eight runs.

552   For each iteration, two out of the eight runs were selected and left out for testing, resulting in a subset

553   of 3114 training sounds on which the estimation was performed and 1038 test sounds. In this way, we

554   obtained four estimates of each voxel's feature profile which were averaged across iterations.

Model evaluation

556   To evaluate the model's prediction accuracy we performed a sound identification analysis (Kay et

557   al., 2008). To this end, we used the fMRI activity patterns predicted by the model to identify which of

558   the test sounds had been heard. Given the trained model $\tilde{C}$ [$F \times V$], and the feature matrix S_{test} [$N_{test} \times F$]

559   for the test set, the predicted fMRI activity $\hat{Y}_{test}$ [$N_{test} \times V$] for the test sounds was obtained as:

$$\hat{Y}_{test} = S_{test}\,\tilde{C} \qquad (2)$$

561   Then, we computed for each stimulus s_k the correlation r_s between its predicted fMRI activity

562    $\hat{Y}_{test}(s)$ [$1 \times V$] and all measured fMRI responses $Y_{test}(s)$ [$1 \times V$]. The rank of the correlation between

563   predicted and observed activity for stimulus s_k was used as a measure of the model's ability to

564   correctly match $Y_{test}(s)$ with its prediction $\hat{Y}_{test}(s)$. The rank was then normalized between 0 and 1 as

565   follows to obtain the sound identification score m for stimulus s (Santoro et al., 2014):

$$m_s = 1 - \frac{rank(r_s) - 1}{N_{test} - 1} \qquad (3)$$

567   Note that $m = 1$ indicates a correct match; $m = 0$ indicates that the predicted activity pattern for

568   stimulus s_i was least similar to the measured one among all stimuli. Normalized ranks (sound

569   identification scores) were computed for all stimuli in the test set, and the overall model's accuracy

570   was obtained as the mean of the sound identification scores across stimuli.

Topographical best-feature maps

The response profiles for temporal modulation, spectral modulation and frequency were computed as marginal sums of the estimated stimulus-activity mapping function C of the frequency-specific modulation model by summing across irrelevant dimensions. For example, to obtain the temporal modulation transfer function (tMTF), we summed across the spectral modulation and frequency dimension:

$$tMTF(\omega) = \sum_{\Omega}\sum_{f} C(\omega, \Omega, f) \qquad (4)$$

To calculate profiles for the spectral modulation transfer function (sMTF) and frequency transfer function (fTF), we correspondingly summed across irrelevant dimensions. The voxels' characteristic values (CTM, CSM, CF) were defined as the maximum of the tMTF, sMTF and fTF, respectively. Cortical maps were generated by color-coding the voxels' preferred values and projecting them onto an inflated representation of the cortex. To obtain group maps, individual maps were transformed to FreeSurfer's *fsaverage5* space and averaged.

Multivariate decoding analysis: Model estimation

In the multivariate decoding analysis, we evaluated the fidelity with which regions of interest (ROIs) in auditory cortex encode acoustic features by estimating decoders. In addition to a region including the whole auditory cortex, we selected five ROIs based on anatomical criteria (using the FreeSurfer labels): Heschl's gyrus/sulcus (HG/HS), planum temporale (PT), planum polare (PP), superior temporal gyrus (STG) and sulcus (STS). In a control ROI analysis, we selected visual cortex (V1, calcarine sulcus), middle frontal (MFG) and superior parietal gyrus (SPG). For each individual in each ROI, a linear decoder was trained for every feature of the modulation space based on the training data only

592   (Santoro et al., 2017). Each stimulus feature S_j [$N_{train} \times 1$] was modelled as a linear transformation of the

593   multi-voxel response pattern Y_{train}[$N_{train} \times V$] plus a bias term b_j and a noise term n [$N_{train} \times 1$] as follows:

$$S_j = Y_{train} C_j + b_j 1 + n \qquad (5)$$

where N_{train} is the number of sounds in the training set, V is the number of voxels, 1 is a [$N_{train} \times 1$]

vector of ones, and C_j is a [$V \times 1$] vector of model parameters, whose elements c_{ji} quantify the

contribution of voxel i to the encoding of feature j. All parameters in Equation (5) were estimated

within the same ridge-regression cross-validation scheme as Equation (1). Here, the regularization

parameter λ was determined independently for each feature by generalized cross validation using the

identical search grid as in the univariate encoding model (see above).

Estimation of multi-voxel modulation transfer functions (MTFs)

Decoders were estimated on the train runs (see above) and tested on the test runs. Given the

trained model $\tilde{C}$ [$V \times F$], and the patterns of fMRI activity for the test sounds Y_{test} [$N_{test} \times V$], the predicted

feature matrix activity $\hat{S}_{test}$ [$N_{test} \times F$] for the test sounds was calculated as:

$$\hat{S}_{test} = Y_{test} \, \tilde{C} \qquad (6)$$

Predictions for all features from the test sets were concatenated and decoders were assessed

individually by computing the Pearson's correlation coefficient (r) between a predicted and a given

actual stimulus feature. For decoding from the whole auditory cortex (ROIs), this resulted in 288 (125)

correlation coefficients, which represented the MTF. To obtain marginal profiles of the MTFs, we

averaged across the momentarily irrelevant dimensions, respectively.

Statistical significance of the reconstruction accuracy was assessed with permutation testing. For

each stimulus feature we computed the null-distribution of accuracies at the single-subject level by

randomly permuting (1, 000 times) the stimulus labels of the test sounds and computing the

correlation coefficient with the predicted features for each permutation. Note that to reduce

computing time and resources, we limited the number of permutations to 1,000 whenever encoding

and decoding models had to be estimated for each permutation at the single-subject level. For less

computationally intense calculations (group comparisons), we used the default of 10,000

permutations.

To preserve the temporal correlations among sounds, the predictors were phase scrambled by

using the fast Fourier transform (fft), randomly permuting the phase angles of the fft and calculating

the inverse fft before convolution with the hemodynamic response function. The empirical chance

level of correlation r_{chance} was defined as the mean of the null distribution. The p-value was computed

as the proportion of permutations that yielded a correlation equal to or more extreme than the

empirical one. The reconstruction accuracies were z-scored relative to the empirical null distribution.

Post-hoc statistical analysis of marginal MTFs

Group marginal profiles of MTFs were obtained as the mean of all individual marginal MTFs. To

assess the statistical significance of the observations on the MTF's marginal profiles, we performed the

following post-hoc analyses.

Selectivity Index. Based on the marginal profiles of the MTFs, we quantified how selectively a brain

region is tuned towards processing the acoustic feature with the highest reconstruction accuracy

relative to all other features using a standard index

632
$$SI = \frac{\max\limits_{j\in[k]}(r_j) - \underset{j\neq jmax}{\mathrm{mean}}(r_j)}{\max\limits_{j\in[k]}(r_j) + \underset{j\neq jmax}{\mathrm{mean}}(r_j)} \qquad (7)$$

Where r_j is the reconstruction accuracy at feature j, k is the number of features, j_{max} is the index of

the maximal reconstruction accuracy. SI was calculated for each acoustic dimension separately (i.e.,

rate, scale or frequency) and compared between age groups using an exact permutation test by

calculating the mean group difference and permuting the age group labels 10,000 times.

Acknowledgements

This research was funded by an ERC consolidator Grant (ERC-CoG-2014-646696 "AUDADAPT" to

J.O.) and the German Research Foundation (DFG; OB 352/2-1). Martin Göttlich helped with MR

sequences. Anne Herrmann, Malte Naujokat, Clara Mergner, and Anne Ruhe helped acquire the data.

We thank the members of the Auditory Cognition group as well as the editors and reviewers for very

constructive feedback on the present study.

Competing interests

The authors declare no competing interests.

Data availability

MRI data and custom code to reproduce all essential findings will be publicly available on the Open

Science Framework (OSF) upon publication (https://osf.io/zbuah/).

References

Alavash M, Tune S, Obleser J (2019) Modular reconfiguration of an auditory control brain network
supports adaptive listening behavior. Proceedings of the National Academy of Sciences of the
United States of America 116:660-669.
Anderson S, Parbery-Clark A, White-Schwoch T, Kraus N (2012) Aging affects neural precision of speech
encoding. The Journal of neuroscience : the official journal of the Society for Neuroscience
32:14156-14164.
Anderson S, Parbery-Clark A, Yi HG, Kraus N (2011) A neural basis of speech-in-noise perception in
older adults. Ear and hearing 32:750-757.
Backoff PM, Shadduck Palombi P, Caspary DM (1999) Gamma-aminobutyric acidergic and glycinergic
inputs shape coding of amplitude modulation in the chinchilla cochlear nucleus. Hearing
research 134:77-88.
Baumann S, Joly O, Rees A, Petkov CI, Sun L, Thiele A, Griffiths TD (2015) The topography of frequency
and time representation in primate auditory cortices. eLife 4.
Baumann S, Petkov CI, Griffiths TD (2013) A unified framework for the organization of the primate
auditory cortex. Front Syst Neurosci 7:11.

Bendor D, Wang X (2008) Neural response properties of primary, rostral, and rostrotemporal core fields
in the auditory cortex of marmoset monkeys. Journal of neurophysiology 100:888-906.
Bitterman Y, Mukamel R, Malach R, Fried I, Nelken I (2008) Ultra-fine frequency tuning revealed in
single neurons of human auditory cortex. Nature 451:197-201.
Camalier CR, D'Angelo WR, Sterbing-D'Angelo SJ, de la Mothe LA, Hackett TA (2012) Neural latencies
across auditory cortex of macaque support a dorsal stream supramodal timing advantage in
primates. Proceedings of the National Academy of Sciences of the United States of America
109:18168-18173.
Caspary DM, Ling L, Turner JG, Hughes LF (2008) Inhibitory neurotransmission, plasticity and aging in
the mammalian central auditory system. The Journal of experimental biology 211:1781-1791.
Chi T, Ru P, Shamma SA (2005) Multiresolution spectrotemporal analysis of complex sounds. The
Journal of the Acoustical Society of America 118:887-906.
Da Costa S, van der Zwaag W, Marques JP, Frackowiak RS, Clarke S, Saenz M (2011) Human primary
auditory cortex follows the shape of Heschl's gyrus. The Journal of neuroscience : the official
journal of the Society for Neuroscience 31:14067-14075.
Destrieux C, Fischl B, Dale A, Halgren E (2010) Automatic parcellation of human cortical gyri and sulci
using standard anatomical nomenclature. NeuroImage 53:1-15.
Erb J, Armendariz M, De Martino F, Goebel R, Vanduffel W, Formisano E (2019a) Homology and
Specificity of Natural Sound-Encoding in Human and Monkey Auditory Cortex. Cerebral cortex
29:3636-3650.
Erb J, Ludwig AA, Kunke D, Fuchs M, Obleser J (2019b) Temporal Sensitivity Measured Shortly After
Cochlear Implantation Predicts 6-Month Speech Recognition Outcome. Ear and hearing 40:27-
33.
Erb J, Obleser J (2013) Upregulation of cognitive control networks in older adults' speech
comprehension. Front Syst Neurosci 7:116.
Esteban O, Markiewicz CJ, Blair RW, Moodie CA, Isik AI, Erramuzpe A, Kent JD, Goncalves M, DuPre E,
Snyder M, Oya H, Ghosh SS, Wright J, Durnez J, Poldrack RA, Gorgolewski KJ (2019) fMRIPrep: a
robust preprocessing pipeline for functional MRI. Nature methods 16:111-116.
Flinker A, Doyle WK, Mehta AD, Devinsky O, Poeppel D (2019) Spectrotemporal modulation provides a
unifying framework for auditory cortical asymmetries. Nat Hum Behav 3:393-405.
Formisano E, Kim DS, Di Salle F, van de Moortele PF, Ugurbil K, Goebel R (2003) Mirror-symmetric
tonotopic maps in human primary auditory cortex. Neuron 40:859-869.
Garrett DD, Lindenberger U, Hoge RD, Gauthier CJ (2017) Age differences in brain signal variability are
robust to multiple vascular controls. Scientific reports 7:10149.
Gates GA, Mills JH (2005) Presbycusis. Lancet 366:1111-1120.
Giraud AL, Lorenzi C, Ashburner J, Wable J, Johnsrude I, Frackowiak R, Kleinschmidt A (2000)
Representation of the temporal envelope of sounds in the human brain. Journal of
neurophysiology 84:1588-1598.
Golub G, Heath M, Wahba G (1979) Generalized Cross-Validation as a Method for Choosing a Good
Ridge Parameter. Technometrics 21:215-223.
Gordon-Salant S, Yeni-Komshian GH, Fitzgibbons PJ, Barrett J (2006) Age-related differences in
identification and discrimination of temporal cues in speech segments. The Journal of the
Acoustical Society of America 119:2455-2466.
Hamilton LS, Huth AG (2018) The revolution will not be controlled: natural stimuli in speech
neuroscience. Language, cognition and neuroscience 1-10.
Henry MJ, Herrmann B, Kunke D, Obleser J (2017) Aging affects the balance of neural entrainment and
top-down neural modulation in the listening brain. Nature communications 8:15801.
Herrmann B, Henry MJ, Johnsrude IS, Obleser J (2016) Altered temporal dynamics of neural adaptation
in the aging human auditory cortex. Neurobiology of aging 45:10-22.

Hoerl AE, Kennard RW (1970) Ridge regression: Biased estimation for nonorthogonal problems.
Technometrics 12:55-67.
Holmes E, Griffiths TD (2019) 'Normal' hearing thresholds and fundamental auditory grouping
processes predict difficulties with speech-in-noise perception. Scientific reports 9:16771.
Hullett PW, Hamilton LS, Mesgarani N, Schreiner CE, Chang EF (2016) Human Superior Temporal Gyrus
Organization of Spectrotemporal Modulation Tuning Derived from Speech Stimuli. The
Journal of neuroscience : the official journal of the Society for Neuroscience 36:2014-2026.
Humphries C, Liebenthal E, Binder JR (2010) Tonotopic organization of human auditory cortex.
NeuroImage 50:1202-1211.
Joly O, Baumann S, Balezeau F, Thiele A, Griffiths TD (2014) Merging functional and structural
properties of the monkey auditory cortex. Frontiers in neuroscience 8:198.
Joris PX, Schreiner CE, Rees A (2004) Neural processing of amplitude-modulated sounds. Physiol Rev
84:541-577.
Kaas JH, Hackett TA (2000) Subdivisions of auditory cortex and processing streams in primates.
Proceedings of the National Academy of Sciences of the United States of America 97:11793-
11799.
Kay KN, Naselaris T, Prenger RJ, Gallant JL (2008) Identifying natural images from human brain activity.
Nature 452:352-355.
Koch U, Grothe B (1998) GABAergic and glycinergic inhibition sharpens tuning for frequency
modulations in the inferior colliculus of the big brown bat. Journal of neurophysiology 80:71-
82.
Kosaki H, Hashikawa T, He J, Jones EG (1997) Tonotopic organization of auditory cortical fields
delineated by parvalbumin immunoreactivity in macaque monkeys. The Journal of
comparative neurology 386:304-316.
Kusmierek P, Rauschecker JP (2014) Selectivity for space and time in early areas of the auditory dorsal
stream in the rhesus monkey. Journal of neurophysiology 111:1671-1685.
Lalwani P, Gagnon H, Cassady K, Simmonite M, Peltier S, Seidler RD, Taylor SF, Weissman DH, Polk TA
(2019) Neural distinctiveness declines with age in auditory cortex and is associated with
auditory GABA levels. NeuroImage 201:116033.
Langers DR, van Dijk P (2012) Mapping the tonotopic organization in human auditory cortex with
minimally salient acoustic stimulation. Cerebral cortex 22:2024-2038.
Leaver AM, Rauschecker JP (2016) Functional Topography of Human Auditory Cortex. The Journal of
neuroscience : the official journal of the Society for Neuroscience 36:1416-1428.
Liang L, Lu T, Wang X (2002) Neural representations of sinusoidal amplitude and frequency
modulations in the primary auditory cortex of awake primates. Journal of neurophysiology
87:2237-2261.
McDermott JH, Simoncelli EP (2011) Sound texture perception via statistics of the auditory periphery:
evidence from sound synthesis. Neuron 71:926-940.
Merzenich MM, Brugge JF (1973) Representation of the cochlear partition of the superior temporal
plane of the macaque monkey. Brain research 50:275-296.
Moerel M, De Martino F, Formisano E (2012) Processing of natural sounds in human auditory cortex:
tonotopy, spectral tuning, and relation to voice sensitivity. The Journal of neuroscience : the
official journal of the Society for Neuroscience 32:14205-14216.
Moerel M, De Martino F, Formisano E (2014) An anatomical and functional topography of human
auditory cortical areas. Frontiers in neuroscience 8:225.
Moerel M, De Martino F, Santoro R, Ugurbil K, Goebel R, Yacoub E, Formisano E (2013) Processing of
natural sounds: characterization of multipeak spectral tuning in human auditory cortex. The
Journal of neuroscience : the official journal of the Society for Neuroscience 33:11888-11898.

Morel A, Garraghty PE, Kaas JH (1993) Tonotopic organization, architectonic fields, and connections of
762   auditory cortex in macaque monkeys. The Journal of comparative neurology 335:437-459.
763   Ng CW, Recanzone GH (2018) Age-Related Changes in Temporal Processing of Rapidly-Presented
764   Sound Sequences in the Macaque Auditory Cortex. Cerebral cortex 28:3775-3796.
765   Overath T, Zhang Y, Sanes DH, Poeppel D (2012) Sensitivity to temporal modulation rate and spectral
766   bandwidth in the human auditory system: fMRI evidence. Journal of neurophysiology
767   107:2042-2056.
768   Park DC, Polk TA, Park R, Minear M, Savage A, Smith MR (2004) Aging reduces neural specialization in
769   ventral visual cortex. Proceedings of the National Academy of Sciences of the United States of
770   America 101:13091-13095.
771   Peelle JE, Wingfield A (2016) The Neural Consequences of Age-Related Hearing Loss. Trends Neurosci
772   39:486-497.
773   Poeppel D, Assaneo MF (2020) Speech rhythms and their neural foundations. Nature reviews
774   Neuroscience.
775   Presacco A, Simon JZ, Anderson S (2016) Evidence of degraded representation of speech in noise, in
776   the aging midbrain and cortex. Journal of neurophysiology 116:2346-2355.
777   Pruim RHR, Mennes M, van Rooij D, Llera A, Buitelaar JK, Beckmann CF (2015) ICA-AROMA: A robust
778   ICA-based strategy for removing motion artifacts from fMRI data. NeuroImage 112:267-277.
779   Rauschecker JP, Tian B, Pons T, Mishkin M (1997) Serial and parallel processing in rhesus monkey
780   auditory cortex. The Journal of comparative neurology 382:89-103.
781   Recanzone G (2018) The effects of aging on auditory cortical function. Hearing research 366:99-105.
782   Sachs MB (1984) Neural coding of complex sounds: speech. Annual review of physiology 46:261-273.
783   Santoro R, Moerel M, De Martino F, Goebel R, Ugurbil K, Yacoub E, Formisano E (2014) Encoding of
784   natural sounds at multiple spectral and temporal resolutions in the human auditory cortex.
785   PLoS computational biology 10:e1003412.
786   Santoro R, Moerel M, De Martino F, Valente G, Ugurbil K, Yacoub E, Formisano E (2017) Reconstructing
787   the spectrotemporal modulations of real-life sounds from fMRI response patterns. Proceedings
788   of the National Academy of Sciences of the United States of America 114:4799-4804.
789   Schonwiesner M, Zatorre RJ (2009) Spectro-temporal modulation transfer function of single voxels in
790   the human auditory cortex measured with high-resolution fMRI. Proceedings of the National
791   Academy of Sciences of the United States of America 106:14611-14616.
792   Shannon RV, Zeng FG, Kamath V, Wygonski J, Ekelid M (1995) Speech recognition with primarily
793   temporal cues. Science 270:303-304.
794   Snell KB, Mapes FM, Hickman ED, Frisina DR (2002) Word recognition in competing babble and the
795   effects of age, temporal processing, and absolute sensitivity. The Journal of the Acoustical
796   Society of America 112:720-727.
797   Striem-Amit E, Hertz U, Amedi A (2011) Extensive cochleotopic mapping of human auditory cortical
798   fields obtained with phase-encoding FMRI. PloS one 6:e17832.
799   Talavage TM, Sereno MI, Melcher JR, Ledden PJ, Rosen BR, Dale AM (2004) Tonotopic organization in
800   human auditory cortex revealed by progressions of frequency sensitivity. Journal of
801   neurophysiology 91:1282-1296.
802   Theunissen FE, Sen K, Doupe AJ (2000) Spectral-temporal receptive fields of nonlinear auditory
803   neurons obtained using natural sounds. The Journal of neuroscience : the official journal of the
804   Society for Neuroscience 20:2315-2331.
805   Vaden KI, Jr., Eckert MA, Dubno JR, Harris KC (2019) Cingulo-opercular adaptive control for younger
806   and older adults during a challenging gap detection task. Journal of neuroscience research.
807   Vaden KI, Jr., Kuchinsky SE, Ahlstrom JB, Dubno JR, Eckert MA (2015) Cortical activity predicts which
808   older adults recognize speech in noise and when. The Journal of neuroscience : the official
809   journal of the Society for Neuroscience 35:3929-3937.

810   Voytek B, Kramer MA, Case J, Lepage KQ, Tempesta ZR, Knight RT, Gazzaley A (2015) Age-Related
811   Changes in 1/f Neural Electrophysiological Noise. The Journal of neuroscience : the official
812   journal of the Society for Neuroscience 35:13257-13265.
813   Wiley TL, Chappell R, Carmichael L, Nondahl DM, Cruickshanks KJ (2008) Changes in hearing
814   thresholds over 10 years in older adults. Journal of the American Academy of Audiology
815   19:281-292; quiz 371.
816   Woods DL, Herron TJ, Cate AD, Yund EW, Stecker GC, Rinne T, Kang X (2010) Functional properties of
817   human auditory cortical fields. Front Syst Neurosci 4:155.

Supplementary Information for

Temporal selectivity declines in the aging human auditory cortex

Julia Erb[1], Lea-Maria Schmitt[1], Jonas Obleser[1]

[1]Department of Psychology, University of Lübeck, Maria-Goeppert-Str. 9A, 23562 Lübeck, Germany

*Correspondence: Julia Erb, Maria-Goeppert-Str. 9a, 23562 Lübeck, Germany

Phone: +49 451 3101 3631, e-mail: julia.erb@uni-luebeck.de

Figure 1 – figure supplement 1. (a) **Audiograms** for older participants (averaged over left and right ear). (b)

Average behavioral performance (mean ± SEM) in the scanner in a four-choice task (3 questions on the story

content after each run) for young (green) and older participants (violet). Chance level was 0.25 [proportion

correct]. Young participants performed significantly better than older participants (mean proportion correct

difference =0.1, p = 0.009 [permutation test], Cohen's d = 0.65).

Figure 1 – figure supplement 2. Modulation spectra of the stimuli. (a) Average of the marginalized
modulation spectra for the stimulus streams separately and their mixture. (b) Standard deviation (SD) of the
marginalized modulation spectra. (c) Pairwise correlations (Pearson's *r*) between the different streams in the
modulation representation. While speech and texture are mostly uncorrelated over time, the speech–texture
mixture is highly correlated with the speech stream in the low frequency range and with textures in the high
frequency range. Note that the left plot has a different scaling than the middle and right plot.

Figure 5 – figure supplement 1. Decoding from auditory cortex with ICA(AROMA)-cleaned data. (a)
Mean ± SEM of the MTFs' marginal profiles for rate, scale and frequency for young (green) and older (violet)
participants separately. (b) The selectivity index *SI* (see Eq. [7]) was compared per acoustic dimension between
age groups using an exact permutation test. Note that irrespective of preprocessing (with or without ICA-
cleaning, Fig. 5), reconstruction accuracies peak at temporal rates of 4 – 8 Hz. Likewise, for AROMA-cleaned data
SI for temporal rates is higher in young compared to older participants. Removing the young-subject outlier for
rate *SI* (exceeding the grand average by ± 2 SD) does not alter the *SI* mean age difference qualitatively (*p* =
0.027).

Figure 7 – figure supplement 1. Z-scored reconstruction accuracies from Heschl's gyrus/sulcus. Single
subject and median reconstruction accuracy profiles for temporal rate, spectral scale and frequency z-scored
with respect to the empirical null distribution for young (a) and older (b) participants. Note that all z-values are >
1.96 and are thus significant at $p < 0.05$.

Figure 7 – figure supplement 2. Topographic maps for the rate selectivity index. Maps were derived by
calculating the *SI* across the rate profile for a given voxel and averaged in the young group (upper panel) and
older group (bottom panel). Higher *SI* (red) indicates higher selectivity. Black outlines indicate Heschl's gyrus.

Figure 7 – figure supplement 3. Decoding of spectral scale and frequency in ROIs. Mean (±SE)

reconstruction accuracy profiles (left) and selectivity index (right) per age group for spectral scale and frequency

in (a) Heschl's gyrus and sulcus (HG/HS), (b) planum temporale (PT), (c) planum polare (PP), (d) superior temporal

gyrus (STG), (e) superior temporal sulcus (STS) and the control regions (f) calcarine sulcus (V1), (g) superior

parietal gyrus (SPG) and middle frontal gyrus (MFG). Selectivity index was compared per ROI between age

groups using exact permutation test. Black line indicates the median *SI*. Note that removing the young outlier

participant (> 2 SD away from the mean) for scale in PT reduces the mean age difference to a still significant *SI*

difference = 0.015, $p = 0.026$.