Functional diversity metrics using kernel density *n*-dimensional hypervolumes

Running title: Kernel-based functional diversity metrics

Stefano Mammola[1,2,†,*], Pedro Cardoso[2,†,**]

1) Molecular Ecology Group (MEG), IRSA –Water Research Institute, National Research Council, Verbania

Pallanza, Italy

2) Laboratory for Integrative Biodiversity Research (LIBRe), Finnish Museum of Natural History, University

of Helsinki, Finland

[†] Equal contribution

* Corresponding author: stefano.mammola@cnr.it

** Corresponding author: pedro.cardoso@helsinki.fi

Author contribution

Equal contribution: both authors conceived the idea, developed the code, contributed critically to

the drafts, and gave final approval for publication.

Summary

1. The use of kernel density n-dimensional hypervolumes [*Global Ecol. Biogeogr.* 23(5):595–609] in trait-

based ecology is rapidly increasing. By representing the functional space of a species or community as a

Hutchinsonian niche space, this relatively new approach is showing great potential for the advance of

functional ecology theory.

2. Functions for calculating the standard set of functional diversity (FD) indexes—richness, divergence, and

regularity—have not been developed in the context of kernel density hypervolumes yet. This gap is delaying

a full exploitation of the kernel density n-dimensional hypervolumes framework in functional ecology,

meanwhile preventing the possibility to compare its performance with that of other FD methods.

3. We develop a set of functions to calculate FD indices based on kernel density n-dimensional

hypervolumes, including alpha (richness), beta (and respective components), dispersion, evenness,

contribution, and originality. Altogether, these indexes provide a coherent framework to explore the primary

mathematical components of FD within a kernel density setting. These new functions can work either with

Hypervolumes objects, HypervolumeList objects, or raw data (species presence or abundance and their traits)

as input data, and are versatile in terms of computation speed, which is achieved by controlling the number

of stochastic points to be used in the different estimations.

4. These newly developed functions are implemented within the R package BAT, an open platform for

biodiversity assessments. As a coherent corpus of functional indexes based on a common algorithm, it opens

up the possibility to fully explore the strengths of the Hutchinsonian niche concept in community ecology

research.

Keywords: alpha diversity, BAT R package, beta diversity, functional metrics, Huntchinsonian hypervolume,

hyperspace, fundamental niche, traits

Introduction

The past two decades have seen an increasing interest on different facets of biodiversity, namely on the way species interact among them and with their environment, with the resulting exponential growth of functional ecology studies. This push was primarily driven by a few milestone theoretical contributions illustrating a way of reshaping entire areas in ecology from a functional perspective, including population and community ecology (Petchey & Gaston, 2002; McGill, Enquist, Weiher, & Westoby, 2006; Bolnick et al., 2011; Violle et al., 2012), conservation biology (Wellnitz & Poff, 2001; Cadotte, Carscadden, & Mirotchnick, 2011), and biogeography (Violle, Reich, Pacala, Enquist, & Kattge, 2014). The fast development of functional ecology theory has led to a proliferation of approaches for estimating functional diversity (FD) (Legras, Loiseau, & Gaertner, 2018), as well as indexes for measuring different mathematical facets underlying FD (Mouillot, Mason, Dumay, & Wilson, 2005; Villéger, Mason, & Mouillot, 2008; Mouchet, Villéger, Mason, & Mouillot, 2010; Schleuter, Daufresne, Massol, & Argillier, 2010; Cardoso, Rigal, Borges, & Carvalho, 2014).

Three main mathematical objects are typically used to represent FD: functional trees (Petchey & Gaston, 2002), convex hull hypervolumes (Cornwell, Schwilk, & Ackerly, 2006), and kernel density hypervolumes (Blonder, Lamanna, Violle, & Enquist, 2014; Blonder et al., 2018) (Figure 1). In the programming language R—likely the main analytical environment used in ecology and evolution (Tippmann, 2015)—these three frameworks have been implemented, respectively, in the packages *BAT* (Cardoso, Rigal, & Carvalho, 2015; Cardoso, Mammola, Rigal, & Carvalho, 2020), *FD* (Laliberté, Legendre, & Shipley, 2014), and *hypervolume* (Blonder, 2018). The *BAT* package is an open platform that provides an assortment of functions for biodiversity assessment (Cardoso et al., 2015), with a special focus on the estimation of the three main components of biological diversity—taxonomic (TD), phylogenetic (PD) and functional (FD) diversity. For calculating FD, this package uses a functional tree in the sense of Petchey & Gaston

(2002, 2006). The *FD* package is possibly the most widely used R package for functional analyses

(over 750 citations as of November 2019 based on Google Scholar metrics). *FD* primarily relies on

convex hull for delineating the functional space (Laliberté & Legendre, 2010), although alternative

formulations and indexes are implemented as well. Finally, the *hypervolume* package relies on the

concept of multi-dimensional niche in the sense of Hutchinson (1957). This is achieved by

delineating the shape and volume of the niche hyperspace, whether functional or bioclimatic, using

different high-dimensional kernel density estimation methods (Blonder et al., 2018)

Both BAT and FD include a vast collection of functional indexes, allowing a user to explore

a functional community based on different mathematical aspects of FD, namely the richness,

divergence, and regularity components (Pavoine & Bonsall, 2011; Tucker et al., 2017). As of yet,

analogous indexes have not been developed in the context of kernel density n-dimensional

hypervolumes. This gap is hindering the use of the hypervolumes framework in functional ecology

studies and is preventing the possibility to fully compare the performance of these alternative

representations of FD. Here, we develop and describe a new set of R functions to calculate the

analogous of standard functional metrics with kernel density n-dimensional hypervolumes, all

available through the R package *BAT* (Cardoso et al., 2020).

New functionalities in BAT

The new functions presented here represent the analogous of the functional indexes that were

originally developed in *BAT* for functional trees (Cardoso et al., 2015). Each new function is briefly

illustrated below.

INPUT DATA

The calculation of the new indexes requires the user to provide either: i) a 'Hypervolume' class R

object; ii) a list of hypervolumes ('HypervolumesList' class R object); or iii) raw data matrices with

species composition of the communities and their functional traits. 'Hypervolume' and

'HypervolumeList' class objects can be constructed using the respective functions of the

hypervolume package (Blonder, 2018), namely one of the three algorithms currently implemented

for hypervolume delineation (box, gaussian, or svm). Full details on these approaches have been

discussed elsewhere (Blonder et al., 2014, 2018; Blonder, 2016). Alternatively, hypervolumes can

be constructed directly within the package BAT by feeding the functions with raw data, namely a

sites x species matrix with incidence or abundance data about the species in the community and a

species x traits matrix.

Note that, although specifically developed for functional hypervolumes, these new functions

can also be applied to other input data. For example, the functions *kernel.beta* and *kernel.similarity*

could be used to explore the bioclimatic niche overlap between two or more species.

FUNCTIONS FOR ESTIMATING FD IN A KERNEL DENSITY CONTEXT

kernel.alpha

This function allows the user to calculate functional richness in a kernel context. Alpha FD is

simply the total volume of the functional niche hyperspace—but see Carmona et al. (2016b, 2016a,

2019) for an alternative estimation based on trait probability density (TPD).

kernel.beta

This function can be used to estimate beta diversity based on the framework recently proposed by

Carvalho & Cardoso (2018). The function computes a pairwise decomposition of the overall

differentiation among kernel hypervolumes into two components: the replacement (shifts) of space

between hypervolumes and net differences between the amount of space enclosed by each

hypervolume. This can be used for decomposing beta diversity into its drivers (Carvalho, Cardoso,

& Gomes, 2012; Carvalho, Cardoso, Borges, Schmera, & Podani, 2013) in a similar way as already

proposed for functional trees (Cardoso, Rigal, Carvalho, et al., 2014).

kernel.contribution

This function evaluates the contribution of each observation to the total volume of the kernel *n*-

dimensional hypervolume. It does so via a leave-one-out approach, whereby the contribution of

each observation to the total functional space is calculated as the difference in volume between the

full hypervolume and a second hypervolume constructed without that specific observation. Note

that, interestingly, the contribution of an individual observation could be negative, if the removal of

this observation increases the volume of the total hypervolume (Figure 2). This might happen,

although not always, in cases when the presence of a given species decreases the average distance

between all the species in the community, i.e., when a given species is close to the "average"

species of that community, making that community less diverse in some sense. By definition, this

does not happen in the case of functional trees, since contribution can only be positive.

kernel.dispersion

In the kernel context, dispersion can be represented as the average dissimilarity between any two

random points of a kernel density hypervolume. Note that this function is the analogous of the

function *dispersion* in BAT, based on functional trees. Conversely, in the package FD, the name

"dispersion" [Fdis; (Laliberté et al., 2014)] is used to represent the average distance of all functional

elements to the centroid.

kernel.evenness

This function measures the functional evenness of an hypervolume, namely the regularity of

functional elements distribution within the total functional space (Schleuter et al., 2010). Following

a similar approach to that of Carmona et al. (2016b), evenness is calculated as the overlap between

the input hypervolume and a second, imaginary hypervolume where traits and abundances are

evenly distributed within their possible range.

kernel.originality

A measure of the originality of each observation used to construct the hypervolume (Pavoine,

Ollier, & Dufour, 2005). This is calculated as the average distance between each observation to a

sample of stochastic points within the boundaries of the hypervolume.

kernel.similarity

A versatile function to compare multiple hypervolumes objects, by calculating their pairwise

centroid distance, minimum distance, intersection, and Jaccard and Sørensen-Dice similarity. This

function is based on a recent simulation study comparing the correlation and performance among

these indexes (Mammola, 2019).

Example analysis

To illustrate the usage of the new functions, we provide an example analysis based on a recent study

of ground-dwelling arthropods in sampling plots characterized by different levels of urbanization

within the municipality of Turin (NW-Italy) (Piano, Giuliano, & Isaia, 2020). R code and databases

to generate the analysis are available in Figshare (doi: 10.6084/m9.figshare.11708277). In order to

test the effects of patch isolation within the urban matrix on the functional diversity of arthropods

assemblages, the authors identified two types of subplots within each plot, where they used pitfall

traps to sample the ground-dwelling fauna (spiders and ground beetles). In particular:

i) isolated subplots are patches within a traffic roundabout, surrounded by roads; and

ii) connected subplots are patches in green areas, connected with the surrounding environmental

matrix and thus more easily colonized by ground-dwelling arthropods.

We extracted from this database a random subset of 60 subplots, 30 isolated and 30 connected. For

each subplot, we extracted abundance data for the spiders community and three functional traits for

each species: body length, dispersal strategy, and hunting strategy. Categorization of hunting

strategies is based on Cardoso et al. (2011), whereas three type of dispersal (1= non- or sporadic

ballooners; 2= ballooners at juvenile stages only; 3= ballooners at all life stages) were assigned by

Piano et al. (2020) based on multiple literature sources (Bell, Bohan, Shaw, & Weyman, 2005;

Blandenier, 2009; Simonneau, Courtial, & Pétillon, 2016). Note that we analyzed here only spider

communities since carabids showed only weak functional signal in the study (Piano et al., 2020).

We used the new kernel-based functions to explore variations in functional diversity

between spider communities in isolated versus connected plots, using *BAT* version 2.0.0. (Cardoso

et al., 2020) and *hypervolume* version 2.0.11. (Blonder, 2018). Since spiders' functional traits

included two categorical traits (Piano et al., 2020), we used the approach proposed by Carvalho &

Cardoso (2018) to incorporate categorical variables in hypervolume estimation. We applied a

Gower dissimilarity measure (Gower, 1971) to the complete trait matrix and then, we analyzed the

resulting distance matrix through Principal Coordinate Analysis (PCoA) in order to extract

orthogonal morphological axes for hypervolume delineation. We retained the first three PCoA axes

to construct hypervolumes using the box method and a default bandwidth (Blonder et al., 2018). We

then estimated α functional diversity of each community with the function *kernel.alpha*.

While the functional spaces of spider community in isolated versus connected plot were

highly interspersed (Figure 3), we observed that, on average, connected plots connected plots had a

higher functional alpha diversity (Figure 4a). This is in accordance with the fact that connected plot

are more easily colonized by a larger number of spider species, that in turn accounts for a greater

diversity of traits. Isolated subplots, on the other hand, are colonized by a subset of specialized

species possessing specific dispersal traits that enhance the colonization of these isolated habitats.

In fact, traits of spider communities in connected plots were more dispersed (Figure 4b) and

homogeneous (Figure 4c) than those in isolated subplots, as estimated using the functions

kernel.dispersion and *kernel.originality*.

Furthermore, we used the *kernel.beta* function to explore variations in traits among

communities inhabiting the two different types of plots. Total β functional diversity values were

higher in pairwise comparisons between communities in connected plots, whereas communities in

isolated plots were in general more similar one another (Figure 4d). This result further confirms

that only spiders with specific traits—in particular those with higher dispersal potential and

probably higher resistance to highly disturbed habitats—are able to reach and survive in isolated

plots. In isolated subplots, the replacement component of β diversity was more preponderant than in

connected subplots (Figure 4e), whereas the richness component was higher in connected plots

(Figure 4f). Differential distributions of β diversity values between the two plot types are

attributable to differences in their effective area. Isolated plots (roundabouts) are always about the

same size, and their area is smaller than those of connected subplots (Piano et al., 2020). Connected

subplots, on the other hand, can be in the middle of both small and large areas. There areas are

characterized by a greater variety of environmental conditions (Piano E., pers. Comm. 2020), hence

the higher differences in β diversity among connected subplots.

By using the function *kernel.contribution*, we further inferred that *Nurscia albomaculata*

(Lucas, 1846) (Araneae: Titanoecidae) is the species contributing the most to the functional space in

isolated subplots, and *Micaria coarctata* (Lucas, 1846) (Gnaphosidae) the most contributing in

connected plots. This is apparently because *N. albomaculata* was the only species of large size (>10

mm) and with low dispersal (no ballooning) that fell in pitfall traps in isolated plots in the study by

Piano et al. (2020), thus it contributed unique traits to roundabouts communities. We finally used

the *kernel.originality* to estimate the functional originality of each spider species included in the

dataset (result not shown), finding that average functional originality was not significantly different

between isolated and connected plots (t-test: −1.23, df= 68.2, p= 0.22), which is not unexpected

since the pool of spider species found in urban environments are already filtered for a small

diversity of traits.

Conclusions

By developing robust functions for estimating the primary facets of FD—the richness, divergence,

and regularity components (Pavoine & Bonsall, 2011; Tucker et al., 2017),—we open up the

potential to fully exploit the strengths of the Hutchinsonian niche concept in functional ecology

research (Blonder, 2019). As in the case of other frameworks for estimating FD (e.g., Mouchet et

al., 2010; Junker, Kuppler, Bathke, Schreyer, & Trutschnig, 2016; Carmona et al., 2019), these new

functions can be used to answer a large assortment of questions. Alpha and beta FD allow us to

understand the richness of a functional hyperspace, and in which components multiple functional

hyperspaces differ, namely the pairwise richness or replacement difference in term of functional

traits. Contribution and originality allow to understand which are the most unique functional

elements, i.e. the species or individuals contributing the most to the overall functional richness of

the hyperspace. Therefore, these indexes yield potential to help in targeting keystone species or

those of high conservation concern. Dispersion and evenness provide a quantification of the

distribution of elements in the functional space enabling, for example, to compare whether multiple

assemblages are differentially harmonious/unbalanced in their functional composition.

CITATION

Researchers relying on these functions should cite this article and in addition can also cite the BAT

package directly. Updated citation information can be obtained by typing in R the command

citation('BAT'). The method and parameters used for constructing kernel density *n*-dimensional

hypervolumes have also to be specified, referring to the publications by Blonder et al. (2014, 2018).

Acknowledgement

Special thanks to Elena Piano for providing data and informations to test the new functions. We are

grateful to Benjamin Blonder for creating the hypervolume framework upon which our methods

heavily rely.

Data availability

Data and R scripts to generate the analyses are available in figshare (doi:

10.6084/m9.figshare.11708277).

Literature cited

Bell, J. R., Bohan, D. A., Shaw, E. M., & Weyman, G. S. (2005). Ballooning dispersal using silk: world fauna, phylogenies, genetics and models. *Bulletin of Entomological Research*, *95*(2), 69–114. doi:10.1079/ber2004350

Blandenier, G. (2009). Ballooning of spiders (Araneae) in Switzerland: General Results from an Eleven-Year Survey. *Arachnology*, *14*(7), 308–316. doi:10.13156/arac.2009.14.7.308

Blonder, B. (2016). Do hypervolumes have holes? *The American Naturalist*, *187*(4), 93–105. doi:10.1086/685444

Blonder, B. (2018). hypervolume: High Dimensional Geometry and Set Operations Using Kernel Density Estimation, Support Vector Machines, and Convex Hulls. Retrieved from https://cran.r-project.org/package=hypervolume

Blonder, B. (2019). Hypervolume concepts in niche- and trait-based ecology. *Ecography*, *41*(9), 1441–1455. doi:10.1111/ecog.03187

Blonder, B., Lamanna, C., Violle, C., & Enquist, B. J. (2014). The n-dimensional hypervolume. *Global Ecology and Biogeography*, *23*(5), 595–609. doi:10.1111/geb.12146

Blonder, B., Morrow, C. B., Maitner, B., Harris, D. J., Lamanna, C., Violle, C., … Kerkhoff, A. J. (2018). New approaches for delineating n-dimensional hypervolumes. *Methods in Ecology and Evolution*, *9*(2), 305–319. doi:10.1111/2041-210X.12865

Bolnick, D. I., Amarasekare, P., Araújo, M. S., Bürger, R., Levine, J. M., Novak, M., … Vasseur, D. A. (2011). Why intraspecific trait variation matters in community ecology. *Trends in Ecology & Evolution*, *26*(4), 183–192. doi:https://doi.org/10.1016/j.tree.2011.01.009

Cadotte, M. W., Carscadden, K., & Mirotchnick, N. (2011). Beyond species: functional diversity and the maintenance of ecological processes and services. *Journal of Applied Ecology*, *48*(5), 1079–1087. doi:10.1111/j.1365-2664.2011.02048.x

Cardoso, P., Mammola, S., Rigal, F., & Carvalho, J. C. (2020). BAT: Biodiversity Assessment Tools. Retrieved from https://cran.r-project.org/package=BAT

Cardoso, P., Pekár, S., Jocqué, R., & Coddington, J. A. (2011). Global patterns of guild composition and functional diversity of spiders. *PLoS ONE*, *6*(6), e21710. doi:10.1371/journal.pone.0021710

Cardoso, P., Rigal, F., Borges, P. A. V., & Carvalho, J. C. (2014). A new frontier in biodiversity inventory: a proposal for estimators of phylogenetic and functional diversity. *Methods in Ecology and Evolution*, *5*(5), 452–461. doi:10.1111/2041-210X.12173

Cardoso, P., Rigal, F., & Carvalho, J. C. (2015). BAT – Biodiversity Assessment Tools, an R package for the measurement and estimation of alpha and beta taxon, phylogenetic and

functional diversity. *Methods in Ecology and Evolution*, *6*(2), 232–236. doi:10.1111/2041-
210X.12310

Cardoso, P., Rigal, F., Carvalho, J. C., Fortelius, M., Borges, P. A. V., Podani, J., & Schmera, D.
(2014). Partitioning taxon, phylogenetic and functional beta diversity into replacement and
richness difference components. *Journal of Biogeography*, *41*(4), 749–761.
doi:10.1111/jbi.12239

Carmona, C. P., de Bello, F., Mason, N. W. H., & Lepš, J. (2016a). The density awakens: A reply to
Blonder. *Trends in Ecology & Evolution*, *31*(9), 667–669. doi:10.1016/j.tree.2016.07.003

Carmona, C. P., de Bello, F., Mason, N. W. H., & Lepš, J. (2016b). Traits Without Borders:
Integrating Functional Diversity Across Scales. *Trends in Ecology & Evolution*, *31*(5), 382–
394. doi:10.1016/J.TREE.2016.02.003

Carmona, C. P., de Bello, F., Mason, N. W. H., & Lepš, J. (2019). Trait probability density (TPD):
measuring functional diversity across scales based on TPD with R. *Ecology*, *100*(12), e02876.
doi:10.1002/ecy.2876

Carvalho, J. C., & Cardoso, P. (2018). Decomposing the causes for niche differentiation between
species using hypervolumes. *BioRxiv*, 485920. doi:10.1101/485920

Carvalho, J. C., Cardoso, P., Borges, P. A. V., Schmera, D., & Podani, J. (2013). Measuring fractions
of beta diversity and their relationships to nestedness: a theoretical and empirical comparison
of novel approaches. *Oikos*, *122*(6), 825–834. doi:10.1111/j.1600-0706.2012.20980.x

Carvalho, J. C., Cardoso, P., & Gomes, P. (2012). Determining the relative roles of species
replacement and species richness differences in generating beta-diversity patterns. *Global
Ecology and Biogeography*, *21*, 760–771. doi:10.1111/j.1466-8238.2011.00694.x

Cornwell, W. K., Schwilk, D. W., & Ackerly, D. D. (2006). A trait-based test for habitat filtering:
Convex hull volume. *Ecology*, *87*(6), 1465–1471. doi:10.1890/0012-
9658(2006)87[1465:ATTFHF]2.0.CO;2

Gower, J. C. (1971). A General Coefficient of Similarity and Some of Its Properties. *Biometrics*,
27(4), 857–871. doi:10.2307/2528823

Hutchinson, G. E. (1957). Concluding Remarks. *Cold Spring Harbor Symposia on Quantitative
Biology*, *22*, 415–427. doi:10.1101/sqb.1957.022.01.039

Junker, R. R., Kuppler, J., Bathke, A. C., Schreyer, M. L., & Trutschnig, W. (2016). Dynamic range
boxes – a robust nonparametric approach to quantify size and overlap of n-dimensional
hypervolumes. *Methods in Ecology and Evolution*, *7*(12), 1503–1513. doi:10.1111/2041-
210X.12611

Laliberté, E., & Legendre, P. (2010). A distance-based framework for measuring functional diversity
from multiple traits. *Ecology*, *91*(1), 299–305. doi:10.1890/08-2244.1

Laliberté, E., Legendre, P., & Shipley, P. (2014). FD: measuring functional diversity from multiple
traits, and other tools for functional ecology.

Legras, G., Loiseau, N., & Gaertner, J.-C. (2018). Functional richness: Overview of indices and
underlying concepts. *Acta Oecologica*, *87*, 34–44. doi:10.1016/J.ACTAO.2018.02.007

Mammola, S. (2019). Assessing similarity of n-dimensional hypervolumes: Which metric to use?
Journal of Biogeography, *46*(9), 2012–2023. doi:10.1111/jbi.13618

McGill, B. J., Enquist, B. J., Weiher, E., & Westoby, M. (2006). Rebuilding community ecology
from functional traits. *Trends in Ecology & Evolution*, *21*(4), 178–185.
doi:10.1016/J.TREE.2006.02.002

Mouchet, M. A., Villéger, S., Mason, N. W. H., & Mouillot, D. (2010). Functional diversity
measures: an overview of their redundancy and their ability to discriminate community
assembly rules. *Functional Ecology*, *24*(4), 867–876. doi:10.1111/j.1365-2435.2010.01695.x

Mouillot, D., Mason, W. H. N., Dumay, O., & Wilson, J. B. (2005). Functional regularity: a
neglected aspect of functional diversity. *Oecologia*, *142*(3), 353–359. doi:10.1007/s00442-004-
1744-7

Pavoine, S., & Bonsall, M. B. (2011). Measuring biodiversity to explain community assembly: a
unified approach. *Biological Reviews*, *86*(4), 792–812. doi:10.1111/j.1469-185X.2010.00171.x

Pavoine, S., Ollier, S., & Dufour, A.-B. (2005). Is the originality of a species measurable? *Ecology*
Letters, *8*(6), 579–586. doi:10.1111/j.1461-0248.2005.00752.x

Petchey, O. L., & Gaston, K. J. (2002). Functional diversity (FD), species richness and community
composition. *Ecology Letters*, *5*(3), 402–411. doi:10.1046/j.1461-0248.2002.00339.x

Petchey, O. L., & Gaston, K. J. (2006). Functional diversity: back to basics and looking forward.
Ecology Letters, *9*(6), 741–758. doi:10.1111/j.1461-0248.2006.00924.x

Piano, E., Giuliano, D., & Isaia, M. (2020). Islands in cities: urbanization and fragmentation drive
taxonomic and functional variation in ground arthropods. Basic and Applied Ecology. *Basic*
and Applied Ecology.

Schleuter, D., Daufresne, M., Massol, F., & Argillier, C. (2010). A user's guide to functional
diversity indices. *Ecological Monographs*, *80*(3), 469–484. doi:10.1890/08-2225.1

Simonneau, M., Courtial, C., & Pétillon, J. (2016). Phenological and meteorological determinants
of spider ballooning in an agricultural landscape. *Comptes Rendus - Biologies*, *339*(9–10),
408–416. doi:10.1016/j.crvi.2016.06.007

Tippmann, S. (2015). Programming tools: Adventures with R. *Nature*, *517*(7532), 109–110.
doi:10.1038/517109a

Tucker, C. M., Cadotte, M. W., Carvalho, S. B., Davies, T. J., Ferrier, S., Fritz, S. A., … Mazel, F.
(2017). A guide to phylogenetic metrics for conservation, community ecology and
macroecology. *Biological Reviews*, *92*(2), 698–715. doi:10.1111/brv.12252

Villéger, S., Mason, N. W. H., & Mouillot, D. (2008). New multidimensional functional diversity
indices for a multifaceted framework in functional ecology. *Ecology*, *89*(8), 2290–2301.
doi:10.1890/07-1206.1

Violle, C., Enquist, B. J., McGill, B. J., Jiang, L., Albert, C. H., Hulshof, C., … Messier, J. (2012).
The return of the variance: intraspecific variability in community ecology. *Trends in Ecology*
& Evolution, *27*(4), 244–252. doi:10.1016/J.TREE.2011.11.014

Violle, C., Reich, P. B., Pacala, S. W., Enquist, B. J., & Kattge, J. (2014). The emergence and
promise of functional biogeography. *Proceedings of the National Academy of Sciences*,
111(38), 13690–13696. doi:10.1073/pnas.1415442111

Wellnitz, T., & Poff, N. L. (2001). Functional redundancy in heterogeneous environments:
implications for conservation. *Ecology Letters*, *4*(3), 177–179. doi:10.1046/j.1461-
0248.2001.00221.x

 Figure captions

Figure 1. Schematic representation of the three main frameworks for calculating FD – examples are based

on two hypothetical traits and two communities, represented in turquoise and orange. **a**) The functional space

can be represented as a functional tree, whose construction is based on the distance matrix of the species

traits. The alpha diversity of a community can be estimated as the total branch length of a tree linking all

species represented in such community—in the example, the FD of a community with 5 species (in orange)

is shown. **b**) The functional space can be represented as the convex hull comprising the species occupying

the functional space. The alpha diversity of a community can be estimated as the smallest convex volume

that contains the set of points. **c**) The functional space can be constructed using kernel density hypervolumes,

whereby the functional space is approximated as a cloud of stochastic points sampled based on a set of

observations (e.g. the functional elements in the community). The alpha diversity of a community can be

estimated as the total volume of the hypervolume.

Figure 2. Scheme of the procedure to estimate contribution. A functional 2-dimensional hypervolume is

constructed via five observations (A, B, C, D, and E). The contribution of each observation can be computed

with a leave-one-out approach, whereby a new hypervolume is constructed leaving out one observation. In

turn, the contribution value is quantified as the difference in volume between the original hypervolume and

the new one. In this example, the contribution of observation D is positive, because the hypervolume

constructed with observations A, B, C, and E is less voluminous than the initial one. Conversely the

contribution of C is negative, because the observation C is clustered with A and B, which leads a higher

proportion of stochastic points to cluster in the vicinity of these three observations. By removing C, the

stochastic points are "freer" to spread in the functional space, which results in a slight increase in the

volume. Note that contour lines bounding the stochastic point are only drawn for graphical representation,

whereas the real hypervolume volume is based on the stochastic points only.

Figure 3. Pairplots representing the estimated 3-dimensional hypervolumes for communities in Isolated and

Connected subplots, based on the first three axes of the Principal Coordinate Analysis (PCoA) performed on

the Gower dissimilarity matrix of the functional traits. Random points are coloured based on the type of plot

of each hypervolume.

Figure 4. a–c) Functional richness, dispersion, and evenness of spider communities in connected and

isolated plots. **d–f)** Density of *β* functional diversity values for pairwise comparison of communities in

connected and isolated subplots. Total functional *β* diversity is split in two components: βrichness is the

turnover in functional composition explained by species loss/gain alone, and βreplacement is turnover in

functional composition explained by replacement of species alone.