Divergent effects of Western and Mediterranean diets on behavior and monocyte polarization

Corbin S.C. Johnson[1], Carol A. Shively[2], Kristofer T. Michalson[2], Amanda J. Lea[3,4], Ryne J. DeBo[2], Timothy D. Howard[5], Gregory A. Hawkins[5], Susan E. Appt[2], Yongmei Liu[6], Charles E. McCall[7], David Herrington[8], Thomas C. Register[2]*, Noah Snyder-Mackler[1,9-13]*†

Affiliations:

[1]Department of Psychology, University of Washington, Seattle, WA 98195 (CSCJ, NSM)

[2]Department of Pathology, Section on Comparative Medicine, Wake Forest School of Medicine, Winston-Salem, NC 27101 (TCR, CAS, KTM, RJB, SEA)

[3]Lewis-Sigler Institute for Integrative Genomics, Princeton University, Princeton, NJ 08544 (AJL)

[4]Department of Ecology and Evolutionary Biology, Princeton University, Princeton, NJ 08544 (AJL)

[5]Department of Biochemistry, Wake Forest School of Medicine, Winston-Salem, NC 27101 (TDH, GAH)

[6]Division of Cardiology, Duke University School of Medicine (YL)

[7]Section of Molecular Medicine, Wake Forest School of Medicine, Winston-Salem, NC 27101 (CEM)

[8]Section on Cardiovascular Medicine, Wake Forest School of Medicine, Winston-Salem, NC 27101 (DH)

[9]Center for Studies in Demography and Ecology, University of Washington, Seattle, WA 98195 (NSM)

[10]Department of Biology, University of Washington, Seattle, WA 98195 (NSM)

[11]Washington National Primate Research Center, University of Washington, Seattle, WA 98195

(NSM)

[12]School of Life Sciences, Arizona State University, Tempe, AZ 85287 (NSM)

[13]Center for Evolution & Medicine, Arizona State University, Tempe, AZ 85287 (NSM)

*Authors contributed equally

†Lead contact

<u>Correspondence:</u>

Thomas C. Register (register@wakehealth.edu)

Noah Snyder-Mackler (nsnyderm@asu.edu)

Summary

Monocytes and macrophages—important mediators of innate immunity, health, and disease—are

sensitive to diet and stress. Western diets promote inflammation and disease, while

Mediterranean diets reduce inflammation and promote health, although the mechanisms through

which diet alters immune function are unknown. Here, we conducted the first randomized, long-

term diet trial in which macaques were fed either a Western- or Mediterranean-like diet to

determine how diet and behavior interact to influence monocyte polarization–a precursor to

subsequent health outcomes. Monocyte gene expression profiles differed markedly between the

two diets, with over 40% of expressed genes being differentially expressed (FDR<0.05). The

Western diet induced a more proinflammatory monocyte phenotype with changes in monocyte

polarization-associated genes. Diet also shifted the co-expression of gene pairs, including small

RNAs and transcription factors that are associated with metabolism and adiposity in humans. We

also found that diet altered a suite of affiliative and anxiety-associated behaviors, and these

behavioral changes mediated ~20% of the effect of diet on monocyte gene expression. Together,

these findings suggest that diet-induced effects on monocyte polarization are due, in part, to both

direct and behaviorally-mediated effects on monocyte gene expression.

Keywords

Immunogenomics, diet, behavior, monocytes, polarization

Introduction

Modern human diets vary across geography, cultures, and socioeconomic strata, and profoundly impact our health and survival. For example, Western diets–characterized by protein and fat from primarily animal sources, simple sugars, and saturated and n-6 fatty acids–are associated with increased risk for metabolic syndrome, type II diabetes, cardiovascular disease, autoimmune disorders, and depression (Jacka et al., 2010; Manzel et al., 2014), as well as increased mortality (Cordain et al., 2005; Drake et al., 2018; Smil, 1989; Smyth and Heron, 2006). Conversely, Mediterranean diets–characterized by protein and fat from primarily vegetable sources, raw fruits and vegetables, and monounsaturated and n-3 fatty acids–lower the incidence of chronic disease and increase longevity (Farchi et al., 1994; Osler and Schroll, 1997; Romagnolo and Selmin, 2017; Trichopoulou et al., 1995). Recent findings implicate inflammation as a key factor in diet-driven chronic disease, as components of the Western diet can arouse the sympathetic nervous system, increase oxidative stress, and elevate levels of inflammatory markers (Giugliano et al., 2006; Lopez-Garcia et al., 2004). In contrast, components of Mediterranean diets such as n-3 fatty acids and antioxidants promote an anti-inflammatory phenotype (O'Keefe et al., 2008). Together, these findings suggest that diets modulate inflammation, which may underlie many of the subsequent health consequences.

Mounting evidence suggests that dietary effects on inflammatory pathologies are mediated by changes in immune cell regulation (Devêvre et al., 2015; Drescher et al., 2019; Holt et al., 2009; Nanri et al., 2007; Nettleton et al., 2006). For instance, components of Western diets increase circulating inflammatory cytokine concentrations in mouse models (Drescher et al., 2019) and in humans (Holt et al., 2009; Nanri et al., 2007; Nettleton et al., 2006). The majority of these

inflammatory cytokines are secreted by monocytes, an innate immune cell. Monocytes and

monocyte–derived macrophages are phenotypically variable along a spectrum. In the circulation,

monocytes are generally defined as classical, intermediate, and non-classical, which upon

activation may become polarized along this spectrum which ranges broadly from

proinflammatory (M1-like) to regulatory/reparative (M2-like) phenotypes. An appropriate

balance of monocyte phenotypes is essential for a healthy immune system. Classically-activated

"M1" monocytes respond to proinflammatory cytokines such as tumor necrosis factor (TNF)-α

and interferon (IFN)-γ by becoming macrophages which propagate the inflammatory response to

infection (Mosser and Edwards, 2008). In contrast, M2 activated monocytes mobilize tissue

repair processes and release anti-inflammatory cytokines in response to interleukin (IL)-4, IL-13,

and transforming growth factor (TGF)-β (Mosser and Edwards, 2008). Thus, diet may alter

disease propensity by reprogramming the balance between these proinflammatory and anti-

inflammatory monocyte subsets (Devêvre et al., 2015).

Diet is only one aspect of the environment that impacts health and survival by altering immune

function. Components of the social environment, such as social status and social integration, may

also affect these fitness outcomes by activating inflammatory programs in primary white blood

cells (Cole, 2013; Snyder-Mackler and Lea, 2018; Tung and Gilad, 2013). For instance, socially

subordinate rhesus macaques exhibit increased activation of the inflammatory response through

MyD88-dependent Th1-mediated inflammatory activation in response to a bacterial infection

(Snyder-Mackler et al., 2016). Given the similar molecular pathways through which diet and the

social environment impact health, these two environmental stimuli could compound or mitigate

one another. Moreover, given that changes in food intake can directly alter social behaviors

themselves (Kaplan et al., 1991; Warden and Fisler, 2008), dietary effects could plausibly act

through (or interact with) social environmental effects.

Earlier investigations of diet effects on immune function primarily focused on the effects of

single nutrient modifications, which precluded potentially important synergistic effects of

multiple nutrients in a given diet (Hu, 2002). Indeed, whole-diet modifications are more effective

at lowering heart rate than single-nutrient supplementation (Whelton et al., 1992), and dietary

supplement trials have shown limited efficacy (Kimmig and Karalis, 2013; Ohlow et al., 2017;

Steinhubl, 2008), emphasizing the need to study the effects of the entire diet rather than single

nutrients. Clinical studies of long-term diet effects and social influences on health are

challenging in humans as diet is difficult to control, and social influences are difficult to

document; thus both are limited by their reliance on self-report (Stice and Durant, 2014;

Suchanek et al., 2011). Further, no studies of whole diet manipulation in primates have probed

the molecular mechanisms through which diet can alter immune function–data that are critical to

identifying targets of future therapies and interventions. To address these gaps, we conducted a

long-term diet intervention study in nonhuman primates by feeding them either Mediterranean-

or Western-like diets. After 15 months on the diet, we assessed monocyte polarization and gene

regulation by measuring genome-wide gene expression. As expected, proinflammatory genes

were more highly expressed in animals fed a Western diet relative to a Mediterranean diet,

indicative of shifts in monocyte phenotypes. Diet also affected gene co-expression patterns,

altering large modules of co-regulated genes, and influenced monkey behavioral phenotypes.

Western-fed monkeys became more socially isolated and exhibited more anxiety-associated

behaviors. Interestingly, these behavioral changes mediated some of the effects of diet on

monocyte gene expression. Together, these results suggest both direct and behaviorally-mediated

effects of diet on monocyte polarization that may contribute to chronic inflammatory diseases.

Results

Dietary intervention alters behavior

We fed adult female cynomolgus macaques either a Western-like (hereafter, "Western") or a Mediterranean-like (hereafter, "Mediterranean") diet for 15 months (the equivalent of ~4 years in a human lifespan; Fig. 1A) prior to blood collection and processing. The experimental diets were nutritionally matched with respect to caloric content of macronutrients and formulated to model human diet patterns, as previously described (Shively et al., 2019). Protein and fat were derived primarily from animal sources in the Western diet and plant sources in the Mediterranean diet. Consequently, the two diets differed in their composition of key micronutrients, including fatty acids, polyunsaturated fatty acid ratios, fiber, and sodium (Fig. 1B; see methods and SI Table 1 for a detailed comparison). As expected, the Western diet resulted in significantly increased body weight, caloric intake, body fat, insulin resistance, and hepatosteatosis relative to the Mediterranean diet (Fig. 1C; (Shively et al., 2019)).

Figure 1. Experimental design and diet effects on body weight. **A)** Monkeys were housed in groups of 3-4 animals (*n* = 35 monkeys) and fed standard monkey chow diet for 8 months before being fed experimental diets. Behavioral data were collected during the last 6 weeks of the baseline phase and the first twelve months of the experimental phase. Body weight was measured 5 months prior to, and 14 months after the start of the experimental phase. Monocytes were isolated from blood collected 15 months after the start of the experimental phase. **B)** Experimental diets were isocaloric with respect to macronutrients, but differed in food sources and relative amounts of micronutrients. Orange bars indicate nutrients with higher concentration in the Western diet formulation, while blue bars indicate higher levels of a given nutrient in the Mediterranean diet. See SI Table 1 for a more detailed comparison of the two diets. **C)** Percent change in body weight from baseline after 14 months on the diet ($t_{(23.0)}$ = -3.02, p = 0.0023).

Monkeys fed the Western and Mediterranean diets differed in a suite of behaviors. Monkeys fed

the Mediterranean diet spent significantly more time in body contact (Mann-Whitney U = 280,

p_{BH} = 5.3 x 10$_{-5}$) and resting (U = 267, p_{BH} = 0.0016), whereas those fed the Western diet spent

significantly more time alone (U = 48, p_{BH} = 0.0042; Fig. 2A). There was no significant

difference in behavior between the two diet groups in all other measured behaviors (SI Fig. 1, SI

Table 2), or in any behaviors during the baseline phase (SI Table 2).

We next leveraged the fact that some behaviors were correlated with one another (SI Fig. 2), and

conducted a principal component analysis to reduce the dimensionality to a few behavioral

factors (Benito et al., 2018; Seltmann et al., 2018). Behaviors associated with dominance

interactions—including aggression, submission, and agonisms—all loaded heavily onto the first

principal component, which explained 26.8% of the overall variance in behavior and did not

differ between diets (Welch-Satterthwaite $t_{(29.2)}$= 0.0362, p = 0.97; SI Fig. 3, SI Note 1).

The second principal component, which explained 15.2% of the variance in behavior, differed

significantly between the two diets, and thus represented a composite of diet-altered behaviors

(hereafter, DAB). Mediterranean-fed monkeys had significantly higher DAB phenotype scores

than their Western-fed counterparts ($t_{(32.6)}$= -4.63, p = 5.6 x 10-5; Fig. 2B). No other principal

component was significantly correlated with diet and thus DAB captures the variance in behavior

altered by diet (Fig. 4B, SI Table 3). DAB captured a number of anxiety- and socially-relevant

behaviors (SI Fig. 5, SI Table 4). Behaviors associated with social integration (e.g., percent of

time in body contact, rate of initiating friendly interactions) were positively correlated with

DAB, while behaviors related to social isolation and anxiety (e.g., percent of time alone, percent

of time grooming self) were associated with lower DAB values (Fig. 2C). Thus, DAB captured a

measure of social integration associated with consuming a Mediterranean-like diet, and social

isolation and anxiety associated with consuming a Western-like diet.

Figure 2. Diet alters behavioral phenotype. A) Three behaviors were significantly different between the two

diet groups. Monkeys fed the Mediterranean diet spent more time in body contact (p_{BH} = 5.3 x 10-5) and resting (p_{BH} = 1.6 x 10-3) than Western-fed monkeys. Monkeys eating the Western diet spent more time alone than Mediterranean-fed monkeys (p_{BH} 4.2 x 10-3). **B)** Composite measures of diet-altered behavior (DAB) were significantly higher in Mediterranean diet compared to Western diet animals ($t_{(32.0)}$= 5.30, p = 8.2 x 10-6). **C)** Eleven of the 32 behaviors observed are significantly correlated with DAB score (p < 0.05). Here, significant correlations with DAB in which behaviors are more frequent in Mediterranean diet or Western diet monkeys are indicated with blue or orange bars, respectively.

Diet alters monocyte gene expression and regulation

To test how diet affected circulating monocytes, we used RNA sequencing to measure genome-

wide gene expression of purified CD14+ monocytes after 15 months on the experimental diets.

The first principal component of genome-wide gene expression, which explained 59.2%

variance, was significantly associated with diet ($t_{(25.1)}$ = -4.41, p = 1.7 x 10-4; Fig. 3A), and 40%

of the 12,240 expressed genes (SI Table 5A) were significantly differentially expressed between

the two diets (n = 4,900 genes, FDR < 0.05; SI Table 5B).

The number of diet-responsive genes was roughly balanced between those that were more highly

expressed in monkeys fed the Mediterranean diet (hereafter "Mediterranean genes"; n = 2,664)

and those that were more highly expressed in monkeys fed the Western diet (hereafter "Western

genes"; n = 2,236). While balanced in direction, the effects of the two diets differed significantly

in magnitude ($t_{(3124.9)}$ = -26.0, p = 4.9 x 10-135). The effect size of diet on Western genes was, on

average, 1.6-fold larger than on Mediterranean genes (Fig. 3B). Thus, the Western diet induced

stronger perturbations in monocyte gene expression than the Mediterranean diet.

Figure 3. Diet affects monocyte gene expression. A) Diet significantly predicts the first principal component of gene expression (59.3% variance explained, $t_{(25.0)}$ = -4.41, p = 1.72 x 10-4). **B)** The effect of diet on genes that were more highly expressed in Western-fed monkeys ("Western genes") was, on average, 1.6-fold larger than "Mediterranean genes" ($t_{(3124.9)}$ = -26.0, p = 4.9 x 10-135; mean effect sizes indicated by dashed lines). The effect sizes for differentially expressed genes are shown (FDR < 0.05, n = 4,900 genes). **C)** The proportion of Western genes (orange) and Mediterranean genes (blue) in the subsets of proinflammatory (top) and regulatory (bottom) genes are presented as log2 fold change from the null expectation (proportion of all genes that are proinflammatory or regulatory). Error bars were calculated through 10,000 permutations of the data. Western genes were enriched for proinflammatory (M1-like) genes (fold-enrichment = 1.27, *95% CI* = 1.09, 1.46), whereas in the Mediterranean genes the same proinflammatory M1-like gene set was under-represented (fold-enrichment = 0.74, *95% CI* = 0.61, 0.88). Regulatory (M2-like) genes were also under-represented in Mediterranean genes (fold-enrichment = 0.67, *95% CI* = 0.40, 0.97), but not in Western genes (fold-enrichment = 0.95, *95% CI* = 0.60, 1.35).

The most strongly affected Western genes included well-known inflammatory-related genes,

such as including interleukin-6 (β_{diet} = -1.66, FDR = 8.9x10-3; Fig. 3B), interleukin-1α (β_{diet} = -

1.22, FDR = 0.03; SI Table 5B), and two subunits of the NF-κB protein (*NFKB1* β_{diet} = -0.30,

FDR = 0.017; *NFKB2* β_{diet} = -0.42, FDR = 0.012). Western genes were enriched for biological

processes critical to anabolic regulation of biomass content and replication, including response to

growth factor (GO:0070848, *p* = 4.6×10^{-3}) and response to insulin (GO:0032868, *p* = 1.4×10^{-3};

SI Table 6A), suggesting that the Western diet reprogrammed proinflammatory immune, and

oxidative metabolic aspects of monocyte gene regulation. Meanwhile, Mediterranean diet

promoted antioxidation reductive processes (GO:0055114, *p* = 4.3×10^{-3}; SI Table 6B), a critical

function in muting proinflammatory monocytes.

Next, to identify putative upstream gene regulatory mechanisms, we examined if diet-induced

changes in gene expression were accompanied by specific transcription factor binding sites. We

identified 34 distinct transcription factor-binding motifs enriched within 2 kilobases of the

transcription start sites of Mediterranean genes and one that was enriched near the transcription

start sites of Western genes (FDR < 0.05; SI Table 7). Among the transcription factors with

motifs enriched in Mediterranean gene promoters were key transcription factors—such as PU.1,

SP1, and MEF2C—that drive monocyte fate and functional differentiation in response to

extracellular signals (Schuler et al., 2008; Scott et al., 1994; Zhang et al., 1994). The binding

motifs of three interferon-regulatory factors (IRF) that reprogram monocyte differentiation and

polarization were enriched near transcription start sites of Mediterranean genes (Chistiakov et al.,

2018). In addition to enrichment of their binding motifs, IRF3, which promotes M2 macrophage

polarization and anti-viral response to TLR4 stimulation, and IRF8, which supports monocyte

maturation (Günthner and Anders, 2013), also had higher gene expression in Mediterranean-fed

monkeys. The sole transcription factor with binding sites enriched in Western diet-associated

genes, ATF2, is a key mediator of inflammatory pathways and diseases, including response to

bacterial endotoxin, atherosclerosis, and obesity (Fledderus et al., 2007; Miyata et al., 2012;

Reimold et al., 2001). Interestingly, Western genes were enriched for activation of the MAPKK

pathway, which lies upstream of ATF2 transcription factor activity (GO:0000186, $p = 4.7 \times 10^{-3}$,

SI Table 6A) (Herlaar and Brown, 1999). Thus, some of the diet-associated changes in monocyte

gene regulation may be mediated by changes in the activity and binding of key transcription

factors.

We next conducted a more targeted analysis of monocyte polarization, by focusing on genes that

were previously reported to be differentially expressed between induced proinflammatory (M1)

and regulatory (M2) monocyte polarization as defined in (Schmidl et al., 2014) (see SI Table

5A,B for polarization categories). Western genes were enriched for M1-associated genes ($n =$

162 genes, 27.1% more than expected, *95% CI = +9.0%, +45.9%*; Fig. 3C), but not M2-

associated genes ($n = 24$ genes, 4.8% fewer than expected, *95% CI = -59.5%, +34.9%*).

Conversely, the Mediterranean gene set was depleted for both M1-associated genes ($n = 112$

genes, 26.2% fewer than expected, *95% CI = -38.9%, -14.4%*) and M2-associated genes ($n = 20$

genes, 33.3% fewer than expected, *95% CI = -60.0%, -3.3%*). Together, these observations

indicate that a Western diet induces a more proinflammatory (M1-like) phenotype thus providing

a target for mechanistic follow-up studies.

Finally, we explored the potential for diet to alter the magnitude or direction of pairwise gene

expression correlations among the most strongly diet-affected genes. If so, this could reveal gene

regulatory networks that are altered by diet (de la Fuente, 2010; Gaiteri et al., 2014). Many gene

pairs were identified that were significantly associated with each other in both diets, both

positively ($n = 714$) and negatively ($n = 332$, $p < 0.05$; SI Table 8A). Drawing on a newly

developed approach, correlation by individual level product (CILP; (Lea et al., 2019)), we

identified 445 gene pairs that were differentially correlated in Mediterranean- versus Western-

fed monkeys at a relaxed FDR < 20% (SI Table 8B). The majority (97%) of these gene pairs

exhibited positive associations in one diet and negative associations in the other, as we have the

most power to detect differential correlation in gene pairs in which the direction of correlation is

opposite in the two diets. Some of these gene pairs were significantly correlated in reversed

directions ($n = 23$ gene pairs; nominal p-value < 0.05), suggesting that diet can reverse the co-

expression relationship between two genes (Figure 4A). Importantly, we identified 16 "hub"

genes that exhibited differential correlations with partner genes more so than expected by chance

(Fig. 4B, SI Table 8C). These hub genes were enriched for genes encoding transcription factors

(Fisher's Exact Test, FET, OR = 7.40, $p = 7.0$ x 10_{-3}), including SOX4 (essential for normal

insulin secretion and glucose tolerance) and NR4A2 (involved in lipid, carbohydrate, and energy

metabolism; (Goldsworthy et al., 2008; Pearen and Muscat, 2010)), thereby implicating immune

and metabolic reprogramming by the Western and Mediterranean diet. Interestingly, the hub

gene involved in the greatest number of differentially-correlated gene pairs was *RF00283*–a non-

coding RNA that has been associated withed BMI and HDL cholesterol in human genome-wide

association studies ((Davis et al., 2017; Kanai et al., 2018; Tachmazidou et al., 2017); Fig. 4B-

D).

Figure 4. Diet affects monocyte gene co-expression. A) The Pearson correlation between a given pair of genes within each of the experimental diets are plotted against each other. Gene pairs that are significantly differently correlated between diets are highlighted in blue (n = 445 significant pairs, FDR < 20%). **B)** Of the genes involved in significant pairs, some were paired with more genes than expected by chance (n = 16 "hub" genes; dotted black line is the maximum number of significant pairs expected by chance). The strongest hub gene was the non-coding RNA *RF00283*. **C)** Residual normalized expression of *RF00283* is significantly greater in Western- than Mediterranean-fed monkeys (β_{diet} = -0.507, FDR = 2.3 x 10-6). **D)** Example of a differential correlation involving *RF00283*. Residual normalized expression of *RF00283* is plotted against expression of *KLF11*, a differentially-expressed transcription factor that regulates insulin and has been associated with type II diabetes in humans (Neve et al., 2005). The two genes are positively correlated in Western-fed monkeys (r = 0.61, p<0.005) and negatively correlated in Mediterranean-fed monkeys (r = -0.63, p<0.01).

Diet-altered behaviors mediate some changes in monocyte gene expression

The strong effects of diet on both behavior and monocyte gene expression led us to examine if

the effect of diet on gene expression was mediated by diet-induced changes in behavior. Of the

4,900 diet-affected genes, 34% were also significantly associated with DAB in a univariate

model ($n = 1,679$, $p < 0.05$). Of these, DAB significantly mediated the effect of diet on the

expression of 872 genes (18% of all diet-associated genes, bootstrapped $p < 0.05$; Fig. 5A).

These DAB-mediated genes were significantly more likely to be Western genes ($n = 482$, 55%,

two-sided binomial test $p = 0.0020$), and were enriched in biological processes such as regulation

of leukocyte migration (FET $p_{BH} = 0.0042$; SI Table 9A-C). We also tested the possibility that

diet effects on behavior were mediated through changes in monocyte gene expression but found

no genes that significantly mediated the relationship between diet and DAB (bootstrapped $p <$

0.05; Fig. 5B).

Figure 5. Behavior partially mediates the effect of diet on gene expression for 20% of diet-associated genes.
A) Behavior significantly mediated the effect of diet on gene expression in 872 genes ($n = 482$ higher in Western,
390 higher in Mediterranean). **B)** Conversely, gene expression did not significantly mediate the effect of diet on
DAB in any genes.

Discussion

Here, we showed for the first time that a controlled, whole-diet manipulation exerts profound

effects on monocyte gene regulation and behavior in a primate. Forty percent of the expressed

genes were differentially expressed between monkeys fed Western or Mediterranean diets,

indicating that diet dramatically altered monocyte programming. Beyond differential gene

expression, we identified differences in gene co-expression and enrichment of transcription

factor binding motifs, suggesting that diet alters gene regulatory networks. More specifically, we

found that the Western diet promoted a more proinflammatory monocyte phenotype, while the

Mediterranean diet had a more benign effect suggesting better balance in monocyte polarization.

We identified the enrichment of binding motifs for numerous transcription factors that appear to

be involved in diet-regulated gene expression. Of note, members of the E26 transformation-

specific (ETS), specificity protein (Sp)/Krüppel-like family (KLF), myocyte-specific enhancer

factor (MEF), and interferon-regulatory factor (IRF) families of transcription factors have all

been linked to myeloid differentiation (Chistiakov et al., 2018; Schuler et al., 2008; Scott et al.,

1994; Zhang et al., 1994) and were overrepresented in regulatory regions of genes with higher

expression in monkeys fed the Mediterranean diet in this study. Broadly, this suggests that

transcriptional networks involved in monocyte differentiation were differentially affected by our

diet manipulation. IRF-1 and IRF-8 are linked to M1 monocyte polarization, while IRF-3 is

associated with M2 polarization, and all three had binding motifs enriched in genes more highly

expressed in monkeys fed the Mediterranean diet (i.e., more lowly expressed in monkeys fed a

Western diet).

The Western diet promoted a more proinflammatory monocyte polarization relative to

Mediterranean diet, which is consistent with that seen in brain myeloid cells of mice fed a

Western diet (Yang et al., 2019). Conversely, Mediterranean diet could reduce proinflammatory

gene expression, as has been suggested in at least one study in humans (Camargo et al., 2012).

Future comparisons of each diet to the current standard diet for captive primates will help to

identify which of these two non-mutually exclusive hypotheses is most likely. It is also worth

noting that the M1/M2 paradigm of monocyte polarization is a simplification of the

heterogeneity that exists within organisms (Martinez and Gordon, 2014; Nahrendorf and Swirski,

2016) and the patterns of gene expression following the diet manipulation described here may

indicate alternative phenotypes that fall outside of the simplified M1/M2 binary.

In addition to altering monocyte gene expression, this diet manipulation also led to differences in

a suite of important behaviors. Monkeys consuming the Western diet exhibited more anxiety and

social isolation, a phenotype remarkably similar to that observed in juvenile Japanese macaques

born to mothers consuming a high-fat Western diet (Thompson et al., 2018). In that study,

offspring behavior was associated with maternal levels of macrophage-derived chemokine

(MDC), which showed higher expression in Western-diet fed animals in our study (β_{diet} = -0.243,

FDR = 0.059). Our findings extend the behavioral effects of a Western diet into adulthood.

Future work will investigate if the direct diet effects on adult behavior are mechanistically

similar to the transgenerational diet effects identified in Japanese macaques born to high-fat-fed

mothers.

For a subset (18%) of genes, the diet-altered behavior (DAB) phenotype mediated the effect of

diet on monocyte gene expression. Notably, the Western diet induced both a proinflammatory

monocyte phenotype, as well as social isolation and anxiety behaviors. It is, therefore, possible

that a Western diet contributes to inflammation by producing a more socially isolated or anxious

animal. Indeed, monocytes have been shown to be responsive to social stimuli, including social

isolation (Cole, 2019) and anxiety (Cole et al., 2015), suggesting that part of the effect of diet on

monocyte function may be due to diet-induced changes in socially relevant behaviors. In support

of this hypothesis, animals fed a Western diet exhibited significantly higher expression of pro-

inflammatory genes involved in the "conserved transcriptional response to adversity" (CTRA;

Wilcox T-test $p = 0.016$; (Cole et al., 2015), and lower expression of antiviral- and antibody-

related CTRA genes (Wilcox T-test $p = 0.023$; SI Fig. 6).

The mechanistic pathways through which diet affects behavior are still unclear. It is possible that

diet induces changes in the central nervous system in myriad ways: including infiltration of

cytokines from the periphery (Raison et al., 2006), stimulation of resident immune cells in the

brain (Yang et al., 2019), alteration of membrane lipids (Du et al., 2016), and direct anti-

inflammatory action of n-3 fatty acids in the brain (Layé et al., 2018). We previously showed

that diet had a strong effect on the microbiome (Nagpal et al., 2018), which is a key component

of the gut-brain axis (Sharon et al., 2016), representing another possible route through which diet

could alter behavior. Diet could also alter behavior through other physiological systems,

including those that monitor satiety and thermoregulation (Aydin and Gordon, 2013; Diekmann

et al., 2019). Future work on the neurobiological responses to diet manipulations could help

clarify these putative mechanisms.

Interestingly, Western diet induced substantial variation in multiple phenotypes, including body

weight, gene expression, and behavior. This heterogeneity in response to diet is consistent with

previous studies demonstrating that some individuals may be more resistant (or susceptible) to

the effects of a Western diet (Shively et al., 2009), presumably due to genetic variation or past

environmental experiences. However, we were unable to identify any consistencies in individual

responsiveness across the phenotypes (SI Fig. 7). For instance, monkeys that exhibited a strong

gene regulatory response to the Western diet did not also exhibit the largest increase in body

weight. Change in body weight was significantly correlated with diet-altered behavior (DAB)

within Western diet fed monkeys ($r_{Western}$ = -0.584, p = 0.0069; SI Fig. 7A), but change in body

weight did not significantly predict the expression of any genes at an FDR < 20%. Western diet

fed individuals thus exhibited a mosaic response to diet across multiple phenotypes, of which the

underlying mechanisms remain unknown but presumably involve interactions between diet,

environment, and the genome.

In summary, we found that diet significantly alters behavior and monocyte polarization. The

Western diet increased the proinflammatory phenotype relative to a diet, which supports the role

of monocyte polarization in diet-associated chronic inflammatory diseases. Thus, avoiding a

Western-style diet and/or consuming a Mediterranean-style diet could be beneficial in preventing

or treating chronic inflammation and disease. The majority of the effects of diet are presumably

mediated through direct or combined actions of saturated/polyunsaturated fats, n-6:n-3 ratios,

pro- and anti-antioxidant characteristics, and other unique features of the protein and

carbohydrate constituents in the two diets. Monocyte reprogramming was also partially mediated

by the diet-induced changes in behavior, although the mechanisms by which this occurred are

unknown. Ongoing and future work will address interactions between social behavior (e.g.,

social status) and diet to further understand how environmental stressors may impact

inflammation in the periphery and in the central nervous system.

Methods

Subjects

At study initiation, subjects were 43 adult (age: mean = 9.0, range = 8.2-10.4 years, estimated by

dentition), female cynomolgus macaques (*Macaca fascicularis*), obtained from Shin Nippon

Biomedical Laboratories (SNBL USA SRC, Alice, TX) and housed at the Wake Forest School of

Medicine Primate Center (Winston-Salem, NC) as described previously (Shively et al., 2019).

Briefly, monkeys were socially housed in groups of 3-4 and consumed standard monkey chow

(SI Table 1) during an eight-month baseline phase, after which they were fed either the Western

(5 groups, *n* = 21) or Mediterranean (6 groups, *n* = 22) diet, balanced on pretreatment

characteristics that reflected overall health, including body weight, body mass index, and plasma

triglyceride concentrations ((Shively et al., 2019); Fig. 1A). Two monkeys did not tolerate the

experimental diet, and were switched to standard monkey chow, three animals died during the

course of the study, and a further three were identified as outliers (see "Removal of Batch

Effects" below), resulting in a sample size of 35 (Western *n* = 20, Mediterranean *n* = 15). All

animal manipulations were performed according to the guidelines of state and federal laws, the

US Department of Health and Human Services, and the Animal Care and Use Committee of

Wake Forest School of Medicine.

Experimental Diets

Experimental diets were formulated to be isocaloric with respect to protein, fat, and carbohydrate

macronutrients, and identical in cholesterol content (~ 320mg / 2000 kilocalories (Cals)/day) as

described previously (Shively et al., 2019). The Western diet was formulated to be similar to that

consumed by American women age 40-49 as reported by the US Dept. Agriculture, with protein

and fat derived mainly from animal sources. The Western diet was relatively high in saturated fat

and sodium, and low in monounsaturated fat and n-3 fatty acids. The Mediterranean diet was

formulated to mimic key aspects of the traditional Mediterranean diet, with an n-6:n-3 fatty acid

ratio similar to a traditional hunter-gatherer type diet (Bédard et al., 2012; Cordain et al., 2005;

Kafatos et al., 2000). Protein and fats were derived mainly from plant sources, fish and dairy,

and monounsaturated fatty acids were relatively high. Mediterranean diet contained more

complex carbohydrates and fiber, and less sodium and refined sugars than Western diet. Key

ingredients included English walnut powder and extra-virgin olive oil which were the primary

components provided to participants in the PREDIMED study, a landmark dietary intervention

study that illustrated the role of the Mediterranean diet on cardiovascular disease prevention

(Estruch et al., 2018). For details of the composition of both diets and how they relate to monkey

chow, see SI Table 1.

Behavioral Characterization

Behavioral data were collected weekly during two 10-minute focal observations, randomly

ordered and balanced for time of day, for 6 weeks during the baseline phase (2 total observation

hours/monkey) and for 12 months during the experimental phase (15 total observation

hours/monkey). Behaviors were collected as previously described (Shively, 1998), and combined

into summary behaviors (e.g., "aggression" was a combination of all total, noncontact, contact

aggressive events). A complete table of all behavioral data can be found in SI Table 2. In order to

quantify the overall impact of diet on behavior, we conducted a principal component analysis

using the R package *FactoMineR* (Lê et al., 2008).

Blood Sample Collection

We briefly removed monkeys from their social group, drew blood via venipuncture within 9
minutes of entering the building, and returned the animal to their group. Blood was collected into
EDTA-containing tubes, mixed with an equal amount of PBS without calcium or magnesium,
and overlaid on a 90% Ficoll-Paque Plus/10% PBS solution in LeucoSep tubes followed by
centrifugation at 800 x g for 20 min. Isolated PBMCs were then immediately used for the
collection of CD14+ monocytes by positive selection using a Miltenyi bead-based protocol
following manufacturer's instructions (Miltenyi Biotec, Bergisch Gladbach, Germany). After
assessing cell viability and numbers, CD14+ monocytes were stored in 85% FBS, 15% DMSO
sterile freezing media at -80°C and transferred to liquid nitrogen for storage until RNA
extraction.

RNA extraction and sequencing

RNA was extracted from monocytes using the AllPrep DNA/RNA Mini Kit (Qiagen, Inc.,
Hilden, Germany), and quantified using a NanoDrop spectrophotometer and Agilent 2100
Bioanalyzer with RNA 6000 Nano chips (Agilent Technology, Inc., Santa Clara, CA). RNA
libraries were prepared for sequencing by the Cancer Genomics Shared Resource (Wake Forest
University, Winston-Salem, NC) using the TruSeq-stranded total RNA kit (Illumina), which
includes a ribosomal depletion step. The RNA-seq libraries were then sequenced using single-
end 76-bp reads on an Illumina NextSeq 500 to an average read depth of 34.5 million reads per
sample (range 25.9 – 41.6 million reads). Reads were mapped to the *Macaca fascicularis*
reference genome (Macaca_fascicularis_5.0, v 93, Ensembl) (Kersey et al., 2018; Kinsella et al.,
2011) using HiSat2 (Kim et al., 2015) and then converted to a sample-by-gene read count matrix

using featureCounts (Liao et al., 2014) (median = 38.0%; range 24.5 - 50.4% of reads mapped to

exons).

Read Count Normalization and Removal of Batch Effects

First, we removed lowly expressed genes (median reads per kilobase per million reads mapped <

1), which resulted in 12,240 genes for our downstream analyses. We normalized read counts

using the *voom* function of the R package *limma* (Ritchie et al., 2015). While investigating

monocyte purity, three samples differed in CD3 gene expression from the rest by several orders

of magnitude. We concluded that these samples were contaminated with CD3+ cells (i.e.,

inefficient CD14 purification, see SI Fig. 8) and excluded them from all analyses, leaving a final

sample size of 35 monkeys (n = 20 WEST, n = 15 MED). To control for batch effects related to

RNA quality and monocyte purity, we calculated the residual gene expression from a model of

normalized gene expression as a function of CD14 expression, CD3 expression, RNA integrity,

and RNA concentration. These residual gene expression values were used for all subsequent

analyses.

Modeling Effect of Diet on Gene Expression

In order to determine which genes were significantly affected by diet, we modeled the residual

expression of each gene as a function of diet using a linear mixed effects model controlling for

relatedness among monkeys using the R package *EMMREML* (Akdemir and Godfrey, 2015).

Relatedness was estimated using the ngsRelate program (Hanghøj et al., 2019) with SNP

genotypes inferred from the RNA-seq reads using bcftools mpileup (Li et al., 2009). We

calculated an empirical false discovery rate (FDR) for each gene using a permutation-based

approach (Snyder-Mackler et al., 2016). Genes that passed a threshold of FDR < 0.05 were

considered differentially expressed between the two diets. To examine global patterns of

variation in gene expression, we conducted principal component analysis on the correlation

matrix of normalized residual gene expression using the *prcomp* function in R.

Enrichment analyses

Gene ontology (GO) enrichment analyses were conducted using Fisher's Exact Tests using the

weight01 algorithm to test for enrichment implemented in the R package *topGO* (Alexa and

Rahnenfuhrer, 2019). For a more targeted analysis of M1 and M2 specific genes, we identified a

set of differentially expressed genes in our data set that were previously found to be involved in

monocyte polarization (Schmidl et al., 2014) (638 proinflammatory and 138 regulatory), which

we used to explore monocyte polarization in the current study. We calculated the proportion of

genes more highly expressed in the Mediterranean- and Western-fed animals in each polarization

category and tested for significance using a permutation test ($n = 100,000$ permutations).

Transcription Factor Binding Site Analysis

We tested for enrichment of transcription factor binding motifs within 2 kb (upstream or

downstream) of the transcription start sites of differentially expressed "Western genes" or

"Mediterranean genes" (FDR < 0.05) using the program HOMER (Heinz et al., 2010) and

equivalent regions around the transcription start sites of all genes expressed in these data as the

background set for enrichment testing. We searched for known vertebrate transcription factor

binding motifs and report the TF motifs passing a threshold of FDR < 0.05.

Gene-gene co-expression analysis

In addition to testing whether diet led to mean differences in gene expression between Western

and Mediterranean animals, we also tested whether diet impacted the correlation structure among

expressed genes (i.e., gene co-expression). Specifically, we used 'correlation by individual level

product' (CILP) (Lea et al., 2019), to test whether diet affected the magnitude or direction of

pairwise gene expression correlations among the top 140 most differentially expressed genes (n

= 9730 gene-gene pairs tested, equivalent to $_{140}C_2$). To test whether a given pair of genes was

differentially co-expressed as a function of diet, we first obtained a vector of products for each

gene pair by multiplying the normalized gene expression values for two genes together.

Normalization was performed by scaling expression values to mean 0 and unit variance within

Mediterranean and Western subsets of the data respectively, to ensure that distributional

differences between sample groups did not bias our results, following previously described

procedures (Lea et al., 2019). Each of these vectors of products were used as the outcome

variable in a linear mixed effects model implemented in the R package *EMMREML* (Akdemir

and Godfrey, 2015), which included a fixed effect of diet and a random effect to control for

genetic relatedness. To assess significance, we extracted the p-value associated with the diet

effect for all 9730 gene pairs. We then reran each linear mixed effects model 100 times after

permuting diet, extracted the p-value associated with the diet effect, and used these values to

calculate an empirical FDR distribution (Snyder-Mackler et al., 2016).

Using this approach, we identified 445 gene pairs that were significantly differentially co-

expressed as a function of diet at a 20% empirical FDR. Next, we performed two follow up

analyses to understand their biological import. First, we tested for the existence of 'hub genes',

defined as genes that displayed differential co-expression to their tested partner genes more so

than expected by chance. To define the null distribution for identifying hub genes, we randomly

sampled 445 gene pairs from the set of all 9730 tested gene pairs 1000 times and calculated the

number of partners a focal gene had in each sample; we considered a gene to be a significant

'hub gene' if it fell outside the 95th percentile of this distribution, which was equivalent to a focal

gene that displayed significant differential co-expression with 13 or more of its tested partner

genes. Second, we asked whether the set of 'hub genes' we identified were enriched for

transcription factors, relative to the background set of all 140 genes tested for differential co-

expression. We performed this analysis because many of the proposed mechanisms to generate

large scale changes in gene co-expression patterns involve changes in transcription factor

function or activity (de la Fuente, 2010; Gaiteri et al., 2014). To implement the enrichment

analysis, we used the TRRUST database of known mammalian transcription factors for

annotation (Han et al., 2018) paired with hypergeometric tests.

Mediation

To test if DAB mediated the effect of diet on gene expression, we conducted mediation analyses.

We used a bootstrapping approach where we conducted 10,000 bootstrap iterations of two

models: (Model 1) the expression of each gene as a function of diet, and (Model 2) the

expression of each gene as a function of diet and DAB (Preacher and Hayes, 2004). For each

bootstrap iteration, we then calculated the mediation effect (i.e., the indirect effect) of DAB as

the difference between the effect size of diet in Model 1 (β_{diet}) and Model 2 (β'_{diet}). We

considered there to be a mediation effect when the 95% confidence interval for the indirect effect

(β_{diet}-β'_{diet}) did not include zero.

A similar method was used to calculate the mediation of gene expression on DAB, testing the

difference between the effect size of diet in two models: (Model 3) DAB as a function of diet,

and (Model 4) DAB as a function of diet and the expression of each gene.

Data Availability

All data and code used to complete these analyses can be found at

https://github.com/cscjohns/diet_behavior_immunity. The raw data can be accessed from the

gene expression omnibus repository from accession # GSE144314.

Acknowledgments

We thank Beth Uberseder, Maryanne Post, JD Bottoms, Edison Floyd, Jason Lucas, Joshua

Long, Diane Wood, and Sherri Samples for technical support. We thank Nicholas Lozier,

Tiffany Pan, and Marina Watowich for their helpful feedback on previous versions of this

manuscript. This work was funded by R01HL087103 (CAS), R01HL122393 (TCR),

U24DK097748 (TCR) from NIH and intramural funding from the Department of Pathology,

Wake Forest School of Medicine (CAS). NSM was supported by R00 AG051764 from NIH, and

AJL was supported by a postdoctoral fellowship from the Helen Hay Whitney Foundation. The

Wake Forest Comprehensive Cancer Center Cancer Genomics Shared Resource is supported by

P30 CA012197 and by a NIH Shared Instrumentation Grant 1S10OD023409-01 to GAH.

Author Contributions

Conceptualization, C.A.S. and T.C.R.; Methodology, C.A.S., T.C.R., N.S.-M., and C.S.C.J.;

Formal Analysis – C.A.S., T.C.R., N.S.-M., A.J.L., and C.S.C.J; Investigation, M.E., A.N.V.,

N.A.V., S.C.P., and S.Y.W.; Writing – Original Draft, C.A.S., T.C.R., N.S.-M., and C.S.C.J.;

Writing – Review & Editing, all authors; Visualization – N.S.-M., A.J.L., and C.S.C.J; Funding

Acquisition, C.A.S. and T.C.R.; Supervision, C.A.S., T.C.R., and N.S.-M.

Declaration of Interests

The authors declare no competing interests.

References

Akdemir, D., and Godfrey, O.U. (2015). EMMREML: Fitting Mixed Models with Known
Covariance Structures.

Alexa, A., and Rahnenfuhrer, J. (2019). topGO: Enrichment Analysis for Gene Ontology.

Aydin, C., and Gordon, C.J. (2013). Thermoregulatory, cardiovascular, and metabolic responses
to mild caloric restriction in the Brown Norway rat. Physiol. Rep. *1*.

Bédard, A., Riverin, M., Dodin, S., Corneau, L., and Lemieux, S. (2012). Sex differences in the
impact of the Mediterranean diet on cardiovascular risk profile. Br. J. Nutr. *108*, 1428–1434.

Benito, X., Fritz, S.C., Steinitz-Kannan, M., Vélez, M.I., and McGlue, M.M. (2018). Lake
regionalization and diatom metacommunity structuring in tropical South America. Ecol. Evol. *8*,
7865–7878.

Camargo, A., Delgado-Lista, J., Garcia-Rios, A., Cruz-Teno, C., Yubero-Serrano, E.M., Perez-
Martinez, P., Gutierrez-Mariscal, F.M., Lora-Aguilar, P., Rodriguez-Cantalejo, F., Fuentes-
Jimenez, F., et al. (2012). Expression of proinflammatory, proatherogenic genes is reduced by
the Mediterranean diet in elderly people. Br. J. Nutr. *108*, 500–508.

Chistiakov, D.A., Myasoedova, V.A., Revin, V.V., Orekhov, A.N., and Bobryshev, Y.V. (2018).
The impact of interferon-regulatory factors to macrophage differentiation and polarization into
M1 and M2. Immunobiology *223*, 101–111.

Cole, S.W. (2013). Social regulation of human gene expression: mechanisms and implications
for public health. Am. J. Public Health *103 Suppl 1*, S84-92.

Cole, S.W. (2019). The Conserved Transcriptional Response to Adversity. Curr. Opin. Behav.
Sci. *28*, 31–37.

Cole, S.W., Levine, M.E., Arevalo, J.M.G., Ma, J., Weir, D.R., and Crimmins, E.M. (2015).
Loneliness, eudaimonia, and the human conserved transcriptional response to adversity.
Psychoneuroendocrinology *62*, 11–17.

Cordain, L., Eaton, S.B., Sebastian, A., Mann, N., Lindeberg, S., Watkins, B.A., O'Keefe, J.H.,
and Brand-Miller, J. (2005). Origins and evolution of the Western diet: health implications for
the 21st century. Am. J. Clin. Nutr. *81*, 341–354.

Davis, J.P., Huyghe, J.R., Locke, A.E., Jackson, A.U., Sim, X., Stringham, H.M., Teslovich,
T.M., Welch, R.P., Fuchsberger, C., Narisu, N., et al. (2017). Common, low-frequency, and rare
genetic variants associated with lipoprotein subclasses and triglyceride measures in Finnish men
from the METSIM study. PLOS Genet. *13*, e1007079.

Devêvre, E.F., Renovato-Martins, M., Clément, K., Sautès-Fridman, C., Cremer, I., and Poitou,
C. (2015). Profiling of the Three Circulating Monocyte Subpopulations in Human Obesity. J.
Immunol. *194*, 3917–3923.

Diekmann, C., Wagner, M., Huber, H., Preuß, M., Preuß, P., Predel, H.-G., Stoffel-Wagner, B.,
Fimmers, R., Stehle, P., and Egert, S. (2019). Acute Impact of Dietary Pattern and Walking on
Postprandial Attention, Mood, and Satiety in Older Adults: A Randomized Crossover Trial.
Nutrients *11*, 2294.

Drake, I., Sonestedt, E., Ericson, U., Wallström, P., and Orho-Melander, M. (2018). A Western
dietary pattern is prospectively associated with cardio-metabolic traits and incidence of the
metabolic syndrome. Br. J. Nutr. *119*, 1168–1176.

Drescher, H.K., Weiskirchen, R., Fülöp, A., Hopf, C., de San Román, E.G., Huesgen, P.F., de
Bruin, A., Bongiovanni, L., Christ, A., Tolba, R., et al. (2019). The Influence of Different Fat
Sources on Steatohepatitis and Fibrosis Development in the Western Diet Mouse Model of Non-
alcoholic Steatohepatitis (NASH). Front. Physiol. *10*.

Du, J., Zhu, M., Bao, H., Li, B., Dong, Y., Xiao, C., Zhang, G.Y., Henter, I., Rudorfer, M., and
Vitiello, B. (2016). The Role of Nutrients in Protecting Mitochondrial Function and
Neurotransmitter Signaling: Implications for the Treatment of Depression, PTSD, and Suicidal
Behaviors. Crit. Rev. Food Sci. Nutr. *56*, 2560–2578.

Estruch, R., Ros, E., Salas-Salvadó, J., Covas, M.-I., Corella, D., Arós, F., Gómez-Gracia, E.,
Ruiz-Gutiérrez, V., Fiol, M., Lapetra, J., et al. (2018). Retraction and Republication: Primary
Prevention of Cardiovascular Disease with a Mediterranean Diet. N Engl J Med 2013;368:1279-
90. N. Engl. J. Med. *378*, 2441–2442.

Farchi, G., Fidanza, F., Mariotti, S., and Menotti, A. (1994). Is diet an independent risk factor for
mortality? 20 year mortality in the Italian rural cohorts of the Seven Countries Study. Eur. J.
Clin. Nutr. *48*, 19–29.

Fledderus, J.O., van Thienen, J.V., Boon, R.A., Dekker, R.J., Rohlena, J., Volger, O.L., Bijnens,
A.-P.J.J., Daemen, M.J.A.P., Kuiper, J., van Berkel, T.J.C., et al. (2007). Prolonged shear stress
and KLF2 suppress constitutive proinflammatory transcription through inhibition of ATF2.
Blood *109*, 4249–4257.

de la Fuente, A. (2010). From "differential expression" to "differential networking" -
identification of dysfunctional regulatory networks in diseases. Trends Genet. TIG *26*, 326–333.

Gaiteri, C., Ding, Y., French, B., Tseng, G.C., and Sibille, E. (2014). Beyond modules and hubs:
the potential of gene coexpression networks for investigating molecular mechanisms of complex
brain disorders: Beyond modules and hubs. Genes Brain Behav. *13*, 13–24.

Giugliano, D., Ceriello, A., and Esposito, K. (2006). The effects of diet on inflammation:
emphasis on the metabolic syndrome. J. Am. Coll. Cardiol. *48*, 677–685.

Goldsworthy, M., Hugill, A., Freeman, H., Horner, E., Shimomura, K., Bogani, D., Pieles, G.,
Mijat, V., Arkell, R., Bhattacharya, S., et al. (2008). Role of the Transcription Factor Sox4 in
Insulin Secretion and Impaired Glucose Tolerance. Diabetes *57*, 2234–2244.

Günthner, R., and Anders, H.-J. (2013). Interferon-Regulatory Factors Determine Macrophage
Phenotype Polarization. Mediators Inflamm. *2013*, 1–8.

Han, H., Cho, J.-W., Lee, S., Yun, A., Kim, H., Bae, D., Yang, S., Kim, C.Y., Lee, M., Kim, E.,
et al. (2018). TRRUST v2: an expanded reference database of human and mouse transcriptional
regulatory interactions. Nucleic Acids Res. *46*, D380–D386.

Hanghøj, K., Moltke, I., Andersen, P.A., Manica, A., and Korneliussen, T.S. (2019). Fast and
accurate relatedness estimation from high-throughput sequencing data in the presence of
inbreeding. GigaScience *8*.

Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y.C., Laslo, P., Cheng, J.X., Murre, C.,
Singh, H., and Glass, C.K. (2010). Simple combinations of lineage-determining transcription
factors prime cis-regulatory elements required for macrophage and B cell identities. Mol. Cell
38, 576–589.

Herlaar, E., and Brown, Z. (1999). p38 MAPK signalling cascades in inflammatory disease. Mol.
Med. Today *5*, 439–447.

Holt, E.M., Steffen, L.M., Moran, A., Basu, S., Steinberger, J., Ross, J.A., Hong, C.-P., and
Sinaiko, A.R. (2009). Fruit and vegetable consumption and its relation to markers of
inflammation and oxidative stress in adolescents. J. Am. Diet. Assoc. *109*, 414–421.

Hu, F.B. (2002). Dietary pattern analysis: a new direction in nutritional epidemiology. Curr.
Opin. Lipidol. *13*, 3–9.

Jacka, F.N., Pasco, J.A., Mykletun, A., Williams, L.J., Hodge, A.M., O'Reilly, S.L., Nicholson,
G.C., Kotowicz, M.A., and Berk, M. (2010). Association of Western and traditional diets with
depression and anxiety in women. Am. J. Psychiatry *167*, 305–311.

Kafatos, A., Verhagen, H., Moschandreas, J., Apostolaki, I., and Van Westerop, J.J. (2000).
Mediterranean diet of Crete: foods and nutrient content. J. Am. Diet. Assoc. *100*, 1487–1493.

Kanai, M., Akiyama, M., Takahashi, A., Matoba, N., Momozawa, Y., Ikeda, M., Iwata, N.,
Ikegawa, S., Hirata, M., Matsuda, K., et al. (2018). Genetic analysis of quantitative traits in the
Japanese population links cell types to complex human diseases. Nat. Genet. *50*, 390–400.

Kaplan, J.R., Manuck, S.B., and Shively, C. (1991). The effects of fat and cholesterol on social
behavior in monkeys.: Psychosom. Med. *53*, 634–642.

Kersey, P.J., Allen, J.E., Allot, A., Barba, M., Boddu, S., Bolt, B.J., Carvalho-Silva, D.,
Christensen, M., Davis, P., Grabmueller, C., et al. (2018). Ensembl Genomes 2018: an integrated
omics infrastructure for non-vertebrate species. Nucleic Acids Res. *46*, D802–D808.

Kim, D., Langmead, B., and Salzberg, S.L. (2015). HISAT: a fast spliced aligner with low
memory requirements. Nat. Methods *12*, 357–360.

Kimmig, L.M., and Karalis, D.G. (2013). Do Omega-3 Polyunsaturated Fatty Acids Prevent
Cardiovascular Disease? A Review of the Randomized Clinical Trials. Lipid Insights *6*,
LPI.S10846.

Kinsella, R.J., Kähäri, A., Haider, S., Zamora, J., Proctor, G., Spudich, G., Almeida-King, J.,
Staines, D., Derwent, P., Kerhornou, A., et al. (2011). Ensembl BioMarts: a hub for data retrieval
across taxonomic space. Database J. Biol. Databases Curation *2011*, bar030.

Layé, S., Nadjar, A., Joffre, C., and Bazinet, R.P. (2018). Anti-Inflammatory Effects of Omega-3
Fatty Acids in the Brain: Physiological Mechanisms and Relevance to Pharmacology.
Pharmacol. Rev. *70*, 12–38.

Lê, S., Josse, J., and Husson, F. (2008). **FactoMineR** : An *R* Package for Multivariate Analysis.
J. Stat. Softw. *25*.

Lea, A., Subramaniam, M., Ko, A., Lehtimäki, T., Raitoharju, E., Kähönen, M., Seppälä, I.,
Mononen, N., Raitakari, O.T., Ala-Korpela, M., et al. (2019). Genetic and environmental
perturbations lead to regulatory decoherence. ELife *8*.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G.,
Durbin, R., and 1000 Genome Project Data Processing Subgroup (2009). The Sequence
Alignment/Map format and SAMtools. Bioinforma. Oxf. Engl. *25*, 2078–2079.

Liao, Y., Smyth, G.K., and Shi, W. (2014). featureCounts: an efficient general purpose program
for assigning sequence reads to genomic features. Bioinforma. Oxf. Engl. *30*, 923–930.

Lopez-Garcia, E., Schulze, M.B., Fung, T.T., Meigs, J.B., Rifai, N., Manson, J.E., and Hu, F.B.
(2004). Major dietary patterns are related to plasma concentrations of markers of inflammation
and endothelial dysfunction. Am. J. Clin. Nutr. *80*, 1029–1035.

Manzel, A., Muller, D.N., Hafler, D.A., Erdman, S.E., Linker, R.A., and Kleinewietfeld, M.
(2014). Role of "Western diet" in inflammatory autoimmune diseases. Curr. Allergy Asthma
Rep. *14*, 404.

Martinez, F.O., and Gordon, S. (2014). The M1 and M2 paradigm of macrophage activation:
time for reassessment. F1000Prime Rep. *6*.

Miyata, Y., Fukuhara, A., Otsuki, M., and Shimomura, I. (2012). Expression of Activating
Transcription Factor 2 in Inflammatory Macrophages in Obese Adipose Tissue. Obesity.

Mosser, D.M., and Edwards, J.P. (2008). Exploring the full spectrum of macrophage activation.
Nat. Rev. Immunol. *8*, 958–969.

Nagpal, R., Shively, C.A., Appt, S.A., Register, T.C., Michalson, K.T., Vitolins, M.Z., and
Yadav, H. (2018). Gut Microbiome Composition in Non-human Primates Consuming a Western
or Mediterranean Diet. Front. Nutr. *5*, 28.

Nahrendorf, M., and Swirski, F.K. (2016). Abandoning M1/M2 for a Network Model of
Macrophage Function. Circ. Res. *119*, 414–417.

Nanri, A., Moore, M.A., and Kono, S. (2007). Impact of C-reactive protein on disease risk and
its relation to dietary factors. Asian Pac. J. Cancer Prev. APJCP *8*, 167–177.

Nettleton, J.A., Steffen, L.M., Mayer-Davis, E.J., Jenny, N.S., Jiang, R., Herrington, D.M., and
Jacobs, D.R. (2006). Dietary patterns are associated with biochemical markers of inflammation
and endothelial activation in the Multi-Ethnic Study of Atherosclerosis (MESA). Am. J. Clin.
Nutr. *83*, 1369–1379.

Neve, B., Fernandez-Zapico, M.E., Ashkenazi-Katalan, V., Dina, C., Hamid, Y.H., Joly, E.,
Vaillant, E., Benmezroua, Y., Durand, E., Bakaher, N., et al. (2005). Role of transcription factor
KLF11 and its diabetes-associated gene variants in pancreatic beta cell function. Proc. Natl.
Acad. Sci. U. S. A. *102*, 4807–4812.

Ohlow, M.J., Sohre, S., Granold, M., Schreckenberger, M., and Moosmann, B. (2017). Why
Have Clinical Trials of Antioxidants to Prevent Neurodegeneration Failed? - A Cellular
Investigation of Novel Phenothiazine-Type Antioxidants Reveals Competing Objectives for
Pharmaceutical Neuroprotection. Pharm. Res. *34*, 378–393.

O'Keefe, J.H., Gheewala, N.M., and O'Keefe, J.O. (2008). Dietary strategies for improving post-
prandial glucose, lipids, inflammation, and cardiovascular health. J. Am. Coll. Cardiol. *51*, 249–
255.

Osler, M., and Schroll, M. (1997). Diet and mortality in a cohort of elderly people in a north
European community. Int. J. Epidemiol. *26*, 155–159.

Pearen, M.A., and Muscat, G.E.O. (2010). Minireview: Nuclear Hormone Receptor 4A
Signaling: Implications for Metabolic Disease. Mol. Endocrinol. *24*, 1891–1903.

Preacher, K.J., and Hayes, A.F. (2004). SPSS and SAS procedures for estimating indirect effects
in simple mediation models. Behav. Res. Methods Instrum. Comput. J. Psychon. Soc. Inc *36*,
717–731.

Raison, C.L., Capuron, L., and Miller, A.H. (2006). Cytokines sing the blues: inflammation and
the pathogenesis of depression. Trends Immunol. *27*, 24–31.

Reimold, A.M., Kim, J., Finberg, R., and Glimcher, L.H. (2001). Decreased immediate
inflammatory gene induction in activating transcription factor-2 mutant mice. Int. Immunol. *13*,
241–248.

Ritchie, M.E., Phipson, B., Wu, D., Hu, Y., Law, C.W., Shi, W., and Smyth, G.K. (2015). limma
powers differential expression analyses for RNA-sequencing and microarray studies. Nucleic
Acids Res. *43*, e47.

Romagnolo, D.F., and Selmin, O.I. (2017). Mediterranean Diet and Prevention of Chronic
Diseases. Nutr. Today *52*, 208–222.

Schmidl, C., Renner, K., Peter, K., Eder, R., Lassmann, T., Balwierz, P.J., Itoh, M., Nagao-Sato,
S., Kawaji, H., Carninci, P., et al. (2014). Transcription and enhancer profiling in human
monocyte subsets. Blood *123*, e90-99.

Schuler, A., Schwieger, M., Engelmann, A., Weber, K., Horn, S., Muller, U., Arnold, M.A.,
Olson, E.N., and Stocking, C. (2008). The MADS transcription factor Mef2c is a pivotal
modulator of myeloid cell fate. Blood *111*, 4532–4541.

Scott, E., Simon, M., Anastasi, J., and Singh, H. (1994). Requirement of transcription factor
PU.1 in the development of multiple hematopoietic lineages. Science *265*, 1573–1577.

Seltmann, M.W., Helle, S., Adams, M.J., Mar, K.U., and Lahdenperä, M. (2018). Evaluating the
personality structure of semi-captive Asian elephants living in their natural habitat. R. Soc. Open
Sci. *5*, 172026.

Sharon, G., Sampson, T.R., Geschwind, D.H., and Mazmanian, S.K. (2016). The Central
Nervous System and the Gut Microbiome. Cell *167*, 915–932.

Shively, C.A. (1998). Social subordination stress, behavior, and central monoaminergic function
in female cynomolgus monkeys. Biol. Psychiatry *44*, 882–891.

Shively, C.A., Register, T.C., and Clarkson, T.B. (2009). Social Stress, Visceral Obesity, and
Coronary Artery Atherosclerosis in Female Primates. Obesity *17*, 1513–1520.

Shively, C.A., Appt, S.E., Vitolins, M.Z., Uberseder, B., Michalson, K.T., Silverstein-Metzler,
M.G., and Register, T.C. (2019). Mediterranean versus Western Diet Effects on Caloric Intake,
Obesity, Metabolism, and Hepatosteatosis in Nonhuman Primates. Obes. Silver Spring Md *27*,
777–784.

Smil, V. (1989). Coronary Heart Disease, Diet, and Western Mortality. Popul. Dev. Rev. *15*,
399.

Smyth, S., and Heron, A. (2006). Diabetes and obesity: the twin epidemics. Nat. Med. *12*, 75–80.

Snyder-Mackler, N., and Lea, A.J. (2018). Functional genomic insights into the environmental
determinants of mammalian fitness. Curr. Opin. Genet. Dev. *53*, 105–112.

Snyder-Mackler, N., Sanz, J., Kohn, J.N., Brinkworth, J.F., Morrow, S., Shaver, A.O., Grenier,
J.-C., Pique-Regi, R., Johnson, Z.P., Wilson, M.E., et al. (2016). Social status alters immune
regulation and response to infection in macaques. Science *354*, 1041–1045.

Steinhubl, S.R. (2008). Why Have Antioxidants Failed in Clinical Trials? Am. J. Cardiol. *101*,
S14–S19.

Stice, E., and Durant, S. (2014). Elevated objectively measured but not self-reported energy
intake predicts future weight gain in adolescents. Appetite *81*, 84–88.

Suchanek, P., Poledne, R., and Hubacek, J.A. (2011). Dietary intake reports fidelity--fact or
fiction? Neuro Endocrinol. Lett. *32 Suppl 2*, 29–31.

Tachmazidou, I., Süveges, D., Min, J.L., Ritchie, G.R.S., Steinberg, J., Walter, K., Iotchkova, V.,
Schwartzentruber, J., Huang, J., Memari, Y., et al. (2017). Whole-Genome Sequencing Coupled
to Imputation Discovers Genetic Signals for Anthropometric Traits. Am. J. Hum. Genet. *100*,
865–884.

Thompson, J.R., Gustafsson, H.C., DeCapo, M., Takahashi, D.L., Bagley, J.L., Dean, T.A.,
Kievit, P., Fair, D.A., and Sullivan, E.L. (2018). Maternal Diet, Metabolic State, and
Inflammatory Response Exert Unique and Long-Lasting Influences on Offspring Behavior in
Non-Human Primates. Front. Endocrinol. *9*.

Trichopoulou, A., Kouris-Blazos, A., Wahlqvist, M.L., Gnardellis, C., Lagiou, P.,
Polychronopoulos, E., Vassilakou, T., Lipworth, L., and Trichopoulos, D. (1995). Diet and
overall survival in elderly people. BMJ *311*, 1457–1460.

Tung, J., and Gilad, Y. (2013). Social environmental effects on gene regulation. Cell. Mol. Life
Sci. CMLS *70*, 4323–4339.

Warden, C.H., and Fisler, J.S. (2008). Comparisons of Diets Used in Animal Models of High-Fat
Feeding. Cell Metab. *7*, 277.

Whelton, P.K., Appel, L., Charleston, J., Dalcin, A.T., Ewart, C., Fried, L., Kaidy, D., Klag,
M.J., Kumanyika, S., Steffen, L., et al. (1992). The Effects of Nonpharmacologic Interventions
on Blood Pressure of Persons With High Normal Levels: Results of the Trials of Hypertension
Prevention, Phase I. JAMA *267*, 1213–1220.

Yang, H., Graham, L.C., Reagan, A.M., Grabowska, W.A., Schott, W.H., and Howell, G.R.
(2019). Transcriptome profiling of brain myeloid cells revealed activation of Itgal, Trem1, and
Spp1 in western diet-induced obesity. J. Neuroinflammation *16*.

Zhang, D.E., Hetherington, C.J., Tan, S., Dziennis, S.E., Gonzalez, D.A., Chen, H.M., and
Tenen, D.G. (1994). Sp1 is a critical factor for the monocytic specific expression of human
CD14. J. Biol. Chem. *269*, 11425–11434.