Loss of function of the mitochondrial peptidase PITRM1 induces

proteotoxic stress and Alzheimer's disease-like pathology in human

cerebral organoids

Dina Ivanyuk[1,2#], María José Pérez[1,2#], Vasiliki Panagiotakopoulou[1,2], Gabriele Di

Napoli[1,2], Dario Brunetti[3], Rawaa Al-Shaana[1,4], Stephan A. Kaeser[1,4], Mathias

Jucker[1,4], Massimo Zeviani[3], Carlo Viscomi[3], Michela Deleidi[1,2*].

1, German Center for Neurodegenerative Diseases (DZNE), Tübingen, Germany.

2, Department of Neurodegenerative Diseases, Hertie Institute for Clinical Brain Research,

University of Tübingen, Tübingen, Germany.

3, MRC-Mitochondrial Biology Unit, Cambridge CB2 0XY, UK.

4, Department of Cellular Neurology, Hertie Institute for Clinical Brain Research, University of

Tübingen, Tübingen, Germany.

[#], These authors contributed equally to the work

Manuscript correspondence:

Michela Deleidi, MD, PhD

German Center for Neurodegenerative Diseases (DZNE) Tübingen within the Helmholtz

Association, Department of Neurodegenerative Diseases, University of Tübingen

Otfried-Müller. Str 23

72076 Tübingen-Germany

Tel.: +49 7071 9254200

Fax: +49 7071 9254074

Abstract

Mutations in pitrilysin metallopeptidase 1 (*PITRM1*), a mitochondrial protease involved in mitochondrial precursor processing and degradation, result in a slow-progressive syndrome, characterized by cerebellar ataxia, psychotic episodes and obsessive behavior as well as cognitive decline. To investigate the pathogenetic mechanisms of mitochondrial presequence processing, we employed cortical neurons and cerebral organoids generated from PITRM1 knockout human induced pluripotent stem cells (iPSCs). PITRM1 deficiency strongly induced mitochondrial unfolded protein response (UPR^{mt}) and enhanced mitochondrial clearance in iPSC-derived neurons. Furthermore, we observed increased levels of amyloid precursor protein and amyloid β in PITRM1 knockout neurons. However, neither cell death nor protein aggregates were observed in 2D iPSC-derived cortical neuronal cultures. On the contrary, cerebral organoids generated from PITRM1 knockout iPSCs spontaneously developed over time pathological features of Alzheimer's disease (AD), including accumulation of protein aggregates, tau pathology, and neuronal cell death. Importantly, we provide evidence for a protective role of UPR^{mt} and mitochondrial clearance against impaired mitochondrial presequence processing and proteotoxic stress. In summary, we propose a novel concept of PITRM1-linked neurological syndrome whereby defects of mitochondrial presequence processing induce an early activation of UPR^{mt} that, in turn, modulates cytosolic quality control pathways. Thus our work supports a mechanistic link between mitochondrial function and common neurodegenerative proteinopathies.

Introduction

Mitochondrial dysfunction has been described as a common hallmark of neurological diseases [1]. However, mitochondria are often considered to be a secondary target, rather than the actual disease driver in these conditions. We have recently described three independent families carrying missense loss of function mutations in *pitrilysin metallopeptidase 1* (PITRM1), resulting in an age-dependent, progressive, neurological syndrome [2, 3]. Patients suffer from progressive cerebellar dysfunction leading to cerebellar atrophy, and psychiatric manifestations including obsessive behavior, anger attacks, and psychosis [2, 3]. Interestingly, some of these patients showed a deterioration of their cognitive functions with a slow progression till their late sixties [2, 3]. Human PITRM1, also known as presequence peptidase (hPreP), is a nuclear-encoded mitochondrial gene, expressed in a number of tissues, including muscles and different brain regions, e.g. cortex, hippocampus, cerebellum, and tectum [4]. PITRM1 was initially identified in *Arabidopsis thaliana* as a protease that degrades targeting peptides in mitochondria and chloroplasts [5]. Human PITRM1 is a mitochondrial matrix enzyme that digests the mitochondrial-targeting sequences (MTS) of proteins imported across the inner mitochondrial membrane, after their cleavage from protein precursors by the mitochondrial matrix presequence peptidase (MPP). When the MTS are not properly degraded, they accumulate within the mitochondrial matrix causing dissipation of the mitochondrial membrane potential and mitochondrial dysfunction [6, 7]. The incomplete processing of mitochondrial preproteins leads to their destabilization, resulting in alterations of mitochondrial proteostasis [8]. *In vitro* studies using recombinant PITRM1 have shown that, besides MTS, the enzyme is also involved in the degradation of short, unstructured peptides and amyloid beta (Aβ) peptides [8-11]. Interestingly, Aβ peptides inhibit the activity of CYM1, the PITRM1 orthologue in yeast, leading to accumulation of precursor proteins [8]. Experimental work in mouse models supports a causal role for PITRM1 in neurodegenerative dementia, whereby loss of PITRM1 function leads to a progressive, neurodegenerative phenotype characterized by hindlimb clasping, impairment in motor coordination, and basal-ganglia related movement control [2]. Interestingly, PITRM1 deficient mice show an age-dependent accumulation of amyloid precursor protein (APP) and Aβ deposits within the brain [2], suggesting a link between defects of mitochondrial proteostasis and adult-onset neurodegenerative dementia. However, due to the embryonic lethality observed in complete PITRM1 knockout mice, the exact role of PITRM1 in brain homeostasis and disease could not be studied [2]. Since PITRM1 is involved in the degradation of MTS, as well as Aβ peptides [8], the pathomechanisms of neurodegeneration linked to loss of PITRM1

function could in principle be due to either accumulation of Aβ peptides in mitochondria, MTS-driven toxicity, or a combination of both. Identifying the mechanisms that lead to neurodegeneration in primary mitochondrial diseases characterized by defects of mitochondrial proteostasis, such as PITRM1-linked neurological syndrome, may help elucidate the long-debated, still unresolved, involvement of altered mitochondrial function in neurodegenerative dementia. To examine PITRM1-related pathogenetic mechanisms, we generated PITRM1-knockout human induced pluripotent stem cells (iPSCs) and examined the role of mitochondrial function and proteostasis using 2D neuronal and 3D brain-organoid model systems.

Methods

Generation of PITRM1 knockout human iPSCs

Control iPSCs (from a female non affected control, 80 years) used in this study were previously generated and characterized [12]. All cells used in the study were derived from patients who signed an informed consent approved by The Ethics Committee of the Medical Faculty and the University Hospital Tübingen. iPSCs were kept in culture in hESC medium [12]. SgRNAs targeting exon 3 and 4 of PITRM1 gene were designed using CRISPR Guide Design Tools (former www.crispr.mit.edu) and purchased from Metabion International AG (exon 3 top AGGAGCCAGGTATTTACACC, exon 3 bottom GGTGTAAATACCTGGCTCCT, exon 4 top TTGAGCATACCGTCCTTTGT, exon 4 bottom ACAAAGGACGGTATGCTCAA). SgRNAs were cloned into the pSpCas9(BB)-2A-Puro plasmid containing the sgRNA scaffold and puromycine-resistance under the U6 promoter (Addgene plasmid #48139). Colonies that successfully integrated sgRNA into backbone plasmid were screened and confirmed by Sanger sequencing using U6 promoter region primer. SgRNA/Cas9 plasmid was delivered into cells using Nucleofector Amaxa system (Lonza Biosciences). In brief, iPS cells were dissociated with Accutase (Sigma-Aldrich) and 10^6 iPSCs were nucleofected with 6μg of each sgRNA plasmid. Cells were then replated on MEF cells in hESC medium, without P/S, supplemented with 10μM ROCK Inhibitor Y-27632 2HCl. After recovery, cells were replated at density 500 cells/cm^2 for single cell subcloning. After recovery, iPSCs were clonally expanded and the genomic deletion was assessed by PCR and Sanger sequencing (exon 3 Fw TTCAGGCAGAAAAGCCAGTT, exon 4 Rv ACTGAATTCCAGTGGGTGTGC). The screening of possible off-target effects was performed using CRISPR Design Tools. Sequencing primers for off-target effects: (5'-3'): PPIL2 NM_148175 Fw CCTCATGCCCTGCTTGACTC, Rv CAGGGAGCACTGTCCCAATTT; NR1D1

NM_021724 Fw CAAACGAGCACACACCACAG, Rv GCTGCCCCCTTGTACAGAAT;

KDEL2   NM_001100603   Fw   TTGGTGGTGGTTATGCCTCA,   Rv

ACCACCAGAAACTCCACTCG; FAM120A NM_014612   Fw

TCCTGCGGTTCTTGTCCTCTA,   Rv   GCATGAATGTGTCTTCTCTGGC;   TTLL2

NM_031949 Fw GTGGGAGGCTGTGTGGTATT, Rv TCAAGTCCCTACCTGTGCCA;

SUCO   NM_014283   Fw   AATCTGGTACTATTCCGATAGCCAA,   Rv

CCATTCAAACAGGACACTGCTG.

Cortical neuronal differentiation

For the induction of cortical neurons, we used an embryoid body (EB)-based

differentiation protocol, with minor modifications [13]. iPSC colonies were manually

picked and grown for 4 d in EB media (20% KO serum replacement in DMEM/F12

medium, 1% NEAA, 1% P/S). On day 5, EBs were plated onto Matrigel-coated

(Corning) plate and grown for 4 more days in N2B27 media (DMEM/F12, 1X N2, 1X

B27-RA, 1% NEAA, 1% P/S, 20 ng/mL bFGF). For the first 8 differentiation days,

the cells were grown in the presence of 10 μM SB431542 (Ascent Scientific) and

2.5 μM dorsomorphin (Sigma Aldrich). After 8 d, neural rosettes were lifted with

Dispase and replated onto matrigel-coated plate and grown in N2B27 media.

Secondary or tertiary rosettes were manually dissected to purify neural progenitor

cells (NPCs). For cortical differentiation, NPCs were dissociated with Accutase and

seeded at a density of 1,000 cells per mm2 on Matrigel-coated plates in neuronal

differentiation medium consisting of DMEM/F12, 1X Glutamax, 1X N2, 1X B27-RA,

20 ng/mL BDNF (Peprotech), 20 ng/mL GDNF, 1 mM dibutyryl-cyclic AMP

(Sigma), 200 nM ascorbic acid (Sigma). Medium was replaced every other day.

Cerebral organoids culture and immunohistochemistry

Cerebral organoids were generated and maintained using the protocol described by

Lancaster et al. [14]. Where indicated, cerebral organoids were treated with 500nM

ISRIB or 500μM NMN (both from Sigma Aldrich) daily, from DIV 20 to DIV 50 or from

DIV45 to DIV50 respectively. For immunostaining, organoids were washed with PBS,

fixed in 4% PFA for 15 min and then equilibrated in 30% sucrose in PBS overnight at

4°C. Next day, organoids were embedded in blocks with a mixture of 10%

sucrose/7.5% gelatin, snap frozen, and kept at –80°C until cryosections were

prepared using Leica CM 1900 instrument with 20 μm thickness. Sections were then

permeabilized and blocked with 10% normal goat serum in PBS in 0.5% Triton X-100.

Primary antibody incubations were performed at 4°C overnight, followed by three 10

min washes in PBS, and staining with AlexaFluor secondary antibodies (Invitrogen, 1:1000) at room temperature for 1h. Primary antibodies included rat anti-CTIP2 (1:500, Abcam ab18465), rabbit anti-TBR1 (1:500, Abcam ab31940), mouse anti-β-III-tubulin (1:1000, Sigma-Aldrich T8328), rabbit anti-β-III-tubulin (1:1000, Biolegend, Previously Covance # PRB-435P), chicken anti-MAP2 (1:3000, Biolegend PCK-554P), mouse anti phospho-tau (AT8 1:500, Thermofisher #MN1020), mouse anti-APP (1:500, Santa Cruz Biotechnology sc53822), mouse anti-Ubiquitin (1:100, Merck MAB1510), mouse anti-Caspase 3 (1:500, Cell Signaling Technology 9664T). For thioflavin T immunostaining, sections were stained with 10µM thioflavin T (Sigma-Aldrich T3516) for 15 min at room temperature and washed 3 times in PBS. Cell nuclei were stained with DAPI and slides were mounted with DAKO mounting medium (Agilent S302380-2). Images were acquired using Leica TCS SP8 confocal microscope (Leica Microsystems). For image analysis, mean of fluorescence intensity per image was calculated with Image J. For each condition, 10-15 images were acquired from at least 5 organoids from three independent experiments (cultures).

Measurement of mitochondrial membrane potential

For measurement of mitochondrial membrane potential, neurons were split at a concentration of 2x10^5 cells/well on 96-wells pre-coated with Matrigel. Cells were washed once with HBSS (Invitrogen) following incubation with 200 nM Tetramethylrhodamine Methylester Perchlorat (TMRM) (Invitrogen) in HBSS with 1% BSA for 30 min at 37°C. Cells were then washed twice with HBSS, detached using Accumax (Invitrogen) and kept in HBSS with 1% BSA (Sigma) for FACS analysis using MaxQuant Analyzer 10 (Miltenyi Biotec). Analysis was performed using the MACSQuantify 2.6 software (Miltenyi Biotec).

Measurement of mitochondrial Reactive Oxygen Species (mtROS)

For measurement of mtROS production of iPSC-derived neurons by flow cytometry, cells were pre-incubated with N2 medium for 48 hrs. Then, 1×10^6 cells were washed once with HBSS, incubated with 5µM of the superoxide indicator MitoSOX Red (Invitrogen) for 30 min at 37°C, washed twice with HBSS, dissociated using Trypsin (Invitrogen) and resuspended in 200µl of HBSS and 1% BSA. Cytofluorimetric analysis was performed using MACSQuant Analyzer 10 (Miltenyi Biotec).

Seahorse XF^e96 Metabolic Flux Analysis

Oxygen consumption rate (OCR) was analyzed using an XFp Extracellular Flux

Analyzer (Agilent). iPSC-derived neurons were plated on XFp microplates (Agilent) at

a density of 70,000 per well and grown in N2 medium for 48 hrs before the

experiment. Measurement of neuronal oxidative consumption rate was performed in

freshly prepared medium consisting of phenol-free DMEM,1 mM Natrium Pyruvate,

2mM Glutamine and 10mM glucose with pH adjusted to 7,4. Mitochondrial function

was evaluated after subsequent injection of 10μM oligomycin, 10 μM carbonyl

cyanide p-trifluoromethoxyphenylhydrazone (CCCP) and 2μM Antimicyn A / 1μM

Rotenone (all Sigma-Aldrich). For each condition 3 measurements, lasting 5 minutes

each, were performed: after each injection, OCR was measured for 2 minutes, the

medium was mixed for 2 min, let rest for 1 min, and then respiration was measured

again. After measurement, values were normalized to cell number by counting DAPI

stained nuclei using a high-content cell analyzer (BD Bioscience, Pathway 855).

Quantitative RT-PCR

mRNA was isolated using an RNA isolation kit (Qiagen). Following the reverse

transcription reaction using the QuantiTect Reverse Transcription kit, quantitative

PCR reaction was performed using SYBR GREEN (all Qiagen) and monitored with a

Viia7 Real time PCR system (Applied Biosystems). The expression level of each

gene was normalized to the housekeeping gene ribosomal protein large P0 (Rplp0).

Fold-changes in gene expression were calculated using the 2^{-DDCT} method, based on

biological reference samples and housekeeping genes for normalization.

qPCR primers (5'-3')

GAPDH	Fw	AGGGGAGATTCAGTCTGG
	Rv	CGACCACTTTGTCAAGCT
HSP60	Fw	TGACCCAACAAAGGTTGTGA
	Rv	CATACCACCTCCCATTCCAC
LONP1	Fw	CCCGCGCTTTATCAAGATT
	Rv	AGAAAGACGCCGACATAAGG
ATF4	Fw	GTCCCTCCAACAACAGCAAG
	Rv	CTATACCCAACAGGGCATCC
DDIT3	Fw	AGCCAAAATCAGAGCTGGAA
	Rv	TGGATCAGTCTGGAAAAGCA
ERO1A	Fw	AGCGGCACAGAGGTGCT
	Rv	TGTAGTCTTGGGAAAAGCCTG
RPLP0	Fw	CCTCATATCCGGGGGAATGTG
	Rv	GCAGCAGCTGGCACCTTATTG
HSPA9	Fw	GGAAGCTGCTGAAAAGGCTA
	Rv	CTTGGGTCCAGAAGAATCCA
CLPP	Fw	CTCTTCCTGCAATCCGAGAG
	Rv	GGATGTACTGCATCGTGTCG

16S (mtDNA)	Fw	GCCTTC CCCCGTAAATGATA
	Rv	TTATGCGATTAC CGGGCTCT
PITRM1	Fw	ATCTGTTCCCGAGCTGTTCC
	Rv	GAAAGGCTCTCTGCACGGAT

Mitochondrial isolation

247 The mitochondrial isolation was performed using Qproteome® Mitochondria Isolation

248 Kit (Qiagen) according to manufacturer's protocol.

Western blot

Proteins were extracted using Tris-buffered Saline (TBS) with 0.5% NP40 protein

extraction buffer containing protease and phosphatase inhibitors (Roche), on ice,

following centrifugation at 14.000 rpm and 4°C for 15min. The protein concentration

of the supernatant was determined by BCA (Pierce). In total, 15–30 µg of the

protein lysate was loaded on polyacrylamide gel (density ranged from 7.5% to 15%,

depending on respective protein molecular weight) and transferred on a PVDF

membrane (Millipore). Blots were blocked with 5% milk powder or 5% BSA in TBS +

0.1% Tween-20 (TBST) and incubated with primary antibodies in milk or BSA

blocking solution overnight at 4 °C. This step was followed by incubation with

corresponding HRP-conjugated secondary antibodies (Sigma Aldrich) for 1 h at room

temperature. Visualization of proteins was done by using Amersham ECL Western

Blotting Detection Reagent and Amersham Hyperfilm (both GE Healthcare).

Densitometric analysis of proteins was performed by ImageJ software. Primary

antibodies included: rabbit anti-LC3B (1:500, Cell Signaling Technologies #2775);

mouse anti-APP 6E10 (Abeta 1-16) (1:1000, Biolegend #803004); rabbit anti-

PITRM1 (1.1000; Atlas Antibodies #HPA006753); rabbit anti-Frataxin (1:1000,

Abcam #ab175402); mouse anti-Total OXPHOS human cocktail (1:1000, Abcam

#ab110411); mouse anti-ubiquitin (1:5000, Millipore #MAB1510); mouse anti-HSPA9

(1:3000, Santa Cruz Biotechnologies #sc-133137); mouse anti-HSP60 (1:3000,

Santa Cruz Biotechnologies #sc-271215); rabbit anti-LONP1 (1:3000, Proteintech

#15440-1-AP); mouse anti-tau (1:1000, HT7, Thermofisher #MN1000); mouse anti

phospho-tau PHF-6 (Thr231) (1:1000, Thermofisher #35-5200); mouse anti-β-Actin

(1:20000, Sigma Aldrich #A5441); mouse anti-VDAC1 (1:1000, Santa Cruz

Biotechnologies #sc-390996).

Autophagy studies

277 Where indicated, cells were treated with NH_4Cl (20 mM) and leupeptin (200 µM)

278 (EMD, Millipore) for 4 h. LC3-II and LC-I levels were quantified by densitometry and

279   normalized to β-actin. LC3 flux was quantified by dividing the levels of LC3-II after

280   treatment with lysosomal inhibitor for 4h, by the levels of LC3-II without treatment.

Immunofluorescence

Cells were fixed in 4% paraformaldehyde (PFA) in PBS (w/v) for 10 min, rinsed with

PBS and blocked by 10% normal goat or donkey serum (NGS/NDS) in PBST (PBS +

0.1% TritonX-100) for 60 min. Cells were then incubated with primary antibodies in

10% NGS/NDS in PBST over night at 4°C following 1h incubation at 24°C with

appropriate Alexa488/568 coupled secondary antibodies (1:1000, Invitrogen). Cell

nuclei were stained with DAPI and final images were acquired using Leica TCS SP8

confocal microscope (Leica Microsystems) and analyzed using Fiji software. Primary

antibodies included: rabbit anti-LC3B (1:500, Cell Signaling Technologies #2775);

mouse anti-CTIP2 (1:500, Abcam ab18465), rabbit anti-TBR1 (1:500, Abcam

ab31940), mouse anti-β-III-tubulin (1:1000, Sigma-Aldrich T8328-25UL), and anti-

Caspase 3 (1:500, Cell Signaling Technology 9664T). LC3 particle number in β-III-

tubulin positive cells was quantified with the "analyse particles" plug-in in ImageJ

(NIH). Caspase 3 was quantified with mean fluorescence intensity in Image J (NIH).

Quantification was carried out on, at least, 50 cells per condition, from three

independent experiments.

LDH assay

300   The LDH assay (Promega) was performed as per the manufacturer's instructions.

Amyloid-beta species measurement

For amyloid-beta species measurement in iPSC-cortical neurons, 10^6 cells were

plated into a one well of 12-well plate and cell supernatant was collected after 5 days,

snap frozen and stored at -80 °C until analysis. Cell pellet was collected and lysed for

protein concentration determination and used for value normalization. Conditioned

medium from at least 3 technical replicates was collected in each experiment. For

amyloid-beta species measurement in cerebral organoids, individual organoids were

re-plated into low attachment 96-well plates on an orbital shaker. Supernatant was

collected after 5 days, snap frozen in liquid nitrogen and stored at -80 °C until

analysis. After collection of supernatant, individual organoids were lysed and protein

concentration measured with BCA for value normalization. The concentrations of

Aβ40 (Aβx–40) and Aβ42 (Aβx–42) in the samples were measured on a Sector

Imager 6000 using an electrochemiluminescence-based immunoassay, V-PLEX Ab

Peptide Panel 1 (6E10) Kits (Meso Scale Discovery, Gaithersburg, MD, USA)

according to manufacturer's instructions. All samples were thawed on ice and diluted
1:2 in buffer (Diluent 35, Meso Scale Discovery) before incubation. Neural
differentiation medium without B27/N2 supplements was used as negative control for
each measurement. Human CSF samples were used as internal references on each
plate. At least 8 organoids were used for each measurement. Every sample was
tested in duplicate (and excluded if the coefficient of variance (CV) was above 20%).
Data analysis was run with the MSD DISCOVERY WORKBENCH software version
2.0.
324
Elisa
Individual cerebral organoids were homogenized in ice-cold RIPA buffer containing
protease and phosphatase inhibitors (Roche), following centrifugation at 14.000 rpm
and 4°C for 10min. The protein concentration of the supernatant was determined by
BCA (Pierce). Total and phospho-tau levels were assessed, in equal protein amounts,
using ELISA assays (both Invitrogen, KHB0041 and KHO0631, respectively)
according to manufacturer's instructions.
332
333 **Statistical analysis**
334 The Statistical Package GraphPad Prism version 7 and 8.3 (GraphPad Software)
335 was used to analyze the data. Statistical significance was evaluated using two-tailed
336 Student's t-test. Data are expressed as mean + S.E.M. or S.D. as indicated.
337
Results
Mitochondrial dysfunction in PITRM1 knockout iPSC-derived neurons
In order to overcome the limitation of the embryonic lethality previously observed in
PITRM1 knockout mice [2] and examine the mechanistic link between PITRM1
deficiency and neurodegeneration in a model that more closely resembles human
disease, we generated PITRM1 knockout (PITRM1$^{-/-}$) human iPSCs using
CRISPR/Cas9 endonuclease-mediated gene editing. Appropriate sgRNAs targeting
exon 3 and exon 4 were designed to introduce a frameshift deletion resulting in the
complete knockout of PITRM1 protein (Supplementary Fig. 1A). Several
homozygous clones were generated, and two fully characterized clones were
selected and used for further analysis (Supplementary Fig. 1B-D). To address the
impact of PITRM1 on the function of human neurons, we differentiated PITRM1$^{+/+}$
and PITRM1$^{-/-}$ iPSCs into neuronal cultures that were enriched for cortical neurons
and we assessed neuronal cultures at 35 and 65 days *in vitro*. Both PITRM1$^{+/+}$ and

PITRM1[-/-] iPSCs efficiently generated cortical neurons (Fig. 1A). Western blot

analysis confirmed the complete absence of PITRM1 protein in PITRM1[-/-] iPSC-

derived neurons (Fig. 1B). No overt cell death was observed in neuronal cultures, as

assessed by LDH assay at 35 and 65 days *in vitro* (Supplementary Fig. 1E) and

cleaved Caspase 3 staining (data not shown). PITRM1 deficiency leads to the

accumulation of non-degraded MTS sequences that in turn impair the processing of

presequence proteins by the peptidase MPP [8]. Since MPP is also involved in the

maturation of the human frataxin precursor [15], we examined frataxin maturation by

immunoblotting. A decreased ratio of processed, mature to immature frataxin was

detected in PITRM1[-/-] iPSC-derived neural precursor cells (NPCs) and neurons

(Supplementary Fig. 1F, G), indicating the impaired function of MPP and defects of

mitochondrial of mitochondrial presequence processing. Since MTS peptides can

bind to the membrane and perturb the mitochondrial electrochemical gradient, we

evaluated the effect of PITRM1 deficiency on mitochondrial membrane potential and

respiratory oxidative activity. Mitochondrial membrane potential was significantly

reduced in PITRM1[-/-] neurons compared to isogenic PITRM1[+/+] neurons (Fig. 1C).

However, mitochondrial reactive oxygen species (mtROS) were not significantly

altered in PITRM1[-/-] neurons (Fig. 1D). Similarly, no significant difference in the

respiratory oxidative activity was observed between PITRM1[+/+] and PITRM1[-/-]

neurons (Fig. 1E). Western blot analysis revealed a significant increase in the level of

Complex II respiratory chain complex subunits in PITRM1[-/-] neurons (Fig. 1F, G).

However, no significant difference for the levels of all other respiratory chain complex

subunits was detected in PITRM1[-/-] neurons (Fig. 1F, G).

PITRM1[-/-] iPSC-derived cortical neurons show induction of mitochondrial

stress response and enhanced mitophagy

Mitochondrial stress response has been identified as a common signature in several

neurodegenerative as well primary mitochondrial diseases [16-19]. Thus, we examined

the expression levels of genes involved in mitochondrial unfolded protein response

(UPR[mt]) and, more generally, in the mitochondrial integrated stress response

pathway (mtISR). PITRM1[-/-] iPSC-derived cortical neurons exhibited a significant

induction of UPR[mt]/mtISR transcripts (*ATF4, DDIT3, HSP60, HSPA9, ERO1*) (Fig.

2A). Moreover, gene expression of the mitochondrial proteases, *LONP1 and CLPP*,

was significantly upregulated in PITRM1[-/-] neurons compared to controls (Fig. 2A). In

line with gene expression data, we found an increase in protein expression of the

chaperones HSPA9 and HSP60, and the mitochondrial protease LONP1 (Fig. 2B).

These data indicate that the accumulation of MTS, due to PITRM1 deficiency, leads

to a strong upregulation of the ISR pathway. Since, the ISR has been shown to

activate autophagy [20, 21], we assessed the autophagosome content by

immunostaining for endogenous light chain type 3 protein (LC3), a marker of

autophagosomes. Our analysis revealed a significant decrease of LC3-positive

vesicles in PITRM1$^{-/-}$ neurons compared to isogenic PITRM1$^{+/+}$ neurons (Fig. 2C, D).

These results were confirmed by Western blot, showing decreased basal levels of

LC3-II in PITRM1$^{-/-}$ neurons (Fig. 2E). However, inhibition of lysosomal degradation

by leupeptin and ammonium chloride revealed that the autophagic flux was

significantly increased in PITRM1$^{-/-}$ neurons, thus confirming autophagy activation

(Fig. 2E, F). To assess whether the observed increase in autophagic flux leads to an

enhanced turnover rate of mitochondria by autophagy, we purified mitochondria from

PITRM1$^{+/+}$ and PITRM1$^{-/-}$ iPSC-derived cortical neurons. Immunoblotting of purified

mitochondria showed increased ubiquitination of PITRM1$^{-/-}$ neurons compared to

PITRM1$^{+/+}$ neurons (Fig. 2G, H), suggesting their targeting to lysosomes and

increased mitochondrial clearance [22]. In line with these results, mitochondrial content

was lower in PITRM1$^{-/-}$ neurons, as shown by a reduced mitochondrial to nuclear

DNA ratio (Fig. 2I). Taken together, these results suggest that loss of PITRM1

function enhances mitochondrial clearance.

408   PITRM1$^{-/-}$ iPSC-derived neurons show accumulation of APP and increase in

409   extracellular Aβ peptides levels

To examine the impact of PITRM1 activity on Aβ pathology, we first assessed the

levels of APP by Western blot. APP protein levels were found significantly increased

in PITRM1$^{-/-}$ neurons compared to PITRM1$^{+/+}$ neurons (Fig. 3A). Next, supernatant

from PITRM1$^{+/+}$ and PITRM1$^{-/-}$ cortical neurons was analyzed using the Meso Scale

Discovery immunoassay for human Aβ peptides. PITRM1$^{-/-}$ neurons showed

significantly higher levels of Aβ40 and Aβ42 peptides as well as an increased

Aβ42/Aβ40 ratio compared to control samples (Fig. 3B). Remarkably, Aβ peptides

were not detected in mitochondrial extracts from PITRM1 deficient neurons. Next, we

explored the mechanisms that link loss of PITRM1 function with alteration of APP

metabolism. Since UPR^{mt} may have an impact on ubiquitin-dependent protein

turnover [23], we examined the levels of ubiquitinated proteins by Western blot. We

observed that PITRM1$^{-/-}$ display increased levels of ubiquitinated proteins (Fig. 3C, D),

suggesting defects in cellular proteostasis.

424   PITRM1$^{-/-}$ cerebral organoids exhibit main features of AD pathology and

425   induction of mitochondrial stress response

Despite APP accumulation and increased Aβ42/Aβ40 ratio, 2D iPSC-derived
neuronal cultures did not show Aβ aggregates nor tau pathology or overt cell death.
To further investigate the mechanisms of PITRM1 neurotoxicity in a model that better
resembles the human disease, we developed cerebral organoids from PITRM1$^{+/+}$ and
PITRM1$^{-/-}$ iPSCs and cultured them over a broad time range (1-6 months). Cerebral
organoids derived from PITRM1$^{+/+}$ and PITRM1$^{-/-}$ iPSCs displayed similar sizes and
cortical layering characteristics (Fig. 4A and Supplementary Fig. 2A). Next, we
examined whether PITRM1$^{-/-}$ cerebral organoids develop AD-like neurodegenerative
features. Western blotting reveled increased APP levels and tau
hyperphosphorylation in 2-months old PITRM1$^{-/-}$ cerebral organoids (Fig. 4B-E).
Furthermore, immunoassay measurements showed a higher Aβ40, Aβ42 and
Aβ42/Aβ40 ratio in PITRM1$^{-/-}$ cerebral organoids compared to controls
(Supplementary Fig. 2B). Similarly, immunofluorescence staining showed increased
APP and phospho-tau levels in PITRM1$^{-/-}$ organoids compared to PITRM1$^{-/-}$ controls
starting at 2 months (Fig. 4F, G, I). No further increase of APP levels or tau
hyperphosphorylation were observed at later time points (Supplementary Figure 2C).
Next, we stained 1-, 2-, and 6-month old PITRM1$^{+/+}$ and PITRM1$^{-/-}$ cerebral organoids,
for cleaved caspase-3. The number of cleaved caspase-3 positive cells in the
neuroepithelial layers was higher in PITRM1$^{-/-}$ cerebral organoids than controls,
starting at 2 months (Fig. 4H, I and Supplementary Fig. 2D), suggesting a higher
extent of cell death. No further increase of cell death was observed at later time
points (Supplementary Figure 2D). To analyze ubiquitin-dependent protein turnover,
an immunostaining against ubiquitinated proteins was performed. PITRM1$^{-/-}$
organoids display increased levels of ubiquitinated proteins (Fig 4J, K). Thioflavin T
positive aggregates were detected in cerebral organoids generated from PITRM1$^{-/-}$
iPSCs, indicating that protein deposits are organized into amyloid-like aggregates,
similar to the ones observed in AD plaques (Supplementary Fig. 2E). Next, we
examined the expression levels of genes involved in UPR^{mt} in 2-month old PITRM1$^{+/+}$
and PITRM1$^{-/-}$ cerebral organoids. PITRM1$^{-/-}$ cerebral organoids exhibited a
significant induction of UPR^{mt} transcripts (*ATF4, DDIT3, HSP60, HSPA9, ERO1*)
(Fig.4L). Moreover, gene expression of the mitochondrial proteases, *LONP1* and
CLPP, was significantly upregulated in PITRM1$^{-/-}$ cerebral organoids compared to
controls (Fig.4L).
459

Inhibition of UPR^{mt} exacerbates Aβ proteotoxicity
Given that UPR^{mt} can extend lifespan in a variety of organisms [24-27], we asked
whether the induction of UPR^{mt} observed in PITRM1$^{-/-}$ cerebral organoids could act as

a protective mechanism against defects of mitochondrial protein maturation and Aβ

proteotoxicity. To this end, cerebral organoids were treated daily with ISRIB, a global

ISR inhibitor [28]. First, we examined APP levels and phospho-tau/tau ratio by

immunostaining. Consistent with a protective role of UPR^{mt} in our model, ISRIB-

treated cerebral organoids showed higher APP and phospho-tau levels compared to

controls (Fig. 5A-C). Remarkably, the effect was similar in both PITRM1$^{+/+}$ and

PITRM1$^{-/-}$ organoids (Fig. 5A-C). In parallel, ISRIB treatment significantly increased

the Aβ42/Aβ40 ratio (Fig. 5D) in both PITRM1$^{+/+}$ and PITRM1$^{-/-}$ organoids.

Interestingly, ISRIB-treated cerebral organoids also showed an increase of

mitochondrial DNA, suggesting that inhibition of UPR^{mt} leads to a decrease of

mitochondrial clearance (Fig. 5E).

Enhancing mitophagy via NAD+ precursors ameliorates Aβ proteotoxicity

Since defects in mitophagy have been shown to contribute to AD [29] and ISRIB-

treated cerebral organoids showed decreased mitochondrial clearance and

exacerbation of AD-like phenotypes, we investigated whether enhancing

mitochondrial clearance ameliorates AD-like phenotype in PITRM1-related

mitochondrial neurodegeneration. To this end, PITRM1$^{+/+}$ and PITRM1$^{-/-}$ cerebral

organoids were treated with the NAD+ precursor nicotinamide mononucleotide

(NMN), which has been shown to ameliorate mitochondrial function and clearance [30].

While NMN supplementation resulted in increased mtDNA/nDNA ratio in PITRM1$^{+/+}$

organoids, a significant decrease of mitochondrial content was observed in PITRM1$^{-/-}$

organoids after treatment (Fig. 5F). These data suggest a differential role of NAD+

boosters in the physiological and pathological conditions, namely induction of

mitochondrial biogenesis in physiological condition and enhancement of

mitochondrial clearance in the disease condition. Furthermore, NMN treatment

significantly decreased the Aβ42/Aβ40 ratio and phospho-tau/tau levels in PITRM1$^{-/-}$

cerebral organoids as revealed by Meso Scale and Elisa measurements (Fig. 5G, H).

Discussion

Since the brain is the organ with the highest demand for energy, it comes as no

surprise that it also represents the major disease target, both in genetically driven

primary mitochondrial diseases, as well as in common age-related

neurodegenerative disorders. Despite this evidence, the causal link between

mitochondrial demise and neurodegeneration still remains elusive. We have recently

reported that pathogenic variants in the nuclear-encoded mitochondrial peptidase

PITRM1 result in childhood-onset recessive cerebellar disease leading to a slowly
progressive syndrome, characterized by spinocerebellar ataxia, mild intellectual
disability, psychiatric manifestations, and cognitive decline [2, 3]. The clinical picture of
these patients is unusual for mitochondrial disease, with a very slow progression of
cognitive and psychiatric symptoms from childhood to their late sixties [2].

PITRM1[+/-] mice show a neurological phenotype with presence of Aβ positive plaques
in the neuropilum [2]. However, due to the embryonic lethality of complete PITRM1[-/-2],
the exact mechanisms of PITRM1 in brain function and disease could not be entirely
studied. To address these quetions, we have generated a novel human stem cell-
based model of loss of PITRM1 function that recapitulates several pathological
aspects of human PITRM1-related mitochondrial and adult onset neurodegenerative
diseases. Employing iPSC-derived cortical neurons, we found that loss of PITRM1
function leads to a strong induction of mitochondrial stress responses, enhanced
autophagic flux and mitochondrial clearance, as well as increased levels of APP, Aβ
peptides, and increased Aβ42/40 ratio. Several works have shown the uptake and
accumulation of Aβ within the mitochondria in post-mortem brains of AD patients as
well as in later stages of disease in APP transgenic mice [31, 32]. Furthermore, *in vitro*
and yeast-based modeling has shown that loss of PITRM1 function results in the
incomplete degradation of Aβ in the mitochondria [2, 8]. Using sub-fractionation
methods coupled with a highly sensitive immunoassay, we were unable to detect Aβ
peptides in mitochondrial extracts from PITRM1 deficient cortical neurons. We
cannot exclude the accumulation of low amount of Aβ peptides within the
mitochondria, below the detection limit of this study. However, PITRM1[-/-] neurons did
not show an increase in mitochondrial ROS production or defects in oxidative
phosphorylation, which have been shown to be a direct effect of the accumulation of
Aβ within mitochondria [33, 34]. Our data suggests that MTS toxicity and mitochondrial
proteostasis imbalance alone can be the first event of the pathogenetic cascade in
PITRM1-related neurological syndrome. The mechanisms whereby PITRM1
dysfunction causes APP accumulation may involve proteasome saturation, in
response to mitochondrial protein misfolding. The ubiquitin proteasome system
(UPS) is also involved in the quality control of mitochondrial proteins, especially the
mitochondrial precursor proteins and proteins of the outer membrane [35]. Supporting
an overload of the UPS system, we detected an accumulation of ubiquitinated
proteins in PITRM1-deficient neurons and cerebral organoids. Based on these data,
we propose that mitochondrial proteotoxic stress, possibly due to the accumulation of
non-degraded MTS as a result of PITRM1 dysfunction and accumulation of

unprocessed mitochondrial proteins, triggers a cytosolic response with overload and

saturation of the proteasome and defects in cytosolic protein degradation.

PITRM1 deficiency led to a strong induction of UPR^{mt} in both 2D and brain organoids

model systems. UPR^{mt} is a transcriptional response involving mitochondrial

chaperones and proteases activated by mitochondrial dysfunction and defects in

mitochondrial protein folding [36]. The UPR^{mt} is a key cellular quality control

mechanism that promotes the maintenance of mitochondrial health and ensures

proper cellular functions [37]. Despite the evidence of UPR^{mt} activation in the ageing

and disease brain [16], whether and how UPR^{mt} contributes to neurodegenerative

processes is unclear. The UPR^{mt} has been proposed to be a double edge sword,

with its chronic activation leading to detrimental consequences for cellular and

organismal function [18, 38]. A detrimental role of UPR^{mt} has been demonstrated in

animal models of mitochondrial diseases [18, 38]. However, mitochondrial stress

responses have been documented in AD and recent work has shown that enhancing

UPR^{mt} provides protections against Aβ proteotoxicity [17, 39]. In line with a beneficial role

of UPR^{mt}, PITRM1$^{-/-}$ cerebral organoids treated with ISRIB, an inhibitor of the ISR,

showed a significant increase of APP levels, increased Aβ42/Aβ40 ratio, and tau

hyperphosphorylation. Based on these data, we propose that PITRM1-related

induction of UPR^{mt} is a protective mechanism against proteotoxic stress both at the

mitochondrial and cytosolic level. Importantly, our findings indicate that the

consequences of chronic mtISR upregulation may vary substantially among different

mitochondrial diseases and the underlying molecular defect should be carefully taken

into consideration for therapeutic decisions.

Even though UPR^{mt} was activated in both 2D and 3D PITRM1 KO models, only

PITRM1 KO cerebral organoids displayed the typical abnormalities observed in the

brain of AD patients, including neuronal cell death, tau pathology, and accumulation

of protein aggregates, similar to Aβ plaques. On the contrary, despite APP

accumulation and increased Aβ42/Aβ40 ratio, we did not detect overt cell death, nor

Aβ aggregates or tau pathology in 2D iPSC-derived neuronal cultures. These

findings indicate that 3D systems provide a more relevant, compared to 2D, disease

model, advantageous in investigating the link between cellular proteostasis and

disease. Due to the prolonged culturing conditions, as well as the presence and

interaction among different cell types, including glial cells, 3D model systems may

promote the development of disease relevant phenotypes, such as protein

aggregation and neuronal death [40]. In respect to the mechanisms, these data also

suggest that PITRM1 deficiency triggers compensatory quality control mechanisms

both at the cytosolic and mitochondrial level (i.e. induction of UPR^{mt} and
autophagy/mitophagy) that ensure the maintenance of cellular proteostasis. However,
over time, these mechanisms may be not sufficient to protect neuronal cells against
mitochondrial proteotoxicity, as observed in long-term culture cerebral organoids.

Several findings, including the induction of autophagic flux, decreased mtDNA levels,
and increase of mitochondrial protein ubiquitination suggest that PITRM1 deficiency
leads to increased mitochondrial clearance. It is known that defects in PITRM1
activity lead to impaired MTS processing and accumulation of MTS and precursor
proteins that have a toxic effect on mitochondria. In line with this evidence, we report
that PITRM1$^{-/-}$ neurons show defects in the maturation of the human frataxin
precursor and decreased mitochondrial membrane potential. Enhanced mitophagy
could be triggered by mitochondrial depolarization in response to MTS accumulation
within mitochondria. Furthermore, UPR^{mt}, and in general the ISR, activates the
autophagic pathway [20, 21]. Interestingly, inhibition of ISR, resulted in an increase in
mtDNA suggesting that the activation of UPR^{mt} is linked to the enhanced
mitochondrial clearance in our model.

Interestingly, mitochondrial stress response and mitophagy transcripts have been
found to be upregulated in mild cognitive impairment as well as in mild and moderate
AD patients, whereas defective mitophagy may play a role in the disease progression
at later stages [29, 41]. Fang et al have recently shown that the enhancement of
mitophagy is able to rescue AD-related pathology in different AD model systems [29].
In line with this finding, we show that stimulating mitophagy with NMN, a NAD+
booster, significantly improves mitochondrial clearance, with a reduction of
Aβ42/Aβ40 ratio and tau hyperphosphorylation. On the contrary, inhibition of UPR^{mt}
with ISRIB led to decreased mitochondrial clearance and aggravation of Aβ and tau
pathology. Thus, our data suggest a protective role of mitophagy against
mitochondrial proteotoxicity induced by PITRM1 deficiency. Interestingly, NMN-
related induction of mitophagy was evident only in PITRM1$^{-/-}$ cerebral organoids,
while induction of mitochondrial biogenesis was detected in PITRM1$^{+/+}$ organoids
upon treatment.

In conclusion, we report a novel cellular model of human PITRM1 deficiency that
recapitulates several fundamental pathological aspects of PITRM1-related
mitochondrial disease. Using human iPSC-derived cortical neurons and cerebral
organoids, we show that loss of PITRM1 function leads to pathological features

similar to the ones observed in AD, namely protein aggregation, tau
hyperphosphorylation, and neuronal death. We report that PITRM1 deficiency
induces impairment of mitochondrial proteostasis and activation of UPR^{mt} that
activates cytosolic quality control pathways, such as the UPS and autophagy. The
overload of the UPS causes, on the long run, a reduced capacity of degrading
cytosolic proteins leading to APP accumulation, increased level of Aβ species,
increased Aβ42/40 ratio, and extracellular protein aggregation. Furthermore, we
show that, similar to what has been described in AD, enhancing UPR^{mt} and
mitophagy ameliorates neuropathological features in primary mitochondrial disease-
related neurodegeneration. Importantly, although PITRM1 mutations are relatively
rare, the disease mechanisms described in the present study may apply to both
primary mitochondrial diseases and more common adult-onset neurological diseases.
Thus, our data support a mechanistic link between primary mitochondrial disorders
and common neurodegenerative proteinopathies.
623
Figure legend.
Figure 1. Mitochondrial dysfunction in PITRM1$^{-/-}$ iPSC-derived cortical neurons.
Control PITRM1$^{+/+}$ (WT) and isogenic PITRM1$^{-/-}$ (KO) iPSCs were differentiated into
cortical neurons. **(A)** Immunostaining of indicated differentiated iPSC cultures at day
in vitro 35. Cells were stained for TBR1 (green) and β-III Tubulin (β-TUBIII, red).
Nuclei were counterstained with DAPI (blue). Scale bars, 50 μm. **(B)** Representative
western blot for PITRM1, showing absence of PITRM1 in human PITRM1$^{-/-}$ iPSC-
derived cortical neurons. **(C)** Mitochondrial membrane potential in isogenic
PITRM1$^{+/+}$ and PITRM1$^{-/-}$ iPSC-derived neurons, as determined by a
tetramethylrhodamine methyl ester (TMRM) assay and flow cytometry analysis. Data
are expressed as percentage of TMRM positive cells (mean + SEM; * $p<0.05$, two-
tailed t test, n=4). **(D)** Mitochondrial reactive oxidative species level as analyzed by
MitoSOX labeling in isogenic PITRM1$^{+/+}$ and PITRM1$^{-/-}$ iPSC-derived neurons (mean
+ SEM, n=5). **(E)** Oxygen consumption rate (OCR) of PITRM1$^{+/+}$ and PITRM1$^{-/-}$ iPSC-
derived neurons. Data are normalized to protein content (mean ± SD, n=6). **(F, G)**
Western blot analysis of OXPHOS complex protein levels in PITRM1$^{+/+}$ and PITRM1^{-}
$^{/-}$ iPSC-derived neurons. Representative blot is shown in (F) and the quantification is
shown in (G) (mean + SEM; * $p<0.05$, two-tailed t test, n=4).
642
Figure 2. PITRM1$^{-/-}$ iPSC-derived neurons show induction of UPR^{mt} and
enhanced mitophagy. (A) Gene expression levels of mitochondrial stress response

genes in PITRM1$^{+/+}$ and PITRM1$^{-/-}$ iPSC-derived cortical neurons (mean + SEM; **

p<0.01, * p<0.05, two-tailed t test, n=5). **(B)** Representative Western blots of the

mitochondrial chaperones HSPA9, HSP60 and the mitochondrial protease LONP1 in

PITRM1$^{+/+}$ and PITRM1$^{-/-}$ iPSC-derived cortical neurons. **(C)** Immunostaining of

PITRM1$^{+/+}$ and PITRM1$^{-/-}$ iPSC-derived cortical neurons for LC3 (green) and β-TUBIII

(red). Nuclei were counterstained with DAPI (blue). Scale bars, 10 μm. **(D)** Number

of LC3-positive vesicles per β-TUBIII positive cell relative to control neurons (mean +

SEM; *** p<0.001, two-tailed t test, n=3). **(E)** Western blot analysis for LC3 in

PITRM1$^{+/+}$ and PITRM1$^{-/-}$ iPSC-derived neuronal cultures, untreated (–) or treated

with 200 μM leupeptin and 20 mM NH$_4$Cl for 4 h (+). **(F)** Quantification of LC3 flux

normalized to WT (mean + SEM; * p<0.05, two-tailed t test, n=4). **(G)** Representative

Western blot of isolated mitochondria from PITRM1$^{+/+}$ and PITRM1$^{-/-}$ iPSC-derived

neurons with an antibody for ubiquitination and VDAC as loading control. **(H)**

Quantification of mitochondrial protein ubiquitination levels in PITRM1$^{+/+}$ and

PITRM1$^{-/-}$ iPSC-derived neurons (mean + SEM; ** p<0.01, two-tailed t test, n=3). **(I)**

mtDNA content was measured as mitochondrial (*16S*) to nuclear (*RPLP0*) DNA ratio

by qRT-PCR (mean + SEM; * p<0.05, two-tailed t test, n=3).

Figure 3. PITRM1$^{-/-}$ iPSC-derived neurons show accumulation of APP and

increased levels of Aβ peptides. (A) Representative Western blots of APP in

PITRM1$^{+/+}$ and PITRM1$^{-/-}$ iPSC-derived cortical neurons. Quantification of APP levels

normalized to loading control (mean + SEM; * p<0.05, two-tailed t test, n=5). **(B)**

Quantification of Aβ species and Aβ42/Aβ40 ratio in the supernatant of PITRM1$^{+/+}$

and PITRM1$^{-/-}$ iPSC-derived cortical neurons at day in vitro 35, as performed by

Meso Scale immunoassay (mean + SEM; ** p<0.01, *** p<0.001, two-tailed t test,

n=4). **(C)** Representative Western blot of PITRM1$^{+/+}$ and PITRM1$^{-/-}$ iPSC-derived

neurons of total ubiquitinated proteins levels. **(D)** Quantification of protein

ubiquitination level in PITRM1$^{+/+}$ and PITRM1$^{-/-}$ iPSC-derived neurons (mean + SEM;

* p<0.05, two-tailed t test, n=4).

Figure 4. PITRM1$^{-/-}$ cerebral organoids display main pathological features of AD

pathology and induction of mitochondrial stress response. (A) Generation and

characterization of cerebral organoids from PITRM1$^{+/+}$ and PITRM1$^{-/-}$ iPSCs.

Immunostaining for β-TUBIII (green), TBR1 (red), and CTIP2 (red) of 2-month old

cerebral organoids. Cell nuclei were counterstained with DAPI (blue). Scale bar,

100μm. **(B)** Western blot of APP in 2-month old PITRM1$^{+/+}$ and PITRM1$^{-/-}$ cerebral

organoids. **(C)** Quantification of APP protein levels in cerebral organoids (mean +

SEM; * p<0.05, two-tailed t test, n=5). **(D)** Representative Western blot of phospho-

tau (p-tau) and total tau in PITRM1$^{+/+}$ and PITRM1$^{-/-}$ cerebral organoids; total tau and

β-Actin were used as loading controls. **(E)** Quantification of phospho-tau protein

levels in cerebral organoids relative to loading control total tau/β-Actin (mean + SEM;

* p<0.05, two-tailed t test, n=5). **(F, G)** MAP2 (green), APP (red, left panel), and

phospho-tau (red, right panel) immunostaining in cerebral organoids. Representative

confocal images are shown. Cell nuclei were counterstained with DAPI (blue). Scale

bars, 100 μm. **(H)** Immunostaining for β-TUBIII (green) and cleaved Caspase 3

(cCASP3, red) in PITRM1$^{+/+}$ and PITRM1$^{-/-}$ 2-month old cerebral organoids. Cell

nuclei were counterstained with DAPI (blue). Scale bar, 100μm. **(I)** Quantification of

APP, phospho-tau fluorescent intensity and analysis of the ratio of cCASP3 positive

cells relative to the total number cells, measured by DAPI staining, in 2-month old

cerebral organoids (mean + SEM; * p<0.05, *** p<0.001, two-tailed t test, n=3-4). **(J)**

MAP2 (green) and Ubiquitin (red) immunostaining in cerebral organoids.

Representative confocal images are shown. Cell nuclei were counterstained with

DAPI (blue). Scale bars, 100 μm. **(K)** Quantification of Ubiquitin fluorescent intensity

(mean + SEM; *** p<0.001, two-tailed t test, n=3). **(L)** Gene expression levels of

mitochondrial stress response genes in PITRM1$^{+/+}$ and PITRM1$^{-/-}$ cerebral organoids

(mean + SEM; * p<0.05, ** p<0.01, two-tailed t test, n=5).

Figure 5. UPR^{mt} and mitophagy exert a protective role in PTRM1$^{-/-}$ cerebral

organoids. PITRM1$^{+/+}$ and PITRM1$^{-/-}$ cerebral organoids were treated with 500 nM

ISRIB or vehicle. **(A, B)** MAP2 (green), APP (red, in A), and phospho-tau (red, in B)

immunostaining in PITRM1$^{+/+}$ and PITRM1$^{-/-}$ cerebral organoids treated with ISRIB or

vehicle. Representative confocal images are shown. Cell nuclei were counterstained

with DAPI (blue). Scale bars, 100 μm. **(C)** Quantification of APP and phospho-tau

fluorescent intensity (mean + SEM; *** p<0.001, ** p<0.01, * p<0.05, two-tailed t test,

n=3-4). **(D)** Quantification of Aβ species in supernatant of PITRM1$^{+/+}$ and PITRM1$^{-/-}$

cerebral organoids treated with ISRIB or vehicle, as measured by Meso Scale

immunoassay (mean + SD; ** p<0.01, *p<0.05, two-tailed t test, n=5). **(E)** mtDNA

content was measured in PITRM1$^{+/+}$ and PITRM1$^{-/-}$ cerebral organoids treated with

ISRIB or vehicle as mitochondrial (*16S*) to nuclear (*RPLP0*) DNA ratio by qRT-PCR

(mean + SD; * p<0.05, two-tailed t test, n=3). **(F-H)** PITRM1$^{+/+}$ and PITRM1$^{-/-}$ cerebral

organoids were treated with 500 μM NMN or vehicle. (F) mtDNA content was

measured as mitochondrial (*16S*) to nuclear (*RPLP0*) DNA ratio by qRT-PCR (mean

+ SEM; *** p<0.001, * p<0.05 two-tailed t test, n=3). (G) Aβ1–42/Aβ1–40 ratio in

cerebral organoids measured by Meso Scale immunoassay (mean + SEM; * p<0.05,

two-tailed t test, n=3). (H) ELISA assay measuring the levels of total and phospho-

tau levels in cerebral organoid homogenates. Protein concentration was measured

by BCA and equal amounts of total protein were used (mean + SEM; * p<0.05, ***

p<0.001, two-tailed t test, n=3).

Acknowledgements

We acknowledge the funding support of CoEN Pathfinder II (Ref. 3038, to M.D.,

M.Z.), the Helmholtz Association (to M.D.), DAAD (PKZ 91723383, to M.J.P.), and

Fondazione Umberto Veronesi 2018-2019 (to D.B.) for this project.

Author contributions

M.D., D.I. and M.J.P. conceived the study. M.D., D.I., M.J.P., V.P., D.B., S.A.K., M.J.,

M.Z., C.V.contributed to experimental design. D.I. performed gene editing. M.J.P.,

V.P., G.D.N. performed the most of the experiments with iPSCs and cerebral

organoids., R.S performed Meso Scale assay, M.D., D.I., M.J.P., V.P. analysed data.

M.D. wrote the manuscript with input and approval from all the authors.

Conflict of interest
737
738 The authors declare no competing interests
739

740 **References**

741 1.   Johri A, Beal MF. Mitochondrial dysfunction in neurodegenerative diseases. *J*
742 *Pharmacol Exp Ther* 2012; **342**(3)**:** 619-630.
743

744 2.   Brunetti D, Torsvik J, Dallabona C, Teixeira P, Sztromwasser P, Fernandez-Vizarra E
745 *et al.* Defective PITRM1 mitochondrial peptidase is associated with Abeta amyloidotic
746 neurodegeneration. *EMBO Mol Med* 2016; **8**(3)**:** 176-190.
747

748 3.   Langer Y, Aran A, Gulsuner S, Abu Libdeh B, Renbaum P, Brunetti D *et al.*
749 Mitochondrial PITRM1 peptidase loss-of-function in childhood cerebellar atrophy. *J Med*
750 *Genet* 2018; **55**(9)**:** 599-606.
751

752 4.   Town L, McGlinn E, Fiorenza S, Metzis V, Butterfield NC, Richman JM *et al.* The
753 metalloendopeptidase gene Pitrm1 is regulated by hedgehog signaling in the developing
754 mouse limb and is expressed in muscle progenitors. *Dev Dyn* 2009; **238**(12)**:** 3175-3184.
755

756 5.   Stahl A, Nilsson S, Lundberg P, Bhushan S, Biverstahl H, Moberg P *et al.* Two novel
757 targeting peptide degrading proteases, PrePs, in mitochondria and chloroplasts, so similar
758 and still different. *J Mol Biol* 2005; **349**(4)**:** 847-860.
759

760 6.   van 't Hof R, Demel RA, Keegstra K, de Kruijff B. Lipid-peptide interactions between
761 fragments of the transit peptide of ribulose-1,5-bisphosphate carboxylase/oxygenase and
762 chloroplast membrane lipids. *FEBS Lett* 1991; **291**(2)**:** 350-354.
763

7.   Zardeneta G, Horowitz PM. Analysis of the perturbation of phospholipid model
membranes by rhodanese and its presequence. *The Journal of biological chemistry* 1992;
267(34)**:** 24193-24198.

8.   Mossmann D, Vogtle FN, Taskin AA, Teixeira PF, Ring J, Burkhart JM *et al.* Amyloid-
beta peptide induces mitochondrial dysfunction by inhibition of preprotein maturation. *Cell*
Metab 2014; **20**(4)**:** 662-669.

9.   Falkevall A, Alikhani N, Bhushan S, Pavlov PF, Busch K, Johnson KA *et al.*
Degradation of the amyloid beta-protein by the novel mitochondrial peptidasome, PreP. *The*
Journal of biological chemistry 2006; **281**(39)**:** 29096-29104.

10.   Teixeira PF, Pinho CM, Branca RM, Lehtio J, Levine RL, Glaser E. In vitro oxidative
inactivation of human presequence protease (hPreP). *Free Radic Biol Med* 2012; **53**(11)**:**
2188-2195.

11.   Pinho CM, Teixeira PF, Glaser E. Mitochondrial import and degradation of amyloid-
beta peptide. *Biochimica et biophysica acta* 2014; **1837**(7)**:** 1069-1074.

12.   Reinhardt P, Schmid B, Burbulla LF, Schondorf DC, Wagner L, Glatza M *et al.*
Genetic correction of a LRRK2 mutation in human iPSCs links parkinsonian
neurodegeneration to ERK-dependent changes in gene expression. *Cell stem cell* 2013;
12(3)**:** 354-367.

13.   Brennand KJ, Simone A, Jou J, Gelboin-Burkhart C, Tran N, Sangar S *et al.*
Modelling schizophrenia using human induced pluripotent stem cells. *Nature* 2011;
473(7346)**:** 221-225.

14.   Lancaster MA, Renner M, Martin CA, Wenzel D, Bicknell LS, Hurles ME *et al.*
Cerebral organoids model human brain development and microcephaly. *Nature* 2013;
501(7467)**:** 373-379.

15.   Branda SS, Cavadini P, Adamec J, Kalousek F, Taroni F, Isaya G. Yeast and human
frataxin are processed to mature form in two sequential steps by the mitochondrial processing
peptidase. *The Journal of biological chemistry* 1999; **274**(32)**:** 22763-22769.

16.   Pellegrino MW, Haynes CM. Mitophagy and the mitochondrial unfolded protein
response in neurodegeneration and bacterial infection. *BMC Biol* 2015; **13:** 22.

17.   Sorrentino V, Romani M, Mouchiroud L, Beck JS, Zhang H, D'Amico D *et al.*
Enhancing mitochondrial proteostasis reduces amyloid-beta proteotoxicity. *Nature* 2017;
552(7684)**:** 187-193.

18.   Anderson CJ, Bredvik K, Burstein SR, Davis C, Meadows SM, Dash J *et al.* ALS/FTD
mutant CHCHD10 mice reveal a tissue-specific toxic gain-of-function and mitochondrial stress
response. *Acta neuropathologica* 2019; **138**(1)**:** 103-121.

19.   Forsstrom S, Jackson CB, Carroll CJ, Kuronen M, Pirinen E, Pradhan S *et al.*
Fibroblast Growth Factor 21 Drives Dynamics of Local and Systemic Stress Responses in
Mitochondrial Myopathy with mtDNA Deletions. *Cell Metab* 2019.

20.   B'Chir W, Maurin AC, Carraro V, Averous J, Jousse C, Muranishi Y *et al.* The
eIF2alpha/ATF4 pathway is essential for stress-induced autophagy gene expression. *Nucleic*
Acids Res 2013; **41**(16)**:** 7683-7699.

21.   Rzymski T, Milani M, Pike L, Buffa F, Mellor HR, Winchester L *et al.* Regulation of
autophagy by ATF4 in response to severe hypoxia. *Oncogene* 2010; **29**(31)**:** 4424-4435.

22.   Grumati P, Dikic I. Ubiquitin signaling and autophagy. *The Journal of biological*
chemistry 2018; **293**(15)**:** 5404-5413.

23.   Segref A, Kevei E, Pokrzywa W, Schmeisser K, Mansfeld J, Livnat-Levanon N *et al.*
Pathogenesis of human mitochondrial diseases is modulated by reduced activity of the
ubiquitin/proteasome system. *Cell Metab* 2014; **19**(4)**:** 642-652.
827

828   24.   Durieux J, Wolff S, Dillin A. The cell-non-autonomous nature of electron transport
829 chain-mediated longevity. *Cell* 2011; **144**(1)**:** 79-91.
830

831   25.   Houtkooper RH, Mouchiroud L, Ryu D, Moullan N, Katsyuba E, Knott G *et al.*
832 Mitonuclear protein imbalance as a conserved longevity mechanism. *Nature* 2013;
833 **497**(7450)**:** 451-457.
834

835   26.   Merkwirth C, Jovaisaite V, Durieux J, Matilainen O, Jordan SD, Quiros PM *et al.* Two
836 Conserved Histone Demethylases Regulate Mitochondrial Stress-Induced Longevity. *Cell*
837 2016; **165**(5)**:** 1209-1223.
838

839   27.   Borch Jensen M, Qi Y, Riley R, Rabkina L, Jasper H. PGAM5 promotes lasting FoxO
840 activation after developmental mitochondrial stress and extends lifespan in Drosophila. *eLife*
841 2017; **6**.
842

843   28.   Sidrauski C, McGeachy AM, Ingolia NT, Walter P. The small molecule ISRIB
844 reverses the effects of eIF2alpha phosphorylation on translation and stress granule assembly.
845 *eLife* 2015; **4**.
846

847   29.   Fang EF, Hou Y, Palikaras K, Adriaanse BA, Kerr JS, Yang B *et al.* Mitophagy
848 inhibits amyloid-beta and tau pathology and reverses cognitive deficits in models of
849 Alzheimer's disease. *Nature neuroscience* 2019; **22**(3)**:** 401-412.
850

851   30.   Fang EF, Kassahun H, Croteau DL, Scheibye-Knudsen M, Marosi K, Lu H *et al.*
852 NAD(+) Replenishment Improves Lifespan and Healthspan in Ataxia Telangiectasia Models
853 via Mitophagy and DNA Repair. *Cell Metab* 2016; **24**(4)**:** 566-581.
854

855   31.   Hansson Petersen CA, Alikhani N, Behbahani H, Wiehager B, Pavlov PF, Alafuzoff I
856 *et al.* The amyloid beta-peptide is imported into mitochondria via the TOM import machinery
857 and localized to mitochondrial cristae. *Proceedings of the National Academy of Sciences of
858 the United States of America* 2008; **105**(35)**:** 13145-13150.
859

860   32.   Manczak M, Anekonda TS, Henson E, Park BS, Quinn J, Reddy PH. Mitochondria
861 are a direct site of A beta accumulation in Alzheimer's disease neurons: implications for free
862 radical generation and oxidative damage in disease progression. *Human molecular genetics*
863 2006; **15**(9)**:** 1437-1449.
864

865   33.   Casley CS, Canevari L, Land JM, Clark JB, Sharpe MA. Beta-amyloid inhibits
866 integrated mitochondrial respiration and key enzyme activities. *Journal of neurochemistry*
867 2002; **80**(1)**:** 91-100.
868

869   34.   Aleardi AM, Benard G, Augereau O, Malgat M, Talbot JC, Mazat JP *et al.* Gradual
870 alteration of mitochondrial structure and function by beta-amyloids: importance of membrane
871 viscosity changes, energy deprivation, reactive oxygen species production, and cytochrome c
872 release. *J Bioenerg Biomembr* 2005; **37**(4)**:** 207-225.
873

874   35.   Livnat-Levanon N, Glickman MH. Ubiquitin-proteasome system and mitochondria -
875 reciprocity. *Biochimica et biophysica acta* 2011; **1809**(2)**:** 80-87.
876

877   36.   Nargund AM, Pellegrino MW, Fiorese CJ, Baker BM, Haynes CM. Mitochondrial
878 import efficiency of ATFS-1 regulates mitochondrial UPR activation. *Science* 2012;
879 **337**(6094)**:** 587-590.
880

881   37.   Shpilka T, Haynes CM. The mitochondrial UPR: mechanisms, physiological functions
882 and implications in ageing. *Nat Rev Mol Cell Biol* 2018; **19**(2)**:** 109-120.
883

884 38. Khan NA, Nikkanen J, Yatsuga S, Jackson C, Wang L, Pradhan S *et al.* mTORC1
885 Regulates Mitochondrial Integrated Stress Response and Mitochondrial Myopathy
886 Progression. *Cell Metab* 2017; **26**(2)**:** 419-428 e415.
887
888 39. Beck JS, Mufson EJ, Counts SE. Evidence for Mitochondrial UPR Gene Activation in
889 Familial and Sporadic Alzheimer's Disease. *Curr Alzheimer Res* 2016; **13**(6)**:** 610-614.
890
891 40. Gerakis Y, Hetz C. Brain organoids: a next step for humanized Alzheimer's disease
892 models? *Mol Psychiatry* 2019; **24**(4)**:** 474-478.
893
894 41. Kerr JS, Adriaanse BA, Greig NH, Mattson MP, Cader MZ, Bohr VA *et al.* Mitophagy
895 and Alzheimer's Disease: Cellular and Molecular Mechanisms. *Trends in neurosciences*
896 2017; **40**(3)**:** 151-166.
897
898

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5