Article type: Original Research Article

Title: Physical-biological interaction underlying phylogeographic pattern of coral-
dependent fishes around the Arabian Peninsula

Running Head: Biophysical model of reef fishes

Authors (names, affiliation, e-mail addresses):
Felipe Torquato[(1)]*, felipe_o_torquato@hotmail.com (correspondence author)

(1) *Section for Evolutionary Genomics, Natural History Museum of Denmark, University*
of Copenhagen, Denmark.

Title: Physical-biological interaction underlying phylogeographic pattern of coral-

dependent fishes around the Arabian Peninsula

Abstract

For marine taxa with freely mobile larvae, oceanographic features can act as soft barriers

by disrupting the larval movements, and ultimately shaping both species and genetic

diversity distributional. This study focused on three putative oceanographic barriers

around the Arabian Peninsula: the Bab el-Mandeb Strait, the Strait of Hormuz and the

upwelling off Oman. We aimed to assess the effects of larval fish traits and ocean

circulation on the permeability of these barriers, and thus provide insights on the

phylogeographic pattern of coral-dependent fish species along the peninsula. To do so,

we simulated the larval dispersal through the putative barriers using an open-source

program, where three-dimensional movements of particles are tracked by using ocean

circulation data. We explored the range of dispersal capabilities across reef fish species

by creating 72 hypothetical strategies, each representing a unique combination of five

biological traits: pelagic larval duration, spawning periodicity, mortality rate, productivity

and vertical migration. Our results showed that the direction and distance travelled by the

virtual larvae were highly variable as a function of both the geographic position of

releasing site and biannual monsoonal winds. On average, larvae released during the

summer exhibited a higher potential for dispersal than larvae released in wintertime. Our

biophysical models complement the few existing empirical research on population

genetics, and the predictions presented here serve as testable hypotheses to future

phylogeographic studies around the Arabian Peninsula.

Keywords: Biophysical modeling, connectivity, upwelling off Oman, Bab el-Mandeb Strait, Strait of Hormuz, connectivity modeling system

Introduction

The Arabian Peninsula lies on a hyper-arid region in the Southwest Asia at the junction of this continent with Africa (Fig. 1). The water-mass distribution and upper-ocean circulation surrounding the peninsula change in correspondence with biannual wind reversals, which creates seasonality in oceanographic conditions (Cutler and Swallow 1984; Shetye et al. 1994). During winter monsoon (November - March), the wind blows away from Asian continent (southwest), and the ocean surface circulation in the Arabian Sea is approximately counter-clockwise. On the opposite, during the summer monsoon (May-September) the wind reverses and blows strongly northeastward, so that the circulation in the Arabian Sea is clockwise. March-April and October are transition periods and winds are weak (see Cutler and Swallow 1984).

Large seasonal variation of such magnitude, which is generally not found elsewhere (Shetye et al. 1994), plays a critical role in fish biodiversity distribution by transporting or disrupting the movements of larvae produced by the adults (Cowen 2002). Thereby, the direction and magnitude of prevailing currents can spread out species distribution by aiding the dispersal of the propagules to future habitats. Nevertheless, currents can also act as a barrier by minimizing long-distance larval dispersal (Cowen et al. 1994, Paris and Cowen 2004) that ultimately may explain either the existence of endemic species (Cowen et al. 2000) or genetic discontinues across species range (Bowen et al. 2016). Although large expanses of ocean waters are the most notable permeable

barriers to larval dispersal (Bowen et al. 2016), mesoscale oceanographic processes such

as fronts (Galarza et al. 2009), river runoff (Rocha 2003), and upwelling (Bakun 1996;

Lett et al. 2007) also act as barriers to marine faunal exchange. The permeability of these

so-called semi-permeable barriers is taxon-specific and, therefore, affected by biological

traits (Ayre et al. 2009).

In the Arabian Peninsula, zoogeographic and population genetic studies on coral-

dependent fishes have shown discontinuities in both species and genetic distribution

along seas surrounding the peninsula (Burt et al. 2011; DiBattista et al. 2016a; Berumen

et al. 2017). The most remarkable of these discontinuities, distinguishes the fish fauna

from each side of the peninsula (DiBattista et al., 2016b). Two other discontinuities in

species compositions are observed, one between the Red Sea and the adjacent Gulf of

Aden through the Bab el-Mandeb Strait, and another between the Arabian Gulf and

adjacent Gulf of Oman through the Strait of Hormuz (DiBattista et al. 2016a). These

boundaries described by multispecies distribution records, are also logical places to

present-day barriers to gene flow between populations (Baums et al. 2006). Indeed,

population genetic studies carried out in the Northwester Indian Ocean (NIO) have

revealed similarities in the geographical position of barriers previously proposed

(DiBattista et al. 2013, 2015, 2017; Nanninga et al. 2014; Saenz-Agudelo et al. 2015;

Priest et al. 2016; Torquato et al. 2019).

Hypotheses to explain the distribution of genetic diversity in the Arabian

Peninsula usually rely on either parapatric speciation outcomes, which resulted from

either repeated vicariance events caused by lowering sea level during the last glaciation

or ecological speciation due to the large spatial gradients and temporal fluctuations in

physical conditions across the peninsula (Nanninga et al. 2014; DiBattista et al. 2016a,b).

An alternative hypothesis has been attributed to the seasonal upwelling in the Arabian

Sea off Oman, which creates unsuitable condition for coral-habitat growth (Sheppard and

Salm 1988) and restrict stepping-stone connectivity between both sides of the Arabian

Peninsula. In turn, little attention has been given to test hypotheses where the

combination of seascape features, ocean circulation and larval traits underling the genetic

patterns observed among coral-dependent fishes inhabiting the Arabian Peninsula.

In numerous cases, hypotheses in marine ecology are prohibitively time

consuming and expensive to be empirically tested, largely owing to the impossibility of

capturing the full range of temporal and spatial fine-scale resolution required to make

inferences (Cowen and Sponaugle 2009). Although a more comprehensive picture is

emerging in the Arabian Peninsula with respect to marine phylogeographic and

population genetic patterns (DiBattista et al. 2016a,b; Berumen et al. 2017), the processes

affecting larval dispersal across the putative barriers are not yet fully understood due to

the paucity of empirical studies. In addition, political realities of some countries

bordering the Western Indian Ocean (WIO) have limited access to scientists, hindering a

more complete perspective of general phylogeography in the region (Berumen et al.

2017). In such cases, if limited empirical data are available, reliable computational

models can be used to make field predictions and advance our knowledge to designing

future experiments for hypothesis testing (Cowen and Sponaugle 2009).

Advances on physical circulation models enabled the investigation of population

connectivity by running semi-realistic simulations of virtual particles. Here we used high-

resolution ocean circulation model (Hybrid Coordinate Ocean Model - HYCOM) to

design a biophysical model in a Lagrangian stochastic scheme (Paris et al. 2013). The

main goal of this study was to simulate multitaxon larval dispersal through the marine

barriers, and thus provide insights on processes and patterns of connectivity leading to the

distribution of genetic diversity of coral-dependent fishes around the Arabian Peninsula.

Specifically, we shed light on three questions: (1) what biological attributes affect the

larval dispersal through the Bab el-Mandeb Strait and Strait of Hormuz? (2) What is

impact of the upwelling off Oman on the larval connectivity between both sides of the

peninsula? (3) How does oceanographic variability, due to the seasonal monsoon, affect

larval dispersal pattern (i.e. direction and magnitude travelled by the particle)? Our

results provide detailed predictions that can be compared to previous and future empirical

studies on the distribution of biodiversity in the Arabian Peninsula.

Material and Methods

Biophysical Model and Larval Dispersal Simulation

Idealized dispersal of fish larvae was modeled using an open-source program,

Connectivity Modelling System (CMS v. 2.0, Paris et al. 2013), which is a biophysical

modeling system based on stochastic Lagrangian framework and Individual-Based Model

(IBM) that couple ocean current, GIS-based habitat, and biological traits. In brief, CMS

uses information on currents and environmental conditions to simulate both deterministic

fourth-order Runge-Kutta and/or stochastic displacements of a large number of virtual

particles (hereafter called larvae), through space and time. In order to explore the range of

dispersal capabilities across reef fishes species, we created 72 hypothetical strategies each

of those representing a unique combination of five biological traits that may influence the

connectivity: pelagic larval duration (PLD), spawning periodicity, mortality rate,

productivity and vertical migration (Table 1).

Hydrodynamic model

We used the CMS package *getdata* to download daily ocean current velocity from

the three-dimensional and eddy-resolving Hybrid Coordinate Ocean Model (HYCOM,

GLBa0.08) from 2014 to 2016. The model had a horizontal resolution of ca. 9 km grid

(1/12˚), and was set up in a nested domain (0˚ - 30˚ N, 31˚ - 70˚ E), comprising all layers

from the surface up to 100 meters depth. In studies carried out around the Arabian

Peninsula, HYCOM has been used to investigate mesoscale variability in the Arabian Sea

(L'Hégaret et al. 2015) as well as water circulation and exchange in the Persian Gulf

(Yao and Johns 2010).

GIS-based Seascape module

The GIS served to delineate the source and recruitment habitats. Suitable

releasing (source habitat) and settlement (sink habitat) locations were delineated in QGIS

v.2.18 by creating a vector grid that overlaid the distribution of coral reefs data from the

UNEP-WCMC (2010). A total of 181 polygons (~ 18km x 18km) representing the coral

reef habitats were placed along coastal areas within our model domain.

Biological module

We assumed spatial homogenous productivity across the habitats and, therefore,

all the 181 polygons released the same number of larvae every 'spawning' event.

However, we contrasted scenarios where productivity varies among species. In scenario

of high productivity, a hundred of larvae were released from each polygon, whereas in

low productivity conditions only ten were released. Here, three seasonal spawning

preferences were accounted by simulating the larval dispersal annually and accordingly

the two dominant seasonal oceanographic conditions around the Arabian Peninsula. After

being released, the larvae were let to drift over a period of 20, 30 or 40 days,

corresponding to the typical pelagic duration (PLD) of most coral reef fish

species/families (Thresher and Brothers 1985; Lindeman et al. 2005).

In this study the PLD was equally divided into the three ontogenetic larval stages

termed preflexion, flexion and postflexion. Due to the paucity of basic data regarding fish

larvae distribution within our study area, the model incorporated an idealized pattern of

ontogenetic vertical migration for the three larval stages (Fig. S1). This idealization

assumes the existence of a global ontogenetic trend of the fish larvae to display

downward ontogenetic shift in vertical distribution (Irisson et al. 2010). During the

postflexion stage the larvae were considered competent to settle if they are inside one of

the 181 reef sites. To assess the importance of vertical migration, we also modeled

epipelagic ichthyoplankton moving only along the sub-surface layer to then contrast with

models where larvae also moved vertically.

Little is known regarding larval mortality in the ocean. To accommodate this

uncertainty, our study included two levels of mortality based on the half-life, such that

approximately 50% of unsettled larvae would be surviving after half the maximum PLD

(Paris et al. 2013; Holstein et al. 2014). Thus, we determined that in high mortality

scenario half of the larvae died by the end of the preflexion stage, while in low mortality

condition half of the larvae died by the end of the postflexion stage (Table 1).

Particle-tracking module

Stochastic IBM Lagrangian model tracked offline over 157 millions larvae around

the Arabian Peninsula for the 72 model taxa. A total of 86,011,200 larvae were released

to mimic taxa spawning throughout year, whereas 35,838,000 larvae were tracked to

simulate taxa spawning on either winter monsoon (November 2014 – March 2015) or

summer monsoon (May 2015 – September 2015).

Preliminary sensitive analysis showed no significant difference in the settlement

proportion when seeding 100 larvae from the 181 polygons at every either 3 or 24 hours

(see Appendix S1 in Supplemental Information). Therefore, the larvae were released

from each reef at every 24 hours, which represents a spawning event. This uniform

temporal distribution of larvae allows us to assess the effects of the hydrographic

variability conditions on larval dispersal (e.g., extreme events, perturbation, instability).

Throughout the PLD, the position of each larva was updated every 6 h time-step, and the

trajectory information (i.e. longitude, latitude, depth) were saved to output in intervals of

24 hours. We accounted for diffusive turbulent motion by adding a horizontal diffusion

coefficient. The value of 50 m^2s^{-1} was chosen from a sensitivity test where 100 larvae

were released from the 181 polygons at every 24 hours (see Appendix S1 in

Supplemental Information).

Analyses

The proportion of survivor larvae that were released from each region i and

successfully settled in a downstream habitat patch (i.e., sink habitat) at region j,

were plotted as a connectivity matrix. Self-recruitment, i.e. larvae settling within its

release location, was represented by the diagonal of the matrix. In order to evaluate

the strength and direction of the potential connections on regional scale (e.g.,

between Red Sea and Gulf of Aden), all cells from each region were merged.

Biological traits vs. biogeographic barriers

To evaluate the effect of the biological traits highlighted in Table 1 on the

putative barriers, we relied on a multiple regression approach. Specifically, the

effects of both Bab-el-Mandeb and Hormuz strait were measured in terms of

permeability, which considers the proportion of surviving larvae that were released

from a source habitat, passed through the strait, and successfully settled on the

other side. In turn, the effect of the upwelling off Oman was measured in terms of

self-recruitment proportion along the Arabian Sea.

Provided that the proportions are a continuous variable that can take on

values restricted to the interval between 0 and 1, a beta regression model as

proposed by Ferrari and Cribari-Neto (2004) was used through the betareg R-

package (Neto and Zeilis 2010). The beta regression is essentially similar to a

Generalized Linear Model (GLM), where it describes the relationship between the

response variable Y_i (hereby, proportions) and the predictors X_i (Table 1) through a

linear predictor η_i. This linear predictor is then linked to the mean of the response

$E(Y_i) = \mu_i$ by means of a link function g, such that $g(\mu_i) = \eta_i$. In this way, the applied

model can be summarized as:

$$Y_i \sim B(\mu_i, \phi)$$

$$logit(\mu_i) = \eta_i$$

$$\eta_i = \beta_0 + \beta_1 PLD + \beta_2 SP + \beta_3 M + \beta_4 Pr + \beta_5 VM + \varepsilon$$

Where ϕ is the precision parameter; η_i the linear predictor expressed on a logit scale; $\beta 0$

the intercept; $\beta 1$-5 the repressors related to each explanatory variable (PLD=Pelagic

larval duration, SP = Spawning periodicity, M = Mortality, Pr = Productivity, VM =

Vertical migration); and ε represents the error term. Model assumptions like residual

homocedasticity, independence and normality were accessed through conventional

graphical checks as mentioned in Zuur et al. (2010).

Larval dispersal pattern and oceanographic conditions

The maximum distance travelled by the larvae around the Arabian Peninsula was

calculated setting up a model with the lowest mortality rate and 40-days PLD. We used

the program QGIS v.2.18 to measure this distance and compare the results in both

summer and winter, as well as in each of the regions within the model domain.

Results

Connectivity matrix

All the matrices representing the 72 model taxa revealed that connectivity occurred exclusively between neighbor regions, though the strength and symmetry of these connections varied either seasonally or due to larval traits (Fig. S2 – S4). The hardest barrier was observed in the south of the Arabian Peninsula, separating both sides of the peninsula, in other words, larvae that were released from the Arabian Sea, the Gulf of Oman or the Arabian Gulf, did not reach the reefs in the Gulf of Aden or the Red Sea, and vice-versa. This barrier was observed especially in simulations in which the larvae drifted for 20 days. When released during either winter or summer monsoon, 30-days PLD larvae that performed vertical migrations were also not able to cross this barrier. Importantly, the connectivity through the Strait of Hommuz was not symmetric, such that the Gulf acts as a sink habitat specially when the larvae drift for 20 days, with high mortality rate and perform vertical migration (Fig. S2).

Intra-annual variability was evident in the potential connectivity (Fig. S2 – S4), suggesting that symmetry of the connectivity was affected by the spawning periodicity of the taxa. For example, in the summer 40-PLD days larvae released from the west side of the peninsula reached the reefs located on the east side. In contrast, during the winter any of these larvae crossed the barrier in this direction.

Effects of the biological traits on the permeability of biogeographic barriers

Graphical evaluation of the model assumptions revealed that all tested models fitted well to the data, and did not violate neither the residual homocedasticity nor

independence. Table S1 summarizes the numerical output of the models, and discussed in

more detail below.

Bab el-Mandeb Strait

For the larvae that were released in the Red Sea and successfully settled in the

Gulf of Aden, only the larval productivity was not significant (Table S1, Fig. 2). The beta

regression showed that this connectivity was significantly higher during the summer than

both during the winter and throughout year. Also, the permeability through the strait was

higher for those larvae that performed vertical migration and were left to drift for longer

period, though no significant difference was observed between larvae that drift for 30 and

40 days.

On the other hand, larvae that moved in the opposite direction, from the Gulf of

Aden to the Red Sea, showed higher success in the connection during the wintertime but

larval productivity did not change the barrier's permeability (Table S1, Fig. 2). Moreover,

a positive relationship was observed between the permeability and PLD, and epipelagic

larvae that moved at the surface increased their permeability through the strait compared

to those that performed vertical migration. Mortality rate was significant and showed the

same tendency of the larvae moving from the Red Sea and to the Gulf of Aden.

Strait of Hormuz

The proportion of larvae released in the Arabian Gulf that successfully reached

the reefs in the Gulf of Oman, increased when they were released throughout year or

during the wintertime, performed vertical migration, had low mortality rate, and travelled

for longer period (Table S1, Fig. 3). Nevertheless, the chance to cross the strait did not

depend on the larval productivity.

Likewise, among all biological attributes, only productivity was not significant for

larvae that were released in the Gulf of Oman, crossed the Strait of Hormuz and

successfully settled in the Arabian Gulf. The probability of this connectivity was higher

in the scenarios where the larvae were release in winter, had low mortality rate, did not

perform vertical migration and remained in the planktonic environment for longer period.

(Table S1, Fig. 3).

Upwelling off Oman

All biological traits, except larval productivity, significantly affected the

proportion of self-recruitment in the Arabian Sea (Table S1, Fig. 4). The scenario in

which a taxon had higher success in self-recruitment occurred during the wintertime or

throughout year, and when the 20-days PLD larvae performed ontogenetic migration

under low mortality scenario. In turn, larvae produced during the summer, under

upwelling conditions, had a slightly more chance to migrate.

Particles trajectories: seasonal variability of spatial scale and direction

The direction and distance travelled by the larvae were highly variable as a

function of both the geographic position of releasing site and the biannual monsoonal

winds (Fig. 5 and 6). On average, larvae released during the summer exhibited a higher

potential for dispersal, especially in areas that are not enclosed (Fig. 5f, 6f). It was in

summer that the movement of larvae southward from the Red Sea was more pronounced,

and it was also in this season that many larvae released in the Gulf of Aden reached

Omani waters in the Arabian Sea. In turn, during the winter the vast majority of larvae

originating in the Red Sea were retained within this area, while larvae released within the

Gulf of Aden either moved eastward, though rarely reaching the Omani coast, or moved

southward along the Somali coast.

In the Arabian Sea, surface circulation exhibited a strong seasonal cycle. During

the summer, larvae released along the Omani coast between 54.6˚ E and 56.1˚ E, either

moved eastward alongshore toward the Gulf of Oman or sharply turned toward the open

ocean. Whereas, larvae released between 57.8˚ E and 59˚ E were exclusively transported

alongshore toward the Gulf of Oman (Fig. S5). In the winter, even though the larvae

along the entire Omani coast moved toward offshore, on average they travelled shorter

distances compared to the summer (Fig. 5f, 6f).

On the east side of the Arabian Peninsula, the temporal variability of both the

direction and the distance travelled by the larvae was less evident. In the Gulf of Oman,

larvae released from the eastern Omani coast tended to move southward and few larvae

reached the Arabian Gulf. However, those larvae originating on the top of the Gulf of

Oman were able to pass through the Strait of Hormuz, though they did not travel far

away. In the Arabian Gulf the patterns of dispersal observed in both seasons were quite

similar, larvae tended not to move far away and hence most of them remained in the Gulf.

Discussion

Measuring the strength of barriers has been a great challenge to understand

biogeographic patterns and population genetic structure (Treml et al. 2015). In our study,

a series of individual-based simulations were performed to assess the permeability of

larvae of 72 fish species through putative barriers within the Arabian Peninsula (i.e., the

Bab el-Mandeb Strait, the Strait of Hormuz and the upwelling off Oman). This multitaxa

comparison demonstrated how biological-physical interaction determine the success of

the larvae being transported through the Bab el-Mandeb Strait and the Strait of Hormuz,

as well as of those larvae being self-replenished in the Arabia Sea.

Here we took advantages of a Lagrangian three-dimensional approach to assess

the effects of the hydrographic variability conditions (e.g., extreme events, perturbation,

instability) on larval dispersal, and thus measuring the strength of putative barriers.

Although realistic, the real number of larvae released per species per site greatly exceeds

what is computationally possible for a Lagrangian approach. Alternatively, Eulerian

advection-diffusion methods, that though are not spatially realistic, have a large impact

on dispersal kernel and hence suitable in quantifying evolutionarily significant tails in

evolutionary connectivity studies (Treml et al. 2012).

Bab el-Mandeb Strait

Our results showed that the strength of this barrier is in accordance with the

seasonal water exchange pattern between the Red Sea and the Gulf of Aden, which

changes from a two-layer surface flow in the winter to a three-layer flow in the summer

(Smeed 2004). During the winter season warmer and fresher water from the Gulf of Aden

flows into the Red Sea in the surface layer, while cooler and saltier water from the Red

Sea (RSOW) flows into the Gulf of Aden in the lower layer. The summer regime at the

Bab el Mandab Strait is very different, the flow in the surface layer is reversed while

intermediate water layer from the Gulf of Aden flows into the Red Sea. In turn, the

RSOW forms the deepest and tiny layer (Smeed 2004).

This seasonal variation in the water exchange explains the consequences of

vertical migration and spawning periodicity on the permeability of the Bab el Mandab

showed in the Fig. 2. According to our biophysical model, a taxon from the Gulf of Aden

that has epipelagic larvae moving horizontally and spawn during the summer has more

chance to reach the reefs in the Red Sea. In contrast, a taxon moving in the same

direction will decrease its probability to cross the strait if the spawning occurs in winter

and the larvae perform vertical migration. Under this condition more larvae will be

positioned within the RSOW at deeper layers that flows from the Red Sea to the Gulf of

Aden.

To date, the strength of this barrier has been debated based on the species

distribution and phylogeographic studies. For example, Kemp (1998) suggested that

although the Bab el-Mandeb Strait was a site of a significant Pleistocene vicariance

event, it does not act as a present-day barrier. According to the author, the paucity of

information about the reef fish assemblage inhabiting the south of the Red Sea and the

adjacent Gulf of Aden is the main reason to hypothesize that the strait acts as a barrier. In

contrast, DiBattista et al. (2016a) recorded high endemism rates for a variety of

taxonomic groups within the Red Sea, ranging from 5.5% for scleractinian coral to 16.5%

for ascidians, and suggested that the Red Sea biota is isolated due to the limited water

exchange between the Red Sea and the Gulf of Aden.

Strait of Hormuz

There are few evidences of the Strait of Hormuz acting as a barrier for coral-

dependent fishes. Substantial differences in the structure and composition of reef fish

assemblages between the Arabian Gulf and the Gulf of Oman (Burt et al. 2011), has been

explained by the large spatial gradients in physical conditions (e.g. seawater temperature

and salinity) along these two areas (Smith and Saleh 1987; Feary et al. 2010). Here we

suggest that the spawning strategies and larval traits may also contribute to this

difference.

The relatively high evaporation rate over the Arabian Gulf combined with the

limited water exchange with the open ocean drives an inverse estuarine system at the

Strait of Hormuz, with a shallow inflow of the Indian Ocean Surface Water (IOSW) and a

deep outflow of the Persian Gulf Water (PGW, Swift and Bower 2003). The currents in

the surface layers are highly variable and exhibit large amplitude variation, especially in

wintertime (Johns et al. 2003). The surface inflow of IOSW occurs mainly in the northern

part of the strait, and continues northward along the Iranian coast (Reynolds 1993),

however in the spring the inflow is wider and fills the entire width of the strait (Chao et

al. 1992). On the other hand, the deep outflow is relatively steady throughout year with

little seasonal variance (Johns et al. 2003, Swift and Bower 2003).

The stability of deep-water outflow and the seasonal variability at the surface

corroborates with the results obtained in our simulations (Fig. 3). It explains, for instance,

the fact that the proportion of larvae leaving the Arabian Gulf did not change seasonally,

and increased when the larvae performed vertical migration. From the opposite direction,

epipelagic larvae released in the Gulf of Oman had a higher chance to reach the Arabian

Gulf, especially during the winter when the inflow surface is highly variable.

Upwelling off Oman

Our simulations showed that the proportion of self-recruitment in reefs along the

southern Omani coast was slightly higher in the winter, during the period of no upwelling

event. In this season the larvae moved away from the coast but travelled shorter distances

compared to those larvae released in the summer. Under upwelling conditions, larvae

released between 54.6˚ E and 56.1˚ E were transported to either open sea or eastward

along the coast. This result coincides with the remote sensing images (Weaks 1983) and

empirical data (Elliott and Savidge 1990) of intense upwelling in the vicinities of

headlands off Dhofar and Kuria Muria Island. In this area, coral colonies are weakly

developed and present a reduced number of species (Sheppard and Salm 1988; Burt et al.

2016). The lack of coral reef habitat is also seen as a potential causes faunal and genetic

difference between both sides of the Peninsula (Priest et al. 2015).

Although the effects of offshore transport on coral reef organisms are rarely

investigated (but see Roughan et al. 2011), the effect of offshore Ekman transport on the

larval recruitment has been well documented for small pelagic fishes. These studies have

described that offshore drift does not favor retention in costal areas and act as a hindrance

to larval connectivity (Parrish et al. 1981). The Benguela upwelling system, for example,

acts as a barrier to some species of phytoplankton, copepods and pelagic fishes between

North and South (Lett et al. 2007).

The effect of Ekman transport on the larval dispersal is stronger at the surface

(Lett et al. 2007). Therefore, larvae that performed vertical migration during the summer

presented a higher change to find suitable habitats in coastal areas along the Omani coast,

supporting the hypothesis that vertical migration is a viable retention mechanism for coral

reef fish larvae (Paris and Cowen, 2007). Simulations carried out in the Benguela

upwelling system, also showed that anchovy larvae at the surface travelled longer

distances (Lett et al. 2007). Another biological attribute that was significant to the self-

recruitment under upwelling conditions off Oman was the PLD. Shorter periods in the

planktonic environment enhanced the self-recruitment along the southern Omani coast.

This result reflects the assumption of our biophysical models in which lower PLD also

means early competence period and vertical migration.

The connectivity matrices showed in the southern Arabian Peninsula a hard west-

east barrier for several combinations of biological traits. The position and strength of this

barrier corroborates to all previous results obtained of phylogeography and fauna

composition studies carrier out in the Arabian Peninsula (Kemp 2000; DiBattista et al.

2016a). Specifically, matrices visualization indicates that symmetry and strength of

connectivity largely varied as a function of biological traits (Fig. 4). For example, our

simulations showed that higher connectivity could be achieved with longer duration of

the larval stage, but empirically the importance of PLD to explain biogeographical

patterns and population genetic structure is debatable (Weersing and Toonen 2009;

Selkoe and Toonen 2011). Because the simple notion that the longer a larva spends in the

water column, the higher its potential for dispersal by ocean currents, PLD is a common

proxy for dispersal potential in coral reef fish. This expectation is based on the

assumption that larvae are largely passive particles, however, fish larvae exhibit strong

behavioral preferences including swimming capabilities during their late ontogenetic

stages. In addition, variation in PLD can occur among and within marine taxa, especially

due to larval responses to environmental stochasticity (Woodson and McManus 2007).

Moreover, spawning periodicity was largely affected by oceanographic variability

due to the seasonal monsoon. During the winter counter-clockwise ocean circulation in

the Arabian Sea restricted eastward dispersal, and hence larvae released in the Gulf of

Aden only reached the Omani coast during the summer monsoon. During the summer, in

turn, the ocean circulation in the Arabian Sea restricted the westward dispersal.

Nevertheless, empirical data that support the hypothesis relating symmetry in

connectivity (e.g., genetic studies) along the Arabian Peninsula are lacking.

Conclusion

Our biophysical models complement the existing research on comparative fish

population genetic by providing a snapshot of the present-day seascape permeability

around the Arabian Peninsula. The comparative and cross-taxon models could identify

key biological traits and biophysical interactions that limit the transport of

ichthyoplankton in our study area. Therefore, we could show that the strength of putative

barriers depending on species spawning strategies. The predictions presented here serve

as testable hypotheses for future studies on population genetics and fish larval

distributions around the Arabian Peninsula.

References

References

Ayre, D. J, Minchinton, T. E., and Perrin, C. 2009. Does life history predict past and
current connectivity for rocky intertidal invertebrates across a marine
biogeographic barrier? Mol. Ecol. 18: 1887–1903.
Bakun, A. 1996. Patterns in the ocean, Ocean processes and marine population dynamics.
University of California Sea Grant in cooperation with Centro de Investigaciones
Biologicas de Noroeste.
Baums, L. B., Paris, C. B., and Chérubin, L. M. 2006. A bio-oceanographic filter to larval
dispersal in a reef-building coral. Limnol. Oceanogr. 51: 1969-1981.
Berumen, M. L., DiBattista, J. D., and Rocha, L. A. 2017. Introduction to virtual issue on
Red Sea and Western Indian Ocean biogeography. J. Biogeogr. 44: 1923–1926.
Bowen, B. W., Gaither, M. R., DiBattista, J. D., Iaccheia, M., Andrews, K. R., Grant, W.
S., Toonen, R. J., and Briggs, J. C. 2016. Comparative phylogeography of the
ocean planet. PNAS 113: 7962–7969.
Burt, J. A., Feary, D. A., Bauman, A. G., Usseglio, P., Cavalcante, G. H., Sale, and P. F.
2011. Biogeographic patterns of reef fish community structure in the northeastern
Arabian Peninsula. ICES J. Mar. Sci. 68: 1875–1883.
Burt, J. A., Coles, S., van Lavieren, H., Taylor, O., Looker, E., and Samimi-Namin, K.
2016. Oman's coral reefs: A unique ecosystem challenged by natural and man-
related stresses and in need of conservation. Mar. Pollut. Bull. 105: 498-506.
Carton, X., L'Hegaret, P., and Baraille, R. 2012. Mesoscale variability of water masses in
the Arabian Sea as revealed by ARGO floats. Ocean. Sci. 8: 227–248.
Chao, S-Y., Kao, T. W., and Al-Hajri, K. R. 1992. A numerical investigation of
circulation in the Arabian Gulf. J. Geophys. Res. 11: 219–236.
Cowen, R. K., Lwiza, K. M. M., Sponaugle, S., Paris, C. B., and Olson, D. B. 2000.
Connectivity of Marine Populations: Open or Closed? Science 287: 857-859.
Cowen, R. K. 2002. Larval dispersal and retention and consequences for population
connectivity, p. 149-170. In Sale, P. F. [ed.], Coral reef fishes: dynamics and
diversity in a complex ecosystem. Academic Press.
Cowen, R. K., and Sponaugle, S. 2009. Larval Dispersal and Marine Population
Connectivity. Annu. Rev. Mar. Sci. 1: 443–466.
Cowen, R. K., Paris, C. B., and Srinivasan, A. 2006. Scaling of Connectivity in Marine
Populations. Science 311: 522 – 527.
Cowen, R. K., and Castro, L. R. 1994. Relation of coral reef fish larval distribution to
island scale circulation around barbados, west indies. B. Mar. Sci. 54: 228-244.
Cutler, A. N., and Swallow, J. C. 1984. Surface currents of the Indian Ocean (to 25S,
100E): compiled from historical data archived by the Meteorological Office,
Institute of Oceanographic Sciences.
Cribari-Neto, F., and Zeileis, A. 2010. Beta Regression in R. J. Stat. Softw. 34: 1–24.
DiBattista, J. D., Berumen, M. L., Gaither, M. R., Rocha, L. A., Eble, J. A., Choat, J. H.,
Craig, M. T., Skillings, D. J., and Bowen, B. W. 2013. After continents divide:
comparative phylogeography of reef fishes from the Red Sea and Indian Ocean. J.
Biogeogr. 40: 1170 – 1181.
DiBattista, J. D., Roberts, M. B., Bouwmeester, J., Bowen, B. W., Coker, D. J., Lozano-
Cortes, D. F., Choat, J. H., Gaither, M. R., Hobbs, J. A., Khalil, M. T., Kochzius,
M., Myers, R. F., Paulay, G., Robitzch, V. S. N., Saenz-Agudelo, P., Salas, E.,

Sinclair-Taylor, T. H., Toonen, R. J., Westneat, M. W., Williams, S. T., and
Berumen, M. L. 2016a. A review of contemporary patterns of endemism for
shallow water reef fauna in the Red Sea. J. Biogeogr. 43: 423–439.
DiBattista, J. D., Choat, J. H., Gaither, M. R., Hobbs, J. A., Lozano-Cortes, D. F., Myers,
R. F., Paulay, G., Rocha, L. A., Toonen, R. J., Westneat, M. W., and Berumen, M.
L. 2016b. On the origin of endemic species in the Red Sea. J. Biogeogr. 43: 13–
30.
DiBattista, J. D., Gaither, M. R., Hobbs, J. A., Saenz-Agudelo, P., Piatek, M. J., Bowen,
B. W., Rocha, L. A., Choat, J. H., McIlwain, J. H., Priest, M. A., Sinclair-Taylor,
T. H., and Berumen, M. L. 2017. Comparative phylogeography of reef fishes
from the Gulf of Aden to the Arabian Sea reveals two cryptic lineages. Coral
Reefs 36: 625–638.
Elliott, A. J., and Savidge, G. 1990. Some features of the upwelling off Oman. J. Mar.
Res. 48: 319-333.
Ferrari, S., and Cribari-Neto, F. 2004. Beta Regression for Modelling Rates and
Proportions. J. Appl. Stat. 31: 799-815.
Galarza, J. A., Carreras-Carbonell, J., Macpherson, E., Pascual, M., Roques, S., Turner,
G. F., and Rico, C. 2009. The influence of oceanographic fronts and early-life-
history traits on connectivity among littoral fish species. PNAS 106: 1473-1478.
Griffa, A., Piterbarg, L. I., and Ozgokmen, T. 2004. Predictability of lagrangian particle
trajectories: effects of smoothing of the underlying Eulerian flow. J. Mar. Res. 62:
1–35.
Irisson, J. O., Paris, C. B., Guigand, C., and Planes, S. 2010. Vertical distribution and
ontogenetic ''migration'' in coral reef fish larvae. Limnol. Oceanogr. 55: 909–
919.
Johns, W. E., Yao, F., Olson, D. B. 2003. Observations of seasonal exchange through the
Straits of Hormuz and the inferred heat and freshwater budgets of the Persian
Gulf. J. Geophys. Res. 108: C12, 3391.
Kemp, J. M. 1998. Zoogeography of the coral reef fishes of the Socotra Archipelago. J.
Biogeogr. 25: 919–933.
L'Hégaret, P., Duarte, R., Carton, X., Vic, C., Ciani, D., Baraille, R., and Corréard, S.
2015. Mesoscale variability in the Arabian Sea from HYCOM model results and
observations: impact on the Persian Gulf Water path. Ocean. Sci. 11: 667–693.
Lett, C., Veitch, J., van der Lingen, C. D., and Hutchings, L. 2007. Assessment of an
environmental barrier to transport of ichthyoplankton from the southern to the
northern Benguela ecosystems. Mar. Ecol. Prog. Ser. 347: 247–259.
Lindeman, K. C., Richards, W. J., Lyczkowski-Shultz, J., Drass, D. M., Paris, C. B., Leis,
J. M., Lara, M., and Comyns, B. H. 2005. Lutjanidae: snappers, p. 1549–1586. In
Richards, W. J. [ed.], Early stages of Atlantic fishes. CRC Press.
Nanninga, G. B., Saenz-Agudelo, P., Manica, A., and Berumen, M. L. 2014.
Environmental gradients predict the genetic population structure of a coral reef
fish in the Red Sea. Mol Ecol 23: 591–602.
North, E. W., Gallego, A., and Petitgas, P. 2009. Manual of recommended practices for
modeling physical–biological interactions during fish early life, ICES
Cooperative Research Report.
Okubo, A. 1971. Oceanic diffusion diagrams. Deep-Sea Res. 18: 789-802.

Paris, C. B., and Cowen, R. K. 2004. Direct evidence of a biophysical retention
mechanism for coral reef fish larvae. Limnol. Oceanogr. 49: 1964–1979.
Paris, C. B., Helgers, J., Sebille, E., and Srinivasan, A. 2013. Connectivity Modeling
System: A probabilistic modeling tool for the multi-scale tracking of biotic and
abiotic variability in the ocean. Environ. Modell. Softw. 42: 47-54.
Parrish, H. R., Nelson, C. S., and Bakun, A. 1981. Transport mechanisms and
reproductive success of fishes in the California current. Biolog. Oceanog. 1: 175-
203.
Priest, M. A., DiBattista, J. D., McIlwain, J. L., Taylor, B. M., Hussey, N. E., and
Berumen, M. L. 2016. A bridge too far: dispersal barriers and cryptic speciation
in an Arabian Peninsula grouper (*Cephalopholis hemistiktos*). J. Biogeogr. 43:
820–832.
Reynolds, R. M. 1993. Physical oceanography of the Persian Gulf, Strait of Hormuz, and
the Gulf of Oman – Results from the Mt. Mitchell Expedition. Mar. Pollut. Bull.
27: 35-59.
Rocha, L. A. 2003. Patterns of distribution and processes of speciation in Brazilian reef
fishes. J. Biogeogr. 30: 1161–1171.
Ross, R. E., Nimmo-Smith, W. A. M., and Howell, K. L. 2016. Increasing the Depth of
Current Understanding: Sensitivity Testing of Deep-Sea Larval Dispersal Models
for Ecologists. PLoS ONE 11: e0161220.
Roughan, M., Macdonald, H. S., Baird, M. E., and Glasby, T. M. 2011. Modelling coastal
connectivity in a Western Boundary Current: Seasonal and inter-annual
variability. Deep-Sea Res II. 58: 628-644.
Saenz-Agudelo, P., DiBattista, J. D., Piatek, M. J., Gaither, M. R., Harrison, H. B.,
Nanninga, G. B., and Berumen, M. L. 2015. Seascape genetics along
environmental gradients in the Arabian Peninsula: insights from ddRAD
sequencing of anemonefishes. Mol. Ecol. 24: 6241–6255.
Selkoe, K. A., and Toonen, R. J. 2011. Marine connectivity: a new look at pelagic larval
duration and genetic metrics of dispersal. Mar. Ecol. Prog. Ser. 436: 291-305.
Sheppard, C. R. C., and Salm, R. V. 1988. Reef and coral communities of Oman, with a
description of a new coral species (Order Scleractinia, genus Acanthastrea). J.
Nat. Hist. 22: 263-279.
Shetye, S. R., Gouveia, A. D., and Shenoi, S. S. C. 1994. Circulation and water masses of
the Arabian Sea. P. Indian. Acad. Sci. 103: 107-123.
Simons, R. D., Siegel, D. A., and Brown, K. S. 2013. Model sensitivity and robustness in
the estimation of larval transport: A study of particle tracking parameters. J. Mar.
Syst. 119–120: 19–29.
Smeed, D. A. 2004. Exchange through the Bab el Mandab. Deep-Sea Res. II 51: 455–
474.
Swift, S. A., and Bower, A. S. 2003. Formation and circulation of dense water in the
Persian/Arabian Gulf. J. Geophys. Res. 108: C1 3004.
Thresher, R. E., and Brothers, E. B. 1985. Reproductive ecology and biogeography of
Indo-West Pacific angelfishes. Evolution 39: 878-887.
Torquato, F. O., Range, P., Ben-Hamadou, R., Sigsgaard, E. E., Thomsen, P. F., Riera,
R., Berumen, M. L., Burt, J. A., Feary, D. A., Marshell, A., D'Agostino, D.,
DiBattista, J. D. and Møller, P. R. 2019. Consequences of marine barriers on

genetic diversity of the coral-specialist yellowbar angelfish from the western
Indian Ocean. Ecol. & Evol. 9: 11215-11226.

Treml, E. A., Roberts, J. J., Chao, Y., Halpin, P. N., Possingham, H. P. and Riginos, C.
2012. Reproductive Output and Duration of the Pelagic Larval Stage Determine
Seascape-Wide Connectivity of Marine Populations. Integr. Comp. Biol. 52: 525–
537.

Treml, E. A., Roberts, J., Halpin, P. N., Possingham, H. P., and Riginos, C. 2015. The
emergent geography of biophysical dispersal barriers across the Indo-West
Pacific. Divers. Distrib. 21: 465–476.

UNEP-WCMC, WorldFish Centre, WRI, TNC (2010). Global distribution of warm-water
coral reefs, compiled from multiple sources including the Millennium Coral Reef
Mapping Project. Version 3.0. Includes contributions from IMaRS-USF and IRD
(2005), IMaRS-USF (2005) and Spalding et al. (2001). UN Environment World
Conservation Monitoring Centre, Cambridge/UK. URL: http://data.unep-
wcmc.org/datasets/1.

Weaks, M. L. 1983. Satellite image of the month, Arabian Sea upwelling. Oceanogr.
Month. Summ. 3: 13.

Weersing, K., and Toonen, R. J. 2009. Population genetics, larval dispersal, and
connectivity in marine systems. Mar. Ecol. Progr. Ser. 393: 1–12.

Woodson, C. B., and McManus, MA (2007) Foraging behavior can influence dispersal of
marine organisms. Limnol Oceanogr 52: 2701 – 2709.

Yao, F., and Johns, W. E. 2010. A HYCOM modeling study of the Persian Gulf: 1.
Model configurations and surface circulation. J. Geophys. Res. 115: 1-17.

Zuur, A. F., Ieno, E. N., and Elphick, C. S. 2010. A protocol for data exploration to avoid
common statistical problems. Methods Ecol. Evol. 1: 3-14.

649
650

Acknowledgements

The author F.T. is supported by a CNPq/Brazil fellowship through the programme

Science Without Borders (Proc. 232875/2014-6). I also thank Peter Rask Møller

(University of Copenhagen) for proving infrastructure, as well as Dr Robert Cowen

(Oregon University) and Dr Luiz Rocha (California Academy of Sciences) for their

helpful comments.

TABLES

Table 1. Range in biological parameter values to characterize the 72 hypothetical model taxa.

PLD (days)	Spawning periodicity	Larval mortality	Vertical migration	Productivity
20	Annual	High	Yes	High
30	Summer)	Low	No	Low
40	Winter)			

FIGURES

Legends

Fig. 1. Study region. Northwestern Indian Ocean with previously described barriers depicted shown as shaded line. (a) Bab-el-Mandeb Strait, (b) Upwelling off Oman and (c) Strait of Hormuz.

Fig. 2. Results on the beta regression models showing the effects of the biological traits on the permeability of the Bab-el-Mandeb Strait. The red triangles represent the mean value. Different letters indicate signicant differences ($p < 0.05$).

Fig. 3. Results on the beta regression models showing the effects of the biological traits on the permeability of the Strait of Hormuz. The red triangles represent the mean value. Different letters indicate signicant differences ($p < 0.05$).

Fig. 4. Results on the beta regression models showing the effects of the biological traits on the self-recruitment along Omani coast. The red triangles represent the mean value. Different letters indicate significant differences ($p < 0.05$).

Fig. 5. Modeled dispersal paths of epipelagic virtual larvae with larval duration of 40 days released in the wintertime from (a) the Red Sea, (b) Gulf of Aden, (c) Arabian Sea, (d) Gulf of Oman and (e) Arabian Gulf. One larva was released from each habitat cell (total of 181) per day in 2015. The distances travelled by the larvae are represented in (f).

Fig. 6. Modeled dispersal paths of epipelagic virtual larvae with larval duration of 40 days released in the summertime from (a) the Red Sea, (b) Gulf of Aden, (c) Arabian Sea, (d) Gulf of Oman and (e) Arabian Gulf. One larva was released from each habitat cell (total of 181) per day in 2015. The distances travelled by the larvae are represented in (f).

704   **Fig. 1.**

Bab-el-Mandeb Strait

Strait of Hormuz

771   **Fig. 4.**
772

Upwelling off Oman

790   **Fig. 5.**
791

792
793
794

795   **Fig. 6.**
796