Evaluation of DNA extraction protocols from liquid-based cytology specimens for studying cervical microbiota

Takeo Shibata,[a,b] Mayumi Nakagawa,[a] Hannah N. Coleman,[a] Sarah M. Owens,[c] William W. Greenfield,[d] Toshiyuki Sasagawa,[b] Michael S. Robeson II[e]

[a]Department of Pathology, University of Arkansas for Medical Sciences, Little Rock, AR, USA

[b]Department of Obstetrics and Gynecology, Kanazawa Medical University, Uchinada, Ishikawa, Japan

[c]Biosciences Division, Argonne National Laboratory, Lemont, IL, USA

[d]Department of Obstetrics and Gynecology, University of Arkansas for Medical Sciences, Little Rock, AR, USA

[e]Department of Biomedical Informatics, University of Arkansas for Medical Sciences, Little Rock, AR, USA

Takeo Shibata: TShibata@uams.edu; Mayumi Nakagawa: MNakagawa@uams.edu; Hannah N. Coleman: ColemanHannahN@uams.edu; Sarah M. Owens: sarah.owens@anl.gov; William W. Greenfield: GreenfieldWilliamW@uams.edu; Toshiyuki Sasagawa: tsasa@kanazawa-med.ac.jp; Michael S. Robeson II: MRobeson@uams.edu.

Corresponding author: Michael S. Robeson II

Tel: 501-526-4242, Fax: 501-526-5964

Email: MRobeson@uams.edu

Abstract

Background: Cervical microbiota (CM) are considered an important factor affecting the

progression of cervical intraepithelial neoplasia (CIN) and are implicated in the persistence of

human papillomavirus (HPV). Collection of liquid-based cytology (LBC) samples is routine for

cervical cancer screening and HPV genotyping, and can be used for long-term cytological

biobanking. Herein, we investigate the feasibility of leveraging LBC specimens for use in CM

surveys by amplicon sequencing. As methodological differences in DNA extraction protocols

can potentially bias the composition of microbiota, we set out to determine the performance of

four commonly used DNA extraction kits (ZymoBIOMICS DNA Miniprep Kit; QIAamp

PowerFecal Pro DNA Kit; QIAamp DNA Mini Kit; and IndiSpin Pathogen Kit) and their ability

to capture the diversity of CM from LBC specimens.

Results: LBC specimens from 20 patients (stored for 716 ± 105 days) with cervical

intraepithelial neoplasia (CIN) 2/3 or suspected CIN2/3 were each aliquoted for the four kits. We

observed that, regardless of the extraction protocol used, all kits provided equivalent accessibility

to the cervical microbiome, with some minor differences. For example, the ZymoBIOMICS kit

appeared to differentially increase access of several microbiota compared to the other kits.

Potential kit contaminants were observed as well. Approximately 80% microbial genera were

shared among all DNA extraction protocols. The variance of microbial composition per

individual was larger than that of the DNA extraction protocol used. We also observed that

HPV16 infection was significantly associated with community types that were not dominated by

Lactobacillus iners.

Conclusions: Collection of LBC specimens is routine for cervical cancer screening and HPV

genotyping, and can be used for long-term cytological biobanking. We demonstrated that LBC

samples, which had been under prolonged storage prior to DNA extraction, were able to provide

a robust assessment of the CM and its relationship to HPV status, regardless of the extraction kit

used. Being able to retroactively access the CM from biobanked LBC samples, will allow

researchers to better interrogate historical interactions between the CM and its relationship to

CIN and HPV. This alone has the potential to bring CM research one-step closer to the clinical

practice.

Keywords; cervical microbiota, DNA extraction, HPV, CIN, liquid-based cytology

Background

High-throughput sequencing (HTS) technology of 16S rRNA gene amplicon sequences has made

it possible to better understand the relationships between cervicovaginal microbiota and human

papillomavirus (HPV) infection [1] [2] [3] [4] [5] and HPV-related diseases [6] [7] [8] [9] [10].

Cervicovaginal microbiota are considered to be an important factor affecting the progress of

cervical intraepithelial neoplasia (CIN) [6] [7] [8] [9] and are implicated in the persistence of

high-risk HPV (HR-HPV) [1] [2] and low-risk HPV (LR-HPV) [3]. For example, the phyla

Actinobacteria and *Fusobacteria* were enriched in HR-HPV positive environment [4] while the

phyla *Actinobacteria*, *Proteobacteria*, and *Fusobacteria* in low-risk HPV (LR-HPV) [3].

Additionally, *Lactobacillus iners*-dominant samples are associated with both HR-HPV and LR-

HPV [5]. Moreover, it has been shown that CIN risk was increased when the cervical microbes

Atopobium vaginae, *Gardnerella vaginalis*, and *Lactobacillus iners* were present with HR-HPV

[10]. The cervicovaginal microbiome specified by *Lactobacillus*-dominant type or non-

Lactobacillus-dominant type has been shown to interact with the immune system [7] [11].

Inflammatory cytokines, such as Interleukin (IL)-1α and IL-18, were increased in non-

Lactobacillus-dominant community types of reproductive-aged healthy women [11]. In the

analysis of patients with cervical cancer, non-*Lactobacillus*-dominant community types were

positively associated with chemokines such as interferon gamma-induced protein 10 (IP-10) and

soluble CD40-ligand activating dendric cells (DCs) [7]. The metabolism of the cervicovaginal

microbiome may be a substantial contributing factor to maternal health during pregnancy,

although the mechanism is still unclear [12].

Little has been reported on the utility of liquid-based cytology (LBC) samples for use in

cervical microbiome studies. Conventionally, microbiome sample collection methods entail the

use of swabs [13] or self-collection of vaginal discharge [14]. To obtain a non-biased and broad

range of cervical microbiota, DNA extraction should be optimized for a range of difficult-to-

lyse-bacteria, *e.g.* *Firmicutes*, *Actinobacteria*, and *Lactobacillus* [13] [15] [16] [17] [18].

LBC samples are promising for cervicovaginal microbiome surveys, as they are an

already established method of long-term cytological biobanking [19]. In clinical practice,

cervical cytology for cervical cancer screening or HPV genotyping is widely performed using a

combination of cervical cytobrushes and LBC samples such as ThinPrep (HOLOGIC) or

SurePath (BD). An LBC specimen can be used for not only cytological diagnosis but also

additional diagnostic tests such as HPV, *Chlamydia*, *Neisseria gonorrhoeae*, and *Trichomonas*

infection [20] [21] [22].

The ability to characterize microbial communities, as commonly assessed by 16S rRNA

gene sequencing, can be biased as a result of methodological differences of cell lysis and DNA

extraction protocols [23] [24] [25]. Herein, we compare four different commercially available

DNA extraction kits in an effort to assess their ability to characterize the cervical microbiota of

LBC samples. Additionally, we examine the relationship between HPV infection and the

composition of cervical microbiota.

Results

Patient characteristics

The age of the patients (n = 20) was 31.4 ± 5.0 years. The distribution of race was 15% African

American (n = 3), 50% Caucasian (n = 10), and 35% Hispanic (n = 7). Cervical histology was

40% CIN2 (n = 8), 50% CIN3 (n = 10), and 10% benign (n = 2). HPV genotypes were 50%

HPV16 positive (n = 10), 10% HPV18 positive (n = 2), and 90% HR-HPV positives (n = 18).

Patient characteristics were summarized in Table 2.

DNA yield

DNA yield per 100 μL ThinPrep solution was 0.09 ± 0.06 μg in ZymoBIOMICS, 0.04 ± 0.01 μg

in PowerFecalPro, and 0.21 ± 0.23 μg in QIAampMini. DNA yield was not calculated for

IndiSpin, as Poly-A Carrier DNA was used. The DNA yield of PowerFecalPro was significantly

lower than that of ZymoBIOMICS (adjusted p value < 0.001) and QIAampMini (adjusted p

value < 0.001) based on Dunn's test with Benjamini-Hochberg-adjustment (Figure S1).

Number of reads and Operational Taxonomic Units (OTUs) before rarefying

We obtained a total of 11,149,582 reads for 80 DNA extractions. A positive control of mock

sample produced 127,142 reads and ThinPrep solution as the negative control produced 1,773

reads. IndiSpin (168,349 ± 57,451 reads) produced a significantly higher number of reads

compared to PowerFecalPro (115,610 ± 68,201 reads, p value = 0.020, Dunn's test with

Benjamini-Hochberg-adjustment) as shown in Table 3. Approximately 90% of reads were

assigned to gram-positive bacteria and about 10% of reads were assigned to gram-negative

bacteria across all kits.

Prior to rarefying, the ZymoBIOMICS kit captured a greater representation of gram-

negative bacterial OTUs (total 346, 17.3 ± 9.8) compared to PowerFecalPro (total 209, 10.5 ±

10.3, p value = 0.012, Dunn's test with Benjamini-Hochberg-adjustment, ratio of gram-negative

bacteria: 41.9% vs 33.7%) as shown in Table 3. No significant differences in the number of

OTUs before rarefying was detected for the entire bacterial community or gram-positive

bacteria.

Microbiome composition per DNA extraction protocol

We analyzed whether differences in DNA extraction methods affect our ability to assess cervical

microbiota composition. The patients can be identified by whether or not they displayed a

Lactobacillus-dominant community type (Figure 2A). Variation between individuals was a

significantly greater influence on the observed microbial composition than was the method of

DNA extraction (Figure 2A).

The following top 10 abundant families are shown in Figure 2A (left) and constituted

approximately 95.7% of cervical bacteria in all kits (80 DNA extractions); *Lactobacillaceae*

(58.9%), *Bifidobacteriaceae* (13.7%), *Veillonellaceae* (4.8%), *Prevotellaceae* (4.3%), *Family XI*

(3.9%), *Atopobiaceae* (3.0%), *Leptotrichiaceae* (2.5%), *Streptococcaceae* (2.0%),

Lachnospiraceae (1.6%). *Ruminococcaceae* (0.9%). The following top 10 abundant genera are

shown in Figure 2A (right) and constituted approximately 92% of cervical bacteria;

Lactobacillus (58.9%), *Gardnerella* (13.6%), *Prevotella* (4.2%), *Megasphaera* (3.7%),

Atopobium (3.0%), *Sneathia* (2.5%), *Streptococcus* (1.9%), *Parvimonas* (1.7%), *Shuttleworthia*

(1.4%), and *Anaerococcus* (1.1%).

Shared and unique microbiota among DNA extraction protocols

All DNA extraction methods were generally commensurate with one another, there were 31 of 41 shared microbes at the family level (Figure 2B left) and 45 of 57 shared microbes at the genus level (Figure 2B right) among the DNA extraction protocols.

However, four gram-negative taxa were uniquely detected by ZymoBIOMICS and one taxon was uniquely detected by QIAampMini both at the genus level (Figure 2B right). Of the uniquely detected ZymoBIOMICS OTUs, *Methylobacterium* was detected in 5 of the 80 DNA extractions, consisting of 912 reads; 0.01% of all kit extractions. A member of this genus, *Methylobacterium aerolatum*, has been reported to be more abundant in the endocervix than the vagina of healthy South African women [26]. *Bacteroidetes*, which are often reported as enriched taxa in an HIV positive cervical environment [27], was detected in 12 of the 80 DNA extractions (1,028 reads; 0.01%). *Meiothermus* was detected in 9 of the 80 DNA extractions (882 reads; 0.01%) and *Hydrogenophilus* was detected in 14 of the 80 DNA extractions (2,488 reads, 0.02%). *Meiothermus* and *Hydrogenophilus* [28] are not considered to reside within the human environment, and are likely kit contaminants, as previously reported [29]. A unique gram-positive taxa obtained from the QIAampMini, *Streptomyces*, which was reported to be detected from the cervicovaginal environment in the study of Kenyan women [30], was detected in 20 of 80 DNA extractions (6,862 reads; 0.06%). No unique taxa were detected in PowerFecalPro and IndiSpin. Although less than 0.005% of the total data set, two samples of IndiSpin also detected potential kit contaminant, *Tepidiphilus* (*Hydrogenophilaceae*).

Venn diagrams at family levels also exhibited that ZymoBIOMICS detected slightly more bacterial taxa (four unique taxa) as shown in Figure 2B (left). These results showed that

major bacteria were commonly detected among all extraction protocols, with only slightly more

uniquely detected microbiota using ZymoBIOMICS.

Alpha and beta diversity

Significantly higher Species richness (`q2-breakaway`) was observed from the

ZymoBIOMICS (56.1 ± 19.4) protocol compared to that of PowerFecalPro (43.2 ± 32.9, p =

0.025), QIAampMini (54.9 ± 29.8, not significant), and IndiSpin (63.6 ± 38.3, not significant)

using Dunn's test with Benjamini-Hochberg-adjustment (Figure 3). Similarly, Faith's

Phylogenetic Diversity was observed to be higher with the ZymoBIOMICS protocol (6.6 ± 2.2),

compared to PowerFecalPro (4.5 ± 1.9, p = 0.012), QIAampMini (5.0 ± 1.8, not significant), and

IndiSpin (5.4 ± 1.7, not significant) using Dunn's test with Benjamini-Hochberg-adjustment

(Figure 3). The use of IndiSpin also resulted significantly higher alpha diversity than that of

PowerFecalPro in an analysis of Species richness (p = 0.042, Dunn's test with Benjamini-

Hochberg-adjustment). Non-phylogenetic alpha diversity metrics such as Observed OTUs,

Shannon's diversity index, and Pielou's Evenness did not show differences among the four

methods.

178   ZymoBIOMICS was able to significantly increase access to several taxonomic groups

compared to the other DNA extraction methods. Additionally, as shown in Table 4,

ZymoBIOMICS did capture a different microbial composition compared to other DNA

extraction methods in the index of Unweighted UniFrac distances (PowerFecalPro: q = 0.002;

QIAampMini: q = 0.002; and IndiSpin: q = 0.002) and in Jaccard distances (QIAampMini: q =

0.018 and IndiSpin: q = 0.033).

Differential accessibility of microbiota by DNA extraction protocol

Linear discriminant analysis (LDA) effect size (LEfSe) analysis [31] identified

taxonomic groups, defined with an LDA score of 2 or higher, for differential accessibility by

extraction kit: 23 in ZymoBIOMICS, 0 in PowerFecalPro, 3 in QIAampMini, and 3 in IndiSpin

(Figure 4A). The following taxa were found to be highly accessible (LDA score > 3) with the use

of the ZymoBIOMICS kit: Phylum *Proteobacteria*, Class *Gammaproteobacteria*, Order

Betaproteobacteriales, Family *Bacillaceae*, and Genus *Anoxybacillus*. Whereas the Order

Streptomycetales was highly enriched with the use of the QIAampMini (LDA score > 3). As

shown in the cladogram (Figure 4B), despite the detection of a potential kit contaminants

(*Meiothermus, Hydrogenophilaceae, and Hydrogenophilus*), ZymoBIOMICS was able to

increase the accessibility to additional microbiota compared to the other extraction protocols.

Microbial community type and HPV16

Dirichlet Multinomial Mixtures (DMM) model [32] detected two cervical microbial community

types across all four DNA extraction protocols (Figure S2). Community type I was composed of

the following: *Gardnerella sp.* (ZymoBIOMICS: 17.1%; PowerFecalPro: 20%; QIAampMini:

23%; IndiSpin: 20%), *Lactobacillus iners* (ZymoBIOMICS: 6.3%; PowerFecalPro: 5%;

QIAampMini: 6%; IndiSpin: 5%), *Atopobium vaginae* [10] (ZymoBIOMICS: 3.5%;

PowerFecalPro: 3%; QIAampMini: 4%; IndiSpin: 5%), *Clamydia trachomatis* (ZymoBIOMICS:

1.9%; PowerFecalPro: 2%; QIAampMini: 3%; IndiSpin: 2%), *Shuttleworthia sp.*

(ZymoBIOMICS: 1.8%; PowerFecalPro: 2%; QIAampMini: 2%; IndiSpin: 2%). Some members

of *Shuttleworthia* are considered to be bacterial vaginosis-associated bacterium (BVAB) [33],

further investigation is required to determine if this OTU is indeed a BVAB. We determined this

community type "high diversity type". Community type II was is dominated by *Lactobacillus*

iners at 88%, 85%, 83%, and 85% respectively for ZymoBIOMICS, PowerFecalPro,

QIAampMini, and IndiSpin.

The relationship between HPV16 infection and community type was observed to be

significantly associated with community type I (HPV16 positive patients [n = 9], HPV16

negative patients [n = 1]) and not community type II (HPV16 positive patients [n = 1], HPV16

negative patients [n = 9], p = 0.001, Fisher's exact test) regardless of the DNA extraction kit used

(Figure S2A). In support of this result, analysis of differentially abundant microbiota using `q2-`

`aldex` (Benjamini-Hochberg corrected p value of Wilcoxon test: p < 0.001, standardized

distributional effect size: −1.2) revealed that *Lactobacillus iners* were differentially enriched in

the cervical environment without HPV16. LEfSe analysis also detected that genus *Lactobacillus*

were enriched in the cervical environment without HPV16 (p < 0.001, LDA score: 5.38, Figure

S2B). No significant differences were observed in the relationship between community type and

HPV18 (p = 0.474, Fisher's exact test), HR-HPV (p = 0.474, Fisher's exact test), results of

cervical biopsy (p = 0.554, Fisher's exact test), and race (African Americans vs not-African

Americans: p = 1; Caucasian vs not-Caucasian: p = 0.656; Hispanic vs not-Hispanic: p = 0.350,

Fisher's exact test, Figure S2A).

Discussion

In this study, we evaluated the utility of LBC specimens for the collection and storage of cervical

samples for microbiome surveys based on the 16S rRNA marker gene. We simultaneously

compared the efficacy of several commonly used DNA extraction protocols on these samples in

an effort to develop a standard operating procedure/protocol (SOP) for such work. We've also

been able to show that there are two cervical microbial community types, which are associated

with the dominance or non-dominance of *Lactobacillis iners* and HPV16 status. The relationship

between community types and HPV16 was detected regardless of the DNA extraction protocol

used.

This study evaluated the composition of microbiota across all DNA extraction methods.

These findings document the importance of selecting DNA extraction methods in cervical

microbiome studies from the LBC samples. All kits were commensurate in their ability to

capture the microbial composition of each patient and the two observed cervical microbial

community state types, making all of these protocols viable for discovering broad patterns of

microbial diversity. However, we did observe that the ZymoBIOMICS protocol was better able

to access additional cervical microbiota (Figure 2B, 4A & B). Coincidentally, we detected

potential DNA contamination with the ZymoBIOMICS and IndiSpin kits. The number of OTUs

prior to rarefying revealed that the ZymoBIOMICS protocol detected more gram-negative OTUs

than the PowerFecalPro (Table 3 & Figure 2B). In particular, LEfSe analysis has shown that

phylum *Proteobacteria* can be better detected with the ZymoBIOMICS kit (Figure 4).

Although rarefying microbiome data can be problematic [34], it can still provide robust

and interpretable results for diversity analysis [35], we were able to observe commensurate

findings with non-rarefying approaches such as `q2-breakaway` [36], `q2-deicode` [37], and

LEfSe [31]. Beta-diversity analysis via Unweighted UniFrac also revealed that ZymoBIOMICS

was significantly different from all other kits. There were no differences in non-phylogenic

indices of alpha diversity with rarefying approaches. These findings lead us to surmise that

phylogenetic indices may be more sensitive than the non-phylogenetic indices.

Although we hypothesized that the detection of difficult-to-lyse-bacteria (*e.g.* gram-

positive bacteria) would vary by kit, we observed no significant differences (Table 3). As shown

in Table 3, the number of reads of gram-positive and gram-negative bacteria also showed that

there was no difference in the four kits. This is likely due to several modifications made to the

extraction protocol as outlined in Table 1. That is, we added bead beating and mutanolysin to the

QIAampMini protocol [38]. We also modified the beating time of the ZymoBIOMICS kit down

to 2 minutes from 10 minutes (the latter being recommended by the manufacturer) to minimize

DNA shearing. We may use the extracted DNA from ZymoBIOMICS for long-read amplicon

sequencing platforms such as PacBio (Pacific Biosciences of California, Inc) [39] or MinION

(Oxford Nanopore Technologies) [40] [41]. Excessive shearing can render these samples

unusable for long-read sequencing. It is quite possible that we could have observed even more

diversity with the ZymoBIOMICS kit for our amplicon survey if we conducted bead-beating for

the full 10 minutes.

Community typing and detection of the differentially abundant microbiota revealed that

Lactobacillus iners were more abundant in the cervical ecosystem without HPV16. These

findings are congruent with those of, Usyk *et al*. [42], Lee *et al*. [1], and Audirac-Chalifour *et al*.

[43]. Usyk *et al*., reported that *L. iners* was associated with clearance of HR-HPV infections

[42]. Lee *et al*. reported that *L. iners* were decreased in HPV positive women [1]. Also, the

results indicated that the proportion of *L. iners* was higher in HPV-negative women compared to

HPV-positive women (relative abundance 14.9% vs 2.1%) was reported by Audirac-Chalifour *et*

al [43]. Similarly, Tuominen *et al*. [18] reported that *L. iners* were enriched in HPV negative

samples (relative abundance: 47.7%) compared to HPV positive samples (relative abundance:

18.6%, p value = 0.07) in the study of HPV positive-pregnant women (HPV16 positive rate:

15%) [44]. As established by the seminal study of Ranjeva *et al*. [45], a statistical model

revealed that colonization of specific HPV types including multi HPV type infection depends on

host-risk factors such as sexual behavior, race and ethnicity, and smoking. It is unclear whether

the association between the cervical microbiome, host-specific traits, and persistent infection of

specific HPV types, such as HPV16, can be generalized and requires further investigation.

We focused on LBC samples as this is the recommended method of storage for cervical

cytology [46]. Here, we confirmed that LBC samples can be used for microbial community

surveys by simply using the remaining LBC solution post HPV testing or cervical cytology. We

used a sample volume of 200 or 300 μL ThinPrep solution in this study. The Linear Array HPV

Genotyping Test (Roche Diagnostics) stably detects β-globin with a base length of 268 bp as a

positive control. Therefore, using a similar sample volume as HPV genotyping (250 μL), it was

expected that V4 (250 bp), which is near the base length of β-globin, would be PCR amplified. It

has been pointed out by Ling *et al*. [47] that the cervical environment is of low microbial

biomass. To control reagent DNA contamination and estimate the sample volume, DNA

quantification by qPCR before sequencing is recommended [48]. Mitra *et al* determined a sample

volume of 500 μL for ThinPrep by qPCR in the cervical microbiome study comparing sampling

methods using cytobrush or swab [19]. The average storage period from sample collection via

LBC to DNA extraction was about two years in this study. Kim *et al*. reported that DNA from

the cervix stored in ThinPrep at room temperature or −80°C was stable for at least one year [49].

Meanwhile, Castle *et al*. reported that β-globin DNA fragments of 268 bases or more were

detected by PCR in 90 % (27 of 30 samples) of ThinPrep samples stored for eight years at an

uncontrolled ambient temperature followed by a controlled ambient environment (10–26.7°C)

[50]. Low-temperature storage may allow the analysis of the short DNA fragments of the V4

region after even long-term storage, although further research is needed to confirm the optimal

storage period in cervical microbiome studies using ThinPrep. SurePath LBC specimens are as

widely used as ThinPrep, but the presence of formaldehyde within the SurePath preservation

solution raises concerns about accessing enough DNA for analysis as compared to ThinPrep,

which contains methanol [51] [52]. It should also be noted that other storage solutions, *i.e.* those

using guanidine thiocyanate have been reported for microbiome surveys of the cervix [53] and

feces [54]. A weakness of the current study is that we did not examine the reproducibility of our

results as each sample was extracted using each kit once. However, the use of actual patient

samples rather than mock samples is a strength of our approach.

Conclusions

In conclusion, regardless of the extraction protocol used, all kits provided equivalent broad

accessibility to the cervical microbiome. Observed differences in microbial composition were

due to the significant influence of the individual patient and not the extraction protocol.

However, ZymoBIOMICS was observed to increase the accessibility of DNA from a greater

range of microbiota compared to the other kits, in that the greatest number of significantly

enriched taxa were identified. This was not because of higher DNA yield nor ability to detect

more gram-positive bacteria. We have shown that the ability to characterize cervical microbiota

from LBC specimens is robust, even after prolonged storage. Our data also suggest that it is

possible to reliably assess the relationship between HPV and the cervical microbiome, also

supported by Kim *et al*. [49] and Castle *et al* [50]. Cervical microbiome in patients with HPV16

or HPV18 which causes 70% of cervical cancers and CIN [55] warrants critical future study.

Selection and characterization of appropriate DNA extraction methods are important for

providing an accurate census of cervical microbiota and the human microbiome in general [23]

[24] [25] [38] [49] [50]. Even though we found all four extraction kits to be commensurate in

their ability to broadly characterize the CM, this study lends support to the view that the

selection of a DNA extraction kit depends on the questions asked of the data, and should be

taken into account for any cervicovaginal microbiome and HPV research that leverages LBC

specimens for use in clinical practice [15] [56].

Methods

Sampling of cervical microbiome

LBC specimens were obtained from 20 patients enrolled in a Phase II clinical trial of an HPV

therapeutic vaccine (NCT02481414). In order to be eligible, participants had to have high grade

squamous intraepithelial lesions (HSILs) or cannot rule out HSILs in cervical cytology or

CIN2/3 in cervical biopsy. Those who qualified for the study based on their cervical cytology

underwent cervical biopsy, and they qualified for vaccination if the results were CIN2/3. The

cervical cytology specimens in this current study were collected before the vaccination and

reserved in the vial of the ThinPrep Pap Test (HOLOGIC) as described in Ravilla *et al*. 2019

[57]. The storage period from sample collection to DNA extraction was 716 ± 105 days in this

study.

HPV genotyping

HPV-DNA was detected by Linear Array HPV Genotyping Test (Roche Diagnostics) which can

detect up to 37 HPV genotypes including 13 HR-HPV genotypes (16, 18, 31, 33, 35, 39, 45, 51,

52, 56, 58, 59, and 68) and 24 LR-HPV genotypes (6, 11, 26, 40, 42, 53, 54, 55, 61, 62, 64, 66,

67, 69, 70, 71, 72, 73, 81, 82, 83, 84, IS39, and CP6108) using ThinPrep solution [58].

DNA extraction protocols

We selected four commercially available DNA extraction kits as the candidates for comparison:

ZymoBIOMICS DNA Miniprep Kit (Zymo Research, D4300), QIAamp PowerFecal Pro DNA

Kit (QIAGEN, 51804), QIAamp DNA Mini Kit (QIAGEN, 51304), and IndiSpin Pathogen Kit

(Indical Bioscience, SPS4104). These kits have been successfully used in a variety of human

cervical, vaginal, and gut microbiome surveys [10] [19] [59]. We'll subsequently refer to each of

these kits in abbreviated form as follows: ZymoBIOMICS, PowerFecalPro, QIAampMini, and

IndiSpin. The protocols and any modifications are outlined in Table 1.

Each LBC sample was dispensed into four separate 2 mL sterile collection tubes

(dispensed sample volume = 500 μL) to create four cohorts of 20 DNA extractions (Figure 1).

Each extraction cohort was processed through one of the four kits above. A total of 80

extractions (4 kits × 20 patients) were prepared for subsequent analyses. Applied sample volume

of ThinPrep solution was 300 μL for ZymoBIOMICS, 300 μL for PowerFecalPro, 200 μL for

QIAampMini, and 300 μL for IndiSpin. The sample volume was standardized to 300 μL as long

as the manufacturer's instructions allowed to do so. DNA extraction for all samples was

performed by the same individual who practiced by performing multiple extractions for each kit

before performing the actual DNA extraction on the samples analyzed in this study. Positive

control was mock vaginal microbial communities composed of a mixture of genomic DNA from

the American Type Culture Collection (ATCC MSA1007). Negative control was the ThinPrep

preservation solution without the sample as blank extraction [60].

Measurement of DNA yield

DNA yield for each method was evaluated by spectrophotometer (Nanodrop One, Thermo

Scientific). Analysis of the DNA yield from IndiSpin was omitted as nucleic acid is used as a

carrier for this kit. The mean DNA yields per 100 μL ThinPrep sample volume were compared.

16S rRNA marker gene sequencing

Controls and the extracted DNA were sent to Argonne National Laboratory (IL, USA) for

amplification and sequencing of the 16S rRNA gene on an Illumina MiSeq sequencing platform.

Paired-end reads from libraries with ~250-bp inserts were generated for the V4 region using the

barcoded primer set: 515FB: 5'-GTGYCAGCMGCCGCGGTAA-3' and 806RB: 5'-

GGACTACNVGGGTWTCTAAT-3' [61] [62] [63] [64] [65]. MiSeq Reagent Kit v2 (2 × 150

cycles, MS-102-2002) was used.

Sequence processing and analysis

Initial sequence processing and analyses were performed using QIIME 2 [66], any commands

prefixed by `q2-` are QIIME 2 plugins. After demultiplexing of the paired-end reads by `q2-`

`demux`, the imported sequence data was visually inspected via QIIME 2 View [67], to determine

the appropriate trimming and truncation parameters for generating Exact Sequence Variants

(ESVs) [68] via `q2-dada2` [69]. ESVs will be referred to as Operational Taxonomic Units

(OTUs). The forward reads were trimmed at 15 bp and truncated at 150 bp; reverse reads were

trimmed at 0 bp and truncated at 150 bp. The resulting OTUs were assigned taxonomy through

`q2-feature-classifier classify-sklearn`, by using a pre-trained classifier for the

amplicon region of interest [70]. This enables more robust taxonomic assignment of the OTUs

[71]. Taxonomy-based filtering was performed by using `q2-taxa filter-table` to remove

any OTUs that were classified as "Chloroplast", "Mitochondria", "Eukaryota", "Unclassified"

and those that did not have at least a Phylum-level classification. We then performed additional

quality filtering via `q2-quality-control`, and only retained OTUs that had at least a 90%

identity and 90% query alignment to the SILVA reference set [72]. Then `q2-alignment` was

used to generate a *de novo* alignment with MAFFT [73] which was subsequently masked by

setting `max-gap-frequency 1 min-conservation 0.4`. Finally, `q2-phylogeny`

was used to construct a midpoint-rooted phylogenetic tree using IQ-TREE [74] with automatic

model selection using ModelFinder [75]. Unless specified, subsequent analyses were performed

after removing OTUs with a very low frequency [76], of less than 0.0005% of the total data set

in this case.

Number of reads and OTUs before rarefying

Table 3 highlights the numbers of reads and OTUs among the DNA extraction protocols prior to

rarefying the data. The reads and OTUs assigned to gram-positive and gram-negative are also

shown. The number of "OTUs before rarefying" shown in Table 3 is distinguished from the

"Observed OTUs" after rarefying in Figure 3 for diversity analysis.

Microbiome analysis

To compare the taxonomic profiles among four types of DNA extraction methods (Figure 1 &

Table 1), the following analyses were performed; (I) bacterial microbiome composition, (II)

detection of common and unique taxa, (III) alpha and beta diversity analysis, and (IV)

identification of specific bacteria retained per DNA extraction method.

Microbiome composition

We generated the bar plot to exhibit bacterial microbiome composition per DNA extraction

method at the family (Figure 2A left) and genus (Figure 2A right) taxonomic level. After all

count data of taxonomy were converted to relative abundance, the top 10 abundant taxonomic

groups in each family and genus level were plotted in colored bar plot [77] [78] [79]. Variation

of microbiome composition per DNA extraction method or per individual was assessed by the

Adonis test (`q2-diversity adonis`)[80][81].

Differentially accessible microbiota by DNA extraction protocol

We set out to determine which microbial taxonomic groups were differentially accessible across

the sampling protocols by LEfSe analyses [31]. We further assessed the microbial taxa using

jvenn [82] at family and genus level. The Venn diagram was created after removing OTUs with a

frequency of less than 0.005% [76].

Alpha and beta diversity analyses with or without rarefying

Non-rarefying approaches to determine both alpha (within-sample) and beta (between-sample)

diversity was assessed by Species richness using `q2-breakaway` [36] and Aitchison distance

using `q2-deicode` [37]. These were compared with rarefied data in which we applied Faith's

Phylogenetic Diversity, Observed OTUs, Shannon's diversity index, Pielou's Evenness,

Unweighted UniFrac distance, Weighted UniFrac distance, Jaccard distance, and Bray-Curtis

distances via `q2-diversity` [66]. In order to retain data from at least 15 of the 20 patients

(*i.e.* 75%; four samples from each of the four DNA extraction methods), we set the sampling

depth to 51,197 reads per sample. Overall our subsequence analysis consisted of 3,071,820 reads

(27.6%, 3,071,820 / 11,149,582 reads). All diversity measurements in this study are listed in

Table 5.

Community type and HPV status

In addition to the analysis above, we tested whether the samples clustered by microbiome

composition were related to the patient's clinical and demographic characteristics such as,

cervical biopsy diagnosis, race, and HPV16 status. HPV16 status has been reported to be

associated with both racial differences as well as microbial community types [57] [83] [84] [85].

We employed the DMM [32] model to determine the number of community types for bacterial

cervical microbiome. Then, we clustered samples to the community type [9] [86]. Since vaginal

microbiota were reported to be clustered with different *Lactobacillus sp*. such as *L. crispatus, L.*

gasseri, L. iners, or *L. jensenii* [16] [87], we also collapsed the taxonomy to the species level and

performed a clustering analysis using "microbiome R package" [79]. We then determined which

bacterial taxa were differentially abundant among the patients with or without HPV16 via `q2-`

`aldex2` [88] and LEfSe [31].

General statistical analysis

All data are presented as means ± standard deviation (SD). Comparisons were conducted with

Fisher's exact test or Dunn's test with Benjamini-Hochberg-adjustment [89] or Wilcoxon test

with Benjamini-Hochberg-adjustment or pairwise PERMANOVA when appropriate. A p value <

0.05 or a q value < 0.05 was considered statistically significant.

Declarations

Ethics approval and consent to participate

This study was approved by the Institutional Review Board at the University of Arkansas for

Medical Sciences (IRB number 202790).

Consent for publication

Written informed consent for publication was obtained for all patients.

Availability of data and materials

MIMARKS compliant [90] DNA sequencing data are available via the Sequence Read Archive

(SRA) at the National Center for Biotechnology Information (NCBI), under the BioProject

Accession: PRJNA598197.

Competing interests

M.N. is one of the inventors named in the patents and patent applications for the HPV

therapeutic vaccine PepCan. The remaining authors declare no conflicts of interest.

Funding

This work was supported by the National Institutes of Health (R01CA143130, USA), Drs. Mae

and Anderson Nettleship Endowed Chair of Oncologic Pathology (31005156, USA), and the

Arkansas Biosciences Institute (the major component of the Tobacco Settlement Proceeds Act of

2000, G1-52249-01, USA).

Authors' contributions

Acknowledgements

We thank Togo Picture Gallery [91] for stock images shown in Figure 1.

Authors' information

Affiliations

Department of Pathology, University of Arkansas for Medical Sciences, Little Rock, AR, USA**:**

Takeo Shibata, Mayumi Nakagawa, & Hannah N. Coleman

Department of Obstetrics and Gynecology, Kanazawa Medical University, Uchinada, Ishikawa,

Japan: Takeo Shibata & Toshiyuki Sasagawa

Biosciences Division, Argonne National Laboratory, Lemont, IL, USA: Sarah M. Owens

Department of Obstetrics and Gynecology, University of Arkansas for Medical Sciences, Little

Rock, AR, USA: William W. Greenfield

Department of Biomedical Informatics, University of Arkansas for Medical Sciences, Little

Rock, AR, USA: Michael S. Robeson II

Corresponding author

Correspondence to Michael S. Robeson II

References

1.   Lee JE, Lee S, Lee H, Song YM, Lee K, Han MJ, et al. Association of the vaginal
microbiota with human papillomavirus infection in a Korean twin cohort. PLoS One.
2013;8(5):e63514; doi: 10.1371/journal.pone.0063514.
2.   Huang X, Li C, Li F, Zhao J, Wan X, Wang K. Cervicovaginal microbiota composition
correlates with the acquisition of high-risk human papillomavirus types. Int J Cancer.
2018;143(3):621-34; doi: 10.1002/ijc.31342.
3.   Zhou Y, Wang L, Pei F, Ji M, Zhang F, Sun Y, et al. Patients With LR-HPV Infection
Have a Distinct Vaginal Microbiota in Comparison With Healthy Controls. Front Cell
Infect Microbiol. 2019;9:294; doi: 10.3389/fcimb.2019.00294.
4.   Onywera H, Williamson AL, Mbulawa ZZA, Coetzee D, Meiring TL. The cervical
microbiota in reproductive-age South African women with and without human
papillomavirus infection. Papillomavirus Res. 2019;7:154-63; doi:
10.1016/j.pvr.2019.04.006.
5.   Brotman RM, Shardell MD, Gajer P, Tracy JK, Zenilman JM, Ravel J, et al. Interplay
between the temporal dynamics of the vaginal microbiota and human papillomavirus
detection. J Infect Dis. 2014;210(11):1723-33; doi: 10.1093/infdis/jiu330.
6.   Godoy-Vitorino F, Romaguera J, Zhao C, Vargas-Robles D, Ortiz-Morales G, Vazquez-
Sanchez F, et al. Cervicovaginal Fungi and Bacteria Associated With Cervical
Intraepithelial Neoplasia and High-Risk Human Papillomavirus Infections in a Hispanic
Population. Front Microbiol. 2018;9:2533; doi: 10.3389/fmicb.2018.02533.
7.   Łaniewski P, Barnes D, Goulder A, Cui H, Roe DJ, Chase DM, et al. Linking
cervicovaginal immune signatures, HPV and microbiota composition in cervical
carcinogenesis in non-Hispanic and Hispanic women. In: Sci Rep. 2018.
8.   Mitra A, MacIntyre DA, Lee YS, Smith A, Marchesi JR, Lehne B, et al. Cervical
intraepithelial neoplasia disease progression is associated with increased vaginal
microbiome diversity. Sci Rep. 2015;5:16865; doi: 10.1038/srep16865.
9.   Piyathilake CJ, Ollberding NJ, Kumar R, Macaluso M, Alvarez RD, Morrow CD.
Cervical Microbiota Associated with Higher Grade Cervical Intraepithelial Neoplasia in
Women Infected with High-Risk Human Papillomaviruses. Cancer Prev Res (Phila).
2016;9(5):357-66; doi: 10.1158/1940-6207.CAPR-15-0350.
10.   Oh HY, Kim BS, Seo SS, Kong JS, Lee JK, Park SY, et al. The association of uterine
cervical microbiota with an increased risk for cervical intraepithelial neoplasia in Korea.
Clin Microbiol Infect. 2015;21(7):674 e1-9; doi: 10.1016/j.cmi.2015.02.026.
11.   De Seta F, Campisciano G, Zanotta N, Ricci G, Comar M. The Vaginal Community State
Types Microbiome-Immune Network as Key Factor for Bacterial Vaginosis and Aerobic
Vaginitis. Front Microbiol. 2019;10:2451; doi: 10.3389/fmicb.2019.02451.
12.   Oliver A, LaMere B, Weihe C, Wandro S, Lindsay KL, Wadhwa PD, et al.
Cervicovaginal microbiome composition drives metabolic profiles in healthy pregnancy.
bioRxiv https://doi.org/10.1101/840520. 2019.
13.   Human Microbiome Project Consortium. Structure, function and diversity of the healthy
human microbiome. Nature. 2012;486(7402):207-14; doi: 10.1038/nature11234.
14.   Bik EM, Bird SW, Bustamante JP, Leon LE, Nieto PA, Addae K, et al. A novel
sequencing-based vaginal health assay combining self-sampling, HPV detection and

genotyping, STI detection, and vaginal microbiome analysis. PLoS One.
2019;14(5):e0215945; doi: 10.1371/journal.pone.0215945.

15.   Berman HL, McLaren MR, Callahan BJ. Understanding and Interpreting Community
Sequencing Measurements of the Vaginal Microbiome. BJOG. 2019; doi: 10.1111/1471-
0528.15978.

16.   Ravel J, Gajer P, Abdo Z, Schneider GM, Koenig SS, McCulle SL, et al. Vaginal
microbiome of reproductive-age women. Proc Natl Acad Sci U S A. 2011;108 Suppl
1:4680-7; doi: 10.1073/pnas.1002611107.

17.   Fettweis JM, Serrano MG, Brooks JP, Edwards DJ, Girerd PH, Parikh HI, et al. The
vaginal microbiome and preterm birth. Nat Med. 2019;25(6):1012-21; doi:
10.1038/s41591-019-0450-2.

18.   Tuominen H, Rautava S, Syrjanen S, Collado MC, Rautava J. HPV infection and
bacterial microbiota in the placenta, uterine cervix and oral mucosa. Sci Rep.
2018;8(1):9787; doi: 10.1038/s41598-018-27980-3.

19.   Mitra A, MacIntyre DA, Mahajan V, Lee YS, Smith A, Marchesi JR, et al. Comparison
of vaginal microbiota sampling techniques: cytobrush versus swab. Sci Rep.
2017;7(1):9802; doi: 10.1038/s41598-017-09844-4.

20.   Bentz JS. Liquid-based cytology for cervical cancer screening. Expert Rev Mol Diagn.
2005;5(6):857-71; doi: 10.1586/14737159.5.6.857.

21.   Gibb RK, Martens MG. The impact of liquid-based cytology in decreasing the incidence
of cervical cancer. Rev Obstet Gynecol. 2011;4(Suppl 1):S2-S11.

22.   Donders GG, Depuydt CE, Bogers JP, Vereecken AJ. Association of Trichomonas
vaginalis and cytological abnormalities of the cervix in low risk women. PLoS One.
2013;8(12):e86266; doi: 10.1371/journal.pone.0086266.

23.   Costea PI, Zeller G, Sunagawa S, Pelletier E, Alberti A, Levenez F, et al. Towards
standards for human fecal sample processing in metagenomic studies. Nat Biotechnol.
2017;35(11):1069-76; doi: 10.1038/nbt.3960.

24.   Stinson LF, Keelan JA, Payne MS. Comparison of Meconium DNA Extraction Methods
for Use in Microbiome Studies. Front Microbiol. 2018;9:270; doi:
10.3389/fmicb.2018.00270.

25.   Teng F, Darveekaran Nair SS, Zhu P, Li S, Huang S, Li X, et al. Impact of DNA
extraction method and targeted 16S-rRNA hypervariable region on oral microbiota
profiling. Sci Rep. 2018;8(1):16321; doi: 10.1038/s41598-018-34294-x.

26.   Balle C, Lennard K, Dabee S, Barnabas SL, Jaumdally SZ, Gasper MA, et al.
Endocervical and vaginal microbiota in South African adolescents with asymptomatic
Chlamydia trachomatis infection. Sci Rep. 2018;8(1):11109; doi: 10.1038/s41598-018-
29320-x.

27.   Klein C, Gonzalez D, Samwel K, Kahesa C, Mwaiselage J, Aluthge N, et al. Relationship
between the Cervical Microbiome, HIV Status, and Precancerous Lesions. MBio.
2019;10(1); doi: 10.1128/mBio.02785-18.

28.   Hayashi NR, Ishida T, Yokota A, Kodama T, Igarashi Y. Hydrogenophilus
thermoluteolus gen. nov., sp. nov., a thermophilic, facultatively chemolithoautotrophic,
hydrogen-oxidizing bacterium. Int J Syst Bacteriol. 1999;49 Pt 2:783-6; doi:
10.1099/00207713-49-2-783.

29.   Glassing A, Dowd SE, Galandiuk S, Davis B, Chiodini RJ. Inherent bacterial DNA
592   contamination of extraction and sequencing reagents may affect interpretation of
593   microbiota in low bacterial biomass samples. In: Gut Pathog. 2016.

594   30.   Birse KD, Romas LM, Guthrie BL, Nilsson P, Bosire R, Kiarie J, et al. Genital Injury
595   Signatures and Microbiome Alterations Associated With Depot Medroxyprogesterone
596   Acetate Usage and Intravaginal Drying Practices. J Infect Dis. 2017;215(4):590-8; doi:
597   10.1093/infdis/jiw590.

598   31.   Segata N, Izard J, Waldron L, Gevers D, Miropolsky L, Garrett WS, et al. Metagenomic
599   biomarker discovery and explanation. Genome Biol. 2011;12(6):R60; doi: 10.1186/gb-
600   2011-12-6-r60.

601   32.   Morgan M. DirichletMultinomial: Dirichlet-Multinomial Mixture Model Machine
602   Learning for Microbiome Data.
603   http://bioconductor.org/packages/release/bioc/html/DirichletMultinomial.html. Accessed
604   12 Mar 2020.

605   33.   Lennard K, Dabee S, Barnabas SL, Havyarimana E, Blakney A, Jaumdally SZ, et al.
606   Microbial Composition Predicts Genital Tract Inflammation and Persistent Bacterial
607   Vaginosis in South African Adolescent Females. Infect Immun. 2018;86(1); doi:
608   10.1128/IAI.00410-17.

609   34.   McMurdie PJ, Holmes S. Waste not, want not: why rarefying microbiome data is
610   inadmissible. PLoS Comput Biol. 2014;10(4):e1003531; doi:
611   10.1371/journal.pcbi.1003531.

612   35.   Weiss S, Xu ZZ, Peddada S, Amir A, Bittinger K, Gonzalez A, et al. Normalization and
613   microbial differential abundance strategies depend upon data characteristics.
614   Microbiome. 2017;5(1):27; doi: 10.1186/s40168-017-0237-y.

615   36.   Willis A, Bunge J. Estimating diversity via frequency ratios. Biometrics.
616   2015;71(4):1042-9; doi: 10.1111/biom.12332.

617   37.   Martino C, Morton JT, Marotz CA, Thompson LR, Tripathi A, Knight R, et al. A Novel
618   Sparse Compositional Technique Reveals Microbial Perturbations. mSystems. 2019;4(1);
619   doi: 10.1128/mSystems.00016-19.

620   38.   Yuan S, Cohen DB, Ravel J, Abdo Z, Forney LJ. Evaluation of methods for the
621   extraction and purification of DNA from the human microbiome. PLoS One.
622   2012;7(3):e33865; doi: 10.1371/journal.pone.0033865.

623   39.   Callahan BJ, Wong J, Heiner C, Oh S, Theriot CM, Gulati AS, et al. High-throughput
624   amplicon sequencing of the full-length 16S rRNA gene with single-nucleotide resolution.
625   Nucleic Acids Res. 2019;47(18):e103; doi: 10.1093/nar/gkz569.

626   40.   Calus ST, Ijaz UZ, Pinto AJ. NanoAmpli-Seq: a workflow for amplicon sequencing for
627   mixed microbial communities on the nanopore sequencing platform. Gigascience.
628   2018;7(12); doi: 10.1093/gigascience/giy140.

629   41.   Wongsurawat T, Nakagawa M, Atiq O, Coleman HN, Jenjaroenpun P, Allred JI, et al. An
630   assessment of Oxford Nanopore sequencing for human gut metagenome profiling: A pilot
631   study of head and neck cancer patients. J Microbiol Methods. 2019;166:105739; doi:
632   10.1016/j.mimet.2019.105739.

633   42.   Usyk M, Zolnik CP, Castle PE, Porras C, Herrero R, Gradissimo A, et al. Cervicovaginal
634   microbiome and natural history of HPV in a longitudinal study. PLoS Pathog.
635   2020;16(3):e1008376; doi: 10.1371/journal.ppat.1008376.

43. Audirac-Chalifour A, Torres-Poveda K, Bahena-Roman M, Tellez-Sosa J, Martinez-
Barnetche J, Cortina-Ceballos B, et al. Cervical Microbiome and Cytokine Profile at
Various Stages of Cervical Cancer: A Pilot Study. PLoS One. 2016;11(4):e0153274; doi:
10.1371/journal.pone.0153274.
44. Di Paola M, Sani C, Clemente AM, Iossa A, Perissi E, Castronovo G, et al.
Characterization of cervico-vaginal microbiota in women developing persistent high-risk
Human Papillomavirus infection. Sci Rep. 2017;7(1):10200; doi: 10.1038/s41598-017-
09842-6.
45. Ranjeva SL, Mihaljevic JR, Joseph MB, Giuliano AR, Dwyer G. Untangling the
dynamics of persistence and colonization in microbial communities. ISME J. 2019:1-13;
doi: 10.1038/s41396-019-0488-7.
46. Linder J, Zahniser D. ThinPrep Papanicolaou testing to reduce false-negative cervical
cytology. Arch Pathol Lab Med. 1998;122(2):139-44.
47. Ling Z, Liu X, Chen X, Zhu H, Nelson KE, Xia Y, et al. Diversity of cervicovaginal
microbiota associated with female lower genital tract infections. Microb Ecol.
2011;61(3):704-14; doi: 10.1007/s00248-011-9813-z.
48. Salter SJ, Cox MJ, Turek EM, Calus ST, Cookson WO, Moffatt MF, et al. Reagent and
laboratory contamination can critically impact sequence-based microbiome analyses.
BMC Biol. 2014;12(1):87; doi: 10.1186/s12915-014-0087-z.
49. Kim Y, Choi KR, Chae MJ, Shin BK, Kim HK, Kim A, et al. Stability of DNA, RNA,
cytomorphology, and immunoantigenicity in Residual ThinPrep Specimens. APMIS.
2013;121(11):1064-72; doi: 10.1111/apm.12082.
50. Castle PE, Solomon D, Hildesheim A, Herrero R, Concepcion Bratti M, Sherman ME, et
al. Stability of archived liquid-based cervical cytologic specimens. Cancer.
2003;99(2):89-96; doi: 10.1002/cncr.11058.
51. Rebolj M, Rask J, van Ballegooijen M, Kirschner B, Rozemeijer K, Bonde J, et al.
Cervical histology after routine ThinPrep or SurePath liquid-based cytology and
computer-assisted reading in Denmark. Br J Cancer. 2015;113(9):1259-74; doi:
10.1038/bjc.2015.339.
52. Naeem RC, Goldstein DY, Einstein MH, Ramos Rivera G, Schlesinger K, Khader SN, et
al. SurePath Specimens Versus ThinPrep Specimen Types on the COBAS 4800 Platform:
High-Risk HPV Status and Cytology Correlation in an Ethnically Diverse Bronx
Population. Lab Med. 2017;48(3):207-13; doi: 10.1093/labmed/lmx019.
53. Ritu W, Enqi W, Zheng S, Wang J, Ling Y, Wang Y. Evaluation of the Associations
Between Cervical Microbiota and HPV Infection, Clearance, and Persistence in
Cytologically Normal Women. Cancer Prev Res (Phila). 2019;12(1):43-56; doi:
10.1158/1940-6207.CAPR-18-0233.
54. Hosomi K, Ohno H, Murakami H, Natsume-Kitatani Y, Tanisawa K, Hirata S, et al.
Method for preparing DNA from feces in guanidine thiocyanate solution affects 16S
rRNA-based profiling of human microbiota diversity. Sci Rep. 2017;7(1):4339; doi:
10.1038/s41598-017-04511-0.
55. Human papillomavirus (HPV) and cervical cancer. https://www.who.int/news-room/fact-
sheets/detail/human-papillomavirus-(hpv)-and-cervical-cancer. Accessed 12 Mar 2020.
56. Sarangi AN, Goel A, Aggarwal R. Methods for Studying Gut Microbiota: A Primer for
Physicians. J Clin Exp Hepatol. 2019;9(1):62-73; doi: 10.1016/j.jceh.2018.04.016.

57.   Ravilla R, Coleman HN, Chow CE, Chan L, Fuhrman BJ, Greenfield WW, et al. Cervical
682   Microbiome and Response to a Human Papillomavirus Therapeutic Vaccine for Treating
683   High-Grade Cervical Squamous Intraepithelial Lesion. Integr Cancer Ther.
684   2019;18:1534735419893063; doi: 10.1177/1534735419893063.
685   58.   Roche Molecular Diagnostics. LINEAR ARRAY® HPV Genotyping.
686   https://diagnostics.roche.com/global/en/products/params/linear-array-hpv-
687   genotyping.html. Accessed 12 Mar 2020.
688   59.   Virtanen S, Kalliala I, Nieminen P, Salonen A. Comparative analysis of vaginal
689   microbiota sampling using 16S rRNA gene analysis. PLoS One. 2017;12(7):e0181477;
690   doi: 10.1371/journal.pone.0181477.
691   60.   Kim D, Hofstaedter CE, Zhao C, Mattei L, Tanes C, Clarke E, et al. Optimizing methods
692   and dodging pitfalls in microbiome research. Microbiome. 2017;5(1):52; doi:
693   10.1186/s40168-017-0267-5.
694   61.   Thompson LR, Sanders JG, McDonald D, Amir A, Ladau J, Locey KJ, et al. A
695   communal catalogue reveals Earth's multiscale microbial diversity. Nature.
696   2017;551(7681):457-63; doi: 10.1038/nature24621.
697   62.   Apprill A, McNally S, Parsons R, Weber L. Minor revision to V4 region SSU rRNA
698   806R gene primer greatly increases detection of SAR11 bacterioplankton. Aquat Microb
699   Ecol. 2015;75(2):129-37; doi: 10.3354/ame01753.
700   63.   Parada AE, Needham DM, Fuhrman JA. Every base matters: assessing small subunit
701   rRNA primers for marine microbiomes with mock communities, time series and global
702   field samples. Environ Microbiol. 2016;18(5):1403-14; doi: 10.1111/1462-2920.13023.
703   64.   Walters W, Hyde ER, Berg-Lyons D, Ackermann G, Humphrey G, Parada A, et al.
704   Improved Bacterial 16S rRNA Gene (V4 and V4-5) and Fungal Internal Transcribed
705   Spacer Marker Gene Primers for Microbial Community Surveys. mSystems. 2016;1(1);
706   doi: 10.1128/mSystems.00009-15.
707   65.   Earth Microbiome Project. 16S Illumina amplicon protocol.
708   http://www.earthmicrobiome.org/protocols-and-standards/16s/. Accessed 12 Mar 2020.
709   66.   Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, et al.
710   Reproducible, interactive, scalable and extensible microbiome data science using QIIME
711   2. Nat Biotechnol. 2019;37(8):852-7; doi: 10.1038/s41587-019-0209-9.
712   67.   QIIME 2 View. https://view.qiime2.org/. Accessed 12 Mar 2020.
713   68.   Callahan BJ, McMurdie PJ, Holmes SP. Exact sequence variants should replace
714   operational taxonomic units in marker-gene data analysis. ISME J. 2017;11(12):2639-43;
715   doi: 10.1038/ismej.2017.119.
716   69.   Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJ, Holmes SP. DADA2:
717   High-resolution sample inference from Illumina amplicon data. Nat Methods.
718   2016;13(7):581-3; doi: 10.1038/nmeth.3869.
719   70.   Bokulich NA, Kaehler BD, Rideout JR, Dillon M, Bolyen E, Knight R, et al. Optimizing
720   taxonomic classification of marker-gene amplicon sequences with QIIME 2's q2-feature-
721   classifier plugin. Microbiome. 2018;6(1):90; doi: 10.1186/s40168-018-0470-z.
722   71.   Werner JJ, Koren O, Hugenholtz P, DeSantis TZ, Walters WA, Caporaso JG, et al.
723   Impact of training sets on classification of high-throughput bacterial 16s rRNA gene
724   surveys. ISME J. 2012;6(1):94-103; doi: 10.1038/ismej.2011.82.

72. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, et al. The SILVA
ribosomal RNA gene database project: improved data processing and web-based tools.
Nucleic Acids Res. 2013;41(Database issue):D590-6; doi: 10.1093/nar/gks1219.
73. Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. Mol Biol Evol. 2013;30(4):772-80; doi:
10.1093/molbev/mst010.
74. Nguyen LT, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: a fast and effective
stochastic algorithm for estimating maximum-likelihood phylogenies. Mol Biol Evol.
2015;32(1):268-74; doi: 10.1093/molbev/msu300.
75. Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. ModelFinder:
fast model selection for accurate phylogenetic estimates. Nat Methods. 2017;14(6):587-9;
doi: 10.1038/nmeth.4285.
76. Bokulich NA, Subramanian S, Faith JJ, Gevers D, Gordon JI, Knight R, et al. Quality-
filtering vastly improves diversity estimates from Illumina amplicon sequencing. Nat
Methods. 2013;10(1):57-9; doi: 10.1038/nmeth.2276.
77. McMurdie PJ, Holmes S. phyloseq: an R package for reproducible interactive analysis
and graphics of microbiome census data. PLoS One. 2013;8(4):e61217; doi:
10.1371/journal.pone.0061217.
78. Bisanz JE. qiime2R: Importing QIIME2 artifacts and associated data into R sessions.
https://github.com/jbisanz/qiime2R. Accessed 12 Mar 2020.
79. Lahti L, Shetty S. microbiome R package. http://microbiome.github.io. Accessed 12 Mar
2020.
80. Anderson MJ. A new method for non-parametric multivariate analysis of variance.
Austral Ecol. 2001;26(1):32-46; doi: DOI 10.1111/j.1442-9993.2001.01070.pp.x.
81. Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, et al. vegan:
Community Ecology Package. R package version 2.5-3. https://CRAN.R-
project.org/package=vegan. Accessed 12 Mar 2020.
82. Bardou P, Mariette J, Escudie F, Djemiel C, Klopp C. jvenn: an interactive Venn diagram
viewer. BMC Bioinformatics. 2014;15(1):293; doi: 10.1186/1471-2105-15-293.
83. Gao W, Weng J, Gao Y, Chen X. Comparison of the vaginal microbiota diversity of
women with and without human papillomavirus infection: a cross-sectional study. BMC
Infect Dis. 2013;13(1):271; doi: 10.1186/1471-2334-13-271.
84. Montealegre JR, Peckham-Gregory EC, Marquez-Do D, Dillon L, Guillaud M, Adler-
Storthz K, et al. Racial/ethnic differences in HPV 16/18 genotypes and integration status
among women with a history of cytological abnormalities. Gynecol Oncol.
2018;148(2):357-62; doi: 10.1016/j.ygyno.2017.12.014.
85. Xi LF, Kiviat NB, Hildesheim A, Galloway DA, Wheeler CM, Ho J, et al. Human
papillomavirus type 16 and 18 variants: race-related distribution and persistence. J Natl
Cancer Inst. 2006;98(15):1045-52; doi: 10.1093/jnci/djj297.
86. Holmes I, Harris K, Quince C. Dirichlet multinomial mixtures: generative models for
microbial metagenomics. PLoS One. 2012;7(2):e30126; doi:
10.1371/journal.pone.0030126.
87. DiGiulio DB, Callahan BJ, McMurdie PJ, Costello EK, Lyell DJ, Robaczewska A, et al.
Temporal and spatial variation of the human microbiota during pregnancy. Proc Natl
Acad Sci U S A. 2015;112(35):11060-5; doi: 10.1073/pnas.1502875112.

88.   Fernandes AD, Macklaim JM, Linn TG, Reid G, Gloor GB. ANOVA-like differential
expression (ALDEx) analysis for mixed population RNA-Seq. PLoS One.
2013;8(7):e67019; doi: 10.1371/journal.pone.0067019.
89.   Dinno A. dunn.test: Dunn's Test of Multiple Comparisons Using Rank Sums.
https://CRAN.R-project.org/package=dunn.test. Accessed 12 Mar 2020.
90.   Yilmaz P, Kottmann R, Field D, Knight R, Cole JR, Amaral-Zettler L, et al. Minimum
information about a marker gene sequence (MIMARKS) and minimum information about
any (x) sequence (MIxS) specifications. Nat Biotechnol. 2011;29(5):415-20; doi:
10.1038/nbt.1823.
91.   Togo Picture Gallery. http://togotv.dbcls.jp/pics.html. Accessed 12 Mar 2020.
92.   Microbial Isolation | ZYMO RESEARCH.
https://www.zymoresearch.com/pages/microbial-isolation. Accessed 12 Mar 2020.
93.   PowerBead Tubes - QIAGEN Online Shop.
https://www.qiagen.com/us/products/discovery-and-translational-research/lab-
essentials/plastics/powerbead-tubes/#orderinginformation. Accessed 12 Mar 2020.
94.   QIAGEN. Pathogen Lysis Tubes - QIAGEN.
https://www.qiagen.com/dk/shop/pcr/pathogen-lysis-tubes/. Accessed 12 Mar 2020.
95.   Silhavy TJ, Kahne D, Walker S. The bacterial cell envelope. Cold Spring Harb Perspect
Biol. 2010;2(5):a000414; doi: 10.1101/cshperspect.a000414.
789
790

Table 1: Characteristics of four different DNA extraction protocols

Kit (Cat. No.)	Manufacturer	Sample volume	Enzyme	Beads	Beating	DNA carrier	Others
ZymoBIOMICS DNA Miniprep Kit (D4300)	Zymo Research	300 μL	No	Ceramic^a	2 min^b	No	^{c}
QIAamp PowerFecal Pro DNA Kit (51804)	Qiagen	300 μL	No	Ceramic^d	10 min^b	No	^{c}
QIAamp DNA Mini Kit (51304)	Qiagen	200 μL	Mutanolysin^e	No	No	No	$^{c, f, g}$
IndiSpin Pathogen Kit (SPS4104)	Indical Bioscience	300 μL	No	Ceramic^h	10 min^b	Yes	$^{c, i}$

a: [92]. b: Disruptor Genie (USA Scientific, Inc.) was used under the maximum speed. c: Nuclease free water (85 μL) as DNA elution buffer was used. d: PowerBead Pro Tubes [93]. e: Instead of lysozyme or lysostaphin, mutanolysin was used as per Yuan *et al*, 2012 [38]. f: DNA Purification from Blood or Body Fluids; Protocols for Bacteria; Isolation of genomic DNA from gram-positive bacteria in QIAamp DNA Mini and Blood Mini Handbook fifth edition was referenced. g: Heating at 56°C for 30 min and 95°C for 15 min was performed. h: Pathogen Lysis Tubes S [94]. i: Pretreatment B2 as per QIAamp cador Pathogen Mini Handbook.

Table 2. Patient characteristics

Characteristics		Values
Number of patients, n		20
Total number of DNA extracts, n		80
Age, mean (SD)		31.4 (5.0)
Race	African American, n (%)	3 (15)
	Caucasian, n (%)	10 (50)
	Hispanic, n (%)	7 (35)
Cervical biopsy	CIN2, n (%)	8 (40)
	CIN3, n (%)	10 (50)
	Benign, n (%)	2 (10)
HPV typing	HPV positive, n (%)	19 (95)
	HPV16 positive, n (%)	10 (50)
	HPV18 positive, n (%)	2 (10)
	HPV16 or 18 positives, n (%)	10 (50)
	HR-HPV positives, n (%)	18 (90)

SD: standard deviation. CIN: cervical intraepithelial neoplasia. HR-HPV: high-risk HPV (HPV16 18, 31, 33, 35, 39, 45, 51, 52, 56, 58, 59, and 68)

Table 3. Reads and OTUs before rarefying assigned to all, gram-, and gram-negative bacteria per DNA extraction protocols

Parameters	Community	Methods	Values	Ratio of GP or GN	p value
Number of reads (mean ± SD)	All	Zy	2,705,044 (135,252 ± 66,011)		[a]
		Pro	2,312,207 (115,610 ± 68,201)		
		QIA	2,765,343 (138,267 ± 49,781)		
		IN	3,366,988 (168,349 ± 57,451)		
	GP	Zy	2,430,380 (121,519 ± 56,209)	89.8%	NS
		Pro	2,116,458 (105,823 ± 57,590)	91.5%	
		QIA	2,503,578 (125,179 ± 46,073)	90.5%	
		IN	2,985,941 (149,297 ± 46,936)	88.7%	
	GN	Zy	274,664 (13,733 ± 29,162)	10.2%	NS
		Pro	195,749 (9,788 ± 23,070)	8.5%	
		QIA	261,765 (13,088 ± 22,638)	9.5%	
		IN	381,047 (19,052 ± 33,038)	11.3%	
Number of OTUs (mean ± SD)	All	Zy	825 (41.3 ± 16.8)		NS
		Pro	621 (31.1 ± 19.4)		
		QIA	778 (38.9 ± 22.4)		
		IN	792 (39.6 ± 22.7)		
	GP	Zy	479 (24.0 ± 9.2)	58.1%	NS
		Pro	412 (20.6 ± 12.7)	66.3%	
		QIA	513 (25.7 ± 13.7)	65.9%	
		IN	531 (26.6 ± 14.9)	67.0%	
	GN	Zy	346 (17.3 ± 9.8)	41.9%	[b]
		Pro	209 (10.5 ± 10.3)	33.7%	
		QIA	265 (13.3 ± 9.2)	34.1%	
		IN	261 (13.1 ± 8.3)	33.0%	

Community of gram-positive bacteria was defined as phylum *Actinobacteria* and *Firmicutes*, which are composed of thick peptidoglycan layers without outer membrane [95]. Community of gram-negative bacteria was defined as a community of bacteria other than phylum *Actinobacteria* and *Firmicutes* in this study. a: I - P: 0.0199; I - Q: 0.1590; P - Q: 0.1436; I - Z: 0.1495; P - Z: 0.1712; and Q - Z: 0.4059. b: I - P: 0.2116; I - Q: 0.4837; P - Q: 0.1143; I - Z: 0.0938; P - Z: 0.0116; Q - Z: 0.1448. Dunn's test with Benjamini-Hochberg-adjustment were performed for comparison of the number of reads and OTUs by DNA extraction method. Zy: ZymoBIOMICS DNA Miniprep Kit, Pro: QIAamp PowerFecal Pro DNA Kit, QIA: QIAamp DNA Mini Kit, IN: IndiSpin Pathogen Kit. SD: standard deviation. All: all bacteria, GP: gram-positive bacteria, GN: gram-negative bacteria. NS: not significant.

Table 4. Beta diversity among DNA extraction methods

Index	Protocol	Protocols compared	p values	q values
Aitchison distance	ZymoBIOMICS	PowerFecalPro	NS	NS
		QIAampMini	NS	NS
		IndiSpin	NS	NS
	PowerFecalPro	QIAampMini	NS	NS
		IndiSpin	NS	NS
	QIAampMini	IndiSpin	NS	NS
Unweighted UniFrac distance	ZymoBIOMICS	PowerFecalPro	0.001	0.002
		QIAampMini	0.001	0.002
		IndiSpin	0.001	0.002
	PowerFecalPro	QIAampMini	NS	NS
		IndiSpin	0.015	0.023
	QIAampMini	IndiSpin	NS	NS
Weighted UniFrac distance	ZymoBIOMICS	PowerFecalPro	NS	NS
		QIAampMini	NS	NS
		IndiSpin	NS	NS
	PowerFecalPro	QIAampMini	NS	NS
		IndiSpin	NS	NS
	QIAampMini	IndiSpin	NS	NS
Jaccard distance	ZymoBIOMICS	PowerFecalPro	0.037	NS
		QIAampMini	0.003	0.018
		IndiSpin	0.011	0.033
	PowerFecalPro	QIAampMini	NS	NS
		IndiSpin	NS	NS
	QIAampMini	IndiSpin	NS	NS
Bray-Curtis distance	ZymoBIOMICS	PowerFecalPro	NS	NS

	QIAampMini	NS	NS
	IndiSpin	NS	NS
PowerFecalPro	QIAampMini	NS	NS
	IndiSpin	NS	NS
QIAampMini	IndiSpin	NS	NS

Pairwise PERMANOVA was tested for comparing beta diversity of DNA extraction method. NS: not significant.

Table 5. Diversity analysis in this study

No.	Parameter	Alpha or Beta diversity	Used data with/without rarefying	Input data with/without phylogenetic information	Plugin of QIIME 2
1	Species richness	Alpha	Not rarefied	Non-phylogenetic	`q2-breakaway` [36]
2	Faith's Phylogenetic Diversity	Alpha	Rarefied	Phylogenetic	`q2-diversity`
3	Observed OTUs	Alpha	Rarefied	Non-phylogenetic	`q2-diversity`
4	Shannon's diversity index	Alpha	Rarefied	Non-phylogenetic	`q2-diversity`
5	Pielou's Evenness	Alpha	Rarefied	Non-phylogenetic	`q2-diversity`
6	Aitchison distance	Beta	Not rarefied	Non-phylogenetic	`q2-deicode` [37]
7	Unweighted UniFrac distance	Beta	Rarefied	Phylogenetic	`q2-diversity`
8	Weighted UniFrac distance	Beta	Rarefied	Phylogenetic	`q2-diversity`
9	Jaccard distance	Beta	Rarefied	Non-phylogenetic	`q2-diversity`
10	Bray-Curtis distances	Beta	Rarefied	Non-phylogenetic	`q2-diversity`
11	Adonis	Beta	Rarefied	Non-phylogenetic	`q2-diversity adonis` [80] [81]

Figure legends

Figure 1. Overview of the study design using 16S rRNA gene to compare the DNA

extraction protocol. (A) Liquid-based cytology (LBC) specimens from 20 patients with CIN2/3

or suspected CIN2/3. (B) A total of 80 DNA extractions were performed. (C) The four DNA

extraction methods. (D) DNA of mock vaginal community as a positive control and preservation

solution as a negative control. (E) Sequencing using Illumina MiSeq. (F) Analysis of the

taxonomic profiles among the DNA extraction protocols. Images form Togo Picture Gallery [91]

were used to create this figure.

Figure 2. Taxonomic resolution among DNA extraction protocols. (A) Relative abundance of

microbe at family level (left) and genus level (right) per DNA extraction method showed the

pattern that variance of microbe composition per patient was higher than that per DNA extraction

protocol. These patterns were confirmed by values of Adonis test (`q2-diversity adonis`);

F.Model: 199.4, R2: 0.982, and p value: 0.001 for patients and F.Model: 2.9, R2: 0.003, and p

value: 0.002 for DNA extraction [80] [81]. After all count data of taxonomy were converted to

relative abundance as shown in the y-axis, the top ten taxonomy at each family and genus level

were plotted in colored bar plot and other relatively few taxonomies were not plotted. The 20

patients ID were described in the x-axis. (B) Venn diagrams, considering only those OTUs with

a frequency greater than 0.005% shown, revealed that ZymoBIOMICS had four unique taxa at

family (left) and genus (right) taxonomic level. Thirty-one of 41 families and 45 of 57 genera

were detected with all DNA extraction protocols.

Figure 3. Comparisons of alpha diversity between different DNA extraction protocols. The

alpha diversity indices determined by Species richness and Phylogenetic diversity are

significantly higher with ZymoBIOMICS in comparison with PowerFecalPro (p = 0.025 and

0.012, respectively, Dunn's test with Benjamini-Hochberg-adjustment). IndiSpin also showed

significantly higher diversity than that of PowerFecalPro using analysis of Species richness (p =

0.042, Dunn's test with Benjamini-Hochberg-adjustment). No significant differences were

observed in other alpha diversity indexes such as observed OTUs, Shannon's diversity index,

and Pielou's Evenness. Zy: ZymoBIOMICS DNA Miniprep Kit, Pro: QIAamp PowerFecal Pro

DNA Kit, QIA: QIAamp DNA Mini Kit, IN: IndiSpin Pathogen Kit.

Figure 4. Distinct detections of microbe among the DNA extraction protocols. (A) A bar

graph showing 23 significantly enriched taxa with ZymoBIOMICS, 3 with QIAamp DNA Mini

Kit, and 3 with IndiSpin Pathogen Kit determined by the linear discriminant analysis (LDA)

effect size (LEfSe) analyses [31]. (B) A taxonomic cladogram from the same LEfSe analyses

showing that the significantly enriched microbiota in ZymoBIOMICS were composed of phylum

Proteobacteria. Also note that *Meiothermus* (a member of the phylum *Deinococcus-Thermus*)

Hydrogenophilaceae (a member of the phylum *Proteobacteria*), and *Hydrogenophilus* (a

member of the phylum *Proteobacteria*) are likely an extraction kit contaminant. Zy:

ZymoBIOMICS DNA Miniprep Kit, Pro: QIAamp PowerFecal Pro DNA Kit, QIA: QIAamp

DNA Mini Kit, IN: IndiSpin Pathogen Kit. g_: genus, f_: family, o_: order, c_: class, p_:

phylum.

Figure S1. Comparison of DNA yields by DNA extraction protocols. DNA yield of

QIAampMini was significantly higher than that of PowerFecalPro ($p < 0.001$, Dunn's test with

Benjamini-Hochberg-adjustment). Also, the DNA yield of ZymoBIOMICS was significantly

higher than that of PowerFecalPro ($p < 0.001$, Dunn's test with Benjamini-Hochberg-

adjustment). The amount of DNA was calculated based on the absorbance of nucleic acids

measured by Nanodrop One. By the protocol recommended by the manufacturer, nucleic acid

(Poly-A carrier) was used in IndiSpin. Therefore, IndiSpin was excluded from the analysis of

DNA yield. The amount of DNA yield per 100 μL ThinPrep sample volume were compared. The

bar graph shows the mean and standard deviation. Zy: ZymoBIOMICS DNA Miniprep Kit, Pro:

QIAamp PowerFecal Pro DNA Kit, QIA: QIAamp DNA Mini Kit.

Figure S2. Community type and HPV 16 assessed by using 4 kits (A) Community types were

classified into two types in all DNA extraction kits, mainly based on the percentage of

Lactobacillus iners. HPV16 infection was negatively associated with the dominance of *L. iners*

(community type I; $p = 0.001$, Fisher's exact test) regardless of DNA extraction method.

Although, we observed slight variation in the abundance of microbiota across the extraction kits

(even within the same individual patient), the ability to detect two community types was

identical across all DNA extraction kits. No significant differences were observed in the

relationship of other phenotypes of patients (HPV18, HR-HPV, Biopsy, and Race). The top 15

bacteria detected for each DNA extraction kit are shown. Samples were clustered by the

Dirichlet component. Narrow columns show each sample and a broader column shows averages

of samples. Rows show taxa at the species level. Dark or thin colors correspond to larger or

smaller counts of OTUs, respectively. CT: community type. (B) LEfSe analysis using combined

data from all four kits detected a significant enrichment of 66 taxa in the cervical environment

with HPV16 infection and 17 taxa without HPV16 infection. Genus Lactobacillus were enriched

in the HPV16 negative patients (p < 0.001, LDA score: 5.38).

(A) Liquid-based cytology on cervix

Cervical microbiome

LBC

(B) Dispensing of each sample to 4 aliquot

(C) DNA extraction using four protocols

ZymoBIOMICS

PowerFecalPro

QIAampMini

IndiSpin

(D)

Positive control (vaginal mock)

Negative control (preservation solution without samples)

(E) 16S rRNA Illumina amplicon sequencing

(F) Evaluation of taxonomic profiles

ZymoBIOMICS   PowerFecalPro

QIAampMini   IndiSpin