Evaluation of DNA extraction protocols from liquid-based cytology specimens

for studying cervical microbiota

Takeo Shibata,[a,b] Mayumi Nakagawa,[a] Hannah N. Coleman,[a] Sarah M. Owens,[c] William W.

Greenfield,[d] Toshiyuki Sasagawa,[b] Michael S. Robeson II[e]

[a]Department of Pathology, University of Arkansas for Medical Sciences, Little Rock, AR, USA

[b]Department of Obstetrics and Gynecology, Kanazawa Medical University, Uchinada, Ishikawa,

Japan

[c]Biosciences Division, Argonne National Laboratory, Lemont, IL, USA

[d]Department of Obstetrics and Gynecology, University of Arkansas for Medical Sciences, Little

Rock, AR, USA

[e]Department of Biomedical Informatics, University of Arkansas for Medical Sciences, Little

Rock, AR, USA

Corresponding author: Michael S. Robeson II

Tel: 501-526-4242, Fax: 501-526-5964

Email: MRobeson@uams.edu

Abstract

Cervical microbiota (CM) are considered an important factor affecting the progression of

cervical intraepithelial neoplasia (CIN) and are implicated in the persistence of human

papillomavirus (HPV). Collection of liquid-based cytology (LBC) samples is routine for cervical

cancer screening and HPV genotyping, and can be used for long-term cytological biobanking.

Herein, we investigate the feasibility of leveraging LBC specimens for use in CM surveys by

amplicon sequencing. As methodological differences in DNA extraction protocols can

potentially bias the composition of microbiota, we set out to determine the performance of four

commonly used DNA extraction kits (ZymoBIOMICS DNA Miniprep Kit; QIAamp PowerFecal

Pro DNA Kit; QIAamp DNA Mini Kit; and IndiSpin Pathogen Kit) and their ability to capture

the diversity of CM from LBC specimens. LBC specimens from 20 patients (stored for $716 \pm$

105 days) with cervical intraepithelial neoplasia (CIN) 2/3 or suspected CIN2/3 were each

aliquoted for extraction by each of the four kits. We observed that, regardless of the extraction

protocol used, all kits provided equivalent accessibility to the cervical microbiome, with some

minor differences. For example, the ZymoBIOMICS kit appeared to differentially increase

access of several more microbiota compared to the other kits. Potential kit contaminants were

observed as well. Approximately 80% microbial genera were shared among all DNA extraction

protocols. The variance of microbial composition per individual was larger than that of the DNA

extraction protocol used. We also observed that HPV16 infection was significantly associated

with community types that were not dominated by *Lactobacillus iners*.

Importance

Collection of LBC specimens is routine for cervical cancer screening and HPV genotyping, and

can be used for long-term cytological biobanking. We demonstrated that LBC samples, which

had been under prolonged storage prior to DNA extraction, were able to provide a robust

assessment of the CM and its relationship to HPV status, regardless of the extraction kit used.

Being able to retroactively access the CM from biobanked LBC samples, will allow researchers

to better interrogate historical interactions between the CM and its relationship to CIN and HPV.

This alone has the potential to bring CM research one-step closer to the clinical practice.

Keywords; cervical microbiota, DNA extraction, HPV, CIN, liquid-based cytology

Introduction

High-throughput sequencing (HTS) technology of 16S rRNA gene amplicon sequences has made it possible to better understand the relationships between cervicovaginal microbiota and human papillomavirus (HPV) infection (1) (2) (3) (4) (5) and HPV-related diseases (6) (7) (8) (9) (10). Cervicovaginal microbiota are considered to be an important factor affecting the progress of cervical intraepithelial neoplasia (CIN) (6) (7) (8) (9) and are implicated in the persistence of high-risk HPV (HR-HPV) (1) (2) and low-risk HPV (LR-HPV) (3). For example, the phyla *Actinobacteria* and *Fusobacteria* were enriched in HR-HPV positive environment (4) while the phyla *Actinobacteria*, *Proteobacteria*, and *Fusobacteria* in low-risk HPV (LR-HPV) (3). Additionally, *Lactobacillus iners*-dominant samples are associated with both HR-HPV and LR-HPV (5). Moreover, it has been shown that CIN risk was increased when the cervical microbes *Atopobium vaginae*, *Gardnerella vaginalis*, and *Lactobacillus iners* were present with HR-HPV (10). The cervicovaginal microbiome specified by *Lactobacillus*-dominant type or non-*Lactobacillus*-dominant type has been shown to interact with the immune system (7) (11). Inflammatory cytokines, such as Interleukin (IL)-1α and IL-18, were increased in non-*Lactobacillus*-dominant community types of reproductive-aged healthy women (11). In the analysis of patients with cervical cancer, non-*Lactobacillus*-dominant community types were positively associated with chemokines such as interferon gamma-induced protein 10 (IP-10) and soluble CD40-ligand activating dendric cells (DCs) (7). The metabolism of the cervicovaginal microbiome may be a substantial contributing factor to maternal health during pregnancy, although the mechanism is still unclear (12).

Little has been reported on the utility of liquid-based cytology (LBC) samples for use in cervical microbiome studies. Conventionally, microbiome sample collection methods entail the

use of swabs (13) or self-collection of vaginal discharge (14). To obtain a non-biased and broad

range of cervical microbiota, DNA extraction should be optimized for a range of difficult-to-

lyse-bacteria, *e.g. Firmicutes*, *Actinobacteria*, and *Lactobacillus* (13) (15) (16) (17) (18).

LBC samples are promising for cervicovaginal microbiome surveys, as they are an

already established method of long-term cytological biobanking (19). In clinical practice,

cervical cytology for cervical cancer screening or HPV genotyping is widely performed using a

combination of cervical cytobrushes and LBC samples such as ThinPrep (HOLOGIC) or

SurePath (BD). An LBC specimen can be used for not only cytological diagnosis but also

additional diagnostic tests such as HPV, *Chlamydia*, *Neisseria gonorrhoeae*, and *Trichomonas*

infection (20) (21) (22).

The ability to characterize microbial communities, as commonly assessed by 16S rRNA

gene sequencing, can be biased as a result of methodological differences of cell lysis and DNA

extraction protocols (23) (24) (25). Herein, we compare four different commercially available

DNA extraction kits in an effort to assess their ability to characterize the cervical microbiota of

LBC samples. Additionally, we examine the relationship between HPV infection and the

composition of cervical microbiota.

Materials and Methods

Ethics

This study was approved by the Institutional Review Board at the University of Arkansas for

Medical Sciences (IRB number 202790).

Sampling of cervical microbiome

LBC specimens were obtained from 20 patients enrolled in a Phase II clinical trial of an HPV

therapeutic vaccine (NCT02481414). In order to be eligible, participants had to have high grade

squamous intraepithelial lesions (HSILs) or cannot rule out HSILs in cervical cytology or

CIN2/3 in cervical biopsy. Those who qualified for the study based on their cervical cytology

underwent cervical biopsy, and they qualified for vaccination if the results were CIN2/3. The

cervical cytology specimens in this current study were collected before the vaccination and

reserved in the vial of the ThinPrep Pap Test (HOLOGIC) as described in Ravilla *et al*. 2019

(26). The storage period from sample collection to DNA extraction was 716 ± 105 days in this

study.

HPV genotyping

HPV-DNA was detected by Linear Array HPV Genotyping Test (Roche Diagnostics) which can

detect up to 37 HPV genotypes including 13 HR-HPV genotypes (16, 18, 31, 33, 35, 39, 45, 51,

52, 56, 58, 59, and 68) and 24 LR-HPV genotypes (6, 11, 26, 40, 42, 53, 54, 55, 61, 62, 64, 66,

67, 69, 70, 71, 72, 73, 81, 82, 83, 84, IS39, and CP6108) using ThinPrep solution

(https://diagnostics.roche.com/global/en/products/params/linear-array-hpv-genotyping.html).

DNA extraction protocols

We selected four commercially available DNA extraction kits as the candidates for comparison:

ZymoBIOMICS DNA Miniprep Kit (Zymo Research, D4300), QIAamp PowerFecal Pro DNA

Kit (QIAGEN, 51804), QIAamp DNA Mini Kit (QIAGEN, 51304), and IndiSpin Pathogen Kit

(Indical Bioscience, SPS4104). These kits have been successfully used in a variety of human

cervical, vaginal, and gut microbiome surveys (10) (19) (27). We'll subsequently refer to each of

these kits in abbreviated form as follows: ZymoBIOMICS, PowerFecalPro, QIAampMini, and

IndiSpin. The protocols and any modifications are outlined in Table 1.

Each LBC sample was dispensed into four separate 2 mL sterile collection tubes

(dispensed sample volume = 500 μL) to create four cohorts of 20 DNA extractions (Fig. 1). Each

extraction cohort was processed through one of the four kits above. A total of 80 extractions (4

kits × 20 patients) were prepared for subsequent analyses. Applied sample volume of ThinPrep

solution was 300 μL for ZymoBIOMICS, 300 μL for PowerFecalPro, and 200 μL for

QIAampMini, and 300 μL for IndiSpin. The sample volume was standardized to 300 μL as long

as the manufacturer's instructions allowed to do so. DNA extraction for all samples was

performed by the same individual who practiced by performing multiple extractions for each kit

before performing the actual DNA extraction on the samples analyzed in this study. Positive

control was mock vaginal microbial communities composed of a mixture of genomic DNA from

the American Type Culture Collection (ATCC MSA1007). Negative control was the ThinPrep

preservation solution without the sample as blank extraction (28).

Measurement of DNA yield

DNA yield for each method was evaluated by spectrophotometer (Nanodrop One, Thermo

Scientific). Analysis of the DNA yield from IndiSpin was omitted as nucleic acid is used as a

carrier for this kit. The mean DNA yields per 100 μL ThinPrep sample volume were compared.

16S rRNA marker gene sequencing

Controls and the extracted DNA were sent to Argonne National Laboratory (IL, USA) for

amplification and sequencing of the 16S rRNA gene on an Illumina MiSeq sequencing platform.

Paired-end reads from libraries with ~250-bp inserts were generated for the V4 region using the

barcoded primer set: 515FB: 5'-GTGYCAGCMGCCGCGGTAA-3' and 806RB: 5'-

GGACTACNVGGGTWTCTAAT-3' (29) (30) (31) (32)

(http://www.earthmicrobiome.org/protocols-and-standards/16s/). MiSeq Reagent Kit v2 (2 × 150

cycles, MS-102-2002) was used.

Sequence processing and analysis

Initial sequence processing and analyses were performed using QIIME 2 (33), any commands

prefixed by `q2-` are QIIME 2 plugins. After demultiplexing of the paired-end reads by `q2-`

`demux`, the imported sequence data was visually inspected via QIIME 2 View

(https://view.qiime2.org), to determine the appropriate trimming and truncation parameters for

generating Exact Sequence Variants (ESVs) (34) via `q2-dada2` (35). Hereafter, ESVs will be

referred to as OTUs (Operational Taxonomic Units). The forward reads were trimmed at 15 bp

and truncated at 150 bp; reverse reads were trimmed at 0 bp and truncated at 150 bp. The

resulting OTUs were assigned taxonomy through `q2-feature-classifier classify-`

`sklearn`, by using a pre-trained classifier for the amplicon region of interest (36). This enables

more robust taxonomic assignment of the OTUs (37). Taxonomy-based filtering was performed

by using `q2-taxa filter-table` to remove any OTUs that were classified as

"Chloroplast", "Mitochondria", "Eukaryota", "Unclassified" and those that did not have at least

a Phylum-level classification. We then performed additional quality filtering via `q2-quality-`

`control`, and only retained OTUs that had at least a 90% identity and 90% query alignment to

the SILVA reference set (38). Then `q2-alignment` was used to generate a *de novo* alignment

with MAFFT (39) which was subsequently masked by setting `max-gap-frequency 1`

`min-conservation 0.4`. Finally, q2-phylogeny was used to construct a midpoint-rooted

phylogenetic tree using IQ-TREE (40) with automatic model selection using ModelFinder (41).

Unless specified, subsequent analyses were performed after removing OTUs with a frequency of

less than 0.0005% of the total data set (42).

Number of reads and OTUs before rarefying

Table 3 highlights the numbers of reads and OTUs among the DNA extraction protocols prior to

rarefying the data. The reads and OTUs assigned to gram-positive and gram-negative was also

shown. The number of "OTUs before rarefying" shown in Table 3 is distinguished from the

"Observed OTUs" after rarefying in Fig. 3 for diversity analysis.

Microbiome analysis

To compare the taxonomic profiles among four types of DNA extraction methods (Fig. 1 &

Table 1), the following analyses were performed; (I) bacterial microbiome composition, (II)

detection of common and unique taxa, (III) alpha and beta diversity analysis, and (IV)

identification of specific bacteria retained per DNA extraction method.

Microbiome composition

We generated the bar plot to exhibit bacterial microbiome composition per DNA extraction

method at the family (Fig. 2A left) and genus (Fig. 2A right) taxonomic level. After all count

data of taxonomy were converted to relative abundance, the top 10 abundant taxonomic groups

in each family and genus level were plotted in colored bar plot (43) (44) (45). Variation of

microbiome composition per DNA extraction method or per individual was assessed by the

Adonis test (q2-diversity adonis) (46) (47).

Differentially accessible microbiota by DNA extraction protocol

195 We set out to determine which microbial taxonomic groups were differentially accessible across

196 the sampling protocols by linear discriminant analysis (LDA) effect size (LEfSe) analyses (48).

197 We further assessed the microbial taxa using jvenn (49) at family and genus level. The Venn

198 diagram was created after removing OTUs with a frequency of less than 0.005% (42).

Alpha and beta diversity analyses with or without rarefying

Non-rarefying approaches to determine both alpha (within-sample) and beta (between-sample)

diversity was assessed by Species richness using q2-breakaway (50) and Aitchison distance

using q2-deicode (51). These were compared with rarefied data in which we applied Faith's

Phylogenetic Diversity, Observed OTUs, Shannon's diversity index, Pielou's Evenness,

Unweighted UniFrac distance, Weighted UniFrac distance, Jaccard distance, and Bray-Curtis

distances via q2-diversity (33). In order to retain data from at least 15 of the 20 patients (*i.e.*

75%; four samples from each of the four DNA extraction methods), we set the sampling depth to

51,197 reads per sample. Overall our subsequence analysis consisted of 3,071,820 reads (27.6%,

3,071,820 / 11,149,582 reads). All diversity measurements in this study are listed in Table S1.

Community type and HPV status

In addition to the analysis above, we tested whether the samples clustered by microbiome

composition were related to the patient's clinical and demographic characteristics such as,

cervical biopsy diagnosis, race, and HPV16 status. HPV16 status has been reported to be

associated with both racial differences as well as microbial community types (26) (52) (53) (54).

We employed the Dirichlet Multinomial Mixtures (DMM) (55) model to determine the number

of community types for bacterial cervical microbiome. Then, we clustered samples to the

community type (9) (56). Since vaginal microbiota were reported to be clustered with different

Lactobacillus sp. such as *L. crispatus*, *L. gasseri*, *L. iners*, or *L. jensenii* (16) (57), we also

collapsed the taxonomy to the species level and performed a clustering analysis using

"microbiome R package" (45). We then determined which bacterial taxa were differentially

abundant among the patients with or without HPV16 via q2-aldex2 (58) and LEfSe (48).

General statistical analysis

All data are presented as means ± standard deviation (SD). Comparisons were conducted with

Fisher's exact test or Dunn's test with Benjamini-Hochberg-adjustment (59) or Wilcoxon test

with Benjamini-Hochberg-adjustment or pairwise PERMANOVA when appropriate. A p value <

0.05 or a q value < 0.05 was considered statistically significant.

Data availability

MIMARKS compliant (60) DNA sequencing data are available via the Sequence Read Archive

(SRA) at the National Center for Biotechnology Information (NCBI), under the BioProject

Accession: PRJNA598197.

Results

Patients characteristics

The age of the patients was 31.4 ± 5.0 years. The distribution of race was 15% African American

(n = 3) and 50% Caucasian (n = 10), and 35% Hispanic (n = 7). Cervical histology was 40%

CIN2 (n = 8), 50% CIN3 (n = 10), and 10% benign (n = 2). HPV genotype were 50% HPV16

positive (n = 10), 10% HPV18 positive (n = 2), 90% HR-HPV positives (n = 18). Patient

characteristics were summarized in Table 2.

DNA yield

DNA yield per 100 μL ThinPrep solution were 0.09 ± 0.06 μg in ZymoBIOMICS, 0.04 ± 0.01 μg

in PowerFecalPro, and 0.21 ± 0.23 μg in QIAampMini. DNA yield was not calculated for

IndiSpin, as Poly-A Carrier DNA was used. The DNA yield of PowerFecalPro was significantly

lower than that of ZymoBIOMICS (adjusted p value < 0.001) and QIAampMini (adjusted p

value < 0.001) based on Dunn's test with Benjamini-Hochberg-adjustment (Fig. S1).

Number of reads and OTUs before rarefying

We obtained a total of 11,149,582 reads for 80 DNA extractions. The 127,142 reads were

produced from a positive control of mock sample and 1,773 reads from ThinPrep solution as the

negative control. IndiSpin (168,349 ± 57,451 reads) produced a significantly higher number of

reads compared to PowerFecalPro (115,610 ± 68,201 reads, p value = 0.020, Dunn's test with

Benjamini-Hochberg-adjustment) as shown in Table 3. Approximately 90% of reads were

assigned to gram-positive bacteria and about 10% of reads were assigned to gram-negative

bacteria across all kits.

Prior to rarefying, the ZymoBIOMICS kit captured a greater representation of gram-

negative bacterial OTUs (total 346, 17.3 ± 9.8) compared to PowerFecalPro (total 209, 10.5 ±

10.3, p value = 0.012, Dunn's test with Benjamini-Hochberg-adjustment, ratio of gram-negative

bacteria: 41.9% vs 33.7%) as shown in Table 3. No significant differences in the number of

OTUs before rarefying was detected for the entire bacterial community or gram-positive bacteria.

Microbiome composition per DNA extraction protocol

We analyzed whether differences in DNA extraction methods affect our ability to assess cervical

microbiota composition. The patients can be identified by whether or not they displayed a

Lactobacillus-dominant community type (Fig. 2A). Variation between individuals was a

significantly greater influence on the observed microbial composition than was the method of

DNA extraction (Fig. 2A).

The following top 10 abundant families are shown in Fig. 2A (left) and constituted

approximately 95.7% of cervical bacteria in all kits (80 DNA extractions); *Lactobacillaceae*

(58.9%), *Bifidobacteriaceae* (13.7%), *Veillonellaceae* (4.8%), *Prevotellaceae* (4.3%), *Family XI*

(3.9%), *Atopobiaceae* (3.0%), *Leptotrichiaceae* (2.5%), *Streptococcaceae* (2.0%),

Lachnospiraceae (1.6%). *Ruminococcaceae* (0.9%). The following top 10 abundant genera are

shown in Fig. 2A (right) and constituted approximately 92% of cervical bacteria; *Lactobacillus*

(58.9%), *Gardnerella* (13.6%), *Prevotella* (4.2%), *Megasphaera* (3.7%), *Atopobium* (3.0%),

Sneathia (2.5%), *Streptococcus* (1.9%), *Parvimonas* (1.7%), *Shuttleworthia* (1.4%), and

Anaerococcus (1.1%).

Shared and unique microbiota among DNA extraction protocols

All DNA extraction methods were generally commensurate with one another, there were 31 of

41 shared microbes at the family level (Fig. 2B left) and 45 of 57 shared microbes at the genus

level (Fig. 2B right) among the DNA extraction protocols.

However, four gram-negative taxa were uniquely detected by ZymoBIOMICS and one

taxon was uniquely detected by QIAampMini both at the genus level (Fig. 2B right). Of the

uniquely detected ZymoBIOMICS OTUs, *Hydrogenophilus*, which was reported as enriched

taxa in LR-HPV positive environment (3), was detected in 14 of the 80 DNA extractions,

consisting of 2,488 reads (0.02% of all kit extractions). *Methylobacterium* was detected in 5 of

the 80 DNA extractions (912 reads; 0.01%). A member of this genus, *Methylobacterium*

aerolatum, has been reported to be more abundant in the endocervix than the vagina of healthy

South African women (61). *Bacteroidetes*, which are often reported as enriched taxa in an HIV

positive cervical environment (62), was detected in 12 of the 80 DNA extractions (1,028 reads;

0.01%). *Meiothermus* was detected in 9 of the 80 DNA extractions (882 reads; 0.01%).

Meiothermus is not considered to reside within the human environment, and may be an

extraction kit contaminant, as previously reported (63). A unique gram-positive taxa obtained

from the QIAampMini, *Streptomyces*, which was reported to be detected from the cervicovaginal

environment in the study of Kenyan women (64), was detected in 20 of 80 DNA extractions

(6,862 reads; 0.06%). No unique taxa were detected in PowerFecalPro and IndiSpin.

Venn diagrams at family levels also exhibited that ZymoBIOMICS detected slightly

more bacterial taxa (four unique taxa) as shown in Fig. 2B (left). These results showed that major

bacteria were commonly detected among all extraction protocols, with only slightly more

uniquely detected microbiota using ZymoBIOMICS.

Alpha and beta diversity

Significantly higher Species richness (`q2-breakaway`) was observed from the

ZymoBIOMICS (56.1 ± 19.4) protocol compared to that of PowerFecalPro (43.2 ± 32.9, p =

0.025), QIAampMini (54.9 ± 29.8, not significant), and IndiSpin (63.6 ± 38.3, not significant)

using Dunn's test with Benjamini-Hochberg-adjustment (Fig. 3). Similarly, Faith's Phylogenetic

Diversity was observed to be higher with the ZymoBIOMICS protocol (6.6 ± 2.2), compared to

PowerFecalPro (4.5 ± 1.9, p = 0.012), QIAampMini (5.0 ± 1.8, not significant), and IndiSpin

(5.4 ± 1.7, not significant) using Dunn's test with Benjamini-Hochberg-adjustment (Fig. 3). The

use of IndiSpin also resulted significantly higher alpha diversity than that of PowerFecalPro in an

analysis of Species richness (p = 0.042, Dunn's test with Benjamini-Hochberg-adjustment). Non-

phylogenetic alpha diversity metrics such as Observed OTUs, Shannon's diversity index, and

Pielou's Evenness did not show differences among the four methods.

ZymoBIOMICS was able to significantly increase access to several taxonomic groups

compared to the other DNA extraction methods. Additionally, as shown in Table 4,

ZymoBIOMICS did capture a different microbial composition compared to other DNA

extraction methods in the index of Unweighted UniFrac distances (PowerFecalPro: q = 0.002;

QIAampMini: q = 0.002;and IndiSpin: q = 0.002) and in Jaccard distances (QIAampMini: q =

0.018 and IndiSpin: q = 0.033).

Differential accessibility of microbiota by DNA extraction protocol

LEfSe analysis identified taxonomic groups, defined with an LDA score of 2 or higher,

for differential accessibility by extraction kit: 23 in ZymoBIOMICS, 0 in PowerFecalPro, 3 in

QIAampMini, and 3 in IndiSpin (Fig. 4A). The following taxa were found to be highly

accessible (LDA score > 3) with the use of the ZymoBIOMICS kit: Phylum *Proteobacteria*,

Class *Gammaproteobacteria*, Order *Betaproteobacteriales*, Family *Bacillaceae*, and Genus

Anoxybacillus. Whereas the Order *Streptomycetales* was highly enriched with the use of the

QIAampMini (LDA score > 3). As shown in the cladogram (Fig. 4B), despite the detection of a

potential kit contaminant (*Meiothermus sp.* a member of the Phylum *Deinococcus-Thermus*),

ZymoBIOMICS was able to increase the accessibility to additional microbiota compared to the

other extraction protocols.

Microbial community type and HPV16

DMM detected two cervical microbial community types across all four DNA extraction

protocols. Community type I was composed of the following: *Gardnerella sp.* (ZymoBIOMICS:

17.1%; PowerFecalPro: 20%; QIAampMini: 23%; IndiSpin: 20%), *Lactobacillus iners*

(ZymoBIOMICS: 6.3%; PowerFecalPro: 5%; QIAampMini: 6%; IndiSpin: 5%), *Atopobium*

vaginae (10) (ZymoBIOMICS: 3.5%; PowerFecalPro: 3%; QIAampMini: 4%; IndiSpin: 5%),

Clamydia trachomatis (ZymoBIOMICS: 1.9%; PowerFecalPro: 2%; QIAampMini: 3%;

IndiSpin: 2%), *Shuttleworthia sp.* (ZymoBIOMICS: 1.8%; PowerFecalPro: 2%; QIAampMini:

2%; IndiSpin: 2%). Some members of *Shuttleworthia* are considered to be bacterial

vaginosis☐associated bacterium (BVAB) (65), further investigation is required to determine if

this OTU is indeed a BVAB. We determined this community type "high diversity type".

Community type II was is dominated by *Lactobacillus iners* at 88%, 85%, 83%, and 85%

respectively for ZymoBIOMICS, PowerFecalPro, QIAampMini, and IndiSpin.

The relationship between HPV16 infection and community type was observed to be

significantly associated with community type I (HPV16 positive patients [n = 9], HPV16

negative patients [n = 1]) and not community type II (HPV16 positive patients [n = 1], HPV16

negative patients [n = 9], p = 0.001, Fisher's exact test) regardless of the DNA extraction kit used.

In support of this result, analysis of differentially abundant microbiota using `q2-aldex`

(Benjamini-Hochberg corrected p value of Wilcoxon test: p < 0.001, standardized distributional

effect size: −1.2) revealed that *Lactobacillus iners* were differentially enriched in the cervical

environment without HPV16. LEfSe analysis also detected that genus *Lactobacillus* were

enriched in the cervical environment without HPV16 (p < 0.001, LDA score: −5.38). No

significant differences were observed in the relationship between community type and HPV18 (p

= 0.474, Fisher's exact test), HR-HPV (p = 0.474, Fisher's exact test), results of cervical biopsy

(p = 0.554, Fisher's exact test), and race (African Americans vs not-African Americans: p = 1;

Caucasian vs not-Caucasian: p = 0.656; Hispanic vs not-Hispanic: p = 0.350, Fisher's exact test).

Discussion

In this study, we evaluated the utility of LBC specimens for the collection and storage of cervical

samples for microbiome surveys based on the 16S rRNA marker gene. We simultaneously

compared the efficacy of several commonly used DNA extraction protocols on these samples in

an effort to develop a standard operating procedure/protocol (SOP) for such work. We've also

been able to show that there are two cervical microbial community types, which are associated

with the dominance or non-dominance of *Lactobacillis iners*. Both community types were

detected regardless of the DNA extraction protocol used.

This study evaluated the composition of microbiota across all DNA extraction methods.

These findings document the importance of selecting DNA extraction methods in cervical

microbiome studies from the LBC samples. All kits were commensurate in their ability to

capture the microbial composition of each patient and the two observed cervical microbial

community state types: making all of these protocols viable for discovering broad patterns of

microbial diversity. However, we did observe that the ZymoBIOMICS protocol was better able

to access additional cervical microbiota (Fig. 2B, 4A & B). Coincidentally, we detected potential

DNA contamination only with the ZymoBIOMICS kit. The number of OTUs prior to rarefying

revealed that the ZymoBIOMICS protocol detected more gram-negative OTUs than the

PowerFecalPro (Table 3 & Fig. 2B). In particular, LEfSe analysis has shown that phylum

Proteobacteria can be better detected with the ZymoBIOMICS kit (Fig. 4). Although rarefying

microbiome data can be problematic (66), it can still provide robust and interpretable results for

diversity analysis (67), we were able to observe commensurate findings with non-rarefying

approaches such as `q2-breakaway` (50), `q2-deicode` (51), and LEfSe (48). Beta-diversity

analysis via Unweighted UniFrac also revealed that ZymoBIOMICS was significantly different

from all other kits. There were no differences in non-phylogenic indices of alpha diversity with

rarefying approaches. These findings lead us to surmise that phylogenetic indices may be more

sensitive than the non-phylogenetic indices.

Although we hypothesized that the detection of difficult-to-lyse-bacteria (*e.g*. gram-

positive bacteria) would vary by kit, we observed no significant differences (Table 3). As shown

in Table 3, the number of reads of gram-positive and gram-negative bacteria also showed that

there was no difference in the four kits. This is likely due to several modifications made to the

extraction protocol as outlined in Table 1. That is, we added bead beating and mutanolysin to the

QIAampMini protocol (68). We also modified the beating time of the ZymoBIOMICS kit down

to 2 minutes from 10 minutes (the latter being recommended by the manufacturer) to minimize

DNA shearing. We may use the extracted DNA from ZymoBIOMICS for long-read amplicon

sequencing platforms such as PacBio (Pacific Biosciences of California, Inc) (69) or MinION

(Oxford Nanopore Technologies) (70) (71). Excessive shearing can render these samples

unusable for long-read sequencing. It is quite possible that we could have observed even more

diversity with the ZymoBIOMICS kit for our amplicon survey if we conducted bead-beating for

the full 10 minutes.

Community typing and detection of the differentially abundant microbiota revealed that

Lactobacillus iners were more abundant in the cervical ecosystem without HPV16. These

findings are congruent with those of Lee *et al*. (1) and Audirac-Chalifour *et al*. (72). Lee *et al*.

reported that *Lactobacillus iners* were decreased in women with HPV positive (1). Also, the

result that the proportion of *Lactobacillus iners* was higher in HPV-negative women compared to

HPV-positive women (relative abundance 14.9% vs 2.1%) was reported by Audirac-Chalifour *et*

al (72). Similarly, Tuominen *et al*. (18) reported that *Lactobacillus iners* were enriched in HPV

negative samples (relative abundance: 47.7%) compared to HPV positive samples (relative

abundance: 18.6%, p value = 0.07) in the study of HPV positive-pregnant women (HPV16

positive rate: 15%). As established by the seminal study of Ranjeva *et al*. (73), a statistical model

revealed that colonization of specific HPV type including multi HPV type infection depends on

host-risk factors such as sexual behavior, race and ethnicity, and smoking. It is unclear whether

the association between the cervical microbiome, host-specific traits, persistent infection of

specific HPV types, such as HPV16, can be generalized and requires further investigation.

We focused on LBC samples as this is the recommended method of storage for cervical

cytology (74). Here, we confirmed that LBC samples can be used for microbial community

surveys by simply using the remaining LBC solution post HPV testing or cervical cytology. We

used a sample volume of 200 or 300 μL ThinPrep solution in this study. HPV genotyping test

using Linear Array HPV Genotyping Test (Roche Diagnostics) stably detects β-globin with a

base length of 268 bp as a positive control. Therefore, using a similar sample volume as HPV

genotyping (250 μL), it was expected that V4 (250 bp), which is near the base length of β-globin,

would be PCR amplified. It has been pointed out by Ling *et al*. (75) that the cervical

environment is of low microbial biomass. To control reagent DNA contamination and estimate

the sample volume, DNA quantification by qPCR before sequencing is recommended (76). Mitra

et al determined a sample volume of 500 μL for ThinPrep by qPCR in the microbiome study

comparing sampling methods using cytobrush or swab from cervix (19). The average storage

period from sample collection via LBC to DNA extraction was about two years in this study.

Kim *et al*. reported that DNA from cervix stored in ThinPrep at room temperature or −80°C was

stable for at least one year (77). Meanwhile, Castle *et al*. reported that β-globin DNA fragments

of 268 bases or more were detected by PCR in 90 % (27 of 30 samples) of ThinPrep samples

stored for eight years at an uncontrolled ambient temperatures followed by a controlled ambient

environment (10–26.7°C) (78). Low-temperature storage may allow the analysis of the short

DNA fragments of the V4 region after even long-term storage, although further research is

needed to confirm the optimal storage period in cervical microbiome studies using ThinPrep.

SurePath LBC specimens are as widely used as ThinPrep, but the presence of formaldehyde

within the SurePath preservation solution raises concerns about accessing enough DNA for

analysis as compared to ThinPrep, which contains methanol (79) (80). It should also be noted

that other storage solutions, *i.e*. those using guanidine thiocyanate have been reported for

microbiome surveys of the cervix (81) and feces (82). A weakness of the current study is that we

did not examine the reproducibility of our results as each sample was extracted using each kit

once. However, the use of actual patient samples rather than mock samples is a strength of our

approach.

In conclusion, regardless of the extraction protocol used, all kits provided equivalent

accessibility to the cervical microbiome. All kits shared the ability to access 31 of 41 families

and 45 of 57 genera (Fig. 2), approximately 90% of bacteria were gram-positive and 10% were

gram-negative. Observed differences in microbial composition were due to the significant

influence of the individual patient and not the extraction protocol. However, ZymoBIOMICS

was observed to increase the accessibility of DNA from a greater range of microbiota compared

to the other kits, in that the greatest number of significantly enriched taxa were identified (Fig. 3).

This was not because of higher DNA yield nor ability to detect more gram-positive bacteria.

Selection and characterization of an appropriate DNA extraction methods, for providing accurate

census of cervical microbiota, and the human microbiome in general are important (23) (24) (25)

(68) (77) (78). We have shown that the ability to characterize cervical microbiota from LBC

specimens is robust, even after prolonged storage. Our data also suggest that it is possible to

reliably assess the relationship between HPV and the cervical microbiome, also supported by

Kim *et al.* (77) and Castle *et al* (78). Even though we found all four extraction kits to be

commensurate in their ability to broadly characterize the CM, this study lends support to the

view that the selection of a DNA extraction kit depends on the questions asked of the data, and

should be taken into account for any cervicovaginal microbiome and HPV research that

leverages LBC specimens for use in clinical practice (15) (83).

Acknowledgments

We thank Togo Picture Gallery (http://togotv.dbcls.jp/pics.html) for stock images shown in Fig.

1.

This work was supported by the National Institutes of Health (R01CA143130, USA), Drs.

Mae and Anderson Nettleship Endowed Chair of Oncologic Pathology (31005156, USA), and

the Arkansas Biosciences Institute (the major component of the Tobacco Settlement Proceeds

Act of 2000, G1-52249-01, USA).

M.N. designed and supervised this project. T.S. and M.S.R. conducted bioinformatics

analysis and wrote paper. T.S., H.C., and M.N. created the protocol of DNA extraction. M.N.,

H.C., S.O., W.G., and T.S. provided important feedback. Samples in the clinical trial were

collected by W.G. and his associates. DNA extraction was conducted by T.S. Sequencing of 16S

RNA gene was conducted by S.O.

M.N. is one of the inventors named in the patents and patent applications for the HPV

therapeutic vaccine PepCan. The remaining authors declare no conflicts of interest.

References

1. Lee JE, Lee S, Lee H, Song YM, Lee K, Han MJ, Sung J, Ko G. 2013. Association of the vaginal microbiota with human papillomavirus infection in a Korean twin cohort. PLoS One 8:e63514.
2. Huang X, Li C, Li F, Zhao J, Wan X, Wang K. 2018. Cervicovaginal microbiota composition correlates with the acquisition of high-risk human papillomavirus types. Int J Cancer 143:621-634.
3. Zhou Y, Wang L, Pei F, Ji M, Zhang F, Sun Y, Zhao Q, Hong Y, Wang X, Tian J, Wang Y. 2019. Patients With LR-HPV Infection Have a Distinct Vaginal Microbiota in Comparison With Healthy Controls. Front Cell Infect Microbiol 9:294.
4. Onywera H, Williamson AL, Mbulawa ZZA, Coetzee D, Meiring TL. 2019. The cervical microbiota in reproductive-age South African women with and without human papillomavirus infection. Papillomavirus Res 7:154-163.
5. Brotman RM, Shardell MD, Gajer P, Tracy JK, Zenilman JM, Ravel J, Gravitt PE. 2014. Interplay between the temporal dynamics of the vaginal microbiota and human papillomavirus detection. J Infect Dis 210:1723-33.
6. Godoy-Vitorino F, Romaguera J, Zhao C, Vargas-Robles D, Ortiz-Morales G, Vazquez-Sanchez F, Sanchez-Vazquez M, de la Garza-Casillas M, Martinez-Ferrer M, White JR, Bittinger K, Dominguez-Bello MG, Blaser MJ. 2018. Cervicovaginal Fungi and Bacteria Associated With Cervical Intraepithelial Neoplasia and High-Risk Human Papillomavirus Infections in a Hispanic Population. Front Microbiol 9:2533.
7. Łaniewski P, Barnes D, Goulder A, Cui H, Roe DJ, Chase DM, Herbst-Kralovetz MM. 2018. Linking cervicovaginal immune signatures, HPV and microbiota composition in cervical carcinogenesis in non-Hispanic and Hispanic women, Sci Rep, vol 8.
8. Mitra A, MacIntyre DA, Lee YS, Smith A, Marchesi JR, Lehne B, Bhatia R, Lyons D, Paraskevaidis E, Li JV, Holmes E, Nicholson JK, Bennett PR, Kyrgiou M. 2015. Cervical intraepithelial neoplasia disease progression is associated with increased vaginal microbiome diversity. Sci Rep 5:16865.
9. Piyathilake CJ, Ollberding NJ, Kumar R, Macaluso M, Alvarez RD, Morrow CD. 2016. Cervical Microbiota Associated with Higher Grade Cervical Intraepithelial Neoplasia in Women Infected with High-Risk Human Papillomaviruses. Cancer Prev Res (Phila) 9:357-66.
10. Oh HY, Kim BS, Seo SS, Kong JS, Lee JK, Park SY, Hong KM, Kim HK, Kim MK. 2015. The association of uterine cervical microbiota with an increased risk for cervical intraepithelial neoplasia in Korea. Clin Microbiol Infect 21:674 e1-9.
11. De Seta F, Campisciano G, Zanotta N, Ricci G, Comar M. 2019. The Vaginal Community State Types Microbiome-Immune Network as Key Factor for Bacterial Vaginosis and Aerobic Vaginitis. Front Microbiol 10:2451.
12. Oliver A, LaMere B, Weihe C, Wandro S, Lindsay KL, Wadhwa PD, Mills DA, Pride D, Fiehn O, Northen T, de Raad M, Li H, Martiny JBH, Lynch S, Whiteson K. 2019. Cervicovaginal microbiome composition drives metabolic profiles in healthy pregnancy. bioRxiv https://doi.org/10.1101/840520.

13. Human Microbiome Project Consortium. 2012. Structure, function and diversity of the
healthy human microbiome. Nature 486:207-14.
14. Bik EM, Bird SW, Bustamante JP, Leon LE, Nieto PA, Addae K, Alegria-Mera V, Bravo C,
Bravo D, Cardenas JP, Carson GA, Caughey A, Covarrubias PC, Perez-Donoso J, Gass G,
Gupta SL, Harman K, Hongo DMB, Jimenez JC, Kraal L, Melis-Arcos F, Morales EH,
Morton A, Navas CF, Nunez H, Olivares E, Ordenes-Aenishanslins N, Ossandon FJ, Phan R,
Pino R, Soto-Liebe K, Varas I, Vera-Wolf P, Walton NA, Almonacid DE, Goddard AD,
Ugalde JA, Zneimer S, Richman J, Apte ZS. 2019. A novel sequencing-based vaginal
health assay combining self-sampling, HPV detection and genotyping, STI detection, and
vaginal microbiome analysis. PLoS One 14:e0215945.
15. Berman HL, McLaren MR, Callahan BJ. 2019. Understanding and Interpreting Community
Sequencing Measurements of the Vaginal Microbiome. BJOG doi:10.1111/1471-
0528.15978.
16. Ravel J, Gajer P, Abdo Z, Schneider GM, Koenig SS, McCulle SL, Karlebach S, Gorle R,
Russell J, Tacket CO, Brotman RM, Davis CC, Ault K, Peralta L, Forney LJ. 2011. Vaginal
microbiome of reproductive-age women. Proc Natl Acad Sci U S A 108 Suppl 1:4680-7.
17. Fettweis JM, Serrano MG, Brooks JP, Edwards DJ, Girerd PH, Parikh HI, Huang B, Arodz TJ,
Edupuganti L, Glascock AL, Xu J, Jimenez NR, Vivadelli SC, Fong SS, Sheth NU, Jean S, Lee
V, Bokhari YA, Lara AM, Mistry SD, Duckworth RA, 3rd, Bradley SP, Koparde VN, Orenda
XV, Milton SH, Rozycki SK, Matveyev AV, Wright ML, Huzurbazar SV, Jackson EM,
Smirnova E, Korlach J, Tsai YC, Dickinson MR, Brooks JL, Drake JI, Chaffin DO, Sexton AL,
Gravett MG, Rubens CE, Wijesooriya NR, Hendricks-Munoz KD, Jefferson KK, Strauss JF,
3rd, Buck GA. 2019. The vaginal microbiome and preterm birth. Nat Med 25:1012-1021.
18. Tuominen H, Rautava S, Syrjanen S, Collado MC, Rautava J. 2018. HPV infection and
bacterial microbiota in the placenta, uterine cervix and oral mucosa. Sci Rep 8:9787.
19. Mitra A, MacIntyre DA, Mahajan V, Lee YS, Smith A, Marchesi JR, Lyons D, Bennett PR,
Kyrgiou M. 2017. Comparison of vaginal microbiota sampling techniques: cytobrush
versus swab. Sci Rep 7:9802.
20. Bentz JS. 2005. Liquid-based cytology for cervical cancer screening. Expert Rev Mol
Diagn 5:857-71.
21. Gibb RK, Martens MG. 2011. The impact of liquid-based cytology in decreasing the
incidence of cervical cancer. Rev Obstet Gynecol 4:S2-S11.
22. Donders GG, Depuydt CE, Bogers JP, Vereecken AJ. 2013. Association of Trichomonas
vaginalis and cytological abnormalities of the cervix in low risk women. PLoS One
8:e86266.
23. Costea PI, Zeller G, Sunagawa S, Pelletier E, Alberti A, Levenez F, Tramontano M,
Driessen M, Hercog R, Jung FE, Kultima JR, Hayward MR, Coelho LP, Allen-Vercoe E,
Bertrand L, Blaut M, Brown JRM, Carton T, Cools-Portier S, Daigneault M, Derrien M,
Druesne A, de Vos WM, Finlay BB, Flint HJ, Guarner F, Hattori M, Heilig H, Luna RA, van
Hylckama Vlieg J, Junick J, Klymiuk I, Langella P, Le Chatelier E, Mai V, Manichanh C,
Martin JC, Mery C, Morita H, O'Toole PW, Orvain C, Patil KR, Penders J, Persson S, Pons
N, Popova M, Salonen A, Saulnier D, Scott KP, Singh B, et al. 2017. Towards standards for
human fecal sample processing in metagenomic studies. Nat Biotechnol 35:1069-1076.

24.　Stinson LF, Keelan JA, Payne MS. 2018. Comparison of Meconium DNA Extraction
Methods for Use in Microbiome Studies. Front Microbiol 9:270.
25.　Teng F, Darveekaran Nair SS, Zhu P, Li S, Huang S, Li X, Xu J, Yang F. 2018. Impact of DNA
extraction method and targeted 16S-rRNA hypervariable region on oral microbiota
profiling. Sci Rep 8:16321.
26.　Ravilla R, Coleman HN, Chow CE, Chan L, Fuhrman BJ, Greenfield WW, Robeson MS,
Iverson K, Spencer H, 3rd, Nakagawa M. 2019. Cervical Microbiome and Response to a
Human Papillomavirus Therapeutic Vaccine for Treating High-Grade Cervical Squamous
Intraepithelial Lesion. Integr Cancer Ther 18:1534735419893063.
27.　Virtanen S, Kalliala I, Nieminen P, Salonen A. 2017. Comparative analysis of vaginal
microbiota sampling using 16S rRNA gene analysis. PLoS One 12:e0181477.
28.　Kim D, Hofstaedter CE, Zhao C, Mattei L, Tanes C, Clarke E, Lauder A, Sherrill-Mix S,
Chehoud C, Kelsen J, Conrad M, Collman RG, Baldassano R, Bushman FD, Bittinger K.
2017. Optimizing methods and dodging pitfalls in microbiome research. Microbiome
5:52.
29.　Thompson LR, Sanders JG, McDonald D, Amir A, Ladau J, Locey KJ, Prill RJ, Tripathi A,
Gibbons SM, Ackermann G, Navas-Molina JA, Janssen S, Kopylova E, Vazquez-Baeza Y,
Gonzalez A, Morton JT, Mirarab S, Zech Xu Z, Jiang L, Haroon MF, Kanbar J, Zhu Q, Jin
Song S, Kosciolek T, Bokulich NA, Lefler J, Brislawn CJ, Humphrey G, Owens SM,
Hampton-Marcell J, Berg-Lyons D, McKenzie V, Fierer N, Fuhrman JA, Clauset A, Stevens
RL, Shade A, Pollard KS, Goodwin KD, Jansson JK, Gilbert JA, Knight R, Earth Microbiome
Project C. 2017. A communal catalogue reveals Earth's multiscale microbial diversity.
Nature 551:457-463.
30.　Apprill A, McNally S, Parsons R, Weber L. 2015. Minor revision to V4 region SSU rRNA
806R gene primer greatly increases detection of SAR11 bacterioplankton. Aquat Microb
Ecol 75:129-137.
31.　Parada AE, Needham DM, Fuhrman JA. 2016. Every base matters: assessing small
subunit rRNA primers for marine microbiomes with mock communities, time series and
global field samples. Environ Microbiol 18:1403-14.
32.　Walters W, Hyde ER, Berg-Lyons D, Ackermann G, Humphrey G, Parada A, Gilbert JA,
Jansson JK, Caporaso JG, Fuhrman JA, Apprill A, Knight R. 2016. Improved Bacterial 16S
rRNA Gene (V4 and V4-5) and Fungal Internal Transcribed Spacer Marker Gene Primers
for Microbial Community Surveys. mSystems 1.
33.　Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, Alexander H, Alm
EJ, Arumugam M, Asnicar F, Bai Y, Bisanz JE, Bittinger K, Brejnrod A, Brislawn CJ, Brown
CT, Callahan BJ, Caraballo-Rodriguez AM, Chase J, Cope EK, Da Silva R, Diener C,
Dorrestein PC, Douglas GM, Durall DM, Duvallet C, Edwardson CF, Ernst M, Estaki M,
Fouquier J, Gauglitz JM, Gibbons SM, Gibson DL, Gonzalez A, Gorlick K, Guo J, Hillmann B,
Holmes S, Holste H, Huttenhower C, Huttley GA, Janssen S, Jarmusch AK, Jiang L, Kaehler
BD, Kang KB, Keefe CR, Keim P, Kelley ST, Knights D, et al. 2019. Reproducible,
interactive, scalable and extensible microbiome data science using QIIME 2. Nat
Biotechnol 37:852-857.
34.　Callahan BJ, McMurdie PJ, Holmes SP. 2017. Exact sequence variants should replace
operational taxonomic units in marker-gene data analysis. ISME J 11:2639-2643.

35.   Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJ, Holmes SP. 2016. DADA2:
604   High-resolution sample inference from Illumina amplicon data. Nat Methods 13:581-3.
605   36.   Bokulich NA, Kaehler BD, Rideout JR, Dillon M, Bolyen E, Knight R, Huttley GA, Gregory
606   Caporaso J. 2018. Optimizing taxonomic classification of marker-gene amplicon
607   sequences with QIIME 2's q2-feature-classifier plugin. Microbiome 6:90.
608   37.   Werner JJ, Koren O, Hugenholtz P, DeSantis TZ, Walters WA, Caporaso JG, Angenent LT,
609   Knight R, Ley RE. 2012. Impact of training sets on classification of high-throughput
610   bacterial 16s rRNA gene surveys. ISME J 6:94-103.
611   38.   Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glockner FO. 2013.
612   The SILVA ribosomal RNA gene database project: improved data processing and web-
613   based tools. Nucleic Acids Res 41:D590-6.
614   39.   Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7:
615   improvements in performance and usability. Mol Biol Evol 30:772-80.
616   40.   Nguyen LT, Schmidt HA, von Haeseler A, Minh BQ. 2015. IQ-TREE: a fast and effective
617   stochastic algorithm for estimating maximum-likelihood phylogenies. Mol Biol Evol
618   32:268-74.
619   41.   Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. 2017.
620   ModelFinder: fast model selection for accurate phylogenetic estimates. Nat Methods
621   14:587-589.
622   42.   Bokulich NA, Subramanian S, Faith JJ, Gevers D, Gordon JI, Knight R, Mills DA, Caporaso
623   JG. 2013. Quality-filtering vastly improves diversity estimates from Illumina amplicon
624   sequencing. Nat Methods 10:57-9.
625   43.   McMurdie PJ, Holmes S. 2013. phyloseq: an R package for reproducible interactive
626   analysis and graphics of microbiome census data. PLoS One 8:e61217.
627   44.   Bisanz JE. 2018. qiime2R: Importing QIIME2 artifacts and associated data into R sessions.
628   https://github.com/jbisanz/qiime2R.
629   45.   Lahti L, Shetty S. 2012-2019. microbiome R package. http://microbiome.github.io.
630   46.   Anderson MJ. 2001. A new method for non-parametric multivariate analysis of variance.
631   Austral Ecol 26:32-46.
632   47.   Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, Minchin PR, O'Hara
633   RB, Simpson GL, Solymos P, Stevens MHH, Szoecs E, Wagner H. 2019. vegan: Community
634   Ecology Package. 2018. R package version 2.5-3. https://CRAN.R-
635   project.org/package=vegan.
636   48.   Segata N, Izard J, Waldron L, Gevers D, Miropolsky L, Garrett WS, Huttenhower C. 2011.
637   Metagenomic biomarker discovery and explanation. Genome Biol 12:R60.
638   49.   Bardou P, Mariette J, Escudie F, Djemiel C, Klopp C. 2014. jvenn: an interactive Venn
639   diagram viewer. BMC Bioinformatics 15:293.
640   50.   Willis A, Bunge J. 2015. Estimating diversity via frequency ratios. Biometrics 71:1042-9.
641   51.   Martino C, Morton JT, Marotz CA, Thompson LR, Tripathi A, Knight R, Zengler K. 2019. A
642   Novel Sparse Compositional Technique Reveals Microbial Perturbations. mSystems 4.
643   52.   Gao W, Weng J, Gao Y, Chen X. 2013. Comparison of the vaginal microbiota diversity of
644   women with and without human papillomavirus infection: a cross-sectional study. BMC
645   Infect Dis 13:271.

53.   Montealegre JR, Peckham-Gregory EC, Marquez-Do D, Dillon L, Guillaud M, Adler-
Storthz K, Follen M, Scheurer ME. 2018. Racial/ethnic differences in HPV 16/18
genotypes and integration status among women with a history of cytological
abnormalities. Gynecol Oncol 148:357-362.
54.   Xi LF, Kiviat NB, Hildesheim A, Galloway DA, Wheeler CM, Ho J, Koutsky LA. 2006.
Human papillomavirus type 16 and 18 variants: race-related distribution and persistence.
J Natl Cancer Inst 98:1045-52.
55.   Morgan M. 2019. DirichletMultinomial: Dirichlet-Multinomial Mixture Model Machine
Learning for Microbiome Data.
http://bioconductor.org/packages/release/bioc/html/DirichletMultinomial.html.
56.   Holmes I, Harris K, Quince C. 2012. Dirichlet multinomial mixtures: generative models
for microbial metagenomics. PLoS One 7:e30126.
57.   DiGiulio DB, Callahan BJ, McMurdie PJ, Costello EK, Lyell DJ, Robaczewska A, Sun CL,
Goltsman DS, Wong RJ, Shaw G, Stevenson DK, Holmes SP, Relman DA. 2015. Temporal
and spatial variation of the human microbiota during pregnancy. Proc Natl Acad Sci U S
A 112:11060-5.
58.   Fernandes AD, Macklaim JM, Linn TG, Reid G, Gloor GB. 2013. ANOVA-like differential
expression (ALDEx) analysis for mixed population RNA-Seq. PLoS One 8:e67019.
59.   Dinno A. 2017. dunn.test: Dunn's Test of Multiple Comparisons Using Rank Sums.
https://CRAN.R-project.org/package=dunn.test.
60.   Yilmaz P, Kottmann R, Field D, Knight R, Cole JR, Amaral-Zettler L, Gilbert JA, Karsch-
Mizrachi I, Johnston A, Cochrane G, Vaughan R, Hunter C, Park J, Morrison N, Rocca-
Serra P, Sterk P, Arumugam M, Bailey M, Baumgartner L, Birren BW, Blaser MJ, Bonazzi
V, Booth T, Bork P, Bushman FD, Buttigieg PL, Chain PS, Charlson E, Costello EK, Huot-
Creasy H, Dawyndt P, DeSantis T, Fierer N, Fuhrman JA, Gallery RE, Gevers D, Gibbs RA,
San Gil I, Gonzalez A, Gordon JI, Guralnick R, Hankeln W, Highlander S, Hugenholtz P,
Jansson J, Kau AL, Kelley ST, Kennedy J, Knights D, Koren O, et al. 2011. Minimum
information about a marker gene sequence (MIMARKS) and minimum information
about any (x) sequence (MIxS) specifications. Nat Biotechnol 29:415-20.
61.   Balle C, Lennard K, Dabee S, Barnabas SL, Jaumdally SZ, Gasper MA, Maseko V, Mbulawa
ZZA, Williamson AL, Bekker LG, Lewis DA, Passmore JS, Jaspan HB. 2018. Endocervical
and vaginal microbiota in South African adolescents with asymptomatic Chlamydia
trachomatis infection. Sci Rep 8:11109.
62.   Klein C, Gonzalez D, Samwel K, Kahesa C, Mwaiselage J, Aluthge N, Fernando S, West JT,
Wood C, Angeletti PC. 2019. Relationship between the Cervical Microbiome, HIV Status,
and Precancerous Lesions. MBio 10.
63.   Glassing A, Dowd SE, Galandiuk S, Davis B, Chiodini RJ. 2016. Inherent bacterial DNA
contamination of extraction and sequencing reagents may affect interpretation of
microbiota in low bacterial biomass samples, Gut Pathog, vol 8.
64.   Birse KD, Romas LM, Guthrie BL, Nilsson P, Bosire R, Kiarie J, Farquhar C, Broliden K,
Burgener AD. 2017. Genital Injury Signatures and Microbiome Alterations Associated
With Depot Medroxyprogesterone Acetate Usage and Intravaginal Drying Practices. J
Infect Dis 215:590-598.

65. Lennard K, Dabee S, Barnabas SL, Havyarimana E, Blakney A, Jaumdally SZ, Botha G,
Mkhize NN, Bekker LG, Lewis DA, Gray G, Mulder N, Passmore JS, Jaspan HB. 2018.
Microbial Composition Predicts Genital Tract Inflammation and Persistent Bacterial
Vaginosis in South African Adolescent Females. Infect Immun 86.
66. McMurdie PJ, Holmes S. 2014. Waste not, want not: why rarefying microbiome data is
inadmissible. PLoS Comput Biol 10:e1003531.
67. Weiss S, Xu ZZ, Peddada S, Amir A, Bittinger K, Gonzalez A, Lozupone C, Zaneveld JR,
Vazquez-Baeza Y, Birmingham A, Hyde ER, Knight R. 2017. Normalization and microbial
differential abundance strategies depend upon data characteristics. Microbiome 5:27.
68. Yuan S, Cohen DB, Ravel J, Abdo Z, Forney LJ. 2012. Evaluation of methods for the
extraction and purification of DNA from the human microbiome. PLoS One 7:e33865.
69. Callahan BJ, Wong J, Heiner C, Oh S, Theriot CM, Gulati AS, McGill SK, Dougherty MK.
2019. High-throughput amplicon sequencing of the full-length 16S rRNA gene with
single-nucleotide resolution. Nucleic Acids Res 47:e103.
70. Calus ST, Ijaz UZ, Pinto AJ. 2018. NanoAmpli-Seq: a workflow for amplicon sequencing
for mixed microbial communities on the nanopore sequencing platform. Gigascience 7.
71. Wongsurawat T, Nakagawa M, Atiq O, Coleman HN, Jenjaroenpun P, Allred JI, Trammel
A, Puengrang P, Ussery DW, Nookaew I. 2019. An assessment of Oxford Nanopore
sequencing for human gut metagenome profiling: A pilot study of head and neck cancer
patients. J Microbiol Methods 166:105739.
72. Audirac-Chalifour A, Torres-Poveda K, Bahena-Roman M, Tellez-Sosa J, Martinez-
Barnetche J, Cortina-Ceballos B, Lopez-Estrada G, Delgado-Romero K, Burguete-Garcia
AI, Cantu D, Garcia-Carranca A, Madrid-Marina V. 2016. Cervical Microbiome and
Cytokine Profile at Various Stages of Cervical Cancer: A Pilot Study. PLoS One
11:e0153274.
73. Ranjeva SL, Mihaljevic JR, Joseph MB, Giuliano AR, Dwyer G. 2019. Untangling the
dynamics of persistence and colonization in microbial communities. ISME J
doi:10.1038/s41396-019-0488-7:1-13.
74. Linder J, Zahniser D. 1998. ThinPrep Papanicolaou testing to reduce false-negative
cervical cytology. Arch Pathol Lab Med 122:139-44.
75. Ling Z, Liu X, Chen X, Zhu H, Nelson KE, Xia Y, Li L, Xiang C. 2011. Diversity of
cervicovaginal microbiota associated with female lower genital tract infections. Microb
Ecol 61:704-14.
76. Salter SJ, Cox MJ, Turek EM, Calus ST, Cookson WO, Moffatt MF, Turner P, Parkhill J,
Loman NJ, Walker AW. 2014. Reagent and laboratory contamination can critically
impact sequence-based microbiome analyses. BMC Biol 12:87.
77. Kim Y, Choi KR, Chae MJ, Shin BK, Kim HK, Kim A, Kim BH. 2013. Stability of DNA, RNA,
cytomorphology, and immunoantigenicity in Residual ThinPrep Specimens. APMIS
121:1064-72.
78. Castle PE, Solomon D, Hildesheim A, Herrero R, Concepcion Bratti M, Sherman ME,
Cecilia Rodriguez A, Alfaro M, Hutchinson ML, Terence Dunn S, Kuypers J, Schiffman M.
2003. Stability of archived liquid-based cervical cytologic specimens. Cancer 99:89-96.

79.   Rebolj M, Rask J, van Ballegooijen M, Kirschner B, Rozemeijer K, Bonde J, Rygaard C,
732   Lynge E. 2015. Cervical histology after routine ThinPrep or SurePath liquid-based
733   cytology and computer-assisted reading in Denmark. Br J Cancer 113:1259-74.
734   80.   Naeem RC, Goldstein DY, Einstein MH, Ramos Rivera G, Schlesinger K, Khader SN,
735   Suhrland M, Fox AS. 2017. SurePath Specimens Versus ThinPrep Specimen Types on the
736   COBAS 4800 Platform: High-Risk HPV Status and Cytology Correlation in an Ethnically
737   Diverse Bronx Population. Lab Med 48:207-213.
738   81.   Ritu W, Enqi W, Zheng S, Wang J, Ling Y, Wang Y. 2019. Evaluation of the Associations
739   Between Cervical Microbiota and HPV Infection, Clearance, and Persistence in
740   Cytologically Normal Women. Cancer Prev Res (Phila) 12:43-56.
741   82.   Hosomi K, Ohno H, Murakami H, Natsume-Kitatani Y, Tanisawa K, Hirata S, Suzuki H,
742   Nagatake T, Nishino T, Mizuguchi K, Miyachi M, Kunisawa J. 2017. Method for preparing
743   DNA from feces in guanidine thiocyanate solution affects 16S rRNA-based profiling of
744   human microbiota diversity. Sci Rep 7:4339.
745   83.   Sarangi AN, Goel A, Aggarwal R. 2019. Methods for Studying Gut Microbiota: A Primer
746   for Physicians. J Clin Exp Hepatol 9:62-73.
747   84.   Silhavy TJ, Kahne D, Walker S. 2010. The bacterial cell envelope. Cold Spring Harb
748   Perspect Biol 2:a000414.
749

Table 1: Characteristics of four different DNA extraction protocols

Kit (Cat. No.)	Manufacturer	Sample volume	Enzyme	Beads	Beating	DNA carrier	Others
ZymoBIOMICS DNA Miniprep Kit (D4300)	Zymo Research	300 μL	No	Ceramic^a	2 min^b	No	^{c}
QIAamp PowerFecal Pro DNA Kit (51804)	Qiagen	300 μL	No	Ceramic^d	10 min^b	No	^{c}
QIAamp DNA Mini Kit (51304)	Qiagen	200 μL	Mutanolysin^e	No	No	No	$^{c, f, g}$
IndiSpin Pathogen Kit (SPS4104)	Indical Bioscience	300 μL	No	Ceramic^h	10 min^b	Yes	$^{c, i}$

a: https://www.zymoresearch.com/pages/microbial-isolation. b: Disruptor Genie (USA Scientific, Inc.) was used under the maximum speed. c: Nuclease free water (85 μL) as DNA elution buffer was used. d: PowerBead Pro Tubes (https://www.qiagen.com/us/products/discovery-and-translational-research/lab-essentials/plastics/powerbead-tubes/#orderinginformation). e: Instead of lysozyme or lysostaphin, mutanolysin was used as per Yuan *et al*, 2012 (68). f: DNA Purification from Blood or Body Fluids; Protocols for Bacteria; Isolation of genomic DNA from gram-positive bacteria in QIAamp DNA Mini and Blood Mini Handbook fifth edition was referenced. g: Heating at 56°C for 30 min and 95°C for 15 min was performed. h: Pathogen Lysis Tubes S (https://www.qiagen.com/dk/shop/pcr/pathogen-lysis-tubes/). i: Pretreatment B2 as per QIAamp cador Pathogen Mini Handbook.

Table 2. Patients characteristics

Characteristics		Values
Number of patients, n		20
Total number of DNA extracts, n		80
Age, mean (SD)		31.4 (5.0)
Race	African American, n (%)	3 (15)
	Caucasian, n (%)	10 (50)
	Hispanic, n (%)	7 (35)
Cervical biopsy	CIN2, n (%)	8 (40)
	CIN3, n (%)	10 (50)
	Benign, n (%)	2 (10)
HPV typing	HPV positive, n (%)	19 (95)
	HPV16 positive, n (%)	10 (50)
	HPV18 positive, n (%)	2 (10)
	HPV16 or 18 positives, n (%)	10 (50)
	HR-HPV positives, n (%)	18 (90)

SD: standard deviation. CIN: cervical intraepithelial neoplasia. HR-HPV: high-risk HPV (HPV16 18, 31, 33, 35, 39, 45, 51, 52, 56, 58, 59, and 68)

Table 3. Reads and OTUs before rarefying assigned to all, gram-, and gram-negative bacteria per DNA extraction protocols

Parameters	Community	Methods	Values	Ratio of GP or GN	p value
Number of reads (mean ± SD)	All	Zy	2,705,044 (135,252 ± 66,011)		[a]
		Pro	2,312,207 (115,610 ± 68,201)		
		QIA	2,765,343 (138,267 ± 49,781)		
		IN	3,366,988 (168,349 ± 57,451)		
	GP	Zy	2,430,380 (121,519 ± 56,209)	89.8%	NS
		Pro	2,116,458 (105,823 ± 57,590)	91.5%	
		QIA	2,503,578 (125,179 ± 46,073)	90.5%	
		IN	2,985,941 (149,297 ± 46,936)	88.7%	
	GN	Zy	274,664 (13,733 ± 29,162)	10.2%	NS
		Pro	195,749 (9,788 ± 23,070)	8.5%	
		QIA	261,765 (13,088 ± 22,638)	9.5%	
		IN	381,047 (19,052 ± 33,038)	11.3%	
Number of OTUs (mean ± SD)	All	Zy	825 (41.3 ± 16.8)		NS
		Pro	621 (31.1 ± 19.4)		
		QIA	778 (38.9 ± 22.4)		
		IN	792 (39.6 ± 22.7)		
	GP	Zy	479 (24.0 ± 9.2)	58.1%	NS
		Pro	412 (20.6 ± 12.7)	66.3%	
		QIA	513 (25.7 ± 13.7)	65.9%	
		IN	531 (26.6 ± 14.9)	67.0%	
	GN	Zy	346 (17.3 ± 9.8)	41.9%	[b]
		Pro	209 (10.5 ± 10.3)	33.7%	
		QIA	265 (13.3 ± 9.2)	34.1%	
		IN	261 (13.1 ± 8.3)	33.0%	

Community of gram-positive bacteria were defined as phylum *Actinobacteria* and *Firmicutes*, which are composed of thick peptidoglycan layers without outer membrane (84). Community of gram-negative bacteria was defined as a community of bacteria other than phylum *Actinobacteria* and *Firmicutes* in this study. a: I - P: 0.0199; I - Q: 0.1590; P - Q: 0.1436; I - Z: 0.1495; P - Z: 0.1712; and Q - Z: 0.4059. b: I - P: 0.2116; I - Q: 0.4837; P - Q: 0.1143; I - Z: 0.0938; P - Z: 0.0116; Q - Z: 0.1448. Dunn's test with Benjamini-Hochberg-adjustment were performed for comparison of the number of read and OTU by DNA extraction method. Zy: ZymoBIOMICS DNA Miniprep Kit, Pro: QIAamp PowerFecal Pro DNA Kit, QIA: QIAamp DNA Mini Kit, IN: IndiSpin Pathogen Kit. SD: standard deviation. All: all bacteria, GP: gram-positive bacteria, GN: gram-negative bacteria. NS: not significant.

Table 4. Beta diversity among DNA extraction methods

Index	Protocol	Protocols compared	p values	q values
Aitchison distance	ZymoBIOMICS	PowerFecalPro	NS	NS
		QIAampMini	NS	NS
		IndiSpin	NS	NS
	PowerFecalPro	QIAampMini	NS	NS
		IndiSpin	NS	NS
	QIAampMini	IndiSpin	NS	NS
Unweighted UniFrac distance	ZymoBIOMICS	PowerFecalPro	0.001	0.002
		QIAampMini	0.001	0.002
		IndiSpin	0.001	0.002
	PowerFecalPro	QIAampMini	NS	NS
		IndiSpin	0.015	0.023
	QIAampMini	IndiSpin	NS	NS
Weighted UniFrac distance	ZymoBIOMICS	PowerFecalPro	NS	NS
		QIAampMini	NS	NS
		IndiSpin	NS	NS
	PowerFecalPro	QIAampMini	NS	NS
		IndiSpin	NS	NS
	QIAampMini	IndiSpin	NS	NS
Jaccard distance	ZymoBIOMICS	PowerFecalPro	0.037	NS
		QIAampMini	0.003	0.018
		IndiSpin	0.011	0.033
	PowerFecalPro	QIAampMini	NS	NS
		IndiSpin	NS	NS
	QIAampMini	IndiSpin	NS	NS
Bray-Curtis distance	ZymoBIOMICS	PowerFecalPro	NS	NS

	QIAampMini	NS	NS
	IndiSpin	NS	NS
PowerFecalPro	QIAampMini	NS	NS
	IndiSpin	NS	NS
QIAampMini	IndiSpin	NS	NS

Pairwise PERMANOVA was tested for comparing beta diversity of DNA extraction method. NS: not significant.

Table S1. Diversity analysis in this study

No.	Parameter	Alpha or Beta diversity	Used data with/without rarefying	Input data with/without phylogenetic information	Plugin of QIIME 2
1	Species richness	Alpha	Not rarefied	Non-phylogenetic	q2-breakaway (50)
2	Faith's Phylogenetic Diversity	Alpha	Rarefied	Phylogenetic	q2-diversity
3	Observed OTUs	Alpha	Rarefied	Non-phylogenetic	q2-diversity
4	Shannon's diversity index	Alpha	Rarefied	Non-phylogenetic	q2-diversity
5	Pielou's Evenness	Alpha	Rarefied	Non-phylogenetic	q2-diversity
6	Aitchison distance	Beta	Not rarefied	Non-phylogenetic	q2-deicode (51)
7	Unweighted UniFrac distance	Beta	Rarefied	Phylogenetic	q2-diversity
8	Weighted UniFrac distance	Beta	Rarefied	Phylogenetic	q2-diversity
9	Jaccard distance	Beta	Rarefied	Non-phylogenetic	q2-diversity
10	Bray-Curtis distances	Beta	Rarefied	Non-phylogenetic	q2-diversity
11	Adonis	Beta	Rarefied	Non-phylogenetic	q2-diversity adonis (46) (47)

Figure legends

Fig. 1. Overview of the study design using the 16S rRNA gene to compare the DNA

extraction protocol for cervical microbiota. (A) Liquid-based cytology (LBC) specimens from

20 patients with CIN2/3 or suspected CIN2/3. (B) A total of 80 DNA extractions were performed.

(C) The four DNA extraction methods. (D) DNA of mock vaginal community as a positive

control and preservation solution as a negative control. (E) Sequencing using Illumina MiSeq.

(F) Analysis of the taxonomic profiles among the DNA extraction protocols. Images form Togo

Picture Gallery (http://togotv.dbcls.jp/ja/pics.html) were used to create this figure.

Fig. 2. Taxonomic resolution among DNA extraction protocols. (A) Relative abundance of

microbe at family level (left) and genus level (right) per DNA extraction method showed the

pattern that variance of microbe composition per patient was higher than that per DNA extraction

protocol. These pattern were confirmed by values of Adonis test (`q2-diversity adonis`);

F.Model: 199.4, R2: 0.982, and p value: 0.001 for patients and F.Model: 2.9, R2: 0.003, and p

value: 0.002 for DNA extraction (46) (47). After all count data of taxonomy were converted to

relative abundance as shown in the y-axis, the top ten taxonomy at each family and genus level

were plotted in colored bar plot and other relatively few taxonomies were not plotted. The 20

patients ID were described in the x-axis. (B) Venn diagrams showed that ZymoBIOMICS had

four unique taxa at family (left) and genus (right) taxonomic level. Thirty-one of 41 families and

45 of 57 genera were detected with all DNA extraction protocols.

Fig. 3. Comparisons of alpha diversity between different DNA extraction protocols. The

alpha diversity indices determined by Species richness and Phylogenetic diversity are

significantly higher with ZymoBIOMICS in comparison with PowerFecalPro (p = 0.025 and

0.012, respectively, Dunn's test with Benjamini-Hochberg-adjustment). IndiSpin also showed

significantly higher diversity than that of PowerFecalPro using analysis of Species richness (p =

0.042, Dunn's test with Benjamini-Hochberg-adjustment). No significant differences were

observed in other alpha diversity indexes such as observed OTUs, Shannon's diversity index,

and Pielou's Evenness. Zy: ZymoBIOMICS DNA Miniprep Kit, Pro: QIAamp PowerFecal Pro

DNA Kit, QIA: QIAamp DNA Mini Kit, IN: IndiSpin Pathogen Kit.

Fig. 4. Distinct detections of microbe among the DNA extraction protocols. (A) A bar graph

showing 23 significantly enriched taxa with ZymoBIOMICS, 3 with QIAamp DNA Mini Kit,

and 3 with IndiSpin Pathogen Kit determined by the linear discriminant analysis (LDA) effect

size (LEfSe) analyses (48). (B) A taxonomic cladogram from the same LEfSe analyses showing

that the significantly enriched microbiota in ZymoBIOMICS were composed of phylum

Proteobacteria. Also note that *Meiothermus (a member of* the phylum *Deinococcus-Thermus)* is

likely an extraction kit contaminant. Zy: ZymoBIOMICS DNA Miniprep Kit, Pro: QIAamp

PowerFecal Pro DNA Kit, QIA: QIAamp DNA Mini Kit, IN: IndiSpin Pathogen Kit. g_: genus,

f_: family, o_: order, c_: class, p_: phylum.

Fig. S1. Comparison of DNA yields by DNA extraction protocols. DNA yield of

QIAampMini was significantly higher than that of PowerFecalPro (p < 0.001, Dunn's test with

Benjamini-Hochberg-adjustment). Also, the DNA yield of ZymoBIOMICS was significantly

higher than that of PowerFecalPro (p < 0.001, Dunn's test with Benjamini-Hochberg-adjustment).

The amount of DNA was calculated based on the absorbance of nucleic acids measured by

Nanodrop One. By the protocol recommended by the manufacturer, nucleic acid (Poly-A carrier)

was used in IndiSpin. Therefore, IndiSpin was excluded from the analysis of DNA yield. The

amount of DNA yield per 100 μL ThinPrep sample volume were compared. The bar graph shows

the mean and standard deviation. Zy: ZymoBIOMICS DNA Miniprep Kit, Pro: QIAamp

PowerFecal Pro DNA Kit, QIA: QIAamp DNA Mini Kit.

(A) Liquid-based cytology on cervix

Cervical microbiome

LBC

(B) Dispensing of each sample to 4 aliquot

(C) DNA extraction using four protocols

ZymoBIOMICS

PowerFecalPro

QIAampMini

IndiSpin

(D)

Positive control (vaginal mock)

Negative control (preservation solution without samples)

(E) 16S rRNA Illumina amplicon sequencing

Welcome

(F) Evaluation of taxonomic profiles

ZymoBIOMICS   PowerFecalPro

QIAampMini   IndiSpin

(A)

(B)