Title: A derivation error that affects carbon balance models exists in the current implementation of

the modified Arrhenius function

Running Title: Derivation error in modified Arrhenius model

Author: Bridget Murphy[1,2], Joseph R. Stinziano[3,*]

Affiliations: [1] Department of Biology, University of Western Ontario, London, ON, Canada; [2] Current

Affiliation: Department of Cell and Systems Biology, University of Toronto, Toronto, ON, Canada;

[3] Department of Biology, University of New Mexico, Albuquerque, NM, USA

[*] Corresponding author; Email: jstinziano@unm.edu; Tel: +1 (226) 678-1670

Article Type: Modelling/Theory

Word Count: 3,623

Introduction: 559

Description: 1640

Results: 688

Discussion: 710

Acknowledgments: 26

Number of Tables: 3

Number of Figures: 4

Supplementary Files: 1

Summary

- Understanding biological temperature responses is crucial to predicting global carbon fluxes.

The current approach to modelling temperature responses of photosynthetic capacity in large

scale modelling efforts uses a modified Arrhenius equation.

- We rederived the modified Arrhenius equation from the source publication from 1942 and

uncovered a missing term that was dropped by 2002. We compare fitted temperature

response parameters between the correct and incorrect derivation of the modified Arrhenius

equation.

- We find that most parameters are minimally affected, though activation energy is impacted

quite substantially. We then scaled the impact of these small errors to whole plant carbon

balance and found that the impact of the rederivation of the modified Arrhenius equation on

modelled daily carbon gain causes a meaningful deviation of ~18% day^{-1}.

- This suggests that the error in the derivation of the modified Arrhenius equation has impacted

the accuracy of predictions of carbon fluxes at larger scales since >40% of Earth System

Models contain the erroneous derivation. We recommend that the derivation error be

corrected in modelling efforts moving forward.

Keywords: Arrhenius, temperature, photosynthesis, gas exchange, modelling, carbon balance

Introduction

Globally, photosynthesis and autotrophic respiration are the largest biological carbon fluxes, with photosynthesis removing ~120 Gt C year^{-1} from the atmosphere and autotrophic respiration releasing ~60 Gt C year^{-1} back to the atmosphere (Amthor, 2000; Ciais *et al.*, 2013). Given the temperature sensitivity of these large carbon fluxes (Lombardozzi *et al.*, 2015), understanding how photosynthesis and respiration respond on acute, acclimatory, and adaptive timescales is crucial for predicting vegetative and carbon cycle responses to future global climates (Rogers *et al.*, 2017; Stinziano *et al.*, 2018). Biological temperature responses including photosynthesis and respiration are typically assumed to be exponential or peaked (Way & Yamori, 2014; Smith & Dukes, 2017; Kumarathunge *et al.*, 2019). Exponential responses are usually modelled based on an Arrhenius-type curve (Arrhenius, 1915):

$$f(T) = k \exp\left(\frac{-E_a}{RT}\right) \qquad \text{Equation 1,}$$

or equivalently,

$$f(T) = k_{25} \exp\left[\frac{E_a(T-298.15)}{298.15RT}\right] \qquad \text{Equation 2}$$

where f(T) is the rate of the process at temperature T, k is a pre-exponential factor, E_a is the activation energy in J mol^{-1}, R is the universal gas constant of 8.314 J mol^{-1} K^{-1}, T in K, k_{25} is the rate of the process at 298.15 K, and 298.15 is the reference temperature in degrees Kelvin (K). As for peaked responses, while a few options are available (Kruse *et al.,* 2008; Hobbs *et al.*, 2013; Heskel *et al.*, 2016), the most commonly implemented version is the modified Arrhenius model of Johnson *et al.* (1942) as presented in Medlyn *et al.* (2002):

$$f(T) = k_{25} \exp\left[\frac{E_a(T-298.15)}{298.15RT}\right]\left[\frac{1+\exp\left(\frac{298.15\Delta S-H_d}{298.15R}\right)}{1+\exp\left(\frac{T\Delta S-H_d}{RT}\right)}\right] \qquad \text{Equation 3}$$

where H_d is the deactivation energy in J mol^{-1}, and ΔS is the entropy of the process in J mol^{-1} K^{-1}.

Equation 3 (**which will be referred to as M2002 from this point forward as it is conventionally**

referred to in the literature as the source of the equation) is used for modelling the temperature

responses of photosynthetic capacity (i.e. maximum carboxylation capacity of rubisco, V_{cmax},

maximum electron transport capacity, J_{max}, and triose phosphate utilization capacity (TPU); see Table

1 for a review of which Earth System Models use M2002). These parameters are then used in

ecophysiological studies to understand thermal acclimation of photosynthesis (see Kattge & Knorr,

2007; Smith & Dukes, 2017; and Kumarathunge *et al.,* 2019 for examples). Furthermore, this equation

is also used in terrestrial biosphere models to predict future carbon cycling (e.g. Rogers *et al.*, 2017).

Due to the ubiquity of M2002 in modelling temperature responses and the thermal acclimation of

photosynthetic capacity (e.g. Kattge & Knorr, 2007; Rogers *et al.*, 2017; Smith & Dukes, 2017;

Kumarathunge *et al.*, 2019; Table 1), we revisited the original Johnson *et al.* (1942) modified

Arrhenius function to rederive M2002. In the process of this rederivation, we uncovered a term that

was completely dropped sometime between Johnson *et al.* (1942) and Medlyn *et al.* (2002) which

causes a systematic error in the application of the modified Arrhenius for modelling photosynthetic

capacity in individual species (e.g. Medlyn *et al.*, 2002) to modelling global scale carbon uptake (e.g.

Rogers *et al.*, 2017; Table 1). We then refit a freely available dataset (Kumarathunge *et al.*, 2019), with

both versions of the modified Arrhenius model, and fed the temperature response fits through a

carbon balance model to estimate the impact of the derivation error on modelled plant carbon

balance. We predicted that the derivation error would cause substantial variation in fitted

temperature response parameters, and that these differences would propagate through to modelled

daily carbon balance.

<u>Description</u>

Rederivation of the modified Arrhenius equation

Johnson *et al.* (1942, equation 24) describe the temperature response of the light intensity of a

luciferase reaction whereby the enzyme reversibly denatures or inactivates at high temperatures as:

$$96 \quad I = \frac{c'' T \exp\left(\frac{-\Delta H\ddagger}{RT}\right)}{1+\exp\left(\frac{\Delta S}{R}\right)\exp\left(\frac{-\Delta H}{RT}\right)} \qquad\qquad \text{Equation 4}$$

98 where I is the intensity of the luciferase reaction, c'' is not explicitly defined in Johnson *et al.* (1942),

99 but appears to represent a constant based on the derivation of Equation 4, T is the temperature in K,

100 R is the universal gas constant of 8.314 J mol^{-1} K^{-1}, $\Delta H\ddagger$ is the activation energy in J mol^{-1}, ΔH is the

101 deactivation energy in J mol^{-1}, and ΔS is the entropy in J mol^{-1} K^{-1}. First, we harmonize the notation

102 with standard ecophysiological representations such that $\Delta H\ddagger$ becomes E_a, and ΔH becomes H_d:

$$104 \quad f(T) = \frac{c'' T \exp\left(\frac{-E_a}{RT}\right)}{1+\exp\left(\frac{\Delta S}{R}\right)\exp\left(\frac{-H_d}{RT}\right)} \qquad\qquad \text{Equation 5}$$

106 We can relativize the equation to a reference temperature by dividing Equation 5 at a hypothetical

107 temperature by Equation 5 at a standard temperature (i.e. 25 °C):

$$109 \quad \frac{f(T)}{k_{25}} = \frac{\left[\dfrac{c'' T \exp\left(\frac{-E_a}{RT}\right)}{1+\exp\left(\frac{\Delta S}{R}\right)\exp\left(\frac{-H_d}{RT}\right)}\right]}{\left[\dfrac{c'' 298.15 \exp\left(\frac{-E_a}{298.15R}\right)}{1+\exp\left(\frac{\Delta S}{R}\right)\exp\left(\frac{-H_d}{298.15R}\right)}\right]} \qquad\qquad \text{Equation 6}$$

111 Next, rearrange fractions so that there is one numerator and one denominator on each line for clarity:

$$113 \quad \frac{f(T)}{k_{25}} = \left[\frac{c'' T \exp\left(\frac{-E_a}{RT}\right)}{c'' 298.15 \exp\left(\frac{-E_a}{298.15R}\right)}\right]\left[\frac{1+\exp\left(\frac{\Delta S}{R}\right)\exp\left(\frac{-H_d}{298.15R}\right)}{1+\exp\left(\frac{\Delta S}{R}\right)\exp\left(\frac{-H_d}{RT}\right)}\right] \qquad \text{Equation 7}$$

Cancel out the constant c'', use exponent rules to simplify so that E_a is contained within a single

exponential function, and the ΔS and H_d terms are grouped together:

$$\frac{f(T)}{k_{25}} = \frac{T}{298.15}\exp\left(\frac{-E_a}{RT}\ —\ \right)\left[\frac{1+\exp\left(\frac{\Delta S}{R}+\frac{-H_d}{298.15R}\right)}{1+\exp\left(\frac{\Delta S}{R}+\frac{-H_d}{RT}\right)}\right]$$ Equation 8

Simplify further by setting a common denominator within exponential terms:

$$f(T) = k_{25}\frac{T}{298.15}\exp\left(\frac{-298.15E_a+E_aT}{298.15RT}\right)\left[\frac{1+\exp\left(\frac{298.15\Delta S-H_d}{298.15R}\right)}{1+\exp\left(\frac{T\Delta S-H_d}{RT}\right)}\right]$$ Equation 9

Simplify exponential terms further to reduce the number of instances that E_a appears in the equation:

$$f(T) = k_{25}\frac{T}{298.15}\exp\left(\frac{E_a(T-298.15)}{298.15RT}\right)\left[\frac{1+\exp\left(\frac{298.15\Delta S-H_d}{298.15R}\right)}{1+\exp\left(\frac{T\Delta S-H_d}{RT}\right)}\right]$$ Equation 10

Note the difference between M2002 and the rederived equation 10 (**which will be referred to as**

J1942): the term T / 298.15 is missing from M2002. There are several alternative expressions of

Equation 4 (e.g. Farquhar *et al.*, 1980; Harley *et al.*, 1986; Harley *et al.*, 1992; Harley & Baldocchi,

1995; Lloyd, 1995), however when relativized to a common temperature, the equations presented in

Farquhar *et al.* (1980), Harley *et al.* (1992), Harley & Baldocchi (1995), and Lloyd (1995) are all

identical to M2002 when relativized by temperature, while the Harley *et al.* (1986) equation is

identical to our derivation (J1942) when relativized to a common temperature (see Appendix A for the

relativizations of Farquhar *et al.* (1980), Harley *et al.* (1986), and Harley *et al.* (1992); Harley &

Baldocchi (1995) uses an identical equation to Eq. 18 from Medlyn *et al.* (2002), while Lloyd (1995)

uses an identical equation to Eq. 17 from Medlyn *et al.* (2002)).

We traced the origin of the dropped term to two papers: Hall (1979) and Farquhar et al. (1980). Both

papers cite the M2002 equation as belonging to Sharpe and DeMichele (1977) when in fact it belongs

to Johnson et al. (1942). In Hall (1979), the reasoning behind the dropped term is given as: "In

addition their linear T term was omitted because it has little influence on the function and would have

prevented Eq. 22 [the M2002 equation] from approaching Eq. 21 [Equation 1 in the present study] in

the limiting conditions of low temperatures" (Hall, 1979, page 305). Considering J1942, at 5 °C the

error due to the dropped term would be ~278 K / 298 K = ~0.93, and at 45 °C would be 318 K / 298 K =

~1.07, so ~7% error at those temperatures. In context of measurement precision, such an assumption

of "little influence" may have been reasonable at the time. However, the reason regarding the

convergence of the two equations is simply tuning the model equations to fit their expectations.

Meanwhile, Farquhar et al. (1980) report that: "Eq. (36) is a simplified version of an equation

developed by Sharpe and DeMichelle [sic] (1977) to describe the effect of temperature on enzyme

inactivation" (Farquhar et al., 1980, page 84). In both cases, it appears that the intention was to

simplify the equation, which in the process of simplifying, generated an error that was propagated for

over 40 years.

The derivation error introduces multiple systematic errors. First, errors are introduced into the fitted

parameters E_a, H_d, and ΔS. Second, k_{25} is scaled using the wrong equation, introducing an error in f(T).

And third, acclimation equations describing E_a, H_d, and ΔS will then be in error due to errors in the

fitted parameters at each temperature. Here we focus on the impact of the dropped term on fitted

temperature response parameters and modelling whole-plant carbon balance.

Data analysis

Using A-C_i curve data compiled in Kumarathunge *et al.* (2019), available from the Kumarathunge *et al.*

(2018) repository, we used the {fitacis} function from the R package {plantecophys} (Duursma, 2015),

setting fitmethod = "bilinear", Tcorrect = FALSE, and fitTPU = TRUE, to obtain V_{cmax} and J_{max}. We then

fit Equations 3 and 10 to the data, allowing H_d to be fit. To ensure that the curves could be fit (i.e. that

there were enough data to fit 4 parameters), we only used data where A-C_i curves were measured at

5 or more temperatures. This reduced the number of candidate temperature responses from 729 to

403. We used the R package {minpack.lm} (Elzhov *et al.,* 2016), using Equation 2 to obtain starting

values for E_a and k_{25}, while the other initial parameters were ΔS = 0.650 kJ mol^{-1} K^{-1}, and H_d varying

from 1 to 1000 kJ mol^{-1}, followed by the {BIC} function to select the best model based on Bayesian

Information Criteria. We obtained 341 and 337 successful V_{cmax} temperature response curve fits for

M2002 and J1942, respectively, and 241 and 242 successful J_{max} temperature response curve fits for

M2002 and J1942, respectively. This resulted in a total of 547 fitted temperature response curves

which we filtered further, requiring that E_a, ΔS, and H_d were all positive values and that the V_{cmax} and

J_{max} data were paired, resulting in 196 complete pairs of V_{cmax} and J_{max} temperature responses for

analysis. The data covered a temperature range from 1 °C to 50 °C across all curves, with a median

range of 14 °C to 42.6 °C.

Modelling

We modelled the impact of the M2002 and J1942 equations on daily net plant carbon balance. Data

for leaf area, root and shoot masses, as well as leaf dark respiration at 25 °C were taken for white

spruce (*Picea glauca*) from Stinziano & Way (2017), while stomatal conductance model parameters

were calculated with the gas exchange data reported in Stinziano & Way (2017). Briefly, Stinziano and

Way (2017) grew white spruce in growth chambers under a simulated autumn treatment based on

temperatures and photoperiods from Trenton, ON, with weekly harvesting of biomass and gas

exchange measurements across 17 weeks as a control treatment, along with a warming treatment

where temperatures were +5 °C of the control, a constant summer day/night temperature with

declining photoperiod, and a constant summer photoperiod with ambient changes in temperature.

Here we only used data from Stinziano & Way (2017) that were from the control treatment. Mean

data were taken from the control treatment at weeks 0 and 12 to provide contrasting biomass

allocation patterns such that week 0 is a low respiration scenario and week 12 is a high respiration

scenario. These different respiration scenarios were used to reduce bias in any conclusions regarding

the impact of the M2002 and J1942 equations on carbon balance, as the ratio of photosynthesis to

respiration may alter the sensitivity of net carbon balance to the Arrhenius equation used. Root

respiration for white spruce was taken from Weger & Guy (1991) and we assumed that stem

respiration was equal to root respiration (Table 2).

For the full model structure and equations, please see the accompanying R package

{arrhenius.comparison} ("arrhenius.comparison_1.0.2.tar.gz"; Stinziano & Murphy, 2020) (see Table 3

for equations). Briefly, we linked the Medlyn *et al.* (2011) stomatal conductance model with the

Farquhar *et al.* (1980) C_3 photosynthesis model, assuming infinite mesophyll conductance to CO_2 as

these same assumptions were used in fitting the data from Kumarathunge *et al.* (2018).

Photosynthetic capacity, both maximum rubisco carboxylation capacity, V_{cmax}, and maximum electron

transport rate, J_{max}, were scaled to temperature using either M2002 or J1942, while respiration was

scaled according to (Atkin & Tjoelker, 2003). Photosynthesis and respiration were summed across

each modelled day to calculate daily plant carbon assimilation.

Modelling was performed on 18 total days of environmental data, with three days of data from three

months (17[th] – 19[th] of May, August, and October, 2019) obtained from external irradiance sensors at

the Biotron Experimental Climate Change Research Centre at the University of Western Ontario and

the remaining environmental data from Environment Canada historical climate data for South London

(43.01°N, 81.27°W, altitude: 251 m; temperature range: -0.1 – 27.9 °C) and the rooftop greenhouse at

the University of New Mexico (35.08°N, 106.62°W, altitude: 1587 m; temperature range: 7.2 – 42.2

°C) to capture different levels of environmental variability (see *Impacts on modelled net carbon*

balance for environmental data).

Overall, the modelling approach allows us to assess the relative differences between Equations 3 and

10 under a low- and high- respiration scenario across different ranges of seasonal variability.

Statistical analysis

Data were analyzed using the {lm} function in R v.3.6.2 (R Core Team, 2019), regressing the data

obtained from M2002 against the data obtained from J1942 for each of $V_{cmax,25}$, $J_{max,25}$, $E_{a,Vcmax}$, $E_{a,Jmax}$,

ΔS_{Vcmax}, ΔS_{Jmax}, $H_{d,Vcmax}$, $H_{d,Jmax}$, BIC values for the fits of Equations 3 and 10, daily photosynthesis, daily

net carbon balance, and daily photosynthesis : daily respiration ratios. Intercepts in the regression,

when significant, were interpreted as a bias in the parameter, while deviations in slope from a 1:1

relationship where interpreted as percentage over- or under- estimation of the parameter (i.e.

[parameter slope – 1] · 100 = % estimation error). All p-values were corrected for multiple testing

using the p.adjust() function in R with Holm's method. All code and data will be made freely available

on GitHub upon publication in the {arrhenius.comparison} R package (Stinziano & Murphy, 2020).

Results

Equations M2002 and J1942 exhibited similar performance

The performance between M2002 and J1942 for fitting $V_{cmax,25}$ temperature responses were essentially identical when assessed based on BIC, with a slope of 1.001 ± 0.001 ($F_{1,195}$ = 5.52 · 10⁵, R^2 = 0.9996, P < 2.2 · 10⁻¹⁶; Fig. **1a**), as was the case for $J_{max,25}$ with a slope of 1.001 ± 0.001 ($F_{1,195}$ = 2.43 · 10⁶, R^2 = 0.9999, P < 2.2 · 10⁻¹⁶; Fig. **1b**). However, while fitted temperature responses based on BIC appeared identical, the largest discrepancy between fitted responses was observed when comparing $J_{max,25}$ of both plant species scaled across 10 ˚C and 50 ˚C (Figs. **1c-f**; note the differences between the blue and red hatched lines).

Estimates of $V_{cmax,25}$ were essentially identical between the M2002 fitting (y-axis) and the J1942 fitting (x-axis) with a slope of 1.002 ± 0.004 ($F_{1,195}$ = 5.84 · 10⁴, R^2 = 0.9967, P < 2.2 · 10⁻¹⁶; Fig. **2e**), as was the case for $J_{max,25}$ with a slope of 1.001 ± 0.001 ($F_{1,195}$ = 6.56 · 10⁵, R^2 = 0.9997, P < 2.2 · 10⁻¹⁶; Fig. **2f**). $E_{a,Vcmax}$ was generally underestimated with a slope of 0.847 ± 0.024 and a positive bias of 9.73 ± 2.59 kJ mol⁻¹ when using M2002 ($F_{1,194}$ = 1202, R^2 = 0.8610, P < 2.2 · 10⁻¹⁶; Fig. **2a**), while $E_{a,Jmax}$ was underestimated with a slope of 0.832 ± 0.013 and a positive bias of 5.98 ± 1.06 kJ mol⁻¹ ($F_{1,194}$ = 3834, R^2 = 0.9518, P < 2.2 · 10⁻¹⁶; Fig. **2b**). ΔS_{Vcmax} was essentially identical between M2002 and J1942 with a slope of 1.000 ± 0.0233 ($F_{1,195}$ = 1849, R^2 = 0.9046, P < 2.2 · 10⁻¹⁶; Fig. **2c**), as was the case with ΔS_{Jmax} with a slope of 0.988 ± 0.033 ($F_{1,195}$ = 904, R^2 = 0.8226, P < 2.2 · 10⁻¹⁶; Fig. **2d**), and $H_{d,Vcmax}$ with a slope of 1.002 ± 0.005 ($F_{1,195}$ = 4.12 · 10⁴, R^2 = 0.9953, P < 2.2 · 10⁻¹⁶; Fig. **2g**). However, $H_{d,Jmax}$ was underestimated with a slope of 0.952 ± 0.024 and a positive bias of 31.48 ± 8.93 kJ mol⁻¹ ($F_{1,194}$ = 1592, R^2 = 0.8914, P < 2.2 · 10⁻¹⁶; Fig. **2h**).

Impacts on modelled net carbon balance

In general, the differences in thermal response parameters were amplified when integrated at the whole-plant level across a range of environmental conditions (Fig. **3**). For modelled daily photosynthesis (A), the slope for the low respiration model was 0.819 ± 0.013 and the intercept was 0.018 ± 0.002 g plant⁻¹ day⁻¹ (M2002 versus J1942; approximately 18% of modelled A) ($F_{1,3523}$ = 4.168 · 10³, R^2 = 0.5418; P < 2.2 · 10⁻¹⁶) (Fig. **4a**). For the high respiration model of daily A, the slope was 0.819 ± 0.013 and the intercept was 0.031 ± 0.003 g plant⁻¹ day⁻¹ (approximately 18% of modelled A) ($F_{1,3523}$ = 4.168 · 10³, R^2 = 0.5418; P < 2.2 · 10⁻¹⁶) (Fig. **4b**). The low respiration model of total daily carbon (C) gain had a slope of 0.836 ± 0.012 and an intercept of 0.012 ± 0.001 g plant⁻¹ day⁻¹ (approximately 16%

of modelled C gain) (M2002 versus J1942; $F_{1,3523}$ = 4.539 · 10^3, R^2 = 0.5629; P <2.2 · 10^{-16}) (Fig. **4c**). The

high respiration of total C gain similarly had a slope of 0.859 ± 0.012 and an intercept of 0.011 ± 0.002

g plant^{-1} day^{-1} (approximately 14% of modelled C gain) (M2002 versus J1942; $F_{1,3523}$ = 5.364 · 10^3, R^2 =

0.6035; P <2.2 · 10^{-16}) (Fig. **4d**). The ratio of the total daily photosynthesis: respiration (A/R) was also

considered when comparing models. The low respiration model of A/R had a slope of 0.982 ± 0.007

and the intercept was 0.146 ± 0.031 (approximately 2% of modelled A/R) (M2002 versus J1942; $F_{1,3523}$

= 2.096 · 10^4, R^2 = 0.8561; P <2.2 · 10^{-16}) (Fig. **4e**). The high respiration model had a similar slope of

0.982 ± 0.007 and the intercept was 0.082 ± 0.017 (approximately 2% of modelled A/R) (M2002

versus J1942; $F_{1,3523}$ = 2.097 · 10^4, R^2 = 0.8562; P <2.2 · 10^{-16}) (Fig. **4f**).

Discussion

We sought to determine whether the missing term in M2002 had a meaningful impact on fitted

temperature response parameters due to its prevalence in photosynthetic temperature response data

and vegetation modelling (Kattge & Knorr, 2007; Duursma & Medlyn, 2012; Rogers *et al.,* 2017; Smith

& Dukes, 2017; Stinziano *et al.,* 2018; Stinziano *et al.,* 2019; Kumarathunge *et al.,* 2019). Our present

analysis suggests that there is a large impact on E$_a$ for both V$_{cmax}$ and J$_{max}$, however there were

minimal impacts on k$_{25}$, H$_d$, and ΔS (though note that H$_{d,Jmax}$ was underestimated) (Fig. **2**). In general,

fitting M2002 instead of J1942 results in E$_a$ reductions of around 15% with positive bias, even though

fitted responses are *visually* similar (Fig. **1**). These findings are promising in that one of the

parameters to which modelled carbon gain is particularly sensitive, H$_d$ (Stinziano *et al.,* 2018), is

minimally affected by the missing term. However, since temperature responses are non-linear, small

changes in the shape of an accelerating curve can have a strong impact on the integral of the

response (Jensen, 1906; e.g. changes in the shape of the temperature response of carbon assimilation

can have strong impacts on total carbon fixation). Despite fit performance being nearly identical and

visually similar (Fig. **1**), the differences due to E$_a$ values between the equations led to an 18%

reduction in modelled daily photosynthesis for M2002 compared to J1942, leading to a ~14-16%

reduction in modelled net daily carbon gain. Overall, comparisons of low-respiration to high-

respiration scenarios resulted in similar percent changes in daily C balance with similar variation.

Not only did we compare the impact of the missing term in M2002 on different respiration scenarios,

but we also modelled data across a range of environmental conditions to understand the impact on C

balances of plants under different temperature regimes and how these impacts could scale up in

global models. A ~15% reduction in net daily carbon gain is substantial by itself, but this difference

would also be amplified over time for a single plant. Plant growth follows the compound interest law

(Blackman, 1919). As leaf area increases with growth so too does the rate of photosynthesis and thus

the amount of carbon available to further increase growth and metabolism, hence why relative

growth rates are commonly utilized in the literature (e.g. Shipley, 1989; Causton, 1991; Tjoelker *et al.*,

1999; Loveys *et al.,* 2003; Poorter *et al.,* 2012; Pommereng & Anders, 2015). A reduction of 18% in

modelled daily photosynthesis caused by using the modified equation with the missing term (M2002)

would become compounded long-term with plant growth and may result in underestimations of

future carbon uptake. It is thus likely that the differences we observed would accumulate to even

larger carbon flux errors across large spatial and temporal scales with fluctuating temperatures.

Based on the above analysis, the impact of the missing term in the modified Arrhenius equation

substantially alters E_a and net daily carbon balance at a whole-plant level. While our models of plant

carbon uptake were indicative of problems caused by using M2002, it would be worth comparing how

the use of this derivation has affected the interpretation of temperature response data and whole

plant carbon balances collected from previous studies. Future work comparing the impact of using

M2002 opposed to J1942 on interpreting whole tree carbon balance and net primary production data

could lead to better estimations of the temperature sensitivity of carbon uptake. Given that carbon

balance is the time integral of net CO_2 assimilation, this may also lead to substantial impacts over a

long time period. On a larger scale, a correction of this derivation error will also be necessary when

considering Earth system models, as 17 out of 40 models in the CMIP6 simulations are affected by the

derivation error (Table 1). We therefore recommend the switch from M2002 to J1942 because: 1)

M2002 is the result of a derivation error circa 1979, and 2) M2002 and J1942 lead to different net

daily carbon estimates due to the derivation error, which may currently be compensated by other

factors in large-scale models, most notably in the Community Land Model which currently relies on

the M2002 equation (Bonan, 1996; Lawrence et al., 2018) and underlies 30% of Earth System Models

(Table 1).

<u>Acknowledgments</u>

We would like to thank Wesley J. Noe at the University of New Mexico for providing climate data. This research was supported by personal funds.

<u>Author Contributions</u>

Both authors contributed to all aspects of the study.

<u>Additional Information</u>

A version of this manuscript was posted on bioRxiv (manuscript ID: BIORXIV/2020/921973).

332 **Tables**

333 **Table 1. List of land surface models affected by the derivation error in M2002 and the affected Earth Systems Models (ESMs) being used**

334 **in the CMIP6 simulations. Note that 17 of the 40 (42.5%) CMIP6 ESMs are affected by the derivation error. Affected models are in bold.**

Land Surface Model Family		# of ESMs	Contains M2002?	Affected ESMs	References
Atmosphere-Vegetation Interaction Model	AVIM	2	No	BCC-CSM2-MR, BCC-ESM1	Ji, 1995
Community Atmosphere-Biosphere Land Exchange Model	**CABLE**	**2**	**Yes**	**ACCESS-CM2, ACCESS-ESM1-5**	**Haverd et al., 2018**
Canadian Land Surface Scheme	**CLASS**	**1**	**Yes**	**CanESM5**	**Swart et al., 2019**
Community Land Model	**CLM**	**12**	**Yes**	**CAMS-CSM1-0, CESM2, CESM2-FV2, CESM2-WACCM, CESM2-WACCM-FV2, NorESM2-LM, NorESM2-MM, E3SM-1-0, FGOALS-f3-L, FGOALS-g3, TaiESM1, SAM0-UNICON**	**Bonan, 1996; Lawrence et al., 2018**
Goddard Institute for Space Studies Global Circulation Model	GISS-GCM	3	No	GISS-E2-1-G, GISS-E2-1-H, GISS-E2-2-G	Friend & Kiang, 2005
Interaction Soil Biosphere Atmosphere Carbon Cycle	ISBAcc	3	No	CNRM-CM6-1, CNRM-CM6-1-HR, CNRM-ESM2-1	Calvet et al., 1998; Joetzjer et al., 2015
Joint Scheme for Biosphere Atmosphere Coupling in Hamburg	JSBACH	4	No	AWI-CM-1-1-MR, MPI-ESM1-2-HR, MPI-ESM1-2-LR, NESM3	Rogers et al., 2017; Cao et al., 2018
Joint UK Land Environment Simulator	JULES	3	No	HadGEM3-GC31-LL, HadGEM-GC31-MM, UKESM	Clark et al., 2011
Land Model 4	LM4	2	No	GFDL-CM4, GFDL-ESM4	Milly et al., 2014
Lund-Potsdam-Jena Model	LPJ	3	No	EC-Earth3, EC-Earth3-Veg, IPSL-CM6A-LR	Sitch et al., 2003; Arora et al., 2020
Simple Biosphere Model	SiB	3	No*	INM-CM4-8, INM-CM5-0, MRI-ESM2-0	Sato et al., 1989; Volodin & Lykosov, 1998; Hosaka, 2011
Vegetation Integrative SImulator for Trace Gases	**VISIT**	**2**	**Yes**	**MIROC6, MIROC-ESL2**	**Ito, 2010; Arora et al., 2020**

335 *The SiB model does not include photosynthesis.

336   **Table 2. Parameters used in modelling daily carbon gain.**

Parameter	Group	Value	Reference
Respiration	R_{dark}	2.78 µmol m^{-2} s^{-1}	Stinziano & Way, 2017
	R_{day}	0.7 * R_{dark}	Ayub *et al.*, 2011
	R_{root}	0.0095 µmol g^{-1} s^{-1}	Weger & Guy, 1991
	R_{stem}	0.0095 µmol g^{-1} s^{-1}	Assumed
	Q_{10}	2.015	Atkin & Tjoelker, 2003
Γ*	25 °C	42.75 µmol mol^{-1}	Bernacchi *et al.*, 2001
	E_a	37.83 kJ mol^{-1}	Bernacchi *et al.*, 2001
K_m	25 °C	718.4 µmol mol^{-1}	Bernacchi *et al.,* 2001
	E_a	65.51 kJ mol^{-1}	Bernacchi *et al.,* 2001
α		0.8	Norman & Campbell, 1998
φ		0.08	Norman & Campbell, 1998
Leaf Area	Low R	0.015 m^2	Stinziano & Way, 2017
	High R	0.025 m^2	Stinziano & Way, 2017
Stem Mass	Low R	0.496 g	Stinziano & Way, 2017
	High R	2.523 g	Stinziano & Way, 2017
Root Mass	Low R	0.498 g	Stinziano & Way, 2017
	High R	5.072 g	Stinziano & Way, 2017

Q_{10}: thermal sensitivity coefficient; Γ*: photorespiratory CO_2 compensation point; K_m: apparent

Michaelis-Menten constant for rubisco carboxylation in 21% O_2/air (i.e. $K_{c,air}$); α: absorbance of

photosynthetically activation radiation; φ: maximum quantum efficiency of photosynthetic electron

transport; E_a: activation energy; R_{dark}: leaf respiration in the dark; R_{day}: leaf respiration in the light;

R_{root}: root respiration; R_{stem}: stem respiration; Low R: low respiration scenario; High R: high

respiration scenario.

Table 3. Equations used in modelling daily carbon uptake.

Equation	Reference
$$f(T) = k_{25}Q_{10}^{\frac{T-298.15}{10}}$$	Atkin & Tjoelker, 2003
$$f(T) = k_{25}e^{\left[\frac{E_a(T-298.15)}{298.15RT}\right]}$$	Arrhenius, 1915
$$f(T) = k_{25}e^{\left[\frac{E_a(T-298.15)}{298.15RT}\right]}\left[\frac{1+e^{\frac{298.15\Delta S-H_d}{298.15R}}}{1+e^{\frac{T\Delta S-H_d}{RT}}}\right]$$	Medlyn et al., 2002
$$f(T) = k_{25}\frac{T}{298.15}e^{\left[\frac{E_a(T-298.15)}{298.15RT}\right]}\left[\frac{1+e^{\frac{298.15\Delta S-H_d}{298.15R}}}{1+e^{\frac{T\Delta S-H_d}{RT}}}\right]$$	Johnson et al., 1942; this study
$$W_c = V_{cmax}\frac{C_i - \Gamma*}{C_i + K_m}$$	Farquhar et al., 1980
$$W_j = J\left(\frac{C_i - \Gamma*}{C_i + 2\Gamma*}\right)$$	Farquhar et al., 1980; Way et al., 2011

Where J is calculated as the minimum root of: $\theta J^2 - (0.5\alpha\phi Q_{in} + J_{max})J + 0.5\alpha\phi Q_{in}J_{max} = 0$

$$A_{gross} = min(W_c, W_j)$$

$$A_{net} = A_{gross} - R_{day}$$

$$g_s = g_0 + 1.6\left(1 + \frac{g_1}{\sqrt{VPD}}\right)\left(\frac{A_{net}}{C_a}\right)$$	Medlyn et al., 2011

$$C_i = C_a - 1.6\frac{A_{net}}{g_s}$$

$$A_{plant} = A_{net} \times LA - R_{dark} \times LA - R_{stem} \times SM - R_{root} \times RM$$

$$C_{balance} = \sum A_{plant} \times 3600 \times \frac{12.01}{1,000,000}$$

f(T): rate of a process at a given temperature; T: temperature in K; k_{25}: rate of a process at 25 °C;

Q_{10}: thermal sensitivity coefficient; E_a: activation energy in kJ mol^{-1}; ΔS: entropy parameter in kJ

mol^{-1}; H_d: deactivation energy in kJ mol^{-1}; R: universal gas constant in 0.008314 kJ mol^{-1} K^{-1}; W_c: rate

of CO_2-limited carboxylation in µmol m^{-2} s^{-1}; W_j: rate of RuBP regeneration-limited carboxylation in

μmol m^{-2} s^{-1}; V_{cmax}: maximum rate of rubisco carboxylation capacity in μmol m^{-2} s^{-1}; J_{max}: maximum

rate of electron transport in μmol m^{-2} s^{-1}; C_i: intercellular CO_2 concentration in μmol mol^{-1}; Γ*:

photorespiratory CO_2 compensation point in μmol mol^{-1}; K_m: Michaelis-Menten constant for rubisco

in μmol mol^{-1}; α: absorbance of photosynthetically active radiation; ɸ: maximum quantum

efficiency of electron transport; Q_{in}: incident photosynthetically active radiation; A_{gross}: gross CO_2

assimilation in μmol m^{-2} s^{-1}; A_{net}: net CO_2 assimilation in μmol m^{-2} s^{-1}; R_{day}: leaf day respiration in

μmol m^{-2} s^{-1}; g_s: stomatal conductance to water in mol m^{-2} s^{-1}; g_o: intercept for the Medlyn et al.

2011 model; g_1: slope for the Medlyn et al. 2011 model; VPD: vapor pressure deficit in kPa; C_a: CO_2

concentration at the leaf surface in μmol mol^{-1}; A_{plant}: whole plant net CO_2 assimilation; LA: leaf

area in m^{-2}; R_{dark}: leaf dark respiration in μmol m^{-2} s^{-1}; R_{stem}: stem respiration in μmol m^{-2} s^{-1}; R_{root};

root respiration in μmol m^{-2} s^{-1}; SM: stem mass in g; RM: root mass in g; $C_{balance}$: whole plant daily

carbon balance in g plant^{-1} day^{-1}.

Figure Captions

Figure 1. Relative performance and fit for Equations 3 (M2002) and 10 (J1942) for temperature

responses of $V_{cmax\ 25}$ and $J_{max\ 25}$ based on a,b) Bayesian Information Criterion (BIC), and c-f) visual

inspection of curve fits across 10-50 ˚C for c,d) *Elymus canadensis* and 20-50 ˚C of e,f) *Betula*

***alleghaniensis*. Red dashed lines indicate curves fitted with the M2002 equation and blue dotted**

lines indicate curves fitted with the J1942 equation. Parameters for c-f) are as follows: c) M2002:

k_{25}: 46.4 μmol m^{-2} s^{-1}, E_a: 29.0 kJ mol^{-1}, ΔS: 0.692 kJ mol^{-1} K^{-1}, H_d: 221.4 kJ mol^{-1}; J1942: k_{25}: 46.4 μmol

m^{-2} s^{-1}, E_a: 29.0 kJ mol^{-1}, ΔS: 0.655 kJ mol^{-1} K^{-1}, H_d: 209.0 kJ mol^{-1}; d) M2002: k_{25}: 68.9 μmol m^{-2} s^{-1}, E_a:

0.0 kJ mol^{-1}, ΔS: 0.746 kJ mol^{-1} K^{-1}, H_d: 240.5 kJ mol^{-1}; J1942: k_{25}: 69.5 μmol m^{-2} s^{-1}, E_a: 0.0 kJ mol^{-1},

ΔS: 0.625 kJ mol^{-1} K^{-1}, H_d: 201.0 kJ mol^{-1}; e) M2002: k_{25}: 37.6 μmol m^{-2} s^{-1}, E_a: 57.7 kJ mol^{-1}, ΔS: 0.579

kJ mol^{-1} K^{-1}, H_d: 185.4 kJ mol^{-1}; J1942: k_{25}: 37.0 μmol m^{-2} s^{-1}, E_a: 57.7 kJ mol^{-1}, ΔS: 0.557 kJ mol^{-1} K^{-1},

H_d: 178.1 kJ mol^{-1}; f) M2002: k_{25}: 66.7 μmol m^{-2} s^{-1}, E_a: 23.0 kJ mol^{-1}, ΔS: 0.761 kJ mol^{-1} K^{-1}, H_d: 246.6

kJ mol^{-1}; J1942: k_{25}: 65.5 μmol m^{-2} s^{-1}, E_a: 23.0 kJ mol^{-1}, ΔS: 0.668 kJ mol^{-1} K^{-1}, H_d: 216.1 kJ mol^{-1}.

Figure 2. The modified Arrhenius equation missing the term (M2002) compared to J1942 fits

different values for E_a (a,b), but mostly similar values for ΔS (c,d), k_{25} (e,f) and H_d (f, g) for both V_{cmax}

(a, c, e, g) and J_{max} (c, d, f, h). E_a: activation energy, ΔS: entropy parameter, k_{25}: rate of the process

at 25 °C, H_d: deactivation energy, V_{cmax}: maximum capacity of rubisco carboxylation, J_{max}: maximum

rate of electron transport. Black line indicates 1:1 line and grey dashed line indicates respective

modelled slopes and intercepts.

Figure 3. Hourly environmental data used to drive the model in Table 3 covering 3 days (17th, 18th,

and 19th) of 3 months. (a,c,e) Albuquerque, NM, USA (ABQ); (b,d,e) London, ON, Canada (LDN).

Environmental parameters include: a,b) temperature; c,d) vapour pressure deficit; and e,f)

irradiance (Q_{in}).

Figure 4. The modified Arrhenius equation without the missing term (M2002) gives differences in

modelled daily carbon fluxes compared to J1942 for low R (a,c,e) and high R (b,d,f) under scenarios

where H_d is allowed to vary. Modelled carbon fluxes include: a,b) Total daily photosynthesis; c,d)

total daily C gain; and e,f) total daily A/R. A: photosynthesis, R respiration, Low R: low respiration,

High R: high respiration. Black line indicates 1:1 line and grey dashed line indicates respective

modelled slopes and intercepts.

Figures

Figure 1. Relative performance and fit for Equations 3 (M2002) and 10 (J1942) for temperature

responses of $V_{cmax\ 25}$ and $J_{max\ 25}$ based on a,b) Bayesian Information Criterion (BIC), and c-f) visual

inspection of curve fits across 10-50 ˚C for c,d) *Elymus canadensis* and 20-50 ˚C of e,f) *Betula*

***alleghaniensis*. Red dashed lines indicate curves fitted with the M2002 equation and blue dotted**

lines indicate curves fitted with the J1942 equation. Parameters for c-f) are as follows: c) M2002:

k_{25}: 46.4 µmol m^{-2} s^{-1}, E_a: 29.0 kJ mol^{-1}, ΔS: 0.692 kJ mol^{-1} K^{-1}, H_d: 221.4 kJ mol^{-1}; J1942: k_{25}: 46.4 µmol

m^{-2} s^{-1}, E_a: 29.0 kJ mol^{-1}, ΔS: 0.655 kJ mol^{-1} K^{-1}, H_d: 209.0 kJ mol^{-1}; d) M2002: k_{25}: 68.9 µmol m^{-2} s^{-1}, E_a:

0.0 kJ mol^{-1}, ΔS: 0.746 kJ mol^{-1} K^{-1}, H_d: 240.5 kJ mol^{-1}; J1942: k_{25}: 69.5 µmol m^{-2} s^{-1}, E_a: 0.0 kJ mol^{-1},

ΔS: 0.625 kJ mol^{-1} K^{-1}, H_d: 201.0 kJ mol^{-1}; e) M2002: k_{25}: 37.6 µmol m^{-2} s^{-1}, E_a: 57.7 kJ mol^{-1}, ΔS: 0.579

kJ mol^{-1} K^{-1}, H_d: 185.4 kJ mol^{-1}; J1942: k_{25}: 37.0 µmol m^{-2} s^{-1}, E_a: 57.7 kJ mol^{-1}, ΔS: 0.557 kJ mol^{-1} K^{-1},

H_d: 178.1 kJ mol^{-1}; f) M2002: k_{25}: 66.7 µmol m^{-2} s^{-1}, E_a: 23.0 kJ mol^{-1}, ΔS: 0.761 kJ mol^{-1} K^{-1}, H_d: 246.6

kJ mol^{-1}; J1942: k_{25}: 65.5 µmol m^{-2} s^{-1}, E_a: 23.0 kJ mol^{-1}, ΔS: 0.668 kJ mol^{-1} K^{-1}, H_d: 216.1 kJ mol^{-1}.

Figure 2. The modified Arrhenius equation missing the term (M2002) compared to J1942 fits

different values for E_a (a,b), but mostly similar values for ΔS (c,d), k_{25} (e,f) and H_d (f, g) for both V_{cmax}

(a, c, e, g) and J_{max} (c, d, f, h). E_a: activation energy, ΔS: entropy parameter, k_{25}: rate of the process

at 25 °C, H_d: deactivation energy, V_{cmax}: maximum capacity of rubisco carboxylation, J_{max}: maximum

rate of electron transport. Black line indicates 1:1 line and grey dashed line indicates respective

modelled slopes and intercepts.

Figure 3. Hourly environmental data used to drive the model in Table 3 covering 3 days (17th, 18th,

and 19th) of 3 months. (a,c,e) Albuquerque, NM, USA (ABQ); (b,d,e) London, ON, Canada (LDN).

Environmental parameters include: a,b) temperature; c,d) vapour pressure deficit; and e,f)

irradiance (Qin).

Figure 4. The modified Arrhenius equation without the missing term (M2002) gives differences in modelled daily carbon fluxes compared to J1942 for low R (a,c,e) and high R (b,d,f) under scenarios where H_d is allowed to vary. Modelled carbon fluxes include: a,b) Total daily photosynthesis; c,d) total daily C gain; and e,f) total daily A/R. A: photosynthesis, R respiration, Low R: low respiration, High R: high respiration. Black line indicates 1:1 line and grey dashed line indicates respective modelled slopes and intercepts.

References

Amthor JS. 2000. Direct effect of elevated CO_2 on nocturnal in situ leaf respiration in nine temperate deciduous tree species is small. *Tree Physiology* **20**: 139–144.

Arora VK, Katavouta A, Williams RG, Jones CD, Brovkin V, Friedlingstein P, Schwinger J, et al. 2020. Carbon-concentration and carbon-climate feedbacks in CMIP6 models, and their comparison to CMIP5 models. *Biogeosciences* doi: 10.5194/bg-2019-473.

Arrhenius S. 1915. *Quantitative laws in biological chemistry*. Bell: London.

Atkin OK, Tjoelker MG. 2003. Thermal acclimation and the dynamic response of plant respiration to temperature. *Trends in Plant Science* **8**:343-351.

Ayub G, Smith RA, Tissue DT, Atkin OK. 2011. Impacts of drought on leaf respiration in darkness and light in *Eucalyptus saligna* exposed to industrial-age atmospheric CO_2 and growth temperature. *New Phytologist* **190**: 1003-1018.

Bernacchi CJ, Singsaas EL, Pimentel C, Portis Jr AR, Long SP. 2001. Improved temperature response functions for models of rubisco-limited photosynthesis. *Plant Cell and Environment* **24**:253-259.

Blackman, VH. 1919. The compound interest and plant growth. *Annals of Botany* **33:** 353-360.

Bonan GB. 1996. A Land Surface Model (LSM Version 1.0) for Ecological, Hydrological, and Atmospheric Studies: Technical Description and User's Guide (No. NCAR/TN-417+STR). *University Corporation for Atmospheric Research.* doi:10.5065/D6DF6P5X

Calvet J-C, Noilhan J, Roujean J-L, Bessemoulin P, Cabelguenne M, Olioso A, Wigneron J-P. 1998. An interactive vegetation SVAT model tested against data from six contrasting sites. *Agricultural and Forest Meteorology* **92**:73-95

Cao J, Wang B, Yang Y-M, Ma L, Li J, Sun B, Bao Y, He J, Zhou X, Wu L. 2018. The NUIST Earth System Model (NESM) version 3: description and preliminary evaluation. *Geoscientific Model Development* **11**:2975-2993.

Causton, DR. 1991. Plant-growth analysis: the variability of relative growth rate within a sample. *Annals of Botany* **67**(2): 137-144.

Ciais P, Sabine C, Bala G, Bopp L, Brovkin V, Canadell J, Chhabra A, DeFries R, Galloway J, Heimann M. 2013. Carbon and Other Biogeochemical Cycles. In: Heinze C, Tans P, Vesala T, eds. *Climate Change 2013: The Physical Science Basis.* Cambridge, UK, and New York, NY, USA: Cambridge University Press.

Clark DB, Mercado LM, Sitch S, Jones CD, Gedney N, Best MJ, Pryor M, et al. 2011. The Joint UK Land

Environment Simulator (JULES), model description – part 2: carbon fluxes and vegetation

dynamics. *Geoscientific Model Development* **4**:701-722.

Drake JE, Tjoelker MG, Aspinwall MJ, Reich PB, Pfautsch S, Barton CVM. 2019. The partitioning of

gross primary production for young *Eucalyptus tereticornis* trees under experimental warming

and altered water availability. *New Phytologist* **222:** 1298-1312

Duursma RA. 2015. Plantecophys – an R package for analysing and modelling leaf gas exchange data.

PLoS ONE **10**:e0143346.

Duursma RA, Medlyn BE. 2012. MAESPA: a model to study interaction between water limitation,

environmental drivers and vegetation function at tree and stand levels, with an example

application to [CO_2] x drought interactions*. Geoscientific Model Development* **5**:919-940.

Elzhov TV, Mullen KM, Spiess A-N, Bolker B. 2016. minpack.lm: R Interface to the Levenberg-

Marquardt nonlinear least-squares algorithm found in MINPACK, plus support for bounds. R

package version 1.2-1. https://CRAN.R-project.org/package=minpack.lm

Farquhar GD, von Caemmerer S, Berry JA. 1980. A biochemical model of photosynthetic CO_2

assimilation in leaves of C_3 species. *Planta* **149**: 78-90.

Friend AD, Kiang NY. 2005. Land surface model development for the GISS GCM: effects of improved

canopy physiology on simulated climate. *Journal of Climate* **18**:2883-2902.

Hall AE. 1979. A model of leaf photosynthesis and respiration for predicting carbon dioxide

assimilation in different environments. *Oecologia* **43**: 299-316.

Harley PC, Baldocchi DD. 1995. Scaling carbon dioxide and water vapour exchange from leaf to

canopy in a deciduous forest. I. Leaf model parameterization. *Plant, Cell and Environment*

18:1146-1156.

Harley PC, Thomas RB, Reynolds JF, Strain BR. 1992. Modelling photosynthesis of cotton grown in

elevated CO_2. *Plant, Cell and Environment* **15**:271-282.

Harley PC, Tenhunen JD, Lange OL. 1986. Use of an analytical model to study limitations on net

photosynthesis in *Arbutus unedo* under field conditions. *Oecologia* **70**:393-401.

Haverd V, Smith B, Nieradzik L, Briggs PR, Woodgate W, Trudinger CM, Canadell JG, Cuntz M. 2018.

A new version of the CABLE land surface model (Subversion revision r4601) incorporating land

use and land cover change, woody vegetation demography, and a novel optimization-based

approach to plant coordination of photosynthesis. *Geoscientific Model Development* **11**:2995-
3026.

Heskel MA, O'Sullivan OS, Reich PB, Tjoelker MG, Weerasinghe LK, Penillard A, Egerton JJG, et al.
2016. Convergence in the temperature response of leaf respiration across biomes and plant
functional types. *Proceedings of the National Academy of Sciences USA* **113**: 3832-3837.

Hobbs JK, Jiao W, Ester AD, Parker EJ, Schipper LA, Arcus VL. 2013. Change in heat capacity for
enzyme catalysis determines temperature dependence of enzyme catalyzed rates. *ACS Chemical*
Biology **8**:2388-2392.

Hosaka M. 2011. A new MRI land surface model HAL. *AGU Fall Meeting Abstracts* GC43B-0901.

Ito A. 2020. Changing ecophysiological processes and carbon budget in East Asian ecosystems under
near-future changes in climate: implications for long-term monitoring from a process-based
model. *Journal of Plant Research* **123**:577-588.

Jensen JLWV. 1906. Sur les fonctions convexes et les inégalités entre les valeurs moyennes. *Acta*
Mathematica **30**: 175-193.

Ji J. 1995. A climate-vegetation interaction model: simulating physical and biological processes at the
surface. *Journal of Biogeography* **22**:445-451.

Joetzjer E, Delire C, Douville H, Ciais P, Decharme B, Carrer D, Verbeeck H, De Weirdt M, Bonal D.
2015. Improving the ISBA$_{CC}$ land surface model simulation of water and carbon fluxes and stocks
over the Amazon forest. *Geoscientific Model Development* 8:1709-1727.

Johnson FH, Eyring H, Williams RW. 1942. The nature of enzyme inhibitions in bacterial
luminescence: sulfanilamide, urethane, temperature and pressure. *Journal of Cellular and*
Comparative Physiology **20**:247-268.

Kattge J, Knorr W. 2007. Temperature acclimation in a biochemical model of photosynthesis: a
reanalysis of data from 36 species. *Plant, Cell and Environment* **30**:1176-1190.

Kruse J, Hopmans P, Adams MA. 2008. Temperature responses are a window to the physiology of
dark respiration: differences between CO_2 release and O_2 reduction shed light on energy
conservation. *Plant, Cell and Environment* **31**: 901-914.

Kumarathunge DP, Medlyn BE, Drake JE, Tjoelker MG, Aspinwall MJ, Battaglia M, Cano FJ, Carter
KR, Molly AC, Lucas AC, et al. 2019. Acclimation and adaptation components of the

temperature dependence of plant photosynthesis at the global scale. *New Phytologist* **222**:768-784.

Kumarathunge DP, Medlyn BE, Drake JE, Tjoelker MG, Aspinwall MJ, Battaglia M, Cano FJ, Carter KR, Molly AC, Lucas AC, et al. 2018. ACi-TGlob_V1.0: a global dataset of photosynthetic CO_2 response curves of terrestrial plants. doi: 10.6084/m9.figshare.7283567.v1.

Lawrence D, Fisher R, Koven C, Oleson K, Swenson S, Vertenstein M, et al. 2018. Technical Description of version 5.0 of the Community Land Model (CLM). National Center for Atmospheric Research. http://www.cesm.ucar.edu/models/cesm2/land/

Lloyd J, Grace J, Miranda AC, Meir P, Wong SC, Miranda HS, Wright IR, Gash JHC, McIntyre J. 1995. A simple calibrated model of Amazon rainforest productivity based on leaf biochemical properties. *Plant, Cell and Environment* **18**:1129-1145.

Lombardozzi DL, Bonan GB, Smith NG, Dukes JS, Fisher RA. 2015. Temperature acclimation of photosynthesis and respiration: a key uncertainty in the carbon cycle-climate feedback. *Geophysical Research Letters* **42**:8624-8631.

Loveys BR, Atkinson LJ, Sherlock DJ, Roberts RL, Fitter AH, Atkin OK. 2003. Thermal acclimation of leaf and root respiration: An investigation comparing inherently fast- and slow-growing plant species. *Global Change Biology* **9**(6): 895-910.

Medlyn BE, Duursma RA, Eamus D, Ellsworth DS, Prentice IC, Barton CVM, Crous KY, et al. 2011. Reconciling the optimal and empirical approaches to modelling stomatal conductance. *Global Change Biology* **17**:2134-2144.

Medlyn BE, Dreyer E, Ellsworth D, Forstreuter M, Harley PC, Kirschbaum MUF, Le Roux X, et al. 2002. Temperature responses of parameters of a biochemically based model of photosynthesis. II. A review of experimental data. *Plant, Cell and Environment* **25**:1167-1179.

Milly PCD, Malyshev SL, Shevliakova E, Dunne KA, Findell KL, Gleeson T, Liang Z, et al. 2014. An enhanced model of land water and energy for global hydrologic and Earth-system studies. *Journal of Hydrometeorology* **15**:1739-1761.

Norman JM, Campbell GS. 1998. *An introduction to environmental biophysics*. New York NY: Springer.

Pommerening A, Muszta, A. 2015. Methods of modelling relative growth rate. *Forest Ecosystems* **2**:5.

Poorter H, Niklas KJ, Reich PB, Oleksyn J, Poot P, Mommer L. 2012. Biomass allocation to leaves, stem, and roots: meta-analyses of interspecific variation and environmental control. *New Phytologist* **193**: 30-50.

R Core Team. 2019. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Rogers A, Medlyn BE, Dukes JS, Bonan G, von Caemmerer S, Dietze MC, Kattge J, et al. 2017. A roadmap for improving the representation of photosynthesis in Earth system models. *New Phytologist* **213**:22-42.

Sato N, Sellers PJ, Randall DA, Schneider EK, Shukla J, Kinter JL, Hou Y-T, Albertazzi E. 1989. Implementing the Simple Biosphere Model (SiB) in a general circulation model: methodologies and results. *NASA Technical Report* NASA-CR-185509.

Sharpe PJH, DeMichele DW. 1977. Reaction kinetics of poikilotherm development. *Journal of Theoretical Biology* **64**: 649-670.

Shipley, B. 1989. The use of above-ground maximum relative growth-rate as an accurate predictor of whole-plant maximum relative growth-rate. *Functional Ecology* **3**(6): 771-775.

Sitch S, Smith B, Prentice IC, Arneth A, Bondeau A, Cramer W, Kaplan JO, et al. 2003. Evaluation of ecosystem dynamics, plant geography and terrestrial carbon cycling in the LPJ dynamic global vegetation model. *Global Change Biology* **9**:161-185.

Smith NG, Dukes JS. 2017. Short-term acclimation to warmer temperatures accelerates leaf carbon exchange processes across plant types. *Global Change Biology* **23**:4840-4853.

Stinziano JR, Murphy BK. 2020. arrhenius.comparison: comparing versions of the modified Arrhenius equation. R package version 1.0.2. https://github.com/jstinzi/arrhenius.comparison.

Stinziano JR, Way DA. 2017. Autumn photosynthetic decline and growth cessation in seedlings of white spruce are decoupled under warming and photoperiod manipulations. *Plant, Cell and Environment* **40:**1296-1316.

Stinziano JR, Way DA, Bauerle WL. 2018. Improving models of photosynthetic thermal acclimation: which parameters are most important and how many should be modified? *Global Change Biology* **24**:1580-1598.

Stinziano JR, Bauerle WL, Way DA. 2019. Modelled net carbon gain responses to climate change in
boreal trees: impacts of photosynthetic parameter selection and acclimation. *Global Change*
Biology **25**:1445-1465.

Swart NC, Cole JNS, Kharin VV, Lazare M, Scinocca JF, Gillett NP, Anstey J, et al. 2019. The Canadian
Earth System Model version 5 (CanESM5.0.3). *Geoscientific Model Development* **12:**4823-4873.

Tjoelker M, Oleksyn J, Reich P. 1999. Acclimation of respiration to temperature and CO_2 in seedlings
of boreal tree species in relation to plant size and relative growth rate. *Global Change Biology*
49(6): 679-691.

Volodin EM, Lykosov VN. 1998. Parameterization of heat and moisture transfer in the soil-vegetation
system for use in atmospheric general circulation models: 1. Formulation and simulations based
on local observational data. *Izvestiya Atmospheric and Oceanic Physics* **34:**405-416.

Way DA, Yamori W. 2014. Thermal acclimation of photosynthesis: on the importance of adjusting our
definitions and accounting for thermal acclimation of respiration. *Photosynthesis Research*
119:89-100.

Weger HG, Guy RD. 1991. Cytochrome and alternative pathway respiration in white spruce (*Picea*
glauca) roots. Effects of growth and measurement temperature. *Physiologia Plantarum* **83**:675-
681.

 Supplementary information

Appendix A – Alternate derivations of the modified Arrhenius equation

Below are the temperature relativizations of Farquhar et al. (1980), Harley et al. (1986), and Harley et

al. (1992) showing that Farquhar et al. (1980) and Harley et al. (1992) can be transformed to be

identical to Equation 3 in the manuscript, while Harley et al. (1986) can be transformed to be identical

to Equation 10 in the manuscript.

Farquhar et al. (1980)

From Eq. 36 in Farquhar et al. (1980):

$$k = \frac{c \times \exp\left[-E/(RT_k)\right]}{1+\exp\left[(ST_k-H)/(RT_k)\right]} \hspace{4cm} \text{Equation A1}$$

Relativizing to 25 °C:

$$k/k_{25} = \left[\frac{c \times \exp\left[-E/(RT_k)\right]}{1+\exp\left[(ST_k-H)/(298.15R)\right]}\right]\left[\frac{1+\exp\left[(ST_k-H)/(298.15R)\right]}{c \times \exp\left[-E/(T_kR)\right]}\right] \hspace{1cm} \text{Equation A2}$$

Simplifying:

$$k/k_{25} = \exp\left[E/(298.15R)-E/(RT_k)\right]\left[\frac{1+\exp\left[(S298.15-H)/(298.15R)\right]}{1+\exp\left[(ST_k-H)/(RT_k)\right]}\right] \hspace{1cm} \text{Equation A3}$$

$$k = k_{25}\exp\left[E\frac{(T_k-298.15)}{(298.15RT_k)}\right]\left[\frac{1+\exp\left[(298.15S-H)/(298.15R)\right]}{1+\exp\left[(ST_k-H)/(RT_k)\right]}\right] \hspace{1cm} \text{Equation A4}$$

where E is activation energy (E_a), S is deactivation entropy (ΔS), and H is deactivation energy (H_d).

Harley et al. (1986)

From Eq. 7 in Harley et al. (1986):

$$k = \frac{T_k\exp\left[c-\Delta H_a/(RT_k)\right]}{1+\exp\left[(\Delta ST_k-\Delta H_d)/(RT_k)\right]} \hspace{4cm} \text{Equation A5}$$

Relativizing to 25 °C:

$$k/k_{25} = \left[\frac{T_k\exp\left[c-\Delta H_a/(RT_k)\right]}{1+\exp\left[(\Delta ST_k-\Delta H_d)/(RT_k)\right]}\right]\left[\frac{1+\exp\left[(\Delta S298.15-\Delta H_d)/(R298.15)\right]}{298.15\exp\left[c-\Delta H_a/(R298.15)\right]}\right] \hspace{0.5cm} \text{Equation A6}$$

Simplifying:

$k/k_{25} = \dfrac{T_k}{298.15}\exp\left[\Delta H_a/(R298.15) - \Delta H_a/(RT_k)\right]\left[\dfrac{1+\exp\left[(\Delta S298.15-\Delta H_d)/(R298.15)\right]}{1+\exp\left[(\Delta S T_k-\Delta H_d)/(RT_k)\right]}\right]$ Equation A7

$k = k_{25}\dfrac{T_k}{298.15}\exp\left[\Delta H_a\dfrac{(T_k-298.15)}{(298.15 RT_k)}\right]\left[\dfrac{1+\exp\left[(298.15\Delta S-\Delta H_d)/(298.15 R)\right]}{1+\exp\left[(\Delta S T_k-\Delta H_d)/(RT_k)\right]}\right]$ Equation A8

where ΔH_a is activation energy (E_a), and ΔH_d is deactivation energy (H_d).

Harley et al. (1992)

From Eq. 9 in Harley et al. (1992):

$k = \dfrac{\exp\left[c-\Delta H_a/(RT_k)\right]}{1+\exp\left[(\Delta S T_k-\Delta H_d)/(RT_k)\right]}$ Equation A9

Relativizing to 25 °C:

$k/k_{25} = \left[\dfrac{\exp\left[c-\Delta H_a/(RT_k)\right]}{1+\exp\left[(\Delta S T_k-\Delta H_d)/(RT_k)\right]}\right]\left[\dfrac{1+\exp\left[(\Delta S298.15-\Delta H_d)/(R298.15)\right]}{\exp\left[c-\Delta H_a/(R298.15)\right]}\right]$ Equation A10

Simplifying:

$k/k_{25} = \exp\left[\Delta H_a/(R298.15) - \Delta H_a/(RT_k)\right]\left[\dfrac{1+\exp\left[(\Delta S298.15-\Delta H_d)/(R298.15)\right]}{1+\exp\left[(\Delta S T_k-\Delta H_d)/(RT_k)\right]}\right]$ Equation A11

$k = k_{25}\exp\left[\Delta H_a\dfrac{(T_k-298.15)}{(298.15 RT_k)}\right]\left[\dfrac{1+\exp\left[(298.15\Delta S-\Delta H_d)/(298.15 R)\right]}{1+\exp\left[(\Delta S T_k-\Delta H_d)/(RT_k)\right]}\right]$ Equation A12

where ΔH_a is activation energy (E_a), and ΔH_d is deactivation energy (H_d).