Between-subject prediction reveals a shared representational geometry in the rodent
hippocampus

Hung-Tu Chen, Jeremy R. Manning, Matthijs A. A. van der Meer*

Department of Psychological & Brain Sciences, Dartmouth College, Hanover, NH 03755

*Correspondence: mvdm@dartmouth.edu

Summary

The rodent hippocampus constructs statistically independent representations across
environments ("global remapping") and assigns individual neuron firing fields to locations within
an environment in an apparently random fashion, processes thought to contribute to the role of
the hippocampus in episodic memory. This random mapping implies that it should be
challenging to predict hippocampal encoding of a given experience in one subject based on the
encoding of that same experience in another subject. Contrary to this prediction, we find that by
constructing a common representational space across rats ("hyperalignment"), we can predict
data of "right" trials (R) on a T-maze in a target rat based on 1) the "left" trials (L) of the target
rat, and 2) the relationship between L and R trials from a different source rat. These
cross-subject predictions relied on ensemble activity patterns including both firing rate and field
location, and outperformed a number of control mappings, such as those based on permuted
data that broke the relationship between L and R activity for individual neurons, and those
based solely on within-subject prediction. This work constitutes proof-of-principle for successful
cross-subject prediction of ensemble activity patterns in the hippocampus, and provides new
insights in understanding how different experiences are structured, enabling further work
identifying what aspects of experience encoding are shared vs. unique to an individual.

Keywords: place cells, remapping, generalization, pattern completion, pattern separation,
hyperalignment

Word Count: 4411 words

Number of Figures: 4 in main text, 10 in Supplementary Information

A fundamental challenge faced by any memory system is how related experiences should be
organized – storing the details of each individual experience preserves potentially valuable
details, but is storage-inefficient and hampers generalization, whereas treating all experiences
as the same risks ignoring potentially important differences[1]. For instance, learning the common
spatial features of different floors in the same building makes it possible to predict the layout of
a not-yet-visited floor ("similar to the others"); at the same time, each floor also has unique
features, such as the location of a specific colleague's office, that do not generalize. Thus,
memory systems need to balance pattern-completion (treating a new observation the same as a
previous one) and pattern-separation (keeping similar observations as distinct).

The rodent hippocampus is a model system for studying the neural basis of these processes.
Strikingly, the hippocampus can construct statistically independent representations across
environments ("global remapping")[2–5] and assigns individual neuron firing fields to locations
within an environment in an apparently random fashion[6,7]. Similarly, "engram" studies suggest
that the population of neurons allocated to a given experience is determined by a competition
based on randomly fluctuating excitability levels among eligible neurons[8]. Although there are
also examples of hippocampal cells whose firing properties are tied to a particular stimulus
feature (e.g. reward[9]) and therefore transfer across different environments, the received wisdom
is that those cells that do change their firing fields between environments or across different
regions of the same environment, do so randomly[10].

Remapping studies to date have been limited to within-subject comparisons, but it is possible in
principle that what appears random within a single subject in fact obeys a common rule that is
shared across subjects. Consider how two related experiences such as running the left (L) and
right arms (R) of a T-maze may be encoded in a population of hippocampal neurons (Figure
S1). The correlation between L and R activity on a cell-by-cell basis may be zero, but still obey
an underlying structure. For instance, cells that tend to fire at the start of L may be more likely to
fire at the end of R (Figure S1a), or more realistically, how place fields shift between L and R
depends on both their location and firing rate (Figure S1b). If such a rule were to exist, it should
be possible to predict, across subjects, what R activity of a target subject looks like, based on
(1) that subject's L activity and (2) the relationship between L and R activity found in a different
"source" subject. Although there is no way to predict how two different subjects encode a given
experience L (especially when sampling randomly from different numbers of neurons that are
not uniquely identifiable across subjects as in e.g. *C. elegans*), the *relationship* between how
two different experiences L and R are represented may be conserved across subjects.

Such a *representational geometry* has been demonstrated in a number of brain regions in
human cognitive neuroscience studies that use fMRI[11–14], but cross-subject prediction has not
yet been applied to ensemble recording data in the rodent hippocampus. If (re)mapping in the
rodent hippocampus were to show a shared representational geometry, this would not only
challenge a long-held dogma about the randomness of place cell allocation, but potentially also

open up novel lines of research that can elucidate the algorithmic basis of memory assignment
and generalization in a wide variety of settings, while creating a bridge between rodent neural
data and human fMRI work.

Results

The overall goal of this study is to determine if we can predict how hippocampal place cells in a
"target" subject encode a particular experience based on two ingredients: (a) knowledge of how
the target subject encodes a distinct but related experience, and (b) how a different, "source"
subject encodes the same two experiences.

We operationalize this idea using data from T-maze tasks, in which rats run along the left and
right arms of the maze to form the two related experiences under study. Specifically, we can
describe hippocampal activity on this task as two subject-specific matrices with time as the
horizontal dimension, and neuron as the vertical dimension (Figure 1a, leftmost column); one
matrix describing the average activity for left trials (L), and another matrix for right trials (R; see
Figure S2a for a description of how this input data is obtained). We aim to predict the R matrix in
the target subject, based on (a) the target's L matrix and (b) the source's L and R matrices. Note
that the left and right arms differ in a number of respects other than spatial location, so we use
these terms here as descriptive labels, rather than as an interpretation about the nature of their
neural encoding (see *Discussion*). We first apply principal component analysis (PCA) to each
subject's data, so that neural activity during L and R trials can be visualized as trajectories in a
subject-specific reduced space (Step 1 in Figure 1a).

Next, the key step for cross-subject prediction employs a procedure from human cognitive
neuroscience, "hyperalignment"[11,15], that projects each subject's idiosyncratic neural activity
into a common space that minimizes the Euclidean distance between neural activity trajectories
(Step 2 in Figure 1a; see also Figure S1a for a schematic). Working in this common space, we
can identify the relationship between L and R activity in the source subject, and express this
relationship as a transformation matrix ("hypertransform") which can be applied to the target
subject's L trials to obtain a predicted R trajectory $\widehat{R}$. This predicted trajectory is then projected
back to the target-specific neural space to obtain a prediction which is compared to the actual
data (final step in Figure 1a). Figure 1b shows in more detail an example prediction alongside
the actual target activity for a different session; note the overall correspondence between
predicted and actual activity (see Figure S3 for more examples).

Figure 1: **Workflow and example for cross-subject prediction using the hypertransform**.
(A) Our objective is to predict place cell activity on the right arm (R) of a T-maze in a "target"
subject, based on (i) place cell activity in the left arm (L) in the target subject, and (ii) L and R
place cell activity in a different, "source" subject. These input data are shown in the top left panel:
both the source and target subjects have two matrices each that describe, for each recorded
neuron, how its activity varies during left and right trials. Neurons are ordered according to their
peak firing time on the R arm. Note that although the number of time bins is the same across
subjects, the number of recorded cells may be different. Therefore, the first step of the analysis
workflow is to apply principal component analysis (PCA), resulting in neural activity trajectories
for left and right trials (red and blue, respectively) in each subject's own PCA space. Three
principal components are shown here for display purposes, but in the main analysis 10 PCs
were used. Next, these neural activity trajectories are mapped into a common space using a
"hyperalignment" procedure that minimizes the Euclidean distance between the trajectories
across subjects (step 2, see *Methods* for details). In this common space, a Procrustean
transformation[16] (HT in step 3) is derived that maps L to R trajectories for the "source" subject
(step 3), which can then be applied to the L trajectory of the "target" subject (step 4) to obtain its
predicted R trajectory in the common space (step 5). This predicted R trajectory is then projected
back to the "target" PCA space using the inverse of the matrix used in step 2 (step 6) and
expanded back into the target's original neuron space (step 7). Finally, the predicted R neural
activity is compared to the actual R activity to yield an error measure (step 8). The diagonal
pattern apparent in the predicted data indicates similarity between actual and predicted R,
although differences are also visible. **(B)** Close-up of example target L and R activity matrices
("Actual", neurons ordered by temporal fields on R) with example predicted R activity derived
from a different source session. Each matrix column describes ensemble neural activity for a
single time bin, and maps to a corresponding point on a neural activity trajectory in common
space (top panels). The hypertransform (HT) is a mapping from L to R activity that operates on
these (aligned) ensemble activity vectors. As a result, the R prediction for any given cell is not
only based on that cell's L activity alone, but also depends on the activity of the other cells at that
time; see Figure S3 for further examples and a more detailed explanation.

If a given subject encodes L and R trials independently, then it should not be possible to use
one subject's neural activity for L and R to predict anything about how another subject encodes
R trials based on its L trials. On the other hand, if there is some shared structure between
subjects in how L and R trials are encoded, then cross-subject prediction should perform better
than chance. To test this idea, we compare the prediction of a target subject's R trials to shuffled
controls. In the first, "row shuffle", we obtain a distribution of chance predictions for each
source-target pair, based on breaking the relationship between L and R trials in the source
subject by randomly permuting the rows of the R matrix (see Figure S2b for a schematic).
Based on this chance distribution, we define three metrics: (1) a z-score of the actually
observed error compared to chance, (2) the difference between the actually observed prediction
error per cell and the mean of the chance prediction error, and (3) the proportion of chance
prediction errors that were lower than the observed error.

We used two different data sets: the first, "Carey" data set[17,18] is from a T-maze where L and R
arms were deliberately furnished with distinct surface colors and textures. In contrast, the
second, "Gupta" data set[19,20] used a T-maze whose arms had similar surfaces. Starting with
the Carey data, we found that the hypertransform (HT) prediction of R trials in the target subject
was better than chance overall for all metrics used (Figure 2, top and middle rows; green "HT"
bars; z-score: $p < 0.001$ for Wilcoxon signed rank test vs. 0; raw error: $p < 0.001$). Cross-subject
prediction of R activity was better than chance even when the R data was withheld entirely from
the hyperalignment step (see *Methods* for details on the withholding procedure; Figure S4a, top
row; $p < 0.001$ for HT vs. 0), when L and R activity was expressed as tuning curves in space
rather than in time (Figure S4c, middle row; $p < 0.001$ for HT vs. 0), and when putative
interneurons were removed (Figure S5a). We found similar results for the Gupta data, with the
HT prediction consistently better than chance (Figure 2, bottom row; $p < 0.001$ for HT vs. 0).
These results demonstrate that the relationship between L and R trials is not random across
subjects.

To test if the hyperalignment step of the prediction procedure is important, rather than some
other part of the workflow, we repeated the analysis with the hyperalignment step left out (i.e.
we applied the L-R transform obtained from the source subject's PCA space to the target
subject's PCA space, "PCA-only"; see Methods for details). The HT-based prediction
outperformed the PCA-only prediction both for the Carey data (blue "PCA" bars in Figure 2a
inset, middle row; $p < 0.001$, binomial test) and the Gupta data (bottom row; $p < 0.001$, binomial
test), although for the Gupta data the difference was noticeably smaller (we investigate this
further below). Again, the HT predictions outperformed PCA-only predictions when spatial
tuning curves were used (Figure S4c) and with putative interneurons removed (Figure S5a); to
facilitate visualization of the results in the analyses that follow, we used the
interneuron-removed data. Although the HT prediction still outperformed PCA-only when
target-R data was withheld entirely, the HT prediction was much better (i.e. closer to half-split
prediction) if at least one trial of target R was included for alignment (Figure S4a and b; insets).
Crucially however, R data cannot be predicted better than chance even when it is included in

Figure 2: Cross-subject prediction of R trials of a "target" subject based on how a
"source" subject encodes L and R trials outperforms prediction based on shuffled source
data. For each source-target pair in the Carey data, we computed a z-score of the actual
observed error between predicted and actual R trials (based on the hypertransform procedure,
"HT") compared to a shuffled distribution in which the R rows of the source subject were
randomly permuted. Thus, lower z-scores indicate lower error and therefore better prediction
than chance. Across all source-target pairs, this z-scored error varied depending on the pair
used (column **A**, top row), but was lower than chance overall, as indicated by a shift in the
z-score histogram relative to 0 ("HT" green bars in column **A**, middle row; median: -3.10 +/- 0.54,
SEM across unique source-target pairs, p < 0.001 for Wilcoxon signed rank test vs. 0).
Cross-subject predictions based on the L-R transform in common space ("hypertransform")
outperformed predictions based on the L-R transform in PCA space ("PCA" blue bars in column

A, middle row, inset; see *Methods* for details; HT < PCA: 99.62% of source-target pairs, p < 0.001 for binomial test) in terms of raw error between predicted and actual R trials (median of HT raw error: 443.73 +/- 62.44, PCA: 1409.14 +/- 204.22, split-half: 130.99 +/- 46.72, red triangle in inset; mean squared error per cell, see *Methods* for details). Next, we applied the same analysis to a different data set ("Gupta", bottom row, in which the L and R maze arms were more similar to each other than in the "Carey" data), and found that the hypertransform prediction was again significantly better than chance ("HT"; median: -2.59 +/- 1.44, p < 0.001, bottom row) and better than PCA-only (HT < PCA: 79.58% of source-target pairs, p < 0.001; column **A**, bottom row, inset; median of HT raw error: 115.26 +/- 142.05, PCA: 121.42 +/- 208.23, split-half: 37.39 +/- 10.01). Columns **B** and **C** use the same layout as column **A**, but using different metrics to describe prediction accuracy. **B** uses the raw error (between predicted and actual R neural activity; lower error/negative indicates better prediction) compared to the mean of the shuffle distribution, and **C** uses the proportion of the shuffle distribution with smaller error than the actually observed error (lower proportions indicate better prediction). For the raw error measure, the HT prediction was better than chance in both the Carey data (column **B** middle row; median: -652.87 +/- 486.05, p < 0.001 for HT vs. 0; red line indicates split-half prediction) and the Gupta data (**B** bottom row; median: -594.62 +/- 735.37, p < 0.001 for HT vs. 0).

A possible trivial explanation for the better-than-chance cross-subject prediction is that rats represent experiences in L and R similarly, so that a simple duplicate of L activity forms a reasonable prediction of R activity. To test if L-R correlations at the cell-by-cell level (i.e. row-wise correlations, Figure S2c) underlie the cross-subject prediction results in Figure 2, we compared cross-subject predictions based on the hypertransform (L-R mapping in common space) with those based on the *identity transform*: a within-subject prediction that simply takes a duplicate of the L trajectory in common space and uses it as the prediction for R. For the Carey data set, the cross-subject HT prediction was significantly better than that based on the identity transform (ID; left panel in Figure 3a; p < 0.001, binomial test), demonstrating that the better-than-chance prediction of R is not due to cell-by-cell correlations with L activity. In contrast, for the Gupta data, the HT prediction did not outperform the identity prediction (right panel in Figure 3a), suggesting that the better-than-chance cross-subject prediction for this data set (Figure 2, bottom row) can be attributed to row-wise correlations between L and R activity.

Figure 3: Cross-subject prediction outperforms within-subject prediction only in the absence of cell-by-cell correlations. A: Comparison of cross-subject prediction error ("hypertransform", green bars; HT) with within-subject prediction error ("identity transform"; blue bars, ID) for two different data sets. In the "Carey" data (left panel) the left and right arms of the maze had different texture and color patterns; in the "Gupta" data (right panel) the two maze arms were identical. For the Carey data, cross-subject prediction was significantly better than within-subject prediction (HT < ID: 100% of unique source-target pairs, p < 0.001 for binomial test) whereas for the Gupta data, this difference was not significant. **B**: Cell-by-cell correlations of firing rates between L and R arms (i.e. row-wise correlations of the L and R matrices in Figure S2c), were significantly higher in the Gupta data compared to the Carey data (Gupta: median r = 0.21 +/- 0.27, SEM across subjects, Carey: median r = -0.02 +/- 0.12, Wilcoxon rank sum test, p < 0.001). Cell-by-cell correlations in the Carey data were not significantly different from 0 (p = 0.44 for Wilcoxon signed rank test vs. 0). **C**: Population vector (PV) correlations between ensemble activity at each time point and every other time point, i.e. column-wise correlations of the L and R activity matrices, averaged across sessions. Both Carey and Gupta data sets show high correlations around the diagonal, indicating an overall autocorrelation in time; however, the Gupta data additionally shows high off-diagonal correlations between L and R which are barely visible in the Carey data. **D**: Quantification of the median PV correlation between L and R (i.e. the values along the diagonal of the lower left quadrant in **C**). For Gupta data, this correlation is remarkably high (median r = 0.95 +/- 0.16) whereas for Carey data, it is significantly lower (median r = 0.09 +/- 0.10, p < 0.001 for Wilcoxon rank sum test) but significantly different from 0 (p < 0.001 for Wilcoxon signed rank test vs. 0), consistent with previous reports of global remapping[2,4]. The above results explain why the HT is not needed for Gupta data to achieve better than chance predictions (as shown in Figure 2, bottom row): L and R activity is sufficiently similar such that the L trajectory alone in either PCA or common space can predict R activity.

To test this idea, we investigated the correlation structure between the L and R firing rate
matrices using two different measures: the cell-by-cell (row-wise) correlation, averaged across
all cells, and the column-wise population vector (PV) correlation (averaged across sessions, see
Figure S2 for schematic; note that in order to compute these measures, putative interneurons
were removed from the data; see Figure S5a and Methods). The cell-by-cell firing rate
correlations between L and R arms were significantly more correlated in the Gupta data
compared to Carey data (Gupta: median r = 0.21 +/- 0.27, Carey: median r = -0.02 +/- 0.12,
SEM across subjects; $p < 0.001$, Wilcoxon rank sum test; Figure 3b). Similarly, PV correlations
in the Gupta data showed high off-diagonal values between L and R, which were barely visible
in the Carey data (Figure 3c-d). The low correlation values observed in the Carey data are
consistent with those previously reported and characterized as global remapping[2,4], whereas
the Gupta correlations are strikingly high, indicating the presence of "symmetric" cells with
similar firing patterns on the L and R arms. High L-R correlations in the Gupta data imply that
the cross-subject (hypertransform) method cannot outperform the already very good prediction
based on within-subject correlations, whereas for the nearly uncorrelated Carey data, there is
room for cross-subject prediction to improve.

Importantly, the comparison between the two data sets suggests that the cross-subject
prediction on Carey data is not the result of within-subject correlations -- because, if it were,
then the HT prediction would be similar to the ID prediction. So, if cross-subject prediction for
the Carey data is not simply a consequence of within-subject correlations between L and R,
what *is* the prediction based on? In other words, can we identify what features of the L-R
relationship are generalizable across subjects without appearing as within-subject correlations
between L and R activity? To address this question, we generated synthetic neural activity
matrices using 1-D Gaussians with three parameters: time, peak firing rate (FR) and width.
Specifically, three simulated data sets captured different potential place cell properties: (1) each
neuron has an independent probability of having a firing field on L and R, and all parameters are
randomly and independently chosen for L and R (**ind-ind**, top row in Figure 4a), (2) if a neuron
has a firing field on L, it does not have a field on R (and vice versa), and the parameters of the
field are chosen randomly (**x-or**, second row) and (3) each neuron has an independent
probability of having a field on L and R, but if a cell has a field in both, all three parameters are
the same (**ind-same-all**, third row). For all these scenarios, we generated synthetic data
matching the number of recording sessions and the number of neurons recorded in the Carey
data, and applied exactly the same analysis procedure.

The independent (**ind-ind**) simulation serves as a crucial sanity check to verify that our
cross-subject prediction procedure cannot exploit shared structure where none exists; as
expected, cross-subject prediction was not different from chance in this scenario (Figure 4a,
right column; $p = 0.92$ for Wilcoxon signed rank test vs. 0). In contrast, both **x-or** and
ind-same-all showed better-than-chance cross-subject prediction (both $p < 0.001$). If the **x-or** or
ind-same-all rules are potential explanations for better-than-chance predictions in Carey data,
we should see zero cell-by-cell correlations and low PV correlations between L and R in these

two simulations as observed in the Carey data (Figure 3b-d). However, the **x-or** scenario shows
negatively correlated PV correlations inconsistent with the Carey data (Figure 4b-c; r = -0.17).
The **ind-same-all** rule shows both high cell-by-cell correlations (Figure 4d; r = 0.25) and high
PV correlations (r = 0.41), which is again inconsistent with the Carey data, but more in line with
the Gupta data (compare Figure 3b-d). In further simulations, we separately investigated the
role of each parameter of the 1-D gaussian place fields (time, FR and width) as a potential
explanation for cross-subject prediction; only when the same time is shared across L and R are
the cross-subject predictions better than chance (**ind-same-time**, top row in Figure S6; p <
0.001). However, as in the main analysis (Figure 3b) the cell-by-cell correlations are inconsistent
with those observed in the Carey data. Thus, although both rules like x-or and same-time can
support cross-subject prediction, the cell-by-cell correlations in the Carey data are different and
therefore these simple rules cannot be the full explanation.

Figure 4: Simulations demonstrate that representational geometry, but not simple rules
such as exclusive-or and firing rate correlations, result in cross-subject prediction
while being consistent with the data. A: Example L and R activity matrices (left column)
and histogram of z-scores of cross-subject prediction compared to the distribution of shuffle
predictions (right column; z-scores lower than zero indicate better-than-chance predictions) of
four simulated data sets: (1) neurons have a fixed, independent probability (0.5) of having a
1-D Gaussian place field on L and/or R, with the three parameters of time, peak firing rate
(FR) and width randomly and independently chosen for L and R (**ind-ind**, top row), (2)
neurons only have a field on *either* L or R but not both, and parameters of the field are chosen
randomly as in (1) (**x-or**, second row), (3) neurons have a fixed independent probability of
having L and R fields as in (1) but with the additional constraint that neurons with both L and
R fields must have the same three parameters (**ind-same-all**, third row) and (4) the activity on
L is simulated by assigning each neuron an independent probability of having a field whose
parameters are randomly chosen, then the activity on R is obtained by applying a L-R

transform (hypertransform, HT) from the Carey data to the simulated L activity (**sim. HT**, last row). As expected, in the **ind-ind** (independent) case, cross-subject prediction is not possible (z-score vs. chance, median: -0.01 +/- 0.03, SEM across unique source-target pairs, p = 0.70 for Wilcoxon signed rank test vs. 0). In contrast, **x-or**, **ind-same-all** and **sim. HT** all show better-than-chance cross-subject predictions (median: -1.26 +/- 0.30 for **x-or**, median: -2.62 +/- 0.06 for **ind-same-all**, median: -5.23 +/- 0.04 for **sim. HT**, all p < 0.001 for Wilcoxon signed rank test vs. 0), indicating that if there is a non-random L-R relationship in the underlying data, the hypertransform procedure can exploit it. Note that although this synthetic example shows substantially negative predicted firing rates, negative firing rates, when they did occur, tended to be much smaller for predictions using actual data. **B**: Population vector (PV; column-wise) correlations between ensemble activity at each time point and every other time point of the L and R activity matrices. Only **ind-same-all** shows high off-diagonal correlations between L and R, resembling the Gupta data set in which L and R arms were identical and firing activity on both arms is highly correlated (compare with Figure 3c). In **x-or**, off-diagonal correlations are slightly negative. **C**: Quantification of the median PV correlation between L and R (i.e. the values along the diagonal of the lower left quadrant in **B**). PV correlations between L and R in **ind-ind** are zero, but L and R are positively correlated in **ind-same-all** (median r = 0.40 +/- 0.01) since for every time point in L where there is a field, the same ensemble activity appears at the same time point in R with probability 0.5. **X-or** shows a negative correlation (median r = -0.18 +/- 0.02) because for every time point in L where there is a field, the same ensemble activity would deterministically be absent in R, and vice versa. None of these simple rules are consistent with the PV correlation found in the Carey data; however, **sim. HT** does show similar correlations as the data (median r = 0.04 +/- 0.03 for **sim. HT** and median r = 0.09 +/- 0.10 for Carey). **D**: Cell-by-cell (row-wise) correlations of L and R show that **ind-same-all** is more correlated than the data, whereas **sim. HT** yields similar correlations (median r = 0.04 +/- 0.02 for **sim. HT** and median r = -0.02 +/- 0.12 for Carey). Thus, taken across panels C and D, simple rules (**x-or** and **ind-same-all**) are inconsistent with the data, but the representational geometry embodied in the **sim. HT** yields correlations that are similar to the data.

If these predetermined rules cannot be the source of cross-subject prediction in the Carey data, what can? We first sought to rule out a number of possible low-level explanations, verifying that differences between source and target sessions in running speed (Figure S7a) and average firing rates (Figure S7b) do not explain cross-subject prediction performance. Plotting cross-subject prediction error as a function of time did not highlight any particular areas as being especially well or poorly predicted (Figure S8c) ruling out explanations based on shared cues such as the choice point or the existence of a reward site at the end of each arm. One more subtle possibility is a *single-cell mapping rule* such as a cell with a field at the start of L being mapped to the end of R; however, no obvious field mapping pattern between L and R was observed for both actual and predicted data (Figure S9), consistent with the low population vector correlations in the data (Figure 3c-d).

Is there another possibility? The hyperalignment procedure we used to derive the L-R transform ("hypertransform") was originally developed to capture so-called *representational geometry*: a shared rule that specifies how differences in the *ensemble* neural response to a set of stimuli may be preserved across subjects even though each subject may encode a given stimulus quite

differently[11]. To test if such a geometry is consistent with the data, we created another synthetic
data set in which the activity on L is simulated by assigning each neuron an independent
probability of having a 1-D gaussian place field whose parameters are randomly chosen, and
the activity on R is obtained by applying the L-R transform (hypertransform, HT) from the Carey
to the simulated L activity (**sim. HT**, bottom row in Figure 4a, see *Methods*). Not only did **sim.
HT** show significant better-than-chance cross-subject predictions on the simulated data (p <
0.001) but the cell-by-cell (r = 0.04) and PV correlations (r = 0.04) were similar to the Carey
data. Thus, unlike the simple x-or and same-parameter simulated scenarios, and unlike
single-cell mapping rules, a shared representational geometry is consistent with the correlations
observed in the data.

The hypertransform capitalizes on a geometric relationship between ensemble activity patterns,
which can be difficult to relate back to the properties of any individual cell. However, some
insight can be obtained by separating (1) *mean firing rate* relationships between L and R
regardless of their specific firing locations (e.g. "cells with high rates on L tend to have high
rates in R", as suggested by the data in Lee et al. 2019[21]), and (2) *specific firing locations*
regardless of mean firing rate differences (e.g. "a cell that was co-active with cells ABC on L will
be co-active with cells DEF on R"). To determine the contributions of *mean firing rate*
relationships, we applied different firing rate normalization methods to each cell's L and R
activity independently (note that normalizing each cell's L and R joint activity is not appropriate
because it introduces an artificial anticorrelation between L and R that the hypertransform will
exploit). L2 normalization preserved a significant but smaller amount of cross-subject prediction
compared to the unnormalized data (Figure S5b), while Z-scoring firing rates abolished
cross-subject prediction (i.e. did not outperform chance level; Figure S5c, see figure legend for
our interpretation of this difference). Reduced cross-subject prediction after firing rate
normalization shows that firing rate relationships between L and R are an important component
of the prediction. To determine the contributions of specific *firing locations*, we performed an
additional shuffle that circularly shifted the R activity by a different random amount for each cell
(see Figure S2 and *Methods*). The full (non-shuffled) cross-subject prediction outperformed this
circular shuffle (p < 0.001 for Wilcoxon signed rank test vs. 0; Figure S10). Taken together,
these analyses demonstrate that both single-cell firing rate and firing location relationships
contribute to cross-subject prediction.

Discussion

We have shown that it is possible to predict across subjects, better than chance, how a given
experience will be encoded in the hippocampus. In particular, we predict how a "target" subject
will represent the right arm of a T-maze (R), given (1) how that same subject represents the left
arm (L) of the same maze, and (2) the relationship between L and R activity in a different
"source" subject. Control analyses based on within-subject prediction and a comparison of the
properties of various simulated data sets and shuffles with the real data suggests that this
cross-subject prediction is unlikely to be the result of trivial relationships such as cells with

symmetric firing fields, fixed field mapping rules like the start of the L being mapped to the end
of R, or simple rules such as exclusive-or. Thus, our results imply that the hippocampal
encoding of different locations in space, commonly reported to be random *within*
subjects[2–4,6,7], in fact has a shared structure *between* subjects.

There have long been indications in the literature that the space of possible neural activity
patterns in the hippocampus is constrained in a number of ways, perhaps most obviously by a
continuous attractor-type structure that enforces a single, coherent "hill of activity" to exist within
a map (chart) at a time[22]. A different example is how the encoding of a novel experience is
constrained by internally generated activity preceding that experience ("preplay")[23,24], and
most recently, the demonstration that the formation of new place fields in response to
optogenetic stimulation is predictable from activity prior to the stimulation[25]. Our findings are
congruent with these overall notions, but add a new dimension in its ability to test and reveal
structure shared *between* subjects. This approach is attractive because it provides rigorous,
quantifiable tests of how generalizable a given model of neural activity is, and because it can
provide insights into what is shared and what is unique between subjects. Using calcium
imaging data from hippocampal ensembles, Rubin et al. (2019)[26] showed that the location of
one animal could be decoded using a decoder trained on data from another animal. Our
approach is similar in that it also uses cross-subject prediction, but addresses a different
question in that we seek to predict not the location of the animal, but the way different and
potentially not-yet-experienced locations will be encoded (i.e. the tuning curves).

Unlike in the rodent literature, a substantial number of human fMRI and ECoG studies have
used cross-subject prediction[11–13,27,28]. Particularly effective are procedures that do not only
align across subjects anatomically (e.g. by mapping each subject to a reference brain) but
functionally, i.e. by finding structure in how related experiences are represented, even though
across subjects the same experience may be represented very differently. Haxby et al. (2011)[11]
refer to such shared structure as "representational geometry", an idea implemented here as a
rotation and translation (Procrustes transformation) in ensemble neural activity. Because this
notion is abstracted away from the raw data and describes a relatively general class of
relationships, the question arises what specific features of the data make cross-subject
prediction possible. In particular, can this abstract geometric concept be reduced to a simpler
set of rules? Hypertransform predictions do not appear to be reducible to single-cell rules, but
rather are based on a mapping in ensemble activity space (i.e. how a given L cell's activity
changes between L and R depends on what cells are co-active with it). This idea is illustrated by
the examples in Figure S3, and supported more formally by (1) comparison of correlations in the
data with simulations (Figure 4), and (2) outperforming both row (firing rate) and shift (firing
location) shuffles. We note that the contribution of firing location differences is smaller compared
to firing rate differences, as shown by comparing the middle rows of Figure 2 and Figure S10,
and by the inability to outperform row shuffles when data is z-scored (Figure S5c; this tends to
introduce noise that masks underlying regularities); nevertheless, these analyses demonstrate a
contribution of both to an underlying ensemble-level regularity.

What circuit-level or hippocampal input properties are involved in this shared geometry? One possibility is suggested by the relationship between grid cells and place cells: Dordek et al. (2016)[29] showed that applying nonnegative principal component analysis (PCA) to ensemble activity of place cells yields grid cell-like activity patterns. In addition, grid cell firing patterns were known to remain intact but realign linearly (and differently for different subjects) during global remapping[30]. This suggests that our hypertransform procedure may rely on a common coding strategy that transforms grid cell activity from one (part of an) environment to another in a predictable manner (e.g. by aligning to a geometric axis in the environment), a possibility supported by recent work[31,32]. Similarly, a related possible cause of a shared representational geometry is input from the head direction (HD) system, which is expected to be "similarly different" across animals for the initial parts of the left and right trials in our data. Specifically, whatever HD cells are active for "left" trials would explicitly not be active for "right" trials due to the ring attractor organization of the HD system[33]. Although the hippocampal place cells in our data set do not behave in this literal exclusive-or pattern expected for head direction cells (we do not see negative left-right cell-by-cell correlations in Figure 3b), similarities in head direction across identically oriented compartments result in overlapping place cell maps compared to differently oriented compartments[34,35], suggesting that a common head direction cell input may contribute to the cross-subject relationship between left and right trials in our hippocampal data (if not masked by symmetric cells as in the Gupta data set). If this is indeed the case, then our cross-subject hypertransform method would be able to predict, say, the place map of compartment oriented along the 45° axis in Grieves et al. (2018)[35] based on the 135° compartment and the relationship between the 45°-135° compartments in a different subject, whereas in the Spiers et al. (2013)[34] experiment, where the compartments all share the same orientation, it would be difficult to outperform the identity prediction.

In general, hippocampal representational similarity is shaped by many factors including not only the geometry of the environment as mentioned above, but also the identity of specific cues and their spatial configuration, the passage of time, and the animal's internal understanding of the task structure[10,36]. In the two data sets we examined here, left and right trials are associated not only with different locations in space, but also with different head directions (at least for the part immediately after the choice point), different room cues, and different textures and rewards (for the Carey data). Based on these data we do not yet know which of these factors are subject-specific versus generalizable across subjects -- in other words, to what extent these factors contribute to cross-subject prediction. Future studies that systematically manipulate these factors while holding the others constant will reveal under what conditions cross-subject predictions, including those for experiences the subject has not yet had, can be obtained. More generally, uncovering the ways in which hippocampal activity is non-random can ultimately inform how processes such as generalization and structure learning are realized in neural circuits.

Acknowledgements

We thank Jim Haxby and Feilong Ma for technical advice on the hyperalignment procedure;
Youki Tanaka, Caleb Kemere, and the participants of the 2018 Methods in Neuroscience at
Dartmouth summer school for discussion; Neil Burgess and Sam Mckenzie for valuable
comments and suggestions, and A. David Redish for the use of one of the data sets in this
study. This work was supported by HFSP Young Investigator Award RGY0088/2014, NWO
VENI grant 863.10.013 and NSF CAREER award IOS-1844935.

Author contributions

HTC wrote computer code and performed data analysis, in part using tools developed by JRM
and MvdM. HTC and MvdM jointly conceived the analyses and discussed the results, with
suggestions from JRM. HTC and MvdM wrote the paper with comments from JRM.

Declaration of Interests

The authors declare no competing interests.

 Figure Legends

Figure 1: **Workflow and example for cross-subject prediction using the hypertransform**.
(A) Our objective is to predict place cell activity on the right arm (R) of a T-maze in a "target"
subject, based on (i) place cell activity in the left arm (L) in the target subject, and (ii) L and R
place cell activity in a different, "source" subject. These input data are shown in the top left
panel: both the source and target subjects have two matrices each that describe, for each
recorded neuron, how its activity varies during left and right trials. Neurons are ordered
according to their peak firing time on the R arm. Note that although the number of time bins is
the same across subjects, the number of recorded cells may be different. Therefore, the first
step of the analysis workflow is to apply principal component analysis (PCA), resulting in neural
activity trajectories for left and right trials (red and blue, respectively) in each subject's own PCA
space. Three principal components are shown here for display purposes, but in the main
analysis 10 PCs were used. Next, these neural activity trajectories are mapped into a common
space using a "hyperalignment" procedure that minimizes the Euclidean distance between the
trajectories across subjects (step 2, see *Methods* for details). In this common space, a
Procrustean transformation[16] (HT in step 3) is derived that maps L to R trajectories for the
"source" subject (step 3), which can then be applied to the L trajectory of the "target" subject
(step 4) to obtain its predicted R trajectory in the common space (step 5). This predicted R
trajectory is then projected back to the "target" PCA space using the inverse of the matrix used
in step 2 (step 6) and expanded back into the target's original neuron space (step 7). Finally, the
predicted R neural activity is compared to the actual R activity to yield an error measure (step
8). The diagonal pattern apparent in the predicted data indicates similarity between actual and
predicted R, although differences are also visible. **(B)** Close-up of example target L and R
activity matrices ("Actual", neurons ordered by temporal fields on R) with example predicted R
activity derived from a different source session. Each matrix column describes ensemble neural
activity for a single time bin, and maps to a corresponding point on a neural activity trajectory in
common space (top panels). The hypertransform (HT) is a mapping from L to R activity that
operates on these (aligned) ensemble activity vectors. As a result, the R prediction for any given
cell is not only based on that cell's L activity alone, but also depends on the activity of the other
cells at that time; see Figure S3 for further examples and a more detailed explanation.

Figure 2: Cross-subject prediction of R trials of a "target" subject based on how a
"source" subject encodes L and R trials outperforms prediction based on shuffled
source data. For each source-target pair in the Carey data, we computed a z-score of the
actual observed error between predicted and actual R trials (based on the hypertransform
procedure, "HT") compared to a shuffled distribution in which the R rows of the source subject
were randomly permuted. Thus, lower z-scores indicate lower error and therefore better
prediction than chance. Across all source-target pairs, this z-scored error varied depending on
the pair used (column **A**, top row), but was lower than chance overall, as indicated by a shift in
the z-score histogram relative to 0 ("HT" green bars in column **A**, middle row; median: -3.10 +/-
0.54, SEM across unique source-target pairs, $p < 0.001$ for Wilcoxon signed rank test vs. 0).
Cross-subject predictions based on the L-R transform in common space ("hypertransform")

outperformed predictions based on the L-R transform in PCA space ("PCA" blue bars in column **A**, middle row, inset; see *Methods* for details; HT < PCA: 99.62% of source-target pairs, p < 0.001 for binomial test) in terms of raw error between predicted and actual R trials (median of HT raw error: 443.73 +/- 62.44, PCA: 1409.14 +/- 204.22, split-half: 130.99 +/- 46.72, red triangle in inset; see *Methods* for details). Next, we applied the same analysis to a different data set ("Gupta", bottom row, in which the L and R maze arms were more similar to each other than in the "Carey" data), and found that the hypertransform prediction was again significantly better than chance ("HT"; median: -2.59 +/- 1.44, p < 0.001, bottom row) and better than PCA-only (HT < PCA: 79.58% of source-target pairs, p < 0.001; column **A**, bottom row, inset; median of HT raw errorl: 115.26 +/- 142.05, PCA: 121.42 +/- 208.23, split-half: 37.39 +/- 10.01). Columns **B** and **C** use the same layout as column **A**, but using different metrics to describe prediction accuracy. **B** uses the raw error (between predicted and actual R neural activity; lower error/negative indicate better prediction) compared to the mean of the shuffle distribution, and **C** uses the proportion of the shuffle distribution with smaller error than the actually observed error (lower proportions indicate better prediction). For the raw error measure, the HT prediction was better than chance in both the Carey data (column **B** middle row; median: -652.87 +/- 486.05, p < 0.001 for HT vs. 0; red line indicates split-half prediction) and the Gupta data (**B** bottom row; median: -594.62 +/- 735.37, p < 0.001 for HT vs. 0).

Figure 3: Cross-subject prediction outperforms within-subject prediction only in the absence of cell-by-cell correlations. A: Comparison of cross-subject prediction error ("hypertransform", green bars; HT) with within-subject prediction error ("identity transform"; blue bars, ID) for two different data sets. In the "Carey" data (left panel) the left and right arms of the maze had different texture and color patterns; in the "Gupta" data (right panel) the two maze arms were identical. For the Carey data, cross-subject prediction was significantly better than within-subject prediction (HT < ID: 100% of unique source-target pairs, p < 0.001 for binomial test) whereas for the Gupta data, this difference was not significant. **B**: Cell-by-cell correlations of firing rates between L and R arms (i.e. row-wise correlations of the L and R matrices in Figure S2c), were significantly higher in the Gupta data compared to the Carey data (Gupta: median r = 0.21 +/- 0.27, SEM across subjects, Carey: median r = -0.02 +/- 0.12, Wilcoxon rank sum test, p < 0.001). Cell-by-cell correlations in the Carey data were not significantly different from 0 (p = 0.44 for Wilcoxon signed rank test vs. 0). **C**: Population vector (PV) correlations between ensemble activity at each time point and every other time point, i.e. column-wise correlations of the L and R activity matrices, averaged across sessions. Both Carey and Gupta data sets show high correlations around the diagonal, indicating an overall autocorrelation in time; however, the Gupta data additionally shows high off-diagonal correlations between L and R which are barely visible in the Carey data. **D**: Quantification of the median PV correlation between L and R (i.e. the values along the diagonal of the lower left quadrant in **C**). For Gupta data, this correlation is remarkably high (median r = 0.95 +/- 0.16) whereas for Carey data, it is significantly lower (median r = 0.09 +/- 0.10, p < 0.001 for Wilcoxon rank sum test) but significantly different from 0 (p < 0.001 for Wilcoxon signed rank test vs. 0), consistent with previous reports of global remapping[2,4]. The above results explain why the HT is not needed for Gupta data to achieve

better than chance predictions (as shown in Figure 2, bottom row): L and R activity is sufficiently
similar such that the L trajectory alone in either PCA or common space can predict R activity.

Figure 4: Simulations demonstrate that representational geometry, but not simple rules
such as exclusive-or and firing rate correlations, result in cross-subject prediction while
being consistent with the data. A: Example L and R activity matrices (left column) and
histogram of z-scores of cross-subject prediction compared to the distribution of shuffle
predictions (right column; z-scores lower than zero indicate better-than-chance predictions) of
four simulated data sets: (1) neurons have a fixed, independent probability (0.5) of having a 1-D
Gaussian place field on L and/or R, with the three parameters of time, peak firing rate (FR) and
width randomly and independently chosen for L and R (**ind-ind**, top row), (2) neurons only have
a field on *either* L or R but not both, and parameters of the field are chosen randomly as in (1)
(**x-or**, second row), (3) neurons have a fixed independent probability of having L and R fields as
in (1) but with the additional constraint that neurons with both L and R fields must have the
same three parameters (**ind-same-all**, third row) and (4) the activity on L is simulated by
assigning each neuron an independent probability of having a field whose parameters are
randomly chosen, then the activity on R is obtained by applying a L-R transform
(hypertransform, HT) from the Carey data to the simulated L activity (**sim. HT**, last row). As
expected, in the **ind-ind** (independent) case, cross-subject prediction is not possible (z-score
vs. chance, median: -0.01 +/- 0.03, SEM across unique source-target pairs, p = 0.70 for
Wilcoxon signed rank test vs. 0). In contrast, **x-or**, **ind-same-all** and **sim. HT** all show
better-than-chance cross-subject predictions (median: -1.26 +/- 0.30 for **x-or**, median: -2.62 +/-
0.06 for **ind-same-all**, median: -5.23 +/- 0.04 for **sim. HT,** all p < 0.001 for Wilcoxon signed rank
test vs. 0), indicating that if there is a non-random L-R relationship in the underlying data, the
hypertransform procedure can exploit it. Note that although this synthetic example shows
substantially negative predicted firing rates, negative firing rates, when they did occur, tended to
be much smaller for predictions using actual data. **B**: Population vector (PV; column-wise)
correlations between ensemble activity at each time point and every other time point of the L
and R activity matrices. Only **ind-same-all** shows high off-diagonal correlations between L and
R, resembling the Gupta data set in which L and R arms were identical and firing activity on
both arms is highly correlated (compare with Figure 3c). In **x-or**, off-diagonal correlations are
slightly negative. **C**: Quantification of the median PV correlation between L and R (i.e. the
values along the diagonal of the lower left quadrant in **B**). PV correlations between L and R in
ind-ind are zero, but L and R are positively correlated in **ind-same-all** (median r = 0.40 +/-
0.01) since for every time point in L where there is a field, the same ensemble activity appears
at the same time point in R with probability 0.5. **X-or** shows a negative correlation (median r =
-0.18 +/- 0.02) because for every time point in L where there is a field, the same ensemble
activity would deterministically be absent in R, and vice versa. None of these simple rules are
consistent with the PV correlation found in the Carey data; however, **sim. HT** does show similar
correlations as the data (median r = 0.04 +/- 0.03 for **sim. HT** and median r = 0.09 +/- 0.10 for
Carey). **D**: Cell-by-cell (row-wise) correlations of L and R show that **ind-same-all** is more
correlated than the data, whereas **sim. HT** yields similar correlations (median r = 0.04 +/- 0.02
for **sim. HT** and median r = -0.02 +/- 0.12 for Carey). Thus, taken across panels C and D,

simple rules (**x-or** and **ind-same-all**) are inconsistent with the data, but the representational
geometry embodied in the **sim. HT** yields correlations that are similar to the data.

STAR Methods

Raw data

We used two different data sets containing ensemble recordings of hippocampal CA1 neurons in rats performing T-maze tasks.

The first data set ("Carey") is as described in van der Meer et al. (2017)[17] and Carey et al. (2019)[17,18]. Briefly, male rats (n = 4) performed daily sessions on a T-maze where they had free choice between left and right arms. Rats were alternately food- and water-restricted across days; the left arm resulted in food reward (five 45 mg pellets), the right arm resulted in water reward (~0.2 ml sucrose solution). Rats ran 15-20 discrete trials per recording session, with no less than 5 trials for the least preferred choice (left or right). Only sessions with at least 40 simultaneously recorded neurons were included, leaving 19 of 24 total sessions for analysis (range: 50-178 neurons per session).

The second data set ("Gupta") is as described in Gupta et al. (2010)[19] and Gupta et al. (2012)[19,20]. Briefly, male rats (n = 4) performed daily sessions on a continuous Multiple-T maze with free choice between left and right arms. Food pellet reward (four 45 mg pellets) was available either by choosing left only, right only, or alternating between left and right; which reward schedule was in effect was determined pseudorandomly at the start of daily recording sessions. In addition, the reward schedule switched approximately halfway throughout the session. Only sessions with at least 40 simultaneously recorded neurons were included, this left 14 of 42 total sessions for analysis (range: 41-101 neurons per session).

Both data sets consist of left (L) and right (R) trials from the choice point to the first reward site; the analyses in this study are concerned with how the relationship between L and R trials is encoded in hippocampal ensemble activity. To avoid the possibility that neural activity on a common trajectory shared between L and R is the main predictor of L and R relationship, data from the central stem of the maze was excluded.

Data preprocessing

Preparation of input data. Both data sets were preprocessed to obtain two types of neural activity matrices that form the starting point for all analyses (Figure S2). The first and main data type is the "Q-matrix", which describes binned firing rate over time for simultaneously recorded neurons *[nNeurons x nTimeBins]* and is used in all main analyses. The second data type is the "TC-matrix" (place turning curves) matrices of dimension *[nNeurons x nSpaceBins]* for Figure S4 and S5).

Since both data sets contain different numbers of L and R trials within a session, trials were first subsampled so that an equal number of L and R trials were used. Next, because trials differed in length because of variations in running speed, all trials were truncated to the last 2.4 seconds

(the median time between passing the choice point and reaching the reward site) so that only
after-choice-point data was used.

To obtain Q-matrices for L and R trials, binned firing rate matrices (time bin width: 50 ms) were
created for individual trials, smoothed with a window size = 1 s, σ = 50 ms Gaussian kernel, and
then averaged across trials within each session.

To obtain TC-matrices for L and R trials, spike firing data with only running speed > 5 cm/s data
were averaged for each place bin (~3 cm per bin) across each session, then smoothed with a
window size = 11 bins, σ = 1 bin Gaussian kernel. Only data from the last 41 place bins were
included so that only after-choice-point data was used.

Criterion for exclusion of interneurons (analyses used in Figure 3 and Figure S5). Neurons with
mean firing rate > 10 Hz across the entire recording session were classified as putative
interneurons. These were excluded for the correlation analysis in Figure 3, because otherwise
variations in firing rates between putative interneurons and projection neurons would dominate
the population vector correlations (described below). In Figure S5a we verify that inclusion of
interneurons was not required for the main results, and noted that source sessions whose
cross-subject predictions were consistently worse than the others (for instance, row 2 of R1 in
Figure 2a, top row) were sessions without putative interneurons.

Normalization (analyses used in Figure S5). Normalization of the input data was conducted by
dividing the L2-norm of each row (neuron) or z-scoring each row of data matrices,
independently for the L and R parts of the input data matrices. Although it may seem intuitive to
normalize each entire row of the input data (i.e. L and R data together), this actually introduces
an artificial anticorrelation between the L and R parts of the matrix, such that even on data
where no relationship exists between L and R, a relationship is introduced by normalization.
Thus, we avoided normalization across L and R when testing cross-subject prediction. Note that
normalization was only performed on data with putative interneurons removed (Figure S5).

Hypertransform analysis procedure

Overview. The overall objective of the hypertransform procedure is to predict R data in a "target"
subject based on (1) the target subject L data, and (2) the L and R data of a different "source"
subject (see Figure 1 for a complete description and schematic). Each step of the procedure is
described in detail below.

Each recording session was used as source and paired with all sessions from all other subjects
to form cross-subject source-target pairs (260 unique source-target session pairs for the Carey
data and 142 pairs for the Gupta data).

PCA. After preprocessing the data as described above, principal component analysis (PCA; *svd*
function in MATLAB R2018b) was applied to concatenated L and R neural data matrices to

reduce to the same dimensionality because (1) there were unequal numbers of neurons
recorded across sessions, and (2) to aid in generalization across sessions. Ten principal
components (PCs), accounting for approximately 95% of the variance (Figure S8d) were then
used to project L and R matrices into neural activity trajectories in each subject's own PCA
space.

Hyperalignment. Hyperalignment is a procedure that maps neural activity from a number (>1) of
input sessions into a common space. It does so by iteratively finding the set of Procrustes
transformations (one for each input session) that minimize the Euclidean distance between input
trajectories in the common space based on rotation and translation (and scaling, in some
implementations; we did not use scaling in this study). This procedure is commonly used to
align fMRI activity trajectories in subject-specific spaces into a common representational space
(Haxby et al. 2011)[11]. In our specific version of this procedure, we aligned L and R neural
trajectories in the PCA spaces from a source subject and a target subject into a common space
by using the *hyperalign* function in *hypertools*[15] *(matlab version)*. This results in two matrices,
one for each subject, that map each subject's PCA space into the common space.

Hypertransform. In this common space, another Procrustes transformation was derived to find
the mapping between the L and R neural trajectories of the source subject (as above, scaling
was disabled). This transformation (hypertransform, HT) was then applied to the target subject's
L trajectory to yield a *predicted* R trajectory. Note that this transformation is not only
subject-specific but *source-target-pair-specific* since the common space is unique for each
source-target pair used.

Projection back into neural space. Next, the inverse of the subject-specific hyperalignment
matrix (projecting from PCA to common space) was applied to project the predicted R trajectory
back to the target subject's PCA space. Principal components obtained earlier were used to
reconstruct the predicted R trajectory in the PCA space into the predicted R data matrix (Q or
TC) for the target subject.

Shuffles and associated metrics. The prediction error for a specific source-target pair is
calculated by summing squared errors between predicted and actual R data matrices of the
target subject and dividing by the number of neurons. This error is compared against a chance
distribution by permuting (shuffling) the rows of the R data matrix of the source subject (but
keeping the L matrix intact; **row shuffle**, see Figure S2b) and repeating the hypertransform
procedure. This shuffle breaks any relationship between L and R activity at the cell-by-cell level.
For example, if the relationship between L and R firing activity is random, then permuting the
source R data matrix in this manner will make no difference in the ability to predict the target R
data matrix. As a result, for random data (such as our "independent" simulation in top row of
Figure 4a), shuffled predicted errors should not be different from the actual observed predicted
error. Note that row shuffles can create idiosyncratic principal components not present in the
original data, affecting the extent to which source and target sessions can be aligned. Because
hyperalignment (when used on two sessions, as in this study) "anchors" to the source session,

asymmetries in the z-scored error relative to shuffle can arise, i.e. source-target and
target-source can have numerically different z-scores.

Next, the **shift shuffle** preserves the mean firing rates of L and R for each neuron, but breaks
the temporal relationship between L and R by circularly shifting each row of the source R matrix
independently by a length randomly chosen from [1, number of total bins] (while keeping the L
matrix intact; Figure S2b). If cross-subject prediction based on the true data outperforms this
shuffle, then the specific firing fields matter, above and beyond mean firing rate relationships
between L and R.

For each shuffling method, 1000 shuffles were conducted for each source-target pair, resulting
in 1000 shuffled prediction errors which were compared to the prediction error observed for the
true data using three metrics: (1) z-scoring the actually observed error relative to the distribution
of shuffle predictions (lower z-scores indicate better prediction), (2) raw error compared to the
mean error of the shuffle distribution (lower observed error indicates better prediction), and (3)
proportion of the shuffle distribution whose error was smaller than actual observed error (lower
proportion indicate better prediction).

Identity transform (used in Figure 3). To test if a better-than-chance cross-subject prediction is
simply due to the similarity of L and R neural activity within the target subject, a within-subject
prediction is obtained by using a duplicate of target subject's L trajectory in the common space
as the predicted R trajectory, i.e. making the L-R transformation equal to the identity
transformation. This predicted trajectory was then used to reconstruct the predicted target
subject's R data matrix as in the hypertransform procedure.

Split-half prediction (used in Figure 2 and Figure S4 insets). To compare the accuracy of
cross-subject hypertransform prediction to the best possible prediction given the variability in the
target R data, we obtained a split-half prediction by averaging across a randomly-chosen half of
right trials from the target session, resulting in a "lower bound" prediction error.

Variation: withheld data (used in Figure S4). To determine the importance of including the data
to be predicted (R) in the hyperalignment step, we performed a number of analysis variations
withholding different amounts of R data. In the first variation, we withhold all target R data.
Specifically, only source and target L were hyperaligned into the common space, in which the
source R data was projected from the PCA space using the source matrix obtained from L
alignment between the two subjects. The source L-R transformation and the predicted R matrix
were then obtained as in the default hypertransform procedure. (Replacing the target R data
matrix with zeros before hyperalignment yielded similar results to complete withholding, so we
report only complete withholding results.).

In the second variation, a randomly-chosen single right trial from the target session was used for
hyperalignment, and the average of the remaining trials were withheld as target R to be

predicted (Figure S4b). The rest of the procedure follows the above hypertransform procedure.
The split-half prediction in this procedure was obtained from half of the withheld trials.

Variation: PCA only (used in Figure 2 and Figure S4). To test whether a relationship between L
and R already exists in subject-specific PCA spaces, we modified the hypertransform procedure
as follows: instead of aligning neural activity trajectories in the common space through
hyperalignment, a L to R linear (Procrustes) transformation was derived from the source
subject's PCA space and directly applied to the L trajectory in target subject's PCA space to
obtain the predicted R trajectory. This predicted trajectory was then used to reconstruct the
predicted target subject's R data matrix as in hypertransform procedure.

Simulations

As a first step towards understanding the possible explanation for cross-subject prediction,
simulated neural activity matrices were generated using 1-D Gaussians with three parameters:
time (the time bin where a neuron has its maximal firing activity; randomly chosen from [1, 48]
out of 48 time bins), peak firing rate (FR; randomly chosen from [10, 20] Hz) and width
(randomly chosen from [2, 7] bins standard deviation). Several scenarios were created to test
the possibility that different potential place cell properties may account for the observed
cross-subject prediction:

- **ind-ind**: Neurons have a fixed, independent probability (0.5) of having a 1-D Gaussian
place field on L and/or R, with the three parameters of time, peak firing rate (FR) and
width randomly and independently chosen for L and R.
- **x-or**: Neurons only have a field on *either* L or R but not both, and parameters of the field
are chosen randomly as ind-ind
- **ind-same-all**: Neurons have a fixed independent probability of having L and R fields as
in **ind-ind** but with the additional constraint that neurons with both L and R fields must
have the same three parameters.
- **sim. HT**: The activity on L is simulated by assigning each neuron an independent
probability of having a field whose parameters are randomly chosen, then the activity on
R is obtained by applying L-R transform (hypertransform, HT) from real data (Carey) to
the simulated L activity.
- **ind-same-time**: Similar to **ind-same-all**, neurons have a fixed independent probability of
having L and R fields as in **ind-ind,** but if when a cell has fields in both, one of the three
parameters: **time** is the same for L and R, and FR and width are independently chosen.
- **ind-same-FR**: Similar to **ind-same-all**, neurons have a fixed independent probability of
having L and R fields as in **ind-ind,** but if when a cell has fields in both, one of the three
parameters: **FR** is the same for L and R, and time and width are independently chosen.
- **ind-same-width**: Similar to **ind-same-all**, neurons have a fixed independent probability
of having L and R fields as in **ind-ind,** but if when a cell has fields in both, one of the
three parameters: **width** is the same for L and R, and time and FR are independently
chosen.

For each simulation scenario, we generated a matching number of sessions (19) as in Carey
data and a matching number of neurons within each session as a dataset. To avoid the
possibility that one particular randomly generated dataset biases the results, 100 datasets were
created and all statistics were averaged across datasets to match the real data.

For each of the 100 datasets in **sim. HT**, a randomly chosen session from the Carey data was
used to hyperalign with simulated L activity (R activity was replaced with zeros first). The
hypertransform obtained from the real session was then applied to the simulated L trajectory in
the common space to obtain a simulated R trajectory. This simulated trajectory was then used to
reconstruct the simulated R activity as in hypertransform procedure.

Correlation analysis of neural and simulated data

Cell-by-cell (row-wise) correlations. For each cell (a row in the neural data matrix), a correlation
coefficient between L and R was computed (see schematic in Figure S2). Whitening noise
(matrices of same size as L and R, in which each element is a number sampled from *Uniform*(0,
10^{-5})) was added so that a coefficient can be computed when there is no activity on either L and
R trials, which would otherwise result in zero variance.

Population vector (PV; column-wise) correlations. L and R data matrices were (horizontally)
concatenated, and a correlation coefficient was computed between each time point (a column)
and all other columns of the L and R concatenated matrix (see schematic in Figure S2). Higher
off-diagonal values (the values along the diagonal of the lower left quadrant) of this
concatenated matrix indicate higher correlated ensemble neural activity between L and R.

For both cell-by-cell and PV correlations, results were combined across subjects by appending
all cells and sessions into a pooled data set regardless of what subject they came from.

All correlation coefficients were computed using the MATLAB *corrcoef* function.

Statistics

Comparison to shuffles. For the hypertransform prediction error (between predicted and actual
neural activity) of each source-target pair, a number of measures were computed: a z-score
against the shuffled distribution, the proportion of shuffles with lower error, and the raw error
compared to the mean of the shuffle errors. Note that raw errors were normalized by dividing by
the number of neurons in the target session so that the prediction accuracy does not depend on
the number of neurons being predicted. These measures are shown for each individual
source-target pair as an element in a matrix such as one in Figure 2 (top row). Then, results
across all source-target pairs are combined and plotted as a histogram; all summary statistics
(median +/- SEM across all unique source-target subject pairs) and significance tests (Wilcoxon
signed-rank tests against zero) are performed on this pooled data.

Comparison of the HT prediction *to PCA-only (Figure 2 and S4) and identity (Figure 3)*
predictions. For each source-target pair, the hypertransform prediction error is compared to the
ones obtained from PCA-only and identity predictions (described in *Identity transform* and
Variation: PCA only sections respectively). Then, results across all source-target pairs are
combined and plotted as a histogram; all summary statistics (percentage of unique
source-target subject pairs whose HT predictions are better) and significance tests (binomial
test) are performed on this pooled data.

Identification of place fields. For Figure S9 only, we identified neurons as having a place field if
three consecutive bins had a firing rate >5 Hz,on either the left arm (L), the right arm (R), or
both. Only neurons with fields on both arms (6.98% of all neurons) are included in the analysis
investigating the systematic mapping between L and R in Carey data set and in hypertransform
(HT) predictions (Figure S9).

Data and Code Availability

The code used in this study is available on [GitHub](#) (we used commit 654aecb). The "Carey"
data set used in this study is publicly available from DataLad (["MotivationalT" data set,](#)) and the
Gupta data is available from the corresponding author on request.

Supplementary Information

Figure S1: **Schematic of hyperalignment applied to hippocampal place cell activity. (A)**
Schematic example of shared structure in place cells of two subjects on the left (L; locations A,
B, C) and right arms (R; locations D, E, F) of a T-maze (left column). Individual neurons of
different subjects (indexed by 1, 2, 3) have idiosyncratic place field locations for example, neuron
1 of subject 1 has place fields in A and F, whereas neuron 1 of subject 2 has fields in C and E
(note that for visualization purposes, these neurons have place fields on both arms, but this is
not a requirement for our analysis.) The ensemble neural activity during L and R runs can be
plotted in a subject-specific neural space whose x-axis is the firing rate (FR) of neuron 1, y-axis
is the FR of neuron 2, etc. (middle column; red line indicates trajectory of L and blue indicates
R). Although subjects have different trajectories of L and R in their individual neural spaces, the
trajectories for the two subjects can be made to line up by a rotation of the axes (e.g. rotate the
x-axis of subject 1 to line up with subject 2's y-axis). Hyperalignment finds the subject-specific
transformation that minimizes the Euclidean distances of trajectories between subjects, and
projects those neural activity trajectories into a common space (right column). In this common
space, axes are shared between subjects, and trajectories of different subjects are now aligned
to each other. Note that the common relationship between L and R in this example follows an
unrealistically simple, fixed spatial mapping rule (the neuron firing at the start of L always fires at
the end of R etc.), unlike what the real data suggests (Figure 3C, Figure S9). **(B)**: Minimal
example of shared structure across subjects with essentially uncorrelated cell-by-cell and
column-wise correlations that is more consistent with the data. The rule for simulating this
example is as follows: neurons with high firing rates circularly shift their place fields to the left,
whereas half of low-firing-rate neurons shift their fields to the right (arrows) and the other half

stay fixed. Hyperalignment successfully captures this abstract rule and correctly predicts the place fields in subject 2's right activity matrix (rightmost panel; locations of A to F correspond to left column in **A**). The shared structure in the actual data, which has many more neurons and more varied firing patterns, is more complex; this example is intended to provide an intuition about how cross-subject patterns can exist in the absence of single-cell or population vector correlations.

Figure S2: Data preprocessing and analysis schematics. A: Data preprocessing. Neural ensemble activity for left (L) and right trials (R) in each recording session was shaped into two types of input data matrices: the *Q-matrix*, which describes binned firing rate over time for simultaneously recorded neurons (dimension [nNeurons x nTimeBins]), used in the main analyses, and the *TC-matrix* (spatial tuning curves, dimension [nNeurons x nSpaceBins]). Trials were subsampled to obtain an equal number of L and R trials were used, and truncated to the last 2.4 seconds (the median time taken from the choice point to the end of a trial). Times when the animal was deemed to be stationary were excluded from analysis. **B**: Illustrations of the row shuffling procedure used in the main analysis (Figure 2) and shift shuffling procedure used in Figure S10. To obtain a distribution of chance cross-subject predictions, the analysis steps illustrated in Figure 1 were applied, except that for the "source" subject, the relationship between L and R activity was disrupted by either randomly permuting the rows of the R matrix (*row shuffles*) or circularly shifting each row of R matrix independently by a randomly chosen length (*shift shuffles*). **C**: Illustration of the correlation analyses used in Figures 3-4. *Cell-by-cell correlations* are obtained by row-wise correlating L and R activity for each cell, and then averaging across all cells. *Population vector correlations* are obtained by column-wise correlating activity at each time or location with activity at every other time or location. This yields one correlation matrix per session, which are then averaged across sessions.

Figure S3: Example target L and R activity matrices with predictions. Two example target L and R activity matrices ("Actual", neurons ordered by time of maximum activity on R) with predictions obtained from three source sessions ("Predicted"). In general, for each session, the predicted and actual activity are clearly related, as is apparent from the overall diagonal pattern in the Predicted matrices indicating agreement with the Actual R data (we quantify this in insets of Figures 2 and S4 and associated analyses). However, clear deviations from the actual data are also visible, for instance in predicting fields that do not exist in the actual data (red arrows; compare actual and predicted R), and predicting fields in incorrect locations (appearing away from the diagonal, e.g. white arrow for source session 2). Comparing between different source sessions (1, 2, 3) the predictions derived from all source sessions share an overall similarity, but there are also differences, typically in the specific locations of predicted place fields in a subset of neurons (compare the two white arrows, for instance). A further point to note is that even for some cells that do not have a field on L, we can correctly predict where this cell will have a field on R (gray rectangles in lower panel). Understanding this requires two components. First, how is it that even though the L activity is the same at different time points, the predicted R is different over time? This occurs because the mapping from L to R at each time point depends on the activity of all the other cells in L at that time point. Second, how is that for a given time point, even though the L activity is the same for two different cells, the predicted R is different for those cells? This occurs because different cells have different loadings on the principal components that are hyperaligned. In other words, cross-subject prediction does not apply a fixed rule to the activity of a single cell, but rather applies a mapping that depends on the activity of the other cells in the ensemble.

Figure S4: Better-than-chance cross-subject predictions can be observed even when the to-be-predicted target data was withheld, and spatial turning curves were used. A: Histogram of three cross-subject prediction metrics: z-scores of actual observed error compared to the distribution of shuffle predictions (left column), raw prediction error compared to the mean of the shuffle distribution (middle), proportion of the shuffle distribution whose error was smaller than actual observed error (right). For all metrics, lower numbers indicate better cross-subject predictions. Even when the R activity of the target subject, which is the activity to be predicted, is withheld from the hyperalignment procedure, the hypertransform (HT) prediction is significantly better than chance (median: -6.03 +/- 0.84, SEM across unique source-target pairs, p < 0.001 for Wilcoxon signed rank test vs. 0; left column) and better than PCA-only (HT < PCA: 70.77% of source-target pairs, p < 0.001 for binomial test; left column, inset; median of HT raw error: 1962.93 +/- 209.86, PCA: 2505.85

+/- 287.00, split-half: 114.05 +/- 17.48, red triangle in inset). **B:** Histogram of three cross-subject
prediction metrics as in **A**. Different from **A**, one randomly-chosen trial of target R was used for
alignment, with the average of rest R trials withheld as the activity to be predicted. The HT prediction
is again better than chance (median: -3.03 +/- 0.51, $p < 0.001$ for HT vs. 0; left column) and
significantly better than PCA-only (HT < PCA: 91.92% of source-target pairs, $p < 0.001$; left column,
inset). Note that the HT prediction was substantially improved by including one trial of target R data
(median of HT raw error: 809.73 +/- 374.22, PCA: 1698.19 +/- 345.25, split-half: 108.83 +/- 110.81).
C: Histogram of three cross-subject prediction metrics as in **A** and **B** for neural activity matrices
calculated as a function of locations (turning curves; TC) instead of time (see Figure S2 and
Methods for details) were used. The cross-subject predictions are significantly better compared to
shuffles (median: -2.97 +/- 0.46, $p < 0.001$ for HT vs. 0; left column) and significantly better than
PCA-only (HT < PCA: 100% of source-target pairs, $p < 0.001$; left column, inset; median of HT raw
error: 499.05 +/- 50.22, PCA: 1469.76 +/- 170.06, split-half: 83.86 +/- 64.48), suggesting time and
location yield similar results (compare with Figure 2).

Figure S5: Cross-subject prediction is preserved when removing putative interneurons and normalizing L and R independently by L2-norm, but abolished when independently z-scoring L and R firing rates. A: Example L and R activity matrices with putative interneurons (mean firing rate > 10 Hz) removed, and corresponding cross-subject prediction z-score histograms compared to the distribution of shuffled predictions (z-scores lower than zero indicate better-than-chance prediction) for both temporal (**Q**) and spatial (**TC**) tuning curves. Both Q and TC show significantly better-than-chance cross-subject prediction (Q median: -0.42 +/- 0.46, SEM across unique source-target pairs, p < 0.01 for Wilcoxon signed rank test vs. 0; TC median: -0.34 +/- 0.43, p < 0.001 for HT vs. 0), indicating high-firing rate interneurons are not required for cross-subject prediction. **B**: Same layout as A, but L and R activity matrices were obtained by removing interneurons and dividing separately each L and R row by its L2-norm (vector length). Both Q and TC again show significantly better-than-chance prediction (Q median: -0.26 +/- 0.29, p < 0.001 for HT vs. 0; TC median: -0.43 +/- 0.32, p < 0.001 for HT vs. 0). **C**: Same layout as A, but L and R activity matrices were obtained by removing interneurons and z-scoring separately each L and R row. Unlike the L2-norm case, normalization by z-scoring does not show better-than-chance prediction; we speculate that this may be related to the relatively marginal performance of the L2-normed data compared to unnormalized data (compare panels **A** and **B**) combined with the tendency for the z-scoring especially to assign very large values to isolated pixels with a small but non-zero firing rate, introducing noise. Note that for both L2-norm and z-scoring cases, independently normalizing L and R activity (L separately from R), rather than normalizing L and R together (normalizing the entire row), avoids introducing artifactual anti-correlations between L and R which would be exploited by the cross-subject prediction algorithm even for independent data.

Figure S6: Simulations show that either the same place field firing time, peak firing rate, or place field width alone cannot account for cross-subject prediction in the Carey data. A: Example L and R activity matrices (left column) and histogram of z-scores of cross-subject prediction compared to the distribution of shuffle predictions (z-scores lower than zero indicate better-than-chance predictions; right column) of three simulated data sets. All three data sets are generated by assigning each neuron a fixed, independent probability (0.5) of having a field on L and/or R, but when a cell has fields in both, one of the three parameters of its 1-D Gaussian place field (**ind-same-time**, **ind-same-FR (firing rate)** or **ind-same-width**) is the same for L and R; the other two are randomly and independently chosen. **Ind-same-time** shows better-than-chance cross-subject predictions (median: -2.33 +/- 0.06, SEM across unique source-target pairs, p < 0.001 for Wilcoxon signed rank test vs. zero) but **ind-same-FR** and **ind-same-width** do not, indicating cross-subject predictions cannot be better than chance when time points of fields are unrelated between L and R. **B**: Population vector (PV; column-wise) correlations between ensemble activity at each time point and every other time point of the L and R activity matrices. **C**: Quantification of the median PV correlation between L and R (i.e. the values along the diagonal of the lower left quadrant in **B**). **Ind-same-time** shows highly positive correlation (median r = 0.35 +/- 0.01) since for every location of L where there are fields, there would some chance that some (although random) amount of ensemble activity would appear on the same location of R. In contrast, L and R in **ind-same-FR** and **ind-same-width** are nearly uncorrelated since no prediction of ensemble activity of one location on L can be made based on the same location of R (see also **ind-ind i**n Figure 3). The **ind-same-time** case is potentially consistent with the Carey data as measured by PV correlations. **D**: Cell-by-cell (row-wise) correlations of L and R. Here, the **ind-same-time** correlations are higher than in the Carey data (median r = 0.23 +/- 0.02 vs. median r = -0.02 +/- 0.12 for Carey), indicating that this scenario does not accurately capture the source of cross-subject prediction in the data.

Figure S7: Differences in running speeds and firing rates between source-target subject pairs do not explain better-than-chance cross-subject hypertransform (HT) predictions. A: Average running speed across time points on both left trials (L) and right trials (R) for each subject (left column). There are significant differences between L and R ($F_{(1, 14745)}$ = 48.99, p < 0.001), between subjects ($F_{(3, 14745)}$ = 1406.95, p < 0.001) and the interaction of both ($F_{(3, 14745)}$ = 448.37, p < 0.001) in running speed. Given this variability in speed, we tested whether differences in average running speed between source and target sessions were related to prediction accuracy (HT z-score matrix in Figure 2a, top row), reasoning that more similar speeds may enable better predictions. Contrary to this expectation, we did not find a significant correlation between absolute running speed differences and prediction accuracy (r = -0.01, p = 0.97; the specific quantity computed was $|\texttt{speed}_{\texttt{source}}-\texttt{speed}_{\texttt{target}}|$). Since there is an interaction between L/R and subject, we also calculated $\texttt{spd-diff}_{\texttt{session}} = |\texttt{speedL}_{\texttt{session}}-\texttt{speedR}_{\texttt{session}}|$ for each session and a specific quantity $|\texttt{spd-diff}_{\texttt{source}}-\texttt{spd-diff}_{\texttt{target}}|$ for each source-target pair, capturing how similar (or different) L/R speed differences are between source-target pairs (right column). Again, we did not find L/R speed differences significantly correlate with the prediction accuracy (r = -0.05, p = 0.38). **B:** Average firing rates (FR) across time points and neurons on both left trials (L) and right trials (R) for each subject (left column). There are also significant differences between L and R ($F_{(1, 181624)}$ = 16.13, p < 0.001), between subjects ($F_{(3, 181624)}$ = 178.16, p < 0.001) and the interaction of both ($F_{(3, 181624)}$ = 132.31, p < 0.001). Again, we repeated the analysis in **A** and did not find significant correlations between the prediction accuracy and the absolute difference in average FR between source-target pairs (r = 0.06, p = 0.30), and between the prediction accuracy and the absolute L and R difference in FR between source-target pairs (r = -0.06, p = 0.32).

Figure S8: Cross-subject hypertransform (HT) prediction is not due to particular time points such as the start or end of the trial. A: Average firing rates (FR) across neurons for each time point for both left trials (L) and right trials (R) in the Carey dataset. **B:** Comparison of average firing rate across neurons over time with average predicted firing rate. Although FR of both L and R varied across time points, we did not find any time point at which FR is substantially high or low, which could potentially be exploited by the HT prediction, for example a high activity at the start of L is always mapped to the end of R. Similarly, the predicted R activity and its difference with actual R also show varying patterns but not a particular time point is highlighted. **C:** Z-scores of cross-subject prediction errors (normalized within each session) as a function of time, averaged across sessions. Errors varied across time but did not highlight particular time points, indicating that cross-subject prediction is not disproportionately due to certain time points such as the end of the trial. **D:** Explained variance as a function of the number of principal components (PCs). In our hypertransform procedure, 10 PCs were used, which accounts for ~95% of the variance of data.

Figure S9: No obvious single-cell mapping between left (L) and right (R) trials for both actual
and predicted data. To investigate if there is a systematic mapping between in L and R in actual
data and hypertransform (HT) predictions, we plotted temporal field indices of L and R for neurons
which have fields on both (6.98% of all neurons) on a matrix where x-axis are temporal field indices
of L and y-axis are indices of R (see *Methods* for how fields were detected). If a neuron has a
temporal field on the end of L and a field on the start of R, it is counted as 1 on the bottom right
corner of the matrix, and total counts are divided by the number of neurons included. We did not
observe a simple rule characterizing mapping in temporal fields between L and R in actual data; for
example, it is not the case that neurons at the end of L are systematically being mapped to the start
of R. Other than some subtle identity predictions (along the diagonal), we did not find an obvious
single-cell mapping between L and predicted R, indicating that the HT prediction captures
ensemble-level relationships rather than single cell relationships.

Figure S10: Cross-subject hypertransform (HT) outperforms temporally shuffled source data.
For each source-target pair in the Carey data, we computed a z-score of the actual observed error
between predicted and actual R trials (based on HT) compared to a shuffled distribution in which
each R row of the source subject was *circularly shifted independently by a randomly chosen length*.
Lower z-scores indicate lower error and therefore better prediction than chance. Across all
source-target pairs, this z-scored error varied depending on the pair used (column **A**, top row), but
was lower than chance overall, as indicated by a shift in the z-score histogram relative to 0 ("HT"
green bars in column **A**, middle row; median: -0.44 +/- 0.29, SEM across unique source-target pairs,
p < 0.001 for Wilcoxon signed rank test vs. 0). Next, the same analysis was applied to a different
data set ("Gupta", bottom row, in which the L and R maze arms were more similar to each other than
in the "Carey" data), and found that the HT prediction was again significantly better than chance
("HT"; median: -3.20 +/- 1.13, p < 0.001, bottom row). Column **B** and **C** use the same layout as
column **A**, but using different metrics to describe prediction accuracy. **B** uses the raw error (between
predicted and actual R neural activity; lower error/negative indicate better prediction) compared to
the mean of the shuffle distribution, and **C** uses the proportion of the shuffle distribution with smaller
error than the actually observed error (lower proportions indicate better prediction). For the raw error

measure, the HT prediction was better than chance in both the Carey data (column **B** middle row; median: -14.12 +/- 12.27, p < 0.001 for HT vs. 0 and the Gupta data (**B** bottom row; median: -206.13 +/- 119.78, p < 0.001 for HT vs. 0).

References

1041

1. McClelland, J. L., McNaughton, B. L. & O'Reilly, R. C. Why there are complementary

learning systems in the hippocampus and neocortex: insights from the successes and

failures of connectionist models of learning and memory. *Psychol. Rev.* **102**, 419–457

(1995).

2. Leutgeb, S. *et al.* Independent codes for spatial and episodic memory in hippocampal

neuronal ensembles. *Science* **309**, 619–623 (2005).

3. Alme, C. B. *et al.* Place cells in the hippocampus: eleven maps for eleven rooms. *Proc.*

Natl. Acad. Sci. U. S. A. **111**, 18428–18435 (2014).

4. Bostock, E., Muller, R. U. & Kubie, J. L. Experience-dependent modifications of

hippocampal place cell firing. *Hippocampus* **1**, 193–205 (1991).

5. Muller, R. U. & Kubie, J. L. The effects of changes in the environment on the spatial firing of

hippocampal complex-spike cells. *J. Neurosci.* **7**, 1951–1968 (1987).

6. Redish, A. D. *et al.* Independence of firing correlates of anatomically proximate

hippocampal pyramidal cells. *J. Neurosci.* **21**, RC134 (2001).

7. Rich, P. D., Liaw, H.-P. & Lee, A. K. Place cells. Large environments reveal the statistical

structure governing hippocampal representations. *Science* **345**, 814–817 (2014).

8. Josselyn, S. A. & Frankland, P. W. Memory Allocation: Mechanisms and Function. *Annu.*

Rev. Neurosci. **41**, 389–413 (2018).

9. Gauthier, J. L. & Tank, D. W. A Dedicated Population for Reward Coding in the

Hippocampus. *Neuron* **99**, 179–193.e7 (2018).

10. Kubie, J. L., Levy, E. R. J. & Fenton, A. A. Is hippocampal remapping the physiological

basis for context? *Hippocampus* (2019) doi:10.1002/hipo.23160.

11. Haxby, J. V. *et al.* A common, high-dimensional model of the representational space in

human ventral temporal cortex. *Neuron* **72**, 404–416 (2011).

12. Nastase, S. A. *et al.* Attention Selectively Reshapes the Geometry of Distributed Semantic

Representation. *Cereb. Cortex* **27**, 4277–4291 (2017).

13. Chen, P.-H. (cameron) *et al.* A Reduced-Dimension fMRI Shared Response Model. in

Advances in Neural Information Processing Systems 28 (eds. Cortes, C., Lawrence, N. D.,

Lee, D. D., Sugiyama, M. & Garnett, R.) 460–468 (Curran Associates, Inc., 2015).

14. Haxby, J. V., Guntupalli, J. S., Nastase, S. A. & Feilong, M. Hyperalignment: Modeling

shared information encoded in idiosyncratic cortical topographies. *Elife* **9**, (2020).

15. Heusser, A. C., Ziman, K., Owen, L. L. W. & Manning, J. R. HyperTools: a Python toolbox

for gaining geometric insights into high-dimensional data. *J. Mach. Learn. Res.* **18**,

5589–5594 (2017).

16. Schönemann, P. H. A generalized solution of the orthogonal procrustes problem.

Psychometrika **31**, 1–10 (1966).

17. van der Meer, M. A. A., Carey, A. A. & Tanaka, Y. Optimizing for generalization in the

decoding of internally generated activity in the hippocampus. *Hippocampus* **27**, 580–595

(2017).

18. Carey, A. A., Tanaka, Y. & van der Meer, M. A. A. Reward revaluation biases hippocampal

replay content away from the preferred outcome. *Nat. Neurosci.* **22**, 1450–1459 (2019).

19. Gupta, A. S., van der Meer, M. A. A., Touretzky, D. S. & David Redish, A. Hippocampal

Replay Is Not a Simple Function of Experience. *Neuron* vol. 65 695–705 (2010).

20. Gupta, A. S., van der Meer, M. A. A., Touretzky, D. S. & David Redish, A. Segmentation of

spatial experience by hippocampal theta sequences. *Nature Neuroscience* vol. 15

1032–1039 (2012).

21. Lee, J. S., Briguglio, J., Romani, S. & Lee, A. K. The statistical structure of the hippocampal

code for space as a function of time, context, and value. *bioRxiv* 615203 (2019)

doi:10.1101/615203.

22. Samsonovich, A. & McNaughton, B. L. Path integration and cognitive mapping in a

continuous attractor neural network model. *J. Neurosci.* **17**, 5900–5920 (1997).

23. Dragoi, G. & Tonegawa, S. Preplay of future place cell sequences by hippocampal cellular

assemblies. *Nature* **469**, 397–401 (2011).

24. Farooq, U., Sibille, J., Liu, K. & Dragoi, G. Strengthened Temporal Coordination within

Pre-existing Sequential Cell Assemblies Supports Trajectory Replay. *Neuron* **103**,

719–733.e7 (2019).

25. McKenzie, S. *et al.* Preexisting hippocampal network dynamics constrain optogenetically

induced place fields. *Neuron* **109**, 1040–1054.e7 (2021).

26. Rubin, A. *et al.* Revealing neural correlates of behavior without behavioral measurements.

Nat. Commun. **10**, 4745 (2019).

27. Simony, E. *et al.* Dynamic reconfiguration of the default mode network during narrative

comprehension. *Nat. Commun.* **7**, 12141 (2016).

28. Owen, L. L. W., Heusser, A. C. & Manning, J. R. A Gaussian process model of human

electrocorticographic data. doi:10.1101/121020.

29. Dordek, Y., Soudry, D., Meir, R. & Derdikman, D. Extracting grid cell characteristics from

place cell inputs using non-negative principal component analysis. *Elife* **5**, e10094 (2016).

30. Fyhn, M., Hafting, T., Treves, A., Moser, M.-B. & Moser, E. I. Hippocampal remapping and

grid realignment in entorhinal cortex. *Nature* **446**, 190–194 (2007).

31. Mark, S., Moran, R., Parr, T., Kennerley, S. & Behrens, T. Transferring structural knowledge

across cognitive maps in humans and models. *bioRxiv* 19 (2019).

32. Whittington, J. C. R., Muller, T. H., Mark, S., Chen, G. & Barry, C. The Tolman-Eichenbaum

Machine: Unifying space and relational memory through generalisation in the hippocampal

formation. *bioRxiv* (2019).

33. Chaudhuri, R., Gerçek, B., Pandey, B., Peyrache, A. & Fiete, I. The intrinsic attractor

manifold and population dynamics of a canonical cognitive circuit across waking and sleep.

Nat. Neurosci. **22**, 1512–1520 (2019).

34. Spiers, H. J., Hayman, R. M. A., Jovalekic, A., Marozzi, E. & Jeffery, K. J. Place field

repetition and purely local remapping in a multicompartment environment. *Cereb. Cortex*

25, 10–25 (2015).

35. Grieves, R. M., Jenkins, B. W., Harland, B. C., Wood, E. R. & Dudchenko, P. A. Place field

repetition and spatial learning in a multicompartment environment. *Hippocampus* **26**,

118–134 (2016).

36. Sanders, H., Wilson, M. A. & Gershman, S. J. Hippocampal remapping as hidden state

inference. *Elife* **9**, (2020).