The role of humans on Antillean land vertebrate extinctions: new insights from
Cuba

Johanset Orihuela[a,*], Lázaro W. Viñola[b], Osvaldo Jiménez Vázquez[c], Alexis Mychajliw[d]
Odlanyer Hernández de Lara[e], Logel Lorenzo[f], J. Angel Soto-Centeno[g,h,*]

[a] Department of Earth and Environment (Geosciences), Florida International University, Miami,
Florida 33199, USA

[b] Department of Biology, University of Florida, Gainesville, Florida 32611-7800

[c] Gabinete de Arqueología de La Habana, Oficina del Historiador de La Habana, Cuba

[d] Department of Antrhopology, University of Oklahoma, Norman 73069

[e] Cuba Arqueológica, Progressus Heritage & Community Foundation, University of Florida,
Gainesville, FL 32611

[f] Fundación Antonio Núñez Jiménez de la Naturaleza y el Hombre, Jardines Bellamar, carretera
las cuevas km 1½, Matanzas, Cuba

[g] Department of Biological Sciences, Rutgers University, Newark, NJ 07102

[h] Department of Mammalogy, American Museum of Natural History, New York, NY 10024

*Corresponding authors: Jorih003@fiu.edu (JO) and angelo.soto@rutgers.edu (JASC)

Declarations of interest: None

ABSTRACT

The Caribbean archipelago is a hotspot of biodiversity characterized by a high rate of extinction.

Recent studies have examined these losses, but the causes of the Antillean Late Quaternary

vertebrate extinctions, and especially the role of humans, are still unclear. Current results provide

support for climate-related and human-induced extinctions, but often downplaying other

complex bio-ecological factors that are difficult to model or to detect from the fossil and

archaeological record. Here, we disscuss Caribbean vertebrate extinctions and the potential role

of humans derived from new and existing fossil and archaeological data from Cuba. Our results

indicate that losses of Cuba's native fauna occurred in three waves: one during the late

Pleistocene and early Holocene, a second during the middle Holocene, and a third one during the

last 2 ka, coinciding with the arrival of agroceramists and the early Europeans. The coexistence

of now-extinct species with multiple cultural groups in Cuba for over 4 ka implies that Cuban

indigenous non-ceramic cultures exerted far less extinction pressures to native fauna than the

later agroceramists and Europeans that followed. This suggests a determinant value to increased

technological sophistication and demographics as most plausible effective extinction drivers.

KEYWORDS

Anthropogenic factors, biodiversity, Caribbean, chronology, Cuba, extinction, extirpation,

Holocene, island, vertebrates, West Indies

1. Introduction

The Caribbean is a hotspot of biodiversity with some of the highest rates of extinction recorded throughout the Holocene (MacPhee and Flemming, 1999; Dávalos and Turvey, 2012). Studies on insular vertebrate biodiversity and their prehistoric remains have shown that the faunas present on this archipelago today are only a remnant of the richer species assemblages of the past (e.g. Morgan and Woods, 1986; Pregill et al., 1988). Recent biodiversity extinctions in the Caribbean are often attributed to natural phenomena (e.g. global climate change) and human effects, such as hunting, habitat alterations, and the introduction of exotic species (Morgan and Woods, 1986; Stoetzel et al., 2016; Borroto-Páez and Mancina, 2017). Globally, most island extinctions seem to coincide or follow the arrival of humans, although the magnitude and overlap of these events seem to vary depending on island size, human cultures, and the characteristics of the fauna (Turvey, 2009a, b; Steadman et al., 2015; Turvey et al., 2017). Other studies have further suggested that prehistoric and historic island extinctions likely represent a single continuous event linked to human activities (Lyons et al., 2016). Thus, as new fossil data becomes available, pinpointing the cause of these losses becomes more critical to further our understanding of the magnitude and tempo of species loss and examine clues to disentangle the causes of recent extinction. These efforts require fine-scale chronological evidence associated with detailed knowledge of human colonization events and local climatic changes.

The role of humans on Caribbean vertebrate extinctions, especially mammals and birds, has been debated for nearly four decades (e.g. Olson, 1982; Morgan and Woods, 1986; Pregill et al., 1994). While recent data has shed light upon the arguments of the cause, effect, and timing (Díaz-Franco 2004, 2011; Jiménez et al., 2005; Jull et al., 2004; Steadman et al., 2005; Silva et al., 2007; Orihuela, 2010; Orihuela and Tejedor, 2012; Soto-Centeno and Steadman, 2015;

Cooke et al., 2017; Turvey et al., 2017; Borroto-Páez and Mancina, 2017), chronologic

resolution and data consensus are still much needed from islands with the highest concentrations

of biodiversity, such as Cuba.

Cuba, itself a sub-archipelago, is the largest of all Caribbean islands and has one of the

most diverse fossil and modern vertebrate records in the region (MacPhee and Fleming, 1999,

MacPhee et al 1999b). Its high levels of endemism, geological complexity (Iturralde and

MacPhee, 1999; Hedges, 2001), and history of human colonization (Napolitano et al., 2019)

make its consideration crucial to understanding how Amerindian populations and their

modifications to the environment shaped Caribbean vertebrate extinctions (Silva et al., 2007).

Nevertheless, Cuban zooarchaeological data has been rarely incorporated into current Antillean

biogeographical research and extinction syntheses. This has been exacerbated by the poor

circulation and accessibility of relevant Cuban research publications (e.g. Nuñez and Mayo,

1970; Pino and Castellanos, 1985; Díaz-Franco, 2004, 2011; Muñíz y Domínguez, 2014;

Jiménez, 2015; Jiménez and Arrazcaeta, 2015). Illustrative examples from the Cuban literature

that have been neglected include the well-documented cases of hutia "domestication" or captive

husbandry, interisland exchange, or the introduction of exotic species by Amerindians (e.g. Pose

et al., 1988; Jiménez and Milera, 2002; Díaz-Franco, 2004; Silva et al., 2007; Díaz-Franco and

Jiménez, 2008).

Moreover, the lack of direct radiocarbon dates on Cuban extinct fauna, and the dilatory

way some of these data became available, have limited detailed assessments. Prior studies have

compiled LADs for 10 Cuban mammals, but only three have direct dates (the eulipotyphlan

Nesophontes micrus, and the sloths *Parocnus browni* and *Megalocnus rodens*; MacPhee et al.,

1999, 2007; Turvey and Fritz, 2011), and associated date estimates from Cuban

zooarchaeological data has been scantly compiled.

Herein we provide novel details on Cuban vertebrate extinctions and highlight the

potential role of humans through the analysis of new and existing radiometric and zoo-

archeological data. To remedy the chronological knowledge gap and to discuss estimate times of

extinction, we generated 17 new direct AMS radiocarbon dates on both extinct and extant

terrestrial volant and non-flying vertebrates, seven of which represent new direct Last

Appearance Dates (LAD). Moreover, using detailed knowledge of the Cuban archaeological

record and literature, we critically compare and contrast the timing of faunal extinctions and

extirpation on the island and their relationship with the known Amerindian cultures that existed

during those intervals.

2. Materials and methods

2.1 Regional setting

Cuba (22.025˚ N, -78.997˚ W) is the largest island in the Caribbean (Fig. 1). It comprises

an archipelago of about 4,000 islands and occupies about 46% of the total land area of the

Caribbean (Woods and Sergile 2009). The island contains diverse habitats ranging from tropical

dry to mesic forests, semideserts, and mountains that reach elevations no greater than 1974 m;

conditions that contribute to the overall species diversity. We compiled fossil and archaeological

data from 17 unique localities across western and central Cuba (Fig. 1, Table 1).

2.2 AMS Radiocarbon (^{14}C) dated material

We present a dataset comprising seven direct last appearance ^{14}C dates (LADs), cross-

referenced with an additional 80 indirect chronologic taxa associations for extinct Cuban birds,

mammals, and a tortoise extracted from available published sources; including previously

unreported fauna from sites with known associated chronologies (Table 1 and Table S1). The

new directly dated material included here was collected from several fossil-rich localities and

complemented with material from 10 direct and 40 indirect records from the literature (Table 1,

Table S1, for study site details see Supplementary Text S1). Material excavated from these sites

is deposited at the Museo Nacional de Historia Natural (MNHNCu) and at the Gabinete de

Arqueología de la Oficina del Historiador (GAOHCu), both in La Habana, Cuba. Accelerated

Mass Spectrometry (AMS ^{14}C) dating was based on bone collagen and performed at Beta

Analytic (Miami, FL), International Chemical Analysis Inc. (ICA, Miami, FL), the UC Irvine

Keck Accelerator Mass Spectrometry facility (UCIAMS). Methods used at UCIAMS followed a

modified Longin collagen extraction protocol (Brown et al., 1988) on specimens with suitable

collagen yields and C: N ratios and were then ultrafiltered (> 30 kDa) to remove contaminants.

All radiocarbon dates were calibrated using OxCal v. 4.3 (Bronk Ramsey, 2009) and the

IntCal13 calibration curve (Reimer et al., 2013), as no marine samples were included. We report

the median calibrated age (Cal BP) and the 95.4% range. For additional details on AMS

methods, see www.radiocarbon.com and www.radiocdating.com.

2.3 Taxonomy

Systematic taxonomy for Cuban extinct and extant vertebrate species follows, for bats

Silva (1974, 1976) and Balseiro (2011), except in considering *Mormoops magna* a valid extinct

species. *Desmodus puntajudensis* is considered a synonym of *Desmodus rotundus* following

Orihuela (2011). For the rest of the land mammals we follow Silva et al. (2007) and González et

al. (2012) but considering the validity of two species of *Nesophontes* shrews (Condis et al., 2005; Rzebik-Kowalska and Woloszyn, 2012), and adapting the new arrangement of including the black-tailed hutia (*M. melanurus*) within the genus *Mesocapromys* (Upham and Borroto-Páez 2017). For birds, we follow Orihuela (2019) and for squamata Arana (2019).

Abbreviations: AD, anno domini or common era. BC, before Christ or before the common era. BP, before the present (datum AD 1950). Cal., calibrated age. Cal BP, median radiocarbon calibrated age before the present. EED, effective extinction date. eHOL, early Holocene. Ka, kilo-years. LAD, last appearance date. LGM, Last Glacial Maximum. lHOL, late Holocene. PHT, Pleistocene-Holocene transition. Rdcy, radiocarbon years before the present.

2.4 Chronologic intervals and last appearance dates (LAD)

All LADs fit into the five chronologic intervals defined by the International Commission on Stratigraphy (ICS, 2019). For ease of communication and consistency with previous literature, we used chronologic intervals names in Soto-Centeno and Steadman (2015). The first interval is the Last Glacial Maximum or Late Pleistocene (ca. 21–11.7 ka), which culminated with the Pleistocene-Holocene transition (PHT, ca. 11.7 ka; Curtis et al., 2001). The following intervals are the early Holocene (eHOL, ca. 11.7–8.2 ka), the middle Holocene (mHOL, ca. 8.2–4.2 ka), and the late Holocene (lHOL, < 4.2 ka). We also separately demarcate the Amerindian interval, defined as the beginning of the transition between the middle to late Holocene (mHOL to lHOL, < ~6 ka), and spans the entire late Holocene. A final and fifth stage is the post-Columbian interval beginning in the last 500 BP (AD 1492–1500 to the present), interchangeably used in the literature as the European or Historic interval (e.g. Morgan and Woods, 1986; Soto-Centeno and Steadman, 2015; Cooke et al., 2017).

2.4.1 A rationale for a subdivision of the Amerindian interval

Given that humans only arrived in Cuba after the middle Holocene (mHOL, < ~6 ka; Cooper, 2010; Napolitano et al., 2019), we subdivide the Amerindian interval into two subintervals to better discuss LADs in the context of different cultural practices and degrees of environmental alteration. This helps in providing a better resolution of the timing or patterns of extinction and extirpation. These two intervals are the early Amerindian (~6–1.5 ka BP) and the late Amerindian (1500–500 BP). Our use of this specific scheme focuses attention on the cultural groups that could have importantly affected the natural environments of the island at a given chronological interval and provides a better resolution to the timing of extinction or extirpation. An expanded rationale for these subintervals, climate-related information, and a description of the sites and deposits are provided in the Supplementaty Text S2.

3. Results

The direct dates we report represent 27% (16 of 59) of the known extinct land vertebrate fauna, including four extant but critically endangered mammal species. Within these 59 vertebrates are 38 birds and one reptile, which together comprise the record of extinct and extirpated land vertebrate species so far registered from the Quaternary of Cuba. We provide direct and indirect LADs for 95% (20 of 21) of the known Cuban quaternary extinct land mammal fauna, including bats. This overall mammalian fauna is composed of two echimyid rodents, four capromyid rodents, four sloths, three eulipotyphlans, one monkey and seven bats (Fig. 2–4, Table 1). Moreover, we provide new direct radiocarbon dates on Amerindian human remains that contribute to a better understanding of the contexts of association for several species

that lack direct dates (Fig. 2, Table S1), and provide further support to formerly estimated LADs

(e.g. for the extirpated bat *Mormoops megalophylla* from Cueva del Gato Jíbaro in Orihuela and

Tejedor, 2012; see Table 1, Table S1).

Of the chronologically analyzed extinct and extirpated vertebrates, 44.4% (12 of 27)

dated well into the late Amerindian (1500–500 BP; Fig. 2, 4, Table 1, Table S1–S2). Only two

(7.4%) of the 27 currently registered extinct species have LADs within the Late Pleistocene (ca.

21–11.7 ka), 11.1% (3 of 27) occur during the early Holocene (eHOL, ca. 11.7–8.2 ka), 33.3% (9

of 27) occur during the middle Holocene (mHOL, ca. 8.2–4.2 ka), and 48.1% (13 of 27) occur

during the Late Holocene (lHOL, ca. > 4.2 ka; Orihuela, 2019). When examining these 13 late

Holocene species based on Amerindian subintervals, only three species (*Macrocapromys*

acevedo, *Parocnus browni,* and *Megalocnus rodens*) had LADs on the early Amerindian

subinterval (ca. 6–1.5ka), whereas 10 dated within the late Amerindian (ca. 1.5ka–500 BP). At

least four of these 13 species could have had EEDs in the post-Columbian interval (ca. 500 BP–

present): *Boromys torrei*, *B. offella*, *Geocapromys columbianus* and *Nesotrochis picapicensis*

(Fig. 2, 4, Table 1; Tables S1–S3). So far, only 5.4% (2 of 38) currently known extinct and

extirpated Cuban birds, have direct radiometric LADs; one at the LGM and one in the late

Amerindian (Orihuela, 2019). There are no known extinct amphibians described so far, and only

one reptile (*Chelonoidis cubensis*).

For mammals, we provide direct dates from specimens of the extinct eulipotiphlan

insectivores *Nesophontes major* and *Nesophontes micrus*, the extinct fruit bats *Artibeus anthonyi*

and *Phyllops vetus*, the likely extinct *Antrozous koopmani*, plus the pygmy hutia *Mesocarpomys*

nanus and *Solenodon cubanus*, the extinct rodents *Boromys torrei*, *B. offella*, *Macrocapromys*

acevedo, and *Mesocapromys kraglievichi*. We also provide associated age estimates for the

extirpated bats *Mormoops megalophylla* and *Desmodus rotundus*, and the hutia *Geocapromys*

columbianus based on new indirect radiocarbon dates (Table S1). All their LADs are also well

within the late Amerindian subinterval, supporting formerly presumed extinction estimates

within the Late Holocene (Jiménez et al., 2005; Orihuela, 2010; Orihuela and Tejedor, 2012;

Orihuela et al., 2020). New associated LADs are also provided for the sloths *Acratocnus*

antillensis and *Neocnus gliriformis,* and the extinct tortoise *Chelonoidis cubensis* based on

associated date estimates (Table 1, Table S1).

Two extinct bats (*Cubanycteris silvai* and *Pteronotus pristinus*) do not have direct or

indirect radiocarbon dates available to establish LADs (Silva 1974, 1979; Mancina and García-

Rivera, 2005). Their extinction interval is assumed to be late Pleistocene based on the

interpretation of the authors (Suárez and Díaz-Franco, 2003), but likely these may be constrained

within the early Holocene (eHOL, ca. 11.7–8.2 ka; e.g. see data in Condis, 2005; Fiol, 2015).

More data is necessary to evaluate the extinction of the monkey *Paraloutta varonai* and the

eulipotiphlan insectivore *Solenodon arredondoi* (Salgado et al., 1990; MacPhee et al., 1996a;

Morgan and Ottenwalder, 1993; Jiménez, 2015), but an early Holocene extinction is also

plausible for these taxa.

Generally, eulipotyphlan insectivores and native rodents yielded LADs that are

borderline to the post-Columbian interval, which seem to corroborate former assumptions of

their extinction during the historic interval (Guarch, 1982, 1984; Pino, 2012; Fig. 2, Table 1,

Table S1–S2. Direct and indirect dates for extant but critically endangered species, such as

Mesocapromys nanus, *Solenodon cubanus*, and the bats *Antrozous koopmani* and *Natalus*

primus, suggest that these species had a much wider distribution throughout western Cuba up to

the time of European colonization. In this sense, these data could be capturing the initial range

collapse or earlier stages of extinction processes.

4. Discussion

Caribbean extinctions have had a significant role in shaping the living vertebrate diversity

in this archipelago. Detailed radiochronologies estimating last appearance dates (LAD) of large

fossil assemblages help us analyze changes in community composition and the timing of

community turnover in association with the recent historical events that drove them. Studies

debating the cause of extinction of Cuban vertebrates, typically do so despite lacking appropriate

resolution of chronological data. Herein, we provided a broadscale overview of vertebrate loss in

Cuba with chronologic LAD coverage of 56% of all extinct taxa documented for the island. For

the few taxa, we analyzed that are still extant, our results reveal a better context of the history of

these species. For extinct taxa specifically, we detect a pattern of associated loss representing

about 48% of Cuban species primarily associated with an interval of active human presence

(lHOL, ca. > 4.2 ka). Our results reveal that at least five species disappeared between 11–8 ka, a

time interval before the arrival of humans on Cuba. Extinctions increases to 7–8 species during

the middle Holocene, and most of these taxa likely disappeared following human arrival after ~6

ka. Most of the extinct species in Table 1 survived the Pleistocene-Holocene transition (PHT, ca.

11.7 ka) and several thousand years of human inhabitation in Cuba (Fig. 2, Table S3).

Up to 90% (22 of 27) of the vertebrate fauna have direct and indirect LADs younger than

eHOL (ca. 8.2 ky; Fig. 2, Table 1, Table S2–S3). The majority of these (66.6%) have LADs well

into the late Amerindian and post-Columbian intervals, coinciding to a time near or after the

arrival of the agroceramist culture groups in Cuba (ca. ~ 1500 BP). Climate conditions during

this time interval were largely similar to those of today (Soto-Centeno and Steadman 2015).

These data support the hypothesis that lost species persisted for thousands of years after the onset

of warming climate conditions of the Holocene. Furthermore, our results show persistence for

over 4 ky of Amerindian presence in Cuba, which is congruent with estimates from previous

studies on other islands and also supports previous late Holocene extinction estimates in Cuba

(MacPhee et al. 1999; Díaz-Franco 2004, 2011; Jiménez et al., 2005; Orihuela, 2010; Orihuela

and Tejedor, 2012; Soto-Centeno and Steadman, 2015; Borroto-Páez and Mancina, 2017).

The earliest Amerindians to arrive in Cuba were the so-called "archaic" pre-Arawak that

reached the island during the mid-Holocene (Cooper, 2007, 2010; Cooper and Thomas, 2011;

Roksandic et al., 2015; Ulloa and Valcárcel, 2016, 2019). These aboriginal colonists were

preceramic hunter-gatherer groups with estimated low demographics and they exploited a wide

range of ecosystems within the archipelago (Tabío and Rey, 1979; Dacal, 1980; Dacal and

Rivero, 1986; Guarch, 1990; Guarch et al., 1995; Godo, 1997; González Herrera, 2008, 2018;

Ulloa and Valcárcel, 2019).

Early Amerindian culture middens show a preference for coastal and riverine resources,

predominantly mollusks and fish (Pino, 1978, 2012). Rodents were secondarily exploited,

especially the larger hutias such as *Capromys pilorides* and *Geocapromys columbianus*, and the

medium to smaller sized *Mysateles prehensilis*, *Mesocapromys melanurus*, *Boromys* spp.

(Kozlowski, 1974; Pino, 1978, 2012; Córdova-Medina et al., 1997; Reyes, 1997; Jiménez and

Arrazcaeta, 2015). *Solenodon* and *Nesophontes* are documented as likely occasional supplements

of the aboriginal diet (Martínez Arango, 1968; Pino, 1978, 2012; Martínez et al., 1993).

Although aboriginals were dietarily, ecologically and culturally diverse, it is unlikely that

they had a major impact on Cuban fauna due to their incipient technological sophistication and

demographics. Many of the larger mammals and the land tortoise seem to have disappeared

during the mHOL, near the arrival time of the first humans to the island. But even with the

current state of knowledge, it is still difficult to discern how early aboriginal colonists could have

perturbed these vertebrate communities. For cavernicolous bats, for example, frequent human

visits to caves likely could have affected their colonies (Silva, 1979; Mancina et al., 2007). The

level of disturbance on bats could have been further exacerbated if cave fires are considered

because smoke can lead to high mortality or complete extirpation. Nonetheless, direct evidence

for these mass deaths or disturbance is lacking from the current record in Cuba (Orihuela and

Tejedor, 2012).

Sloth remains are rare from midden deposits attributed to early Amerindian groups.

While the context and age of the evidence are still debated, sloths have been reported from at

least a dozen archaeological sites (Pino y Castellanos, 1985; Jiménez and Arredondo, 2011:208;

Díaz-Franco, 2004, 2011; Pino, 2012; Arredondo and Villavicencio, 2011). Taphonomic

evidence of human predation in the form of tool and cut marks has been scant and remains

questionable (Arredondo and Villavicencio, 2011; Jiménez and Arredondo, 2011; Orihuela et al.,

2016). LADs for *Megalocnus rodens* and *Parocnus browni* and several of these other species

(Table 1, Table S1) suggest that they coexisted in time but there is yet no conclusive evidence

that sloths were predated by early Amerindians (Jull et al. 2004; MacPhee et al. 2007).

Other groups of hunter-fisher-gatherer Amerindians are also present in the Cuban

archaeological record with radiocarbon dates between 2500 BC and AD ~1500 (Guarch, 1990;

Guarch et al., 1995; Cooper, 2007, 2010; Chinique and Rodriguez, 2012; Roksandic et al., 2015;

Chinique et al., 2015; Chinique et al., 2019; see Supplementary Text S2). At least in Cuba, these Amerindians had a varied toolkit and likely practiced incipient agriculture (i.e. use of cultigens and wild varieties; see Chinique et al. 2015, 2016, 2019) with comparable adaptations but with slightly larger demographics given that some groups coexisted with the Taíno and Europeans that arrived later around 1500–500 BP, respectively (Fernández Oviedo, 1535; Las Casas, 1875; Guarch,1978, 1982; Dacal, 1980; Rouse, 1992; Torres, 2006). The hunter-fisher-gatherer midden remains, and direct isotope analyses indicate that they had variable diets acquired in diverse habitats. Some groups were dependent on land mammals, in addition to marine/estuarine organisms (Chinique and Rodriguez, 2012; Chinique et al., 2015, 2016, 2019). Rodents, particularly *Capromys pilorides*, *G. columbianus*, and the smaller spiny rats *Boromys* spp., were an important component to these culture's diet (Guarch, 1982; Córdova-Medina, 1993; Reyes, 1997; Díaz-Franco, 2004; Jiménez and Arredondo, 2011; Pino, 2012; Colten and Worthington, 2018). It is likely these rodents survived past the post-Columbian interval (> AD 1500) even though most have LADs within the late Amerindian subinterval (Fig. 2).

The first agroceramists collectively called Taíno arrived in Cuba ~1500 BP (AD 800–900; Guarch, 1978; Rouse, 1992; Guarch et al., 1995; Valcárcel, 2002, 2008; Torres, 2006). Radiocarbon dates between the "archaic" pre-Arawak groups and Taíno groups overlap, supporting coexistence between different cultural groups at least in some areas of the archipelago as observed by early European chronicles (Fernández de Oviedo, 1535; Las Casas, 1875; Dacal, 1980; Rouse, 1992; Godo, 1997; Cooper, 2010; Cooper and Thomas, 2011; Valcárcel, 2012; Roksandic et al., 2015; Chinique et al., 2016, 2019; Orihuela et al., 2017).

The Taíno hunted, fished, gathered, practiced land clearing for agriculture and lived in a wide range of sites that they could have fully exploited (Guarch, 1978; Rouse, 1992; Torres,

2006). They had a strong impact on the environment as evidenced by the introduction of

domestic dogs (Arredondo, 1981; Wing and Scudder, 1983; Jiménez y Fernández-Milera, 2002;

Newsom y Wing, 2004; Borroto-Páez, 2011; Hofman et al., 2011; Grouard et al., 2013; Laffoon

et al., 2013). Chronicler accounts on *Canis lupus familiaris* indicate that it was already widely

dispersed within the island by Cal 1450–1530 AD when more domestic dogs were further

introduced by the Spanish colonists (Rodríguez-Durán y Santiago, 2014). Some taphonomic

evidence supports canid predation on the native vertebrate fauna, including *Solenodon* and large

rodents, and also feeding on human middens (Jiménez and Arrendondo, 2011; Orihuela et al.,

2016).

Perhaps the most important interval for human-wildlife interactions in Cuba began during

Taino's coexistence with Europeans (< 1492 AD; Marrero, 1972; Morgan and Woods, 1986;

MacPhee et al., 1999a/b; Valcárcel, 2012). These two waves of human colonization combined

(i.e. agroceramists followed by Europeans) could have been a major source of continuous

environmental perturbation through direct habitat destruction and exotic species introductions,

increasing the rate of loss of native fauna. Europeans introduced many exotic species between

1511–1890 including house mice (*Mus musculus*), rats (*Rattus* spp.), domestic dogs (*C. lupus*

familiaris), pigs (*Sus scrofa*), rabbits, New World monkeys, mongoose (*Herpestes*

auropunctatus), and others (Borroto-Páez, 2011). These exotics likely intensified pressure on

native vertebrate populations like *S. cubanus*, *Boromys* spp., and *Nesophontes* spp. (Morgan and

Ottenwalder, 1993; MacPhee et al., 1999; Silva et al., 2007; Jiménez and Arredondo, 2011).

Habitat loss, the introduction of exotic species, direct hunting, and human-induced

climate change are some of the most important indirect anthropogenic drivers of modern

extinctions in the Caribbean (IUCN, 2019). The concomitance of human colonization and timing

of extinctions in the Greater Antilles has been widely considered as evidence for anthropogenic extinctions (Steadman et al., 1984; Pregill et al., 1988; 1994; Turvey et al., 2007; Soto-Centeno and Steadman, 2015). However, we hypothesize that even though there were the land vertebrate extinctions during the early Amerindian subinterval, it was the combination of the latter agroceramist and European colonization which seem to coincide with most species' LADs (i.e. disappearances) during the last 2000 years. Evidence that all the species that disappeared during the late Amerindian and post-Columbian intervals had survived over 4000 years of previous human inhabitation on the island, as in other areas of the Greater Antilles (e.g. Soto-Centeno and Steadman, 2015; Stoetzel et al., 2016, Soto-Centeno et al., 2017), thus validates the importance of the combination of threats brought by new colonizers with increased demographics, better environmental knowledge, and the technology to exploit it.

5. Conclusion

We support the hypothesis that extinctions in Cuba occurred in multiple episodes during the Holocene. The majority of these reflected in direct and indirect LADs that coincide with arrival of agroceramist groups and other later on, in the subinterval we have defined as the late Amerindian (> 1.5 ka). Our data suggest that at least in Cuba, nearly half of the extinct land mammals (i.e. 44 %), and at least some avian extinctions, occurred during the last 1500 years. Making the late Amerindian and post-Columbian extinction episodes the most significant in Cuba's late Quaternary. Historically, the first two centuries of colonialism in Cuba were not considered as having a strong impact on the fauna because large deforestation and environmental modifications began after the XVIII century.

Although a seemingly important extinction event is observed from LADs within the early

Amerindian intervals (i.e. 33% of the 27 analyzed taxa have LADs between 6–1.5 ka), there is no

evidence for a megafaunal overkill or blitzkrieg extinction in Cuba. It is possible that this

apparent extinction episode is an artifact of the lack of data because taxa with LADs within this

interval could have persisted later in time. However, it nevertheless suggests that not all

extinctions were driven by humans because some species did go extinct before human arrival to

the archipelago (e.g. the bats *Pteronotus pristinus*, *Cubanycteris silvai*, *Phyllops silvai*, and the

monkey *Paralouatta varonai*). Among the causes during the middle Holocene is a probable

cascade effect combining climatic and anthropogenic factors that resulted in the asynchronous

extinction of native land vertebrates.

The LADs discussed herein indicate that nearly half of the species listed survived over

4000 years of human coexistence in Cuba. At least the earliest human communities likely

affected some groups (e.g. large terrestrial birds and sloths), either directly or indirectly, but

caused little if any major environmental alterations. Half of the species more frequently exploited

and consumed survived. Such condition dramatically changed after the arrival of the first

agroceramist (Taíno) groups and where amplified later by the European colonists. It seems

apparent that agroceramist and European colonizers could have played a decisive role in the

direct or indirect extinction, extirpation or endangerment of Cuba's native fauna. The effect of

climate-related factors cannot be decoupled from anthropogenic ones precisely because they are

concomitant from the onset of any human colonization, and do not disappear thereafter. Before

human arrival, climate and bio-ecological-related factors likely played the most important role in

determining geographical distribution, local or final extinction, and this explains extinctions

during the late Pleistocene to Holocene transition. This changed significantly at the arrival of

Amerindians and later European colonists. Then, climate-related variables likely became

secondary or less of a driver of changes in distribution or extinction in comparison to the more

drastic human-related threats that are still present.

The consideration of the complex combination of climate and anthropogenic factors

combined can provide a powerful and more realistic explanation to the extinction episodes

observed after the Amerindian intervals. While some species may have survived previous

changes in climate by shifting their distributions, they finally came to extirpation or extinction

due to other causes, many of which remain unexplored (Supplementary Text S2). The cause for

Cuban vertebrate mid to late Holocene extinctions cannot be solely attributed to either climate or

human-related drivers, either direct or indirect impacts, but more likely resulted from a

combination of complex factors, including variables that we cannot model for, or that are not yet

detected in the fossil record. The understanding of the timing and the complexity of the causes

leading to the extirpation or global extinction of native fauna is crucial to develop future

protection and conservation programs, which must consider human demographic and

technological growth, deforestation, climate change, and the individual species responses to

these pressures.

Acknowledgments

JO thanks Sigma Xi for providing funding for some of the analyses (grant id:

G20141015720649). To Leonel Perez Orozco, Candido Santana, and Ricardo Viera for field

assistance, discussions and logistics. Ercilio Vento, Stephen Díaz Franco and Abel Hernández

for providing specimens used in this study. The *Consejo Nacional de Patrimonio Cultural,*

Ministerio de Cultura de Cuba for excavation/exportation permits, and Jorge Garcell and Jaime Triana for all their help in securing them. To Adrian Tejedor, Herman Benitez, Nick Czaplewski, Gary Morgan, Jorge Ulloa, Roberto Valcárcel, Mathew Peros, Jago Cooper, Yadira Chinique, Mirjana Roskandic, Bronislaw Woloszyn and Ross MacPhee for literature and guidance. JASC thanks R.D. Barrilito for support. Radiocarbon work was partly funded by a Rutgers University Research Council Award to JASC. In figure 1, images for rat, sloth, solenodon, and vespertilionid bat obtained from phylopic.org and used with permission under a creative commons license; other images produced by JASC.

References

Acevedo González, M., Arredondo, O., and González, N. 1975. *La Cueva del Túnel*. Editorial Pueblo y Educación, La Habana.

Acevedo González, M., and Arredondo, O. 1982. Paleozoografía y geología del cuaternario de Cuba: Características y distribución geográfica de los depósitos con restos de vertebrados. *IX Jornada Científica del Instituto de Geología y Paleontológica de la Academia de Ciencias de Cuba*: 54–70, La Habana, Cuba.

Alonso Alonso, E. M. 1989. Un campamento aborigen en Sierra del Rosario. *Revista Cubana de Ciencias Sociales, Academia de Ciencias de Cuba*, (21):38-58.

Alonso Alonso, E., G. Izquierdo Díaz, U. M. González Herrera, G. Hernández Ramírez, R. Valcárcel Rojas, M. Pino Rodríguez, and E. Blanco Castillo 2015. *Las Comunidades Aborígenes en la Historia de Cuba*. Fundación Fernando Ortiz, Selvi Artes Gráficas, Valencia.

Anderson-Córdova, K. F. 1990. *Hispaniola and Puerto Rico. Indian acculturation and heterogeneity, 1492 - 1550* (PhD Diss.). Yale University, Yale.

Arana Pedroso, E. 2019. *Systematics of Quaternary Squamata from Cuba*. (Master's thesis). University of Sao Paulo, Brazil.

Arredondo, O. 1981. Nuevo genero y especie de mamífero (Carnivora: Canidae) del Holoceno de
Cuba. *Poeyana*, 218:1-28.

Arredondo, C. and Villavicencio, R. 2004. Contribución taxonómica a la interpretación del
depósito arqueológico Solapa del Megalocnus en el noroeste de Villa Clara, Cuba. *Revista*
Biología, 18(2): 20-30.

Balseiro, F. 2011. Los murciélagos extintos p. 170-177. In Borroto-Páez, R., and Mancina C.A.
*(*eds.*)*, *Mamíferos en Cuba*. UPC Print Vaasa, Finland.

Borroto-Páez, R. 2011. Los mamíferos invasores o introducidos, p. 220–241. In Borroto-Páez, R.
and Mancina, C.A. (eds.), *Mamíferos en Cuba*. UPC Print, Vaasa, Finland.

Borroto-Páez, R. and Mancina C.A. 2017. Biodiversity and conservation of Cuban mammals:
past, present, and invasive species. *Journal of Mammalogy*, 98(4):964–985.

Bronk Ramsey, C. 2009. Bayesian analysis of radiocarbon dates. *Radiocarbon*, 51(1), 337-360

Brown, T.A., Nelson, D.E., Vogel, J.S., and Southon, J.R. 1988. Improved collagen extraction by
modified Longin method. *Radiocarbon*, 30(2):171-177.

Burney, D.A., Pigott Burney L. and MacPhee, R.D.E. 1994. Holocene charcoal stratigraphy from
Laguna Tortuguero, Puerto Rico, and the timing of human arrival on the island. *Journal of*
Archaeological Science, 21:273–281.

Casas, B. de las, 1875. *Historia de Las Indias.* Tomo I, Imprenta de Miguel Ginesta, calle de
Campomanes, no. 8, Madrid.

Chinique de Armas, Y., and Rodríguez Suárez, R. 2012. Cambios en las actividades
subsistenciales de los aborígenes del sitio arqueológico Canímar Abajo, Matanzas, Cuba.
Cuba Arqueológica 5 30-48.

Chinique de Armas, Y., Buhay, W. M., Rodríguez Suárez, R., Bestel, S., Smith, D., Mowat, S.
D., and Roksandic, M. 2015. Starch analysis and isotopic evidence of consumption of
cultigens among Fisher-gatherers in Cuba: the archaeological site of Canímar Abajo,
Matanzas. *Journal of Archaeological Science* 58: 121-132.

Chinique de Armas, Y., Roksandic, M., Rodríguez Suárez, R., Smith, D. G., and Buhay, W. M.
2016. Isotopic evidence of variations in subsistence strategies and food consumption
patterns among "Fisher-gatherer" populations of Western Cuba. Chapter 8 pp: 125-146 *En*
Ivan Roksandic (Ed.) *Cuban Archaeology in the Caribbean*. University Press of Florida,
Gainesville: DOI:10.5744/florida/9781683400028.003.0009

Chinique de Armas, Y., Rodriguez Suárez, R., Buhay, W. M. and Roksandic, M. 2019. Chapter
8: Subsistence strategies and food consumption patter of Archaic Age populations from
Cuba: From traditional perspectives to current analytical results *in* Hofman, C.L. and

Antczak, A.T. (eds): *Early Settlers of the Insular Caribbean. Dearchaizing the Archaic.* Sidestone Press (Leiden), pp. 107-118.

Colten, R., Newman, E. T., and Worthington, B., 2009. Pre-Ceramic Faunal Exploitation at the Las Obas Site, Cuba. *The Bulletin of the Peabody Museum of Natural History* 50(1):75-84.

Colten, R. and Worthington, B. 2014. Faunal Remains from the Archaic and Archaic with Ceramic Site of Vega del Palmar, Cuba. *Journal of Caribbean Archaeology* 14:23-49.

Colten, R. and Worthington, B. 2018. Museum Collections and Archaic Era Vertebrate Faunal Remains from Cuba. *Environmental Archaeology*, DOI: 10.1080/14614103.2018.1500157.

Cooke, S.B., Dávalos, L.M., Mychajliw, A.M., Turvey, S.T., and Upham, N.S. 2017. Anthropogenic Extinction Dominates Holocene Declines of West Indian Mammals. *Annual Review of Ecology, Evolution, and Systematics*, 48, 301–327.

Cooper, J. 2007. Registro nacional de arqueología aborígen de Cuba: Una discusión de métodos y prácticas. *El Caribe Arqueológico*, 10: 132-150.

Cooper, J. 2010. Pre-Columbian archaeology of Cuba: A study of site distribution patterns and radiocarbon chronologies. Chapter 4: pp 81-107 *in* Scott M. Fitzpatrick and Ann H. Ross (Eds.) *Island Shores, Distant Pasts: Archaeological and Biological Approaches to the Pre-Columbian Settlement of the Caribbean.* University Press of Florida, Gainesville.

Cooper, J. and K. D. Thomas. 2011. Constructing Caribbean chronologies: comparative radiocarbon dating of shell and wood artifacts from pre-Columbian sites in Cuba. *Archaeometry* 54 (2): 401-425. DOI: 10.1111/j.1475-4754.2011. 00622.x

Condis Fernández, M.M., Jiménez Vázquez, O. and Arredondo, C. 2005. Revisión taxonómica del género Nesophontes (Insectivora: Nesophontidae) en Cuba: análisis de los caracteres diagnósticos p. 95-100. In Alcover, J. A. & Bover, P. (eds.), Proceedings of the International Symposium *"Insular Vertebrate Evolution: the Palaeontological Approach".* *Monografies de la Societat d'Història Natural de les Balears*, 12.

Condis Fernández, M.M. 2005. Inferencias paleoecológicas sobre especies de la mastofauna cuaternaria cubana, conservadas en el depósito superficial de la Caverna GEDA, Pinar del Rio, Cuba. (PhD diss.), Universidad de Pinar del Río, Cuba.

Córdova-Medina, A. P. 1993. Caracterización de la alimentación en una comunidad neolítica. Aspectos zoológicos y paleontológicos. *Carta Informativa 23*, Época III. Academia de Ciencias de Cuba, La Habana.

Córdova-Medina, A., Crespo Díaz, R., and Jiménez Vázquez, O. 1997. Importancia arqueológica y zoológica del sitio Solapa de Sílex. *El Caribe Arqueológico*, 2: 78-83.

Curtis, J.H., Brenner, M. and Hodell, D.A. 2001. Climate change in the Circum-Caribbean (Late Pleistocene to Present) and implication for regional biogeography, p. 35-54. In Woods, C.A. & Sergile, F. (eds.), *Biogeography of the West Indies* (Second Edition). CRC Press, Boca Raton, Florida.

Dacal-Maure, R. 1980. De los Siboneyes del Padre de las Casas a los Siboneyes de 1966. *Revista Universidad de la Habana 211*: 6-4pp.

Dacal-Maure, R., and M. Rivero de la Calle. 1986. *Arqueología aborigen de Cuba.* Gente Nueva, Habana, Cuba 174pp.

Dávalos, L. M., and Russel, A. L. 2012. Deglaciation explains bat extinction in the Caribbean. *Ecology and Evolution*, 2 (12): 3045-3051.

Dávalos, L.M. and Turvey, S.T. 2012. West Indian mammals: the old, the new, and the recently extinct. *Bones, clones and biomes: the history and geography of recent neotropical mammals* (ed. by B.D. Patterson and L.P. Costa), pp. 157–202. University of Chicago Press, Chicago, IL.

Del Risco Rodríguez, E. 1989. Mapa a 1:2, 000,000 vegetación original. Flora y Vegetación 3. In *Nuevo Atlas Nacional de Cuba*. Instituto de Planificación Física, La Habana, Cuba.

Díaz-Franco, S. 2004. Análisis de la extinción de algunos mamíferos Cubanos, sobre la base de evidencias paleonotológicas y arqueológicas. *Revista Biologica 18(2): 147-154pp.*

Díaz-Franco, S. and Jiménez Vázquez, O. 2008. *Geocapromys brownii* (Rodentia: Capromyidae: Capromyinae) en Cuba. *Solenodon*, 7: 41-47.

Díaz-Franco, S. 2011. Los perezosos extintos de Cuba: algunas reflexiones a propósito de los últimos fechados radiocarbónicos. *El Caribe Arqueológico*, 12: 15-29.

Fernández de Oviedo y Valdés, G. 1535 [1851]. *Historia general y natural de Las Indias, islas y tierra-firme del mar océano. Primera parte*, Imprenta de la Real Academia de la Historia, Madrid, 632 pp.

Flemming, C., and R. D. E. MacPhee. 1999. Redetermination of the holotype of *Isolobodon portoricensis* (Rodentia: Capromyidae), with notes on recent mammalian extinctions in Puerto Rico. *American Museum Novitates*, 3278: 1-11.

Fiol González, S. (2015). La fauna de mamíferos fósiles del depósito paleontológico "El Abrón" (Nivel IX) Pinar del Río, Cuba. (Master's Thesis), Universidad de La Habana, Facultada de Biología, La Habana, Cuba.

Garrido, O.H., and Kirkconnell A. 2000. *Field Guide to the Birds of Cuba*. Cornell University Press, Ithaca.

Giovas, C. M. 2018. The beasts at large-Perennial questions and new paradigms for Caribbean translocation research. Part II: Mammalian introductions in cultural context. Environmental Archaeology, DOI: 10.1080/14614103.2018.1470211.

Godo, P. P. 1997. El problema del protoagrícola de Cuba. Discusión y perspectivas. *El Caribe Arqueológico*, 2: 19-30.

González Alonso, H., Rodríguez, L., Rodríguez, A., Mancina, C.A., and Ramos, I. 2012 (eds.). *Libro Rojo de los Vertebrados de Cuba*. Editorial Academia, La Habana.

González Herrera, U. M. 2008. Ciboneyes, guanahatabeyes y cronistas. Discusión en torno a problemas de reconstrucción etnohistórica en Cuba. *El Caribe Arqueológico*, 11:98-105.

González Herrera, U. M. 2018. Expediente del sitio arqueológico Playa El Mango, municipio Río Cauto, provincia Granma. Departamento de Arqueología, Instituto Cubano de Antropología, CITMA, La Habana (inedited).

Grouard S., Perdikaris S. and Debue K. 2013. Dog burials associated with Human burials in the West Indies during the early pre-Columbian Ceramic Age (500 BC-600 AD). *Anthropo-zoologica*, 48 (2): 447-465. http://dx.doi.org/10.5252/az2013n2a17.

Guarch, J. M. 1978. *El Taíno de Cuba*. Academia de Ciencias de Cuba, La Habana.

Guarch, J. M. 1982. Sobre la extinción tardía de *Geocapromys* y *Heteropsomys* en Cuba. *Carta Informativa*, Época II, 34: 1-2.

Guarch, J. M. 1984. Evidencias de la existencia postcolombina de *Geocapromys* y *Heteropsomys* (Mammalia: Rodentia) en Cuba. *Miscelánea Zoológica*, 18: 1.

Guarch, J. M. 1990. *Estructura para las comunidades aborígenes de Cuba*. Editorial Academia, La Habana.

Guarch, J. M., Domínguez, J. M., Rives, A., Calvera, J., Pino, M., Castellanos, N., Rodríguez Arce, C., Jardines, J., Pedroso, R., Sampedro, R. and Izquierdo, G. 1995. *Historia Aborigen de Cuba, según datos arqueológicos*. CD-ROM, *Taino: Arqueología de Cuba*. Centro de Antropología y CEDISAC, Colima.

Hedges, B. 2001. Biogeography of the West Indies: an overview. Pp. 15-33 in Woods, C. A. and F. E. Sergile (eds). *Biogeography of the West Indies: Patterns and Perspectives*. CRC Press, Boca Raton.

International Commission on Stratigraphy. 2019. Visited on 11 April 2019 at http://www.stratigraphy.org/index.php/ics-chart-timescale.

Iturralde-Vinent, M., and MacPhee, R.D.E. 1999. Paleogeography of the Caribbean region, implications for Cenozoic biogeography. *Bulletin of American Museum Natural History*, (238): 1-95.

IUCN 2019. The IUCN Red List of Threatened Species. Version 2019-3. http://www.iucnredlist.org. Accessed on 10 December 2019.

Jiménez Vázquez, O. 1997. Seis nuevos registros de aves fósiles en Cuba. *El Pitirre*, 10(2):49.

Jiménez Vázquez, O. 2001. Registros ornitológicos en residuarios de dieta de los aborígenes precerámicos cubanos. *El Pitirre*, 14(3):120–126.

Jiménez Vázquez, O. 2015. Sobre la coexistencia de los aborígenes precolombinos y los monos en Cuba. *Cuba Arqueológica*, 8(2):33-40.

Jiménez Vázquez, O., and Milera, J. F. 2002. Cánidos precolombinos de Las Antillas: Mitos y verdades*. Gabinete de Arqueología*, 2(2):78-87.

Jiménez Vázquez, O., Condis, M.M., and García Cansío E. 2005. Vertebrados Post-glaciales en un Residuario fósil de *Tyto alba* Scopoli (Aves: Tytonidae) en el Occidente de Cuba. *Revista Mexicana de Mastozoología*, 9:80–101.

Jiménez Vázquez, O. and Arrazcaeta, R. 2008. Las aves en la arqueología histórica de La Habana Vieja. *Boletín del Gabinete de Arqueología*, 7:17–29.

Jiménez Vázquez, O. and C. Arredondo Antúnez. 2011. Los mamíferos en la arqueozoología. Capítulo 5: 206-211pp, en *Mamíferos en Cuba* (R. Borroto-Páez y C. A. Mancina, eds), UPC Print, Finlandia, 276 p.

Jiménez Vázquez, O., and Arrazcaeta, R. 2015. Las aves y su relación con las culturas precolombinas de Cuba. *Boletín del Gabinete de Arqueología*, 11:141–157.

Jiménez Vázquez, O., and Orihuela, J. (in press). Nuevos hallazgos paleobiológicos de aves extintas o en peligro crítico endémicas de cuba, géneros *Ara* (Psittacidae), *Pulsatrix* (Strigidae) y *Campephilus* (Picidae). *Novitates Caribaea* (Scheduled publication on no. 16, 2020).

Jull, A.J.T., Iturralde-Vinent, M., O'Malley, J.M., MacPhee, R.D.E., McDonald, H.G., Martin, P.S., Moddy, J., and Rincón, A. 2004. Radiocarbon dating of extinct fauna in the Americas recovered from tar pits. *Nuclear Instruments and Methods in Physics Research*, B 223/224:668–671.

Keegan, W. F. (1992). *The People Who Discovered Columbus. The Prehistory of the Bahamas.* University Press of Florida, Gainesville.

Koch, Paul L., and A. D. Barnosky. 2006. Late quaternary extinctions: State of debate. *Annual Reviews of Ecology, Evolution and Systematics 37:215-250pp.*

Kozlowski, J. 1974. Preceramic cultures in the Caribbean. *Prace Archeologiczne Zeszyt*, 20.

Laffoon, J. E. 2012. Pattern of paleomobility in the ancient Antilles: an isotopic approach. (PhD Diss). Universidad de Leiden.

Laffoon, J., E. Plomp, G. Davies, M. Hoogland y C. L. Hofman. 2013. The movement and exchange of dogs in the prehistoric Caribbean: an isotopic investigation. *International Journal of Osteoarchaeology*: doi: 101002/oa.2313 (online 18 Apr 2013).

Laffoon, J., Menno L. P. Hoogland, Gareth R. Davies, and Corine L. Hofman. 2017. A multi-isotope investigation of human and dog mobility and diet in the pre-colonial Antilles. *The Journal of Human Palaeoecology*: doi.org/10.1080/14614103.2017.1322831

LeFebvre, M. J., DuChemin, G., deFrance, S. D., Keegan, W. F. and Walczesky, K. (2018a). Bahamian hutia (*Geocapromys ingrahami*) in the Lucayan realm: Pre-Columbian exploitation and translocation. Environmental Archaeology, https://doi.org/10.1080/14614103.2018.1503809.

LeFebvre, M. J., and Giovas, C. M., and Laffoon, J. E. (2018b). Advancing the study of Amerindian ecodynamics in the Caribbean: current perspectives. Environmental Archaeology, https://doi.org/10.1080/14614103.2018.1505224.

Lorenzen, E. D., Nogues-Bravo, D., *et al* (2011). Species-specific responses of late Quaternary megafauna to climate change and humans. *Nature*, 479: 359-364.

Lyons, SK, Miller, JH, Fraser, D, Smith, FA, Boyer, A, Lindsey, E, and Mychajliw, AM. 2016. The changing role of mammal life histories in late Quaternary extinction vulnerability on continents and islands. *Biology Letters*, doi:10.1098/rsbl.2016.0342

MacPhee, R.D.E., Flemming, C., and Lunde D.P. 1999. "Last Occurrence" of the Antillean insectivoran Nesophontes: new radiometric dates and their interpretations. *American Museum Novitates*, 3261:1–20.

MacPhee, R.D.E. & Flemming, C. 1999. *Requiem Aeternam*: the last five hundred years of mammalian species extinctions. *Extinctions in near time: causes, contexts, and consequences* (ed. by R.D.E. MacPhee), pp. 333–367. Kluwer Academic, New York, NY.

MacPhee, R.D.E., Iturralde-Vinent, M.A., and Jiménez Vázquez O. 2007. Prehistoric sloth extinctions in Cuba: Implications of a new "Last" appearance date. *Caribbean Journal of Science*, 43(1):94–98.

Mancina, C. A., and García-Rivera, L. 2005. New genus and species of fossil bat (Chiroptera: Phyllostomidae) from Cuba. *Caribbean Journal of Science*, 41(1):22–27.

Mancina, C. A., and L. M. Echenique-Díaz, Adrián Tejedor, L. García, A. Daniel- Álvarez, M., and A. Ortega-Huerta. 2007. Endemics under threat: an assessment of the conservation status of Cuban bats. *Hystrix International Journal of Mammalogy*, 18(1): 3-15.

Martínez Arango, F. 1968. *Superposición Cultural en Damajayabo.* Ciencia y Técnica, Instituto del Libro, La Habana: 123pp.

Martínez Gabino, A., E. Vento Canosa and C. Roque García. 1993. *Historia Aborigen de Matanzas*. Ediciones Matanzas, Matanzas.

Martínez-López, J. G., C. Arredondo Antúnez, R. Rodríguez Suárez and S. Díaz-Franco. 2009. Aproximación tafonómica en los depósitos humanos del sitio arqueológico Canímar Abajo, Matanzas, Cuba. *Arqueología Iberoamericana*, 4: 5-21pp.

Marrero, L. 1972. *Cuba: Economía y Sociedad Vol. 1*. Editorial Playor, S. A., Madrid

Mickleburgh, H. L., and Pagán-Jiménez, J. 2012. New insights into the consumption of maize and other food plants in the pre-Columbian Caribbean from starch grains trapped in human dental calculus. *Journal of Archaeological Science* 39: 2468-2478.

Miller, G. S., (1916). Bones of mammals from indian sites in Cuba and Santo Domingo. *Smithsonian Miscellaneous Collections*, 66 (12): 1-10.

Miller, G. S. 1929. Mammals eaten by Indians, owls and Spaniards in the coast region of the Dominican Republic. *Smithsonian Miscellaneous Collection 66 (13): 1-3pp.*

Morgan, G. S.1991. Neotropical Chiroptera from the Pliocene and Pleistocene of Florida. *Bull. Am. Mus. Nat. Hist. 206: 176-213pp.*

Morgan, G. S. 2001. Patterns of extinction in West Indian Bats. In C.A. Woods and F. E. Sergile (Editors), *Biogeography of the West Indies: Patterns and Perspectives:* 369-406. Boca Ratón, Florida: CRC Press.

Morgan, G.S., and Woods, C.A. 1986. Extinction and zoogeography of the West Indian land mammals. *Biology Journal of the Linnaean Society*, 28:167–203.

Morgan, G. S., and J. A. Ottenwalder. 1993. A new extinct species of Solenodon (Mammalia: Insectivora: Solenodontidae) from the Late Quaternary of Cuba. *Annals of Carnegie Museum,* 62: 151-164.

Muñiz, C.L. and Domínguez Samalea, Y. 2014. Las cuevas de Paredones y del Túnel y la Caverna de Pío Domingo: Principales yacimientos fosilíferos de vertebrados del Pleistoceno en Cuba occidental. *Geoinformativa*, 8(2):32–43.

Napolitano, M. F. DiNapoli, R.J., Stone, J.H., Levin, M.J., Jew, N.P., Lane, B.G., O'Connor, J.T., and Fitzpatrick, S.M. 2019. Reevaluating human colonization of the Caribbean using chronometric hygiene and Bayesian modeling. *Science Advances*, 5: eaar7806.

Newsom, L. A., and E. S. Wing. 2004. *On land and sea. Native American uses of biological resources in the West Indies*. The University of Alabama Press, 323 pp.

Norr, L. 2002. Bone isotopic analysis and prehistoric diet at the Tutu site. *En*: E. Righter (Ed.), *The Tutu Archaeological Village* Site: 263-273. Routledge, London.

Nuñez Jiménez, A., and Mayo, N. A. 1970. Antigüedad de la fauna vertebrada fósil de Cuba. *Serie Espeleológica y Carsológica*, 7.

Olson, S. L. 1982. Biological archaeology in the West Indies. *Florida Anthropologist*, 35: 162-168.

Orihuela, J. 2010. Late Holocene Fauna from a Cave Deposit in Western Cuba: post Columbian occurrence of the Vampire Bat *Desmodus rotundus* (Phyllostomidae: Desmodontinae). *Caribbean Journal of Science*, 46:297–312.

Orihuela, J. 2011. Skull variation of the vampire bat *Desmodus rotundus* (Chiroptera: Phyllostomidae): taxonomic implication for the Cuban fossil vampire bat *Desmodus puntajudensis*. Chiroptera Neotropical, 17(1):963-976.

Orihuela, J. 2013. Fossil Cuban crow *Corvus* cf. *nasicus* from a Late Quaternary cave deposit in northern Matanzas, Cuba. *Journal of Caribbean Ornithology*, 26:12–16.

Orihuela, J. 2019. An annotated list of Late Quaternary extinct birds of Cuba. *Ornitología Neotropical*, 30: 57–67.

Orihuela, J. and Tejedor A. 2012. Peter's ghost-faced bat *Mormoops megalophylla* (Chiroptera: Mormoopidae) from a pre-Columbian archeological deposit in Cuba. *Acta Chiropterologica*, 14:63–72.

Orihuela, J., Jiménez Vázquez, O., and Garcell, J.F. 2016. Modificaciones tafonómicas en restos óseos: ejemplos arqueológicos y paleontológicos de Mayabeque y Matanzas, Cuba. *Cuba Arqueológica*, IX (2):13–36.

Orihuela, J., Viera Muñoz, R. A., and Pérez Orozco, L. 2018. Contribución a la cronología y la paleodieta de un individuo aborigen excavado en el sitio arqueológico El Morrillo (Matanzas, Cuba). *Cuba Arqueológica*, X (2):16-31.

Orihuela, J., Viñola, L. W., and Viera, R. A. (2020). New bat locality records from Cuba with emphasis on the province of Matanzas. *Novitates Caribaea*, 15:96-116.

Orihuela, J., Pérez Orozco, L., Álvarez, Viera, R. A., and Santana, C. (submitted, April 2019).

Late Holocene land vertebrate fauna from Cueva de los Nesofontes, Western Cuba: last appearance dates and paleoecology. (Target journal *Palaeontologia Electronica*) Biorxiv DOI: https://doi.org/10.1101/2020.01.17.909663.

Pagán-Jiménez, J., M. A. Rodríguez López, A. Chanlatte Baik and Narganes Storde, Y. 2005. La temprana introducción y uso de algunas plantas domésticas, silvestres y cultivos en las Antillas precolombinas. Una primera revaloración desde la perspectiva del 'arcaico' de Viequez y Puerto Rico. *Diálogo Antropológico* 3(10): 7-33.

Pagán-Jiménez, J. and R. Rodríguez Ramos. 2007. Sobre el origen de la agricultura en Las Antillas. *In* B. Reid, H. Petitjean, y A. Curet (Eds.), *Twenty-first Congress of the International Association for Caribbean Archaeology*: 252-259. University of the West Indies, Kingston.

Pagán-Jiménez, J. and J. R. Oliver. 2008. Starch residues on lithic artifacts from two contrasting contexts in northwestern Puerto Rico: Los Muertos Cave and Vega de Nelo Vargas Farmstead. *In* C. Hoffman, M. Hoogland y A. Van Gijn (Eds.), *Crossiing the Borders: New Methods and Techniques in the Study of Archaeological Materials from the Caribbean*: 137-158. University of Alabama Press, Tuscaloosa.

Pestle, W. J. 2010. *Diet and Society in Prehistoric Puerto Rico* (PhD diss). Graduate College of the University of Illinois at Chicago, University of Illinois.

Pino, M. R. 1978. Consideraciones Sobre los Elementos Dietarios del Sitio Levisa, Mayarí. In Pp. 135-148 "*Cuba Arqueológica*" Editorial Oriente, Santiago de Cuba, 270 pp.

Pino, M. R. and Castellanos, N. 1985. Acerca de la asociación de perezosos cubanos extinguidos con evidencias culturales de aborígenes cubanos. *Reporte Investigativo del Instituto de Ciencias Sociales*, 4: 1-29.

Pino, M. R. 2012. Association of Cuban terrestrial mammals with aborigines' cultural evidences. Pp. 357-362 in Borroto-Páez, R., Woods, C. A., and Sergile, F. E. (eds) *Terrestrial Mammals of the West Indies*. Florida Museum of Natural History and Wacahoota Press, Gainesville.

Pose, J., R. Sampedro and M. Celaya. 1988. Contribución al estudio de la domesticación de roedores de la época prehispánica mediante el análisis de tomografía axial computarizada, rayos x y exámenes microscópicos de evidencias óseas. *Anuario de Arqueología* (año 1988): 70–83. La Habana, Cuba.

Pregill, G.K., Steadman, D.W., Olson, S.L., & Grady, F. V. (1988) Late Holocene fossil vertebrates from Burma Quarry, Antigua, Lesser Antilles. *Smithsonian Contributions to Zoology*, 463: 1–27.

Pregill, G. K., D. W. Steadman, and D. R. Watters. 1994. Late Quaternary vertebrate

faunas of the Lesser Antilles: Historic components of Caribbean Biogeography. *Bulletin of the Carnegie Museum of Natural History. 30 :1-51 pp.*

Prescott, G. W., William, D. R., Balmford, A., Green, R. E. and Manica, A. 2012. Quantitative global analysis of the role of climate and people in explaining late Quaternary megafaunal extinctions. *PNAS*, 109(2): 4527-4531.

Reimer, P.J, Bard, E. Bayliss, A., Warren, B. J. et al. 2013. IntCal13 and Marine13 radiocarbon age calibration curves 0-50,000 years Cal BP. *Radiocarbon*, 55(4): 1869-1887, DOI: 10.2458/azu_js_rc.55.16947

Reyes Cardero, J. M. 1997. Estudios dietarios de cinco sitios apropiadores ceramistas (protoagrícola) del sur oriente cubano. *El Caribe Arqueológico*, 2: 41-49.

Rímoli, R. O. 1974. Prueba zooarqueológica de tránsito del hombre precolombino de Cuba hacia La Hispaniola. *Revista Dominicana de Antropología e Historia*, 4:27-28.

Rímoli, R. O. 1976. Roedores fósiles de La Hispaniola. Universidad Central del Este. *Serie Científica*, III: 1-94 pp.

Rodríguez-Durán, A., and Santiago-Valentín, E. 2014. Una historia de la mastozoología en el Caribe insular. Pp. 129-142, in: *Historia de la Mastozoología en Latinoamérica, las Guayanas y el Caribe* (J. Ortega, J. L. Martínez, and D. G. Tiria, eds.). Editorial Murciélago Blanco y Asociación Ecuatoriana de Mastozoología, Ecuador.

Roksandic, M, WM Buhay, Y. Chenique de Armas, R. Rodríguez, M. Ceperos, I. Roksandic, S. Mowat, L.M. Viera, C. Arredondo, A. Martínez Fuentes and D.G. Smith. 2015. Radiocarbon and Stratigraphic Chronology of Canímar Abajo, Matanzas, Cuba. *Radiocarbon*, 57: DOI: 10.2458/azu_rc.57.18313.

Rouse, I. 1992. *The Tainos: Rise and Decline of the People Who Greeted Columbus*. Yale University Press, 211pp.

Rzebik-Kowalska, B., and Woloszyn B.W. 2012. New data on Nesophontes subfossil populations from Cuba and Isla de la Juventud (Cuba). *N. Jb. Geol. Palaont. Abh.*, 263/2:155–166.

Salgado, E.J., Calvache, D.G., MacPhee, R.D.E., and Gould, G.C. 1992. The monkey caves of Cuba. *Cave Sciences*, 19(1):25–28.

Sauer, C. O. 1966. *The Early Spanish Main*. University of California Press, Berkeley.

Silva-Taboada, G. 1974. Fossil Chiroptera from cave deposits in central Cuba, with a description of two new species (genera *Pteronotus* and *Mormoops*) and the first West Indian record of *Mormoops megalophylla*. *Acta Zoologica Cracoviensia*, 19:33–73.

Silva Taboada, G. 1976. Historia y actualización taxonómica de algunas especies antillanas de murciélagos de los géneros *Pteronotus*, *Brachyphylla*, *Lasiurus*, y *Antrozous*. *Poeyana*, 153:1–24.

Silva Taboada, G. 1979. *Los Murciélagos de Cuba.* Editorial Academia, La Habana.

Silva Taboada, G., Suárez, W., and Díaz-Franco, S. 2007. *Compendio de los Mamíferos terrestres autóctonos de Cuba vivientes y extinguidos.* Ediciones Boloña, La Habana.

Socha, R., and Woloszyn, B. W. 2010. Holocene remains of the genus Heteropsomys (= Boromys; Rodentia) from Cueva del Abuelo (Isla de Pinos, Cuba) in *The Nature and Culture of Latin America. Review of Polish Studies* (eds Z. Mirek., A. Flakus, A. Krzanowski, A. Paulo & J. Wojtusiak), pp. 1–9. W. Szafer Institute of Botany, Polish Academy of Sciences, Kraków.

Soto-Centeno, J.A. and Steadman, D.W. 2015. Fossils reject climate change as the cause of extinction of Caribbean bats. *Scientific Reports*, 5, doi: 10.1038/srep07971.

Soto-Centeno, J. A., N. B. Simmons, and D. W. Steadman. 2017. The bat community of Haiti and evidence for its long-term persistence at high elevations. *PLoS ONE* 12: e0178066.

Steadman, D.W., G. K. Pergill, and S. Olson. 1984. Fossil vertebrates from Antigua, Lesser Antilles: Evidence for Late Holocene human caused extinctions in the West Indies. *Proc. Natl. Acad. Sci. U.S.A. 81:4448-4451pp.*

Steadman, D. S., and Franklin, J. 2014. Changes in a West Indian bird community since the late Pleistocene. Journal of Biogeography, doi:10.1111/jbi.12418.

Steadman, D.W., et al. 2005. Asynchronous extinction of late Quaternary sloths on continents and islands. *Proceedings of the National Academy of Science*, 102:11763–11768.

Steadman, D.W., Albury, N.A., Kakuk, B., Mead, J.I., Soto-Centeno, J.A., Singleton, H.M., and Franklin, J. 2015. Vertebrate community on an ice-age Caribbean island. *Proceedings of the National Academy of Sciences USA*, 12, 5963–5971.

Stoetzel, E., Royer, A., Cochard, D., and Lenoble, A. 2016. Late Quaternary changes in bat palaeobiodiversity and palaeobiogeography under climatic and anthropogenic pressure: new insights from Marie-Galante, Lesser Antilles. Quaternary Science Reviews, 143, 150–174.

Suárez, WS and SL Olson. 2001. A remarkable new species of small falcon from the Quaternary of Cuba (Aves: Falconidae: *Falco*). *Proceedings of the Biological Society of Washington* 114: 34–41.

Suárez, W.S. and Díaz-Franco, S. 2003. A new fossil bat (Chiroptera: Phyllostomidae) from a Quaternary cave deposit in Cuba. *Caribbean Journal of Science*, 39: 371–377.

Suárez, W.S., and Olson, S.L. 2015. Systematics and distribution of the giant fossil barn owls of the West Indies (Aves: Strigiformes: Tytonidae). *Zootaxa*, 4020: 533-553.

Tabío, E., and E. Rey. 1979. *Prehistoria de Cuba*. Ciencias Sociales, La Habana, Cuba, 234pp.

Tabío, E. 1984. Nueva periodización para el estudio de las comunidades aborígenes de Cuba. *Islas* 78: 37-52.

Torres Etayo, D. 2006. *Taínos: Mitos y Realidades de un Pueblo Sin Rostro*. Editorial Asesor Pedagógico, S. A., México.

Turvey, S., J. R. Oliver, Y. M. Narganes Storde, and P. Rye. 2007. Late Holocene extinction of Puerto Rican native land mammals. *Biology Letters 3: 193-196pp.*

Turvey, S. T. 2009a. Holocene mammal extinctions. Chapter 3 in S. Turvey (ed.) *Holocene Extinctions*. Oxford University Press, 353pp.

Turvey, S. T. 2009b. In the shadow of the megafauna: prehistoric mammals and bird extinctions across the Holocene. Pp. 17-40 *in* S. T. Turvey (ed). *Holocene Extinctions* Oxford University Press, Oxford, UK.

Turvey, S. T., Kennerley, R. J., Nunez-Mino, J. M, and Young, R. P. 2017. The last survivors: current status and conservation of the non-volant land mammals of the insular Caribbean. *Journal of Mammalogy*, 98(4): 918-936.

Tuttle, M. D. 1995. Saving North America's beleaguered bats. *National Geographic*, 188(2): 37-57.

Ulloa Hung, J. and Valcárcel Rojas, R. 2016. *Indigenas e indios end el Caribe: Presencias, Legado y Estudio.* Editorial Buho, S. R. L., Santo Domingo.

Ulloa Hung, J., and Valcárcel Rojas, R. 2019. Chapter 12: Construction and deconstruction of the "Archaic" in Cuba and Hispaniola in Hofman, C.L. and Antczak, A.T. (eds): *Early Settlers of the Insular Caribbean: Dearchaizing the Archaic*. Sidestone Press (Leiden), pp. 163-176.

Upham, N. S., and Borroto-Páez, R. 2017. Molecular phylogeography of endangered Cuban hutias within the Caribbean radiation of capromyid rodents. Journal of Mammalogy, 98 (4): 950-963. DOI: https://doi.org/10.1093/jmammal/gyx077.

Valcárcel, R. 2002. *Banes Precolombino. La Ocupación Agricultora*. Ediciones Holguín, Holguín, Cuba.

Valcárcel, R. 2008. Las sociedades agriculturas ceramistas en Cuba. Una mirada desde los datos arqueológicos y etnohistóricos. *El Caribe Arqueológico*, 11: 2-19.

Valcárcel Rojas, R. 2012. Interacción colonial en un pueblo de indios encomendados: El Chorro
de Maita, Cuba. Universidad de Leiden, Holanda.

Valcárcel Rojas, R., Ulloa Hung, J., and Feria García, O. 2019. Chapter 13: Levisa 1. Studying
the earliest indigenous peoples of Cuba in multicomponent archaeological sites" in:
Hofman, C.L. and Antczak, A.T. (eds): *Early Settlers of the Insular Caribbean:*
Dearchaizing the Archaic. Sidestone Press (Leiden), pp. 177-190.

Viñola López, L.W., Garrido, O.H., and Bermúdez A. 2018. Note on *Mesocapromys*
sanfelipensis (Rodentia: Capromyidae) from Cuba. *Zootaxa*, 4410 (1):164–176.

Wiley, JW & GM Kirwan. 2013. The extinct macaws of the West Indies, with special reference
to Cuban Macaw *Ara tricolor*. *Bulletin of the British Ornithologists' Club* 132: 125-156.

Wing, E. S. 2001. Native American use of animals in the Caribbean. Pp. 481–518 *In* CA
Woods, FE Sergile (eds) *Biography of the West Indies, Patterns and Perspectives*. CRC
Press: Boca Raton.

Wing, E. S. and Scudder, S. J. 1983. Animal exploitation by prehistoric people living on a
tropical marine edge. Pp. 197-210 in Grigson, C. and Clutton-Brock, J. (eds). *Animals and*
Archaeology: 2 Shell Middens, Fishes and Birds. BAR International Series 183, Oxford.

Woods, C.A., & Sergile, F.E. 2009. Antilles biology. In Encyclopedia of Islands. R.G. Gillespie
& D.A. Clauge (eds.). University of California Press, Berkeley, CA.

3   Figure 1. Map of Cuba highlighting the 17 unique fossil localities described in this study. Numbered labels represent locality names as
4   listed in parenthesis in Table 1 and described in Supplementary Text S2 study site details. Cuba inset shows specific localities within
5   Artemisa, La Habana, and Mayabeque provinces. Marker 8/12 in inset represents two nearby unique localities.

Figure 2. Chronology of extinction of Cuban vertebrates based on calibrated AMS radiocarbon (^{14}C) dates. Taxonomic groups are
indicated by a silhouette (respectively: Testudines, Rodentia, Primates, Pilosa, Eulipotiphla, Chiroptera, Aves). Circles represent
radiocarbon dates: closed circle = direct date, open circle = indirect date, gray = earliest appearance in the fossil record, black = last
appearance date (LAD). Horizontal bars are used to highlight the existence of a taxon over time and arrows indicate extant taxa.
Vertical dashed lines show the Early Amerindian (a = ~6ky), Late Amerindian (b = ~1500y), and post-Columbian (c = <500y)
subintervals (see Methods and Supplementary Text S2).

Figure 3. Representative extinct and endangered vertebrate fauna from Cuba associated with radiocarbon LADs. Clockwise from top left: a = *Desmodus rotundus* skull (Post-Columbian, Cueva de los Nesofontes); b = *Desmodus rotundus* distal humerus (Post-Columbian, Cueva de los Nesofontes; an attempt to directly date this specimen was unsuccessful); c = *Artibeus anthonyi* left hemimandible (paleontological context, ~5190 BP, Cueva del Gato Jíbaro); d = *Antrozous koopmani* left hemimandible (~1150 BP, Cueva de los Nesofontes); e = *Nesophontes micrus* skull (late Amerindian, Cueva de los Nesofontes); f and f1 = *Nesophontes major* skull and hemimandible (~1418 BP, Cueva de los Nesofontes); g = *Pulsatrix arredondoi* tarsus (~1390 BP, Cueva del Muerto); h = *Mesocapromys kraglievichi* left hemimandible (late Amerindian, Cueva de los Nesofontes); i = *Solenodon cubanus*, left hemimandible (~820 BP, Cueva de Calero). All bars scaled to 10 mm, except in b as indicated.

2   Figure 4. Proportion of extinctinct taxa per anthropogenic interval in Cuba. Half of the

3   documented LADs postdate 4ka and about 42% of these date within the later Amerindian

4   interval defined here (see Methods and Supplementary Text S2).

Table 1. Direct Accelerator Mass Spectrometer (AMS) radiocarbon (^{14}C) dates for individual fossil remains from 17 unique localities across Cuba. All estimates represented as conventional ^{14}C dates in chronological order. Intervals definitions: Late Amerindian (l Am), post Columbian (p Col), Amerindian (Am), Early Holocene (e HOL), Middle Holocene (m HOL), and Late Pleistocene (l PL).

Taxa	Interval	Conventional ^{14}C age (yr BP)	Calibrated ^{14}C age (cal BP, 2δ)	Sample No.	Site / Deposit	Source
Nesophontes micrus	l Am–p Col	490 ± 50	640–341	Beta-115695	Cueva Martin, Santi Spíritus (1)	MacPhee et al. 1999a
Solenodon cubanus	l Am	650 ± 15	665–561	UCIAMS-218808	Cueva La Caja/Nesofontes, Mayabeque (2)	This paper
Mesocapromys nanus	l Am	665 ± 20	670–562	UCIAMS-218804	Abra de Figueroa (owl roost), Matanzas (3)	This paper
N. major	l Am	670 ± 15	670–565	UCIAMS-218809	Abra de Figueroa (owl roost), Matanzas (3)	This paper
N. micrus	l Am	770 ± 50	790–653	Beta-115696	Solapa La Jaula (4)	MacPhee et al. 1999a
S. cubanus	l Am	820 ± 15	760–690	UCIAMS-218807	Cueva Calero, Cantel, Matanzas (arch.) (5)	This paper
N. micrus	l Am	840 ± 15	786–703	UCIAMS-218810	Abra de Figueroa (owl roost) (3)	This paper
N. micrus	l Am	860 ± 30	901–695	ICA-18B/1234	Cueva El Gato Jíbaro, Matanzas (6)	This paper
Antrozous koopmani	l Am	1150 ± 30	1174–979	Beta-526491	Cueva La Caja/Nesofontes, Mayabeque (2)	This paper
Boromys torrei	l Am	1180 ± 15	1175–1061	UCIAMS-218802	Cueva La Caja/Nesofontes, Mayabeque (2)	This paper
Homo sapiens	l Am	1190 ± 40	1255–984	ICA-16B/0605	Cueva del Muerto, Mayabeque (7)	Orihuela et al. 2016
A. koopmani	l Am	1280 ± 30	1288–1176	Beta-526493	Cueva La Caja/Nesofontes, Mayabeque (2)	This paper
Artibeus anthonyi	l Am	1290 ± 30	1286–1180	ICA 14B/1102	Cueva La Caja/Nesofontes, Mayabeque (2)	Orihuela et al. unpub.
A. koopmani	l Am	1300 ± 30	1291–1181	Beta-526492	Cueva La Caja/Nesofontes, Mayabeque (2)	This paper
B. offella	l Am	1320 ± 15	1294–1187	UCIAMS-218806	Cuevas Blancas, Mayabeque (arch.) (8)	This paper
Puslatrix arredondoi	l Am	1390 ± 30	1348–1277	ICA-16B/0606	Cueva del Muerto, Mayabeque (7)	Jiménez et al. unpub.
N. major	l Am	1418 ± 20	1348–1294	Beta-392022	Cueva La Caja/Nesofontes, Mayabeque (2)	Orihuela et al. unpub.
H. sapiens	l Am	1570 ± 40	1548–1378	ICA-19B/0508	Cueva El Gato Jíbaro, Matanzas (6)	This paper
Macrocapromys acevedo	l Am	1590 ± 30	1545–1409	Beta-526495	Cueva de Leonel, Trinidad (9)	This paper
M. nanus	l Am	1675 ± 15	1613–1541	UCIAMS-218803	Cueva del Caracol (10)	This paper
M. nanus	l Am	1775 ± 20	1780–1615	UCIAMS-218805	Cueva de los Paredones, Mayabeque (11)	This paper
M. kraglievichi	l Am	1780 ± 50	1822–1569	ICA-15B/0118	Solapa del Megalocnus, Mayabeque (12)	This paper
Phyllops vetus	l Am	1960 ± 30	1989–1830	ICA 18B/0845	Cueva La Caja/Nesofontes, Mayabeque (2)	Orihuela et al. unpub.
Megalocnus rodens	Am–m HOL	4190 ± 40	4844–4584	Beta-206173	Solapa de Silex, Mayabeque (13)	MacPhee et al. 2007
Parocnus browni	Am–m HOL	4960 ± 280	6310–4975	AA35290	Las Breas de San Felipe, Matanzas (14)	Jull et al. 2004
A. anthonyi	Am–m HOL	5190 ± 40	6172–5773	ICA-19B/0450	Cueva El Gato Jíbaro, Matanzas (6)	This paper
M. rodens	Am–m HOL	6250 ± 50	7270–7007	B-115697	Cueva Beruvides, Matanzas (15)	Steadman et al. 2005
P. browni	l PL–e HOL	10520 ± 440	13307–11138	AA35291	Las Breas de San Felipe, Matanzas (13)	Jull et al. 2004
P. browni	l PL–e HOL	11880 ± 420	15236–12973	AA35292	Las Breas de San Felipe, Matanzas (13)	Jull et al. 2004
Tyto noeli	l PL	17406 ± 160	21494–20590	NA	El Abrón, Pinar del Río (16)	Suárez & Franco 2003
Pinus sp.	l PL	>30000 ± 2000	NA	L-130A	Casimbas de Ciego Montero, Cienfuegos (17)	Kulp et al. 1952
Unsp. Wood	l PL	25000 ± 2000	35503–25759	L-130B	Casimbas de Ciego Montero, Cienfuegos (17)	Kulp et al. 1952