Multi-tissue epigenetic analysis of the osteoarthritis susceptibility locus mapping to the plectin gene *PLEC*

A.K. Sorial[†*], I.M.J Hofer[†], M. Tselepi[†], K. Cheung[†], E. Parker[†], D.J. Deehan[‡], S.J. Rice[†], J. Loughlin[†*]

[†]Newcastle University, Biosciences Institute, Newcastle upon Tyne, UK

[‡]Freeman Hospital, Newcastle Upon Tyne, UK

Author email addresses:

Antony Sorial: tony.sorial@newcastle.ac.uk

Ines Hofer: ines.hofer@newcastle.ac.uk

Maria Tselepi: maria.tselepi@newcastle.ac.uk

Kathleen Cheung: kat.cheung@newcastle.ac.uk

Eleanor Parker: eleanor.parker@newcastle.ac.uk

David Deehan: david.deehan@newcastle.ac.uk

Sarah Rice: sarah.rice@newcastle.ac.uk

John Loughlin: john.loughlin@newcastle.ac.uk

[*]Address correspondence and reprint requests to:

Mr. Antony K. Sorial or Professor John Loughlin, Newcastle University, Biosciences Institute, International Centre for Life, Newcastle upon Tyne, NE1 3BZ, United Kingdom.

Tel: +44 (0)191 241 8988; Fax: +44 (0)191 241 8666. E-mail: tony.sorial@newcastle.ac.uk (A.K. Sorial), john.loughlin@newcastle.ac.uk (J. Loughlin)

Running   headline: Epigenetic   analysis   of   the   *PLEC*   OA   risk   locus

Objective

Osteoarthritis (OA) associated single nucleotide polymorphism (SNP) rs11780978 correlates

with differential expression of *PLEC,* and methylation quantitative trait loci (mQTLs) at

PLEC CpGs in cartilage. This implies that methylation links chondrocyte genotype and

phenotype, thus driving the functional effect. *PLEC* encodes plectin, a cytoskeletal protein

that enables tissues to respond to mechanical forces. We sought to assess whether *PLEC*

functional effects were cartilage specific.

Method

Cartilage, fat pad, synovium and peripheral blood were collected from patients undergoing

arthroplasty. *PLEC* CpGs were analysed for mQTLs and allelic expression imbalance (AEI)

was performed. We focussed on previously reported mQTL clusters neighbouring

cg19405177 and cg14598846. Plectin was knocked down in a mesenchymal stem cell (MSC)

line using CRISPR/Cas9 and cells phenotyped by RNA-sequencing.

Results

Novel mQTLs were discovered in fat pad, synovium and peripheral blood at both clusters.

The genotype-methylation effect of rs11780978 was stronger in cg14598846 than in

cg19405177 and stronger in joint tissues than in peripheral blood. We observed AEI in

synovium in the same direction as for cartilage. Knocking-down plectin impacted on

pathways reported to have a role in OA, including Wnt signalling, glycosaminoglycan

biosynthesis and immune regulation.

Conclusions

Synovium is also a target of the rs11780978 OA association functionally operating on *PLEC*.

In fat pad, mQTLs were identified but these did not correlate with *PLEC* expression,

suggesting the functional effect is not joint-wide. Our study highlights interplay between

genetic risk, DNA methylation and gene expression in OA, and reveals clear differences

between tissues from the same diseased joint.

Keywords: osteoarthritis, genetic risk, epigenetics, gene expression, *PLEC*, CRISPR/Cas9,

cytoskeletal

Introduction

Osteoarthritis (OA) is an extremely common musculoskeletal disorder that is characterised by focal loss of articular cartilage leading to chronic pain, disability, co-morbidities including cardiovascular disease, and premature death[1-3]. OA is polygenic and in recent years over 90 genome-wide significant loci have been mapped in Europeans using high-density single nucleotide polymorphism (SNP) arrays[4-10]. An overwhelming majority of the associated polymorphisms reside in non-protein-coding regions of the genome and, as such, OA genetic susceptibility is presumed to act principally via changes to gene expression. The availability of excised tissue following arthroplasty of an OA joint offers the opportunity to experimentally test this in disease-relevant cells, including cartilage chondrocytes, the single cell type in this tissue[11]. As such, many OA risk alleles have been shown to correlate with alteration in the expression of nearby genes in chondrocytes[12]. These effects are known as expression quantitative trait loci (eQTLs). DNA methylation changes at CG dinucleotides (CpGs) also regulate gene expression and OA-associated SNPs have been shown to correlate with methylation levels in cartilage DNA in *cis*[13]. These effects are known as methylation QTLs (mQTLs). Furthermore, several OA risk loci correlate with both methylation and expression, with alteration of the former being shown experimentally to modulate the latter[14]. These effects are known as methylation and expression QTLs (meQTLs). Overall, these studies indicate that DNA methylation links chondrocyte genotype and phenotype, with this epigenetic mechanism underpinning and driving a large proportion of the functional impact of OA genetic risk. Furthermore, in most instances, OA mQTLs localize to predicted enhancers[13], placing OA in the "enhanceropathy" category of common diseases[15].

We previously reported that an OA association marked by SNP rs11780978 (G>A) correlated with methylation of nine CpGs measured using the Illumina Infinium HumanMethylation450 genome-wide CpG array[16]. The SNP and CpGs are all located within the *PLEC* gene on chromosome 8q24.3. We subsequently focused on two of the most significant CpGs from amongst the nine, cg19405177 and cg14598846, and by using pyrosequencing, we replicated the correlation between rs11780978 genotype and CpG methylation in an independent panel of DNAs. The two pyrosequencing assays used to replicate cg19405177 and cg14598846 captured an additional seven CpGs flanking cg19405177 and an additional three CpGs adjacent to cg14598846. The methylation of these additional CpGs also correlated with rs11780978 genotype. Furthermore, using allelic expression imbalance (AEI) analysis (a complementary approach to standard eQTL

analysis[17]) we demonstrated that the OA risk-conferring A allele of rs11780978 correlated

with reduced *PLEC* expression, and that expression of the gene also correlated with

methylation at CpGs in the cg19405177 and cg14598846 clusters[16]. These mQTL, eQTL and

meQTL studies were all performed on cartilage DNA and RNA, and highlighted *PLEC* as a

target of the OA susceptibility mapping to chromosome 8q24.3.

PLEC encodes plectin, a large cytoskeletal protein that regulates signalling from the

extracellular environment to the cell nucleus[18-19]. Plectin enables cells to respond to external

mechanical stimuli and forces, such as those experienced by the chondrocyte during joint

movement[20]. Although cartilage loss is central to the OA disease process, there are

pathological changes to other tissues of the articulating joint that occur concurrent to and

following the loss of cartilage[21]. *PLEC* is expressed widely and in this report, we set out to

test whether the mQTL, eQTL and meQTL effects that we had discovered were cartilage

specific or whether they were active in other joint or non-joint tissues. We also modelled the

transcriptional effect that the OA risk-conferring allele of rs11780978 has on *PLEC* by

knocking-down plectin in a mesenchymal stem cell line using CRISPR/Cas9 and assessing

the impact on global gene expression.

Materials and Methods

OA Patients

Tissue samples were collected from patients undergoing knee or hip arthroplasty for primary

OA at the Newcastle upon Tyne Foundation Trust hospitals. Ethical approval was granted by

the Health Research Authority of the National Health Service (research ethics committee

reference 14/NE/1212) and each donor provided verbal and written informed consent. We

collected macroscopically intact cartilage (36 samples, distal from the lesion and avoiding

areas of fibrillated tissue), infrapatellar fat pad (68 samples) and synovium (81 samples). We

also collected whole peripheral blood samples from 55 of the patients just prior to their

surgery, using EDTA vacutainers for DNA extraction and Tempus[TM] tubes for RNA

extraction (ThermoFisher Scientific). In total, we analysed 240 samples from 202 OA

patients, with some patients donating more than one tissue type for analysis. Further patient

details can be found in Supplementary Table 1.

Nucleic acid extraction from tissue samples

Joint tissue samples were stored frozen and ground to a powder using a mixermill (Retsch Limited) under liquid nitrogen. For cartilage, RNA was extracted using TRIzol (Life Technologies) and DNA was extracted using the E.Z.N.A. Tissue DNA isolation kit (Omega Biotek, VWR). For fat pad and synovium, DNA and RNA were extracted using the E.Z.N.A. DNA/RNA isolation kit (Omega Biotek, VWR). For blood, DNA was extracted using the QIAamp DNA blood mini kit (Qiagen) and RNA was extracted using the Tempus™ Spin RNA isolation kit (ThermoFisher Scientific).

Quantitative gene expression

cDNA synthesis and quantitative PCR (qPCR) were performed as described previously[22]. Predesigned TaqMan assays (Integrated DNA Technologies) were used to quantify expression of the housekeeping genes *HPRT1*, *18S* and *GAPDH*. For *PLEC,* expression primers were designed using the Roche probe library system and are located in exons 19 and 20, which are present in all isoforms of the gene (primers 5'-CTGCGTAGGAAATACAGTT-3' and 5'-CAGCTGTTCCTTCTCGTCCT-3'). The relative expression of *PLEC* was calculated by the $2^{-\Delta Ct}$ method, where ΔCt is the mean Ct value of the three housekeeping genes subtracted from the Ct value of *PLEC*.

mQTL and eQTL analysis

Genotyping of association SNP rs11780978 and *PLEC* transcript SNP rs11783799 (A>G), DNA methylation analysis at the cg19405177 and cg14598846 CpG clusters, and *PLEC* AEI analysis using rs11783799 were all performed as described previously using the same protocols and assays[16]. In the methylation analysis, PCRs were performed in duplicate on each sample and the mean calculated, with samples being excluded from the analysis if the methylation between the PCR replicates differed by >5%. In the AEI analysis, PCRs were performed in triplicate on each tissue cDNA and DNA sample, with samples excluded from the analysis if the values between the PCR replicates differed by >5%.

Generation of a mesenchymal stem cell line constitutively expressing Cas9 (MSC-Cas9)

We purchased the immortalised human mesenchymal stem cell (MSC) line SCRC-4000 (ATCC) and transduced it with the lentiCRISPR v2 plasmid 52961, which contains a Cas9 insert and confers puromycin resistance (Addgene). The cells were expanded in complete growth medium consisting of serum free media (PCS-500-030, ATCC) supplemented with

Mesenchymal Stem Cell Growth Kit containing L-Alanyl-L-Glutamine (PCS-500-040,

ATCC), G418 disulphate salt solution (G8168, Sigma Aldrich) and penicillin-streptomycin

(30-2300, ATCC). Selection of Cas9 expressing cells was achieved by addition of 0.35ug/µl

of puromycin to the complete growth media. Constitutive expression of Cas9 was confirmed

by immunoblot with a Cas9 antibody (7A9-3A3, Cell Signalling Technology).

CRISPR/Cas9

CHOPCHOP (https://chopchop.cbu.uib.no/) was used to design guide RNA (gRNA)

sequences that would generate a 26bp frame-shift deletion of exon 3 of *PLEC* (primers 5'-

CTGTATGAAGACCTCCGCGA-3' and 5'-CCCCGAGAGGACCTCCAGCA-3'). This exon

is present in all isoforms of the gene. The gRNAs were then purchased as synthetic Alt-R®

CRISPR-Cas9 crRNAs alongside Alt-R® CRISPR-Cas9 tracrRNA (Integrated DNA

Technologies). MSC-Cas9 cells were plated at sub-confluence (45,000 cells/well) in 6-well

plates (VWR) and allowed to adhere for 24 hours. For deletions (4 replicates), paired crRNAs

were annealed to tracrRNA in a 1:1:2 ratio at 95°C for 5 minutes, whilst for controls (4

replicates), tracrRNA alone was used. Deletion and control mixes were separately combined

with Dharmafect1 (Dharmacon) in serum free media (PCS-500-030, ATCC) to produce a

transfection mix. 200ul of media/well was aspirated and replaced with transfection mix for 24

hours, after which the media was changed to complete growth medium and the cells

expanded in T182.5 flasks (VWR) in monolayer and until confluent (between 10-14 days),

with two changes of media per week. Cells were detached by trypsinization, washed in

phosphate-buffered saline (PBS), spun and the cell pellets frozen. DNA, RNA and protein

were simultaneously extracted from pellets using the NucleoSpin TriPrep Kit (Macherey-

Nagel, supplied by ThermoFisher). Deletion of the target region was confirmed by end-point

PCR   of   DNA   using   primers   flanking   the   guides   (primers   5'-

GAGGGGTTGAGCTGGATTCC-3' and 5'-CCAGCACCCACTCTGTAGAT-3') and by

Sanger sequencing.

Immunoblotting

Up to 10µg of total protein was resolved on NuPAGE™ 3-8% Tris-Acetate Protein Gels

(Invitrogen) in 1x NuPAGE™ Tris-Acetate SDS Running Buffer (Invitrogen). Protein was

transferred through a semi-dry transfer at 200mA/gel for 3 hours. Blots were probed with

anti-plectin guinea pig polyclonal serum (GP21, Progen)/goat anti-guinea pig (AB6908,

Abcam) and anti-GAPDH (MAB374, Sigma Aldrich)/goat anti-mouse (P0447, Dako) as a

loading control.

RNA-sequencing

For each sample, the TruSeq Stranded mRNA library prep kit (Illumina) was used to prepare

a sequencing library from 1.5µg of total RNA. All samples were then sequenced to a depth of

~52-71 million paired end reads on an Illumina NextSeq platform. Quality control of raw

sequencing reads was performed using FastQC (v0.11.7). All samples passed QC. Reads

were mapped to the hg38 (ENSEMBL version 91) reference genome using Salmon (v0.14.1)

in quasi-mapping mode on default settings[23]. All samples except one achieved alignment

rates of ~88%. Counts were summarised to gene level and imported into R using tximport[24].

Principle component analysis (PCA) revealed the sample with the low alignment rate to be an

outlier and this was removed from further analysis, leaving four *PLEC* deletion and three

controls. Unwanted variation in the dataset was estimated using surrogate variable analysis

(svaseq[25]). The number of surrogate variables was estimated at two and these were included

into the model for differential expression analysis. Differential expression analysis between

deletion and control samples was performed using the Wald test implemented in DESeq2[26].

The false discovery rate (FDR) was controlled using the Benjamini-Hochberg procedure.

Genes with an FDR *P*<0.05 were considered significant. Gene set enrichment analysis

(GSEA[27]) was used to discover overrepresented pathways in the set of up- and down-

regulated genes. The full gene list was ranked by signed log10(*P* value); positive sign if up-

regulated and negative if down-regulated. This pre-ranked list was used as input into the

fgsea package in Bioconductor to query enrichment of Reactome pathways[28]. Pathways with

an FDR *P*<0.25 were considered significant, as recommended by the developers of GSEA[27].

To visualise splice junctions, reads were aligned to hg38 reference genome using STAR

(v2.4.2a) short read aligner using default settings[29] and viewed in Integrative Genomics

Viewer (IGV[30]).

Statistical analyses

When correlating association between rs11780978 genotype and methylation of CpGs, *P*

values were calculated using the Kruskal-Wallis test. For comparison of global methylation

between tissues, Kruskal-Wallis with Dunn's multiple comparisons was used. For

quantitative *PLEC* expression analyses, *P* values were calculated using a Mann-Whitney 2-

tailed exact test. For *PLEC* AEI, *P* values were calculated using Wilcoxon's matched pairs

signed rank test. For heat maps, the correlation between all three genotypes (GG, GA and

AA) and methylation was determined using adjusted r^2 values calculated using a standard

least squares linear regression model and expressed as a percentage.

Results

***PLEC* cartilage mQTLs are present in other tissues**

We previously discovered *PLEC* mQTLs in cartilage DNA[16]. For cg19405177 and its seven

flanking CpGs (located in a 124bp interval), the OA risk-conferring A allele of rs11780978

correlated with increased methylation, whereas for cg14598846 and its three adjacent CpGs

(located in a 21bp interval), the A allele correlated with decreased methylation.

Supplementary Figure 1 highlights the relative positions of rs11780978 and the CpGs in

PLEC. In this report, we first assessed if these mQTLs were also active in DNA from fat pad,

synovium and blood. We also included a new cohort of cartilage DNAs. Figure 1 provides

the data for cg19405177 and cg14598846, whilst the data for all 12 CpGs across the four

tissues can be found in Supplementary Figures 2 (cartilage), 3 (fat pad), 4 (synovium) and 5

(blood). The mQTLs were replicated in the new batch of cartilage DNAs and in the same

directions as before. The mQTLs were also present in the other tissues and also in the same

direction.

When we plotted the methylation data by tissue type but independent of genotype,

there were significant differences in methylation levels (Supplementary Figure 6). For

example, cg19405177 demonstrated higher methylation values in fat pad and synovium

compared to cartilage and blood, whilst cg14598846 demonstrated higher methylation values

in blood compared to joint tissues. To emphasize this variability in CpG methylation in

tissues from the same patient, we plotted the cg19405177 and cg14598846 data for those

patients in whom we had studied two or more tissues. There were 18 such patients with data

generated at cg19405177 and 20 patients with data generated at cg14598846 (Figure 2). The

variability in methylation between the tissues of an individual is particularly clear for patient

6550 at cg19405177, and patient 6599 at cg14598846, with a methylation range of >30%

across the four tissues tested for each CpG.

We next constructed heat maps of the percentage effect of rs11780978 genotype on

methylation at each CpG in each tissue. For cartilage, we included data from the replication

study of our previous report[16] to maximise power. For the cg19405177 CpGs, the average

genotypic effect for all eight CpGs combined was 31.3% in cartilage, 17.4% in fat pad, 21.2 % in synovium and 9.3% in blood (Figure 3A). For the cg14598846 CpGs, the average genotypic effect for all four CpGs combined were comparable across the joint tissues, with values of 85.0% in cartilage, 81.4% in fat pad and 87.5% in synovium (Figure 3B). Although the genotypic effect in blood was lower, it was still high at 43.5%. At the individual CpG level, there were some striking differences between tissues. For example, cg19405177 had a genotypic effect of >30% in cartilage, synovium and blood but only 3.9% in fat pad. These large differences imply intra-tissue differences in the nature of the mQTLs.

Overall, this analysis highlighted that the previously reported *PLEC* mQTLs are active in non-cartilaginous tissues and that the effects are stronger in joint tissues than blood. Although the genotypic effect of rs11780978 is greater in cartilage for the cg19405177 CpGs, this is not the case for the cg14598846 CpGs.

The *PLEC* rs11780978 eQTL is active in synovium

Before testing for the *PLEC* rs11780978 eQTL by AEI analysis, we measured the relative expression of the gene in fat pad, synovium and blood along with cartilage as a comparator (Supplementary Figure 7). The qPCR data revealed that *PLEC* expression was equivalent between cartilage and synovium, was higher in fat pad and lower in blood.

We performed AEI as described previously[16]. In summary, we identified rs11780978 heterozygotes who were also heterozygous for the *PLEC* transcript SNP rs11783799 (r^2=0.93; the OA risk conferring A allele of rs11780978 correlates with the G allele of rs11783799). We then used rs11783799 to quantify the relative allelic expression of *PLEC* in the compound heterozygotes. The blood cDNA samples consistently failed quality control, with the PCR replicates differing by >5% in all samples. This is likely a reflection of the lower expression of *PLEC* in this tissue and we could not therefore generate blood AEI data. We were able to generate AEI data for 19 fat pad and 20 synovium compound heterozygotes (Figure 4). AEI was detectable in both tissues and as for our previous study in cartilage[16], the AEI ratios varied between patients, implying that other genetic polymorphisms can also modulate *PLEC* expression. Only in synovium did the direction of AEI significantly correlate with an allele of rs11783799, with reduced expression of *PLEC* correlating with the G allele of the SNP in the combined analysis (*P*=0.0054; Figure 4B). In fat pad, AEI was observed in a majority of the patients, but this did not correlate with a particular allele (*P*=0.31; Figure 4A).

To emphasize the variability of AEI in tissues from the same patient, we next plotted

267   the data for those patients in whom we had studied fat pad and synovium (Supplementary

268   Figure 8). There were six such patients with clear variability in the G/A ratio between the two

269   tissues tested for most of these, with patients 5680 and 5725 demonstrating particularly clear

270   inter-tissue variability in the ratio; in both, the G allele is more highly expressed in fat pad

271   but more lowly expressed in synovium relative to the A allele.

272   The significant AEI in synovium was in the same direction that we had observed in

273   cartilage[16], with the G allele of rs11783799, which marks the OA risk conferring A allele of

274   rs11780978, demonstrating reduced *PLEC* expression. In that previous report, we had studied

275   19 cartilage compound heterozygotes and the mean G/A ratio was 0.89. The mean synovium

276   G/A ratio we detected here was 0.92.

277   We next plotted the *PLEC* AEI data for the fat pad and synovium compound

heterozygotes against their individual methylation data. In fat pad, there was no evidence of a

significant correlation between methylation and AEI at cg19405177, cg14598846 or the

additional 10 CpGs measured (all *P* values >0.05; Supplementary Figure 9). However, in

synovium, three of the CpGs correlated with AEI; CpG3 from the cg19405177 cluster

(*P*=0.049), and cg14598846 (*P*=0.039) and CpG2 (*P*=0.047) from the cg14598846 cluster

(Supplementary Figure 10). These three CpGs had also shown meQTL activity in our

previous cartilage study[16]. For CpG3 from the cg19405177 cluster, the same direction of

effect is seen in cartilage and synovium, with increased methylation correlating with

decreased AEI (the $\log_2$ AEI ratio moves towards zero with increasing methylation).

However, for cg14598846 and CpG2 from the cg14598846 cluster, the effects are in opposite

directions between cartilage and synovium; in cartilage, increased methylation correlated

with decreased AEI whereas in synovium, increased methylation correlated with increased

AEI (the $\log_2$ AEI ratio moves away from zero with increasing methylation).

291   In summary, the *PLEC* rs11780978 eQTL previously observed in cartilage also

operates in synovium. Furthermore, we observed correlations between *PLEC* expression and

CpG methylation in synovium. These meQTLs are shared with cartilage but are not

necessarily operating in the same manner.

Knock-down of *PLEC* and plectin elicits a response

Our previous data and the results generated here highlight that the OA risk conferring A

allele of rs11780978 correlates with decreased expression of *PLEC* in cartilage and

synovium. We modelled this effect by knocking-down plectin in an immortalised human

MSC cell line. We chose to use this line because: 1) being an MSC line, it is biologically

relevant to joint cells; 2) it expressed *PLEC* at a high level and; 3) it is readily transfectable

and modifiable via CRISPR/Cas9[31]. We transduced it to constitutively express Cas9 and we

then generated a 26bp frame-shift deletion of *PLEC* exon 3, with four replicates for deletion

and four control replicates. The cells were cultured in monolayer for up to 14 days and the

deletion was confirmed by PCR and Sanger sequencing, whilst a near-total plectin knock-out

was observed by immunoblotting using an antibody that binds to the C-terminus of the

protein (Figure 5A-C). The isolated RNA was then subjected to next-generation mRNA

sequencing. This revealed alternate splicing occurring in *PLEC* in the deletion samples, with

exon 3 being completely skipped in a subset of transcripts (Figure 5D). No exon skipping

was seen in the control samples. The level of *PLEC* mRNA was 26% lower in the deletion

samples compared to the controls (log$_2$ fold change of -0.44; FDR *P*=0.001; Supplementary

Table 2). In summary, this data implies that the Cas9-mediated deletion impacted on the

transcription of *PLEC* and on the translation of *PLEC* mRNA into plectin.

We next assessed the effect of this plectin knock-down on the transcriptome. In

addition to *PLEC*, the deletion had a significant effect (FDR *P*<0.05) on the expression of

269 genes (Figure 6A and B, Supplementary Table 2). Pathway analysis revealed

upregulation of processes involving immune regulation and protein post-translation

modification (Figure 6C), and downregulation of Wnt signalling, metabolic pathways and

glycosaminoglycan biosynthesis, amongst others (Figure 6D). In conclusion, this model

system highlighted that depletion of *PLEC* and plectin impacted upon a range of pathways

reported to play a critical role in cartilage biology and OA.

Discussion

The mapping of risk loci for polygenic traits is now a relatively straightforward procedure,

with the next major step in complex trait analysis being the transition from association signal

to functional characterization[32]. For OA, considerable strides have been made in this regard

in recent years[33-35]. Our interest is in the epigenetic dimension of OA genetic risk and

especially the role of DNA methylation. This is the best characterized and most stable

epigenetic modification and modulates gene expression by disrupting the binding of

transcription factors and by facilitating the binding of methyl-binding proteins, which initiate

compaction of chromatin and gene silencing[36]. Last year, we demonstrated that the OA

association signal marked by SNP rs11780978 correlated with methylation of CpGs within

PLEC and with variable expression of the gene[16]. These analyses were chondro-centric and

here, we set out to determine whether these functional effects were more widespread. If so,

an exclusive focus on cartilage as a target tissue of the rs11780978 association signal would

not be merited.

We discovered that the rs11780978-*PLEC* eQTLs, mQTLs and meQTLs that are

active in cartilage are also active in synovium, but with some differences; the genotypic

effect is substantially greater in cartilage for the cg19405177 cluster of CpGs, whilst the

cg14598846 meQTLs act in opposite directions between the two tissues. Further

investigation in additional cohorts should confirm whether both the commonalty and discord

of the effects that we have observed between cartilage and synovium are robust. In fat pad,

we observed the mQTLs but *PLEC* AEI did not correlate with rs11780978 genotype. Instead,

the OA risk-conferring A allele of rs11780978 demonstrated increased and decreased

expression relative to the non-risk allele across the 19 fat pad samples studied, with no

overall significant difference when the data was combined. There are therefore eQTLs

operating on *PLEC* in fat pad, but these are unrelated to rs11780978. These distinctions point

toward regulatory variability in gene expression mediated by the same association between

joint tissues. In blood, the mQTLs were detectable but their genotypic effect was much

weaker in comparison to the three joint tissues. Such variable strengths of mQTLs effects

between cells from different lineages are expected[37]. We were unable to test blood for *PLEC*

eQTLs due to the low expression level of the gene. In summary, we observed the *PLEC*

mQTLs across all four tissues tested with the joint tissues more similar to each other than to

blood. This was exemplified by the mQTL genotypic effects being much stronger in joint

tissues. Since synovium was subject to eQTLs and meQTLs, it too should be investigated as a

target of the rs11780978 association signal.

Finally, we modelled the effect of the rs11780978 risk-conferring allele by knocking

down plectin in a cell line. *PLEC* mRNA was detectable in the RNA-sequencing data but at a

significantly lower level compared to control. The knock-down was very efficient at the

protein level, with a near total absence of plectin observed. The knock-down impacted on a

range of cellular pathways. Of particular note, both the innate and the acquired immune

response showed an upregulation, an effect that has been reported on previously in OA for

both cartilage and synovium[38-39]. Downregulation of Wnt signalling was also of interest, as

this pathway is critical to cartilage homeostasis[33,40]. Furthermore, there is growing evidence

of an interplay between Wnt signalling and inflammation in joint tissues[41]. Our model system

did not include mechanical load, a force which plectin allows cells to respond to[18-20].

Aberrant mechanical load, including only moderate change, can elicit an inflammatory

response that has been termed "mechanoflammation"[42]. Our data suggests that even in the

absence of load, the loss of plectin is eliciting this response. A future model incorporating

load may therefore reveal yet more insight into a potential key role of plectin as a regulator of

inflammation in response to mechanical load and joint tissue injury in OA. This model should

incorporate plectin deficient differentiated joint cells, rather than the undifferentiated MSCs

investigated here, to more accurately recapitulate the articulating joint. Furthermore, and

despite our inability to undertake *PLEC* AEI on blood cells, we should not exclude

circulating immune cells as potential targets of the rs11780978 association signal,

particularly in light of the altered expression of plectin in macrophages in response to

immunomodulatory agents[43].

In conclusion, our investigations have highlighted the interplay between OA genetic

risk, DNA methylation and gene expression, and reveal clear differences in these effects for

tissues from the same diseased joint. Furthermore, by recapitulating the effect of the risk

locus at *PLEC*, we have linked OA genetic susceptibility with a number of cellular pathways,

thus providing mechanistic insight into OA aetiology.

References

1. Glyn-Jones S, Palmer AJ, Agricola R, Price AJ, Vincent TL, Weinans H, et al.

Osteoarthritis. Lancet 2015; 386: 376-387.

2. Kendzerska T, Juni P, King LK, Croxford R, Stanaitis I, Hawker GA. The longitudinal

relationship between hand, hip and knee osteoarthritis and cardiovascular events: a

population-based cohort study. Osteoarthritis Cartilage 2017; 25: 1771-1780.

3. Hawker GA, Croxford R, Bierman AS, Harvey PJ, Ravi B, Stanaitis I, et al. All-cause

mortality and serious cardiovascular events in people with hip and knee osteoarthritis: a

population based cohort study. PLoS One 2014; 9: e91286.

4. arcOGEN Consortium, arcOGEN Collaborators, Zeggini E, Panoutsopoulou K, Southam

L, Rayner NW, et al. Identification of new susceptibility loci for osteoarthritis

(arcOGEN): a genome-wide association study. Lancet 2012; 380: 815-823.

5. Reynard LN. Analysis of genetics and DNA methylation in osteoarthritis: What have we

learnt about the disease? Semin Cell Dev Biol 2017; 62: 57-66.

6. Styrkarsdottir U, Helgason H, Sigurdsson A, Norddahl GL, Agustsdottir AB, Reynard

LN, et al. Whole-genome sequencing identifies rare genotypes in COMP and CHADL

associated with high risk of hip osteoarthritis. Nat Genet 2017; 49: 801-805.

7. Zengini E, Hatzikotoulas K, Tachmazidou I, Steinberg J, Hartwig FP, Southam L, et al.

Genome-wide analyses using UK Biobank data provide insights into the genetic

architecture of osteoarthritis. Nat Genet 2018; 50: 549-558.

8. Styrkarsdottir U, Lund SH, Thorleifsson G, Zink F, Stefansson OA, Sigurdsson JK, et al.

Meta-analysis of Icelandic and UK data sets identifies missense variants in SMO, IL11,

COL11A1 and 13 more new loci associated with osteoarthritis. Nat Genet 2018; 50:

1681-1687.

9. Tachmazidou I, Hatzikotoulas K, Southam L, Esparza-Gordillo J, Haberland V, Zheng J,

et al. Identification of new therapeutic targets for osteoarthritis through genome-wide

analyses of UK Biobank data. Nat Genet 2019; 51: 230-236.

10. Styrkarsdottir U, Stefansson OA, Gunnarsdottir K, Thorleifsson G, Lund SH,

Stefansdottir L, et al. GWAS of bone size yields twelve loci that also affect height, BMD,

osteoarthritis or fractures. Nat Commun 2019; 10: 2054.

11. den Hollander W, Pulyakhina I, Boer C, Bomer N, van der Breggen R, Arindrarto W, et

al. Annotating transcriptional effects of genetic variants in disease-relevant tissue:

transcriptome-wide allelic imbalance in osteoarthritic cartilage. Arthritis Rheumatol

2019; 71: 561-570.

12. Shepherd C, Reese AE, Reynard LN, Loughlin J. Expression analysis of the osteoarthritis

genetic susceptibility mapping to the matrix Gla protein gene MGP. Arthritis Res Ther

2019; 21: 149.

13. Rice SJ, Cheung K, Reynard LN, Loughlin J. Discovery and analysis of methylation

quantitative trait loci (mQTLs) mapping to novel osteoarthritis genetic risk signals.

Osteoarthritis Cartilage 2019; 27: 1545-1556.

14. Reynard LN, Bui C, Syddall CM, Loughlin J. CpG methylation regulates allelic

expression of GDF5 by modulating binding of SP1 and SP3 repressor proteins to the

osteoarthritis susceptibility SNP rs143383. Hum Genet 2014; 133: 1059-1073.

15. Smith E, Shilatifard A. Enhancer biology and enhanceropathies. Nat Struct Mol Biol

2014; 21: 210-219.

16. Rice SJ, Tselepi M, Sorial AK, Aubourg G, Shepherd C, Almarza D, et al. Prioritization

of PLEC and GRINA as osteoarthritis risk genes through the identification and

characterization of novel methylation quantitative trait loci. Arthritis Rheumatol 2019;

71: 1285-1296.

17. Pastinen T. Genome-wide allele-specific analysis: insights into regulatory variation. Nat

Rev Genet 2010; 11: 533-538.

18. Castanon MJ, Walko G, Winter L, Wiche G. Plectin-intermediate filament partnership in

skin, skeletal muscle, and peripheral nerve. Histochem Cell Biol 2013; 140: 33-53.

19. Rezniczek GA, Winter L, Walko G, Wiche G. Functional and genetic analysis of plectin

in skin and muscle. Methods Enzymol 2016; 569: 235-259.

20. Goldmann WH. Intermediate filaments and cellular mechanics. Cell Biol Int 2018; 42:

132-138.

21. Loeser RF, Goldring SR, Scanzello CR, Goldring MB. Osteoarthritis: a disease of the

joint as an organ. Arthritis Rheum 2012; 64: 1697-1707.

22. Gee F, Rushton MD, Loughlin J, Reynard LN. Correlation of the osteoarthritis

susceptibility variants that map to chromosome 20q13 with an expression quantitative

trait locus operating on NCOA3 and with functional variation at the polymorphism

rs116855380. Arthritis Rheumatol 2015; 67: 2923-2932.

23. Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. Salmon provides fast and bias-

aware quantification of transcript expression. Nat Methods 2017; 14: 417-419.

24. Soneson C, Love MI, Robinson MD. Differential analyses for RNA-seq: transcript-level

estimates improve gene-level inferences. F1000Res 2015; 4: 1521.

25. Leek JT. svaseq: removing batch effects and other unwanted noise from sequencing data.

Nucleic Acids Res 2014; 42.

26. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. Genome Biol 2014; 15: 550.

27. Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al.

Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide

expression profiles. Proc Natl Acad Sci U S A 2005; 102: 15545-15550.

28. Croft D, O'Kelly G, Wu G, Haw R, Gillespie M, Matthews L, et al. Reactome: a database

of reactions, pathways and biological processes. Nucleic Acids Res 2011; 39: D691-697.

29. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast

universal RNA-seq aligner. Bioinformatics 2013; 29: 15-21.

30. Thorvaldsdottir H, Robinson JT, Mesirov JP. Integrative Genomics Viewer (IGV): high-

performance genomics data visualization and exploration. Brief Bioinform 2013; 14: 178-

192.

31. Farhang N, Brunger JM, Stover JD, Thakore PI, Lawrence B, Guilak F, et al. CRISPR-

based epigenome editing of cytokine receptors for the promotion of cell survival and

tissue deposition in inflammatory environments. Tissue Eng Part A 2017; 23: 738-749.

32. Gallagher MD, Chen-Plotkin AS. The post-GWAS era: from association to function. Am

J Hum Genet 2018; 102: 717-730.

33. Monteagudo S, Cornelis FMF, Aznar-Lopez C, Yibmantasiri P, Guns LA, Carmeliet P, et

al. DOT1L safeguards cartilage homeostasis and protects against osteoarthritis. Nat

Commun 2017; 8: 15889.

34. Shepherd C, Zhu D, Skelton AJ, Combe J, Threadgold H, Zhu L, et al. Functional

characterization of the osteoarthritis genetic risk residing at ALDH1A2 identifies

rs12915901 as a key target variant. Arthritis Rheumatol 2018; 70: 1577-1587.

35. Klein JC, Keith A, Rice SJ, Shepherd C, Agarwal V, Loughlin J, et al. Functional testing

of thousands of osteoarthritis-associated variants for regulatory activity. Nat Commun

2019; 10: 2434.

36. Zhu H, Wang G, Qian J. Transcription factors as readers and effectors of DNA

methylation. Nat Rev Genet 2016; 17: 551-565.

37. Smith AK, Kilaru V, Kocak M, Almli LM, Mercer KB, Ressler KJ, et al. Methylation

quantitative trait loci (meQTLs) are consistently detected across ancestry, developmental

stage, and tissue type. BMC Genomics 2014; 15: 145.

38. Berenbaum F, van den Berg WB. Inflammation in osteoarthritis: changing views.

Osteoarthritis Cartilage 2015; 23: 1823-1824.

39. Miller RJ, Malfait AM, Miller RE. The innate immune response as a mediator of

osteoarthritis pain. Osteoarthritis Cartilage 2019.

40. Monteagudo S, Lories RJ. Cushioning the cartilage: a canonical Wnt restricting matter.

Nat Rev Rheumatol 2017; 13: 670-681.

41. Claudel M, Jouzeau JY, Cailotto F. Secreted Frizzled-related proteins (sFRPs) in osteo-

articular diseases: much more than simple antagonists of Wnt signaling? FEBS J 2019;

286: 4832-4851.

42. Vincent TL. Mechanoflammation in osteoarthritis pathogenesis. Semin Arthritis Rheum

2019; 49: S36-S38.

43. Kamal AH, Chakrabarty JK, Udden SM, Zaki MH, Chowdhury SM. Inflammatory

proteomic network analysis of statin-treated and lipopolysaccharide-activated

macrophages. Sci Rep 2018; 8: 164.

Author Contributions

504 AKS and JL designed the study. AKS, IMJH, MT and SJR generated the experimental data.

505 AKS, IMJH, KC and EP analysed the data and prepared the manuscript figures. AKS, IMH,

506 and JL prepared the main manuscript text. DJD provided patient samples for analyses. All

507 authors were involved in drafting the article or revising it critically for intellectual content,

508 and all authors approved the final version to be published.

Acknowledgements

511 Dr Colin Shepherd provided input into data interpretation and reviewed the manuscript.

Funding

This work was supported by Versus Arthritis (grant 20771) and by the Medical Research

Council and Versus Arthritis as part of the Centre for Integrated Research into

Musculoskeletal Ageing (CIMA, grant references JXR 10641, MR/P020941/1 and

MR/R502182/1). Funds to purchase the pyrosequencer were provided by The Ruth and

Lionel Jacobson Charitable Trust, which also supported EP. AKS was funded by the Royal

College of Surgeons of Edinburgh Cutner Fellowship and The John George William

Patterson (JGWP) Foundation, IMJH was funded by JGWP.

Competing interest

523 The authors declare that they have no competing interests.

Data Availability

526 All data are available from the corresponding authors upon request. Raw RNA-seq data has

527 been deposited into the NCBI GEO datasets repository with accession number GSE143725.

Legends to figures

Figure 1. Association between rs11780978 genotype and methylation at CpGs cg19405177 and cg14598846 in cartilage, fat pad, synovium and blood DNA. *P* values were calculated using the Kruskal-Wallis test. Horizontal lines and error bars show the mean ± SEM. n = the number of patients providing data per CpG site for each tissue.

Figure 2. Differences in methylation within tissues of the same patient. Methylation at CpGs cg19405177 (**A**) and cg14598846 (**B**) were plotted for those individuals in whom methylation data was generated for two or more tissues. Numbers on the x-axis refer to the anonymised identification number assigned to patients at recruitment. Data is grouped by genotype at rs11780978.

Figure 3. Heat maps displaying the percentage effect of genotype at rs11780978 upon methylation at CpG clusters cg19405177 (**A**) and cg14598846 (**B**) for the four patient tissues. Data analysis was performed using SAS JMP Statistical Data Visualization Software and heat maps produced in PRISM. Patients contributing data to the cg19405177 cluster 1: cartilage, n=81; fat pad, n=39; synovium, n=64; blood, n=43. Patients contributing data to the cg14598846 cluster: cartilage, n=103; fat pad, n=68; synovium, n=79; blood, n=54.

Figure 4. *PLEC* allelic expression imbalance (AEI) in fat-pad (**A**) and synovium (**B**). AEI analysis was conducted using transcript SNP rs11783799. The risk/non-risk allelic ratio is plotted, with a ratio <1 indicating decreased *PLEC* expression from the risk allele. For each patient, the mean of the DNA ratio (black, n=3 technical repeats) and the mean of the cDNA ratio (grey, n=3 technical repeats) is plotted. Numbers on the x-axis refer to the anonymised identification number assigned to patients at recruitment. The values for DNA and cDNA for all patients combined are represented as box plots to the right, in which the lines within the box represent the median, the box represents the 25th to 75th percentile, and the whiskers represent the maximum and minimum values. *P* values were calculated using Wilcoxon's matched pairs signed rank test.

Figure 5. Knock-down of plectin using CRISPR/Cas9. (**A**) The 26bp sequence of *PLEC* exon 3 deleted. (**B**) Representative agarose gel image showing the PCR product for control (non-

deleted) and deletion. (**C**) Representative immunoblot comparing plectin levels in control

versus deletion, with GAPDH as a loading control. (**D**) Sashimi plot showing splicing of

merged *PLEC* deletion (red) and control (blue) samples. Data were viewed on IGV genome

browser. The multiple isoforms of *PLEC* are shown on the bottom of the panel. Read

coverages show the presence of reads within exons. The lines illustrate the splicing between

exons. The values on the lines show the number of junction spanning reads. The merged

deletion track shows the absence of the 26bp region deleted within *PLEC* exon 3 (indicated

by the box).

Figure 6. Differentially expressed genes and pathways. (**A**) Volcano plot illustrating

significantly differentially expressed genes between *PLEC* deletion and control samples. The

x-axis shows the $\log_2$ fold change between condition and the y-axis shows the -$\log_{10}$(FDR).

Genes with an FDR *P*<0.05 are shown in red. (**B**) Heatmap of significantly differentially

expressed genes between *PLEC* deletion and control. Normalized counts were used for

plotting. Hierarchical clustering was used to cluster samples and genes. The heatmap shows

that samples cluster by condition (D0-Del, deletions; D0-Neg, controls). (**C**) Top 25

significantly upregulated Reactome pathways found using GSEA. (**D**) Top 25 significantly

downregulated Reactome pathways found using GSEA. All genes were ranked using signed

$\log_{10}$(*P*-value) and this pre-ranked list used as input into GSEA to query enrichment of

Reactome pathways.

Supplementary Figure 1. The location in *PLEC* of rs11780978 and the analysed CpGs. The

PLEC gene is shown in blue, exons are boxes/vertical lines, and the direction of transcription

is from right to left. The cg19405177 assay captures a total of 8 CpGs, the cg14598846 assay

a total of four. cg19405177 and cg14598846 are marked as red circles, the additional CpGs as

black circles. The physical position of the CpGs on chromosome 8 (hg19 assembly, UCSC

Genome Browser) is shown.

Supplementary Figure 2. Association between rs11780978 genotype and methylation at

CpG clusters cg19405177 (**A**) and cg14598846 (**B**) in cartilage DNA. *P* values were

calculated using the Kruskal-Wallis test. Horizontal lines and error bars show the mean ±

SEM. n = the number of patients providing data per CpG site.

Supplementary Figure 3. Association between rs11780978 genotype and methylation at
CpG clusters cg19405177 (**A**) and cg14598846 (**B**) in fat pad DNA. *P* values were calculated
using the Kruskal-Wallis test. Horizontal lines and error bars show the mean ± SEM. n = the
number of patients providing data per CpG site.

Supplementary Figure 4. Association between rs11780978 genotype and methylation at
CpG clusters cg19405177 (**A**) and cg14598846 (**B**) in synovium DNA. *P* values were
calculated using the Kruskal-Wallis test. Horizontal lines and error bars show the mean ±
SEM. n = the number of patients providing data per CpG site.

Supplementary Figure 5. Association between rs11780978 genotype and methylation at
CpG clusters cg19405177 (**A**) and cg14598846 (**B**) in blood DNA. *P* values were calculated
using the Kruskal-Wallis test. Horizontal lines and error bars show the mean ± SEM. n = the
number of patients providing data per CpG site.

Supplementary Figure 6. Methylation at CpG clusters cg19405177 (**A**) and cg14598846 (**B**)
for the four patient tissues without stratification by rs11780978 genotype. *P* values were
calculated using Kruskal-Wallis with Dunn's multiple comparisons. Horizontal lines and
error bars show the mean ± SEM. n = the number of patients providing data per tissue and at
each CpG site. *$P \leq$ 0.05; **$P \leq$ 0.01; ***$P \leq$ 0.001 ****$P \leq$ 0.0001.

Supplementary Figure 7. Expression of *PLEC* in tissue samples from OA patients. *PLEC*
mRNA levels were measured by qPCR in cartilage (n=10), fat pad (n=9), synovium (n=10)
and blood (n=10). Horizontal lines and error bars show the mean ± SEM. *P* values were
calculated using a Mann-Whitney 2-tailed exact test. **$P \leq$ 0.01; ***$P \leq$ 0.001

Supplementary Figure 8. The six patients with AEI data for fat pad (FP) and synovium
(SYN). The risk/non-risk allelic ratio is plotted, with a ratio <1 indicating decreased *PLEC*
expression from the risk allele. For each patient, the mean of the DNA ratio (black, n=3
technical repeats) and the mean of the cDNA ratio (grey, n=3 technical repeats) is plotted.
Numbers on the x-axis refer to the anonymised identification number assigned to patients at
recruitment.

629 **Supplementary Figure 9.** Methylation expression quantitative trait locus analyses in fat pad.

630 *PLEC* $\log_2$ allelic expression imbalance (AEI) ratios were plotted against methylation at all

631 CpGs in the cg19405177 (**A**) and cg14598846 (**B**) clusters. The square of the correlation

632 coefficient (r^2) and *P* values were calculated by linear regression analysis using a standard

633 least squares model.

635 **Supplementary Figure 10.** Methylation expression quantitative trait locus analyses in

636 synovium. *PLEC* $\log_2$ allelic expression imbalance (AEI) ratios were plotted against

637 methylation at all CpGs in the cg19405177 (**A**) and cg14598846 (**B**) clusters. The square of

638 the correlation coefficient (r^2) and *P* values were calculated by linear regression analysis

639 using a standard least squares model.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

A

26bp deletion

5' CTCCAGCAGGGAGATGAGGTTGTGGCCATCGCGGAG 3'   Control
5' CTCCA   CGGAG 3'   Deletion

B

DNA

C

Plectin

GAPDH

Protein

D

Exon 3

Deletion

Control

PLEC

Figure 6

A

B

C

Upregulated Pathways

PLEC deletion vs control

D

Downregulated Pathways