Integration of polygenic risk scores with modifiable risk factors improves risk prediction: results from a pan-cancer analysis

Linda Kachuri [1], Rebecca E. Graff [1], Karl Smith-Byrne[2], Travis J. Meyers[1], Sara R. Rashkin[1], Elad Ziv[3,4,5], John S. Witte[4,5,6] *, Mattias Johansson[2] *

* Co-senior / corresponding authors

Affiliations:

1. Department of Epidemiology and Biostatistics, University of California, San Francisco, San Francisco, USA
2. Genetic Epidemiology Group, Section of Genetics, International Agency for Research on Cancer, Lyon, France
3. Department of Medicine, University of California, San Francisco, San Francisco, USA
4. Helen Diller Family Comprehensive Cancer Center, University of California, San Francisco, San Francisco, USA
5. Institute for Human Genetics, University of California, San Francisco, San Francisco, USA
6. Department of Urology, University of California, San Francisco, San Francisco, USA

Corresponding Authors:

John S. Witte

Department of Epidemiology and Biostatistics
University of California, San Francisco
1450 3rd Street, Room 388, San Francisco, CA 94158
Email: jwitte@ucsf.edu

Mattias Johansson

Genetic Epidemiology Group, Section of Genetics
International Agency for Research on Cancer
150 Cours Albert Thomas, 69372 Lyon, France
Email: JohanssonM@iarc.fr

ABSTRACT

Cancer risk is determined by a complex interplay of genetic and modifiable risk factors. Combining individual germline risk variants into polygenic risk scores (PRS) creates a personalized genetic susceptibility profile that can be leveraged for disease prediction. Using data from the UK Biobank cohort (413,753 individuals; 22,755 incident cases), we systematically quantify the added predictive value of augmenting conventional cancer risk factors with PRS for 16 cancer types. Our results indicate that incorporating PRS in addition to family history of cancer and modifiable risk factors improves prediction accuracy, but the magnitude of incremental improvement varies substantially between cancers. We also demonstrate the utility of PRS for risk stratification. Individuals with high genetic risk (PRS≥80th percentile) have significantly divergent 5-year absolute risk trajectories across strata based on family history and modifiable risk factors. Finally, we estimate that high genetic risk accounts for 4.0% to 30.3% of new cancer cases, which exceeds the impact of many lifestyle-related risk factors. In summary, we provide novel quantitative data illustrating the importance of integrating PRS into personalized cancer risk assessment.

INTRODUCTION

Cancer susceptibility is inherently complex, but it is well-accepted that heritable genetic factors and modifiable exposures contribute to cancer development. While our knowledge of causal modifiable risk factors has gradually evolved over the past decades, genome-wide association studies (GWAS) have rapidly produced a wealth of germline genetic risk variants for different cancers. These studies have shed light on genetic mechanisms of cancer susceptibility, however, the public health impact of GWAS findings has been modest. In response, GWAS results have been leveraged to create polygenic risk scores (PRS) by combining weighted genotypes for risk alleles into a single, integrated measure of an individual's genetic predisposition to a specific phenotypic profile. Such genetic risk scores are not designed to reflect the complexity of molecular susceptibility mechanisms, but they are highly amenable to phenotypic prediction.

Multiple studies have demonstrated that PRS can generate informative predictions for heritable traits[1,2] and diseases[3,4], prompting many to advocate for increased integration of genetic risk scores into clinical practice[5,6]. An important step towards realizing the promise of PRS in precision medicine lies in systematically assessing the added value of genetic information in comparison to conventional risk factors and examining how it affects lifetime risk trajectories[6]. The recent development of large, prospective cohorts with both genome-wide genotyping and deep phenotyping data, such as the UK Biobank[7], provide an opportunity for integrative analyses of genetic variation and modifiable risk factors. In addition to evaluating PRS predictive performance, these data also provide a unique opportunity to answer etiological questions about the relative contribution of genetic and modifiable risk factors to cancer susceptibility.

Our overarching aim was to quantify the relative contribution of common, low-penetrance risk variants to cancer risk prediction and overall disease susceptibility. To address these aims, we assembled PRS for 16 cancer types, based on results from previously published GWAS, and applied them to 413,870 individuals in the UK Biobank (UKB) cohort. First, we assessed the degree to which PRS can improve risk prediction and stratification based on established cancer risk factors, such as family history and modifiable health-related characteristics. Next, we estimated the proportion of cancer cases at the population-level that can be attributed to high genetic susceptibility, captured by the PRS, and compared this to modifiable determinants of cancer.

RESULTS

Characteristics of the UKB study population are presented in **Supplementary Table 1**. Over the course of the follow-up period a total of 22,755 incident cancers were diagnosed in 413,753 individuals, after excluding participants outside of the age enrollment criteria and those who withdrew consent after enrollment. Established cancer risk factors (listed in **Supplementary Table 2**) exhibited associations of expected magnitude and direction with each cancer (**Supplementary Table 3**). Family history of cancer in first-degree relatives, at the corresponding site, conferred a significantly higher risk of prostate (HR=1.84, 95% CI: 1.68-2.00, p=9.1×10^{-46}), breast (HR=1.56, 1.44-1.69, p=3.0×10^{-29}), lung (HR=1.61, 1.43-1.81, p=7.4×10^{-15}), and colorectal (HR=1.26, 1.14-1.40, p=1.2×10^{-5}) cancers. Metrics of tobacco use, such as smoking status, intensity, and duration were positively associated with risks of lung, colorectal, bladder, kidney, pancreatic, and oral cavity/oropharyngeal cancers. Weekly alcohol intake was associated with higher risks of breast (HR per 70 grams = 1.04, p=2.3×10^{-5}), colorectal (HR=1.04, p=5.9×10^{-9}), and oral cavity/pharyngeal (HR=1.05, p=3.0×10^{-10}) cancers. Adiposity was associated with cancer risk at multiple sites, including endometrium (BMI: HR per 1-unit = 1.09, 1.08-1.10, p=1.6×10^{-49}), colon/rectum (waist-to-hip ratio: HR per 10% increase = 1.17, 1.11-1.24, p=2.2×10^{-8}), and kidney (BMI: HR=1.04, 1.02-1.05, p=1.7×10^{-6}). Particulate matter ($PM_{2.5}$) was associated with lung cancer risk[8] ($PM_{2.5}$: HR per 1 micro-g/m^3 = 1.10, 1.05-1.15, p=1.9×10^{-5}) in the model that included smoking status and intensity.

All PRS associations with the target cancer reached at least nominal statistical significance (**Figure 1; Supplementary Table 4**). We considered three PRS approaches (see Methods for details): standard weights corresponding to reported risk allele effect sizes (PRS_β); unweighted sum of risk alleles (PRS_{unw}); inverse variance weights that incorporate the standard error of the risk effect size (PRS_{IV}). The latter approach resulted in stronger or equivalent (HR $\pm$ 0.01) associations for most cancers, except Non-Hodgkin lymphoma (NHL). Compared to standard PRS_β, substantial differences were observed for prostate (PRS_{IV}: HR=1.77, P=4.3×10^{-366} vs. PRS_β: HR=1.39, P=2.0×10^{-105}), colon/rectum (PRS_{IV}: HR=1.48, P=1.8×10^{-94} vs. PRS_β: HR=1.32, P=5.5×10^{-50}), leukemia (PRS_{IV}: HR=1.70, P=6.3×10^{-23} vs. PRS_β: HR=1.45, P=8.0×10^{-13}), and thyroid (PRS_{IV}: HR=1.75, P=1.9×10^{-15} vs. PRS_β: HR=1.57, P=5.7×10^{-10}). All subsequent analyses use PRS_{IV} since this approach appears to improve PRS performance by appropriately downweighing the contribution of variants with less precisely estimated effects.

Improvement in risk prediction

The predictive performance of each risk model was evaluated based on its ability to accurately estimate risk (calibration) and distinguish cancer cases from cancer-free individuals (discrimination). All cancer-specific risk models were well-calibrated (Goodness of fit p>0.05; **Supplementary Figure 1**). Model discrimination was assessed by Harrell's C-index, estimated as a weighted mean between 1 and 5 years of follow-up time. For completeness, we also report the AUC at 5 years of follow-up time[9]. Proportionality violations (p<0.05) were detected for age in the breast cancer model and PRS_{IV} for cervical cancer. For breast cancer this was resolved by incorporating an interaction term with follow-up time. As a sensitivity analysis for cervical cancer we modelled a time-varying PRS effect (**Supplementary Figure 2**).

The C-index reached 0.60 with age and/or sex, for all cancers except for breast and thyroid (**Supplementary Table 5**). For cancers with available information on family history of cancer at the same site (prostate, breast, colon/rectum, and lung), incorporating this had a modest impact on the C-index (ΔC<0.01). In fact, replacing family history with the PRS resulted in an improvement in discrimination for prostate (C=0.763, ΔC=0.047), breast (C=0.618, ΔC=0.060), and colorectal (C=0.708, ΔC=0.029), but not lung (C=0.711, ΔC=-0.002) cancers.

Next, we assessed the change in the C-index (ΔC) after incorporating the PRS into prediction models with all available risk factors for each cancer (**Figure 2; Supplementary Table 5**). The resulting improvement in prediction performance was variable. The largest increases in the C-index were observed for cancer sites with few available predictors, such as testes (C_{PRS}=0.766, ΔC=0.138), thyroid (C_{PRS}=0.692, ΔC=0.099), prostate (C_{PRS}=0.768, ΔC=0.051) and lymphocytic leukemia (C_{PRS}=0.756, ΔC=0.061). However, adding the PRS also improved prediction accuracy for melanoma (C_{PRS}=0.664, ΔC=0.042), breast (C_{PRS}=0.631, ΔC=0.060), and colorectal (C_{PRS}=0.716, ΔC=0.030) cancers, which have multiple environmental risk factors. The highest overall C-index was observed for lung (C_{PRS}=0.849) and bladder (C_{PRS}=0.814) cancers, which was primarily attributed to non-genetic predictors (C without PRS: lung = 0.846; bladder = 0.808). Changes in the AUC at 5 years of follow-up were of similar magnitude (**Supplementary Table 5**).

As a complementary metric of model performance, Royston's R^2 was calculated to quantify the variation in the time-to-event outcome captured by each risk model[10]. Across all 16 sites, the median change in R^2 (ΔR^2) was 0.066. Large improvements, defined as ΔR^2 >0.10, were observed for cancers of the breast (R^2_{PRS}=0.146; ΔR^2=0.103), pancreas (R^2_{PRS}=0.439; ΔR^2=0.103), leukemia (R^2_{PRS}=0.415; ΔR^2=0.160), prostate (R^2_{PRS}=0.510; ΔR^2 =0.161), thyroid (R^2_{PRS}=0.310; ΔR^2 =0.230), and testis (R^2_{PRS}=0.605; ΔR^2 =0.421). These results parallel the trend in improvement observed based on C-index and AUC.

For 15 out of 16 cancers, incorporating the PRS resulted in significant improvement in reclassification, as indicated by positive percentile-based net reclassification index (NRI)[11] values with 95% bootstrapped confidence intervals excluding 0 (**Supplementary Table 6**). The overall NRI was primarily driven by the event NRI (NRI_e), which is the increase in the proportion of cancer cases reclassified to a higher risk group. Positive NRI_e values >0.25 were observed for prostate, thyroid, breast, testicular, leukemia, melanoma, and colorectal cancers. The largest reclassification improvement in non-event NRI (NRI_{ne}) observed for the lung PRS (NRI_{ne}=0.015) and breast PRS (NRI_{ne}=0.012). Four cancers (testes, leukemia, kidney, oral cavity/pharynx) had significantly negative NRI_{ne} values indicating that adding the PRS decreased classification accuracy in cancer-free individuals.

Refinement of risk stratification

The ability of the PRS to refine risk estimates was assessed by examining 5-year absolute risk trajectories as a function of age, across strata defined by percentiles of PRS (high risk ≥80%, average: >20% to <80%, low risk: ≤20%) and family history of cancer (**Figure 3**). Significantly diverging risk trajectories, overall and at age 60, were observed for prostate ($P<4.5\times10^{-25}$), breast ($P<9.3\times10^{-36}$), colorectal ($P<2.0\times10^{-21}$), and lung cancers ($P<0.031$). For all cancers except lung, risk stratification was primarily driven by PRS. For instance, 60-year-old men with a high PRS but no family history of prostate cancer had a higher mean 5-year disease risk (4.74%) compared to men with a positive family history and an average PRS (3.66%). For lung cancer, on the other hand, participants with a positive family history had higher average 5-year risks, even with a low PRS (0.54%), compared to those without (high PRS: 0.46%; low PRS: 0.29%). There was evidence of interaction between the PRS and family history of cancer for prostate ($P = 9.0\times10^{-128}$), breast ($P = 1.7\times10^{-104}$), colorectal ($P = 8.7\times10^{-14}$) cancers (**Supplementary Table 7**). For lung cancer the interaction with family history was limited to the high PRS group ($P = 5.9\times10^{-3}$).

We also compared 5-year risk projections across strata of PRS and modifiable risk factors. Effects of multiple risk factors were combined into a single score by generating summary linear predictors for each cancer (see Methods for details). For several common cancers, individuals with a high PRS were predicted to have higher cancer risk, even modifiable risk factor scores below the median (**Figure 4**). PRS achieved significant risk stratification for breast cancer (pre-menopausal: $P<5.9\times10^{-12}$; post-menopausal: $P<4.3\times10^{-50}$), colorectal cancer ($P<1.8\times10^{-42}$), and melanoma ($P<4.6\times10^{-105}$) (**Figure 4**). The same pattern of stratification was observed for NHL, leukemia, pancreatic, thyroid, and testicular cancers (**Supplementary Figure 3**). For other phenotypes, lifestyle-related risk factors had a stronger overall influence on risk trajectories than PRS (**Figure 5**). However, the stratifying by levels of PRS still resulted in significantly diverging risk projections for several cancers (lung: $P<1.1\times10^{-13}$; oral cavity/pharynx: $P<1.2\times10^{-12}$; kidney: $P<1.1\times10^{-13}$). For bladder cancer, the risk trajectories for high PRS/reduced modifiable risk and low PRS/high modifiable risk were overlapping ($P=0.98$).

There was evidence of larger than additive risk differences, at age 60 between elevated modifiable risk factor profiles and all ordinal PRS categories for melanoma ($P=3.3\times10^{-122}$), post-menopausal breast ($P=1.3\times10^{-21}$), colorectal ($P=1.3\times10^{-208}$), lung ($P=1.1\times10^{-37}$), bladder ($P=1.5\times10^{-50}$), kidney ($P=5.5\times10^{-29}$), and oral cavity/pharynx cancers ($P=5.2\times10^{-11}$) (**Supplementary Table 7**). For pre-menopausal breast cancer the interaction was limited to women in the high PRS group ($P=4.4\times10^{-4}$).

Quantifying population-level impact

Population attributable fractions (PAF) were used to summarize the relative contribution of genetic susceptibility and modifiable risk factors to cancer risk at the population level. In order to allow comparisons between PAF estimates, the PRS and modifiable risk score distributions were both dichotomized at ≥80[th]

percentile. All risk factors nominally contributed ($P<0.05$) to cancer incidence (**Figure 6; Supplementary Table 8**), with the exception of the PRS for oral cavity/pharynx cancer ($P=0.78$) and $PM_{2.5}$ for lung cancer in never smokers ($P=0.44$).

PAF for high genetic risk exceeded the contribution of modifiable exposures for several cancers, such as thyroid ($PAF_{PRS}=0.268$, $P=1.7\times10^{-9}$), prostate ($PAF_{PRS}=0.232$, $P=5.5\times10^{-158}$), colon/rectum ($PAF_{PRS}=0.167$, $P=9.2\times10^{-50}$), breast ($PAF_{PRS}=0.166$, $P=2.6\times10^{-85}$), and melanoma ($PAF_{PRS}=0.139$, $P=1.3\times10^{-23}$). For testicular cancer ($PAF_{PRS}=0.303$, $P=4.5\times10^{-4}$), leukemia ($PAF_{PRS}=0.269$, $P=4.5\times10^{-4}$), lung cancer in never smokers ($PAF_{PRS}=0.077$, $P=0.045$), and NHL ($PAF_{PRS}=0.053$, $P=1.9\times10^{-3}$), PRS was the only significant risk factor other than demographic factors. Cancers for which modifiable risk factors had a substantially larger impact on disease burden than PRS included oral cavity/pharynx ($PAF_{mod}=0.310$ vs. $PAF_{PRS}=0.006$), lung ($AF_{mod}=0.636$ vs. $PAF_{PRS}=0.040$), endometrium ($PAF_{mod}=0.353$ vs. $PAF_{PRS}=0.043$), kidney ($PAF_{mod}=0.210$ vs. $PAF_{PRS}=0.046$), and bladder cancers ($PAF_{mod}=0.189$ vs. $PAF_{PRS}=0.085$). For other sites, such as pancreas ($PAF_{mod}=0.118$ vs. $PAF_{PRS}=0.133$) and ovary ($PAF_{mod}=0.100$ vs. $PAF_{PRS}=0.082$), the contribution of PRS and modifiable risk factors were more balanced.

DISCUSSION

Cancer is a multifactorial disease with a complex web of etiological factors, from macro-level determinants, such as health policy, to individual-level characteristics, such as health-related behaviors and heritable genetic profiles. Heritable and modifiable risk factors act in concert to influence cancer development, but their relative contributions to disease risk are rarely compared directly in the same population. In this study we provide new insight into the potential utility of PRS for cancer risk prediction and provide insight into the relative of contribution of genetic and modifiable risk factors to cancer incidence at population level.

Our first major finding is that cancer-specific PRS comprised of lead GWAS variants improve risk prediction for all 16 cancers examined. However, the magnitude of the resulting improvement in prediction varies substantially between sites. In evaluating the added predictive value of the PRS it is important to keep in mind that achieving the same incremental increase in the C-index/AUC is more difficult when the baseline model already performs well[12]. This was applicable to most cancers, where age and/or sex alone achieved non-trivial risk discrimination (C-index/AUC>0.60). Expanding the set of predictors to include modifiable risk factors further improved discrimination, as previously shown[13]. By adding the PRS to the most comprehensive risk factor models facilitated by our data, we adopted a conservative approach for quantifying its added predictive value, which provides an informative benchmark for future efforts seeking to incorporate genetic predisposition in cancer risk assessment.

Cancer sites for which the PRS resulted in the largest gains in prediction performance included prostate, testicular, and thyroid cancers, as well as leukemia, and melanoma. This is consistent with high heritability estimates reported for these cancers in twin studies[14] and our analyses in the UK Biobank[15]. Modelling the PRS in addition to established risk factors yielded very modest improvements in risk discrimination for cancers of the lung, endometrium, bladder, oral cavity/pharynx, and kidney. These cancers have strong environmental risk factors, such as smoking, alcohol consumption, obesity, and HPV infection, some of which were captured in our analysis. Limited predictive ability for cervical and endometrial cancers may also be due to a low number of variants included in the PRS (9 and 10, respectively). The association of the lung cancer PRS with cigarettes per day[16] may have diminished its apparent predictive value when added to a model with smoking status and intensity, which already achieved an AUC>0.80 making difficult to elicit further improvement. Furthermore, PRS may be

particularly relevant for assessing lung cancer risk in never smokers, since other risk factors have a limited impact in this population.

Few pan-cancer PRS studies have been conducted in prospective cohorts and none have considered the breadth of modifiable risk factors that we evaluated. Shi et al.[17] tested 11 cancer PRS in cases from The Cancer Genome Atlas and controls from the Electronic Medical Records and Genomics Network. This analysis was limited by fewer risk variants in each PRS, as well as potential for bias due to selection of cases and controls from different populations. A phenome-wide analysis in the Michigan Genomics Initiative cohort by Fritsche et al.[18] examined PRS for 12 cancers and reported similar associations for the target phenotype. However, risk stratification was not formally evaluated. Considering cancer-specific studies, the PRS presented here achieved superior prediction performance for some cancers[19-22], but not others[23,24]. For pancreatic cancer[25] and melanoma[26], our results are consistent with previous analyses using PRS of similar composition. Generally, comparison of prediction performance is complicated by differences in PRS content, population characteristics, and inclusion of different non-genetic predictors. Outside the cancer literature, our conclusions align with a recent study of ischemic stroke, which demonstrated that the PRS is similarly or more predictive than multiple established risk factors, including family history[27].

Our second major finding advances the idea of using germline genetic information to refine individual risk estimates. We show that incorporating PRS improves risk stratification provided by conventional risk factors alone, as illustrated by significantly diverging 5-year risk projections within strata based on family history or modifiable risk factors. For certain cancers, including some with strong environmental risk factors, such as melanoma, breast, colorectal, and pancreatic cancers, PRS was the primary determinant of risk stratification. For others, such as lung and bladder cancers, modifiable risk factors had a stronger impact on 5-year risk trajectories. A consistent finding for all cancers was that individuals in the top 20% of the PRS distribution with an unfavorable modifiable risk factor profile had the highest level of risk, with evidence that the effects of PRS and modifiable risk factors may be synergistic. Taken together, these findings highlight the potential for attenuating high genetic risk by adhering to a healthier lifestyle. Similar risk stratification results based on genetic and modifiable risk factors have also been reported for coronary disease[28] and Alzheimer's[29].

In addition to evaluating predictive performance and risk stratification, our work demonstrates the relevance of common genetic risk variants at the population level. High genetic risk (PRS≥80th percentile) explained between 4.0% and 30.3% of new cancer cases, and for many phenotypes this exceeded PAF estimates for modifiable risk factors or family history. The contribution of genetic variation to disease risk is typically conveyed by heritability, which is an informative metric, although not easily translated into a measure of disease burden useful in a public health context. Recent work on cancer PAF in the UK[30] and a series of publications from the ComPARe initiative in Canada[31,32] examined wide range of modifiable risk factors. Despite providing useful data, these studies overlook the contribution of genetic susceptibility. Our work addresses these limitations by providing a more complete perspective on the determinants of cancer and potential impact of future prevention policies.

In evaluating the contributions of our study, several limitations should be acknowledged. First, we did not account for the impact of workplace exposures and socio-economic determinants of health, thereby underestimating the role of non-genetic risk factors. We also lacked data on several known carcinogens, such as ionizing radiation, and clinical biomarkers, such as prostate-specific antigen, thus limiting the extent to which our results inform risk discrimination for certain cancers. Information on family history was also not available for all cancer types. Second, since the UK Biobank cohort is unrepresentative of the general UK population due to low participation and resulting healthy volunteer bias[33], we may have

underestimated PAFs for modifiable risk factors. Finally, the models presented here are calibrated to the UKB population and we urge caution in extrapolating prediction performance and absolute risk projections to other populations. Since our analytic sample is restricted to individuals of predominantly European ancestry, this limits the applicability of our findings to diverse populations.

This work has several important strengths. The UK Biobank resource enabled us to simultaneously evaluate heritable and modifiable cancer risk factors in a population-based cohort with uniform deep phenotyping. We report a series of metrics that comprehensively characterize different dimensions of predictive performance that can be improved by incorporating genetic risk scores. While our results are promising, we anticipate that the performance of the PRS reported here may be enhanced by adopting less stringent p-value thresholding to include additional risk variants, optimizing subtype-specific weights, and implementing more sophisticated PRS models that incorporate linkage disequilibrium structure, functional annotations, or SNP interactions. Some of these strategies are already being successfully implemented[4,23]. We also provide insight into PRS modelling by showing that accounting for the variance in risk allele effect sizes improves PRS performance. This approach may be particularly advantageous for PRS derived from multiple sources rather than a single GWAS. Throughout this study we consider a relatively lenient definition of high genetic risk, corresponding to the top 20% of the PRS distribution. Exploring other cut-points will be informative, however, our results are valuable for demonstrating that the utility of PRS for stratification is not limited to the most extreme ends of the genetic susceptibility spectrum. This threshold is also compelling from a population-health perspective, as it allows us to quantify the proportion of cases attributed to a risk factor with a 20% prevalence.

Genetic risk scores have the potential to become a powerful tool for precision health, but only if the resulting information can be understood and acted on appropriately. One important consideration is the accuracy and stability of PRS-based risk classifications, especially at clinically actionable risk thresholds that exist for certain cancers. For instance, there are established screening programs for breast and colorectal cancers, and increasing evidence supporting the effectiveness of low-dose computed tomography for lung cancer screening[34,35]. For these cancers PRS could be used to adjust the optimal age for screening initiation and/or intensity. However, to justify this, studies are needed to demonstrate the benefit of using PRS to supplement conventional screening criteria. Such trials are already underway for breast cancer, where genetic risk scores are being incorporated to personalize risk-based screening[36]. For other cancers, such as prostate, screening remains controversial and PRS may prove useful in identifying a subset of high-risk individuals who may benefit the most from screening.

Another area where PRS may prove useful is for prioritizing individuals for targeted health and lifestyle-related interventions. In support of this, our study demonstrates that those with the highest levels of genetic risk, based on the PRS, may also experience larger decreases in risk from shifting to a healthier lifestyle. However, there is also accumulating evidence that simply reporting genetic risk information to individuals does not induce behavior change that could lead to meaningful reductions in risk[37]. Therefore, progress in our ability to construct and apply PRS to identify high-risk individuals must be also accompanied by the development of effective behavioral interventions that can be implemented in response to high disease risk, in addition to early detection and screening protocols.

Ultimately, the impact of PRS on clinical decision-making should be carefully evaluated in randomized trials prior to deployment in healthcare settings. By demonstrating cancer-specific improvements in risk prediction, as well as the substantial proportion of cancer incidence that is captured by known genetic susceptibility variants, we provide novel evidence that contextualizes the potential for using genetic information to improve cancer outcomes.

METHODS

Study Population

The UK Biobank (UKB) is a population-based prospective cohort of individuals aged 40 to 69 years, enrolled between 2006 and 2010. All participants completed extensive questionnaires, in-person physical assessments, and provided blood samples for DNA extraction and genotyping[7]. Health-related outcomes were ascertained via individual record linkage to national cancer and mortality registries and hospital in-patient encounters[7]. Details of the quality control and phenotyping procedures for this dataset have been previously described[15,16]. Briefly, individuals with at least one recorded incident diagnosis of a borderline, in situ, or malignant primary cancer were defined as cases. Cancer diagnoses coded by International Classification of Diseases (ICD)-9 or ICD-10 codes were converted into ICD-O-3 codes using the SEER site recode paradigm in order to classify cancers by organ site.

Participants were genotyped on the UKB Affymetrix Axiom array (89%) or the UK BiLEVE array (11%)[7]. Genotype imputation was performed using the Haplotype Reference Consortium as the main reference panel, supplemented with the UK10K and 1000 Genomes phase 3 reference panels[7]. Genetic ancestry principal components (PCs) were computed using fastPCA[38] based on a set of 407,219 unrelated samples and 147,604 genetic markers[7]. All analyses were restricted to self-reported European ancestry individuals with concordant self-reported and genetically inferred sex. To further minimize potential for population stratification, we excluded individuals with values for either of the first two ancestry PCs outside of five standard deviations of the population mean. Based on a subset of genotyped autosomal variants with minor allele frequency (MAF)≥0.01 and genotype call rate ≥97%, we excluded samples with call rates <97% and/or heterozygosity more than five standard deviations from the mean of the population. With the same subset of SNPs, we used KING[38] to estimate relatedness among the samples. We excluded one individual from each pair of first-degree relatives, preferentially retaining individuals to maximize the number of cancer cases remaining, resulting in a total of 413,870 UKB participants.

Polygenic Risk Scores

In order to derive polygenic risk scores (PRS) for each of the 16 cancers, we extracted previously associated variants by searching the National Human Genome Research Institute (NHGRI)-European Bioinformatics Institute (EBI) Catalog of published GWAS. For every eligible GWAS, both the original primary manuscript and supplemental materials were reviewed. Additional relevant studies were identified by examining the reference section of each article and via PubMed searches of other studies in which each article had been cited. We abstracted all autosomal variants with minor allele frequency MAF≥ 0.01 and $P<5\times10^{-8}$ identified in populations of at least 70% European ancestry and published by June 2018, with the exception of one colorectal cancer GWAS[39] (published in December 2018). For inclusion in the PRS we preferentially selected independent SNPs (LD r^2<0.3) with the highest imputation score and we excluded SNPs with allele mismatches or MAF differences >0.10 relative to the 1000 Genomes reference population, and palindromic SNPs with MAF≥0.45. For associations reported in more than one study of the same ancestry and phenotype, we selected the one with the most information (i.e., which reported the risk allele and effect estimate) with the smallest p-value. Further details of the PRS development approach, including a list of source studies, is described by Graff et al[16].

We considered three approaches for combining risk variants in the PRS. First, we used standard PRS weights, corresponding to the log odds ratio (β) for each risk allele:

$$PRS_{\beta} = \beta_1 \times SNP_1 + \beta_2 \times SNP_2 + \cdots + \beta_n \times SNP_n$$

We compared this to an unweighted score corresponding to the sum of the risk alleles, which is equivalent to assigning all variants an equal weight of 1:

$$PRS_{unw} = SNP_1 + SNP_2 + \cdots + SNP_n$$

Lastly, we applied inverse variance (IV) weights that incorporated the standard error (SE) of the SNP log(OR) to account for uncertainty in risk allele effect sizes and downweigh the contribution of variants with less precisely estimated associations (weights provided in **Supplementary Data 1**):

$$PRS_{IV} = \frac{\beta_1}{SE(\beta_1)} \times SNP_1 + \frac{\beta_2}{SE(\beta_2)} \times SNP_2 + \cdots + \frac{\beta_n}{SE(\beta_n)} \times SNP_n$$

Each PRS was standardized across the entire analytic cohort to have a mean of 0 and standard deviation (SD) of 1.

Statistical Analysis

Development of risk models for each cancer

Cancer-specific prediction models consisting of four classes of risk factors were developed: i) demographic factors (age and sex); ii) family history of cancer in first-degree relatives; iii) modifiable risk factors; and iv) genetic susceptibility, represented by the PRS. Family history of cancer was derived based on self-reported illnesses in non-adopted first-degree relatives, which only listed cancers of the prostate, breast, bowel, or lung. In addition to these four cancer sites, family history of breast cancer was included as a predictor for ovarian cancer[40,41]. Models for pancreatic cancer included a composite variable for family history of cancer at any of these four sites[42,43]. Selection of modifiable risk factors was informed by literature review and reports, such as the European Code Against Cancer[44], with an emphasis on risk factors that are likely to have a causal role. Final models included established environmental and lifestyle-related characteristics that were collected for the entire UK Biobank cohort (**Supplementary Table 1**).

Cause-specific Cox proportional-hazard models were used to estimate the hazard ratios (HR) and corresponding 95% confidence intervals (CI) for genetic and lifestyle factors associated with each incident cancer. Death from any cause, other than cancer site-specific mortality, was treated as a competing event. Information on primary and contributing causes of death was used to identify cancer site-specific mortality. Follow-up time was calculated from the date of enrollment to the date of cancer diagnosis, date of death, or end of follow-up (January 1, 2015). For each cancer, individuals with a past or prevalent cancer diagnosis at that same site were excluded from the analysis, while individuals diagnosed with cancers at other sites were retained in the population. All models including the PRS were also adjusted for genotyping array and the first 15 genetic ancestry PCs. For the PRS, HR estimates correspond to 1 SD increase in the standardized genetic score.

Risk model evaluation

The predictive performance of each risk model was evaluated based on its ability to accurately estimate risk (calibration) and distinguish cancer cases from cancer-free individuals (discrimination). Calibration was assessed with a Hosmer-Lemeshow goodness-of-fit statistic modified for time-to-event outcomes[45], and by plotting the expected event status against the observed event probability[46] across risk deciles. For rarer cancers calibration was assessed across quantiles of risk to ensure a minimum of 5 cases per group. Violation of the proportionality of hazards assumption was assessed by examining the association between standardized Schoenfeld residuals and time.

We evaluated nested models starting with the most minimal set of predictors, such as demographic factors, followed by models including family history of cancer and modifiable risk factors, and finally models incorporating the PRS. Risk discrimination was assessed based on Harrell's C-index, calculated as a weighted average between 1 and 5 years of follow-up time, and Area Under the Curve (AUC) at 5 years. We also report pseudo R^2 coefficients based on Royston's measure of explained variation for survival models[10]. Percentile-based net reclassification improvement (NRI) index[11] was used to quantify improvements in reclassification. NRI summarizes the proportion of appropriate directional changes in predicted risks. Any upward movement in risk categories for cases indicates improved classification, and any downward movement implies worse reclassification. The opposite is expected for non-cases:

$$NRI_e = \frac{P(event|up) \times n_U - P(event|down) \times n_D}{n \times P(event)}$$

$$NRI_{ne} = \frac{(1 - P(event|down)) \times n_D - (1 - P(event|up)) \times n_U}{n \times (1 - P(event))}$$

Where n_U is the number of individuals up-classified and n_D is the number down-classified. Overall NRI is the sum of the NRI in cases and NRI in non-cases: $NRI = NRI_e + NRI_{ne}$. Bootstrapped confidence intervals were obtained based on 1000 replicates.

Risk stratification: genetic vs. modifiable factors

For each individual, we estimated the 5-year absolute risk of being diagnosed with a specific cancer using the formula of Benichou & Gail[47], as implemented by Ozenne et al[48]. Absolute risk trajectories were examined as a function of age across strata defined by genetic and modifiable risk profiles, as well as family history. Individuals in the top 20% of the PRS distribution (PRS≥80th percentile) for a given cancer were classified has having high genetic risk, those in the bottom 20% (PRS≤20th percentile) were classified as low risk, and the middle category (>20th to <80th percentile) classified as average genetic risk.

Modifiable risk factors were summarized by generating summary linear predictors (predicted log-hazard ratios) based on risk factors in **Supplementary Table 1**, excluding age, sex, and family history. Individuals above the median of this risk score distribution were considered to have an unfavorable modifiable risk profile. Risk trajectories in each stratum were visualized by fitting linear models with smoothing splines across individual risk estimates as a function of age. Differences in mean absolute risk at age 60 were tested using a two-sample t-test. We also tested for interaction between the 3-level ordinal PRS variable and the modifiable risk score (dichotomized at the median) in a linear model with the predicted absolute risk as the outcome.

Etiology: contribution of genetic vs. modifiable risk factors

The relative contribution of genetic and modifiable cancer risk factors at the population level was quantified with population attributable fractions (PAF) using the method of Sjölander & Vansteedlandt[49,50] based on the counterfactual framework. To obtain comparable AF estimates, thresholds for high genetic risk and high burden of modifiable risk factors corresponded to the top 20% (≥80th percentile) of each risk score distribution.

DATA AVAILABILITY

The UK Biobank in an open access resource, available at https://www.ukbiobank.ac.uk/researchers/. This research was conducted with approved access to UK Biobank data under application number 14105.

ACKNOWLEDGEMENTS

Disclaimer: Where authors are identified as personnel of the International Agency for Research on Cancer / World Health Organization, the authors alone are responsible for the views expressed in this article and they do not necessarily represent the decisions, policy or views of the International Agency for Research on Cancer / World Health Organization.

This research was supported by funding from the National Institutes of Health (US NCI R25T CA112355 and R01 CA201358; PI: Witte) and Cancer Research UK (C18281/A19169).

COMPETING INTERESTS

The authors declare no competing interests.

References

1. Khera, A.V.*, et al.* Polygenic Prediction of Weight and Obesity Trajectories from Birth to Adulthood. *Cell* **177**, 587-596 e589 (2019).
2. Yengo, L.*, et al.* Meta-analysis of genome-wide association studies for height and body mass index in approximately 700000 individuals of European ancestry. *Hum Mol Genet* **27**, 3641-3649 (2018).
3. Inouye, M.*, et al.* Genomic Risk Prediction of Coronary Artery Disease in 480,000 Adults: Implications for Primary Prevention. *J Am Coll Cardiol* **72**, 1883-1893 (2018).
4. Khera, A.V.*, et al.* Genome-wide polygenic scores for common diseases identify individuals with risk equivalent to monogenic mutations. *Nat Genet* **50**, 1219-1224 (2018).
5. Torkamani, A., Wineinger, N.E. & Topol, E.J. The personal and clinical utility of polygenic risk scores. *Nat Rev Genet* **19**, 581-590 (2018).
6. Lambert, S.A., Abraham, G. & Inouye, M. Towards clinical utility of polygenic risk scores. *Hum Mol Genet* (2019).
7. Bycroft, C.*, et al.* The UK Biobank resource with deep phenotyping and genomic data. *Nature* **562**, 203-209 (2018).
8. Raaschou-Nielsen, O.*, et al.* Air pollution and lung cancer incidence in 17 European cohorts: prospective analyses from the European Study of Cohorts for Air Pollution Effects (ESCAPE). *Lancet Oncol* **14**, 813-822 (2013).
9. Heagerty, P.J. & Zheng, Y. Survival model predictive accuracy and ROC curves. *Biometrics* **61**, 92-105 (2005).
10. Royston, P. Explained variation for survival models. *The Stata Journal* **6**, 83-96 (2006).
11. McKearnan, S.B., Wolfson, J., Vock, D.M., Vazquez-Benitez, G. & O'Connor, P.J. Performance of the Net Reclassification Improvement for Nonnested Models and a Novel Percentile-Based Alternative. *Am J Epidemiol* **187**, 1327-1335 (2018).
12. Pencina, M.J., D'Agostino, R.B. & Massaro, J.M. Understanding increments in model performance metrics. *Lifetime Data Anal* **19**, 202-218 (2013).
13. Usher-Smith, J.A., Sharp, S.J., Luben, R. & Griffin, S.J. Development and Validation of Lifestyle-Based Models to Predict Incidence of the Most Common Potentially Preventable Cancers. *Cancer Epidemiol Biomarkers Prev* **28**, 67-75 (2019).
14. Mucci, L.A.*, et al.* Familial Risk and Heritability of Cancer Among Twins in Nordic Countries. *JAMA* **315**, 68-76 (2016).
15. Rashkin, S.R.*, et al.* Pan-cancer study detects novel genetic risk variants and shared genetic basis in two large cohorts. *bioRxiv* **635367**(2019).
16. Graff, R.E.*, et al.* Cross-Cancer Evaluation of Polygenic Risk Scores for 17 Cancer Types in Two Large Cohorts. *bioRxiv*, 2020.2001.2018.911578 (2020).
17. Shi, Z.*, et al.* Systematic evaluation of cancer-specific genetic risk score for 11 types of cancer in The Cancer Genome Atlas and Electronic Medical Records and Genomics cohorts. *Cancer Med* **8**, 3196-3205 (2019).
18. Fritsche, L.G.*, et al.* Association of Polygenic Risk Scores for Multiple Cancers in a Phenome-wide Study: Results from The Michigan Genomics Initiative. *Am J Hum Genet* **102**, 1048-1061 (2018).
19. Amin Al Olama, A.*, et al.* Risk Analysis of Prostate Cancer in PRACTICAL, a Multinational Consortium, Using 25 Known Prostate Cancer Susceptibility Loci. *Cancer Epidemiol Biomarkers Prev* **24**, 1121-1129 (2015).
20. Hoffmann, T.J.*, et al.* A large multiethnic genome-wide association study of prostate cancer identifies novel risk variants and substantial ethnic differences. *Cancer Discov* **5**, 878-891 (2015).
21. Smith, T., Gunter, M.J., Tzoulaki, I. & Muller, D.C. The added value of genetic information in colorectal cancer risk prediction models: development and evaluation in the UK Biobank prospective cohort study. *Br J Cancer* **119**, 1036-1039 (2018).
22. Garcia-Closas, M.*, et al.* Common genetic polymorphisms modify the effect of smoking on absolute risk of bladder cancer. *Cancer Res* **73**, 2211-2220 (2013).
23. Mavaddat, N.*, et al.* Polygenic Risk Scores for Prediction of Breast Cancer and Breast Cancer Subtypes. *Am J Hum Genet* **104**, 21-34 (2019).
24. Yang, X.*, et al.* Evaluation of polygenic risk scores for ovarian cancer risk prediction in a prospective cohort study. *J Med Genet* **55**, 546-554 (2018).

25.   Klein, A.P*., et al.* Genome-wide meta-analysis identifies five new susceptibility loci for pancreatic cancer. *Nat Commun* **9**, 556 (2018).

26.   Fritsche, L.G*., et al.* Exploring various polygenic risk scores for skin cancer in the phenomes of the Michigan genomics initiative and the UK Biobank with a visual catalog: PRSWeb. *PLoS Genet* **15**, e1008202 (2019).

27.   Abraham, G*., et al.* Genomic risk score offers predictive performance comparable to clinical risk factors for ischaemic stroke. *Nat Commun* **10**, 5819 (2019).

28.   Khera, A.V*., et al.* Genetic Risk, Adherence to a Healthy Lifestyle, and Coronary Disease. *N Engl J Med* **375**, 2349-2358 (2016).

29.   Licher, S*., et al.* Genetic predisposition, modifiable-risk-factor profile and long-term dementia risk in the general population. *Nat Med* **25**, 1364-1369 (2019).

30.   Brown, K.F*., et al.* The fraction of cancer attributable to modifiable risk factors in England, Wales, Scotland, Northern Ireland, and the United Kingdom in 2015. *Br J Cancer* **118**, 1130-1141 (2018).

31.   Brenner, D.R*., et al.* The burden of cancer attributable to modifiable risk factors in Canada: Methods overview. *Prev Med* **122**, 3-8 (2019).

32.   Poirier, A.E*., et al.* The current and future burden of cancer attributable to modifiable risk factors in Canada: Summary of results. *Prev Med* **122**, 140-147 (2019).

33.   Fry, A*., et al.* Comparison of Sociodemographic and Health-Related Characteristics of UK Biobank Participants With Those of the General Population. *Am J Epidemiol* **186**, 1026-1034 (2017).

34.   National Lung Screening Trial Research, T*., et al.* Reduced lung-cancer mortality with low-dose computed tomographic screening. *N Engl J Med* **365**, 395-409 (2011).

35.   De Koning, H., Van Der Aalst, C., Ten Haaf, K. & Oudkerk, M. PL02.05 Effects of Volume CT Lung Cancer Screening: Mortality Results of the NELSON Randomised-Controlled Population Based Trial. *Journal of Thoracic Oncology* **13**, S185 (2018).

36.   Shieh, Y*., et al.* Breast Cancer Screening in the Precision Medicine Era: Risk-Based Screening in a Population-Based Trial. *J Natl Cancer Inst* **109**(2017).

37.   Hollands, G.J*., et al.* The impact of communicating genetic risks of disease on risk-reducing health behaviour: systematic review with meta-analysis. *BMJ* **352**, i1102 (2016).

38.   Manichaikul, A*., et al.* Robust relationship inference in genome-wide association studies. *Bioinformatics* **26**, 2867-2873 (2010).

39.   Huyghe, J.R*., et al.* Discovery of common and rare genetic risk variants for colorectal cancer. *Nat Genet* **51**, 76-87 (2019).

40.   Wooster, R. & Weber, B.L. Breast and ovarian cancer. *N Engl J Med* **348**, 2339-2347 (2003).

41.   Kazerouni, N., Greene, M.H., Lacey, J.V., Jr., Mink, P.J. & Schairer, C. Family history of breast cancer as a risk factor for ovarian cancer in a prospective study. *Cancer* **107**, 1075-1083 (2006).

42.   Olson, S.H. & Kurtz, R.C. Epidemiology of pancreatic cancer and the role of family history. *J Surg Oncol* **107**, 1-7 (2013).

43.   Molina-Montes, E*., et al.* Risk of pancreatic cancer associated with family history of cancer and other medical conditions by accounting for smoking among relatives. *Int J Epidemiol* **47**, 473-483 (2018).

44.   Schuz, J*., et al.* European Code against Cancer 4th Edition: 12 ways to reduce your cancer risk. *Cancer Epidemiol* **39 Suppl 1**, S1-10 (2015).

45.   Demler, O.V., Paynter, N.P. & Cook, N.R. Tests of calibration and goodness-of-fit in the survival setting. *Stat Med* **34**, 1659-1680 (2015).

46.   Gerds, T.A., Andersen, P.K. & Kattan, M.W. Calibration plots for risk prediction models in the presence of competing risks. *Stat Med* **33**, 3191-3203 (2014).

47.   Benichou, J. & Gail, M.H. Estimates of absolute cause-specific risk in cohort studies. *Biometrics* **46**, 813-826 (1990).

48.   Ozenne, B., Lyngholm Sørensen, A., Scheike, T., Torp-Pedersen, C. & Gerds, T.A. riskRegression: Predicting the Risk of an Event using Cox Regression Models. *The R Journal* **9**, 440-460 (2017).

49.   Sjolander, A. & Vansteelandt, S. Doubly robust estimation of attributable fractions in survival analysis. *Stat Methods Med Res* **26**, 948-969 (2017).

50.   Dahlqvist, E., Zetterqvist, J., Pawitan, Y. & Sjolander, A. Model-based estimation of the attributable fraction for cross-sectional, case-control and cohort studies using the R package AF. *Eur J Epidemiol* **31**, 575-582 (2016).

Figure 1: Hazard ratios (HR) per one standard deviation (SD) increase in the standardized polygenic risk score (PRS) estimated using cause-specific Cox proportional hazards models, accounting for mortality as a competing risk. A comparison of three weighting approaches for combining individual risk variants in the PRS is presented: standard weights based on per-allele log odds ratios (PRS$_\beta$), unweighted sum of risk alleles (PRS$_{unw}$), and inverse variance (IV) weights (PRS$_{IV}$).

Figure 2: Assessment of model discrimination based on Harrell's C index, computed as a weighted average between 1 and 5 years of follow-up time. Comparisons are conducted between the most comprehensive risk factor model for each cancer, including all available lifestyle-related risk factors and family history (if applicable), and a nested model that also includes the standardized polygenic risk score (PRS$_{IV}$) for that cancer and the top 15 genetic ancestry principal components.

Figure 3: Comparison of predicted 5-year absolute risk trajectories across strata defined by the presence of family history and the level of genetic risk, based on percentiles of the normalized polygenic risk score (PRS) distribution. Low PRS corresponds to ≤20th percentile, average PRS is defined as >20th to <80th percentile, and high PRS includes individuals in the ≥80th percentile of the genetic risk score distribution. P-values are based on t-tests comparing mean absolute risk in each stratum at age 60.

Figure 4: Predicted 5-year absolute risk trajectories for cancers where risk stratification is driven by genetic factors. Low genetic risk is based on percentiles of the standardized polygenic risk score (PRS). Low PRS corresponds to ≤20th percentile, average PRS is defined as >20th to <80th percentile, and high PRS includes individuals in the ≥80th percentile. Individuals below the median of the modifiable risk factor distribution were considered to have reduced risk, whereas those above the median had elevated risk. P-values are based on t-tests comparing mean absolute risk in each stratum at age 60, except for pre-menopausal breast cancer where differences at age 50 were tested.

Figure 5: Predicted 5-year absolute risk trajectories for cancers where risk stratification is driven by modifiable risk factors. Low genetic risk is based on percentiles of the standardized polygenic risk score (PRS). Low PRS corresponds to ≤20th percentile, average PRS is defined as >20th to <80th percentile, and high PRS includes individuals in the ≥80th percentile. Individuals below the median of the modifiable risk factor distribution were considered to have reduced risk, whereas those above the median had elevated risk. P-values are based on t-tests comparing mean absolute risk in each stratum at age 60.

Figure 6: Population attributable fractions (PAF) estimated at 5 years of follow-up time for the top 20% (≥80th percentile) of the modifiable risk factor and polygenic risk score (PRS) distributions, respectively, and family history of cancer at the relevant site. PAF estimates were derived from Cox proportional hazard regression models that were adjusted for age at enrollment, sex, family history of cancer (if available), genotyping array, and the top 15 genetic ancestry principal components.

Supplementary Table 1: Characteristics of the UK Biobank study population, restricted to participants of predominately European ancestry, stratified by incident cancer status.

| Baseline Characteristic | Status at the end of follow-up | | | |
| | Cancer-free (n=390,998) | | Incident Cancer (n=22,755) | |
	N	(%)	N	(%)
Age at assessment (years)				
Mean (SD)	56.53	(8.02)	60.19	(6.88)
Sex				
Females	211513	(54.10)	11080	(48.69)
Males	179485	(45.90)	11675	(51.31)
Smoking status				
Never	211936	(54.20)	10610	(46.63)
Ever	177685	(45.44)	12049	(52.95)
Former	137493	(35.16)	9174	(40.32)
Current	40192	(10.28)	2875	(12.63)
Pack-years: Mean (SD)	23.17	(15.61)	27.70	(19.19)
Unknown	1377	(0.35)	96	(0.42)
Body-mass index (kg/m^2)				
Normal: 18.5 ≤BMI<25	127817	(32.69)	6614	(29.07)
Underweight: <18.5	1984	(0.51)	96	(0.42)
Overweight: 25 ≤BMI<30	165859	(42.42)	10055	(44.19)
Obese: BMI≥30	94070	(24.06)	5914	(25.99)
Unknown	1268	(0.32)	76	(0.33)
Frequency of alcohol consumption				
Never	25864	(6.61)	1651	(7.26)
Special occasions	41557	(10.63)	2495	(10.96)
1-3 times per month	43627	(11.16)	2312	(10.16)
1-2 times per week	102999	(26.34)	5753	(25.28)
3-4 times per week	94177	(24.09)	5225	(22.96)
Daily	82482	(21.10)	5304	(23.31)
Unknown	292	(0.07)	1651	(7.26)
Ever diagnosed with hypertension?				
Yes	92541	(23.67)	6772	(29.76)
Any first-degree relative diagnosed with cancer[1]?				
Yes	139833	(35.76)	9195	(40.41)
Ever had screening for breast, prostate, or colorectal cancer?				
Yes	258249	(66.05)	16908	(74.30)
Deaths occurring during follow-up				
Death from any cause	5982	(1.53)	4878	(21.44)
Death due to cancer	2103	(0.54)	4696	(20.64)

[1] Based on self-reported cancers of the breast, prostate, lung, or bowel in non-adopted parents and siblings

Supplementary Table 2: Risk factors in addition to age and sex (if applicable), such as environmental exposures, lifestyle factors, and family history, that were included in the most comprehensive model for each cancer. Risk factors were selected based on literature review and availability in the UK Biobank cohort.

Cancer Site	Risk Factors	Model Specification Notes
Prostate	Family history of prostate cancer	
Testis	-	
Breast	Family history of breast cancer, parity ($\geq$1 live birth vs. none), age at menarche (years), menopausal status (pre-menopausal vs. post-menopausal vs. unknown or hysterectomy), ever used hormone replacement therapy (HRT), duration of oral contraceptive use (never used (0) vs. <20 years vs. $\geq$20 years), body mass index (BMI), weekly alcohol intake (grams)[1]	Interactions: (BMI)*(Menopausal status)
Endometrium	Family history of cancer, parity ($\geq$1 live birth vs. none), age at menarche (years), menopausal status (pre-menopausal vs. post-menopausal vs. unknown or hysterectomy), ever used HRT, duration of oral contraceptive use (never used (0) vs. <20 years vs. $\geq$20 years), BMI	
Ovary	Family history of breast cancer, parity ($\geq$1 live birth vs. none), menopausal status (pre-menopausal vs. post-menopausal vs. unknown or hysterectomy), ever used HRT, duration of oral contraceptive use (never used (0) vs. <20 years vs. $\geq$20 years), BMI	Interactions: (BMI)*(Menopausal status)
Cervix	Parity ($\geq$1 live birth vs. none), duration of oral contraceptive use (never used (0) vs. <20 years vs. $\geq$20 years), cigarette pack-years	
Colorectum	Family history of bowel cancer, waist to hip ratio (WHR), cigarette pack-years, frequency of processed meat intake (<1 per week vs. $\geq$1 per week), moderate and/or strenuous physical activity (days per week), weekly alcohol intake (grams)	
Melanoma	Frequency of UV protection use (always vs. most times vs. sometimes vs. never out in the sun vs. never), time spent outside in the summer (hours per day), ease of tanning (very easily vs. moderate vs. mild vs. mostly burn)	
Lung	Family history of lung cancer, cigarettes per day (0 for never smokers), years of smoking (0 for never smokers), smoking status (never vs. former vs. current), $PM_{2.5}$ level in 2010 (micro-g/m^3)	Interactions: (Former smoker)* (cigarettes/day) (Former smoker)*(years of smoking)
Never-smokers	Family history of lung cancer, $PM_{2.5}$ level in 2010 (micro-g/m^3)	
Smokers	Family history of lung cancer, cigarettes per day, years of smoking, smoking status (former vs. current), years since quitting smoking (0 for current smokers), $PM_{2.5}$ level in 2010 (micro-g/m^3)	Interactions: (Smoking status)*(cigarettes/day) (Smoking status)*(years smoking)
NHL	-	
Bladder	BMI, smoking status (never vs. former vs. current), cigarette pack-years	
Kidney	BMI, smoking status (never vs. former vs. current), cigarette pack-years, ever diagnosed with hypertension	
Pancreas	BMI, smoking status (never vs. former vs. current), cigarette pack-years, family history of cancer (prostate, breast, lung or bowel)	
Oral cavity/pharynx	Smoking status (never vs. former vs. current), cigarette pack-years, weekly alcohol intake (grams)	
Lymphocytic leukemia	-	
Thyroid	BMI categories (BMI <25 vs. 25 $\leq$BMI <30, BMI$\geq$30)	

[1]. Weekly alcohol intake was derived by summing up the total number of drinks per week across different types of alcoholic beverages (beer, wine, spirits) and converting to units of alcohol based on values from UK Composition of foods integrated dataset: https://www.gov.uk/government/publications/composition-of-foods-integrated-dataset-cofid

Supplementary Table 3: Hazard ratios (HR) for each cancer risk factor estimated using a cause-specific Cox regression model accounting for death as a competing risk.

Cancer Site and Risk Factors	HR[1]	(95% CI)	P-value
Prostate			
Family history of prostate cancer	1.84	(1.69 - 2.00)	9.1×10^{-46}
Breast			
Family history of breast cancer	1.56	(1.44 - 1.69)	3.0×10^{-29}
Parity (≥1 live birth)	0.91	(0.84 - 0.98)	0.010
Age at menarche (per 1 year)	0.99	(0.97 - 1.00)	0.14
BMI (per 1-unit increase)	0.99	(0.98 - 1.00)	0.11
Menopausal status: pre-menopausal	1.00		
Menopausal status: post-menopausal	0.36	(0.24 - 0.53)	1.8×10^{-7}
Menopausal status: unknown/hysterectomy	0.32	(0.18 - 0.55)	5.0×10^{-5}
Ever used hormone replacement therapy (HRT)	1.09	(1.02 - 1.17)	7.0×10^{-3}
Oral contraceptive use: 0 (never used)	1.00		
Oral contraceptive use: <20 years	1.00	(0.93 - 1.08)	0.94
Oral contraceptive use: ≥20 years	1.10	(0.98 - 1.23)	0.12
Alcohol intake[2] (70 g/week)	1.04	(1.02 - 1.05)	2.3×10^{-5}
BMI * menopausal status (post-menopausal)			2.0×10^{-5}
BMI * menopausal status (unknown/hysterectomy)			7.6×10^{-4}
Endometrium			
Family history of cancer	1.11	(0.95 - 1.30)	0.20
Parity (≥1 live birth)	0.64	(0.53 - 0.77)	3.3×10^{-6}
Age at menarche (per 1-year increase)	0.92	(0.88 - 0.97)	1.8×10^{-3}
BMI (per 1-unit increase)	1.09	(1.08 - 1.10)	1.6×10^{-49}
Menopausal status: pre-menopausal	1.00		
Menopausal status: post-menopausal	1.01	(0.73 - 1.39)	0.97
Menopausal status: unknown/hysterectomy	0.02	(0.00 - 0.08)	6.1×10^{-8}
Ever used HRT	0.84	(0.71 - 0.99)	0.041
Oral contraceptive use: 0 (never used)	1.00		
Oral contraceptive use: <20 years	0.83	(0.70 - 1.00)	0.051
Oral contraceptive use: ≥20 years	0.36	(0.24 - 0.56)	4.9×10^{-6}
Ovary			
Family history of breast cancer	1.30	(1.00 - 1.70)	0.051
Parity (≥1 live birth)	0.72	(0.57 - 0.91)	6.2×10^{-3}
BMI (per 1-unit increase)	1.04	(1.00 - 1.08)	0.036
Menopausal status: post-menopausal	3.19	(0.88 - 11.56)	0.08
Menopausal status: unknown/hysterectomy	1.07	(0.13 - 8.52)	0.95
Ever used HRT	0.97	(0.79 - 1.19)	0.79
Duration of oral contraceptive use: <20 years	0.82	(0.66 - 1.03)	0.09
Duration of oral contraceptive use: ≥20 years	0.57	(0.37 - 0.88)	0.012
BMI * menopausal status (post-menopausal)			0.023
BMI * menopausal status (unknown/hysterectomy)			0.45
Cervix			
Parity: ≥1 live birth	1.81	(1.30 - 2.53)	4.9×10^{-4}

Oral contraceptive use: 0 (never used)			
Oral contraceptive use: <20 years	0.84	(0.58 - 1.21)	0.34
Oral contraceptive use: ≥20 years	1.08	(0.69 - 1.69)	0.74
Cigarette pack-years (per 10 pack-years)	1.15	(1.05 - 1.25)	1.8×10^{-3}
Colon/rectum			
Family history of bowel cancer	1.26	(1.14 - 1.40)	1.2×10^{-5}
Waist to hip ratio (per 10% increase)	1.17	(1.11 - 1.24)	2.2×10^{-8}
Cigarette pack-years (per 10 pack-years)	1.04	(1.02 - 1.06)	2.1×10^{-4}
Processed meat intake: never	1.00		
Processed meat intake: < once a week	0.99	(0.98 - 1.00)	0.20
Processed meat intake: ≥ once a week	1.08	(0.92 - 1.28)	0.34
Physical activity: strenuous or moderate (days/week)	1.15	(0.98 - 1.35)	0.09
Alcohol intake[2] (70 g/week)	1.04	(1.03 - 1.05)	5.9×10^{-9}
Melanoma			
Apply UV protection: never	1.00		
Apply UV protection: sometimes	1.37	(1.10 - 1.69)	4.1×10^{-3}
Apply UV protection: most times	1.82	(1.48 - 2.25)	2.1×10^{-8}
Apply UV protection: always	1.68	(1.34 - 2.09)	5.3×10^{-6}
Apply UV protection: never in the sun	1.06	(0.46 - 2.42)	0.89
Time outdoors in the summer (hours per day)	1.03	(1.01 - 1.05)	7.3×10^{-3}
Ease of tanning: get very tan	1.00		
Ease of tanning: moderate	1.33	(1.16 - 1.53)	6.4×10^{-5}
Ease of tanning: mild	1.60	(1.37 - 1.86)	1.9×10^{-9}
Ease of tanning: mostly burn	1.61	(1.37 - 1.89)	4.0×10^{-9}
Lung			
Family history of lung cancer	1.61	(1.43 - 1.81)	7.4×10^{-15}
$PM_{2.5}$ in 2010 (per 1 micro-g/m^3)	1.10	(1.05 - 1.15)	1.9×10^{-5}
Cigarettes per day	1.00	(0.99 - 1.00)	0.52
Years of smoking	1.07	(1.06 - 1.09)	9.0×10^{-23}
Smoking status: never	1.00		
Smoking status: former	0.34	(0.25 - 0.46)	6.9×10^{-13}
Smoking status: current	0.84	(0.44 - 1.61)	0.60
Smoking status (former) * cigarettes per day			2.2×10^{-8}
Smoking status (former) * years of smoking			0.40
Lung (Never smokers)			
Family history of lung cancer	0.91	(0.61 - 1.38)	0.67
$PM_{2.5}$ in 2010 (per 1 micro-g/m^3)	0.93	(0.81 - 1.07)	0.32
Lung (Current or former smokers)			
Family history of lung cancer	1.71	(1.51 - 1.94)	3.8×10^{-17}
$PM_{2.5}$ in 2010 (per 1 micro-g/m^3)	1.12	(1.07 - 1.18)	1.2×10^{-6}
Cigarettes per day	1.02	(1.02 - 1.03)	1.2×10^{-13}
Years of smoking	1.07	(1.05 - 1.10)	1.4×10^{-8}
Smoking status: former	1.00		
Smoking status: current	2.12	(0.92 - 4.89)	0.08
Years since quitting (per 1 year)	1.01	(0.98 - 1.04)	0.44
Smoking status (current) * cigarettes per day			2.4×10^{-8}

Smoking status (current) * years of smoking			0.51
Bladder			
Cigarette pack-years (per 10 pack-years)	1.07	(1.03 - 1.11)	3.4×10^{-4}
Smoking status: never	1.00		
Smoking status: former	1.52	(1.26 - 1.82)	8.0×10^{-6}
Smoking status: current	2.07	(1.62 - 2.64)	5.7×10^{-9}
BMI (per 1-unit increase)	1.01	(0.99 - 1.02)	0.45
Kidney			
BMI (per 1-unit increase)	1.04	(1.02 - 1.05)	1.7×10^{-6}
Smoking status: never	1.00		
Smoking status: former	1.07	(0.87 - 1.32)	0.52
Smoking status: current	1.36	(1.02 - 1.83)	0.039
Cigarette pack-years (per 10 pack-years)	1.07	(1.02 - 1.12)	7.9×10^{-3}
Diagnosed with hypertension	1.69	(1.44 - 1.98)	2.1×10^{-10}
Pancreas			
Family history of cancer (prostate, breast, lung, bowel)	1.40	(1.17 - 1.67)	1.9×10^{-4}
BMI (per 1-unit increase)	1.03	(1.01 - 1.05)	8.4×10^{-4}
Cigarette pack-years (per 10 pack-years)	1.04	(0.97 - 1.10)	0.27
Smoking status: never	1.00		
Smoking status: former	1.08	(0.84 - 1.39)	0.56
Smoking status: current	2.04	(1.46 - 2.84)	2.8×10^{-5}
Oral cavity/pharynx			
Alcohol intake[2] (70 g/week)	1.05	(1.04 - 1.07)	3.0×10^{-10}
Cigarettes per day	1.01	(1.00 - 1.02)	8.2×10^{-3}
Years of smoking	1.02	(1.01 - 1.04)	2.0×10^{-4}
Smoking status: never	1.00		
Smoking status: former	0.58	(0.37 - 0.91)	0.050
Smoking status: current	1.09	(0.60 - 1.96)	0.43
Thyroid			
BMI: <25	1.00		
BMI: 25 to <30	1.43	(1.01 - 2.02)	0.045
BMI: 30 to <35	1.59	(1.05 - 2.41)	0.028
BMI: ≥35	1.15	(0.61 - 2.15)	0.67

1. In addition to the listed risk factors all Cox regression models were adjusted for age and sex (if applicable)

2. Weekly alcohol intake was derived by summing up the total number of drinks per week across different types of alcoholic beverages (beer, wine, spirits) and converting to units of alcohol based on values from UK Composition of foods integrated dataset: https://www.gov.uk/government/publications/composition-of-foods-integrated-dataset-cofid

Supplementary Table 4: Hazard ratios (HR) per one standard deviation (SD) increase in the standardized polygenic risk score (PRS) for each cancer, estimated using cause-specific Cox proportional hazards models, accounting for mortality as a competing risk. Results comparing three types of weighting approaches for combining individual risk variants in the PRS are presented: standard weights based on log odds ratios (PRS$_\beta$), unweighted sum of risk alleles (PRS$_{unw}$), and inverse variance (IV) weights (PRS$_{IV}$).

Cancer Site	Cases	PRS Description Variants	PRS Description Weights	HR[1]	(95% CI)	P-value	C index[2]	(C SE)	AUC[3]
Prostate	4740	161	PRS$_\beta$	1.39	(1.35-1.43)	2.0×10^{-105}	0.738	(0.004)	0.740
			PRS$_{unw}$	1.66	(1.62-1.71)	3.4×10^{-266}	0.759	(0.004)	0.761
			PRS$_{IV}$	1.77	(1.72-1.82)	4.3×10^{-336}	0.768	(0.004)	0.769
Testis	52	52	PRS$_\beta$	2.18	(1.66-2.87)	2.3×10^{-8}	0.749	(0.034)	0.783
			PRS$_{unw}$	1.96	(1.49-2.58)	1.4×10^{-6}	0.745	(0.035)	0.769
			PRS$_{IV}$	2.26	(1.71-2.99)	1.0×10^{-8}	0.766	(0.033)	0.787
Breast	4760	187	PRS$_\beta$	1.25	(1.21-1.28)	1.8×10^{-53}	0.594	(0.005)	0.598
			PRS$_{unw}$	1.41	(1.37-1.45)	2.1×10^{-123}	0.616	(0.005)	0.621
			PRS$_{IV}$	1.51	(1.47-1.56)	3.1×10^{-179}	0.632	(0.005)	0.637
Endometrium	643	9	PRS$_\beta$	1.19	(1.10-1.29)	1.1×10^{-5}	0.749	(0.011)	0.755
			PRS$_{unw}$	1.18	(1.09-1.28)	2.4×10^{-5}	0.749	(0.011)	0.754
			PRS$_{IV}$	1.18	(1.09-1.27)	3.5×10^{-5}	0.749	(0.011)	0.754
Ovary	445	36	PRS$_\beta$	1.13	(1.04-1.24)	6.2×10^{-3}	0.655	(0.015)	0.656
			PRS$_{unw}$	1.18	(1.07-1.29)	5.8×10^{-4}	0.652	(0.016)	0.658
			PRS$_{IV}$	1.20	(1.10-1.32)	9.0×10^{-5}	0.654	(0.015)	0.660
Cervix	282	10	PRS$_\beta$	1.22	(1.09-1.37)	7.3×10^{-4}	0.750	(0.017)	0.745
			PRS$_{unw}$	1.20	(1.07-1.35)	1.5×10^{-3}	0.749	(0.017)	0.745
			PRS$_{IV}$	1.21	(1.07-1.35)	1.5×10^{-3}	0.749	(0.017)	0.745
Colon/rectum	2725	103	PRS$_\beta$	1.32	(1.27-1.37)	5.5×10^{-50}	0.704	(0.006)	0.704
			PRS$_{unw}$	1.46	(1.41-1.52)	9.2×10^{-87}	0.714	(0.006)	0.714
			PRS$_{IV}$	1.48	(1.43-1.54)	1.8×10^{-94}	0.716	(0.006)	0.716
Melanoma	1805	24	PRS$_\beta$	1.43	(1.36-1.49)	5.7×10^{-51}	0.663	(0.008)	0.652
			PRS$_{unw}$	1.43	(1.36-1.49)	1.2×10^{-50}	0.662	(0.008)	0.652
			PRS$_{IV}$	1.44	(1.37-1.50)	2.4×10^{-53}	0.664	(0.008)	0.654
Lung	1541	109	PRS$_\beta$	1.16	(1.11-1.22)	1.5×10^{-9}	0.849	(0.006)	0.846
			PRS$_{unw}$	1.15	(1.09-1.20)	1.5×10^{-8}	0.849	(0.006)	0.846
			PRS$_{IV}$	1.17	(1.12-1.23)	1.2×10^{-10}	0.849	(0.006)	0.846
NHL	970	19	PRS$_\beta$	1.16	(1.09-1.24)	1.0×10^{-6}	0.676	(0.010)	0.677
			PRS$_{unw}$	1.18	(1.11-1.25)	2.9×10^{-7}	0.675	(0.010)	0.678
			PRS$_{IV}$	1.15	(1.08-1.22)	1.1×10^{-5}	0.674	(0.010)	0.677
Bladder	890	15	PRS$_\beta$	1.28	(1.20-1.37)	2.1×10^{-13}	0.813	(0.008)	0.803
			PRS$_{unw}$	1.30	(1.21-1.39)	7.6×10^{-15}	0.814	(0.008)	0.803
			PRS$_{IV}$	1.30	(1.22-1.39)	1.5×10^{-15}	0.814	(0.008)	0.804
Kidney	612	19	PRS$_\beta$	1.16	(1.08-1.26)	1.0×10^{-4}	0.724	(0.011)	0.722
			PRS$_{unw}$	1.13	(1.05-1.22)	1.5×10^{-3}	0.723	(0.011)	0.721

			PRS$_{IV}$	1.15	(1.07-1.24)	2.6×10^{-4}	0.723	(0.011)	0.722
Pancreas	493	22	PRS$_\beta$	1.49	(1.36-1.62)	1.3×10^{-18}	0.742	(0.012)	0.745
			PRS$_{unw}$	1.44	(1.31-1.57)	1.1×10^{-15}	0.738	(0.012)	0.741
			PRS$_{IV}$	1.49	(1.37-1.63)	5.2×10^{-19}	0.743	(0.012)	0.745
Oral cavity/ pharynx	481	14	PRS$_\beta$	1.11	(1.01-1.21)	2.3×10^{-2}	0.686	(0.015)	0.702
			PRS$_{unw}$	1.11	(1.02-1.22)	1.9×10^{-2}	0.686	(0.015)	0.702
			PRS$_{IV}$	1.12	(1.02-1.23)	1.3×10^{-2}	0.687	(0.015)	0.702
Lymphocytic leukemia	340	75	PRS$_\beta$	1.45	(1.31-1.61)	8.0×10^{-13}	0.735	(0.016)	0.719
			PRS$_{unw}$	1.67	(1.51-1.86)	1.2×10^{-21}	0.755	(0.015)	0.736
			PRS$_{IV}$	1.70	(1.53-1.88)	6.3×10^{-23}	0.756	(0.015)	0.738
Thyroid	191	12	PRS$_\beta$	1.57	(1.36-1.82)	5.7×10^{-10}	0.666	(0.023)	0.679
			PRS$_{unw}$	1.55	(1.34-1.78)	1.9×10^{-9}	0.671	(0.023)	0.676
			PRS$_{IV}$	1.75	(1.53-2.01)	1.9×10^{-15}	0.692	(0.022)	0.701

1. Hazard ratio estimates are adjusted for age at assessment (years), sex (if applicable), family history of cancer (for sites with available self-reported information), genotyping array, the first 15 genetic ancestry principal components, and any additional risk factors applicable to each cancer listed in Supplementary Table 1

2. Harrell's C-index was calculated as a weighted average between 1 and 5 years of follow-up

3. AUC values were estimated at 5 years of follow-up

Supplementary Table 5: Assessment of model discrimination for each cancer comparing different combinations of conventional risk factors and polygenic risk scores (PRS).

Cancer Site	Cases	Model specification	C index[2]	(C SE)	AUC[3]	Pseudo R^2
Prostate	4740	Age	0.710	(0.004)	0.713	0.349
		Age + family history	0.716	(0.004)	0.720	0.366
		Age + PRS$_{IV}$ (instead of family history)	0.763	(0.004)	0.766	0.496
		Age + family history + PRS$_{IV}$	0.768	(0.004)	0.769	0.510
Testis	52	Age	0.627	(0.045)	0.658	0.184
		Age + PRS$_{IV}$	0.766	(0.033)	0.787	0.605
Breast	4760	Age	0.543	(0.005)	0.548	0.017
		Age + family history	0.559	(0.005)	0.562	0.031
		Age + PRS$_{IV}$ (instead of family history)	0.619	(0.005)	0.626	0.125
		Age + family history + other predictors	0.572	(0.005)	0.573	0.043
		Age + family history + other predictors + PRS$_{IV}$	0.632	(0.005)	0.637	0.146
Endometrium	643	Age	0.631	(0.012)	0.631	0.127
		Age + family history	0.629	(0.012)	0.632	0.129
		Age + family history + other predictors	0.744	(0.011)	0.747	0.463
		Age + family history + other predictors + PRS$_{IV}$	0.749	(0.011)	0.754	0.486
Ovary	445	Age	0.607	(0.016)	0.620	0.106
		Age + family history	0.611	(0.016)	0.622	0.111
		Age + family history + other predictors	0.641	(0.015)	0.643	0.151
		Age + family history + other predictors + PRS$_{IV}$	0.654	(0.015)	0.660	0.193
Cervix[4]	282	Age	0.729	(0.017)	0.719	0.346
		Age + other predictors	0.736	(0.018)	0.731	0.386
		Age + other predictors + PRS$_{IV}$	0.749	(0.017)	0.745	0.437
Colon/rectum	2725	Age + sex	0.678	(0.006)	0.680	0.235
		Age + sex + family history	0.679	(0.006)	0.681	0.239
		Age + sex + PRS (instead of family history)	0.708	(0.006)	0.708	0.319
		Age + sex + family history + other predictors	0.686	(0.006)	0.688	0.258
		Age + sex + family history + other predictors + PRS$_{IV}$	0.716	(0.006)	0.716	0.345
Melanoma	1805	Age + sex	0.597	(0.008)	0.592	0.063
		Age + sex + other predictors	0.622	(0.008)	0.616	0.100
		Age + sex + other predictors + PRS$_{IV}$	0.664	(0.008)	0.654	0.180
Lung	1541	Age + sex	0.706	(0.007)	0.704	0.307
		Age + sex + family history	0.713	(0.007)	0.714	0.333
		Age + sex + PRS (instead of family history)	0.711	(0.007)	0.710	0.322
		Age + sex + family history + other predictors	0.846	(0.006)	0.843	0.789
		Age + sex + family history + other predictors + PRS$_{IV}$	0.849	(0.006)	0.846	0.799
Never smokers	207	Age + sex + family history + other predictors	0.709	(0.020)	0.712	0.320
		Age + sex + family history + other predictors + PRS$_{IV}$	0.723	(0.020)	0.723	0.354
Smokers	1334	Age + sex + family history + other predictors	0.805	(0.007)	0.804	0.641

	N	Model	C-index	(SE)	AUC	R^2
		Age + sex + family history + other predictors + PRS$_{IV}$	0.809	(0.007)	0.808	0.657
NHL	970	Age + sex	0.667	(0.010)	0.669	0.207
		Age + sex + PRS$_{IV}$	0.674	(0.010)	0.677	0.227
Bladder	890	Age + sex	0.792	(0.008)	0.784	0.548
		Age + sex + other predictors	0.808	(0.008)	0.796	0.595
		Age + sex + other predictors + PRS$_{IV}$	0.814	(0.008)	0.804	0.628
Kidney	612	Age + sex	0.685	(0.012)	0.687	0.253
		Age + sex + other predictors	0.716	(0.011)	0.713	0.338
		Age + sex + other predictors + PRS$_{IV}$	0.723	(0.011)	0.722	0.366
Pancreas	493	Age + sex	0.692	(0.014)	0.695	0.273
		Age + sex + family history + other predictors	0.714	(0.013)	0.715	0.336
		Age + sex + family history + other predictors + PRS$_{IV}$	0.743	(0.012)	0.745	0.439
Oral cavity / pharynx	481	Age + sex	0.610	(0.014)	0.627	0.117
		Age + sex + other predictors	0.681	(0.015)	0.693	0.332
		Age + sex + other predictors + PRS$_{IV}$	0.687	(0.015)	0.702	0.356
Lymphocytic leukemia	340	Age + sex	0.695	(0.016)	0.688	0.255
		Age + sex + PRS$_{IV}$	0.756	(0.015)	0.738	0.415
Thyroid	191	Age + sex	0.577	(0.024)	0.590	0.060
		Age + sex + other predictors	0.592	(0.024)	0.604	0.079
		Age + sex + other predictors + PRS$_{IV}$	0.692	(0.022)	0.701	0.310

1. Harrell's C-index was calculated as a weighted average between 1 and 5 years of follow-up

2. AUC values were estimated at 5 years of follow-up

3. R^2 values correspond to the measure of explained variation by Royston P (2006) [Reference 10]

4. Incorporating time-varying PRS effects did not have an impact on predictive performance (AUC=0.745). As an additional sensitivity analysis we estimated the AUC using scaled Schoenfeld residuals, to account for non-proportionality in the PRS effects, which yielded AUC=0.772.

Supplementary Table 6: Percentile net reclassification improvement (NRI) index comparing the most comprehensive conventional risk factor model for each cancer with the model that incorporates the polygenic risk score (PRS) in addition to these risk factors. Event NRI (NRI$_e$) and non-event NRI (NRI$_{ne}$) quantify reclassification improvement in cases and event-free individuals, respectively.

Cancer Site	NRI[1]	(95% CI)	NRI$_e$[2]	(95% CI)	NRI$_{ne}$[3]	(95% CI)
Prostate	0.444	(0.420, 0.467)	0.441	(0.417, 0.463)	0.003	(-0.0002, 0.007)
Testis	0.368	(0.174, 0.538)	0.385	(0.192, 0.555)	-0.018	(-0.021, -0.014)
Breast	0.379	(0.361, 0.397)	0.367	(0.349, 0.385)	0.012	(0.008, 0.016)
Endometrium	0.058	(0.019, 0.098)	0.060	(0.021, 0.099)	-0.002	(-0.004, 0.0003)
Ovary	0.173	(0.097, 0.236)	0.168	(0.093, 0.233)	0.005	(0.001, 0.008)
Cervix	0.073	(-0.001, 0.152)	0.075	(0.001, 0.156)	-0.002	(-0.005, 0.001)
Colon/rectum	0.273	(0.244, 0.301)	0.271	(0.242, 0.298)	0.003	(0.0003, 0.005)
Melanoma	0.293	(0.260, 0.323)	0.285	(0.250, 0.315)	0.008	(0.005, 0.011)
Lung	0.044	(0.020, 0.070)	0.029	(0.004, 0.055)	0.015	(0.013, 0.017)
NHL	0.119	(0.078, 0.161)	0.116	(0.074, 0.157)	0.004	(0.002, 0.006)
Bladder	0.116	(0.069, 0.164)	0.116	(0.069, 0.164)	0.0003	(-0.002, 0.002)
Kidney	0.065	(0.026, 0.103)	0.069	(0.030, 0.107)	-0.004	(-0.006, -0.003)
Pancreas	0.228	(0.174, 0.284)	0.228	(0.173, 0.284)	0.001	(-0.002, 0.003)
Oral cavity / pharynx	0.065	(0.022, 0.107)	0.069	(0.026. 0.111)	-0.004	(-0.006, -0.002)
Lymphocytic leukemia	0.307	(0.227, 0.388)	0.312	(0.232, 0.394)	-0.004	(-0.007, -0.002)
Thyroid	0.415	(0.330, 0.507)	0.409	(0.324, 0.501)	0.006	(0.003, 0.008)

[1.] NRI = NRI$_e$ + NRI$_{ne}$

[2.] Difference in proportions of subjects with events (incident cancer) correctly reclassified to a higher-risk category minus those reclassified to a lower-risk category

[3.] Difference in proportions of subjects without events (cancer-free) correctly reclassified to a lower-risk category minus those reclassified to a higher-risk category

Supplementary Table 7: Assessment of interaction on the absolute risk scale between ordinal polygenic risk score (PRS) categories (average: 20th to <80th percentile; high: ≥80th percentile vs. low: ≤20th percentile) and family history of cancer (yes vs. none) or elevated modifiable risk factor profile (summary score >50th percentile vs. ≤50th percentile). Interaction was assessed using linear regression models with predicted absolute risk of cancer at age 60 (age 50 for pre-menopausal breast cancer) as the outcome.

Cancer Site	Interaction Combination	Interaction		Overall Interaction P-value[1]
		Coefficient	P-value	
Prostate	PRS(average) * family history(yes)	7.8×10^{-3}	7.0×10^{-15}	9.0×10^{-128}
	PRS(high) * family history(yes)	0.026	7.4×10^{-104}	
Breast	PRS(average) * family history(yes)	3.1×10^{-3}	2.2×10^{-9}	1.7×10^{-104}
	PRS(high) * family history(yes)	0.012	1.0×10^{-82}	
Colorectal	PRS(average) * family history(yes)	3.3×10^{-4}	0.030	8.7×10^{-14}
	PRS(high) * family history(yes)	1.2×10^{-3}	2.8×10^{-12}	
Lung	PRS(average) * family history(yes)	-5.4×10^{-5}	0.88	7.0×10^{-4}
	PRS(high) * family history(yes)	1.11×10^{-3}	5.9×10^{-3}	
Breast (Pre-menopausal)	PRS(average) * modifiable(elevated)	2.3×10^{-3}	0.063	1.5×10^{-3}
	PRS(high) * modifiable(elevated)	5.3×10^{-3}	4.0×10^{-4}	
Breast (Post-menopausal)	PRS(average) * modifiable(elevated)	1.7×10^{-3}	2.1×10^{-5}	1.3×10^{-21}
	PRS(high) * modifiable(elevated)	4.6×10^{-3}	1.3×10^{-21}	
Endometrium	PRS(average) * modifiable(elevated)	5.0×10^{-4}	4.0×10^{-5}	5.3×10^{-16}
	PRS(high) * modifiable(elevated)	1.2×10^{-3}	1.2×10^{-16}	
Ovary	PRS(average) * modifiable(elevated)	1.5×10^{-4}	5.9×10^{-7}	1.4×10^{-19}
	PRS(high) * modifiable(elevated)	3.5×10^{-4}	1.9×10^{-20}	
Cervix	PRS(average) * modifiable(elevated)	7.7×10^{-5}	0.072	9.1×10^{-5}
	PRS(high) * modifiable(elevated)	2.2×10^{-4}	2.6×10^{-5}	
Colorectum	PRS(average) * modifiable(elevated)	1.0×10^{-3}	3.0×10^{-38}	1.3×10^{-208}
	PRS(high) * modifiable(elevated)	2.9×10^{-3}	1.3×10^{-194}	
Melanoma	PRS(average) * modifiable(elevated)	5.1×10^{-4}	3.8×10^{-28}	3.3×10^{-122}
	PRS(high) * modifiable(elevated)	1.3×10^{-3}	3.9×10^{-118}	
Lung	PRS(average) * modifiable(elevated)	1.2×10^{-3}	3.6×10^{-8}	1.1×10^{-37}
	PRS(high) * modifiable(elevated)	3.3×10^{-3}	4.3×10^{-37}	
Bladder	PRS(average) * modifiable(elevated)	3.4×10^{-4}	3.3×10^{-11}	1.5×10^{-50}
	PRS(high) * modifiable(elevated)	9.3×10^{-4}	1.2×10^{-49}	
Kidney	PRS(average) * modifiable(elevated)	1.8×10^{-4}	9.7×10^{-9}	5.5×10^{-29}
	PRS(high) * modifiable(elevated)	4.2×10^{-4}	1.6×10^{-29}	
Pancreas	PRS(average) * modifiable(elevated)	1.6×10^{-4}	3.7×10^{14}	2.1×10^{-91}
	PRS(high) * modifiable(elevated)	5.2×10^{-4}	6.5×10^{-85}	
Oral cavity/pharynx	PRS(average) * modifiable(elevated)	1.1×10^{-4}	9.6×10^{-4}	5.2×10^{-11}
	PRS(high) * modifiable(elevated)	2.7×10^{-4}	1.1×10^{-11}	
Thyroid	PRS(average) * modifiable(elevated)	4.7×10^{-5}	2.1×10^{-4}	5.4×10^{-30}
	PRS(high) * modifiable(elevated)	1.7×10^{-4}	6.6×10^{-28}	

1. Calculated based on a chi-square test with 2 degrees of freedom

Supplementary Table 8: Population attributable fractions (PAF) were estimated at 5 years of follow-up time for the top 20% (≥80th percentile) of the modifiable risk factor and polygenic risk score (PRS) distributions, respectively, and family history of cancer at the relevant site. PAF estimates were derived from Cox proportional hazard regression models that were adjusted for age at enrollment, sex, family history of cancer (if available), genotyping array, and the top 15 genetic ancestry principal components.

	High Genetic Risk			High Modifiable Risk			Family History		
	PAF	(95% CI)	P-value	PAF	(95% CI)	P-value	PAF	(95% CI)	P-value
Prostate	0.232	(0.215 - 0.249)	5.5×10^{-158}	-	-	-	0.055	(0.045 - 0.065)	1.4×10^{-25}
Testis	0.303	(0.135 - 0.472)	4.5×10^{-4}	-	-	-	-	-	-
Breast	0.166	(0.149 - 0.182)	2.6×10^{-85}	0.042	(0.026 - 0.057)	2.0×10^{-7}	0.052	(0.040 - 0.063)	1.9×10^{-18}
Pre-menopausal	0.184	(0.147 - 0.221)	1.7×10^{-22}	0.061	(0.039 - 0.083)	9.6×10^{-8}	0.044	(0.020 - 0.068)	3.1×10^{-4}
Post-menopausal	0.155	(0.135 - 0.175)	1.3×10^{-51}	0.008	(0.001 - 0.015)	0.029	0.054	(0.040 - 0.069)	9.2×10^{-14}
Endometrium	0.043	(0.002 - 0.084)	0.039	0.353	(0.303 - 0.404)	1.5×10^{-43}	0.042	(-0.020 - 0.103)	0.18
Ovary[1]	0.082	(0.031 - 0.134)	1.6×10^{-3}	0.100	(0.038 - 0.161)	1.4×10^{-3}	0.025	(-0.012 - 0.063)	0.19
Cervix	0.123	(0.057 - 0.190)	2.7×10^{-4}	0.065	(-0.001 - 0.130)	0.053	-	-	-
Colon/rectum	0.167	(0.145 - 0.190)	9.2×10^{-50}	0.111	(0.085 - 0.136)	1.6×10^{-17}	0.027	(0.012 - 0.042)	5.3×10^{-4}
Melanoma	0.139	(0.112 - 0.166)	1.3×10^{-23}	0.066	(0.039 - 0.093)	1.1×10^{-6}	-	-	-
Lung	0.040	(0.013 - 0.066)	3.1×10^{-3}	0.636	(0.606 - 0.666)	1.4×10^{-376}	0.089	(0.065 - 0.114)	1.0×10^{-12}
Never smokers[2]	0.077	(0.002 - 0.151)	0.045	0.049	(-0.076 - 0.174)	0.44	-0.013	(-0.066 - 0.039)	0.62
Smokers	0.035	(0.007 - 0.063)	0.015	0.663	(0.620 - 0.706)	3.2×10^{-200}	0.105	(0.078 - 0.132)	2.8×10^{-14}
NHL	0.053	(0.020 - 0.087)	1.9×10^{-3}	-	-	-	-	-	-
Bladder	0.085	(0.048 - 0.121)	4.7×10^{-6}	0.189	(0.140 - 0.237)	4.1×10^{-14}	-	-	-
Kidney	0.046	(0.005 - 0.087)	0.026	0.210	(0.160 - 0.260)	2.4×10^{-16}	-	-	-
Pancreas[3]	0.133	(0.082 - 0.184)	2.9×10^{-7}	0.118	(0.064 - 0.172)	1.9×10^{-5}	0.134	(0.063 - 0.205)	2.3×10^{-4}
Oral cavity / pharynx	0.006	(-0.038 - 0.051)	0.78	0.310	(0.253 - 0.368)	4.0×10^{-26}	-	-	-
Lymphocytic leukemia	0.269	(0.204 - 0.334)	7.4×10^{-16}	-	-	-	-	-	-
Thyroid	0.268	(0.180 - 0.355)	1.7×10^{-9}	0.202	(0.039 - 0.366)	0.015	-	-	-

1. Family history variable refers to self-reported breast cancer in a first-degree relative
2. The only modifiable risk factor is air pollution, modeled here as a categorical variable corresponding to $PM_{2.5}$ levels above the median
3. Family history variable refers to self-reported breast, prostate, lung, or bowel cancer in a first-degree relative

Supplementary Figure 1: Calibration plots comparing predicted and observed event probabilities for each of the 16 cancers examined. The most comprehensive risk factor model available is plotted in black and the same model with the addition of the polygenic risk score (PRS) is overlaid in red.

Supplementary Figure 2: Plots of Schoenfeld residuals and corresponding p-values for the cervical cancer polygenic risk score (PRS$_{IV}$). Below is a comparison of PRS$_{IV}$ effects estimated using a time-varying model (blue) and hazard ratio estimated under the proportionality assumption (red).

Supplementary Figure 3: Predicted 5-year absolute risk trajectories across strata defined by PRS and modifiable risk factors, where applicable. Low genetic risk is based on percentiles of the standardized polygenic risk score (PRS). Low PRS corresponds to ≤20th percentile, average PRS is defined as >20th to <80th percentile, and high PRS includes individuals in the ≥80th percentile. Individuals below the median of the modifiable risk factor distribution were considered to have reduced risk, whereas those above the median had elevated risk. P-values are based on t-tests comparing mean absolute risk in each stratum at age 60 or age 50 for cervical and testicular cancers.

Note: for cervical cancer estimates of absolute risk were derived from the Cox proportional hazards model without time-varying PRS effects