1   Classification: Biological Sciences

2   Title: Horizontal gene transfer in the human and skin commensal *Malassezia*: a bacterially-
3   derived flavohemoglobin is required for NO resistance and host interaction

8   Giuseppe Ianiri[1#], Marco A. Coelho[1], Fiorella Ruchti[2], Florian Sparber[2], Timothy J. McMahon[3],
9   Ci Fu[1], Madison Bolejack[4,6], Olivia Donovan[5,6], Hayden Smutney[5,6], Peter Myler[6,7,8], Fred
10   Dietrich[1], David Fox III[4,6], Salomé LeibundGut-Landmann[2], and Joseph Heitman*[1]

[1] Department of Molecular Genetics and Microbiology, Duke University Medical Center, Durham,
NC, 27710, USA. [2]Section of Immunology, Vetsuisse Faculty, University of Zürich, Zürich,
Switzerland. [3]Department of Medicine, VA Medical Center, Durham, NC, 27705, USA.[4]UCB
Pharma, Bainbridge Island, WA, USA;[5]UCB Pharma, Bedford, MA, USA;[6]Seattle Structural
Genomics Center for Infectious Disease (SSGCID), Seattle, WA; [7]Center for Global Infectious
Disease Research, Seattle Children's Research Institute, Seattle, WA USA; [8]Department of
Biomedical Information and Medical Education and Department of Global Health, University of
Washington, Seattle, WA USA.
23
24
25   [#]Present address: Department of Agricultural, Environmental and Food Sciences, University of
26   Molise, 86100, Campobasso, Italy

30   [*]Address correspondence to Joseph Heitman, Department of Molecular Genetics and
31   Microbiology, Duke University Medical Center, Durham, North Carolina, United States of
32   America. Phone +1 (919) 684-2824. Fax (919) 684-2790. Email heitm001@duke.edu.

Abstract

The skin of humans and animals is colonized by commensal and pathogenic fungi and

bacteria that share this ecological niche and have established microbial interactions. *Malassezia*

are the most abundant fungal skin inhabitant of warm-blooded animals, and have been implicated

in skin diseases and systemic disorders, including Crohn's disease and pancreatic cancer.

Flavohemoglobin is a key enzyme involved in microbial nitrosative stress resistance and nitric

oxide degradation. Comparative genomics and phylogenetic analyses within the *Malassezia* genus

revealed that flavohemoglobin-encoding genes were acquired through independent horizontal

gene transfer events from different donor bacteria that are part of the mammalian microbiome.

Through targeted gene deletion and functional complementation in *M. sympodialis*, we

demonstrated that bacterially-derived flavohemoglobins are cytoplasmic proteins required for

nitric oxide detoxification and nitrosative stress resistance under aerobic conditions. RNAseq

analysis revealed that endogenous accumulation of nitric oxide resulted in upregulation of genes

involved in stress response, and downregulation of the MalaS7 allergen-encoding genes. Solution

of the high-resolution X-ray crystal structure of *Malassezia* flavohemoglobin revealed features

conserved with both bacterial and fungal flavohemoglobins. *In vivo* pathogenesis is independent

of *Malassezia* flavohemoglobin. Lastly, we identified additional 30 genus- and species- specific

horizontal gene transfer candidates that might have contributed to the evolution of this genus as

the most common inhabitants of animal skin.

Key words: *Malassezia*; Flavohemoglobin; Horizontal gene transfer

Significance statement

Malassezia species are the main fungal components of the mammalian skin microbiome

and are associated with a number of skin disorders. Recently, *Malassezia* has also been found in

association with Crohn's Disease and with pancreatic cancer. The elucidation of the molecular

bases of skin adaptation by *Malassezia* is critical to understand its role as commensal and pathogen.

In this study we employed evolutionary, molecular, biochemical, and structural analyses to

demonstrate that the bacterially-derived flavohemoglobins acquired by *Malassezia* through

horizontal gene transfer resulted in a gain of function critical for nitric oxide detoxification and

resistance to nitrosative stress. Our study underscores horizontal gene transfer as an important

force modulating *Malassezia* evolution and niche adaptation.

Introduction

The skin microbiome includes numerous microorganisms that establish a variety of direct and indirect interactions characterized by the exchange of genetic material that impact microbial biology contributing to their speciation and evolution. *Malassezia* is the most abundant fungal genus resident on human skin, representing more than 90% of the skin mycobiome (1). This genus presently consists of 18 diverse species (2), each with an unusually compact genome that underwent extensive gene turnover events as a result of evolutionary adaptation and colonization to a nutrient-limited ecological niche such as the skin (3). Although commensals, *Malassezia* species are also associated with a number of clinical skin disorders, including pityriasis versicolor, dandruff, and atopic dermatitis (AD) (4). A recently-developed epicutaneous murine model revealed that the host responses to *Malassezia* are dominated by pro-inflammatory cytokine IL-17 and related factors that prevent fungal overgrowth and exacerbate inflammation under atopy-like conditions (5). Furthermore, *Malassezia* species have also been implicated recently as causal agents of Crohn's Disease/Inflammatory Bowel Disease in patients with *CARD9* mutations, and in accelerating the progression of Pancreatic Adenocarcinoma in murine models and in humans, and cystic fibrosis pulmonary exacerbation (6-8).

Nitric Oxide (NO) is a reactive compound of central importance in biological systems and it functions both as a signaling and toxic molecule. While little is known about NO synthesis in fungi, in mammals NO is synthesized by NO synthases (NOS isoforms). Nos1 and Nos3 are constitutively expressed in neurons and endothelium, respectively, and produce NO to promote S-nitrosylation and transcriptional regulation. S-nitrosylation is a post-translational mechanism involving oxidative modification of cysteine by NO, and this is the central NO-mediated signaling mechanism that affects myriad of cellular physiological and pathophysiological processes (9). On the other hand, the Nos2 is not constitutively expressed but is induced in inflammatory cells in response to infection and is involved in wound healing, immune regulation, and host defense (10).

In fungi, NO is synthesized through a reductive denitrification pathway from nitrite, and through an oxidative pathway from L-arginine, although the detailed biochemical mechanisms have not yet been fully elucidated (11-13). Compared to mammals, plants, and bacteria, the role of NO in fungal biology is understudied. In *S. cerevisiae* NO is important for activation of transcription factors that are involved in resistance to a variety of environmental stress conditions, such as oxidative stress, heat shock, and hydrostatic pressure (11). Other studies report an

involvement of NO in pathogenesis of *Botrytis cinerea* and *Magnaporthe oryzae*, in
morphogenesis and reproduction in *Aspergillus nidulans*, and in the yeast-to-hyphae dimorphic
transition in *Candida albicans* (12, 14, 15).

Imbalance in cellular NO levels leads to altered redox homeostasis, resulting in the
production of reactive nitrogen species that are responsible for nitrosative stress (10). NO
dioxygenases are enzymes that living cells use to actively consume poisonous NO by converting
it to inert nitrate, a source of nitrogen (16). Red blood cell hemoglobin is the main mammalian
dioxygenase that metabolizes NO in the vascular lumen, whereas a type I flavohemoglobin
constitutes the main enzyme deployed by microbes to counteract NO toxicity (10, 11). Some fungi
within the *Aspergillus* genus have two type I flavohemoglobins, one that is cytosolic and protects
the cells against exogenous NO, and another that is mitochondrial and is putatively involved in
detoxification of NO derived from nitrite metabolism (17, 18). A type II flavohemoglobin has also
been identified in *Mycobacterium tuberculosis* and other actinobacteria, but it lacks NO consuming
activity and it utilizes D-lactate as an electron donor to mediate electron transfer (19, 20).

Evolution of flavohemoglobins in microbes has been previously investigated, revealing a
dynamic distribution across bacteria and eukaryotes characterized by frequent gene loss, gene
duplication, and horizontal gene transfer (HGT) events (21-23). An interesting finding of these
phylogenetic studies was the HGT-mediated acquisition of a bacterial flavohemoglobin-encoding
gene, *YHB1*, by *M. globosa* and *M. sympodialis* (21, 22). Here, we employed evolutionary,
molecular, biochemical, and structural analyses to demonstrate that the HGT of the bacterial
flavohemoglobin in *Malassezia* resulted in a gain of function critical for resistance to nitrosative
stress and detoxification of NO under aerobic conditions. Moreover, analysis of the available
Malassezia genomes revealed that extant flavohemoglobin-encoding genes are present as a single
copy in different species, and resulted from a complex pattern of differential retention/loss of one
of two genes (*YHB1* and *YHB101*), which were ancestrally acquired by *Malassezia* through HGT
from different donor bacterial lineages. Through trans-species complementation we demonstrated
that the second bacterially-derived flavohemoglobin identified, Yhb101, restores resistance of the
*yhb1*Δ mutant to nitrosative stress-inducing agents and also is able to consume NO. RNAseq
analysis revealed that endogenous accumulation of NO results in upregulation of genes involved
in stress responses and transport, and downregulation of the allergen-encoding genes. The
characterization of the X-ray crystal structure of the *Malassezia* flavohemoglobin revealed features

shared with bacterial and fungal flavohemoglobins. Moreover, *ex vivo* and *in vivo* experiments
suggest that *Malassezia* flavohemoglobins are dispensable for pathogenesis under our tested
experimental conditions. Because the HGT-mediated acquisition of the flavohemoglobin-
encoding genes conferred the ability to metabolize NO, *Malassezia* genomes were searched for
other HGT that could represent important gain of function events. Thirty additional genus- and
species-specific HGT events were identified, with the donors being predominantly Actinobacteria
and Proteobacteria. Similar to *Malassezia*, these donor species are some of the most common
members of human and mammalian microbiomes, suggesting that niche overlap may have
enhanced the opportunity for inter-kingdom HGT.

Results

***Malassezia* flavohemoglobin-encoding genes were ancestrally acquired from bacteria through independent HGT events.** Flavohemoglobins are critical for nitric oxide (NO) detoxification and counteract nitrosative stress (10). Previous studies reported that the flavohemoglobin-encoding gene *YHB1* was acquired by *M. globosa* and *M. sympodialis* through HGT from *Corynebacterium*, a bacterial genus within the Actinobacteria that includes species that are part of the human microbiome (21, 22).

Because flavohemoglobins are widespread in both bacteria and eukaryotes, we examined whether the remaining 13 sequenced *Malassezia* species also contain a flavohemoglobin-encoding gene, and whether it had a fungal or bacterial origin. BLAST analyses with the *M. globosa* Yhb1 sequence as query identified a single copy of *YHB1* in all *Malassezia* species and strains with sequenced genomes. Intriguingly, this comparative search revealed that the best hits in *M. yamatoensis* and *M. slooffiae* had lower *E*-values (5e-64 and 1e-59, respectively) compared to the remaining *Malassezia* species (*E*-values ranging from 0.0 to 7e-161), possibly suggesting different origins or modifications of these flavohemoglobin-encoding genes. To elucidate the evolutionary trajectory of flavohemoglobin within the whole *Malassezia* genus, a Maximum Likelihood (ML) phylogenetic tree was reconstructed using >2000 Yhb1 bacterial and fungal sequences retrieved from GenBank. This phylogenetic analysis revealed two clades of flavohemoglobin within *Malassezia* genus: clade 1 that includes 13 species and clusters together with *Brevibacterium* species belonging to Actinobacteria; and clade 2 that includes *M. yamatoensis* and *M. slooffiae* and clusters together with different Actinobacteria with the closest relative being *Kocuria kristinae* (Fig. 1 A-C). To evaluate the statistical phylogenetic support for the two *Malassezia* flavohemoglobin clades whose distribution was not monophyletic, we performed approximately unbiased (AU) comparative topology tests. The constrained ML phylogeny, in which all *Malassezia* flavohemoglobins were forced to be monophyletic was significantly rejected (AU test, *P* value = 0.001, Fig. 1D), thus not supporting the null hypothesis that all flavohemoglobin genes in *Malassezia* have a single origin.

To validate this further, a region of ~30 kb surrounding the flavohemoglobin encoding gene in all sequenced *Malassezia* species was subjected to synteny comparison (Fig. 2). Overall, this region was highly syntenic across species, with the exception of some lineage-specific rearrangements located upstream of *YHB1* (Fig. 2B). Remarkably, while the same regions were

also highly conserved in *M. yamatoensis* and *M. slooffiae*, they both lacked the flavohemoglobin-

encoding gene, which is instead located in a different, non-syntenic, regions of their genomes (Fig.

2B and Fig. S1). Therefore, both phylogenic and synteny comparisons strongly support the

hypothesis that *Malassezia* flavohemoglobin genes were acquired through independent HGT

events from different bacterial donor species. We named the flavohemoglobin of clade 1 Yhb1

following the *S. cerevisiae* nomenclature (24), and that of *M. yamatoensis* and *M. slooffiae* Yhb101

(clade 2). The two different flavohemoglobins protein sequences share 38% identity (Fig. S2).

Because several studies suggest that genomic regions flanking horizontally-acquired genes

are enriched in DNA transposons and retrotransposons (25, 26), a 5 kb region surrounding the

flavohemoglobin-encoding genes was analyzed in two *Malassezia* species representative of clade1

(*M. sympodialis*) and clade 2 (*M. slooffiae*). Dot plot comparisons revealed overall high co-

linearity with the common flanking genes encoding a hypothetical protein and Nsr1 (Fig. S3A).

Interestingly, a highly repetitive sequence that shares similarity with the long terminal repeat

(LTR) Gypsy was identified in the *NSR1* gene flanking *YHB1* (Fig. S3A-B), and we speculate that

this LTR-like region might have facilitated the non-homologous end joining (NHEJ) integration

of the bacterial *YHB1* gene into the *Malassezia* common ancestor.

The flavohemoglobin-encoding gene *YHB101* in *M. yamatoensis* and *M. slooffiae* seems

to have been acquired in a single HGT event from the same bacterial donor lineage (Fig. 1A-C),

although its genomic location is not syntenic in these two species (Fig. 2B). In *M. slooffiae*,

YHB101 is located in a region that is otherwise highly conserved across *Malassezia* (Fig. S1A-B)

and is devoid of transposable elements or repetitive regions that could have facilitated NHEJ of

YHB101 (Fig. S1B). In contrast, in *M. yamatoensis YHB101* is located at the end of a chromosome

and the adjacent genes are not syntenic in other *Malassezia* species, with the exception of a more

distant group of five genes (from *JLP1* to *MSS1*) (Fig. S1C). In other *Malassezia* species (e.g. *M.*

japonica, M. slooffiae, M. sympodialis) these five genes are subtelomeric, suggesting that

chromosomal reshuffling might have contributed to generate the unique arrangement of genes

surrounding *M. yamatoensis YHB101* (Fig. 2; Fig. S1C).

Based on the analyses performed and on the availability of *Malassezia* genomes, we

propose the following evolutionary model of flavohemoglobin-mediated HGT in *Malassezia*.

First, the *YHB1* and *YHB101* genes were independently acquired by the *Malassezia* common

ancestor via HGT from a *Brevibacterium*-related and a *Kocuria*-related bacterial donor,

respectively. An early loss of the *YHB101* subsequently occurred in the common ancestor of the
lineages that include *M. sympodialis* and *M. globosa* (Clades A and B, Fig. 2A), which retained
the *YHB1* gene in its ancestral location; lack of synteny in the region upstream of the *YHB1* gene
in *M. globosa* and *M. restricta* represents a more recent chromosomal rearrangement (Fig. 2B). In
the early-branching *Malassezia* lineage (Clade D, Fig. 2A), the *YHB1* gene was lost in *M. slooffiae,*
which instead retained the *YHB101* gene, presumably in its ancestral location; conversely, *M.*
cuniculi lost the *YHB101* gene and retained the *YHB1* gene in the ancestral location. Lastly, all
species within the *M. furfur* lineage (Clade C, Fig. 2A) have the *YHB1* gene in its ancestral
location, with the exception of *M. yamatoensis* that has lost this gene and retained instead the
YHB101 gene, which was then relocated from its original position to a subtelomeric region (Fig.
2B). This model implies that *M. vespertilionis, M. japonica, M. obtusa*, and *M. furfur* have
independently lost the *YHB101* gene during their evolution (Fig. 2A). Because none of the
Malassezia species has the two flavohemoglobin genes (*YHB1* and *YHB101*) in their genomes, we
posit that loss of one or the other flavohemoglobin may be a consequence of different selection
pressures across descendant lineages of the HGT recipient.

Bacterially-derived flavohemoglobin-encoding genes are required for nitrosative stress
resistance and NO detoxification in *Malassezia*. Flavohemoglobins are critical for NO
detoxification and counteract nitrosative stress (10). To assess whether this HGT event in
Malassezia resulted in a gain of function, we deleted the *YHB1* ORF (*MSYG_3741*) of *M.*
sympodialis ATCC42132 through targeted mutagenesis using our recently developed
transformation protocol based on transconjugation mediated by *Agrobacterium tumefaciens* (27,
28) (Fig. S4A).
The *M. sympodialis yhb1Δ* mutant exhibits hypersensitivity to the NO-donors DETA
NONOate and sodium nitrite (NaNO$_2$), but not to hydrogen peroxide (H$_2$O$_2$) (Fig. 3A). The two
identified *Malassezia* flavohemoglobins Yhb1 and Yhb101 were used to generate GFP fusion
proteins whose expression was driven by the respective endogenous promoter to complement the
M. sympodialis yhb1Δ mutant phenotype and to assess protein localization (Fig. S4B).
Reintroduction of either flavohemoglobin in the *M. sympodialis yhb1Δ* mutant restored resistance
to nitrosative stress at the WT level (Fig. 3A). In agreement, fusion protein expression in
complemented strains was confirmed by qPCR (Fig. S4C-D), FACS, and fluorescence microcopy

imaging of GFP expression, which revealed that *Malassezia* flavohemoglobins are cytoplasmic

(Fig. 3B).

Next, we tested whether *Malassezia* flavohemoglobins were able to actively detoxify NO,

which could potentially account for its involvement in nitrosative stress resistance. To this aim,

we adapted a biochemical assay used for evaluating NO consumption by hemoglobin in red blood

cells and plasma (29) to the commensal yeast *Malassezia* (Fig. S5). As shown in Figure 2C, while

the *M. sympodialis* WT strain exhibited robust and dose-dependent NO degradation, the *yhb1Δ*

mutant showed no NO consumption. Complemented strains were able to actively consume NO,

although *M. sympodialis yhb1Δ + YHB101-GFP* displayed a lower NO consumption,

corroborating the results of the phenotypic assay, GFP expression, and FACS analysis (Fig. 3A-

C). These differences might be due to less efficient cross-species complementation of the *M.*

yamatoensis Yhb101-GFP fusion protein in the *yhb1Δ* mutant of *M. sympodialis*, which was a

strategy chosen because of the lack of protocols for gene deletion in *M. slooffiae* and *M.*

yamatoensis. Taken together these genetic and biochemical analyses show that the bacterially-

derived flavohemoglobins protect *Malassezia* from nitrosative stress by decreasing toxic levels of

NO.

To assess intracellular production of NO by *M. sympodialis*, cells were stained with the

NO-specific dye 4-Amino-5-methylamino- 2′,7′-diaminofluorescein diacetate (DAF-FM DA),

which passively diffuses across membranes and emits increased fluorescence after reacting with

NO. Fluorescent microscopy revealed intracellular accumulation of NO in both the WT and *yhb1Δ*

mutant of *M. sympodialis* (Fig. 3D). NO-staining was quantified by FACS analysis, revealing

significantly higher NO accumulation in the flavohemoglobin mutant *yhb1Δ* compared to the *M.*

sympodialis WT (Fig. 3E-F). Because DAF-FM DA and GFP have similar excitation/emission

spectra, complemented strains could not be tested for NO accumulation via flow cytometry, and

therefore an independent *M. sympodialis yhb1Δ* mutant was tested and yielded similar results.

These results indicate that the lack of a functional flavohemoglobin leads to intracellular

accumulation of NO.

Finally, a broader analysis was performed to assess other functions of the *Malassezia*

flavohemoglobins in response to a variety of environmental stresses and clinical antifungals, but

in all cases the *M. sympodialis yhb1Δ* mutant phenotype was not significantly different from the

WT (Fig. S6). These phenotypic results are in agreement with those obtained for the

basidiomycetous yeast *Cryptococcus neoformans* (30, 31), but contrast with the studies carried out in ascomycetous fungi, in which in addition to NO and nitrosative stress sensitivity, flavohemoglobin mutants exhibited higher resistance to hydrogen peroxide in *A. nidulans* (32), and hyper-filamentation in *C. albicans* (33). Several studies also report the protective role of both bacterial and fungal flavohemoglobins against NO under anaerobic conditions (10, 24, 34). However, we could not confirm this function for the *Malassezia* flavohemoglobin in our anaerobic experiments because no phenotypic differences were observed between the WT, the *yhb1*Δ mutant, and the complemented strains (Fig. S7).

A recent inactivation of *YHB1* in *M. nana* results in compromised NO enzymatic consumption. Analysis of Yhb1 protein prediction across species revealed that *M. nana YHB1* underwent pseudogenization [i.e. loss of gene function by disruption of its coding sequence with generation of a pseudogene, which is usually indicated as ψ (35)] following a G-to-T transversion in the glycine codon GGA, generating a premature TGA stop codon at the 29[th] amino acid (Fig. S8A). Literature search revealed that the sequenced strain of *M. nana* CBS9557 was isolated in Japan from a cat with otitis externa (36), while the other four known *M. nana* strains were collected in Brazil: *M. nana* CBS9558 and CBS9559 from cows with otitis externa, and CBS9560 and CBS9561 from healthy cows (36).

The *M. nana* strains CBS9557, CBS9559, and CBS9560 were used to investigate whether the pseudogenization event occurred in a *M. nana* ancestor, and whether it impacts nitrosative stress resistance and NO consumption. Because no genomes are available for the *M. nana* strains CBS9559 and CBS9560, their *YHB1* gene was amplified by PCR and Sanger sequenced using primers designed on the *YHB1* of *M. nana* CBS9557 (Table S2). *YHB1* sequence comparison confirmed a premature stop codon present in only CBS9557, with both Brazilian *M. nana* isolates having a full-coding *YHB1* gene (Fig. S8A). Phenotypic analysis revealed no significant difference in resistance to nitrosative stress by the three *M. nana* strains, with only a modest increased sensitivity displayed by CBS9557 exposed to 10 mM of sodium nitrite (Fig. S8B). Strikingly, *M. nana* CBS9557 displayed undetectable NO consumption activity as observed for the *M. sympodialis yhb1*Δ mutant, while *M. nana* CBS9559 and CBS9560 showed regular dose-dependent NO consumption (Fig. S8C). These data suggest that the inactivation of flavohemoglobin in *M. nana* CBS9557 impaired the ability to consume NO, but this does not

impact the resistance to nitrosative stress, which might be compensated by other stress responsive
pathways.

Intriguingly, another pseudogenization event of a bacterial gene encoding an aliphatic
amidase was also identified in *M. nana* CBS9557 (represented in Fig. 6). These nonsense
mutations were identified only in the *M. nana* CBS9557 isolated in Japan, suggesting that the
different origin of the *M. nana* strains might contribute to this intraspecies diversity. This
hypothesis is further supported by different phenotypic traits displayed by the *M. nana* isolates
(Fig. S9). Exposure to several stress conditions revealed different responses to the most common
antifungal drugs by *M. nana* strains, with strain CBS9557 displaying increased sensitivity to
amphotericin B and resistance to fluconazole, and the geographically-related strains CBS9559 and
CBS9560 displaying an opposite phenotype (Fig. S9).

NO accumulation in *M. sympodialis* leads to upregulation of genes involved in nitrogen
metabolism, ergosterol biosynthesis, and protein folding, and downregulation of predicted
pathogenicity factors. Because the *M. sympodialis* flavohemoglobin mutant *yhb1Δ* accumulates
higher amounts of NO than the WT (Fig. 3 D-E), we compared their transcriptomic profile to
elucidate any potential signaling role of endogenous NO. RNAseq analysis revealed 36
differentially expressed genes for false discovery rate (FDR) <0.05, of which 14 were upregulated
and 22 were downregulated; using an additional threshold of $\log_2$FC +/- 0.5 we found 3
upregulated and 9 downregulated genes (Fig. 4A; Dataset S1-S2). Of these, the only upregulated
gene with $\log_2$FC > 1 encodes an uncharacterized protein (*MSYG_1280*), while two others with
0.5 < $\log_2$FC < 1 encode Nop56 (or Sik1), a nucleolar protein involved in pre-rRNA processing,
and an uncharacterized protein (*MSYG_0148*) predicted to be involved in magnesium transport.
Other known upregulated genes with $\log_2$FC <0.5 are involved in response to stresses and transport
(Dataset S1). The majority of the downregulated genes include those encoding hypothetical
proteins (5 out of 9), the regulator of phospholipase D Srf1, two MalaS7 allergens, and an
uncharacterized allergen (Dataset S2). It is worth noting that a large number of differentially
expressed genes (DEGs) are predicted to encode unknown proteins, suggesting novel and unknown
signaling pathways regulated by endogenous NO in *Malassezia* (Dataset S1-S2).

Next, to elucidate the global transcriptomic response of *M. sympodialis* exposed to
nitrosative stress, RNAseq analysis for *M. sympodialis* WT cells treated with sodium nitrite was

performed. Compared to the untreated control, 112 genes were upregulated and 50 were
downregulated (FDR<0.05; $\log_2$FC +/-0.5) (Fig. 4B). The most expressed genes included *HEM1*
encoding a 5-aminolevulinate synthase involved in heme biosynthesis, *MSYG_3126* encoding a
hypothetical secreted lipase, the allantoicase encoding gene *DAL2, MSYG_3153* encoding an
uncharacterized NAD(P)/FAD-dependent oxidoreductase, and *DCG1* encoding a protein with
unknown function predicted to be related to nitrogen metabolism (Fig. 4B; Dataset S3). The
flavohemoglobin-encoding gene *YHB1* was significantly upregulated for FDR <0.05, but it had
low expression level ($\log_2$FC=0.33). Low expression of *YHB1* was also observed in *S. cerevisiae*
cells exposed to nitrosative stress (37), although its role in NO consumption has been well
characterized (24). The most represented classes of upregulated genes are involved in stress
resistance, cellular detoxification and transport, and metabolism (Fig. 4C). Functional protein
association network analysis revealed enrichment of genes involved in nitrogen metabolism and
regulation, ergosterol biosynthesis, and heat shock response (Fig. S10); we speculate that among
the upregulated transcription factor encoding genes (*HSF1, UPC2, BAS1*, and *HMS1*), the heat
shock factor Hsf1 might be the key candidate that activates nitrosative stress responsive genes,
given its known role in response to stresses in other fungi (38, 39). Conversely, response to
nitrosative stress is mediated by the transcription factors Yap1 and Msn2/Msn4 in *S. cerevisiae*
and *Schizosaccharomyces pombe* (37, 40), and by the transcription factor Cta4 and the Hog1
kinase in *C. albicans* (41), with the consequent activation of genes known to be required for
oxidative stress response, such as those involved in glutathione turnover and other anti-
oxidant/detoxification systems. *M. sympodialis CTA1* and *CCP1* are the only oxidative stress
responsive genes activated in response to nitrosative stress (Fig. S10; Dataset S3).
The most represented GO category of downregulated genes encodes integral components
of membrane, which includes transporters and putative *Malassezia* allergens; other downregulated
genes are involved in calcium metabolism, protein folding, and proteolysis. Two transcription
factors were downregulated, and they include the pH responsive Rim101, and an uncharacterized
bZIP transcription factor (Fig. 4C, Dataset S4).
Comparison of the two different RNAseq datasets revealed two common upregulated
genes, encoding the glycerol dehydrogenase Gcy1 and the catalase Cta1, and 10 downregulated
genes that include 4 *Malassezia* allergens, a putative secreted lipase, and 5 hypothetical proteins

(Fig. 4D). While it is not surprising to find upregulation of a detoxifying enzyme such as catalase,

it is intriguing to find downregulation of genes encoding predicted pathogenicity factors.

In conclusion, our transcriptomic data indicate that the response of *Malassezia* to NO and

nitrosative stress is mostly different from other studied fungi and it involves metabolic pathways

and genes that were not known to be relevant to overcome nitrosative stress.

***Malassezia* flavohemoglobin has characteristic features of both bacterial and fungal**

flavohemoglobins. We hypothesized that the structure of a protein acquired by HGT will likely

remain similar to that of the donor organism in order to retain its original function. Attempts to

resolve the crystal structures of both *Malassezia* flavohemoglobins were carried out, but only the

M. yamatoensis flavohemoglobin Yhb101 formed crystals to be analyzed. The structure was

determined *de novo* by SAD phasing off the heme-iron bound to the globin domain of the protein

(Table S3). The flavohemoglobin structure is highly conserved with previously characterized

structures of this enzyme family, and it consists of an N-terminal globin domain coordinating an

iron-bound (Fe^{2+}) heme and a C-terminal reductase domain with both FAD- and NAD-binding

sub-domains, of which only FAD is bound (Fig. 5A). An overlay of a flavohemoglobin structure

from *E. coli* and *S. cerevisiae* on the *M. yamatoensis* crystal structure highlights conserved binding

sites between the proteins (Fig. 5B). Alignment of the globin domains between literature and

experimental structures resulted in an RMSD value of 1.532Å for *E. coli* and 1.434Å for *S.*

cerevisiae, mostly resulting from slight shifts in the D-loop and E-helix between the structures

compared. Common to all structures analyzed is the histidine residue coordinating with the heme

iron from the proximal side. This member of the catalytic triad is supported by tyrosine (Tyr98)

and glutamate (Glu140) residues conserved in sequence and structure between bacterial, and

fungal/yeast flavohemoglobins (42, 43). In *M. yamatoensis*, as also observed in *E. coli*, the heme

iron is ligated by 5 atoms: 4 from the heme and His88 from the F-helix. Substrates commonly bind

on the distal side of the heme and lead to a conformational change in the planarity of the heme

molecule. The E-Helix on the distal side of the heme molecule contributes Leu58, a conserved

residue which approaches the heme-bound iron from 3.7Å away. At this position, the 6[th]

coordination site for the iron is occluded, again similar to the *E. coli* crystal structure, but unlike

the yeast structure where a three-atom small molecule co-crystallized.

In *M. yamatoensis* Yhb101, the D-loop acts as a bridge between the C- and E-helices and
the interface between the bound FAD and heme. Comparison of the D-loops from these structures
shows the *M. yamatoensis* D-loop adopts a nearly identical helical structure as that of *S. cerevisiae*,
in contrast to the *E. coli* D-loop**,** which is more extended. The *M. yamatoensis* E-helix also adopts
a ~30° bend immediately following Leu58, which may straighten out once a substrate is bound.
This structural adjustment likely communicates substrate binding near the heme to the reductase
domain through movements in the D-loop as the heme B pyrrole proprionate forms a hydrogen
bond with the main-chain NH of Ser45, the first residue in the D-loop (Fig. S11).

Lastly, in Fig. S12 a detailed comparison of the functional residues is shown between the
Malassezia flavohemoglobins with those of the closer HGT donor bacteria *B. ravenspurgense*, *K.*
kristinae, R. nasimurium, and with the model yeast *S. cerevisiae.*

***Malassezia* flavohemoglobins are not required for survival on the host.** Previous studies in
human fungal pathogens indicate that flavohemoglobins are required for pathogenesis (30, 33). In
our experiments we found that *M. sympodialis* WT, *yhb1Δ* mutant, and *yhb1Δ + YHB1* and *yhb1Δ*
+ YHB101 complemented strains have similar levels of survival within activated macrophages
(Fig. S13A-B). This result is in contrast with previous findings in *C. neoformans* (30), but it
corroborates results obtained in *A. fumigatus* (18). Furthermore, the recently developed murine
model for *Malassezia* skin infection (5) was utilized to test pathogenicity of the flavohemoglobin
strains and the induction of host response. Corroborating *ex vivo* data, we found no differences
both in terms of host tissue colonization and host inflammatory response for the *yhb1Δ* mutant
compared to the complemented strains (Fig. S13C-E, Fig. S14). In agreement, there were no
differences between WT and *Nos2-/-* mice when challenged with *M. sympodialis* WT (Fig. S13F-
H). These results suggest that flavohemoglobin is not required for pathogenesis of *Malassezia* in
an experimental skin model.

Lastly, several attempts were also carried out to test survival of *M. sympodialis* WT and
flavohemoglobin strains within the GI tract of WT and *Nos2-/-*mice. Because of the high amount
of NO produced in the GI tract of mice during inflammation (44), and the recently-reported
involvement of *M. restricta* in Crohn disease (6), we hypothesized that the flavohemoglobin would
be required for *M. sympodialis* survival in GI tract during inflammation. We followed the protocol
developed for GI tract colonization by *M. restricta* (6), but unfortunately we could never recover

any *M. sympodialis* colony. For this reason, whether *Malassezia* flavohemoglobin is required for
survival within the GI tract could not be determined.
456
Analysis of *Malassezia* genomes revealed extensive HGT events from bacteria. Given the gain
of function due to acquisition of the bacterially-derived flavohemoglobins by *Malassezia* species,
we sought to identify additional HGT candidate genes in the *Malassezia* genus. In a previous study,
8 HGT events were identified in *M. sympodialis*, and then their presence was assessed in other
species within the genus (3). In the present study we applied a previously described analytical
pipeline (45) based on three HGT metrics - the HGT index (46), the Alien Index (AI) (47) and the
Consensus Hit Support (CHS) (48) - to identify novel genus and species-specific HGT events. Our
goal was not to explicitly establish the evolutionary history of individual genes, but rather to
estimate bacteria-derived HGT candidates for the complete set of *Malassezia* genomes. Besides
recovering the *YHB1* and *YHB101* genes as HGT candidates, in addition this analysis identified a
total of 30 HGT candidate genes (Fig. 6 and Dataset S5), seven of which in common with the
previous study. HGT candidates found in the majority of the *Malassezia* species include genes
involved in broad resistance to stresses, including three that were upregulated in *M. sympodialis*
exposed to nitrosative stress (Dataset S3), such as the NAD(P)/FAD-dependent oxidoreductase-
encoding gene *MSYG_3153*, the catalase-encoding gene *MSYG_3147*, and the sorbitol
dehydrogenase-encoding gene *MSYG_0932* (Fig. 6). Other HGT candidates include a
deoxyribodipyrimidine photo-lyase predicted to be involved in repair of UV radiation-induced
DNA damage, and a class I SAM-dependent methyltransferase potentially modifying a variety of
biomolecules, including DNA, proteins and small-molecule secondary metabolites. Another
interesting HGT candidate is the gene encoding a septicolysin-like protein, which is known as a
pore-forming bacterial toxin that might play a role as virulence factor (49, 50). This gene is absent
in all *Malassezia* species phylogenetically related to *M. sympodialis*, and is present as five copies
in *M. globosa*. Furthermore, a large number of HGT events unique to *Malassezia* species of clade
A were found, and the acquired genes encode a variety of proteins with different functions, such
as hydrolysis, protein transport and folding, detoxification of xenobiotics, and resistance to
stresses. Finally, 12 of the HGT candidates identified were unique to certain *Malassezia* species.
An intriguing case is *M. japonica* for which we found four unique HGT candidates, one of them

in 3 copies. These genes encode orthologs of the fungal Gre2 protein, which is known to be
involved in responses to a variety of environmental stresses (51, 52).

Discussion

In the present study we report the functional characterization of two *Malassezia*
flavohemoglobin encoding genes that were independently acquired through HGT from different
Actinobacteria donors. Our experimental analyses demonstrate that both bacterially-derived
flavohemoglobins are involved in nitrosative stress resistance and NO degradation, consistent
with its known functions in bacteria and fungi (10).

We propose an evolutionary HGT model in which extant flavohemoglobin-encoding genes
in *Malassezia* result from a complex pattern of gene retention/loss after being both acquired by a
Malassezia common ancestor. Nevertheless, other evolutionary scenarios could also be
hypothesized, such as: 1) the acquisition of the *YHB1* in a *Malassezia* common ancestor via HGT
from a *Brevibacterium*-related donor; 2) followed by more recent acquisitions of *YHB101* by *M.
yamatoensis* and *M. slooffiae* via independent HGT events from a common, or closely related,
bacterial donor(s) (*Kocuria*). In this scenario, the "resident" *YHB1* in *M. yamatoensis* and *M.
slooffiae* could have been displaced upon secondary acquisition of the *YHB101* gene [a
phenomenon termed as xenolog gene displacement (53)], or the acquisition of *YHB101* by HGT
could have been preceded by the loss of the cognate *YHB1* copy. The identification of novel
Malassezia species and the analysis of their genomes will be key for the elucidation of these
complex models of gene evolution in *Malassezia*.

Although the mechanisms of HGT in fungi are not fully understood, several possible
mechanisms have been reported (25, 54). One such mechanism is gene acquisition through
conjugation, which requires contact between bacterial donor and fungal recipient (54). For the
HGT events that mediated flavohemoglobin acquisition by *Malassezia*, the closest phylogenetic
donors are Actinobacteria that are part of the mammalian microbiome and hence share the same
ecological niche with *Malassezia.* A dilemma that is common to all HGT events is that if a gene
is required for survival in a certain condition, its transfer under that condition might in theory
be difficult if not impossible (55). Because NO is synthesized by mammals, including by the
skin (56), we speculate that the presence of NO enhanced the HGT transfer of bacterial

flavohemoglobins to a fungal *Malassezia* ancestor that acquired the ability to actively consume
NO.

Notably, a large number of eukaryotic organisms including fungi lack a
flavohemoglobin-encoding gene, suggesting the existence of alternative pathways for
nitrosative stress resistance and NO utilization. For example, in *Histoplasma capsulatum*, the
etiologic agent of histoplasmosis, Yhb1 is replaced by a P450-type NO reductase (57), whereas
in other cases, such as for the basidiomycetous fungi *Moniliella, Ustilago,* and *Puccinia*, NO
metabolism in the absence of flavohemoglobin has yet to be elucidated. The evolution and
diversification of flavohemoglobin-encoding genes has been a dynamic and complex process
characterized by several prokaryote-prokaryote and prokaryote-eukaryote HGT events (22),
hence suggesting its significant contribution for habitat colonization by a species but likely a
dispensable role in evolutionary divergence.

Is the bacterial flavohemoglobin required for *Malassezia* interaction with the host?
While a number of studies in bacteria and fungi reported a role for flavohemoglobin in microbial
pathogenesis (10), we surprisingly found that *Malassezia* flavohemoglobins are dispensable for
survival within macrophages and for skin infection in our experimental conditions. Conversely,
we propose that *Malassezia* flavohemoglobins are important for the commensal lifestyle of
Malassezia through regulation of NO homeostasis, a hypothesis corroborated by
downregulation of genes-encoding putative virulence factors (i.e. allergens and lipases) in our
transcriptomic analyses. Another hypothesis is that the HGT-mediated acquisition of
flavohemoglobins might be important to mediate *Malassezia* response to NO that is produced
by sympatric microbial communities and acts as a quorum signaling molecule, as reported in
bacteria (58, 59) and in *S. cerevisiae* (60).

Horizontal gene transfer is thought to occur much less frequently in eukaryotes than in
prokaryotes (61, 62), but there are notable cases that invoke HGT as a prominent mechanism
of eukaryotic evolution, such as in the transition of green plants from aquatic to terrestrial
environments (63), and in the colonization of the animal digestive tracts by rumen fungi and
ciliates (55, 64, 65). Analysis of *Malassezia* genomes revealed a large number of HGT events,
suggesting that they may also have played a substantial contribution in *Malassezia* evolution
and niche adaptation. Donor bacteria include those that are part of the microbiota of animals, but
also others that are known to inhabit a variety of terrestrial and marine habitats, raising questions

about a possible wider environmental distribution of a *Malassezia* ancestor. This could be
correlated with the presence of *Malassezia* DNA in a number of unexpected areas, such as in
association with corals and sea sponges in the ocean (66). Moreover, most of the HGT candidate
genes identified in *Malassezia* operate as a self-contained metabolic unit, which has been
proposed to facilitate HGT (22). Intriguingly, the high number of HGT events suggests also a
predisposition of *Malassezia* to bacterial conjugation, in line with our previous findings that *A.*
tumefaciens-mediated transformation is the only effective technique for molecular manipulation
of *Malassezia* (67). There are a number of identified HGT that are predicted to be important for
Malassezia pathophysiology and that can be characterized using the methodologies reported in
the present study.

Materials and Methods

Strains used in the present study. The *M. sympodialis* strains ATCC42132 was utilized as a

model species for genetic manipulations. In addition, *M. yamatoensis* CBS9725 (or MY9725) and

M. nana strains CBS9557, CBS9559, and CBS9560 were employed for NO consumption assays.

These strains were grown on modified Dixon's media (mDixon), which is the medium routinely

used for culturing *Malassezia* species (3).

Identification of *YHB1* genes in *Malassezia* genomes and synteny analyses. Protein-coding

sequences were obtained using the *ab initio* gene predictor AUGUSTUS v3.2.2 (68) for each

Malassezia species that lacked a genome annotation (as of September 2017). To search for *YHB1*

homologs, local BLAST databases were set up for both the genome assemblies and the translated

coding sequences, and the *M. globosa* Yhb1 protein sequence (GenBank XP_001730006.1) was

queried against each database using BLASTP or TBLASTN. Top BLAST-identified protein

sequences were retrieved, and the presence of the typical FHbs domains, consisting of an N-

terminal globin domain fused with a C-terminal FAD- and NAD-binding oxidoreductase modules,

were inspected using InterProScan 5 (69). For the analysis of relative gene-order conservation

(synteny) between *Malassezia* species, a region of ~ 32 kb surrounding the *YHB1* gene (~7-8 genes

on each side) was carefully examined and compared using top scoring BLASTP against the

reference genome annotations of *M. sympodialis*, *M. globosa* and *M. pachydermatis*.

Yhb1 phylogeny and topology tests. The *Malassezia globosa* Yhb1 protein was queried against

the Genbank non-redundant (nr) protein database (last accessed in August 2017) with BLASTP

and an e-value inclusion threshold of 1e-10. Protein sequences corresponding to the top 5000 hits

were extracted and utilized for downstream analyses. In addition, all putative *Malassezia* Yhb1

proteins identified from the genome data, as well as the functionally characterized Yhb1 gene of

S. cerevisiae, were included in the dataset. Highly similar sequences were collapsed with CD-HIT

v4.7 (70) using a sequence identity threshold of 0.95 (-c 0.95) and word length of 5 (-n 5) to remove

redundancy and correcting the bias within the dataset. Sequences were aligned with MAFFT

v7.310 using the FFT-NS-i strategy (71) and poorly aligned regions were trimmed with TrimAl

v1.4. (-gappyout) (72). The Yhb1 phylogenetic tree was constructed using IQ-TREE v1.5.5 (73)

and the LG+F+I+G4 amino acid model of substitution as determined by ModelFinder (74).

Because the dataset consisted of a large number of relatively short sequences, we reduced the
perturbation strength (-pers 0.2) and increased the number of stop iterations (-nstop 500) during
tree search. Branch support was accessed with the ultrafast bootstrap approximation (UFboot) (75)
and the SH-like approximate likelihood ratio (SH-aLRT) test (76) both with 10,000 replicates (-
bb 10000 –alrt 10000). Tests of monophyly were performed in CONSEL version 1.2 (77) using
the approximately unbiased (AU) test (78) to determine whether the maximum likelihood (ML)
estimates of the best tree given a constrained topology differed significantly from the
unconstrained best ML tree. To produce a constrained topology, *Malassezia* sequences were forced
to be monophyletic, and all other branches were resolved to obtain the maximum -log likelihood
using RAxML v8.2.11, with five alternative runs on distinct starting trees (-# 5) and the same
amino acid model of substitution. Site-wise log-likelihood values were estimated for both trees
(option: -f g in RAxML) and the resulting output was analyzed in CONSEL.

Species phylogeny. To reconstruct the phylogenetic relationship among the 15 *Malassezia* species
selected, top pair-wise BLAST results of whole proteomes were clustered by a combination of the
bidirectional best-hit (BDBH), COGtriangles (v2.1), and OrthoMCL (v1.4) algorithms
implemented in the GET_HOMOLOGUES software package (79), to construct homologous gene
families. The proteome of *M. sympodialis* ATCC42132 served as reference and clusters containing
inparalogs (i.e. recent paralogs defined as sequences with best hits in its own genome) were
excluded. A consensus set of 246 protein sequences was computed out of the intersection of the
orthologous gene families obtained by the three clustering algorithms using the perl script
compare_clusters.pl included in the package. These single copy orthologous gene families were
individually aligned with MAFFT v7.310 using the L-INS-i strategy and trimmed with TrimAl (-
gappyout). The resulting alignments were concatenated with the python script ElConcatenero (80)
to obtain a final supermatrix consisting of a total of 134,437 amino acid sites (47,942 parsimony-
informative). The phylogenetic tree was constructed with IQ-TREE v1.5.5 and the LG+F+R5
amino acid model of substitution and branch support values were obtained from 10,000 replicates
of both UFBoot and SH-aLRT.

Horizontal gene transfer analyses. To assess the extent of horizontal transfer into *Malassezia*
genomes, we applied a previously described pipeline (45) to the set of 15 available *Malassezia*

proteomes, adjusting the parameters accordingly. In brief, protein sequences were first aligned to

the UniRef100 database (last accessed September 2017) using Diamond 'blastp' (81) with the

following parameters: '--sensitive --index-chunks 1 --max-target-seqs 500 --Evalue 1e-5'. Three

metrics were defined for each protein sequence used as query: (a) the HGT Index (46), (b) the

Alien Index (AI) (47), and (c) the Consensus Hit Support (CHS) (48). Hits in Malasseziales were

omitted (NCBI taxid 162474), and we specified 'Fungi' as the ingroup lineage (NCBI taxid 4751)

and 'non-Fungi' as outgroup. Proteins receiving an HGT index $\geq$ 30, AI index > 45, CHS $\geq$ 90,

and that have Bacteria as the donor lineage, were considered well supported HGT candidates and

analyzed further. When a given HGT candidate was detected in more than one, but not all the

species, TBLASTN searches were used to distinguish between *bona fide* gene losses or genome

mis-annotation.

Molecular manipulation of *M. sympodialis*. A detailed procedure of the cloning procedures and

transformation technique used is reported in SI Appendix, SI Material and Methods. The plasmid

for targeted mutagenesis of *M. sympodialis YHB1* was generated by cloning regions of 1.5 kb

flanking the *YHB1* gene fused with the *NAT* marker within the T-DNA (transfer DNA) of the

binary plasmid pGI3 as previously reported (67, 82). For *yhb1*Δ functional complementation, two

different GFP-fusion plasmids were generated. The *M. sympodialis YHB1* ORF, the endogenous

M. sympodialis YHB1 gene including its native promoter and terminator, or the endogenous *M.*

yamatoensis YHB101 gene including its native promoter, were amplified by PCR using chimeric

primers that generated recombination sites that allowed T-DNAs assembly in the following order:

plasmid pGI35 was p*YHB1, YHB1, GFP, t*YHB1, NEO, and it is referred as *YHB1-GFP*; plasmid

pGI31 was p*YHB101, YHB101, GFP*, t*YHB1, NEO*, and it is referred as *YHB101-GFP*. Plasmids

were recombined through *in vivo* recombination in *S. cerevisiae* (67). Correct plasmids were

identified by PCR and introduced into the *A. tumefaciens* EHA105 strain by electroporation, and

the transformants selected on LB + 50 µg/ml kanamycin.

M. sympodialis was transformed through *A.-tumefaciens* mediated transformation

following our previously published method (27, 28). Transformants were selected on mDixon

supplemented with nourseothricin (100 µg/ml) or neomycin G418 (100 µg/ml), and cefotaxime

(350 µg/ml) to inhibit *Agrobacterium* growth. Transformants were purified to single colonies, and

subjected to phenol-chloroform-isoamyl alcohol (25:24:1) DNA extraction (83). The correct

replacement of the *YHB1* target locus, as well as integration of the reconstitute versions of the

YHB1 and *YHB101* genes, were assessed by PCR. Primer sequences are listed in Table S2.

NO quantification and GFP expression analysis. Intracellular levels of NO was measured by

flow cytometry adapted from previous method (14). In brief, *M. sympodialis* WT and *yhb1Δ*

mutant were grown ON in mDixon media, washed once with PBS, and then incubated in PBS with

agitation overnight. For each sample, ~ 2 x 10^8 cells/mL were equally divided, and were stained

with DAF-FM DA at a final concentration of 10 µM, and the other half not stained and used as

background control. Cells were washed once with PBS and analyzed using a BD FACSCantoTM

II.

To measure intracellular GFP intensity by flow cytometry, *M. sympodialis yhb1Δ* mutants,

and complemented strains *yhb1Δ + YHB1-GFP* and *yhb1Δ + YHB101-GFP* were grown in

mDixon media ON, washed in PBS, and ~1 x 10^8 cells/mL used for flow cytometry analysis on

the Becton-Dickinson FACScan at Duke Cancer Institute Flow Cytometry Shared Resource. The

results were analyzed using FlowJo. *M. sympodialis yhb1Δ* mutants, and complemented strains

yhb1Δ + YHB1-GFP and *yhb1Δ + YHB101-GFP* prepared in the same way were also used for GFP

microscopy analysis carried out at the Duke Light Microscopy Core Facility using a Zeiss 710

inverted confocal microscope.

NO consumption assay. The NO consumption assay was performed as previously reported (84,

85). A custom-made glass reaction cell was filled with PBS (4 mL, pH 7.4, 37°C) and connected

with a flow meter and a TEA NO analyzer. Under near-vacuum conditions, NO generation was

achieved by injecting DETA NONOate (dissolved in 0.01 N NaOH at a concentration of 30 mM)

into the reaction cell. Under a helium flow, NO was passed through an inline condenser (removing

water vapor) to a NO chemiluminescence analyzer (TEA 810, Ellutia) to generate an

electrochemical NO baseline signal (expressed as mV) (Fig. S5).

We established the optimum amount of DETA NONOate to be injected as 30 µL (225 µM),

leading to an increase of ~60 mV in the baseline. *Malassezia* yeasts were grown ON in mDixon,

washed twice with PBS pH = 7.4, and adjusted in PBS to OD_{600}=1. Because *Malassezia* yeasts

form clumps in liquid culture, cell counts using a hemocytometer was inaccurate and the number

of viable cells in cultures was instead determined by plating an aliquot (100 µL) of the cellular

suspensions on mDixon agar and counting colonies. The density of the *Malassezia* cultures was

~2.3 x 10^8 CFU/mL, with the exception of *M. nana* cultures that were 1.5 x 10^8 CFU/mL. For NO

consumption assays, injection of a volume of PBS pH = 7.4 alone greater than 50 µL interfered

with the NO signal baseline, and therefore amounts ranging from 10 µL (2.3 x 10^6 cells injected)

to 50 µL (1.15 x 10^7 cells injected) of *Malassezia* cellular suspensions were utilized. For *M. nana*

injections of 15 µL (2.25 x 10^6 cells injected), 30 µL (4.50 x 10^6 cells injected) and 45 µL (6.75 x

10^6 cells injected) were performed. For technical reasons (i.e. presence of clumps, non-

resuspended cells, and cultures too dense for the syringe needle) we found that the indicated range

of cell counts were optimal to obtain results that were clean and reproducible.

Our standardized NO consumption assay was performed as follows: once a steady baseline

was obtained (approximately 1 min), 30 µL of DETA NONOate was injected, and after a stable

NO signal was achieved, different volumes of *Malassezia* cellular suspensions were injected into

the buffer solution (PBS + DETA NONOate), resulting in a decrease in NO signal that indicates

NO consumption. Following the injection of each *Malassezia* species, the reaction chamber was

flushed, and refilled with PBS and DETA NONOate. A maximum of 5 injections was performed

using the same buffer solution, after which it became cloudy, generating an unstable signal.

***In vitro* phenotypic characterization of *Malassezia* strains.** Phenotypic analysis of the

Malassezia strains was performed on mDixon agar by spotting 1.5 µL of 1:10 dilutions of each

cellular suspension in the following conditions: DETA NONOate (0.5 mM and 1 mM), $NaNO_2$ (5

mM and 10 mM), hydrogen peroxide (1 mM), UV (200 µJ x 100), 37°C, benomyl (20 µM), 5-

flucytosine (5FC, 50 and 100 µg/mL), amphotericin B (AmB, 50 µg/mL), caspofungin (2.5 µg/mL

and 5 µg/mL), fluconazole (FLC, 0.5 µg/mL and 1 µg/mL), NaCl (1M), LiCl (100 mM), Congo

red (0.4 µg/mL), pH = 7.5, or pH = 4.When hypoxic conditions were required, the GasPak EZ

Container System was used (BD Diagnostics). Plates of mDixon supplemented with DETA

NONOate (0.1 nM, 1nM, 10 nM and 0.1 mM) and $NaNO_2$ (10 nM, 50 nM, 0.1 mM and 1 mM)

were spotted as reported above and placed in the GasPak Large Incubation Chamber with three

anaerobe sachets added prior to sealing the chamber. Maintenance of hypoxic conditions (less than

1% O2 and greater than 13% CO2) was monitored using Dry Anaerobic Indicator Strips (BD).

RNAseq analysis. 10 mL liquid mDixon cultures of ~ 1×10^7 cells/mL of *M. sympodialis* WT ATCC42132 with and without 10 mM NaNO$_2$, and *M. sympodialis yhb1Δ* mutant were grown for ON on shaking culture at 30°C. Cells were pelleted at 3,000 rpm in a table top centrifuge for 3 minutes, washed with 10 mL dH2O, and the cell pellet was lyophilized and stored at -80°C until RNA extraction. Cells pellets was broken with sterile beads and RNA extracted with TRIzol (ThermoFisher) according to the manufacturer's instructions. Final RNA pellets were treated with TURBO DNase (ThermoFisher, catalog # AM2238) according to manufacturer's instructions, and RNA was resuspended in 50 µl of nuclease free water. Illumina 50 bp single end libraries were prepared using the TruSeq Stranded Total RNA-seq Kit and subjected to Illumina sequencing. Library preparation and RNA sequencing was performed at the Duke University Center for Genomic and Computation Biology. Three biological replicates were performed for each sample.

After sequencing, Illumina raw reads were trimmed with Trimmomatic to remove Illumina adaptors (86) and mapped to the most recent *M. sympodialis* reference genome (87) using HiSat. Generated .bam files were used to run StringTie with the *M. sympodialis* annotation as guide, and the –e option to provide the correct output for the statistical analysis (88). Read count information for statistical analysis were extracted using a provided python script (*prepDE.py*). DESeq2 was used to determine the differentially expressed genes (DEGs) as having FDR < 0.05 and log$_2$FC ± 0.5, which are common parameters used to define relevant genes in RNAseq experiments (89). StringTie and DEseq2 were run on Galaxy (90). Functional annotation of the DEGs was performed using the Blast2GO pipeline, which includes the BLASTx against the NCBI non-redundant protein database, gene ontology (GO) annotation and InterProScan (91). Venn diagram to identify DEGs in common between the two comparisons were generated using the following web server (http://bioinformatics.psb.ugent.be/webtools/Venn/). Gene network interactions were determined using STRING (https://string-db.org/cgi/input.pl) with the *S. cerevisiae* protein set as reference.

Flavohemoglobin purification and crystal structure. A detailed procedure for the flavohemoglobin purification and crystal structure is reported in SI Appendix, SI Material and Methods. Briefly, a construct expressing His-Tev-*YHB101* was cloned into *E. coli* BL21(DE3) cells for expression studies. Protein expression was induced with 1 mM Isopropyl β-D-1-thiogalactopyranoside (IPTG), and 5-aminoluevulinic acid 0.3 mM was also added to facilitate heme biosynthesis. The cellular pellet was collected and lysed via microfluidization with two

passes at 15,000 PSI on ice and clarified via centrifugation at 200 rcf for 45 minutes at 4°C, and
filtered with a 0.2 µm filter. The supernatant was applied to a Ni^{2+} charged HiTrap Chelating HP
(GE Healthcare) columns and the protein eluted with a 500 mM imidazole gradient. The fractions
of interest (which were visibly red-brown) were pooled, the His-TEV tag removed while dialyzing
overnight at 4°C, and fractions of the cleaved protein eluted from the column were concentrated
for size exclusion chromatography via centrifugal concentration.

Apo *M. yamatoensis* flavohemoglobin protein was set up for crystallization. Crystals were
obtained with Morpheus B12: 12.5% (w/v) PEG1000, 12.5% (w/v) PEG3350, 12.5% (v/v) MPD,
0.03 M each sodium fluoride, sodium bromide, and sodium iodide, and 0.1 M bicine/Trizma base
pH 8.5. After 35 days, red-brown crystals were harvested and analyzed on a Rigaku FR-E+ 007
SuperBright rotating anode equipped with Rigaku Varimax optics and a Saturn 944+ detector using
several packages described in detail in SI Appendix, SI Material and Methods. The
flavohemoglobin structures of *E. coli* (PDB ID 1GVH) and *Saccharomyces cerevisiae* (PDB ID
4G1V) were used for comparison with that of *M. yamatoensis*.

Interaction of *M. sympodialis* strains with the host. The ability of *M. sympodialis* WT, *yhb1Δ*
mutant, and complemented strains *yhb1Δ + YHB1-GFP* and *yhb1Δ + YHB101-GFP* to survive
within macrophage was carried out according to previous protocol (30, 92). J774 A.1 cells
(1×10^5/well) were added to 96-well plates and activated by addition of either 1) 10 nM phorbol
myristate acetate (PMA) and incubated for 1 h at 37 °C with 5% CO_2, 2) 100 U/mL interferon-γ
and LPS (0.6 µg/mL) and incubated overnight at 37˚C 5% CO_2. Two days old cultures of *M.*
sympodialis were washed twice with sterile water, and resuspended in RPMI + human serum 20
% for opsonization for 2 h at 37˚C. *M. sympodialis* cells were washed three times with water,
resuspended in DMEM and added to the macrophages at a 1:1 yeast: macrophage ratio. Plates were
incubated for 2 h at 37 °C in 5% CO2, then the co-cultures were washed three times with PBS to
remove yeasts that were not internalized, and the plate was incubated at 37 °C in 5% CO2 for 18 h.
Yeast cells were collected by lysing macrophages, and from each condition a 1:20 dilution was
plated on mDixon agar and incubated at 30 °C for 3-5 days to determine yeast survival.

For *in vivo* infection experiments, WT C57BL/6j mice were purchased from Janvier
Elevage. *Nos2*-/- mice (93) were obtained from Nicolas Fasel (Lausanne). Mice were maintained
at the Laboratory Animal Science Center of University of Zurich, Zurich, Switzerland and used at

6-12 weeks in sex- and age-matched groups. Epicutaneous infection of the mouse ear skin was

performed as described previously (5, 94). Briefly, *Malassezia* strains were grown for 3-4 days at

30°C, 180 rpm in liquid mDixon medium. Cells were washed in PBS and suspended in native olive

oil at a density of 20 OD_{A600}/mL. 100 µl suspension (corresponding to 2 OD_{A600}) of yeast cells

was applied topically onto the dorsal ear skin that was previously barrier-disrupted by mild tape

stripping while mice were anaesthetized. For determining the fungal loads in the skin, tissue was

transferred in water supplemented with 0.05% Nonidet P40 (AxonLab), homogenized and plated

on mDixon agar and incubated at 30°C for 3-4 days. For quantification of cellular infiltrates, single

cell suspensions of ear skin were stained with antibodies directed against CD45 (clone 104),

CD11b (clone M1/70), Ly6G (clone 1A8), Ly6C (clone HK1.4) in PBS supplemented with 1%

FCS, 5 mM EDTA and 0.02% NaN_3. LIVE/DEAD Near IR stain (Life Technologies) was used

for exclusion of dead cells. Cells were acquired on a FACS Cytoflex (Beckman Coulter) and the

data were analyzed with FlowJo software (FlowJo LLC). The gating of the flow cytometric data

was performed according to the guidelines for the use of flow cytometry and cell sorting in

immunological studies (95), including pre-gating on viable and single cells for analysis. For

transcript expression analysis, total RNA was isolated from ear skin according to standard

protocols using TRI Reagent (Sigma Aldrich). cDNA was generated by RevertAid reverse

transcriptase (Thermo Fisher). Quantitative PCR was performed using SYBR Green (Roche) and

a QuantStudio 7 Flex (Life Technologies) instrument for *Il17a* (96), *Defb 3* (97), and *Nos2*. All

RT-qPCR assays were performed in duplicates and the relative expression (rel. expr.) of each gene

was determined after normalization to *Actb* transcript levels. The primers used for qPCR are listed

in Table S2.

Data Availability. The sequence data generated in this study were submitted to National Center

for Biotechnology Information under BioProject accession number PRJNA626605. Individual

accession numbers are SRR11574550 for RNA-seq reads of *Malassezia* WT untreated control

samples, SRR11574549 for RNA-seq reads of *Malassezia* WT NO-treated samples and

SRR11574548 for RNA-seq *Malassezia yhb1*Δ mutant. The final structure factors and coordinates

of the flavohemolgobin Yhb101 of *M. yamatoensis* were deposited in the PDB with code 6O0A.

ACKNOWLEDGMENTS

We thank Stephen Rogers for assistance with the NO consumption assay, Nicolas Fasel for *Nos2-*

/- mice, Ellen Wallace for her contribution to cloning the *Malassezia* crystallography constructs,

Jan Abendroth for his contributions to solving the structure of *Malassezia* by phasing off the bound

iron, Jason Yano and Rana Sidhu for consultation on improving heme incorporation during

recombinant expression of the crystallography constructs, and Tom Edwards and Don Lorimer for

their overall support of the project.

FUNDING INFORMATION

This work was in part supported by NIH/NIAID R01 grant AI50113-15, and by NIH/NIAID R37

MERIT award AI39115-22 (to J.H.) and by SNF grant 310030_189255 (to S.L.L.). VA Merit BX-

003478 supported TJM's work. Crystallization work was funded by the NIH/NIAID (contract nos.

HHSN272200700057, HHSN272201200025C, and HHSN272201700059C to Peter J. Myler).

Joseph Heitman is Co-Director and Fellow of the CIFAR program "Fungal Kingdom: Threats &

Opportunities".

     Figures legend

Figure 1. Evidence for independent HGT events of the flavohemoglobin-encoding genes in

***Malassezia* from the Actinobacteria.** (A) Maximum likelihood phylogeny of consisting of 2,155

flavohemoglobin protein sequences. Two groups (clade 1 and 2) of horizontally transferred

flavohemoglobin genes (*YHB1* and *YHB101*) in *Malassezia* are colored in orange. Other tree

branches are colored according to the key on the top left, representing other major groups of

organisms. The phylogeny was visualized using iTOL v3.6.1 (98) and was rooted at the midpoint.

(B and C) Zoomed views of the ML phylogeny showing in more detail the position of *Malassezia*

flavohemoglobins from clades 1 and 2, and their putative bacterial donor lineages. (D) Results of

topological constraints tests that significantly rejected the monophyletic origin for both *Malassezia*

flavohemoglobins clades, providing additional support for independent HGT events.

Figure 2. Evolutionary trajectory of flavohemoglobin-encoding genes in *Malassezia* after

their acquisition via HGT from different donor bacteria lineages. (A) Phylogenetic

relationship of *Malassezia* species with available genome sequence inferred from the

concatenation of 246 single-copy proteins. Color codes assigned to the different phylogenetic

clades (named A to D) are kept consistent in all figures. The tree was rooted at the midpoint and

white circles in the tree nodes indicate full UFboot and SH-aLRT branch support. The proposed

evolutionary events that led to the final arrangement of the flavohemoglobin-encoding genes

shown in panel B are shown in the phylogenetic tree, as given in the key; double arrows indicate

relocation of the *YHB101* gene in subtelomeric position. (B) Chromosomal regions encompassing

the *YHB1* gene in *Malassezia*. Genes are shown as arrows denoting the direction of transcription

and orthologs are represented in the same color. Non-syntenic genes are shown in white, and small

arrows in black represent tRNAs. The *YHB1* gene is shown as red arrows outlined in bold in the

center. The end of a scaffold is represented by a forward slash. For *M. yamatoensis* and *M.*

slooffiae, yellow bars indicate the absence of the *YHB1* gene in otherwise syntenic regions, and

those in green indicate instances where another flavohemoglobin-encoding gene, named *YHB101*

and represented as orange arrows outlined in bold, was acquired by an independent HGT event. A

defective *YHB1* gene in *M. nana* CBS9557 is denoted by the Greek symbol ψ. Gene codes in red

or blue are as they appear in *M. globosa* (prefix "MGL_") or *M. sympodialis* (prefix "MSYG_")

genome annotations, respectively, those in black were named based on top BLASTp hits in *S.*

cerevisiae, and "hyp" represent hypothetical proteins. Black circle represents the end of a

chromosome. Scaffold/chromosomal locations and accession numbers are given for each region

in Table S1.

Figure 3. *M. sympodialis* flavohemoglobins are involved in nitrosative stress resistance and

NO degradation. (A) Stress sensitivity assay of *M. sympodialis* WT, *yhb1*Δ mutant, and

complementing strains *yhb1*Δ + *YHB1* and *yhb1*Δ + *YHB101* on mDixon agar supplemented with

the NO-donor agent DETA NONOate and $NaNO_2$, and with hydrogen peroxide. (B) GFP

expression in the *M. sympodialis yhb1*Δ mutant, and complementing strains *yhb1*Δ + *YHB1* and

*yhb1*Δ + *YHB101*, and respective GFP signal analyzed trough FACS. (C) NO consumption assay

by *M. sympodialis* WT, *yhb1*Δ mutant, and complementing strains *yhb1*Δ + *YHB1* and *yhb1*Δ +

YHB101; the blue trace indicates the NO level over a period of 15 min. NO and *Malassezia* (Y =

yeast) injections are indicated by purple arrows. In this experiment, 10 µL (Y), 20 µL (x 2) and 30

µL (x 3), 40 µL (x 4) and 50 µL (x 5) of *Malassezia* cellular suspensions were injected. (D)

Representative fluorescent staining of intracellular NO with DAF-FM DA in *M. sympodialis* WT

and two independent *yhb1*Δ mutants, and (E-F) quantification of the NO signal by flow cytometry;

spontaneous fluorescence of *M. sympodialis* was used as background to detect specifically DAF-

FM DA signal. Asterisks indicates statistically significant differences (* = $p<0.05$, ** = $p<0.01$)

according to the unpaired student's t-test with Welch's correction.

Figure 4. Transcriptomic profile of *M. sympodialis* strains under NO-accumulation

conditions. (A) MA-plot displaying the transcriptomic changes of the *M. sympodialis yhb1*Δ

mutant compared to the WT *M. sympodialis* strain. Red dots indicate differentially expressed genes

for FDR < 0.05. The most upregulated and downregulated genes (*MSYG_1280* and *MSYG_0901*,

respectively) are indicated, along with the *YHB1* gene, which represents an internal control as its

downregulation is expected because the gene is deleted. (B) MA-plot displaying the transcriptomic

changes of *M. sympodialis* WT grown in the presence of $NaNO_2$ compared to the untreated control.

Red dots indicate differentially expressed genes for FDR < 0.05; the most upregulated and

downregulated genes are indicated. (C) Gene ontology classification relative to the RNAseq

condition reported in B. Upregulated genes are indicated in red, and downregulated genes are

indicated in green. (D) Venn diagrams comparison of the upregulated and downregulated genes

relative to RNAseq conditions reported in A and B; the panel on the right shows a heatmap of the
log2FC of the shared upregulated (red) and downregulated (green) genes. Predicted allergens are
indicated with one asterisk, and two asterisks indicate a predicted secreted lipase.

Figure 5. 3-dimensional X-ray crystal structure of the *M. yamatoensis* flavohemoglobin
Yhb101. (A) The globin domain (cyan) binds a heme molecule. The reductase domain consists of
a FAD-binding domain (gray) and a NAD-binding domain (tan) that bind a FAD molecule. (B)
An overlay of flavohemoglobin globin domains from fungus, bacteria, and yeast: globin domains
of *M. yamatoensis* (PDB ID 6O0A; blue), *S. cerevisiae* (PDB ID 4G1V; yellow), and *E. coli* (PDB
ID 1GVH; green) show structural similarity.

Figure 6. *Malassezia* genes acquired through HGT from bacteria. HGT candidates identified
in the genomes of the 15 *Malassezia* species (represented on the top according to their phylogenetic
classification) are shown as different lines in the presence-absence matrix, with the closest
ortholog in *S. cerevisiae* indicated in parenthesis, where available. For each HGT candidate, the
presence of the gene in a genome is indicated by orange square, and the intensity of the color is
correlated with the gene copy number (numbers in white). HGT candidates occurring in multiple
Malassezia species are shown in the top half of the matrix, whereas those that are species-specific
HGT candidates are shown in the bottom half of the matrix, and color-coded as shown in the key.
Asterisks indicate HGT candidate genes identified in the previous study (3). The bacterially-
derived gene encoding an aliphatic amidase identified in *M. nana* CBS9557 seems to be another
instance of a pseudogene in this strain (indicated as ψ).

References

1. K. Findley *et al.*, Topographic diversity of fungal and bacterial communities in human skin.
Nature **498**, 367-370 (2013).
2. B. Theelen *et al.*, *Malassezia* ecology, pathophysiology, and treatment. *Medical Mycology*
56, S10-S25 (2018).
3. G. Wu *et al.*, Genus-wide comparative genomics of *Malassezia* delineates its phylogeny,
physiology, and niche adaptation on human skin. *PLOS Genetics* **11**, e1005614 (2015).
4. G. Gaitanis, P. Magiatis, M. Hantschke, I. D. Bassukas, A. Velegraki, The *Malassezia*
genus in skin and systemic diseases. *Clin Microbiol Rev* **25**, 106-141 (2012).
5. F. Sparber *et al.*, The skin commensal yeast *Malassezia* triggers a type 17 response that
coordinates anti-fungal immunity and exacerbates skin inflammation. *Cell Host & Microbe*
25, 389-403.e386 (2019).
6. J. J. Limon *et al.*, *Malassezia* is associated with Crohn's disease and exacerbates colitis in
mouse models. *Cell Host & Microbe* **25**, 377-388.e376 (2019).
7. B. Aykut *et al.*, The fungal mycobiome promotes pancreatic oncogenesis via activation of
MBL. *Nature* **574**, 264-267 (2019).
8. P. Soret *et al.*, Respiratory mycobiome and suggestion of inter-kingdom network during
acute pulmonary exacerbation in cystic fibrosis. *Scientific Reports* **10**, 3589 (2020).
9. C. T. Stomberski, D. T. Hess, J. S. Stamler, Protein S-nitrosylation: Determinants of
specificity and enzymatic regulation of S-nitrosothiol-based signaling. *Antioxid Redox*
Signal **30**, 1331-1351 (2019).
10. M. T. Forrester, M. W. Foster, Protection from nitrosative stress: a central role for
microbial flavohemoglobin. *Free radical biology & medicine* **52**, 1620-1633 (2012).
11. R. I. Astuti, R. Nasuno, H. Takagi, Nitric oxide signaling in yeast. *Appl Microbiol*
Biotechnol **100**, 9483-9497 (2016).
12. D. Cánovas, J. F. Marcos, A. T. Marcos, J. Strauss, Nitric oxide in fungi: is there NO light
at the end of the tunnel? *Current genetics* **62**, 513-518 (2016).
13. Y. Zhao, J. Lim, J. Xu, J. H. Yu, W. Zheng, Nitric oxide as a developmental and metabolic
signal in filamentous fungi. *Mol Microbiol* 10.1111/mmi.14465 (2020).
14. B. Koch *et al.*, A metabolic checkpoint for the yeast-to-hyphae developmental switch
regulated by endogenous nitric oxide signaling. *Cell reports* **25**, 2244-2258.e2247 (2018).
15. A. T. Marcos, M. S. Ramos, T. Schinko, J. Strauss, D. Canovas, Nitric oxide homeostasis
is required for light-dependent regulation of conidiation in *Aspergillus*. *Fungal Genet Biol*
137, 103337 (2020).
16. P. R. Gardner, Nitric oxide dioxygenase function and mechanism of flavohemoglobin,
hemoglobin, myoglobin and their associated reductases. *Journal of inorganic biochemistry*
99, 247-266 (2005).
17. S. Zhou *et al.*, Functional analysis and subcellular location of two flavohemoglobins from
Aspergillus oryzae. *Fungal Genet Biol* **48**, 200-207 (2011).
18. K. Lapp *et al.*, Characterization of the *Aspergillus fumigatus* detoxification systems for
reactive nitrogen intermediates and their impact on virulence. *Frontiers in microbiology* **5**,
469-469 (2014).
19. S. Gupta, S. Pawaria, C. Lu, S. R. Yeh, K. L. Dikshit, Novel flavohemoglobins of
mycobacteria. *IUBMB life* **63**, 337-345 (2011).

20. N. Thakur, A. Kumar, K. L. Dikshit, Type II flavohemoglobin of *Mycobacterium*
smegmatis oxidizes d-lactate and mediate electron transfer. *International Journal of*
Biological Macromolecules **112**, 868-875 (2018).
21. D. Hoogewijs, S. Dewilde, A. Vierstraete, L. Moens, S. N. Vinogradov, A phylogenetic
analysis of the globins in fungi. *PLoS One* **7**, e31856 (2012).
22. J. H. Wisecaver, W. G. Alexander, S. B. King, C. T. Hittinger, A. Rokas, Dynamic
evolution of nitric oxide detoxifying flavohemoglobins, a family of single-protein
metabolic modules in bacteria and eukaryotes. *Mol Biol Evol* **33**, 1979-1987 (2016).
23. S. N. Vinogradov *et al.*, A phylogenomic profile of globins. *BMC Evolutionary Biology* **6**,
31 (2006).
24. L. Liu, M. Zeng, A. Hausladen, J. Heitman, J. S. Stamler, Protection from nitrosative stress
by yeast flavohemoglobin. *Proc Natl Acad Sci U S A* **97**, 4672-4676 (2000).
25. F. Husnik, J. P. McCutcheon, Functional horizontal gene transfer from bacteria to
eukaryotes. *Nature reviews. Microbiology* **16**, 67-79 (2018).
26. J. Ropars *et al.*, Adaptive horizontal gene transfers between multiple cheese-associated
fungi. *Curr Biol* **25**, 2562-2569 (2015).
27. G. Ianiri, S. Applen Clancey, S. C. Lee, J. Heitman, FKBP12-dependent inhibition of
calcineurin mediates immunosuppressive antifungal drug action in *Malassezia*. *mBio* **8**,
e01752-01717 (2017).
28. G. Ianiri, G. Dagotto, S. Sun, J. Heitman, Advancing functional genetics through
Agrobacterium-mediated insertional mutagenesis and CRISPR/Cas9 in the commensal and
pathogenic yeast *Malassezia*. *Genetics* **212**, 1163-1179 (2019).
29. X. Wang *et al.*, Biological activity of nitric oxide in the plasmatic compartment. *Proc Natl*
Acad Sci U S A **101**, 11477-11482 (2004).
30. M. de Jesus-Berrios *et al.*, Enzymes that counteract nitrosative stress promote fungal
virulence. *Curr Biol* **13**, 1963-1968 (2003).
31. A. Idnurm, J. L. Reedy, J. C. Nussbaum, J. Heitman, *Cryptococcus neoformans* virulence
gene discovery through insertional mutagenesis. *Eukaryot. Cell* **3**, 420-429 (2004).
32. S. Zhou *et al.*, *Aspergillus oryzae* flavohemoglobins promote oxidative damage by
hydrogen peroxide. *Biochem Biophys Res Commun* **394**, 558-561 (2010).
33. B. S. Hromatka, S. M. Noble, A. D. Johnson, Transcriptional response of *Candida albicans*
to nitric oxide and the role of the *YHB1* gene in nitrosative stress and virulence. *Mol Biol*
Cell **16**, 4814-4826 (2005).
34. M. M. Nakano, Essential role of flavohemoglobin in long-term anaerobic survival of
Bacillus subtilis. *Journal of bacteriology* **188**, 6415-6418 (2006).
35. Z. Zhang, M. Gerstein, Large-scale analysis of pseudogenes in the human genome. *Current*
opinion in genetics & development **14**, 328-335 (2004).
36. A. Hirai *et al.*, *Malassezia nana* sp. nov., a novel lipid-dependent yeast species isolated
from animals. *Int J Syst Evol Microbiol* **54**, 623-627 (2004).
37. S. Horan, I. Bourges, B. Meunier, Transcriptional response to nitrosative stress in
Saccharomyces cerevisiae. *Yeast* **23**, 519-535 (2006).
38. D.-H. Yang *et al.*, Rewiring of signaling networks modulating thermotolerance in the
human pathogen *Cryptococcus neoformans*. *Genetics* **205**, 201-219 (2017).
39. A. O. Veri *et al.*, Tuning Hsf1 levels drives distinct fungal morphogenetic programs with
depletion impairing Hsp90 function and overexpression expanding the target space. *PLOS*
Genetics **14**, e1007270 (2018).

40.   P. Biswas, S. Ghosh, Global transcriptomic profiling of *Schizosaccharomyces pombe* in
response to nitrosative stress. *Gene* **558**, 241-253 (2015).
41.   C. Herrero-de-Dios *et al.*, Redox regulation, rather than stress-induced phosphorylation, of
a Hog1 mitogen-activated protein kinase modulates its nitrosative-stress-specific outputs.
mBio **9**, e02229-17 (2018).
42.   E. El Hammi *et al.*, Active site analysis of yeast flavohemoglobin based on its structure
with a small ligand or econazole. *FEBS J* **279**, 4565-4575 (2012).
43.   A. Ilari, A. Bonamore, A. Farina, K. A. Johnson, A. Boffi, The X-ray structure of ferric
Escherichia coli flavohemoglobin reveals an unexpected geometry of the distal heme
pocket. *J Biol Chem* **277**, 23725-23732 (2002).
44.   S. E. Winter *et al.*, Host-derived nitrate boosts growth of *E. coli* in the inflamed gut. *Science*
339, 708-711 (2013).
45.   R. W. Nowell *et al.*, Comparative genomics of bdelloid rotifers: Insights from desiccating
and nondesiccating species. *PLOS Biology* **16**, e2004830 (2018).
46.   C. Boschetti *et al.*, Biochemical diversification through foreign gene expression in bdelloid
rotifers. *PLOS Genetics* **8**, e1003035 (2012).
47.   W. G. Alexander, J. H. Wisecaver, A. Rokas, C. T. Hittinger, Horizontally acquired genes
in early-diverging pathogenic fungi enable the use of host nucleosides and nucleotides.
Proceedings of the National Academy of Sciences **113**, 4116-4121 (2016).
48.   G. Koutsovoulos *et al.*, No evidence for extensive horizontal gene transfer in the genome
of the tardigrade *Hypsibius dujardini*. *Proceedings of the National Academy of Sciences*
113, 5053-5058 (2016).
49.   A. Beceiro, M. Tomás, G. Bou, Antimicrobial resistance and virulence: a successful or
deleterious association in the bacterial world? *Clinical Microbiology Reviews* **26**, 185-230
(2013).
50.   N. Mosqueda *et al.*, Characterization of plasmids carrying the blaOXA-24/40
carbapenemase gene and the genes encoding the AbkA/AbkB proteins of a toxin/antitoxin
system. *The Journal of antimicrobial chemotherapy* **69**, 2629-2633 (2014).
51.   J. M. Tkach *et al.*, Dissecting DNA damage response pathways by analysing protein
localization and abundance changes during DNA replication stress. *Nat Cell Biol* **14**, 966-
976 (2012).
52.   M. Rep *et al.*, The *Saccharomyces cerevisiae* Sko1p transcription factor mediates HOG
pathway-dependent osmotic regulation of a set of genes encoding enzymes implicated in
protection from oxidative damage. *Mol Microbiol* **40**, 1067-1083 (2001).
53.   E. V. Koonin, Orthologs, paralogs, and evolutionary genomics. *Annual Review of Genetics*
39, 309-338 (2005).
54.   B. Lacroix, V. Citovsky, Transfer of DNA from bacteria to eukaryotes. *mBio* **7**, e00863-
00816 (2016).
55.   C. L. Murphy *et al.*, Horizontal gene transfer as an indispensable driver for
Neocallimastigomycota evolution into a distinct gut-dwelling fungal lineage. *Applied and
environmental microbiology* 10.1128/aem.00988-19, AEM.00988-00919 (2019).
56.   M. M. Cals-Grierson, A. D. Ormerod, Nitric oxide function in the skin. *Nitric oxide :
biology and chemistry* **10**, 179-193 (2004).
57.   M. P. Nittler, D. Hocking-Murray, C. K. Foo, A. Sil, Identification of *Histoplasma
capsulatum* transcripts induced in response to reactive nitrogen species. *Molecular biology
of the cell* **16**, 4792-4813 (2005).

58.   I. Heckler, E. M. Boon, Insights into nitric oxide modulated quorum sensing pathways. *Frontiers in microbiology* **10**, 2174-2174 (2019).

59.   D. E. Williams, E. M. Boon, Towards understanding the molecular basis of nitric oxide-regulated group behaviors in pathogenic bacteria. *Journal of Innate Immunity* **11**, 205-215 (2019).

60.   L. Yang *et al.*, Nitric oxide increases biofilm formation in *Saccharomyces cerevisiae* by activating the transcriptional factor Mac1p and thereby regulating the transmembrane protein Ctr1. *Biotechnology for Biofuels* **12**, 30 (2019).

61.   P. J. Keeling, J. D. Palmer, Horizontal gene transfer in eukaryotic evolution. *Nat Rev Genet* **9**, 605-618 (2008).

62.   J. O. Andersson, Gene transfer and diversification of microbial eukaryotes. *Annu Rev Microbiol* **63**, 177-193 (2009).

63.   J. Yue, X. Hu, H. Sun, Y. Yang, J. Huang, Widespread impact of horizontal gene transfer on plant colonization of land. *Nature Communications* **3**, 1152 (2012).

64.   S. Garcia-Vallve, A. Romeu, J. Palau, Horizontal gene transfer of glycosyl hydrolases of the rumen fungi. *Mol Biol Evol* **17**, 352-361 (2000).

65.   G. Ricard *et al.*, Horizontal gene transfer from Bacteria to rumen Ciliates indicates adaptation to their anaerobic, carbohydrates-rich environment. *BMC Genomics* **7**, 22 (2006).

66.   A. Amend, From dandruff to deep-sea vents: *Malassezia*-like fungi are ecologically hyper-diverse. *PLoS pathogens* **10**, e1004277 (2014).

67.   G. Ianiri, A. F. Averette, J. M. Kingsbury, J. Heitman, A. Idnurm, Gene function analysis in the ubiquitous human commensal and pathogen *Malassezia* genus. *mBio* **7**, e01853-01816 (2016).

68.   M. Stanke, B. Morgenstern, AUGUSTUS: a web server for gene prediction in eukaryotes that allows user-defined constraints. *Nucleic Acids Res* **33**, W465-467 (2005).

69.   P. Jones *et al.*, InterProScan 5: genome-scale protein function classification. *Bioinformatics (Oxford, England)* **30**, 1236-1240 (2014).

70.   L. Fu, B. Niu, Z. Zhu, S. Wu, W. Li, CD-HIT: accelerated for clustering the next-generation sequencing data. *Bioinformatics (Oxford, England)* **28**, 3150-3152 (2012).

71.   K. Katoh, D. M. Standley, MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol Biol Evol* **30**, 772-780 (2013).

72.   S. Capella-Gutierrez, J. M. Silla-Martinez, T. Gabaldon, trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**, 1972-1973 (2009).

73.   L. T. Nguyen, H. A. Schmidt, A. von Haeseler, B. Q. Minh, IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol Evol* **32**, 268-274 (2015).

74.   S. Kalyaanamoorthy, B. Q. Minh, T. K. F. Wong, A. von Haeseler, L. S. Jermiin, ModelFinder: fast model selection for accurate phylogenetic estimates. *Nature methods* **14**, 587-589 (2017).

75.   B. Q. Minh, M. A. Nguyen, A. von Haeseler, Ultrafast approximation for phylogenetic bootstrap. *Mol Biol Evol* **30**, 1188-1195 (2013).

76.   S. Guindon *et al.*, New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Systematic Biology* **59**, 307-321 (2010).

77. H. Shimodaira, M. Hasegawa, CONSEL: for assessing the confidence of phylogenetic tree selection. *Bioinformatics* **17**, 1246-1247 (2001).

78. H. Shimodaira, An approximately unbiased test of phylogenetic tree selection. *Syst Biol* **51**, 492-508 (2002).

79. B. Contreras-Moreira, P. Vinuesa, GET_HOMOLOGUES, a versatile software package for scalable and robust microbial pangenome analysis. *Appl Environ Microbiol* **79**, 7696-7701 (2013).

80. ODiogoSilva, ElConcatenero (Version v1.0). *Zenodo.* http://doi.org/10.5281/zenodo.10588 (2014).

81. B. Buchfink, C. Xie, D. H. Huson, Fast and sensitive protein alignment using DIAMOND. *Nature methods* **12**, 59-60 (2015).

82. G. Ianiri, K. J. Boyce, A. Idnurm, Isolation of conditional mutations in genes essential for viability of *Cryptococcus neoformans*. *Curr Genet* **63**, 519-530 (2017).

83. C. S. Hoffman, "Preparation of yeast DNA" in Current Protocols in Molecular Biology. (John Wiley & Sons, Inc., 2001), 10.1002/0471142727.mb1311s39.

84. A. G. Pinder, S. C. Rogers, A. Khalatbari, T. E. Ingram, P. E. James, The measurement of nitric oxide and its metabolites in biological samples by ozone-based chemiluminescence. *Methods in molecular biology (Clifton, N.J.)* **476**, 11-28 (2008).

85. S. C. Rogers *et al.*, Effect of plasma processing and storage on microparticle abundance, nitric oxide scavenging, and vasoactivity. *Transfusion* **59**, 1568-1577 (2019).

86. A. M. Bolger, M. Lohse, B. Usadel, Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**, 2114-2120 (2014).

87. Y. Zhu *et al.*, Proteogenomics produces comprehensive and highly accurate protein-coding gene annotation in a complete genome assembly of *Malassezia sympodialis*. *Nucleic Acids Research* **45**, 2629-2643 (2017).

88. M. Pertea, D. Kim, G. M. Pertea, J. T. Leek, S. L. Salzberg, Transcript-level expression analysis of RNA-seq experiments with HISAT, StringTie, and Ballgown. *Nat Protoc* **11**, 1650-1667 (2016).

89. N. J. Schurch *et al.*, How many biological replicates are needed in an RNA-seq experiment and which differential expression tool should you use? *RNA (New York, N.Y.)* **22**, 839-851 (2016).

90. E. Afgan *et al.*, The Galaxy platform for accessible, reproducible, and collaborative biomedical analyses: 2018 update. *Nucleic Acids Research* **46**, W537-W544 (2018).

91. A. Conesa *et al.*, Blast2GO: a universal tool for annotation, visualization, and analysis in functional genomics research. *Bioinformatics* **21**, 3674-3676 (2005).

92. T. R. O'Meara, S. M. Holmer, K. Selvig, F. Dietrich, J. A. Alspaugh, *Cryptococcus neoformans* Rim101 is associated with cell wall remodeling and evasion of the host immune responses. *mBio* **4**, e00522-12 (2013).

93. V. E. Laubach, E. G. Shesely, O. Smithies, P. A. Sherman, Mice lacking inducible nitric oxide synthase are not resistant to lipopolysaccharide-induced death. *Proc Natl Acad Sci U S A* **92**, 10688-10692 (1995).

94. F. Sparber, S. LeibundGut-Landmann, Infecting mice with *Malassezia* spp. to study the fungus-host interaction. http://dx.doi.org/10.3791/60175.

95. A. Cossarizza, e. al, Guidelines for the use of flow cytometry and cell sorting in immunological studies (second edition). *European Journal of Immunology* **49**, 1457-1973 (2019).

96. L. Overbergh *et al.*, The use of real-time reverse transcriptase PCR for the quantification
of cytokine gene expression. *Journal of Biomolecular Techniques* **14**, 33-43 (2003).
97. K. Trautwein-Weidner, A. Gladiator, S. Nur, P. Diethelm, S. LeibundGut-Landmann, IL-
17-mediated antifungal defense in the oral mucosa is independent of neutrophils. *Mucosal*
immunology **8**, 221-231 (2015).
98. I. Letunic, P. Bork, Interactive tree of life (iTOL) v3: an online tool for the display and
annotation of phylogenetic and other trees. *Nucleic Acids Res* **44**, W242-245 (2016).

1215   **Figure 1**

D

		AU test of monophyly	
Tree	−lnL	*Diff −lnL	*P* value
Unconstrained	-607815.172		
Constrained	-608369.863	554.6	0.001

*Diff −lnL, difference in −log likelihood between the best topology and
the constrained topology that forced *Malassezia* to be monophyletic.

1234   **Figure 2**

Figure 3

1263   **Figure 4**

1279 **Figure 5**

1297 **Figure 6**

1298