Comparative genomics reveals different population structures associated with

host and geographic origin in antimicrobial-resistant *Salmonella enterica*

Jingqiu Liao[1, 2], Renato Hohl Orsi[1], Laura Carroll[1†], Martin Wiedmann[1*]

[1]Department of Food Science, Cornell University, Ithaca, New York, 14853, USA

[2]Graduate Field of Microbiology, Cornell University, Ithaca, New York, 14853, USA

[†]Present address: Structural and Computational Biology Unit, EMBL, Heidelberg, Germany

*For correspondence: Martin Wiedmann (martin.wiedmann@cornell.edu)

Running title: Population structure of AMR *S. enterica*

Originality/Significance Statement

Salmonella enterica, which is the causative agent of salmonellosis, poses a growing public health

concern due to the emergence and spread of antimicrobial resistant (AMR) strains. The mechanisms

underlying the population structure associated with different hosts and geographic origins of AMR *S.*

enterica are underexplored due to limited genome-wide studies assessing the impact of ecological

niches on genetic variations. By employing comparative genomics, our study provided insights into

the genomic profiles of AMR *S. enterica* associated with two distinct hosts and two distant

geographic locations, improving the mechanistic understanding of how bacterial population structure

is shaped by different ecological niches. Our findings have broad implications for elucidating the

impact of ecological and evolutionary forces on the adaptation, antimicrobial resistance, and

pathogenicity of bacteria. Also, specific genetic markers we identified may help predict host or

geographic origin of AMR *Salmonella* isolates, which could benefit the source tracking (e.g., host

and geographic origins) of human disease cases and contamination events caused by AMR *S.*

enterica.

Summary

Genetic variation in a pathogen, including the causative agent of salmonellosis, *Salmonella enterica*, can occur as a result of eco-evolutionary forces triggered by dissimilarities of ecological niches. Here, we applied comparative genomics to study 90 antimicrobial resistant (AMR) *S. enterica* isolates from bovine and human hosts in New York state and Washington state to understand host- and geographic-associated population structure. Results revealed distinct presence/absence profiles of functional genes and pseudogenes (e.g., virulence genes) associated with bovine and human isolates. Notably, bovine isolates contained significantly more transposase genes but fewer transposase pseudogenes than human isolates, suggesting the occurrence of large-scale transposition in genomes of bovine and human isolates at different times. The high correlation between transposase genes and AMR genes, as well as plasmid replicons, highlights the potential role of horizontally transferred transposons in promoting adaptation to antibiotics. By contrast, a number of potentially geographic-associated single-nucleotide polymorphisms (SNPs), rather than geographic-associated genes, were identified. Interestingly, 38% of these SNPs were in genes annotated as cell surface protein-encoding genes, including some essential for antibiotic resistance and host colonization. Overall, different evolutionary forces and limited recent inter-population transmission appear to shape AMR *S. enterica* population structure in different hosts and geographic origins.

Introduction

Dissimilarity of ecological niches formed by different hosts or geographic origins is an important

factor that can lead to genetic variation (e.g., gene presence/absence and polymorphism level

divergence) and shape population structure, thus contributing to the evolution of pathogens (Strachan

et al., 2015; Hoberg and Brooks, 2008; Richards *et al.*, 2011; Strachan *et al.*, 2015). Besides

providing insights in adaptation, pathogenesis and host range of pathogens, population structure

associated with specific attributes (e.g., disease, host or geographic origin) also provides clues for

evolutionary processes (e.g., selection, genetic drift and gene flow) that shaped pathogen diversity (Li

et al., 2019; Yue *et al.*, 2015). In addition, understanding of ecological niche-associated population

structure is advancing the field of epidemiology by interrogating complex genome information to

accurately predict contamination source and transmission patterns of pathogens including *Salmonella*

enterica (Lupolova *et al.*, 2016; Lupolova *et al.*, 2017; Morgan *et al.*, 2004; Moura *et al.*, 2017).

The Gram-negative facultative anaerobe *S. enterica* is one of the major causes of human

gastroenteritis, bacteraemia and enteric fever worldwide (Eng *et al.*, 2015; Ao *et al.*, 2015). The

emergence and increasing prevalence of antimicrobial-resistant (AMR) *S. enterica* have further

raised public health concerns and may increase the mortality rate of infections caused by this

pathogen (Hong *et al.*, 2016; Eng *et al.*, 2015). AMR *S. enterica* has been detected in a wide range of

animals in the majority of states in the US (Eng *et al.*, 2015; Centers for Disease Control and

Prevention, 2017). Among >2,500 recognized serotypes of *S. enterica*, the broad host range serotypes

Typhimurium and Newport are frequently associated with the human *Salmonella* infections, while

the bovine adapted serotype Dublin is frequently associated with cattle infections. The emergence of

AMR strains has been described for all three serotypes (Foley and Lynne, 2008). In our previous

study of AMR *S.* Typhimurium, *S.* Newport and *S.* Dublin isolated from human and bovine sources

in New York state (NY) and Washington state (WA), we found that it is likely that antimicrobial

resistance has emerged independently in multiple *S.* Typhimurium lineages, as compared to single

lineages of *S.* Newport and *S.* Dublin (Liao *et al.*, 2019). Also, our initial study of these isolates

revealed a strong source and geographic association with antimicrobial resistance (Carroll *et al.*,

2017). For example, resistance to sulfamethoxazole-trimethoprim resistance was only observed in

human isolates, while resistance to quinolones and fluoroquinolones was only observed in *S.*

Typhimurium isolated from humans in WA. However, the mechanisms underlying the population

structure associated with different hosts and geographic origins of AMR *S. enterica* are still

underexplored at a genomic scale.

Comparative genomics is now widely used as a tool to study the evolution of bacteria (Chen *et*

al., 2006; Richards *et al.*, 2011; Zheng *et al.*, 2017) and can be used to identify genome-wide genetic

variants that may be associated with host and geographic origin. Identification of such genetic

variants in AMR *S. enterica* allows for a better understanding of genetic basis of pathogenicity and

environmental adaptation. Humans and bovines are identified as two major hosts for AMR *S.*

enterica (Dandekar *et al.*, 2015; Rodriguez-Rivera *et al.*, 2016; Zhao *et al.*, 2003). These two

mammals differ in many aspects, e.g., physiology and diet, and hence form two distinct niches for

associated microorganisms, including the pathogen *Salmonella*. While geographical origins can also

represent distinct habitats and environmental conditions, definition of distinct niches based on

geographical origins is more challenging since complex ecological factors (e.g., climate,

anthropological activities) shape these niches. In order to assess the effects of both host and

geographical origin on AMR *Salmonella* population structure, we selected to study isolates from two

distant locations in the US: NY and WA. These two states showed similar incidence rate of reported

Salmonella cases with 14.87 and 12.12 cases of salmonellosis per 100,000 population in NY and WA

in 2015, respectively (Centers for Disease Control and Prevention, 2017). However, they represent

distinct environments; NY has a humid continental climate, while the WA climate varies greatly

from west to east with a rainy oceanic climate in the western half to an arid climate in the eastern

half, where the majority of cattle operations are located. In addition, the management of dairy

operations is different between NY and WA (United States Department of Agriculture, 2014). For

example, the average dairy cattle herd size of WA is much larger (335 cows) than that of NY (113

cows). Consequently, NY and WA may represent distinct niches, particularly for bovine associated

Salmonella. Hence, genomic comparison of AMR *Salmonella* isolated from bovine and human hosts

in NY and WA will not only provide insights into the understanding of the adaption and evolution of

this pathogen, but also has practical implications as genetic variation associated with different

sources may allow for improved source tracking (e.g., human vs. bovine sources, different

geographic sources) of human disease cases and contamination events. Thus, in this study, we used a

previously reported WGS data set for 90 AMR *S.* Dublin, *S.* Newport, and *S.* Typhimurium isolates

obtained from dairy cattle and humans from NY and WA (Carroll *et al.*, 2017; Liao *et al.*, 2019) to

characterize population structure associated with host and geographic origin using comparative

genomics.

Materials and Methods

Isolates and whole genome sequence data

A previously reported set of 90 *S. enterica* AMR isolates representing three serotypes including

Dublin (n=21), Newport (n=32) and Typhimurium (n=37) collected between 2008 and 2012 (Carroll

et al., 2017; Liao *et al.*, 2019) was used for this study. Among those 90 isolates, 45 were collected

from NY (Dublin (n=8), Newport (n=19) and Typhimurium (n=18)) and 45 were from WA (Dublin

(n=13), Newport (n=13) and Typhimurium (n=19)). Also, among these 90 isolates, 41 were isolated

from fecal samples of bovine (Dublin (n=10), Newport (n=14) and Typhimurium (n=17)) and 49

were isolated from the stool samples of human patients (Dublin (n=11), Newport (n=18) and

Typhimurium (n=20)).

Genome sequence data for all 90 isolates have previously been reported (Carroll *et al.*, 2017)

and deposited in the National Center for Biotechnology Information's (NCBI) Sequence Read

Archive (SRA) under accession number SRP068320. Genome assembly has also been reported

previously (Carroll *et al*., 2017, Liao *et al*., 2019) and assembled genomes are available at NCBI

DDBJ/ENA/GenBank under the accession numbers listed in Additional file 1: Table S4 in Liao *et al*.,

2019. Previously reported data (Liao *et al*., 2019) for (i) genome annotation including identification

of pseudogenes by NCBI Prokaryotic Genome Annotation Pipeline (Tatusova *et al*., 2016), (ii)

identification of orthologous genes (non-pseudogenes) by OrthoMCL (Li *et al*., 2003), (iii)

assignment of gene ontology (GO) and enzyme commission (EC) terms by Blast2GO v1.2.1 (Conesa

et al. 2005), and (iv) high-quality core SNPs for *S.* Dublin, *S.* Newport, and *S.* Typhimurium called

by Cortex variant caller (Iqbal *et al*., 2012) were used here.

Statistics

The Mann-Whitney test was carried out in Prism 7 to determine if the numbers of all pseudogenes as

well as pseudogenes and functional genes annotated as encoding transposase differed significantly

between bovine isolates and human isolates and if the numbers of all pseudogenes differed

significantly between isolates from NY and WA as well. The false-discovery rate procedure (FDR) of

Benjamini and Hochberg (BH) (1995) was employed in R version 3.6.0 to correct for multiple testing

among isolate groups.

Non-metric multidimensional scaling (NMDS) (Kruskal, 1964) was performed using Bray-

Curtis dissimilarities and the metaMDS function in R version 3.6.0's vegan package to compare the

dissimilarity of gene presence/absence pattern among isolates across serotypes and within each

serotype. Permutational multivariate analysis of variance (PERMANOVA) (Anderson and Walsh,

2013) was employed using the adonis function in R version 3.6.0's vegan package to test whether the

centroids of isolate groups as defined by host or geographic origin are equivalent for all groups based

on gene presence/absence among isolates across serotypes and within each serotype. *P*-values and

PERMANOVA test statistics (F) were obtained using Bray-Curtis dissimilarities and 999

permutations. The FDR of BH was employed in R version 3.6.0 to correct for multiple testing among

isolate groups.

Enrichment analyses of pseudogene functions, orthologous genes, and GO terms were

conducted using Fisher's exact test in R version 3.3.1 for (i) bovine versus human isolates and (ii)

isolates from NY versus isolates from WA. FDR of BH was employed using R version 3.6.0 to

correct for multiple testing for Fisher's exact tests for pseudogene functions, orthologous genes, and

GO terms. Items were defined as significantly enriched in one host or geographic origin when FDR

was < 0.05 and odds ratio was > 6.71. These cut-offs were based on a previous report that odds ratio

$= 6.71$ is equivalent to Cohen's $d = 0.8$ (large), indicating strong association (Chen *et al*., 2010).

Correlations between presence/absence of orthologous genes annotated as encoding

transposase in this study and AMR genes as well as plasmid replicons previously reported by Carroll

et al. (2017), using the same dataset, were assessed by Phi coefficient using the Python

sklearn.metrics module. Transposase genes, AMR genes, and plasmid replicons that are present in all

90 genomes and less than 5 genomes were excluded from the Phi coefficient analyses. Items with Phi

coefficient $r > 0.25$ were defined as having a strong positive correlation and included in heatmap

visualization using R version 3.6.0.

Custom Python scripts were used to identify SNPs in (i) core genes present in all three

serotypes and (ii) core genes defined separately for each of the serotypes. Fisher's exact tests were

employed to identify SNPs at which the distribution of nucleotides is dependent on host and/or

geographic origin of isolates; these tests were performed in R version 3.3.1 followed by FDR of BH.

Potentially geographic- or host-associated SNPs were defined as significant when FDR was < 0.05.

Results

Orthologous gene and pseudogene profiles differed significantly between bovine and human

AMR *Salmonella* isolates

Overall, the genomes of the 90 AMR *S. enterica* isolates included 3,637 core genes (i.e., genes

present in all 90 isolates) as well as 3,440 accessory genes. NMDS plots showed that across serotypes

(Figure 1a) and within each serotype (Figure 1b, 1c, 1d), isolates clustered according to their host

based on gene presence/absence, with no evidence for clustering by geographic location.

PERMANOVA further showed that bovine and human cluster centroids, based on gene

presence/absence, were not equivalent across serotypes and within each serotype (FDR < 0.01; Table

1), while no significant difference was observed between isolates from NY and WA in all tests (FDR

> 0.05; Table 1).

The number of pseudogenes among bovine AMR *S. enterica* isolates were significantly lower

than human isolates (Figure 2a, *P* < 0.0001). Across all serotypes, isolates from bovine and human

sources had an average of 107 and 160 pseudogenes, respectively. Similarly, for each serotype

bovine isolates had lower average numbers of pseudogenes; bovine isolates had 154, 89, and 94

pseudogenes, while human isolates had 225, 134 and 148 pseudogenes on average within *S.* Dublin,

S. Newport and *S.* Typhimurium, respectively. By contrast, for all serotypes and within each

serotype, the number of pseudogenes was not significantly different among isolates from NY and

WA (Figure 2b); the AMR *S. enterica* isolates from NY and WA had an average of 132 and 140

pseudogenes, respectively. Within *S.* Dublin, *S.* Newport and *S.* Typhimurium, isolates from NY had

an average number of 207, 109, and 123 pseudogenes, while isolates from WA had 181, 122 and 124

pseudogenes on average, respectively.

Enrichment analysis identified 55, 35, 33, and 34 pseudogene functions significantly enriched

in human isolates across all serotypes, within *S.* Dublin, *S.* Newport, and *S.* Typhimurium,

respectively (FDR < 0.05, odds ratio > 6.71; Table S1). Among the human-associated pseudogene

functions, 8 (e.g., IS3, IS4, IS5/IS1182, IS630 family transposase), 27 (e.g., MarR family

transcriptional regulator, effector protein YopJ), 20 (e.g., type VI secretion lipoprotein/VasD), and 14

(e.g., IS66 family transposase) were exclusively present in all human isolates of all serotypes, *S.*

Dublin, *S.* Newport, and *S.* Typhimurium, respectively (Table S1). By comparison, enrichment

analysis identified fewer pseudogene functions significantly overrepresented among bovine isolates

across all serotypes (35 functions), within *S.* Dublin (20 functions), *S.* Newport (20 functions), and *S.*

Typhimurium (23 functions) (FDR < 0.05, odds ratio > 6.71; Table S1). Among the bovine-

associated pseudogene functions, 13 (e.g., arginine:ornithine antiporter, multidrug transporter subunit

MdtG), 9 (e.g., acyl carrier protein), and 4 (e.g., PTS fructose transporter subunit EIIBC) were

exclusively present in all bovine isolates representing Dublin, Newport, and Typhimurium,

respectively (Table S1). By contrast, no significant geographic-associated pseudogene functions were

identified except for two (alpha-ketoglutarate transporter and peptide transporter) found enriched in

S. Newport isolates from NY.

A number of genes were associated with bovine hosts, including 30 genes found exclusively in

all bovine isolates

Among the 3,440 accessory genes for all 90 isolates, a total of 118 genes were significantly enriched

in bovine isolates (FDR < 0.05, odds ratio > 6.71), including 30 genes found among all 41 bovine

isolates regardless of serotype, but not present in any human isolates (Figure S1a; see Table S2 for a

list of all genes). Annotated functions associated with these 30 genes predominantly represented the

category "hypothetical proteins" (25 of the 30 genes); the remaining 5 genes were annotated as

encoding transposase, small toxic polypeptide LdrA/LdrC, aspartate ammonia-lyase,

permidine/putrescine ABC transporter substrate-binding protein, and racemase (Table S2).

Within *S.* Dublin, *S.* Newport, and *S.* Typhimurium, a total of 79, 64 and 82 genes,

respectively, were significantly enriched in bovine isolates (FDR < 0.05, odds ratio > 6.71); these

genes included 72, 54, and 46 genes exclusively found in all bovine isolates classified into a given

serotype, but not identified in any of the human isolates classified into the same serotype (Figure S1a,

Table S2). Based on these data we identified an overall number of 143 bovine-associated genes,

which included the 118 genes identified as bovine-associated across all serotypes as well as

additional bovine-associated genes found within each serotype. Among all these 143 bovine-

associated genes, 60 genes were annotated with known functions; 19 and 11 of these 60 genes were

annotated as encoding transposases and cell surface proteins, respectively (Figure 3, Table S3). The

annotated functions for 15 bovine-associated genes matched pseudogene functions found here to be

over-represented among human isolates; this group included some virulence factors (e.g., effector

protein YopJ, type IV secretion protein Rhs) (Table S3), suggesting that these virulence functions are

more important in bovine than human hosts.

Enrichment analysis of GO/EC terms identified 8 terms that were significantly enriched among the

41 bovine isolates representing all three serotypes, including the terms "transposition"

(GO:0032196), "double-stranded methylated DNA binding" (GO:0010385), and "hemi-methylated

DNA-binding" (GO:0044729) (Figure S1b, Table S4). Within *S. Dublin, S. Newport, and S.

Typhimurium, a total of 9 (e.g., GO:0050545:endopeptidase inhibitor activity, GO:0046148: pigment

biosynthetic process, GO:0032196:transposition), 1 (GO:0032196: transposition), and 1

(GO:0032196: transposition) GO/EC terms were significantly enriched in bovine isolates,

respectively (Table S4).

Bovine isolates exhibited significantly more transposase genes but fewer transposase

pseudogenes than human isolates.

As a large number of transposases from different IS families were identified as human-associated

pseudogene functions (Table S1) and many active transposase genes were enriched among bovine

isolates (Figure 3), further analysis was conducted on the comparison of number of pseudogenes as

well as orthologous genes (which represent "non-pseudogenes") annotated as encoding a transposase

between bovine and human isolates. Results showed that there were significantly more transposase

pseudogenes in human isolates (14, 20, 11, and 14 on average) than bovine isolates (4, 4, 5, and 4 on

average) across all serotypes, within *S. Dublin, S. Newport, and S. Typhimurium, respectively

(Figure 4a, $P < 0.0001$). By contrast, orthologous genes annotated as transposase showed an opposite

pattern. There were significantly more genes annotated as transposase among bovine isolates with an

average of 19, 29, 13, and 17 than human isolates with an average of 13, 16, 9, and 15 across all

serotypes $(P < 0.0001)$, within *S.* Dublin $(P < 0.0001)$, *S.* Newport $(P < 0.0001)$, and *S.*

Typhimurium $(P < 0.05)$, respectively (Figure 4b). Hence, both pseudogenes and genes annotated as

transposases exhibited strong host associations.

Among all 69 orthologous transposase genes (i.e., genes present in at least one of the 90

isolates studied), 39 showed strong positive correlation with AMR genes, including 16 identified as

bovine-associated transposase genes (Phi coefficient r > 0.25, Figure 5a; Table S5). Specific

examples of high correlations (Phi coefficients r > 0.8) between transposase genes and AMR genes

include (i) one IS91 family transposase gene (Cluster_3939) and *floR* (resistance to phenicols); (ii)

one IS6 family transposase genes (Cluster_3940) and *sulII* (resistance to sulfonamides) as well as

strA-B (resistance to aminoglycosides); (iii) one IS6 family transposase genes (Cluster_5580) and

tetRG (resistance to tetracyclines), *tetG* (resistance to tetracyclines) as well as *blaCARB* (resistance to

beta-lactam); and (iv) one IS1380 family transposase gene (Cluster_4114) and *sulII* (resistance to

sulfonamides).

Phi coefficient analysis was further performed to assess the correlation between transposase

genes and plasmid replicons. A total of 38 transposase genes showed strong positive correlation with

plasmid replicons, including 17 identified as bovine-associated transposase genes (Phi coefficient r >

0.25, Figure 5b; Table S6). Specific examples of high correlations (Phi coefficients r > 0.8) between

transposase genes and plasmid replicons include (i) one IS1380 family transposase gene

(Cluster_4114) and IncA/C2; (ii) one IS110 family transposase gene (Cluster_5385) and IncX1; (iii)

one IS6 family transposase gene (Cluster_5580) and IncFIB(S); (iv) three transposase genes with no

specific family annotated (Cluster_4603, Cluster_5335, and Cluster_5596) and IncFII(S), IncX1, and

IncI1, respectively. The majority (36) of these 38 transposase genes positively correlated with

plasmid replicons were also positively correlated with AMR genes (Figure 5), suggesting plasmid-

mediated transposons potentially carrying AMR genes. Notably, 9 transposase genes showing the

same pattern in the correlation with AMR genes (all positively correlated with *cmlA* and *aadB*) were

very likely located on plasmid IncX1 (these genes are shown on the right-hand side of Figure 5).

A larger number of genes and GO terms were associated with human isolates, as compared to

bovine isolates

With the same types of analyses described above for identification of bovine-associated genes, a

larger number of human-associated genes were found (Figure S1a, Table S2). As compared to 118

bovine-associated genes identified in a combined analysis of all 90 isolates representing the three

serotypes, a total of 390 genes were significantly enriched in human isolates across all serotypes

(FDR < 0.05, odds ratio > 6.71), including 119 genes present in all 49 human isolates, but absent

from all 41 bovine isolates (Figure S1a; see Table S2 for a list of all genes). Annotated functions

associated with these 119 genes predominantly represented the category "hypothetical proteins" (99

of the 119 genes); examples of annotated functions in this group of genes included genes encoding an

arsenic transporter, a ferredoxin, and a secreted effector protein SteB (Table S2).

Within *S.* Dublin, *S.* Newport, and *S.* Typhimurium, a total of 249, 244, and 244 genes,

respectively, were significantly enriched in human isolates (FDR < 0.05, odds ratio > 6.71), including

222, 195, and 188 genes found in all human isolates representing a given serotype, but not found in

any bovine isolates with a given serotype (Figure S1a, Table S2). Based on these data, we identified

an overall number of 413 human-associated genes, which included the 390 genes identified as

human-associated across all serotypes as well additional human-associated genes found within each

serotype. These 413 human-associated genes included 278 annotated as "hypothetical proteins" as

well as 135 genes with known functions; a large number of genes (n= 42) were annotated as

encoding cell surface proteins (e.g., transporters, secretion system proteins), phage-associated

proteins (e.g., integrase, phage tail protein), and transcriptional regulators (e.g., LysR, Cro/CI, AraC,

LuxR) (Figure 6, Table S7). The annotated functions for 43 human-associated genes matched

pseudogene functions found here to be over-represented among bovine isolates. As this group

includes genes previously suggested to be involved in host infection (e.g., fimbrial protein FimI,

deubiquitinase SseL, heme ABC transporter ATP-binding protein CcmA) (Table S7), our data

suggest that these genes may be more important for human than bovine infections.

At the gene ontology level, more human-associated GO and EC terms were identified

compared to bovine-associated ones (Figure S1b, Table S4). A total of 30 GO/EC terms were

significantly enriched in human isolates among all three serotypes, including the GO term "provirus

excision" (GO:0032359). While two GO terms were exclusively detected in all 49 human isolates

(GO:0071287 cellular response to manganese ion; GO:0030026 cellular manganese ion

homeostasis), these GO terms only represent a single protein (the small protein MntS, which is

classified into both GO terms) (Table S4). Within *S*. Dublin, *S*. Newport, and *S*. Typhimurium, a total

of 17, 11, and 11 GO/EC terms were significantly enriched in human isolates, respectively.

Potentially geographic-and host-associated SNPs were identified within each serotype

No geographic-associated genes or GO/EC terms were identified across all three serotypes and within

each serotype. Thus, genetic variants at SNP level were further investigated. A total of 47,913

polymorphic sites were identified among all 3,317 core genes of the 90 AMR *S. enterica* isolates. No

SNPs were shown to have a distribution significantly dependent on the geographic origin and host of

isolates across all serotypes based on Fisher's exact tests (FDR > 0.05). Potentially geographic- and

host-associated SNPs were subsequently investigated within each serotype.

Within *S*. Dublin, 342 polymorphic sites were identified in 186 Dublin core genes. Fisher's

exact test identified 13 SNPs at which the distribution of nucleotides was significantly dependent on

the geographic origin of isolates (FDR < 0.05). The most prevalent nucleotide differed between

isolates from NY and WA for each of these 13 SNPs, and 10 of them resulted in nonsynonymous

substitutions (Table 2). Five potentially geographic-associated SNPs in *S*. Dublin were located in

genes annotated as encoding cell surface proteins such as conjugal transfer protein, porin, and

fimbrial protein SefA (Table 2). By contrast, no potential host-associated SNPs were found within *S.*

Dublin (FDR > 0.05).

Within *S.* Newport, 480 polymorphic sites were identified in 204 Newport core genes. Fisher's

exact test identified 11 SNPs at which the distribution of nucleotides was significantly dependent on

the geographic origin of isolates (FDR < 0.05). The most prevalent nucleotide differed between

isolates from NY and WA for each of these 11 SNPs, and 7 of them resulted in nonsynonymous

substitutions (Table 2). Among these 11 potentially geographic-associated SNPs in *S.* Newport, 4

SNPs were located in the genes annotated as encoding cell surface proteins (e.g., transporter and 4-

hydroxyphenylacetate permease; Table 2). By contrast, no potential host-associated SNPs were found

within *S.* Newport as well (FDR > 0.05).

Within *S.* Typhimurium, 2,590 polymorphic sites were identified in 1,425 Typhimurium core

genes. While no geographic-associated SNPs were identified within *S.* Typhimurium (FDR > 0.05),

537 SNPs were found to have a distribution of nucleotides significantly dependent on the isolate host

(FDR < 0.05). All 537 potentially host-associated SNPs of *S.* Typhimurium had the same most

prevalent nucleotide among bovine and human isolates, except for one SNP located in a gene

annotated as encoding a hypothetical protein. The majority of those SNPs (77%) were found in 3

genes annotated as encoding hypothetical proteins and 2 genes annotated as encoding transcriptional

regulator and NAD(P)-dependent oxidoreductase, respectively. For a total of 523 of the 537 SNPs

(97%) a monophyletic group of 8 human isolates (Figure 5; Liao *et al* (2019)) harbored a different

nucleotide compared to 12 other human isolates and all 17 bovine isolates, indicating that the

significant host association of these SNPs likely represents a single human host-associated

Typhimurium lineage.

In summary, 24 SNPs potentially associated with geographic origin were identified within *S.*

Dublin and *S.* Newport; 9 SNPs (38%) were located in genes annotated as encoding cell surface

proteins (Table 2). The majority of these SNPs (71%) resulted in nonsynonymous substitutions

(Table 2), including one located in a gene (annotated as encoding E3 ubiquitin--protein ligase)

showing evidence for positive selection as reported by Liao *et al.*, 2019 using the same set of isolates.

While a number of potentially host-associated SNPs were identified within AMR *S.* Typhimurium,

those were likely just associated with one lineage of AMR *S.* Typhimurium.

Few closely related isolates were identified between bovine and human and between NY and

WA sources based on high-quality core SNPs

Our results provided evidence for associations between genetic variations (e.g., gene

presence/absence, SNPs) and isolate sources (e.g. human or bovine), suggesting limited transmission

of AMR *Salmonella* between bovine and human populations, and between NY and WA. We thus

used high-quality (hq) core SNP data to identify closely related isolates in order to further

characterize the frequency of recent transmission events between different sources among the 90

isolates characterized here. Overall, isolates from different geographic origins as well as isolates from

different hosts were not very closely related. Only 6, 4, and 0 human isolates for the three serotypes

(Dublin, Newport, Typhimurium) showed <10 SNP difference to at least one bovine isolate; 7 human

S. Typhimurium isolates showed <50 SNP difference to at least one bovine *S.* Typhimurium isolate

(see Figure S2a, Figure S2c, Figure S2e for hqSNP distance matrices for comparison between the

two hosts for each serotype). For comparison between the two geographic locations, only 2, 1, and 0

isolates from NY for the three serotypes (Dublin, Newport, Typhimurium) showed <10 SNP

difference to at least one WA isolate; 2 *S.* Typhimurium isolates from NY showed <50 SNP

difference to at least one *S.* Typhimurium isolate from WA (see Figure S2b, S2d, S2f for hqSNP

distance matrices for comparison between the two geographic locations for each serotype).

Discussion

Salmonella has a broad host range including humans, other mammals, birds, and reptiles (Eng *et al.*,

2015; Schleker *et al.*, 2012). *Salmonella* can also survive for a long period of time in water (e.g.,

sewage, freshwater, marine coastal water, and groundwater) (Baudart *et al.*, 2000). This ability of

Salmonella to survive and multiply in a wide range of ecological niches and to cause disease in

diverse hosts is not only practically significant (as many sources can be responsible for a human

infection), but also makes this organism a relevant model for studying the population genetics and

evolution of environmentally transmitted zoonotic pathogens. In addition, the fact that antimicrobial

resistance, including multi-drug resistance, has emerged multiple times in different *Salmonella*

serotypes, makes studies on the population structure of different AMR *Salmonella* serotypes

important to help inform design of control and prevention strategies and to better understand the

evolution and potential for inter-species transmission of AMR zoonotic pathogens with wide host

ranges. A number of previous studies have identified genetic markers associated with host adaptation

in *S. enterica* (Langridge *et al.*, 2015; Lupolova *et al.*, 2017; Thomson *et al.*, 2008; Yue *et al.*, 2015).

For example, Thomson *et al.* (2008) reported that chicken-restricted *S. enterica* serovar Gallinarum

isolates showed extensive genome degradation through deletion and pseudogene formation, including

the loss of flagella or fimbriae. Lupolova *et al.* (2017) demonstrated that, based on gene content, the

source hosts including human, bovine, and swine of *S.* Typhimurium can be accurately predicted

using machine learning approach. However, only a few studies, typically focused on non-AMR

strains, have explored genetic markers associated with geographic origins of *S. enterica* isolates

(Sangal *et al.*, 2010, Zheng *et al.*, 2017). While an initial study has shown that AMR genes of *S.*

enterica exhibited a strong association with both host source and geographic origin (Carroll *et al.*,

2017), our study reported here indicates that different evolutionary forces appear to drive AMR *S.*

enterica population structure in human and bovine hosts and geographic regions. Specifically, our

data not only suggests that transposition plays a more important role in genome evolution and

antibiotic resistance transmission of contemporary bovine AMR isolates, as compared to human

isolates, but also suggests that association of different virulence-associated genes and pseudogenes

with bovine and human AMR *Salmonella* could represent a barrier to inter-host transmission.

Importantly, our findings also suggest that specific genetic markers may be able to help predict host

or geographic origin of AMR *Salmonella* isolates.

The host-transposase association suggests that the occurrence time of large-scale transposition

and adaptation strategies differ by AMR *S. enterica* populations from bovine and human.

Transposase is a DNA-binding enzyme that binds to the end of a transposon and catalyzes the

movement of the transposon to another location of the genome via 'cut-and-paste' or 'copy-and-

paste' mechanisms (Rice and Baker, 2001). The significant enrichment of intact genes annotated as

encoding transposase among bovine isolates suggests ongoing large-scale transposition and genomic

rearrangement events in bovine AMR *S. enterica* populations. By contrast, those events probably

occurred earlier in human AMR *S. enterica* populations, since a substantial number of genes

annotated as encoding transposase represent pseudogenes in genomes of human AMR *S. enterica* and

were not functional anymore. Consistent with our findings, bovine *S. agalactiae* genomes showed a

significantly elevated number of transposable elements compared to human *S. agalactiae* genomes

(Richards *et al*., 2011), indicating that such distribution patterns of transposase influenced by host is

not restricted to *S. enterica*. This phenomenon might be caused by the different stages of host

adaptation/restriction pathogens are experiencing in different hosts. As reviewed by Moran and

Plague (2004), when bacteria switch from a free-living to host restriction lifestyle, IS elements in

genomes are expected to proliferate during the short-term process of host dependence, with the

purpose of reducing genome size by disrupting existing regulatory regions and inactivating genes via

transposons (Feschotte, 2008); in the long run, mobile elements are gradually deleted or mutated by

forming pseudogenes due to lack of exposure to novel element transposition. As such, facultative

intracellular species including *S. enterica* normally harbor dramatically more transposase genes than

obligate intracellular bacteria (e.g., *Wigglesworthia glossinidia, Buchnera aphidicola,*

Chlamydophila caviae) (Bordenstein and Reznikoff, 2005). Additionally, high numbers of

transposase genes inactivated by forming pseudogenes have been observed in the luminous bacterial

symbionts of deep-sea ceratiod anglerfishes (Hendry *et al.*, 2018). Based on these mechanisms of

genomic changes and previous findings, our data suggests that AMR *S. enterica* in human

populations may be at a later stage of host adaptation than those in bovine populations. Adaptation to

a human host has also been observed in other host generalist serotypes of *Salmonella*. Klemm *et al.*

(2016) detected substantial genome degradation via pseudogene formation and an elevated

substitution rate in *S*. Enteritidis which had infected an immunocompromised human patient for 15

years, highlighting the fast within-human evolution of *Salmonella*.

Besides potentially taking advantage of transposons to degrade the genome during a likely

early stage of host adaptation, it is also possible that bovine AMR *S. enterica* are maintaining a large

number of transposase genes in their genomes due to some beneficial mutations responsible for

adaptive phenotypic changes brought by transposons (Casacuberta and González, 2013). One such

benefit may be promoting bacterial adaptation to antibiotics used in clinical practice and agriculture

(Blot, 1994). Our data showed that more than half of the transposase genes identified in this study

showed a significant positive correlation with AMR genes. Congruous with our finding, a variety of

transposons carrying antimicrobial resistance genes have been identified in *Salmonella* as

summarized in Miriagou *et al.*, (2006). For example, the tetracycline resistance gene *tet*(A), which

encodes a membrane-associated efflux protein, has been reported to be linked to transposon Tn1721

(Pezzella *et al.*, 2004). bla_{TEM-1} which encodes β-lactamases that mediate resistance to β-lactam

antimicrobial agents, has been shown to be carried by Tn3 (Pasquali *et al.*, 2005). The *strA-strB*

genes, which confer streptomycin resistance, have been described as part of a particular Tn5393-

derivative transposon (Carattoli *et al.*, 2002). The *armA* gene, which encodes aminoglycoside

resistance methylase and confers resistance to 4,6-disubstituted deoxystreptamines and fortimicin,

has been described as linked to a composite transposon Tn1548 (Galimand *et al.*, 2005). Class 1 and

class 2 integrons, which often harbor gene cassettes encoding aminoglycoside modifying enzymes,

are commonly associated with various transposons of the Tn3 family (e.g., Tn21, Tn1696, and

Tn1412) and Tn7 transposon, respectively (Miriagou *et al.*, 2006). In our study, *tet*(A) was found to

be highly correlated with IS1380 family transposase genes; bla_{TEM-1D} was highly correlated with

IS200/IS605 family transposase, IS630 family transposase, and other two non-specific family

annotated transposase genes; *strA-strB* were highly correlated with IS6 family transposase and

IS1380 family transposase genes. Selection of antibiotic resistance genes carried by transposons is a

strong force in maintaining transposons in populations (Blot, 1994). Thus, keeping a large number of

transposase genes could be a strategy employed by bovine AMR *S. enterica* to better adapt to the

antibiotics used in cows.

Identification of host associated genes and pseudogenes encoding presumptive virulence factors

suggests adaptation of AMR *S. enterica* to human and bovine hosts.

Our analyses identified a number of host-associated genes, including genes found exclusively in

isolates from one host (i.e., human or bovine) as well as genes that more frequently presented

premature stop codons (hence forming pseudogenes) in isolates from one host. These findings

provide initial evidence for adaptation of AMR *S. enterica* to bovine and human populations, at least

in the isolate set analyzed here. More specifically, identification of host-association in genes

annotated as encoding potential virulence factors supports such adaptation. We specifically identified

a number of putative virulence factor-encoding genes that were overrepresented among human AMR

S. enterica relative to bovine isolates, suggesting not only a specific role of these genes in human

infections, but also indicating the possibility of reduced ability to infect human hosts among at least

some bovine isolates. For example, the genes encoding the deubiquitinase SseL and fimbrial protein

FimI were found to be significantly enriched in human isolates; on the other hand, *sseL* and *fimI*

pseudogenes were significantly enriched in bovine isolates. Deubiquitinase SseL, one of the

Salmonella SPI-2 Type III secretion system effectors, shows deubiquitinating activity during

infection of human and muriune cell lines (Rytkönen *et al.*, 2007). Jennings *et al*. (2017) furthermore

reported that SseL inhibits accumulation of lipid droplets, prevents autophagic clearance of cytosolic

aggregates, and induces late macrophage cell death. *fimI* is required for production of normal type 1

fimbriae and is located in the *fim* gene cluster (Knight and Bouckaert, 2009). It is currently not clear

whether FimI constitutes a subunit of type 1 fimbriae or is a protein regulating the assembly of type 1

fimbriae (Knight and Bouckaert, 2009). It has been reported that mannose-sensitive type 1 fimbriae

of *S.* Typhimurium promote invasion and appear to play a critical role as an accessory virulence

factor (Ernst *et al.*, 1990). As mutations in *fimI* lead to dysfunctional type 1 fimbriae (Valenski *et al.*,

2003), absence of functional FimI (e.g., through formation of pseudogenes, as found here among

bovine isolates) may lead to reduced virulence at least in some hosts. We also found that the gene

encoding SteB is exclusively found in all human isolates but absent from all bovine isolates in this

study. SteB is an effector that requires *Salmonella* SPI-1or SPI-2-encoded T3SS for its translocation

(McGhie *et al.*, 2009). SteB has been reported to act as a putative picolinate reductase which is

required for efficient mouse spleen colonization in *Salmonella* (McGhie *et al.*, 2009). Identification

of putative and well-documented virulence factors encoded by human AMR *S. enterica*, but absent or

inactivated in bovine isolates not only indicates that these genes encode host specific virulence

factors, but also suggests adaptation of *Salmonella* to human hosts.

In addition, we also identified some virulence genes associated with bovine hosts. For example,

the genes annotated as encoding the type IV secretion protein Rhs and the effector protein YopJ were

not only found to be significantly enriched in bovine isolates, but *rhs* and *yopJ* pseudogenes were

also overrepresented among human isolates. *rhs* genes were first described in *E. coli*, and

subsequently found in a wide range of Gram-negative bacteria, including other members of the

Enterobacteriaceae (e.g., *Salmonella*) as well as *Pseudomonadaceae*. Rhs has been shown to be a

mammalian virulence determinant in *Pseudomonas aeruginosa*; Rhs was found to be induced during

infection of monocyte/macrophage-like cells, was translocated into these cells, and subsequently

caused inflammasome-mediated cell death (Kung *et al.*, 2012). YopJ effectors have been found in a

variety of animal pathogens including *Yersinia* spp., *S. enterica*, *Vibrio parahaemolyticus* and

Aeromonas salmonicida (Ma and Ma, 2016). These effectors target proteins in hosts by acetylating

specific serine, threonine, and/or lysine residues, thus influencing the function and/or stability of their

target proteins and eventually dampening innate immunity of hosts (Ma and Ma, 2016). Our findings

suggest that the type IV secretion protein Rhs and the effector protein YopJ may play a more

important role in AMR *S. enterica* during infection of bovine hosts than they do in humans.

Consistent with our results, *S.* Typhimurium has been shown to use host-specific bacterial factors

(e.g., type III secretion systems, cell surface polysaccharides, cell envelope proteins) to colonize

calves and chicks (Morgan *et al.*, 2004).

Furthermore, we identified a number of genes without likely virulence-associated functions that

were associated with either bovine or human isolates. While many annotated bovine-associated genes

represented transposases, a number of human-associated genes represented other functions. For

example, the small protein MntS, which helps to enlarge the manganese pool when manganese is

scarce in *E. coli* (Martin *et al.*, 2015), was encoded by a gene exclusively found in all human AMR *S.*

enterica isolates but absent from all bovine isolates in this study. *ccmA* was also significantly

associated with human isolates, while a *ccmA* pseudogene was overrepresented among bovine

isolates; *ccmA* encodes the heme ABC transporter ATP-binding protein CcmA, a putative ATPase

essential for TMA/TMAO and nitrite/nitrate as terminal electron acceptors under conditions of

reduced oxygen (Batista *et al.*, 2015). Interestingly, all *S.* Pullorum genomes analyzed by Batista *et*

al. (2015) also carried a *ccmA* pseudogene; *S.* Pullorum is highly adapted to poultry. In addition, a

large proportion of human-associated genes in our study reported here were annotated as encoding

transcriptional regulators (e.g., AraC, LuxR, Rha family transcriptional regulator) and phage-

associated proteins; Lupolova *et al.* (2016) previously also reported that human *E. coli* O157 isolates

showed a higher frequency of phage-associated genes relative to bovine isolates. The specific roles of

those host-associated genetic variants in differential adaptation in *S. enterica* remain unknown and

further exploration is needed.

In summary, our findings suggest that AMR *S. enterica* serotypes Dublin, Newport, and

Typhimurium show evidence of adaptation to human and bovine hosts, including through

development of repertoires of host-specific virulence genes. While this raises the intriguing

possibility that genetic mechanisms may restrict the transmission of AMR *Salmonella*, at least for the

three serotypes studied here, between human and bovine hosts, further work on larger isolate sets,

including those from different regions, are needed to confirm this hypothesis. Importantly, however,

our findings are consistent with other studies which show the development of host-specific subgroups

within closely related pathogen taxa. For example, different alleles of *fimH* encoding type 1 fimbrial

adhesin showed different binding preference to human cells and bovine cells (Yue *et al.*, 2015).

Evolution of host specificity has also been well established in a number of other bacterial pathogens,

as reviewed by Bäumler and Fang (2013).

The geographic-associated SNPs highlight the role of geographic origin in influencing

population structure of AMR *S. enterica* and evolution of cell surface proteins

Potential geographic-associated SNPs were found within AMR *S.* Dublin and AMR *S.* Newport. *S.*

Newport has been reported to display a geographic structure (Sangal *et al.*, 2010). Specifically,

Lineage I strains were more frequently isolated from Europe, while Lineages II and III strains were

more likely isolated from North America. Such geographic association of lineages was suggested to

be caused by differences in prophage regions, pathogenicity islands and fimbrial operons among

strains from different geographic locations (Zheng *et al.*, 2017). AMR *S.* Newport analyzed here

belong to Lineage IIC as previously reported by Liao *et al.* (2019). Our findings suggest that within

this lineage, *S.* Newport also has a geographic structure at the SNP level.

A large proportion of geographic-associated SNPs found in *S.* Dublin and *S.* Newport (38%)

were located in genes annotated as encoding cell surface proteins (e.g., cardiolipin synthase B, porin,

permease, transporter, fimbrial protein SefA), higher than the proportion of cell surface proteins

among the total amount of proteins in eukaryotic and prokaryotic cells in general (20%-30%).

Interestingly, cardiolipin synthase B, which catalyzes the synthesis of cardiolipin in bacteria, has

been found to be important in modulating the physical properties of membranes in response to

environmental stress such as osmotic stress (Mileykovskaya and Dowhan, 2009). Mutation of

cardiolipin synthase has been shown to be associated with daptomycin resistance in enterococci.

Davlieva *et al*. (2013) reported that when challenging *Enterococcus faecium* and *Enterococcus*

faecalis with daptomycin, the pool of phosphatidylglycerol was substantially reduced. However,

mutations observed in gene encoding cardiolipin synthase were able to compensate for the decreased

phosphatidylglycerol and lead to a restoration of the cardiolipin synthase pool. In addition, Peleg *et*

al., 2012 found genomic evidence that point mutations in gene coding for cardiolipin synthase were

responsible for developing reduced susceptibility to daptomycin in *Staphylococcus aureus*.

Daptomycin has been a widely used antibiotic in the treatment of complicated skin and skin structure

infections in humans (Steenbergen *et al*., 2009). While highly speculative, the geographically

associated SNPs identified in genes encoding cardiolipin synthase B for *S*. Newport could suggest

possible follow-up studies on use of daptomycin in NY and WA.

Another interesting gene, annotated as encoding an exported protein, E3 ubiquitin--protein

ligase, contained a geographic-associated SNP resulting in a nonsynonymous substitution, and this

gene showed evidence of positive selection in a previous study using this same set of isolates (Liao *et*

al., 2019). Genes encoding E3 ubiquitin--protein ligase are present in several species of pathogenic

bacteria, including *Salmonella* (Quezada *et al.*, 2009), and this protein has been reported to be used

as a virulence factor (Maculins *et al.*, 2016). Multiple studies have shown that E3 ubiquitin ligase

was used by *Salmonella* to hijack host cells and subvert its ubiquitination pathway (Maculins *et al.*,

2016; Quezada *et al*., 2009). Ubiquitination systems function in a wide range of cellular processes

(e.g., cell cycle and division, immune response and inflammation) in eukaryotic organisms, and can

lead to cancer and neurodegenerative disorder when ubiquitination is defective (Finley and Chau,

1991). Thus, expressing and releasing of E3 ubiquitin ligase by *Salmonella* could promote bacterial

survival and pathogenicity in the host. A geographically associated SNP found in a gene encoding E3

ubiquitin--protein ligase might be a consequence of positive selection triggered by specific ecological

niches associated with NY or WA.

Overall, our findings suggest that, in addition to host, geographic origin is also a critical factor

contributing to the population structure, antimicrobial resistance, and pathogenicity of AMR *S.*

enterica, likely due to distinct ecological niches (e.g., represented by climate, dairy operation

management, antibiotics use) associated with different geographic origins.

AMR *S. enterica* population structure is likely driven by different evolutionary forces and a low

potential of recent transmission among hosts and geographic origins.

It has been demonstrated that ecological niches can change population structure and affect the

evolution of bacteria (Vos, 2011; Li *et al*., 2019). A recent study (Behringer *et al*., 2018) on the long-

term evolution of *E. coli* cultures proposed that the evolution of genetic differentiation resulting in

subpopulation structure was facilitated by spatial differentiation followed by metabolic

differentiation. To cope with different ecological niches, adaptive evolutionary processes mediated

by a mixture of mechanisms - mainly gene flow, mutation, natural selection, and genetic drift – will

occur in bacteria, resulting in gene gain/loss and functional divergence of existing genes (Toft and

Andersson, 2010; Li *et al*., 2019; Slatkin, 1987; Strachan *et al*., 2015). Our data provides evidence in

strong association of gene gain/loss with host as well as evidence for SNP level divergence of core

genes associated with geographic origin (in at least some AMR *S. enterica* serotypes), suggesting

different mechanisms are underlying adaptation of AMR *S. enterica* to host and to geographic

location. For example, the strong association between bovine-associated transposases and plasmids

(e.g., IncX2 and IncFII(S)) observed in this study indicates that some transposons in bovine AMR *S.*

enterica were likely obtained via horizontal gene transfer. Consistent with our finding, transposons

have been frequently detected on conjugative or mobilizable plasmids in various *Salmonella* obtained

from different sources (Miriagou *et al*., 2006). Most of human-associated pseudogenes are the result

of frameshifts and/or premature stop codons caused point mutations, which are common mechanisms

in the formation of pseudogenes in bacterial genomes (Lerat and Ochman, 2005). While

geographically associated SNPs were not exclusive under positive selection, nonsynonymous SNPs

in genes encoding cell surface proteins are potential targets of positive selection, as cell surface

proteins have been found to be a major group driven by positive selection in bacteria (Chen *et al*.,

2006; Petersen *et al*., 2007; Wachter and Hill, 2016). By contrast, synonymous geographic-associated

SNPs may be a consequence of genetic drift, as genetic drift is the driving force for neutral mutations

(Bromham and Penny, 2003). Overall, our findings suggest that gene flow and mutation are playing

important but not exclusive roles in shaping host-associated population structure of AMR *S. enterica*,

while natural selection and genetic drift are essential, but not exclusive, in shaping geographically

associated population structure of AMR *S. enterica*.

In addition to different evolutionary forces influencing the population structure of AMR *S.*

enterica, our data together with previously reported associations between AMR genes and isolate

source (Carroll *et al*., 2017) also suggests a low potential for recent transmission of AMR

populations between bovine and human populations and between NY and WA. Transmission of

bacteria can promote genetic exchanges, so a more homogenized gene pool and closely related

individuals are expected (Slatkin, 1987). In this study, AMR *S. enterica* showed high degree of host-

genome and geographic-genome associations at gene, GO terms, pseudogene, and SNP levels, and

absence of a large number of closely related isolates between bovine and human populations and

between NY and WA. Similarly, Mather *et al*. (2013) demonstrated that *S.* Typhimurium DT104 and

its antimicrobial resistance genes were maintained largely separately within animal and human

populations in Scotland, with little transmission between animals and humans and vice versa.

Overall, these data suggest that transmission of *Salmonella* between bovine and human hots may be

less frequent than sometimes assumed, even though there have been cases of AMR *S.* Dublin

(Harvey *et al.*, 2017), *S.* Newport (Centers for Disease Control and Prevention, 2002; Plumb *et al*.,

2019), *S*. Typhimurium (McLaughlin *et al*., 2006; Olsen *et al*., 2004) outbreaks in humans linked to

raw milk, raw beef or cheese. Our findings are also consistent with some studies that indicated

geographical association with different *Salmonella* clonal groups; for example, Kovac *et al*. (2017)

reported that *S*. Cerro found in Texas and NY represent largely distinct clonal groups. As NY and

WA are about 3,500 km apart, it is plausible that such a distance could generate a barrier for

Salmonella transmission, since microbial dispersal is typically a passive process, which does not

likely result in long-distance dispersal events (Nemergut *et al*., 2013). While contemporary complex

food distribution systems may reduce the effectiveness of distance as a barrier for transmission, our

data suggest that geographical distance may still represent a barrier to long-distance *Salmonella*

dispersal.

Conclusion

The population structure of antimicrobial resistant bacterial pathogens which differs in host and

geographic origin could reveal key elements in understanding their evolution in adaptation,

pathogenesis, and antimicrobial resistance. Collectively, our study identified a number of genetic

variants in AMR *S. enterica* associated with host and geographic origin using comparative genomics.

It highlights a vital role of host in shaping genome arrangement and promoting antimicrobial

resistance via potentially plasmid-mediated transposons and in developing host adaption via

preferred virulence factors. It also highlights a critical role of geographic origin in driving the

evolution of cell surface proteins and antimicrobial resistance potentially mediated by positive

selection and genetic drift. Even though the advancement of sequencing techniques has allowed for

rapid progress in identifying niche-associated genetic variants, our understanding of the evolutionary

mechanisms underlying the generation, fixation and function of those variants in populations is still

limited. Our study provides a large number of candidate genes, such as secreted effector protein

SteB, fimbrial protein FimI, E3 ubiquitin--protein ligase, for future investigations to better

understand the adaptation and pathogenesis of AMR *S. enterica*. As AMR *S. enterica* represent a

substantial public health concern, focusing on AMR populations allowed us to identify specific

genetic markers that could be potentially used in source tracking (e.g., host and geographic origins)

of human disease cases and contamination events caused by AMR *S. enterica*. Our data also suggest

a low potential for inter-host and long distance-geographic transmission events of AMR *Salmonella*.

Despite the benefits of focusing on AMR populations, this study is limited by a relatively small

dataset. To fully understand the genetic basis of environmental adaptation and differential virulence

associated with host and geographic origin in *S. enterica*, it is necessary for future studies to include

non-AMR populations as well.

Author Contributions

JL and MW conceived and designed the study. JL wrote the manuscript with input from MW, RHO

and LC. JL performed the computational analyses and statistical analyses.

Conflict of Interests Statement

The authors declare that they have no competing interests.

Acknowledgement

This work was partially supported by the USDA National Institute of Food and Agriculture Hatch

project NYC-143436. Any opinions, findings, conclusions, or recommendations expressed in this

publication are those of the author(s) and do not necessarily reflect the view of the National Institute

of Food and Agriculture (NIFA) or the United States Department of Agriculture (USDA). We are

grateful to Dr. Rachel A. Cheng for her assistance in interpreting functions of host-associated genes.

Figures

Figure 1. Gene presence/absence based-Non-metric multidimensional scaling ordination (NMDS) of

antimicrobial-resistant (AMR) *S. enterica* isolates (a) across all serotypes; (b) within *S.* Dublin, (c)

within *S.* Newport, and (d) within *S.* Typhimurium. Isolates from New York state (NY) are indicated

by symbol circle; isolates from Washington state (WA) are indicated by symbol triangle. Bovine

isolates (BOV) are in blue; human isolates (HUM) are in red. For panel (a), *S.* Dublin are indicated

by "+"; *S.* Newport are indicated by "×"; *S.* Typhimurium are indicated by "◊". Dotted lines indicate

clustering of isolates correspond to host.

Figure 2. Box and whisker plot of the number of pseudogenes in AMR *S. enterica* isolates from (a)

bovine and human populations, and (b) NY and WA among all three serotypes and within each

serotype. DUBN - *S.* Dublin; NEWP - *S.* Newport; TYPH - *S.* Typhimurium; *P* values were

determined by Mann-Whitney test after FDR correction. Minimum and maximum values are

depicted by short horizontal lines above and below the box; the box signifies the upper and lower

quartiles, and the median is represented by a short line within the box.

Figure 3. Heatmap of the 60 bovine-associated of AMR *S. enterica* orthologous genes annotated

with known functions. Genes encoding transposases, cell surface proteins, and phage-associated

proteins are highlighted in blue, purple, and red, respectively; genes encoding other functions are

indicated in dark grey. Light grey boxes indicate absence of genes (see key). Isolates (shown on y

axis) are color coded by serotype, with green indicating *S.* Dublin, yellow indicating *S.* Newport, and

orange indicating *S.* Typhimurium; bovine isolates are highlighted in bold. Details of these 60

bovine-associated genes are shown in Table S3.

Figure 4. Box and whisker plot of number of (a) pseudogenes and (b) orthologous genes annotated

as encoding transposases in AMR *S. enterica* isolates from bovine and human populations. Bovine

isolates are indicated by blue boxes; human isolates are indicated by red boxes; *P* values were

determined by Mann-Whitney test after FDR correction. Minimum and maximum values are

depicted by short horizontal lines above and below the box; the box signifies the upper and lower

quartiles, and the median is represented by a short line within the box.

Figure 5. Heatmaps of Phi coefficient between (a) transposase genes and AMR genes, and (b) transposase genes and plasmid replicons. Dark grey boxes indicate that the positive correlation is not strong (Phi coefficient < 0.25). Bovine-associated transposase genes are highlighted in bold. Gray parallelograms between panels a and b link clusters (e.g., Cluster_3939) that showed strong positive correlations with both AMR genes and plasmid replicons.

Figure 6. Heatmap of the 135 human-associated AMR *S. enterica* orthologous genes annotated with

known functions. Genes encoding transposases, cell surface proteins, phage-associated proteins, and

transcriptional factors are highlighted in blue, purple, red, and green, respectively; genes encoding

other functions are indicated in dark grey. Light grey boxes indicate absence of genes (see key).

Isolates (shown on y axis) are color coded by serotype, with green indicating *S.* Dublin, yellow

indicating *S.* Newport, and orange indicating *S.* Typhimurium; human isolates are highlighted in

bold. Details of these 135 human-associated genes are shown in Table S7.

757   **Tables**

758   Table 1 PERMANOVA of gene presence/absence for isolate groups of AMR *S. enterica*[a]

Serotypes	Grouping factor[b]	F Statistic	Uncorrected *p*-value	FDR[c]
All	**Host**	**13.98**	**< 0.001**	**< 0.001**
All	Geographic origin	0.86	0.48	0.48
***S.* Dublin**	**Host**	**21.37**	**< 0.001**	**< 0.001**
S. Dublin	Geographic origin	2.75	0.04	0.06
***S.* Newport**	**Host**	**17.686**	**< 0.001**	**< 0.001**
S. Newport	Geographic origin	2.24	0.05	0.06
***S.* Typhimurium**	**Host**	**10.37**	**< 0.001**	**< 0.001**
S. Typhimurium	Geographic origin	1.21	0.26	0.29

759   [a] Rows in boldface indicate that FDR test was significant (FDR < 0.05).

760   [b] Host – bovine vs human, Geographic origin – NY vs WA.

761   [c] FDR was determined by a Benjamini and Hochberg correction.

762 Table 2 Annotated functions of genes which contain potentially geographic-associated SNPs within *S.* Dublin and *S.* Newport

Gene	Function[a]	Codon with nucleotide enriched in NY (AA) [b]	Codon with nucleotide enriched in WA (AA) [b]
S. Dublin			
Cluster_82	integral component of membrane[#]	CAG (**Gln**)	CGG (**Arg**)
Cluster_208	integral component of membrane[#]	AAA(**Lys**)	GAA(**Glu**)
Cluster_104	4-(cytidine 5'-diphospho)-2-C-methyl-D-erythritol kinase	CTC (**Leu**)	CCC (**Pro**)
Cluster_4108	DNA polymerase III subunit epsilon	CCC(**Pro**)	TCC(**Ser**)
Cluster_5206	hypothetical protein	CCA(**Pro**)	ACA(**Thr**)
Cluster_5312	DUF4755 domain-containing protein	AGT(**Ser**)	GGT(**Gly**)
Cluster_3703	ATP-dependent protease	GCC(Ala)	GCT(Ala)
Cluster_4136	restriction endonuclease subunit M	GCA(Ala)	GCC(Ala)
Cluster_4288	RNA-splicing ligase RtcB	TAT(**Tyr**)	GAT(**Asp**)
Cluster_4195	conjugal transfer protein[#]	GTA(**Val**)	GCA(**Ala**)
Cluster_195	cytosol nonspecific dipeptidase	GCG(**Ala**)	ACG(**Thr**)
Cluster_476	porin[#]	GCA(Ala)	GCT(Ala)
Cluster_5407	fimbrial protein SefA[#]	CAG(**Gln**)	AAG(**Lys**)
S. Newport			
Cluster_1979	transporter[#]	ATC(**Ile**)	GTC(**Val**)
Cluster_3964	phage tail protein	TTC(**Phe**)	TGC(**Cys**)
Cluster_3624	acetyl-CoA carboxylase subunit beta	GCC(Ala)	GCG(Ala)
Cluster_2245	penicillin-binding protein activator	CTA(Leu)	CTG(Leu)
Cluster_3735	cytoplasmic protein[#]	TAC(**Tyr**)	CAC(**His**)
Cluster_3207	cardiolipin synthase B[#]	GGT(Gly)	GGC(Gly)
Cluster_1005	AraC family transcriptional regulator	TGT(**Cys**)	TAT(**Tyr**)
Cluster_3568	2-succinyl-5-enolpyruvyl-6-hydroxy-3- cyclohexene-1-carboxylic-acid synthase	GTG(**Val**)	GCG(**Ala**)
Cluster_815	DUF179 domain-containing protein	TGC(**Cys**)	CGC(**Arg**)
Cluster_2311	E3 ubiquitin--protein ligase	TGG(**Trp**)	CGG(**Arg**)
Cluster_3830	4-hydroxyphenylacetate permease[#]	ACA(Thr)	ACC(Thr)

[a] # indicates that genes were annotated as encoding cell surface proteins.

[b] SNPs resulting in nonsynonymous substitutions (i.e., change in amino acid (AA)) and relevant AA are in bold; A-Adenine, T-Thymine, G-Guanine, C-Cytosine; Ala-

Alanine, Arg-Arginine, Asp-Aspartate, Cys-Cysteine, Gln-Glutamine, Glu-Glutamate, Gly-Glycine, His-Histidine, Ile-Isoleucine, Leu-Leucine, Lys-Lysine, Phe-

Phenylalanine, Pro-Proline, Ser-Serine, Thr-Threonine, Trp-Tryptophan, Tyr-Tyrosine, Val-Valine.

Reference

Abeles, S.R. and Pride, D.T. (2014) Molecular bases and role of viruses in the human microbiome. J
Mol Biol 426: 3892-3906.

Anderson, M.J. and Walsh, D.C.I. (2013) PERMANOVA, ANOSIM, and the Mantel test in the face
of heterogeneous dispersions: What null hypothesis are you testing? Ecol Monogr 83: 557-574.

Ao, T.T., Feasey, N.A., Gordon, M.A., Keddy, K.H., Angulo, F.J., and Crump, J.A. (2015) Global
burden of invasive nontyphoidal *Salmonella* disease 2010. Emerg Infect Dis 21: 941.

Batista, D.F.A., Neto, O.C.F., Barrow, P.A., de Oliveira, M.T., Almeida, A.M., Ferraudo, A.S., and
Berchieri Jr, A. (2015) Identification and characterization of regions of difference between the
Salmonella Gallinarum biovar Gallinarum and the *Salmonella* Gallinarum biovar Pullorum
genomes. Infect Genet Evol 30: 74-81.

Baudart. J., Lemarchand, K., Brisabois, A., Lebaron, P. (2000) Diversity of *Salmonella* strains
isolated from the aquatic environment as determined by serotyping and amplification of the
ribosomal DNA spacer regions. Appl Environ Microbiol 66: 1544-52.

Bäumler, A. and Fang, F.C. (2013) Host specificity of bacterial pathogens. Cold Spring Harb
Perspect Med 3: a010041.

Behringer, M.G., Choi, B.I., Miller, S.F., Doak, T.G., Karty, J.A., Guo, W., and Lynch, M. (2018)
Escherichia coli cultures maintain stable subpopulation structure during long-term evolution. Proc
Nat Acad Sci USA 115: E4642-E4650.

Blot, M. (1994) Transposable elements and adaptation of host bacteria. Genetica 93: 5-12.

Bordenstein, S.R. and Reznikoff, W.S. (2005) Mobile DNA in obligate intracellular bacteria. Nat
Rev Microbiol 3: 688.

Bromham, L. and Penny, D. (2003) The modern molecular clock. Nature Rev Genet 4: 216-224.

Carattoli, A., Filetici, E., Villa, L., Dionisi, A.M., Ricci, A., and Luzzi, I. (2002) Antibiotic resistance

genes and *Salmonella* genomic island 1 in *Salmonella enterica* serovar Typhimurium isolated in

Italy. Antimicrob Agents Chemother 46: 2821-2828.

Carroll, L.M., Wiedmann, M., den Bakker, H., Siler, J., Warchocki, S., Kent, D., *et al.* (2017) Whole-

genome sequencing of drug-resistant *Salmonella enterica* isolates from dairy cattle and humans in

New York and Washington States reveals source and geographic associations. Appl Environ

Microbiol 83: e00140-17.

Casacuberta, E. and González, J. (2013) The impact of transposable elements in environmental

adaptation. Mol Ecol 22: 1503-17.

Centers for Disease Control and Prevention. (2017) National *Salmonella* surveillance annual report

and appendices, 2015.

Centers for Disease Control and Prevention (2002). Outbreak of multidrug-resistant *Salmonella*

Newport--United States, January-April 2002. Morbidity and Mortality Weekly Report. 51: 545.

Chen, H., Cohen, P., and Chen, S. (2010) How big is a big odds ratio? Interpreting the magnitudes of

odds ratios in epidemiological studies. Commun Stat Simul Comput 39: 860-864.

Chen, S.L., Hung, C.S., Xu, J., Reigstad, C.S., Magrini, V., Sabo, A., *et al.* (2006) Identification of

genes subject to positive selection in uropathogenic strains of *Escherichia coli*: a comparative

genomics approach. Proc Nat Acad Sci USA 103: 5977-5982.

Dandekar, T., Fieselmann, A., Fischer, E., Popp, J., Hensel, M., and Noster, J. (2015) *Salmonella*-

how a metabolic generalist adopts an intracellular lifestyle during infection. Front Cell Infect

Microbiol 4: 191.

Davlieva, M., Zhang, W., Arias, C.A., and Shamoo, Y. (2013) Biochemical characterization of

cardiolipin synthase mutations associated with daptomycin resistance in enterococci. Antimicrob

Agents Chemother 57: 289-296.

Ellermeier, C.D. and Slauch, J.M. (2006) The genus *Salmonella*. The Prokaryotes: Volume 6:

Proteobacteria: Gamma Subclass, 123-158.

Eng, S.K., Pusparajah, P., Ab Mutalib, N.S., Ser, H.L., Chan, K.G., and Lee, L.H. (2015) *Salmonella*:

a review on pathogenesis, epidemiology and antibiotic resistance. Front Life Sci 8: 284-293.

Ernst, R.K., Dombroski, D.M., and Merrick, J.M. (1990) Anaerobiosis, type 1 fimbriae, and growth

phase are factors that affect invasion of HEp-2 cells by *Salmonella* typhimurium. Infect Immun

58: 2014-2016.

Feschotte, C. (2008) Transposable elements and the evolution of regulatory networks. Nat Rev Genet

9: 397-405.

Finley, D. and Chau, V. (1991) Ubiquitination. Annu Rev Cell Dev Biol 7: 25-69.

Foley, S. L. and Lynne, A. M. (2008) Food animal-associated *Salmonella* challenges: pathogenicity

and antimicrobial resistance. J Anim Sci 86: E173-E187.

Galimand, M., Sabtcheva, S., Courvalin, P., and Lambert, T. (2005) Worldwide disseminated *armA*

aminoglycoside resistance methylase gene is borne by composite transposon Tn1548. Antimicrob

Agents Chemother 49: 2949-2953.

Harvey, R.R., Friedman, C.R., Crim, S.M., Judd, M., Barrett, K.A., Tolar, B., *et al*. (2017)

Epidemiology of *Salmonella enterica* serotype Dublin infections among humans, United States,

1968–2013. Emerg. Infect. Dis. 23:1493.

Hendry, T.A., Freed, L.L., Fader, D., Fenolio, D., Sutton, T.T., and Lopez, J.V. (2018) Ongoing

transposon-mediated genome reduction in the luminous bacterial symbionts of deep-sea ceratioid

anglerfishes. mBio 9: e01033-18.

Hoberg, E.P. and Brooks, D.R. (2008) A macroevolutionary mosaic: episodic host‐switching,

geographical colonization and diversification in complex host–parasite systems. J Biogeogr 35:

1533-50.

Hong, S., Rovira, A., Davies, P., Ahlstrom, C., Muellner, P., Rendahl, A., *et al*. (2016) Serotypes and

antimicrobial resistance in *Salmonella enterica* recovered from clinical samples from cattle and

Swine in Minnesota, 2006 to 2015. PloS One 11: e0168016.

Hufbauer, R.A. and Roderick, G.K. (2005) Microevolution in biological control: mechanisms,

patterns, and processes. Biol Control 35: 227-239.

Iqbal, Z., Caccamo, M., Turner, I., Flicek, P., and McVean, G. (2012) *De novo* assembly and

genotyping of variants using colored de Bruijn graphs. Nat Genet 44: 226-32.

Jennings, E., Thurston, T.L., and Holden, D.W. (2017) *Salmonella* SPI-2 type III secretion system

effectors: molecular mechanisms and physiological consequences. Cell Host Microbe 22:217-231.

Klemm, E.J., Gkrania-Klotsas, E., Hadfield, J., Forbester, J.L., Harris, S.R., Hale, C., *et al*. (2016)

Emergence of host-adapted *Salmonella* Enteritidis through rapid evolution in an

immunocompromised host. Nat Microbiol 1: 15023.

Knight, S.D. and Bouckaert, J. (2009) Structure, function, and assembly of type 1 fimbriae. In

Glycoscience and Microbial Adhesion (pp. 67-107). Springer, Berlin, Heidelberg.

Kovac, J., Cummings, K.J., Rodriguez-Rivera, L.D., Carroll, L.M., Thachil, A. and Wiedmann, M.

(2017). Temporal genomic phylogeny reconstruction indicates a geospatial transmission path of

Salmonella Cerro in the United States and a clade-specific loss of hydrogen sulfide production.

Front Microbiol 8: 737.

Kruskal, J.B. (1964) Nonmetric multidimensional scaling: A numerical method. Psychometrika 29:

115-129.

Kung, V.L., Khare, S., Stehlik, C., Bacon, E.M., Hughes, A.J., and Hauser, A.R. (2012) An *rhs* gene

of *Pseudomonas aeruginosa* encodes a virulence protein that activates the inflammasome. Proc

Nat Acad Sci USA 109: 1275-1280.

Langridge, G.C., Fookes, M., Connor, T.R., Feltwell, T., Feasey, N., Parsons, B.N., *et al*. (2015)

Patterns of genome evolution that have accompanied host adaptation in *Salmonella*. Proc Nat

Acad Sci USA 112: 863-868.

Lerat, E. and Ochman, H. (2005) Recognizing the pseudogenes in bacterial genomes. Nucleic Acids

Res 33: 3125-3132.

Li, Y., Pinto-Tomás, A.A., Rong, X., Cheng, K., Liu, M., and Huang, Y. (2019) Population genomics

insights into adaptive evolution and ecological differentiation in *Streptomycetes*. Appl Environ

Microbiol 85: e02555-18.

Liao, J., Orsi, R.H., Carroll, L., Kovac, J., Ou, H., Zhang, H., and Wiedmann, M. (2019) Serotype-

specific evolutionary patterns of antimicrobial-resistant *Salmonella enterica*. BMC Evol Biol 19:

132.

Lupolova, N., Dallman, T.J., Holden, N.J., and Gally, D.L. (2017) Patchy promiscuity: machine

learning applied to predict the host specificity of *Salmonella enterica* and *Escherichia coli*.

Microb Genom 3.

Lupolova, N., Dallman, T.J., Matthews, L., Bono, J.L., and Gally, D.L. (2016) Support vector

machine applied to predict the zoonotic potential of *E. coli* O157 cattle isolates. Proc Nat Acad

Sci USA 113: 11312-7.

Ma, K.W. and Ma, W. (2016) YopJ family effectors promote bacterial infection through a unique

acetyltransferase activity. Microbiol Mol Biol Rev 80: 1011-1027.

Maculins, T., Fiskin, E., Bhogaraju, and S., Dikic, I. (2016) Bacteria-host relationship: ubiquitin

ligases as weapons of invasion. Cell Res 26: 499.

Martin, J.E., Waters, L.S., Storz, G., and Imlay, J.A. (2015) The *Escherichia coli* small protein MntS

and exporter MntP optimize the intracellular concentration of manganese. PLoS Genet 11:

e1004977.

Maury, M.M., Tsai, Y.H., Charlier, C., Touchon, M., Chenal-Francisque, V., Leclercq, A., *et al.*

(2016) Uncovering *Listeria monocytogenes* hypervirulence by harnessing its biodiversity. Nature

Genet 48: 308.

McGhie, E.J., Brawn, L.C., Hume, P.J., Humphreys, D., and Koronakis, V. (2009) *Salmonella* takes

control: effector-driven manipulation of the host. Curr Opin Microbiol 12: 117-124.

McLaughlin, J.B., Castrodale, L.J., Gardner, M.J., Ahmed, R. and Gessner, B.D. (2006). Outbreak of

multidrug-resistant *Salmonella* Typhimurium associated with ground beef served at a school

potluck. J. Food Prot. 69: 666-670.

Mileykovskaya, E. and Dowhan, W. (2009) Cardiolipin membrane domains in prokaryotes and

eukaryotes. Biochim Biophys Acta 1788: 2084– 2091.

Miriagou, V., Carattoli, A., and Fanning, S. (2006) Antimicrobial resistance islands: resistance gene

clusters in *Salmonella* chromosome and plasmids. Microb Infect 8: 1923-1930.

Moran, N.A. and Plague, G.R. (2004) Genomic changes following host restriction in bacteria. Curr

Opin Genet Dev 14: 627-633.

Morgan, E., Campbell, J.D., Rowe, S.C., Bispham, J., Stevens, M.P., Bowen, A.J., *et al*. (2004)

Identification of host-specific colonization factors of *Salmonella enterica* serovar Typhimurium.

Mol Microbiol 54: 994-1010.

Moura, A., Criscuolo, A., Pouseele, H., Maury, M.M., Leclercq, A., Tarr, C., *et al*. (2017) Whole

genome-based population biology and epidemiological surveillance of Listeria monocytogenes.

Nat Microbiol 2: 16185.

Nemergut, D.R., Schmidt, S.K., Fukami, T., O'Neill, S.P., Bilinski, T.M., Stanish, L.F., *et al*. (2013)

Patterns and processes of microbial community assembly. Microbiol Mol Biol Rev 77: 342-356.

Olsen, S.J., Ying, M., Davis, M.F., Deasy, M., Holland, B., Iampietro, L., *et al*. (2004) Multidrug-

resistant *Salmonella* Typhimurium infection from milk contaminated after pasteurization. Emerg.

Infect. Dis. 10: 932.

Parkhill, J., Dougan, G., James, K.D., Thomson, N.R., Pickard, D., Wain, J., *et al.* (2001) Complete

genome sequence of a multiple drug resistant *Salmonella enterica* serovar Typhi CT18. Nature

413: 848-852.

Pasquali, F., Kehrenberg, C., Manfreda, G., and Schwarz, S. (2005) Physical linkage of Tn 3 and part

of Tn 1721 in a tetracycline and ampicillin resistance plasmid from *Salmonella* Typhimurium.

Antimicrob Chemother 55: 562-565.

Peleg, A.Y., Miyakis, S., Ward, D.V., Earl, A.M., Rubio, A., Cameron, D.R., *et al.* (2012) Whole

genome characterization of the mechanisms of daptomycin resistance in clinical and laboratory

derived isolates of *Staphylococcus aureus*. PloS One 7: e28316.

Petersen, L., Bollback, J.P., Dimmic, M., Hubisz, M., and Nielsen, R. (2007) Genes under positive

selection in *Escherichia coli*. Genome Res 17: 1336-1343.

Pezzella, C., Ricci, A., DiGiannatale, E., Luzzi, I., and Carattoli, A. (2004) Tetracycline and

streptomycin resistance genes, transposons, and plasmids in *Salmonella enterica* isolates from

animals in Italy. Antimicrob Agents Chemother 48: 903-908.

Plumb, I.D., Schwensohn, C.A., Gieraltowski, L., Tecle, S., Schneider, Z.D., Freiman, J., *et al.*

(2019) Outbreak of *Salmonella* Newport infections with decreased susceptibility to azithromycin

linked to beef obtained in the United States and soft cheese obtained in Mexico—United States,

2018–2019. Morbidity and Mortality Weekly Report 68:713.

Quezada, C.M., Hicks, S.W., Galán, J.E., and Stebbins, C.E. (2009) A family of *Salmonella*

virulence factors functions as a distinct class of autoregulated E3 ubiquitin ligases. Proc Nat Acad

Sci USA 106: 4864-4869.

Rice, P.A. and Baker, T.A. (2001) Comparative architecture of transposase and integrase complexes.

Nat Struct Biol 8: 302-307.

Richards, V.P., Lang, P., Bitar, P.D., Lefébure, T., Schukken, Y.H., Zadoks, R.N., *et al*. (2011)

Comparative genomics and the role of lateral gene transfer in the evolution of bovine adapted

Streptococcus agalactiae. Infect Genet Evol 11: 1263-75.

Rodriguez-Rivera, L.D., Cummings, K.J., Loneragan, G.H., Rankin, S.C., Hanson, D.L., Leone,

W.M., *et al*. (2016) *Salmonella* prevalence and antimicrobial susceptibility among dairy farm

environmental samples collected in Texas. Foodborne Pathog Dis 13: 205-11.

Rytkönen, A., Poh, J., Garmendia, J., Boyle, C., Thompson, A., Liu, M., *et al.* (2007) SseL, a

Salmonella deubiquitinase required for macrophage killing and virulence. Proc Nat Acad Sci USA

104: 3502-3507.

Schleker, S., Sun, J., Raghavan, B., Srnec, M., Müller, N., Koepfinger, M., *et al*. (2012) The current

Salmonella☐host interactome. Proteomics Clin Appl 6: 117-33.

Slatkin, M. (1987) Gene flow and the geographic structure of natural populations. Science 236: 787-

92.

Strachan, N.J., Rotariu, O., Lopes, B., MacRae, M., Fairley, S., Laing, C., *et al*. (2015) Whole

genome sequencing demonstrates that geographic variation of *Escherichia coli* O157 genotypes

dominates host association. Sci Rep 5: 14145.

Steenbergen, J.N., Mohr, J.F., and Thorne, G.M. (2009) Effects of daptomycin in combination with

other antimicrobial agents: a review of in vitro and animal model studies. J Antimicrob Chemother

64: 1130-1138.

Toft, C. and Andersson, S.G. (2010) Evolutionary microbial genomics: insights into bacterial host

adaptation. Nat Rev Genet 11: 465.

Thomson, N.R., Clayton, D.J., Windhorst, D., Vernikos, G., Davidson, S., Churcher, C., *et al*. (2008)

Comparative genome analysis of *Salmonella* Enteritidis PT4 and *Salmonella* Gallinarum 287/91

provides insights into evolutionary and host adaptation pathways. Genome Res 18: 1624-37.

United States Department of Agriculture. (2014) Nutrient Management Practices on U.S. Dairy

Operations.

Valenski, M.L., Harris, S.L., Spears, P.A., Horton, J.R., and Orndorff, P.E. (2003) The Product of the

fimI gene is necessary for *Escherichia coli* type 1 pilus biosynthesis. J Bacteriol 185: 5007-5011.

Vos, M. (2011) A species concept for bacteria based on adaptive divergence. Trends Microbiol 19: 1-

7.

Wachter, J. and Hill, S. (2016) Positive selection pressure drives variation on the surface-exposed

variable proteins of the pathogenic *Neisseria*. PloS One.11: e0161348.

Yue, M., Han, X., De Masi, L., Zhu, C., Ma, X., Zhang, J., *et al*. (2015) Allelic variation contributes

to bacterial host specificity. Nat Commun 6: 8754.

Zhao, S., Qaiyumi, S., Friedman, S., Singh, R., Foley, S.L., White, D.G, *et al*. (2003)

Characterization of *Salmonella enterica* serotype Newport isolated from humans and food

animals. J Clin Microbiol 41: 5366-71.

Zheng, J., Luo, Y., Reed, E., Bell, R., Brown, E.W., and Hoffmann, M. (2017) Whole-genome

comparative analysis of *Salmonella enterica* serovar Newport strains reveals lineage-specific

divergence. Genome Biol Evol 9: 1047-1050.