Can constraint closure provide a generalized understanding of community dynamics in ecosystems?

Steven L. Peck[1]*, Andrew Heiss[2]

1 Biology Dept, Brigham Young University, Provo, UT, USA
2 Department of Public Management and Policy/Andrew Young School of Policy Studies, Georgia State University, Atlanta, Georgia, USA

* Steven_Peck@byu.edu

Abstract

Understanding causal complexity in ecological communities has been a challenge to ecologists since the inception of the discipline. Here we draw insights from recent work on constraint closure that suggests ways of grappling with ecological complexity and that yield generalizable theoretical insights. Using a set of evolutionary constraints on species flow in ecological communities, including selection, species drift, dispersal, and speciation, combined with multispecies interactions such as mutualistic interactions, and abiotic constraints, we show how theoretical work drawing on constraint closure allows the possibly of the emergence of communities as semi-autonomous structures. Through agent-based modeling we articulate ways ecological pattern formation viewed through the lens of constraint closure, which inform questions about stability and turnover in community ecology.

Introduction

Ecological theorists have generated a number of yet unresolved disputes that try to untangle the difficulty of understanding the nature of complex ecological communities–including the question of whether the idea of a community in such systems is coherent. For example, questions about ecological community structure and how it is maintained have been argued about for over a hundred years. These questions are still taking their current principle divisions from camps first staked out by Clements [1] and Gleason [2] in the early decades of the 20th Century: Are communities assembled non-randomly, conditioned on certain possibilities of co-occurrence and structured by general ecological regularities, or are they more randomly constructed entities brought together by a complex combination of abiotic factors, available species pools, competition, and other ecological processes that suggest a strong element of contingency [3, 4]?

Currently, there does not seem to be a unified body of theory and practice that explains most features of community structure. The hunt for generality has also been fraught (See (Elliott-Graves Manuscript in Review) for a detailed overview). Theories in the discipline often seem a piecemeal confederation of contingent ideas that carve nature mid-bone as it were, often focused on specific natural systems that provided only limited empirical and theoretical support among other differently structured ecological communities. It is clear that the theoretical constructions used by ecologists have been crucially instrumental in the development of research programs and have allowed

progress in understanding community ecology. However, even basic concepts in the discipline are often viewed differently by different research groups that are often divided into different camps, often to the point that there are even debates about the usefulness of the concept of community ecology. For example, the following questions are still being debated. Are they individuals [5–14]? Which ecological theories best describe the dynamics of their structure, maintenance, and development [8,9,11,13,14]? Do they have boundaries [5,15]? Are they stable [16]? How are communities assembled, and are they so contingent that each community must be examined as an individual case study [17–23]? Are ecosystems even real entities [24,25]?

These questions are all interrelated, yet the abundant theories used to tame and understand these ecological systems rarely provide a unified, or coherent, picture among theories. Moreover, in order to make progress in the idea of a ecological community it seems important to discover generalities that hold across the numerous possible biotic configurations that comprise these individual community assemblages. It is also important to view this project from multiple perspectives, in order to tackle the inherent complexity, and provide a unified, or coherent, picture among theories for a more unified conception of community ecology. There have been a number of attempts to do so, and the importance of these efforts has been recognized, [9,11,14,20,26–31], though conditioned on some skepticism about the possibility of finding well supported law-like regularities in community ecology [19,32–34].

The Piecemeal Nature of Ecological Theories

Almost from the inception of ecology as a discipline, piecemeal theories emerge that propose ways that one can get a handle on Ecology's untamed complexity. Table 1 provides a sampling of some of these theories. Note that the structure of these theories is such that each takes a particular, narrowly focused set of ecosystem patterns and processes from which it attempts to reduce complexity by simplifying the ecosystem to a few aspects extracted from the ecosystem as a whole.

For example, food web theory looks at interactions among species that feed in trophic hierarchies; whereas metapopulation theory considers movement among organisms among small populations scattered in a matrix of uninhabitable or poor quality habitat; and neutral theory examines community construction from the perspective of available species pools exploiting contingent opportunities to commandeer niche openings. Many other such ecological theories could be included.

Ecological Frameworks

This abstraction and idealization of common ecological processes are theories in the sense that they are a collection of models (as semantic view of theory construction [24]), that work well with Scheiner and Willigs articulation of theories in ecology [35]. However, as theories in the more traditional sense of the word, they lack a grounding generalization that would inspire more confidence that these are theories qua theories. Most draw only lightly on defining the domain over which they hold sway. For example, in metapopulation theory, organisms that are capable of colonization and extinction within small habitat patches are the pattern generators of interest, but these are not specifically defined except as possible parameterizations of real organisms (including movement rates, extinction probabilities, etc.). Specifics, however, as to what exactly and clearly constitute a set of metapopulations has not been laid out in any detail except to accept that any organism that looks metapopulationish in terms of its structure (e.g., patchy, connected, populations that undergo movement among patches), is allowed in.

Table 1. Sampling of Ecological Theories.

Network Theory [36, 37]
Food webs [38, 39]
Metapopulation Theory [?, 40]
Niches and Niche Construction [41–43]
Island Biogeography [?, 44]
Hubbell's Unified Neutral Theory of Biodiversity and Biogeography [45, 46]

Pragmatically, it should be noted that these constructions work well enough to make these frameworks useful in providing grounding for experimental design and data analysis, but as formal theories, they come up wanting. This may be why community ecology has seemed so elusive. The piecemeal nature of its theories, the lack of formal attributes that allow a clear and complete articulation of a domain, and aspects of ecological theory have all made the discipline poorly supported theoretically compared to other special sciences. This is not to argue that they are not useful, but rather they seem more useful in the sense of general frameworks than as formal theories. That may be why these tend to proliferate with a new one arriving and gaining popularity every few years. These frameworks do much of the work of theories, they organize thought by providing a structure to inform experimental design, data gathering and analysis, and provide predictions that set up expectations and interpretations by using various kinds of baseline models.

However, integrating these frameworks, given the complexity of ecosystems, has been problematic. In part, because we want a generalized structure of an ecological community to contain all of these: food webs, niches, metapopulation, contingent openings in niche resources, and the other framework-based articulations of ecosystem components and processes. All of these frameworks have been successful in providing information about certain aspects of community structure and provide insights into the nature of these systems that allow them to be better understood, analyzed, and predictions made based on the assumptions and relationships captured in the constituent models. However, an overarching and general articulation of ecosystem communities has been hard to come by. This is important because although we want to acknowledge that reductive techniques have done much to open our understanding to some of aspects of ecosystem science, to understand ecosystems as ecosystems we are going to have to include their complexity as complexity if they are to be understood holistically rather than in piecemeal fashion [47].

Integrating Ecological Frameworks

There have been several attempts to integrate community dynamics. By way of illustration, one proposal to capture this generality suggests that ecosystems might best be handled as indexically structured systems. In an indexically structured system, a particular group of organisms is chosen (often based on conservation need or other management goals) as the index species and its relationships with other components of the abiotic and biotic community are identified and articulated. If multiple species are considered together, this framework creates a network of relationships constructed by

overlapping all of the connections among all species present in an ecosystem, in theory giving a sense of the entire community's interactions. By noting how indexically structured organisms allow stabilities to form among the constituent species, a more complete view of a suitably bounded ecosystem could be constructed. A practical problem with such an approach is that given the amount of work required of an ecologist to do for even just a single pair of species relationships to be fully resolved is nearly impossible in practice due to expense, ecosystems disruption, and other logistical and ethical considerations it is unlikely that the full potential of this idea will be implemented.

In short, we are confronted with the problems of studying highly complex systems in which regularities are tamed by conceptual frameworks that while doing real work in establishing structure from which to view ecosystem processes, nevertheless work piecemeal and do not generalize among differing frameworks. This is not intended as a criticism of these frameworks, and we recognize that it may be how ecological regularities are best handled given the limitations of field work. Still, within the discipline of community ecology, researchers desire strongly to forge a more general theory.

Two recent approaches used together suggest a promising way to think more generally about ecosystem dynamics in community ecology and offer potential for a more generalized theory. First, consider the recent call by Vellend (2016) [14]to focus on a small set of processes that provide a base structure for ecosystem theory, similar to the way that theoretical population genetics has been fashioned to focus on a small set of processes that provide a base structure for ecosystem theory similar to the way that theoretical population genetics has been fashioned; and second, by exploring the role of functional constraints in defining autonomous and semi-autonomous systems, of which structured ecological communities may, arguably, be considered [7, 48–52].

Generalizing Ecological Community Theory

Mark Vellend proposed we look at ecological communities in light of four processes found in a wide variety of ecosystems. These activities are analogous to those found in basic population genetics: selection, species drift, dispersal, and speciation [14, 53]. These are chosen to mirror processes found in basic population genetics: selection, genetic drift, dispersal and mutation. These chosen community level processes are thought to capture the means whereby most communities change. Vellend argues that population genetics has made significant advances in understanding how genetic concerns play out in populations by focusing on these activities. By focusing on these fundamental aspects, population genetics has been able to make significant theoretical inroads in understanding how genes in populations are structured and behave. Can community ecology likewise benefit by focusing on a similar set of ecosystem characteristics?

Vellend (2016) [14] suggests that this subset of possible processes covers much of the conceptual territory being explored in community ecology. Additional suggestions may bear on his insights by considering communities as autonomous structures conditioned on the constraints imposed by selection, species drift, dispersal, and speciation; three additional forces may be added: mutualisms, abiotic boundaries, and patch structure and dynamics. This set seems to capture a general and well-studied set of constraints in ecosystem communities. Focusing on constraints may be an especially useful approach for ecological generalization as it provides ways to theoretically consider general outcome characteristics like stability, expected species turnover rates, and other features of community dynamics like those noted by Gleason and Clements about how communities are assembled and maintained. How constraints allow the emergence of autonomous systems, which we turn to next, will suggest how these ideas can work

together to inform ecology.

Autonomous Systems

One way to approach a general idea of ecosystem functioning can be drawn from Moreno and Mossio's (2015) [51] work on biologically autonomous systems. Even through the question of whether ecological communities are autonomous entities is empirically unsettled, they identify three dimensions of general biological autonomy.

The first dimension of biological autonomy is the capacity for self-determination, meaning that the grounding, normativeity, teleology, and functionality come from the system itself. For this to occur, a closure of constituent constraints is required. Fig 1 illustrates a minimal set of requirements for the occurrence of constraint closure such that components modulate the flow of energy and materials within the system. These provide a sustainable feedback loop which allows the system to achieve coherence for a period of time, τ_1.

Fig 1. Constraint Closure. Constraint closure is described as constraints on the flow on matter or energy that form a closed loop. This is a necessary step in the formation of an autonomous system. These provide a sustainable feedback loop which allows the system to achieve coherence for a period of time, τ_1. Redrawn from Figure 1.3, p. 21, Moreno and Mossio's (2015) [51]

The second dimension is found in a community's interaction with the larger environment, which provides the energy and materials used by the ecosystem. It can also be the source of disturbances that might restructure the system through things like direct human influence, invasive species, diseases, or abiotic influence such as fire, weather-related events, and other non-biological influences.

The historical aspects of the system, which include past events and disturbances, ecological influences on evolutionarily forces, contingent factors, co-evolutionary adaptations, and other biotic and abiotic influences, frame the third dimension of biological autonomy.

Given the above, there are reasons to think that ecological communities might be autonomous systems. Energy flow within the system is maintained and channeled within the system itself by its components. For example, long-term community stability has been shown to be structured by a number of well-studied ecosystem processes, e.g., competitive exclusion of species maintaining particular niches, networks of mutualistic relations, typical ecological interactions like predator prey relations, etc.

Likewise, there are also reasons to think they are not. Do the contingent aspects of ecologies affect the possibility of their being autonomous? Many of the structures we see in ecosystems appear to be arbitrary members of a set derived from a contingent species pool, what is autonomous in such a system? Is it, perhaps, the fungible functional role that particular species plays in an ecosystem and that subsequently structure the patterns and processes we observe? Or does it depend on long-term coevolutionary relationships among particular species? Given the debates between ecologists since Clement and Gleason, it seems as if both may be important and therefore some generalities in the form of regularities that span different kinds of ecosystems may qualify these as autonomous systems at least for some interval τ. One particular problem is assigning boundaries to ecological communities. This seems to be an important step in exploring these systems as autonomous systems.

One way to approach this is through work done on clarifying the notion of individuality [54]. As Griesemer notes this is a fraught concept [55], "I think we learn more about the concepts, claims, and consequences of individuality in the sciences by

supposing it is a theoretical term that can mean different, even conflicting, things in different theoretical contexts, such as in different scientific specialties and disciplines with different research questions, projects, and phenomena to consider, rather than a logical or metaphysical term that that should shave a single, best, all-purpose fixed meaning" p.137. One useful perspective on individuality in the context of community ecology is Roberta Millstein's articulation of "Land Community" as envisioned by Aldo Leopold [56, 57].

A land community combines aspects of community ecology and ecosystem ecology that emphasizes both energy flow though the biotic and abiotic components of the system and the relationships among organisms and their abiotic context . Both of these are important in thinking about these as autonomous systems and provide a framework for providing a practical conception of boundaries. Millstein defines a Leopoldean land community as consisting of, "populations of different species interacting with each other and with their abiotic environment; these survival-relevant interactions often produce a flow of energy and materials between biotic components and between biotic components and abiotic components (and vice versa)."

She then articulates a use notion of boundaries for these systems, "Land community boundaries for well-bounded systems are where discontinuities or steep gradients in the flow of material an energy coincide with discontinuities or steep gradients in species interactions. Land Community boundaries for open systems are at a minimum delineated by the smaller of the two types of discontinuities or steep gradients, including the more extensive interactions or matter/energy flows if and only if those interactions or matter/energy flows are stronger or larger than those of the smaller area."

Lastly she sets up three criteria for a land community to be considered an individual that we will uses as a bases to explore such as autonomous systems. They are, (1) spatially and temporally restricted; (2) integrated by causal interactions which produce a shard fate; (3) are temporally bracketed with a beginning and an end; and (4) are continuous in time allowing for change. We read the fourth as considering these as process-based. These set the stage for what further is required to consider these as autonomous and in all that follows by community we mean a Land community as described by Millstein.

Nunes-Neto, et al. [52] make a compelling argument that ecological communities can be seen as autonomous systems because of the functional components that structure most ecosystems. In particular, they look at the function of the components of biodiversity using what is termed the 'organizational approach' (OA).

Following Mossio, Saborido, and Moreno [58], they formally set up the following for definition of biological function, "A trait T has a function in the organization O of a system S if and only if:

C1 : T contributes to the maintenance of the organization O of S;

C2 : T is produced and maintained under some constraints exerted by O;

C3 : S is organizationally differentiated. [58] p. 828

This is contrasted with the Etiological Approach (EA), which uses evolutionary history to define function as doing that which it was selected to do in an adaptionist sense. OA then considers only what the function does in the current ecosystem, not as it is embedded in its historical or evolutionary framing. This, drawing on the work of Moreno and Mossio [51] on constraint closure, leads to an ecological definition of function, by Nunes et al. "An ecological function is a precise (differentiated) effect of a given constraining action on the flow of matter and energy (process) performed by a given item of biodiversity, in an ecosystem closure of constraints" [52].(p. 131)

This provides a handle into seeing land communities as autonomous systems, as maintained by constraint closure. The constraints are modulations induced by the components of biodiversity acting on the flow of matter and energy through the system,

in a way that feeds back into the maintenance of the ecosystem. It is important to point out that the claim of its being an autonomous system is not the same as the claim that they are individuals in the biological sense, although they do seem to exhibit weak individuality [49, 52].

Dussault and Bouchard [48] add a significant consideration to the above account, developing a focus on particular kinds of functions which they call Persistence Enhancing Functions (PEP). PEPs are a class of functions that allow the currently structured ecosystem to persist. The focus is not upon these functions in an adaptive sense, such that their historical grounding has been necessarily worked out. Rather it is future-directed such that it attends to those extant functions that contributed to current stability–defined as the ability to return to equilibrium after a disturbance. They define a PEP function as, "The ¡¡PEP¿¿ function of x in an ecosystem E is to F if, and only if, x is capable of doing F and x's capacity to F contributes to E's propensity to persist." [48](p. 8)

PEPs then allow us to explore the question, what features provide for the ability of this ecosystem, envisioned as a Leopold land community, to to survive disturbances to equilibrium. It pulls away from questions about how particular assemblages come to be constructed and looks for general characteristics of ecosystem functioning and construction that are based upon the roles that species play in the current community.

These two proposals (OA and PEP) offer possible constraints within ecological communities that influence stability and structure and that allow them to be considered autonomous systems. Are there other constraints that might influence a generalized theory of community dynamics?

Vellend's Set of Processes as Constraints

By combining the above ideas about how functions contribute to ecosystem structure and autonomy with Vellend's proposal, we can see a possible space open with which to unite these views. The processes on which Vellend focuses could also be considered a set constraints that are found in nearly all ecosystems, and which are known to structure the 'flow' of species through an ecosystem. In nearly all ecosystems, his proposed focus, i.e., selection, species drift, speciation, and dispersal form a set of constraints on the flow of species through the ecosystem. Since the constraints that Moreno and Mossio discuss are not limited to the kinds of systems typically seen in matter and energy flow (e.g., chemical or cellular systems, and other micro processes), any set of constraints on a given process, including the flow of species through an ecosystem, would seem to be a candidate for the emergence of autonomy in the sense they prefigure. Known to affect the modulation of the flow of species through spatially and temporally bound ecosystems, Vellend's set of proposed processes provides a possible set of constraints. Selection constrains the success of a species by affecting its ability to compete in its current abiotic and biotic situation; providing fitness conditions on a species' success or failure in surviving in an area. Species drift constrains species genetic architecture and stochastic variation in how stable a species is in a bounded region and how it drifts in adjusting or modifying its current niche conditions. Speciation, in addition to the effects of selection, constrains how many new species emerge in an area due to genetic factors, and the interplay between trait differences and the biological fitness of those differences. Likewise, dispersal constrains species flow by influencing a combination of a species' ability to spread beyond its original location, for instance in the sense that geneticist Sewall Wright proposed in his three-step shifting balance theory, i.e., (1) Initial Establishment; (2) Population Growth; and (3) Species dispersal of individuals to new areas [59]. This forms a hypothesis about how constraints influence ecosystem function. Constrains on species flow through the system may set up such communities as

autonomous systems through constraint closure analogous to the way Moreno and Mossio [51] envisioned how autonomous systems work and, in particular, frame a set of PEP functions that influence stability and other aspects in community dynamics. There may be other possible constraints on species flow. For example, things like boundaries over which species cannot cross, may affect which species enter into a particular community; the spatial patch dynamics of particular landscapes such that metapopulation structure creates a constraint on species flow; and mutualistic relationships between species within an ecosystem, which affects survival probabilities of cooperating species.

A Model of Constraint Closure in Autonomous Ecological Communities

A word about modeling in theory construction

Representing an ecological community in a model presents several challenges. Modeling such systems has a rich history, and three types of models have dominated: Analytic, those largely represented by mathematical systems of equations (called analytic even if computation must be used to solve the equation using numerical methods); Simulation or computational models, those represented by algorithms instantiating procedures, rules, and/or manipulating abstract objects or data structures representing individuals or agents; or statistical models that look for relationships among data objects using probability theory to structure inferences and predictions from the data. All three of these are used regularly in community ecology and serve different purposes.

In all of these types of models what is being represented varies over the uses the developer wants target with her models. For example, the target systems my be a specific community with specified entities and for which the modeler hopes to make specific claims or make targeted predictions about this particular system. On the other end of the spectrum, the target might be a generalized ideation of a hypothesized structure. For example in early in the history of population genetics, long before the nature of DNA was worked out, much progress was made modeling the properties that genes must necessarily possess to do the work proposed for what it meant to be a gene.

Given the complexity of ecological systems, simulation seems the method best equipped to handle the complexity of these systems. Simulation is best viewed as a hermeneutical practice [60]. Overview challenges of simulation models. Scenario construction and fictional use [47].

Pattern recognition has played a pivotal role in ecology almost since its inception [61], but especially since Levin's influential paper laying out the important role in ecosystems of spatial patterns and scale [62]. Since that time, ecological researchers have focused on an number of issues related to pattern formation both temporal and spatial scales, including: Scales of interaction in both in trophic relationships; modularity of ecosystems function; patterns of diversity; assemblage rules; and other influences that are dependent on differential scales of influence and interaction [63]. This leads to a number modeling problems orbiting questions about how to best represent these systems, especially in regard to the use of null models. or in trying to sort out assemblage rules, which empirically have proven to be complex and difficult to accomplish adequately [4], making modeling even more challenging. Especially if, as argued above, these have emergent properties and patterns. Modeling emergent phenomena require that we take into account the diachronic aspects of the system which allow the emergence and modeling of individual interactions. This is most easily done using agent-based models. They may be an important way of drawing out

the connection between the patterns we find in nature and those generated by our idealized [64], p. 150.

Grimm and Railsback [65] have offered a rationale and strategy for recognizing and modeling patterns in complex ecological systems, which they call 'Pattern-oriented modeling' (POM). This lends itself well to agent-based models and for exploring the complexity inherent in ecological communities and identifying the right levels and processes to include in the model. They suggest the following set of practices be used after empirically identifying the patterns:

"(i) determine what scales, entities, variables and processes the model needs, (ii) test and select submodels to represent key low-level processes such as adaptive behaviour, and (iii) find useful parameter values during calibration." p. 298.

For example, consider the problem of exploring Vellend's proposal in light of the work on constraint closure. To examine this, we will use mycorrhizal networks as a model system to explore these question.

The mycorrhizal networks that form mutualistic relationships with plants may be a good target system to explore ideas about constraints and stability in communities [66, 67]. Soil Mycrorhizal fungi form symbiotic relationships with tree roots and establish a chemical communication system among a diversity of ecosystem constituents [68]. These networks can enhance plant growth by increasing nutrient availability by facilitating their release from rocky substrate and helping break down organic carbon sources [69]. They also may help improve soil quality by modifying heavy metals and other inorganic pollutants [70, 71]. They have been shown to also affect interactions among competing plants, invasive species, and can have a large impact on community composition [72–75]. This is because these fungal communities can indirectly change the relationships among competing plants by providing resources to a broader spectrum of plants thereby increasing above ground niche breadth [76].

Below we use mycorrhizal networks as a model system to test how the combination of Vellend's suggestions about using a base set of processes as the building blocks for developing a generalized theory of community interactions in ecological systems, and ideas of constraint closure may make some of the patterns found in such communities interpretable. The following is a generalized model without a specific target other than the kind of interactions found in the broad class of forest/mycorrhizal systems. This model system would lend itself to exploring nested mutualistic structures, but it is not included in this paper [43].To put the following in terms of a model mapping schema provided by Weisberg (2013) [77] the model below is a generalized, computational model, using a minimalist idealization of key processes thought to potentially function in ecological communities, including Vellend's [14] proposed set of high-level processes for conceptually framing ecological communities, i.e., selection, species drift, speciation, and dispersal. In addition, the model considers the kinds of constraints thought to capture the dynamics proposed by Moreno and Mossio [51]. These two ideas should allow community stability as recognized by signature patterns to emerge. The fidelity criteria we aim for will be achieved if such patterns are present and suggest that these processes are sufficient to confer known emergent attributes on these model systems. We make no claim that this model will actually represent any specific target system, but will give in principle insight into places for further testable hypotheses about how these ecosystems function.

To explore this possibility we have developed an agent-based model that explores the potential of using these ideas to gesture to the possibility that the ideas of constraint closure can be a way to structure general insights into theoretical ecological community ecology.

Methods and Model Description

The model is designed to explore how constraints in certain circumstances might provide long-term stability in ecological communities.

ODD

We will use the "Overview, Design Concepts, and Details" (ODD) protocol recommended by Grimm et. al. 2010 [78] to describe the model. This protocol has proven to be an important way to describe computer simulation studies and in particular agent based models such as this one. This protocol facilitates understanding how the model has been designed, what assumptions the authors are making, and how various components of such models such as spatial scale, time, stochasticity, and other choices for structuring the model are implemented.

Purpose

The following model has been designed to provide in-principal arguments for how ecological communities maintain stability even in the presence of stochastic turnover. Using Vellend's [14] proposed set of high-level processes for conceptually framing ecological communities, i.e., selection, species drift, speciation, and dispersal, we hypothesize that these provide constraint closure in the way proposed by Moreno and Mossio [51] that contribute to biological autonomy. Moreover we hypothesize specifically that these will allow stable ecological signatures to arise and be maintained through constraint closure. In this way we seek to exemplify the possibility of creating some measure of generalizability in ecosystem community structure.

Entities, state variables, and scales

Table 2 gives a list of the parameters in the model. The model is structured on a spatial toroidal grid composed of NxN patches that form clusters of suitable and unsuitable areas. Fig 2 gives an example of one of the landscapes produced. This procedure is based upon xxxx.

The unsuitable habitat patches form a barrier, or impediment, to migration. Each habitat patch represents a packed set of niches that can hold M species. Each patch contains an environmental niche score, a variable representing a habitat suitability index chosen from a uniform random distribution between 0 and 1. Each species has a similar niche score, which represents a one-dimensional niche measure of the species needs. This corresponds to the patch environmental niche score, which in turn is a measure of what needs that habitat will provide. A species' fitness will be defined as the absolute difference between a species' niche score and its patch niche score.

Fig 2. Model Patches and Paramters The model is written in NetLogo an Agent Based Programing Environment. The variables and process control of the following model is detailed below.

Some species are able to form mutualisms with other species and a directed network is established between $k <= M$ species. The fitness of a linked species, differs from unlinked species and is calculated as a given function of linked species in a given patch, e.g., the maximum or average fitness of the linked species.

Each species also plays a functional role in a patch. This is specified by sp_ν in which $\nu \in \{1 \ldots sp_{max}\}$ and sp_{max} is the maximum number of functional roles in each patch's niches. The functional roles also determine a "link designation" which picks out which

Table 2. Table of Principal Parameters Used in the Model.

Parameter	Name (role)	Range	Default Value
T_{max}	Maximum number of time steps	$\in \{1, 20,000\}$	500
S_c	Species Selection (constraint)	Binary Switch	On
D_c	Species Dispersal (constraint)	Binary Switch	On
C_c	Species Competition (constraint)	Binary Switch	On
Ω_c	Speciation (constraint)	Binary Switch	On
P_c	Perturbation (constraint)	Binary Switch	On
Ψ	Proportion Species which form mutualisms	$\in \{0, 1\}$	0.5
ν	variance in environmental needs (used in fitness)	$\in \{0, 1\}$	0.05
Φ	Frequency of Perturbation	$\in \{0, T_{max}\}$	500
Σ	Proportion of species wiped out in perturbation	$\in \{0, 1\}$	0.2
λ	Landscape Density Factor	$\in \{0, 1\}$	0.25
Λ	Patch Distribution Factor	$\in \{0, 20\}$	5
ξ	Maximum Number of Species per Patch	$\in \{1, 50\}$	10
κ	Number of Mutualisms	$\in \{1, \xi_{max}\}$	5

Table notes: All parameters are coded either integers or random variables

species can form links, i.e., $sp_{numlink}$, designates which species can have links, and which cannot, such that if $sp_\nu \leq numlink$ links can be formed among other species that can also form links. 438 439 440

The specific constraints thought to allow the creation of stable communities are (a) Selection; (b) Dispersal; (c) mutualism (with any eye to mycorrhizal symbiotic relationships); (d) Spatial Competition; and (e) Fitness Competition between linked and unlinked species. In addition there are two assumed constraints that are structural parts of the model and are not tested for effect in this stage of the model analysis but may be explored at a latter time. (a) niche packing, i.e., a finite niche space; and (b) spatial structure limiting dispersal among species. 441 442 443 444 445 446 447

Process overview and scheduling 448

The overall flow of the model is given in fig 3.

Fig 3. Process Flow Within Model. The agent-based model used in this paper cycles through the evolutionary components of Vellend's proposed evolutionary structures in ecological systems (described above). These structures are used as constraints to create semi-autonomous structures. The flow chart shows the order of events and possible flow paths possible in the model.

After establishing the grid (details below in initialization section), M species are created in each patch i. A given number K, are able to form mutualistic links, the rest, $M - K$, are not. Each time step, τ, represents the mean time a species is likely to exist in a community. At time, τ_0, k mutualistic links are established within the patch for the initial population of species in each patch where $k = K\rho$. K is the maximum number of links that can be established, and ρ is a random variable that gives the proportion of potential links that do get established for a given K. The following processes procedures are given in the order they are executed: 450 451 452 453 454 455 456 457

- Select: Fitness is calculated based on the absolute distance of a given species' niche score to the patch's environmental niche score. For linked species the fitness is the highest fitness among those species with which it is linked. The species with the lowest fitness on each patch is eliminated. 458 459 460 461

- Dispersal: The species with the very highest fitness spreads to one of its neighbors if it has an opening, i.e., one of its neighbor patches has fewer than M species. If it is a species that can form links in its new patch it will do so.

- Fitness competition: One of the linkable species and one of the unlinkable species is randomly selected within each patch. The one with the higher fitness reproduces, the one with the lower fitness goes extinct in the patch. The new species will have a niche score with the same mean fitness requirement as its parent's fitness value, but with variance σ.

- Species drift: Whenever there are $G <= M$ openings in the niches within a patch, G new species are created from the G species with the highest fitnesses. The new species will have a niche score with the same mean fitness as its parent's fitness value, but with variance σ. There will also be the possibility the species drifting to a new functional role. The drift can occur across the link designator boundary $sp_{numlink}$, i.e., species can go from a linkable species to unlinkable species and visa versa, but if it drifts such that $\nu < 1$ or $\nu < maxsp$ it reflects back to the numlink boundary without changing its link designation, i.e., If sp_1 where about to drift to the non-existent sp_0, then $sp_1 \implies sp_{numlink-1}$ and likewise if sp_{maxsp} was going to drift out of the range above $maxsp$, then $sp_{maxsp} \implies sp_{numlink+1}$ such that it remains in its current link designation.

- Catastrophes: At given intervals, $I \in 1, t_{max}$ a proportion, ϕ, of the M species in each patch can be randomly wiped out. These perturbations provide a test of stability in terms of how the patch recovers from its loss of species.

Design Concepts

Emergence: We expect stability to emerge from certain combinations of constraints which form a closed loop of interaction and create constraint closure among the ecological processes described above.

- Objectives: We run the model under a number of different parameter combinations to understand how stability is structured.

- Interactions: Among species-agents in the model, interactions can be positive among those forming mutualistic associations, and negative among those that compete.

- Stochasticity: Stochasticity enters into the model in several places. (a) The initial patch structure of the landscape is randomly generated; (b) The proportion of species chosen to actually form links, ρ is a random variable; (c) the species that go extinct in the drift stage are randomly chosen with probability proportional to fitness; species with high fitness chosen to move, move randomly to one of their 8-Moore-neighborhood neighbors; (d) The variation of fitness around newly spectated species is a random variable v; and initially the environmental variable and the species environmental needs variable are randomly chosen; and (e) Species eliminated in a catastrophe are randomly chosen.

Variables of Interest

- *Turnover rate*: The turn over rate is the number of new species created at each time step. The way the model is formulated, new species arise when there is an opening in relation to the packed niche. This happens in competition between the linked and unlinked species, in selection events in which the least fit species is selected for elimination, and in perturbation events.

- *Evenness*: To capture patch evenness the Pielou's evenness index was modified for algorithmic efficiency. Rather than the mathematical equation $J' = \dfrac{H'}{H'_{\max}}$, , where H' is Shannon's Index, $H' = -\sum_{i=1}^{R} \dfrac{p_i}{ln p_I}$ we modify it as follows. Because the number of species per patch is fixed, the algorithem used is computed by the number of unique species in a patch divided by the number of species in the patch. This has the range, $\left(\dfrac{1}{sp_{max}}, 1\right)$ of where the lower range is found in the case where every species in the patch is unique, and the higher, when there is only one species.

- *Landscape Fitness* Fitness is measured as . . .

Model Structure and Testing

- Initialization: Initialization consists of a random landscape being produced using two variables that control the number of patches selected and the distribution of patches.

- Submodels: The only submodels are those described in the Process Overview section.

- Testing: The model Behavior Space in NetLogo was used to test the overall behavior of the model. A complete iteration set of the binary switches turning on and off the constraint submodels and for three values of the possible number of links per species and two values of different landscape types leading to 144 possible combinations which compromise the permutation set used to explore the constraint space, with $t_{max} = 500$ timesteps for each sample from the model hyperspace. A uniform random distribution of the variables propform, varfitness, frqmod, and perwipe, were drawn from the 150 iterations of each of the possible constraint permutation set for each of the These were taken with replacement over the multidimensional parameter space. Mixed model analyses (details are found in Supplemental Materials) were used to explore the influence of each of the parameters thought to be of interest. The results of these analyses need a cautionary warning. As White et al. (2014) (White et al. 2014) note, these results should not be interpreted as a test of statistical significance of each of the parameter set as the likelihood values and statistical tests are easily influenced by the number of iterations used in the model testing. Therefore they were used as a heuristic to chose the set of parameters that might be averaged or representative values chosen that do not influence how the constraints inform the results of the model.

  In all of the model selection analyses, the constraint variables were retained in parameters related to fitness and as expected. For dependent variables involving fitness the full model was usually needed to explain the variation. Exceptions include for within cluster measures of fitness. The variation in turnover rates were explained by the full set of constraints and by the frequency of perturbation and the intensity of the perturbation. This provides evidence that the model's variable and parameter set are adequate.

Statistical Analysis

...

Latin Hypersphere Runs

In the Latin Hypersphere run of the model with random non-constraint independent
variables, the following analyses were used to reconstruct the effects of the constraints
on the model landscape.For each of the constraints, random forests were generated to
explore how different combinations and subsets of the chosen evolutionary constrains
affected the ecological indices [79].

Results and Discussion

The role of the chosen constraints was clear from the simulation results. It took the
shape of both inducing stability and creating conditions for a more dynamic community
with increases in species turnover through time. In fig 4 through fig 8 key ecological
variables showed an overall stability in the landscape structure when plotted against the
number of contraints. For example, fig 4a shows that with an increasing number of
constraints, the overall fitness, and the variance (fig 5) in the fitness was decreasing on
average across the landscape. Implying, by Fisher's fundamental theorem that there
was a lowering of evolvability across the landscape. Likewise, in fig 6a and fig 7a, both
species evenness and species richness were reduced. These measures all indicate that the
number of constraints regardless of which constraint considered simplify the systems
and create a more stable community. This can be further examined by looking at the
purity plots found in fig 4b through fig 8b in which the individual constraints were then
analyzed by constructing a Gini coefficient. The mean decrease in Gini coefficient is a
measure of how each variable contributes to the homogeneity of the nodes and leaves in
the resulting random forest. Each time a particular variable is used to split a node, the
Gini coefficient for the child nodes are calculated and compared to that of the original
node. The Gini coefficient is a measure of homogeneity from 0 (homogeneous) to 1
(heterogeneous). The changes in Gini are summed for each variable and normalized at
the end of the calculation. Variables that result in nodes with higher purity have a
higher decrease in Gini coefficient. A good description of the use of random forests and
their evaluation can be found in [80]. These results explore which constraints are most
effective in creating these structural differences.

Fig 4. Effect of the number of constraints on landscape fitness Along the
horizontal axis, the number of constraints considered over all other variables explored in
the model are plotted against landscape fitness. A. Boxplots of mean fitness considered
over all other variables used in the model for the total number of constraints. B. Forest
plots showing the influence of all possible constrain combinations by use of the Geni
coefficient. See text for more explanation of these results.

**Fig 5. Effect of the number of constraints on mean variance of landscape
fitness** Along the horizontal axis, the number of constraints considered over all other
variables explored in the model are plotted against boxplots of mean landscape variance.

Conclusions

While these evolutionary constraints need much more fleshing out. We understand
enough about the nature of these constraints, that the claim does not seem preposterous.
It might be objected that this type of analogy might not hold in the way that
systems such as biological cells are maintained through constraint closure through

Fig 6. Effect of the number of constraints on ecological evenness Along the horizontal axis, the number of constraints considered over all other variables explored in the model are plotted against ecological evenness (as defined in the text). A. Boxplots of species richness considered over all other variables used in the model for the total number of constraints. B. Forest plots showing the influence of all possible constrain combinations by use of the Geni coefficient. See text for more explanation of these results.

Fig 7. Effect of the number of constraints on species richness Along the horizontal axis, the number of constraints considered over all other variables explored in the model are plotted against species richness. A. Boxplots of species richness considered over all other variables used in the model for the total number of constraints. B. Forest plots showing the influence of all possible constrain combinations by use of the Geni coefficient. See text for more explanation of these results.

Fig 8. Effect of the number of constraints on the difference between turnover in linked and unlinked species Along the horizontal axis, the number of constraints considered over all other variables explored in the model are plotted against difference between turnover in linked and unlinked species. A. Boxplots of difference between turnover in linked and unlinked species considered over all other variables used in the model for the total number of constraints. B. Forest plots showing the influence of all possible constrain combinations by use of the Geni coefficient. See text for more explanation of these results.

specific mechanisms that constrain the flow of energy in material systems. Stretching the notion of constraint closure to something as nebulous as evolutionary 'forces' on individual species and their formation and extinction should be approached with caution. However, we believe it has enough promise to be worth turning over to theorists and modelers who could formalize the structural arguments to demonstrate whether the analogy holds. If these evolutionary forces constrain species in the way we suggest, then the structures of community ecology such as stability and long-term persistence should emerge as constant patterns in such systems allowing for a research program for community ecology that would be able to discover general features of these systems that maintain their continuance through time.

The emergence of particular pattern formation suggests empirical work that may support whether these evolutionary forces form PEP functions in the right way to meet the above criteria defined by Dussault and Bouchard (2016) [48]. Especially, though the use modeling (e.g., agent-based modeling), which provides the structure that would allow the exploration of constraint closure and its effect on ecologically relevant pattern formation, community stability, and turnover.

References

1. Clements FE. Plant Succession: An Analysis of the Development of Vegetation. Carnegie Institution of Washington publication. Carnegie Institution of Washington; 1916.

2. Gleason HA. The Individualistic Concept of the Plant Association. Bulletin of the Torrey Botanical Club. 1926;53(1):7–26. doi:10.2307/2479933.

3. Eliot C. Method and metaphysics in Clements's and Gleason's ecological explanations. Studies in History and Philosophy of Science Part C: Studies in

History and Philosophy of Biological and Biomedical Sciences. 2007;38(1):85–109. doi:http://dx.doi.org/10.1016/j.shpsc.2006.12.006.

4. Götzenberger L, de Bello F, Bråthen KA, Davison J, Dubuis A, Guisan A, et al. Ecological assembly rules in plant communities—approaches, patterns and prospects. Biological Reviews. 2012;87(1):111–127. doi:10.1111/j.1469-185X.2011.00187.x.

5. Bryant R. What If Ecological Communities Are Not Wholes? The Environment: Philosophy, Science, and Ethics. 2012; p. 37–56.

6. Collier J, Cumming G. In: A Dynamical Approach to Ecosystem Identity. vol. 11. Amsterdam: North-Holland; 2011. p. 201–218.

7. Huneman P. Individuality as a theoretical scheme. I. Formal and material concepts of individuality. Biological Theory. 2014;9(4):361–373.

8. Marquet PA, Allen AP, Brown JH, Dunne JA, Enquist BJ, Gillooly JF, et al. On Theory in Ecology. BioScience. 2014;64(8):701–710.

9. Odenbaugh J. Complex ecological systems. Philosophy of complex systems. 2011;.

10. Odenbaugh J. In: Hazlett A, editor. On the Very Idea of an Ecosystem. London: Palgrave Macmillan UK; 2010. p. 240–258.

11. Roughgarden J. Is there a general theory of community ecology? Biology & Philosophy. 2009;24(4):521–529. doi:10.1007/s10539-009-9164-z.

12. Sterelny K. Local Ecological Communities*. Philosophy of Science. 2006;73(2):215–231. doi:10.1086/510819.

13. Sutherland WJ, Freckleton RP, Godfray HCJ, Beissinger SR, Benton T, Cameron DD, et al. Identification of 100 fundamental ecological questions. Journal of Ecology. 2013;101(1):58–67. doi:10.1111/1365-2745.12025.

14. Vellend M. The Theory of Ecological Communities (MPB-57). Monographs in Population Biology. Princeton University Press; 2016.

15. Haber MH, Odenbaugh J. The Edges and Boundaries of Biological Objects. Biological Theory. 2009;4(3):219–224. doi:10.1162/biot.2009.4.3.219.

16. Odenbaugh J. A general unifying theory of ecology. A general theory of ecology. 2011; p. 51–64.

17. Chase JM. Community assembly: when should history matter? Oecologia. 2003;136(4):489–498. doi:10.1007/s00442-003-1311-7.

18. Rosindell J, Hubbell SP, He F, Harmon LJ, Etienne RS. The case for ecological neutral theory. Trends in Ecology & Evolution;27(4):203–208. doi:10.1016/j.tree.2012.01.004.

19. Shrader-Frechette K, McCoy ED. Applied Ecology and the Logic of Case Studies. Philosophy of Science. 1994;61(2):228–249.

20. Simberloff D, Associate Editor: Jonathan BL. Community Ecology: Is It Time to Move On? (An American Society of Naturalists Presidential Address). The American Naturalist. 2004;163(6):787–799. doi:10.1086/420777.

21. Vellend M, Srivastava DS, Anderson KM, Brown CD, Jankowski JE, Kleynhans EJ, et al. Assessing the relative importance of neutral stochasticity in ecological communities. Oikos. 2014;123(12):1420–1430. doi:10.1111/oik.01493.

22. Weiher E, Freund D, Bunton T, Stefanski A, Lee T, Bentivenga S. Advances, challenges and a developing synthesis of ecological community assembly theory. Philosophical Transactions of the Royal Society B: Biological Sciences. 2011;366(1576):2403.

23. Wennekes PL, Rosindell J, Etienne RS. The Neutral—Niche Debate: A Philosophical Perspective. Acta Biotheoretica. 2012;60(3):257–271. doi:10.1007/s10441-012-9144-6.

24. Odenbaugh J. Seeing the Forest ¡em¿and¡/em¿ the Trees: Realism about Communities and Ecosystems. Philosophy of Science. 2007;74(5):628–641. doi:10.1086/525609.

25. Sterelny K. The Reality of Ecological Assemblages: A Palaeo-ecological Puzzle. Biology and Philosophy. 2001;16(4):437–461. doi:10.1023/A:1011972529013.

26. Evans MR, Grimm V, Johst K, Knuuttila T, de Langhe R, Lessells CM, et al. Do simple models lead to generality in ecology? Trends in Ecology & Evolution. 2013;28(10):578–583. doi:10.1016/j.tree.2013.05.022.

27. Heleno R, Garcia C, Jordano P, Traveset A, Gómez JM, Blüthgen N, et al. Ecological networks: delving into the architecture of biodiversity. Biology Letters. 2014;10(1). doi:10.1098/rsbl.2013.1000.

28. Linquist S. Against Lawton's Contingency Thesis; or, Why the Reported Demise of Community Ecology Is Greatly Exaggerated. Philosophy of Science. 2015;82(5):1104–1115. doi:10.1086/684024.

29. Linquist S, Gregory TR, Elliott TA, Saylor B, Kremer SC, Cottenie K. Yes! There are Resilient Generalizations (or "Laws") in Ecology. The Quarterly Review of Biology. 2016;91(2):119–131. doi:10.1086/686809.

30. Loreau M, de Mazancourt C. Biodiversity and ecosystem stability: a synthesis of underlying mechanisms. Ecology Letters. 2013;16:106–115. doi:10.1111/ele.12073.

31. de Mazancourt C, Isbell F, Larocque A, Berendse F, De Luca E, Grace JB, et al. Predicting ecosystem stability from community composition and biodiversity. Ecology Letters. 2013;16(5):617–625. doi:10.1111/ele.12088.

32. Lawton JH. Are There General Laws in Ecology? Oikos. 1999;84(2):177–192. doi:10.2307/3546712.

33. Lockwood D, xa, R. When Logic Fails Ecology. The Quarterly Review of Biology. 2008;83(1):57–64. doi:10.1086/529563.

34. Woodward J. Law and Explanation in Biology: Invariance Is the Kind of Stability That Matters. Philosophy of Science. 2001;68(1):1–20. doi:10.1086/392863.

35. Scheiner SM, Willig MR. A general theory of ecology. Theoretical Ecology. 2008;1(1):21–28. doi:10.1007/s12080-007-0002-0.

36. Proulx SR, Promislow DE, Phillips PC. Network thinking in ecology and evolution. Trends in ecology & evolution. 2005;20(6):345–353.

37. Bascompte J. NNtworks in ecology. Basic and Applied Ecology. 2007;8(6):485–490.

38. Hastings A. Food web theory and stability. Ecology. 1988;69(6):1665–1668.

39. Pimm SL. Food webs. In: Food webs. Springer; 1982. p. 1–11.

40. Hanski I, et al. Metapopulation ecology. Oxford University Press; 1999.

41. Odling-Smee J, Erwin DH, Palkovacs EP, Feldman MW, Laland KN. Niche construction theory: a practical guide for ecologists. The Quarterly review of biology. 2013;88(1):3–28.

42. Odling-Smee FJ, Laland KN, Feldman MW. Niche construction: the neglected process in evolution (MPB-37). vol. 61. Princeton university press; 2013.

43. Krishna A, Guimarães Jr PR, Jordano P, Bascompte J. A neutral-niche theory of nestedness in mutualistic networks. Oikos. 2008;117(11):1609–1618. doi:10.1111/j.1600-0706.2008.16540.x.

44. Losos JB, Ricklefs RE. The theory of island biogeography revisited. Princeton University Press; 2009.

45. Hubbell SP. The unified neutral theory of biodiversity and biogeography (MPB-32). Princeton University Press; 2001.

46. Rosindell J, Hubbell SP, Etienne RS. The unified neutral theory of biodiversity and biogeography at age ten. Trends in ecology & evolution. 2011;26(7):340–348.

47. Peck SL. Agent-based models as fictive instantiations of ecological processes. Philosophy & Theory in Biology. 2012;4.

48. Dussault AC, Bouchard F. A persistence enhancing propensity account of ecological function to explain ecosystem evolution. Synthese. 2016; p. 1–31.

49. Huneman P. About the conceptual foundations of ecological engineering: Stability, individuality and values. Procedia Environmental Sciences. 2011;9:72–82. doi:http://dx.doi.org/10.1016/j.proenv.2011.11.013.

50. Montévil M, Mossio M. Biological organisation as closure of constraints. Journal of Theoretical Biology. 2015;372:179–191. doi:http://dx.doi.org/10.1016/j.jtbi.2015.02.029.

51. Moreno A, Mossio M. Biological Autonomy: A Philosophical and Theoretical Enquiry. History, Philosophy and Theory of the Life Sciences. Springer Netherlands; 2015.

52. Nunes-Neto N, Moreno A, El-Hani CN. Function in ecology: an organizational approach. Biology & Philosophy. 2014;29(1):123–141. doi:10.1007/s10539-013-9398-7.

53. Vellend M. Conceptual Synthesis in Community Ecology. The Quarterly Review of Biology. 2010;85(2):183–206. doi:10.1086/652373.

54. Clarke E. The multiple realizability of biological individuals. The Journal of Philosophy. 2013;110(8):413–435.

55. Griesemer J. Individuation of Developmental Systems. Individuation, Process, and Scientific Practices. 2018; p. 137.

56. Millstein RL. Debunking myths about Aldo Leopold's land ethic. Biological conservation. 2018;217:391–396.

57. Millstein RL. Is Aldo Leopold's 'Land Community'an Individual? Individuation, Process, and Scientific Practices. 2018; p. 279.

58. Mossio M, Saborido C, Moreno A. An Organizational Account of Biological Functions. The British Journal for the Philosophy of Science. 2009;60(4):813–841.

59. Pigliucci M. Sewall Wright's adaptive landscapes: 1932 vs. 1988. Biology & Philosophy. 2008;23(5):591–603. doi:10.1007/s10539-008-9124-z.

60. Peck SL. The hermeneutics of ecological simulation. Biology & Philosophy. 2008;23(3):383–402. doi:10.1007/s10539-008-9109-y.

61. Hutchinson GE. The concept of pattern in ecology. Proceedings of the Academy of Natural Sciences of Philadelphia. 1953;105:1–12.

62. Levin SA. The problem of pattern and scale in ecology: the Robert H. MacArthur award lecture. Ecology. 1992;73(6):1943–1967.

63. Chave J. The problem of pattern and scale in ecology: what have we learned in 20 years? Ecology Letters. 2013;16:4–16. doi:10.1111/ele.12048.

64. Humphreys P. Emergence. Oxford University Press; 2016.

65. Grimm V, Railsback SF. Pattern-oriented modelling: a 'multi-scope' for predictive systems ecology. Philosophical Transactions of the Royal Society B: Biological Sciences. 2012;367(1586):298.

66. Simard SW, Beiler KJ, Bingham MA, Deslippe JR, Philip LJ, Teste FP. Mycorrhizal networks: Mechanisms, ecology and modelling. Fungal Biology Reviews. 2012;26(1):39–60. doi:http://dx.doi.org/10.1016/j.fbr.2012.01.001.

67. Yang G, Liu N, Lu W, Wang S, Kan H, Zhang Y, et al. The interaction between arbuscular mycorrhizal fungi and soil phosphorus availability influences plant community productivity and ecosystem stability. Journal of Ecology. 2014;102(4):1072–1082. doi:10.1111/1365-2745.12249.

68. Oldroyd GED. Speak, friend, and enter: signalling systems that promote beneficial symbiotic associations in plants. Nat Rev Micro. 2013;11(4):252–263.

69. Perry DA. Forest Ecosystems. Johns Hopkins paperback. Life sciences. Johns Hopkins University Press; 1994.

70. Leyval C, Turnau K, Haselwandter K. Effect of heavy metal pollution on mycorrhizal colonization and function: physiological, ecological and applied aspects. Mycorrhiza. 1997;7(3):139–153.

71. Mnasri M, Janoušková M, Rydlová J, Abdelly C, Ghnaya T. Comparison of arbuscular mycorrhizal fungal effects on the heavy metal uptake of a host and a non-host plant species in contact with extraradical mycelial network. Chemosphere. 2017;171:476–484.

72. Booth MG. Mycorrhizal networks mediate overstorey-understorey competition in a temperate forest. Ecology Letters. 2004;7(7):538–546. doi:10.1111/j.1461-0248.2004.00605.x.

73. Gorzelak MA, Asay AK, Pickles BJ, Simard SW. Inter-plant communication through mycorrhizal networks mediates complex adaptive behaviour in plant communities. AoB Plants. 2015;7:plv050.

74. Heijden MGA, Martin FM, Selosse M, Sanders IR. Mycorrhizal ecology and evolution: the past, the present, and the future. New Phytologist. 2015;205(4):1406–1423.

75. Traveset A, Richardson DM. Mutualistic interactions and biological invasions. Annual Review of Ecology, Evolution, and Systematics. 2014;45:89–113.

76. Peay KG. The Mutualistic Niche: Mycorrhizal Symbiosis and Community Dynamics. Annual Review of Ecology, Evolution, and Systematics. 2016;47(1).

77. Weisberg M. Simulation and Similarity: Using Models to Understand the World. Oxford Studies in Philosophy of Science. OUP USA; 2013.

78. Grimm V, Berger U, DeAngelis DL, Polhill JG, Giske J, Railsback SF. The ODD protocol: A review and first update. Ecological Modelling. 2010;221(23):2760–2768. doi:http://dx.doi.org/10.1016/j.ecolmodel.2010.08.019.

79. Kabasenche WP, O'Rourke M, Slater MH. The Environment: Philosophy, Science, and Ethics. Topics in contemporary philosophy. MIT Press; 2012.

80. Maurer BA. Untangling Ecological Complexity: The Macroscopic Perspective. Untangling Ecological Complexity: The Macroscopic Perspective. University of Chicago Press; 1999.

Adapted from: Moreno and Mossio, 2015, *Biological Autonomy: A Philosophical and Theoretical Enquiry*, Fig. 1.3 p. 21

Constraints in Community Ecology

Point = mean value

Point = mean value

Point = mean value

Average Evenness_t

Point = mean value

landscape richness m

Point = mean value

ave turnover dlu