# Biofilms mechanically damage epithelia by buckling.

Alice Cont, Tamara Rossy, Zainebe Al-Mayyah, Alexandre Persat*

Institute of Bioengineering and Global Health Institute, School of Life Sciences, Ecole

Polytechnique Fédérale de Lausanne, Lausanne, Switzerland

*alexandre.persat@epfl.ch

**Abstract**

During chronic infections and in microbiota, bacteria predominantly colonize their hosts as multicellular structures called biofilms. Biofilms form when single cells divide while embedding themselves in an elastic matrix of extracellular polymeric substance, protecting resident cells from external stressors and coordinating collective behaviors[1]. While we understand how biofilms form in vitro, we still lack a basic understanding of how they interact with tissues in vivo, and ultimately how they influence host physiology. A common assumption is that biofilm-dwelling bacteria biochemically interact with their hosts, for example by secreting toxins[2]. However, the contributions of mechanics, while being central to the process of biofilm formation, have been vastly overlooked as a factor influencing host physiology. Here we show that biofilms mechanically damage epithelia by transmitting of internally-generated forces to host tissue. By combining tissue-engineering and biomechanical measurement techniques, we found that forces generated by *Vibrio cholerae* biofilms can disrupt host epithelia by breaking epithelial cell-cell junctions and attachment to extracellular matrices. We demonstrate that biofilms from *V. cholerae* and *Pseudomonas aeruginosa* can similarly induce large deformations of soft synthetic hydrogels. Using traction force microscopy, we demonstrate that friction between the biofilm and the hydrogel surface generates internal mechanical stress that cause biofilms to buckle. Matrix components that maintain the mechanical cohesion of biofilms promote buckling, while those

26 that maintain surface adhesion transmit buckling stress to the substrate. Altogether, our

27 results demonstrate that forces generated by bacterial communities play an important role

28 not only in biofilm morphogenesis but also in host physiology, suggesting a mechanical

29 mode of infection.

32  To explore how biofilms form within their hosts, we engineered epithelial monolayers of

33 Caco-2 cells on a soft extracellular matrix (ECM) (**Fig. 1A**). This produces soft and tight

34 ECM-adherent epithelia that reproduce the mechanical properties of real tissues. We

35 seeded the surface of these epithelia with *V. cholerae* expressing constitutively high levels of

36 cyclic-di-GMP (Vc WT*), which forms robust biofilms and has reduced virulence[3]. Biofilms

37 formed at the epithelial surface within 20 h (**Fig. 1B-D**). Overall, biofilms perturbed the shape

38 of the epithelium. Under biofilms, the cell monolayer detached from its ECM substrate and

39 was often bent (**Fig. 1B-ii**). In other instances, Caco-2 cells lost cohesion and were engulfed

40 by the biofilm. This allowed the biofilm to breach the epithelium and contact the ECM. There,

41 biofilms deformed the ECM substrate, turning the initially flat surface into a dome-like shape

42 (**Fig. 1B-iv**). These disruptions did not depend on host cell type as *V. cholerae* could also

43 damage and bend monolayers of MDCK cells which has strong cell-cell junctions[4] (**Fig. 1C**).

44 Our observations suggest that biofilms apply mechanical forces on host tissue that can

45 perturb the morphology and integrity of epithelia, as well as the underlying ECM.

46  To test whether and how biofilms disrupt epithelia, we sought to decouple the effect of

47 mechanics from host cell biological response by exploring their formation on soft axenic

48 substrates. We generated ~100 µm-thick polyethylene glycol (PEG) hydrogel films in

49 microchannels, thus enabling high resolution live confocal imaging of biofilm formation under

50 flow (**Fig. 2A**). On soft hydrogels, *V. cholerae* formed biofilms whose bottom surfaces

51 appeared bell-shaped, consistent with what we observed on epithelial monolayers and ECM

(**Fig. 2B**), but in striking difference with the typically flat-growing biofilms forming on glass. To test whether this shape was a result of hydrogel deformation, we embedded fluorescent tracer particles within the hydrogel film. We could observe that biofilms in fact deformed hydrogels as the fluorescent tracer particles filled the apparent bell-shaped void at its core (**Fig. 2C**). We also found that biofilms of *P. aeruginosa*, an opportunistic pathogen that forms robust airway epithelial biofilms during chronic pneumonia, could similarly deform soft PEG hydrogels (**Fig. 2D-E, Fig. S1**). Therefore, *V. cholerae* and *P. aeruginosa* both deform the substrate, consistent with a force-driven mechanism that is independent of their respective biofilm architecture and the composition of their extracellular polymeric substance (EPS).

How could biofilms mechanically deform their substrates? Bacterial colonies wrinkle on agar plates due to internal mechanical stress generated by simultaneous expansion and friction[5]. In biofilms, this also causes internal stresses influencing the spatial organization of single cells within *V. cholerae* biofilms[6,7]. Friction force between the microcolony and the surface opposes biofilm expansion, generating an inward internal stress that leads to a buckling instability verticalizing or reorienting contiguous cells[7,8]. We thus reasoned that biofilm could deform soft substrates by transmission of internal stresses to the substrate they grow on. To demonstrate this, we performed dynamic visualizations of hydrogel film deformations and recorded the surface profiles for multiple biofilms. By radial re-slicing and averaging around the biofilm centers, we could extract the deformation profile $\delta$, its maximum deformation amplitude $\delta_{max}$ and full-width at half maximum $\lambda$ (**Fig. 3A**). By reconstructing hydrogel surfaces for biofilms of different sizes, we found that $\delta_{max}$ and $\lambda$ linearly scaled with the diameter $d$ of the biofilm (**Fig. S2**), indicating that biofilm expansion promotes surface deformation. We went further by tracking these deformations over time for single biofilms. Deformations increased as biofilms grew, even showing a slight recess near the biofilm edges reminiscent of higher order buckling modes (**Fig. 3B-C, Movie S1**). In these visualizations, we noticed that there was a lag between the increase in biofilm diameter and the onset of deformation, with a finite deformation only appearing after 7 h of

growth. This was further confirmed by following the deformations generated by many

biofilms, as finite in plane deformations mainly appeared after 6 to 7 h of growth (**Fig. 3D**).

Rescaling by the diameter of the biofilm collapsed $\delta_{max}$ measurements, highlighting a critical

biofilm diameter (35 µm) above which deformations emerged (**Fig. 3E**). The existence of a

critical diameter is reminiscent to buckling instabilities of rigid bodies subject to compressive

stress, as in Euler buckling. Also consistent with an Euler-type buckling instability, the width

of the deformation scales linearly with biofilm diameter[9].

To further support our model where biofilm buckling drives surface deformations, we

investigated how bacterial growth applied forces on the hydrogel. We therefore tracked the

displacements of the fluorescent particle tracers embedded within the hydrogel in 3D with a

digital volume correlation algorithm. Specifically, the hydrogel displacement field shows that

the biofilm pushes the hydrogel radially before reaching the critical diameter, in the outward

direction (**Fig. 3F** and **Fig. S4**). These deformations result from the opposition between

growth and friction between cells and the surface, and generate an internal force within the

biofilm oriented radially towards its center, which ultimately causes the biofilm to buckle.

We then wondered whether forces driving deformations could break epithelial cell-cell

junctions and disrupt monolayers. To answer this question, we directly measured the forces

generated by the biofilm on the hydrogel films by traction force microscopy. Traction forces

were surprisingly large, reaching 5 MPa in the biofilm center after 12 h of growth (**Fig. 3G**).

In comparison, cell-cell junctions break when experiencing a few kPa [10]. Therefore, biofilms

produce sufficient force to mechanically dismantle epithelia. We note that the magnitude of

the stress is relatively large for a biological structure of this size. This stress is in essence

generated by the expansion of single bacterial cell, itself driven by turgor pressures can

reach the MPa range[11].

Given the large forces generated by biofilms on substrates, we wondered to which extent

they could deform different types of tissues. To test this, we reproduced the mechanical

properties of various tissue types by tuning the stiffness of the PEG hydrogel films between

10 kPa and 200 kPa [12,13]. The rate of increase of the deformation amplitude was inversely

correlated with stiffness, resulting in differences in $\delta_{max}$ between colonies of identical

diameter growing on substrates with distinct stiffnesses **(Fig. 3H)**. For each stiffness, the

deformation amplitude $\delta_{max}$ and the width $\lambda$ increased linearly with biofilm diameter (**Fig. 3I**

and **Fig. S3**). Stiff hydrogels only slightly deformed, recapitulating the morphology of glass-

grown biofilms. In contrast, biofilms growing on soft hydrogels displayed large deformations

**(Fig. 3I)**. Rescaling $\delta_{max}$ with the biofilm diameter highlights a power-law relationship

between deformation and substrate stiffness qualitatively consistent with the theory of

buckling of plates coupled to an elastic foundation (**Fig. S5**)[14].

We then identified biofilm components regulating force generation and transmission,

particularly focusing on EPS matrix components. The *V. cholerae* matrix is composed of a

polysaccharide and proteins, including Rbma which is responsible for cell-cell cohesion and

Bap1 which promotes biofilm-surface adhesion[15]. We found that biofilms of *rbmA* deletion

mutants were unable to deform the hydrogel substrate, demonstrating that cell-cell cohesion

is an essential ingredient in force generation (**Fig. 4A**). Interestingly, biofilms of *bap1*

deletion mutants grew slightly bent but delaminated from the substrate which remained flat,

thereby creating a gap between the biofilm and the hydrogel (**Fig. 4A**). This demonstrates

that adhesion between the biofilm and the substrates transmits normal mechanical stress. In

a similar manner, mutation in EPS matrix components of *P. aeruginosa* reduced but did not

abolish deformations (**Fig. 4B**). *P. aeruginosa* secretes the polysaccharides Pel and Psl, and

the protein CdrA, all of which play a role in maintaining viscoelastic properties of the

biofilm[16–18]. Deletion mutants in *psl, pel* and *cdrA* showed a decrease in deformation

amplitude, further demonstrating that mechanical cohesion plays a key role in surface

deformation (**Fig. 4B-C**). Since *P. aeruginosa* does not possess a dedicated matrix

component for substrate adhesion, we grew biofilms on hydrogels with large Young's

modulus to increase the relative contribution of adhesion and elastic energy of the

substrate[19]. Consistent with our buckling-adhesion model, these biofilms delaminated (**Fig.**

**4D**). In summary, cell-cell mechanical cohesion is essential in generating the internal stress

that promotes biofilm buckling, while cell-substrate adhesion transmits this stress to the

underlying substrate (**Fig. 4E**).

We found that biofilms can cause damage to epithelial tissues by buckling, causing

delamination or rupture of intercellular junctions, suggesting that biofilms could mechanically

damage host tissues. In fact, biofilms commonly cause tissue lesions. For example the urine

of vaginosis patients contains desquamated epithelial cells covered with biofilms[20,21]. Also,

commensal biofilms form scabs at the epithelial surface of honeybee's gut, triggering

immune responses [22]. Epithelial integrity is also commonly compromised in intestinal

diseases such as inflammatory bowel disease, while being highly influenced by the

microbiota[23]. Finally, hyper-biofilm forming clinical variants of *P. aeruginosa* causes

significant damage to the surrounding host tissue despite its reduced virulence[24].

Mechanical interactions between bacterial collectives and their host may thus represent an

overlooked contributor of infections and dysbiosis.

**Acknowledgements**

We would like to thank the Fitnat Yildiz, Matt Parsek, Melanie Blokesch and Bonnie Bassler

for strains and plasmids and John Kolinski and Pedro Reis for discussions.

**Funding**

This work was supported by the Swiss National Science Foundation Projects Grant

31003A_169377, the Gabriella Giorgi-Cavaglieri Foundation, the Gebert Rüf Stiftung and the

Fondation Beytout.

**Competing interests**

None

**Materials and Methods**

*Cell culture*

Caco-2 cells and MDCK cells were maintained in T25 tissue culture flasks (Falcon) with

DMEM medium (Gibco) supplemented with 10% fetal bovine serum at 37°C in a CO2

incubator.

*Cell culture on collagen/Matrigel gels*

To resemble the extracellular matrix natural niche, we cultured epithelial cells at the surface

of collagen and Matrigel based hydrogels. Hydrogel solutions were prepared on ice to avoid

premature gelation by mixing 750 µl of neutralized collagen with 250 µl of growth-factor

reduced Matrigel matrix (Corning, 356231). The neutralized collagen was obtained by mixing

800 µl of native type I collagen isolated from the bovine dermis (5mg/ml, Cosmo Bio Co.,

Ltd.) with 10 µl of $NaHCO_3$ (1 M), 100 µl of DMEM-FBS and 100 µl of  DMEM 10X. We then

spread 100 μl of the hydrogel solution in glass bottom dishes (P35G-1.5-20-C, MatTek),

which were kept on ice. Excess solution was removed from the sides of the well to avoid the

formation of a meniscus. To promote collagen adhesion, the wells were previously

functionalized with a 2% polyethyleneimine solution (Sigma-Aldrich) for 10 min and a 0.4%

glutaraldehyde solution (Electron Microscopy Science) for 30 min. We finally placed the

coated dishes at 37°C in a $CO_2$ incubator for 20 minutes to allow gelation.

MDCK and Caco-2 cells were detached from the flask using trypsin (Sigma-Aldrich). We

seeded the cells at a concentration of 1000 cells/mm$^2$ on top of the gels. We let the cells

adhere for 1 day and then we filled the dishes with 2 ml of culture medium. The medium was

changed every 2 days.

*Bacterial strains and culture conditions*

A list of the strains and plasmids is provided in **Table 1**. All strains were grown in LB medium

at 37°C. Deletion of the *V. cholerae* genes *rbmA* and *bap1* were generated by mating a

parental A1552 *V. cholerae* strain, rugose variant, with *E. coli* S17 strains harboring the

deletion constructs according to previously published protocols [25]. *P.aeruginosa* strains

(PAO1 parental strain) are all constitutively expressing GFP (attTn7::miniTn7T2.1-Gm-

GW::PA1/04/03::GFP).

*Infection of tissue-engineered epithelia by Vibrio cholera*

*V. cholerae* was grown in LB medium at 37°C to mid-exponential phase (OD 0.3-0.6).

Bacteria were washed 3 times by centrifugation and resuspension in Dulbecco's phosphate-

buffered saline (D-PBS). The cultures were then diluted to an optical density of 10$^{-7}$ and

filtered (5.00 μm-pore size filters, Millex) to ensure the removal of large bacterial clumps,

thereby isolating planktonic cells. This ensured that biofilms growing on epithelia formed

from single cells. We loaded 200 μL of diluted culture on top of Caco-2 or MDCK cells that

were cultured for 1 to 7 days post-confluence on collagen/Matrigel gels. Bacteria were

allowed to adhere to the surface for 20 minutes, after which cells were rinsed two times with

D-PBS.

For the implementation of the flow on top of Caco-2 cells, we prepared a circular slab of

PDMS with the same dimensions as the dish. We punched 1mm inlet and outlet ports in this

PDMS slab. We then glued it to the rim of the dish, where no cells are present. We then

connected the inlet port to a disposable syringe (BD Plastipak) filled with culture medium

using a 1.09 mm outer diameter polyethylene tube (Instech) and a 27G blunt needle

(Instech). The syringes were mounted onto a syringe pump (KD Scientific) positioned inside

a $CO_2$ incubator at 37°C. The volume flow rate was set to 50 µL·min$^{-1}$.

For stationary biofilm growth on MDCK cells, the glass bottom dishes were filled with 2

mL of culture medium and were incubated at 37°C in a $CO_2$ incubator.

*Fabrication of PEG hydrogels and mechanical characterization*

To generate PEG hydrogels films we prepared solutions of M9 minimal medium containing

poly(ethylene glycol) diacrylate (PEGDA) as the precursor and lithium phenyl-2,4,6-

trimethylbenzoylphosphinate (LAP, Tokio Chemical Industries) as the photoinitiator.

Molecular weight and concentration of PEGDA were tuned to obtain hydrogels with different

stiffnesses (**Table 2**), while the concentration of LAP is kept constant at 2 mM.

To incorporate fluorescent microparticles into the PEG hydrogels, we modified the original

solution by substituting 2 µL of M9 medium with 2 µL of red fluorescent particles solution

(ThermoFischer, FluoSpheres, Carboxylate-modified Microspheres, 0.1 µm diameter, 2%

solids, F8887).

To prepare the samples for mechanical characterization, we filled PDMS wells (5 mm

diameter, 4 mm height) with the hydrogel solution. We covered the wells with a coverslip and

we let them polymerize in a UV transilluminator (Bio-Rad Universal Hood II) for 5 minutes.

The resulting hydrogel cylinders were immersed in M9 overnight and tested with a

rheometer (TA instruments) in compression mode, at a deformation rate of 10 µm/s.

Beforehand, the diameter of the cylinders was measured with a digital caliper, while the

height of the cylinder was defined as the gap distance at which the force starts differing from

zero. The elastic modulus corresponds to the slope of the linear fit of the stress-strain curves

in the range of 15% strain. The final modulus is the average modulus of 3 replicates.

*Fabrication of thin PEG hydrogel layers and implementation with PDMS microfluidic chip*

We fabricated microfluidic chips following standard soft lithography techniques. More

specifically, we designed 2 cm-long, 2 mm-wide channels in Autodesk AutoCAD and printed

them on a soft plastic photomask. We then coated silicon wafers with photoresist (SU8

2150, Microchem), with a thickness of 350 μm. The wafer was exposed to UV light through

the mask and developed in PGMEA (Sigma-Aldrich) in order to produce a mold. PDMS

(Sylgard 184, Dow Corning) was subsequently casted on the mold and cured at 70 °C

overnight. After cutting out the chips, we punched 1 mm inlet and outlet ports. We finally

punched a 3 mm hole right downstream of the inlet port. This hole, after being covered with

a PDMS piece, acts as a bubble trap.

To obtain thin and flat hydrogel layers, a drop of about 80 μL of the hydrogel solution was

sandwiched between two coverslips and incubated in the UV transilluminator for 5 minutes

to allow gelation. The bottom coverslip (25x60 mm Menzel Gläser) was cleaned with

isopropanol and MilliQ water, while the upper one (22x40 mm Marienfeld) was functionalized

with 3-(Trimethoxysilyl)propyl methacrylate (Sigma-Aldrich) following the standard

procedure. In short, cleaned coverslips were immersed in a 200 mL solution of ethanol

containing 1 mL of the reagent and 6ml of dilute acetic acid (1:10 glacial acetic acid:water)

for 5 minutes. They were subsequently rinsed in ethanol and dried. This functionalization

enables the covalent linkage of the hydrogel to the coverslip.

Right after polymerization, the coverslips were separated using a scalpel and thus

exposing the hydrogel film surface. We then positioned the PDMS microfluidic chip on top of

the hydrogel film. This results in a reversible, but sufficiently strong bond between the

hydrogel and the PDMS, allowing us to use the chips under flow without leakage for several

days. The assembled chips were filled with M9 to maintain the hydrogel hydrated.

*Biofilm growth in microfluidic chambers*

All *V. cholerae* and *P. aeruginosa* strains were grown in LB medium at 37°C until mid-

exponential phase (OD 0.3-0.6). The cultures were diluted to an optical density of $10^{-3}$ and

subsequently filtered (5.00 µm-pore size filters, Millex) to ensure the removal of large

bacterial clumps. We then loaded 6.5 µL of the diluted bacterial culture in the channels, from

the outlet port. We let them adhere for 20 minutes before starting the flow. We connected

the inlet port to a disposable LB-filled syringe (BD Plastipak) mounted onto a syringe pump

(KD Scientific), using a 1.09 mm outer diameter polyethylene tube (Instech) and a 27G

needle (Instech). For all conditions, the volume flow rate was 10 µL·min$^{-1}$, which

corresponds to a mean flow speed of about 0.25 mm·s$^{-1}$ inside the channels. The biofilms

were grown at 25°C.

*Staining procedures*

Caco-2 cells and MDCK cells were incubated for 20 minutes in a 10 µM solution of

CellTracker Orange CMRA (Invitrogen, C34551) and washed with DPBS before seeding the

bacteria.

Since *V. cholerae* strains were not constitutively fluorescent, biofilms were incubated for

20 minutes with a 10 µM solution of SYTO9 (Invitrogen, S34854) and washed with M9

minimal medium before visualization. This results in double staining of epithelial cells in the

case of infection experiments.

*Visualization*

For all visualizations, we used an Nikon Eclipse Ti2-E inverted microscope coupled with a

Yokogawa CSU W2 confocal spinning disk unit and equipped with a Prime 95B sCMOS

camera (Photometrics). For low magnification images, we used a 20x water immersion

objective with N.A. of 0.95, while for all the others we used a 60x water immersion objective

with a N.A. of 1.20. We used Imaris (Bitplane) for three-dimensional rendering of z-stack

pictures and Fiji for the display of all the other images.

To obtain the deformation profiles, z-stacks of the hydrogel containing fluorescent

microparticles were performed every 0.5 µm, while a brightfield image of the base of the

biofilm was taken to allow measurement of the diameter of the biofilm. For the visualization

of the full biofilm, z-stacks of the samples were taken every 2-3 µm. For timelapse

experiments, biofilms were imaged as soon as the flow was started, while for all the other

experiments biofilms were imaged between 10 and 24 h post-seeding.

*Image analysis and computation of deformation profiles*

Starting from confocal imaging pictures of the microparticle-containing hydrogel, we aimed at

identifying the gel surface and extracting quantitative information about its deformation

induced by the biofilms. In most cases, we used an automated data analysis pipeline as

described below. To get an average profile of the deformation caused by the biofilms, we

performed a radial reslice in Fiji over 180 degrees around the center of the deformation (one

degree per slice). We then performed an average intensity projection of the obtained stack.

Tocalculate the diameter of the biofilm, we averaged 4 measurements of the biofilm

diameter taken at different angles. The resliced images were then imported in Matlab

R2017a (Mathworks) as two-dimensional (x-y) matrices of intensities. In these images, the

surface was consistently brighter than the rest of the gel. Therefore, we identified the surface

profile as the pixels having the maximal intensity in each column of the matrix. Note that the

bottom of the gel sometimes also comprised bright pixels that introduced noise in the profile.

To reduce this problem, we thus excluded 20 rows at the bottom of each image (~3.7 µm).

We then calculated the baseline position of our gel – namely, the height of the non-deformed

portion of the gel. In our pictures, this corresponds to the height at the left and right

extremities of the profile. Therefore, we defined the baseline as the average of the first 50

and last 50 pixels of the profile (~9 µm on each side of the profile). We then offset the whole

picture so that the baseline position corresponded to y = 0. We undersampled the extracted

surface profiles to further reduce noise, by keeping only the maximal y value over windows

of 40 pixels. Finally, we fitted a smoothing spline to the undersampled profile using the built-

in *fit* function in Matlab, with a smoothing parameter value of 0.9999.

To quantify the deformation that biofilms induced on the hydrogel, we measured the

amplitude ($\delta_{max}$) of the deformed peak and its full width at half maximum (λ). First, we

evaluated the fitted profile described above at a range of points spanning the whole width of

the picture and spaced by 0.0005 µm. We identified the maximal value of the profile at these

points, which corresponds to the amplitude of the peak $\delta_{max}$ (with respect to the baseline,

which is defined as y = 0). We then split the profile in two: one part on the left of the

maximum, and one part on its right. On each side, we found the point on the profile whose y

value was the closest to $0.5 \cdot \delta_{max}$ using the Matlab function *knnsearch*. We then calculated

the distance between their respective x values, which corresponds to the λ of the deformed

peak. Our data analysis program also included a quality control feature, which prompted the

user to accept or reject the computed parameters. When imaging quality was insufficient to

ensure proper quantification with our automated pipeline, we measured the deformation

manually in Fiji.

*Digital volume correlation and traction force microscopy*

We performed particle tracking to measure local deformations and ultimately compute stress

and traction forces within hydrogels as biofilms grew. To do this, we performed timelapse

visualizations of the hydrogel during the formation of a biofilm at high spatial resolution with

a 60X, NA 0.95 water immersion objective. We thus generated 200 µm x 200 µm x 25 µm

(50 stacks of 1200x1200 pixels) volumes at 14 different time points. These images were

subsequently registered to eliminate drift using the Correct 3D Drift function in Fiji. To

compute local material deformations which we anticipated to generate large strains, we used

an iterative Digital Volume Correlation (DVC) scheme [26]. These were performed with

128x128x64 voxel size in cumulative mode, meaning deformations are calculated by

iterations between each time point over the whole 4D timelapse, rather than directly from the

reference initial image. The DVC code computes material deformation fields in 3D which we

subsequently use as input for the associated large deformation traction force microscopy

(TFM) algorithm [26]. The TFM calculates stress and strain fields given the material's Young

modulus ($E$ = 38 kPa in our case) to ultimately generate a traction force map at the hydrogel

surface.

**References**

1.   Flemming, H. C. & Wingender, J. The biofilm matrix. *Nat. Rev. Microbiol.* **8**, 623–633

(2010).

2.   Donlan, R. M. & Costerton, J. W. Biofilms: Survival mechanisms of clinically relevant

microorganisms. *Clinical Microbiology Reviews* **15**, 167–193 (2002).

3.   Tischler, A. D. & Camilli, A. Cyclic diguanylate regulates Vibrio cholerae virulence

gene expression. *Infect. Immun.* **73**, 5873–82 (2005).

4.   Harris, A. R. *et al.* Characterizing the mechanics of cultured cell monolayers. *Proc.*

*Natl. Acad. Sci. U. S. A.* **109**, 16449–16454 (2012).

5.   Yan, J. *et al.* Mechanical instability and interfacial energy drive biofilm

morphogenesis. *Elife* **8**, (2019).

6.   Hartmann, R. *et al.* Emergence of three-dimensional order and structure in growing

biofilms. *Nat. Phys.* **15**, 251–256 (2019).

7.   Beroz, F. *et al.* Verticalization of bacterial biofilms. *Nat. Phys.* **14**, 954–960 (2018).

8.   Duvernoy, M. C. *et al.* Asymmetric adhesion of rod-shaped bacteria controls

microcolony morphogenesis. *Nat. Commun.* **9**, 25–28 (2018).

9.   Timoshenko, S. & Gere, J. M. *Theory of elastic stability*. (Dover Publications, 2009).

10.  Charras, G. & Yap, A. S. Tensile Forces and Mechanotransduction at Cell–Cell

Junctions. *Curr. Biol.* **28**, R445–R457 (2018).

11.  Rojas, E. R. & Huang, K. C. Regulation of microbial growth by turgor pressure. *Curr.*

*Opin. Microbiol.* **42**, 62–70 (2018).

12.  Discher, D. E., Janmey, P. & Wang, Y. Tissue Cells Feel and Respond to the

Stiffness of Their Substrate. **310**, 1139–1144 (2005).

13.  Lee, S., Tong, X. & Yang, F. The effects of varying poly(ethylene glycol) hydrogel

crosslinking density and the crosslinking mechanism on protein accumulation in three-

dimensional hydrogels. *Acta Biomater.* **10**, 4167–4174 (2014).

14.  Wang, C. Y. On the Buckling of a Circular Plate on an Elastic Foundation. *J. Appl.*

*Mech.* **72**, 795–796 (2005).

15.  Teschler, J. K. *et al.* Living in the matrix: assembly and control of Vibrio cholerae

biofilms. *Nat. Rev. Microbiol.* **13**, 255–268 (2015).

16.  Jennings, L. K. *et al.* Pel is a cationic exopolysaccharide that cross-links extracellular

DNA in the Pseudomonas aeruginosa biofilm matrix. **112**, 11353–11358 (2015).

17.  Kovach, K. *et al.* Evolutionary adaptations of bio fi lms infecting cystic fi brosis lungs

promote mechanical toughness by adjusting polysaccharide production. (2016).

doi:10.1038/s41522-016-0007-9

18.  Colvin, K. M. *et al.* The Pel and Psl polysaccharides provide Pseudomonas

aeruginosa structural redundancy within the biofilm matrix. *Environ. Microbiol.* **14**,

1913–28 (2012).

19.  Wang, Q. & Zhao, X. A three-dimensional phase diagram of growth-induced surface

instabilities. *Sci. Rep.* **5**, 8887 (2015).

20.  Swidsinski, A. *et al.* Presence of a Polymicrobial Endometrial Biofilm in Patients with

Bacterial Vaginosis. *PLoS One* **8**, e53997 (2013).

21.   Hardy, L. *et al.* A fruitful alliance: the synergy between Atopobium vaginae and

Gardnerella vaginalis in bacterial vaginosis-associated biofilm. *Sex. Transm. Infect.*

**92**, 487–491 (2016).

22.   Engel, P., Bartlett, K. D. & Moran, N. A. The Bacterium Frischella perrara Causes

Scab Formation in the Gut of its Honeybee Host. *MBio* **6**, e00193-15 (2015).

23.   Chelakkot, C., Ghim, J. & Ryu, S. H. Mechanisms regulating intestinal barrier integrity

and its pathological implications. *Exp. Mol. Med. 2018 508* **50**, 1–9 (2018).

24.   Pestrak, M. J. *et al.* Pseudomonas aeruginosa rugose small-colony variants evade

host clearance, are hyper-inflammatory, and persist in multiple host environments.

*PLOS Pathog.* **14**, e1006842 (2018).

25.   Fong, J. C. N., Karplus, K., Schoolnik, G. K. & Yildiz, F. H. Identification and

characterization of RbmA, a novel protein required for the development of rugose

colony morphology and biofilm structure in Vibrio cholerae. *J. Bacteriol.* **188**, 1049–59

(2006).

26.   Toyjanova, J. *et al.* High Resolution, Large Deformation 3D Traction Force

Microscopy. *PLoS One* **9**, e90976 (2014).

27.   Berk, V. *et al.* Molecular architecture and assembly principles of Vibrio cholerae

biofilms. *Science* **337**, 236–9 (2012).

28.   Yildiz, F. H., Liu, X. S., Heydorn, A. & Schoolnik, G. K. Molecular analysis of rugosity

in a Vibrio cholerae O1 El Tor phase variant. *Mol. Microbiol.* **53**, 497–515 (2004).

29.   Hickman, J. W., Tifrea, D. F. & Harwood, C. S. A chemosensory system that regulates

biofilm formation through modulation of cyclic diguanylate levels. *Proc. Natl. Acad.*

*Sci. U. S. A.* **102**, 14422–7 (2005).

30.    Rybtke, M. T. *et al.* Fluorescence-based reporter for gauging cyclic di-GMP levels in

412           Pseudomonas aeruginosa. *Appl. Environ. Microbiol.* **78**, 5060–9 (2012).

413    31.    Rybtke, M. *et al.* The LapG protein plays a role in Pseudomonas aeruginosa biofilm

414           formation by controlling the presence of the CdrA adhesin on the cell surface.

415           *Microbiologyopen* **4**, 917–30 (2015).

**Fig. 1: Biofilms deform and disrupt epithelial cell monolayer.** (A) Caco-2 and MDCK

cells grow at the surface of a soft ECM into a tight monolayer on which we seed a liquid

inoculum of *V. cholerae*. (B) Confocal images of uninfected (i) and infected (ii-v) monolayers

of Caco-2 cells. Yellow arrow indicates gaps in the epithelial monolayer (ii and iii), blue arrow

shows deformed ECM (iv). (C) Confocal images of uninfected (i) and infected (ii-iii)

425    monolayers of MDCK cells, also showing delamination and rupture as illustrated in (D).

426    Scale bars: 20 μm.

A

Growth medium

*V. cholerae or P. aeruginosa*

Microfluidic chip

Hydrogel film

Coverslip

15 h

Hydrogel film

Biofilm

B

C

Biofilm

Hydrogel film with embedded fluorescent particles

■ *V. cholerae*

■ Hydrogel with fluorescent particles

D

■ *P. aeruginosa*

E

**Fig. 2: Biofilms deform soft substrates.** (A) Thin hydrogel films at the bottom surface of

microchannels reproduce the mechanical properties of host tissues. (B) Confocal

visualizations show that *V. cholerae* biofilms growing on hydrogels display large gaps at their

core. (C) *V. cholerae* biofilms formed at the surface of hydrogels containing fluorescent

particles. (D) *P. aeruginosa* biofilms similarly deform the soft substrates. Hydrogel elastic

modulus: (B and C) E= 12 kPa , (D and E) E = 38 kPa.  Scale bars: (C and D) 100 μm, (B

and E) 20 μm.

**Fig. 3: Biofilm buckle to deform their growth substrate.** (A) Morphological parameters

$\delta_{max}$ (maximum deformation amplitude) and $\lambda$ (half max full width) computed from resliced

deformation profiles. Dashed line indicates the baseline position of the gel surface. (B)

Timelapse visualization of deformation (reslice, bottom) with biofilm growth (brightfield, top).

Dashed lines indicate biofilm position and size on hydrogel profile. (C) Superimposition of

these profiles shows the rapid deformation and the emergence of a recess at biofilm edges

after 15 h. Each color corresponds to a different biofilm. (D) Biofilm age-dependence of

$\delta_{max}$. (E) The dependence of $\delta_{max}$ on biofilm diameter highlights a critical biofilm diameter

$d_c$ above which deformation occurs. (F) Hydrogel deformation field computed between 11 h

and 12 h of growth, superimposed with a brightfield image of the biofilm. Data represented

for the top right quarter of the biofilm shown in inset (dashed lines). (G) traction force

measurements at the deformation surface. The dashed line shows the edge of the biofilm.

(H) Deformation profiles on three hydrogels with different stiffness for biofilms of equal

diameter. (I) Biofilm diameter-dependence of maximum deformation for four different

hydrogel stiffnesses matching tissue properties. Scale bar: 10 µm for inset $t = 0$ h in (B), else

20 µm.

**Fig. 4: Biofilm mechanical properties and adhesion drive substrate deformation.** (A)

Deformations of hydrogel substrates by *V. cholerae* WT\*, $\Delta rbma$ and $\Delta bap1$ biofilms. (B)

Deformations of hydrogels by *P. aeruginosa* WT\* and $\Delta cdrA$ biofilms. (C) Dependence of

maximum deformations on *P. aeruginosa* WT\*, $\Delta cdrA$, $\Delta pel$ and $\Delta psl$ biofilm diameter. (D)

Increasing hydrogel stiffness to 200 kPa induces delamination of biofilms, as observed on

glass. (E) A model for biofilm deformation of soft substrates. Scale bars: 20 µm.

**Table 1.** Plasmids and strains used in this study

| Strain or plasmid | Relevant genotype | Source |
|---|---|---|
| pFY_113 | plasmid for generation of in-frame *rbmA* deletion mutants | [27] |
| pFY_330 | plasmid for generation of in-frame *bap1* deletion mutants | [27] |
| V. cholerae O1 El Tor A1552 (Vc WT*) | rugose variant | [28] |
| V. cholerae ΔrbmA | in frame deletion of Rbma in rugose backgrounds | This study |
| V. cholerae Δbap1 | in frame deletion of Bap1 in rugose backgrounds | This study |
| PAO1 WT | wild-type, Gm$^r$ | [29] |
| PAO1ΔwspF (PAO1 WT*) | in frame deletions of WspF, Gm$^r$ | [30] |
| PAO1ΔwspFΔpel | in frame deletions of WspF, PelA genes, Gm$^r$ | [30] |
| PAO1ΔwspFΔpsl | in frame deletions of WspF, PslBCD genes, Gm$^r$ | [30] |
| PAO1ΔwspFΔcdrA | in frame deletions of WspF, PslBCD, cdrA genes, Gm$^r$ | [31] |

472
473    **Table 2.** Molecular weight and concentrations of the precursors used for the generation of
474    the hydrogels and resulting elastic modulus

| Precursor | Concentration wt/vol | Modulus kPa |
|---|---|---|
| PEGDA MW 10000 (Biochempeg) | 10% | 12.1 ± 0.8 |
| PEGDA MW 6000 (Biochempeg) | 10% | 38.3 ± 1.0 |
| PEGDA MW 3400 (Biochempeg) | 10% | 30.9 ± 2.0 |
| PEGDA MW 700 (Sigma-Aldrich) | 15% | 203.3 ± 13.7 |