Extreme hot weather has stronger impacts on avian reproduction in forests
than in cities

Ivett Pipoly[1,2*], Bálint Preiszner[3], Krisztina Sándor[2,4], Csenge Sinkovics[2], Gábor Seress[1], Ernő
Vincze[1,2,5], Veronika Bókony[6,†], András Liker[1,2,†]

[1] MTA-PE Evolutionary Ecology Research Group, University of Pannonia, Veszprém, Hungary

[2] Behavioral Ecology Research Group, Center for Natural Science, Faculty of Engineering, University
of Pannonia, Veszprém, Hungary

[3] Fish and Conservation Ecology Research Group, Balaton Limnological Research Institute, Eötvös
Loránd Research Network, Tihany, Hungary

[4] MTA-ELTE Comparative Ethology Research Group, Budapest, Hungary

[5] Theoretical Population Ecology and Evolution Group, Department of Biology, Lund University, Lund,
Sweden

[6] Lendület Evolutionary Ecology Research Group, Plant Protection Institute, Centre for Agricultural
Research, Eötvös Loránd Research Network, Budapest, Hungary

[†] these two authors are joint last authors

* corrensponding author: Ivett Pipoly, pipoly.ivett@gmail.com, +3688 624773, and András
Liker, andras.liker@gmail.com. University of Pannonia, H-8200, Veszprém, Egyetem street
10. Building N 14, Hungary

Author Contributions: IP, VB, and AL designed the study. All authors collected data in the
field. IP, VB and AL did the statistical analyzes. IP and VB made the data visualization. IP,
VB and AL wrote the manuscript and all authors took part in finalizing it.

Preprint server: biorXiv, https://www.biorxiv.org/content/10.1101/2020.01.29.924332v1

Classification: Biological Sciences, Ecology, Population Biology

Competing interest statement: The authors declare no competing interests

Key words: climate change, heat waves, urban heat island, offspring size, offspring mortality

Abstract

Climate change and urbanization are among the most salient human-induced changes

affecting Earth's biota. Extreme weather events can have high biological impacts and are

becoming more frequent recently. In cities, the urban heat island can amplify the intensity and

frequency of hot weather events. However, the joint effects of heat events and urban

microclimate on wildlife are unclear, as urban populations may either suffer more from

increased heat stress or become adapted to warmer temperatures. Here we test whether the

effects of hot weather on reproductive success of great tits (*Parus major*) are exacerbated or

dampened in urban environments compared to forest habitats. By studying two urban and two

forest populations over six years, we show that 14-16 days-old nestlings have smaller body

mass and tarsus length, and suffer increased mortality when they experience a higher number

of hot days during the nestling period. The negative effects of hot weather on body mass and

survival are significantly stronger in forests than in urban areas, where these effects are

dampened or even reversed. These results suggest that urban birds are less vulnerable to

extreme hot weather conditions than their non-urban conspecifics, possibly by adaptively

evolving or flexibly adjusting nestling physiology to tolerate heat stress, and/or by adjusting

parental behavior in response to heat. This finding highlights that endothermic vertebrates

may be able to adapt to heat stress, which may help their populations cope with the joint

challenges of climate change and urbanization.

Significance statement

Extreme weather events are becoming more frequent due to climate change and can have

substantial effects on reproduction and survival of wild animals. Urban heat island can

amplify the frequency of extreme hot weather events, making it potentially more harmful for

city-dwelling organisms. Alternatively, urban populations living in warmer environments may

adapt to better tolerate heat. We investigated these alternatives by comparing nestling

development and survival between urban and forest great tit (*Parus major*) populations. We

found that urban populations are less vulnerable to heat: nestling body mass and survival

decreased rapidly with the increasing number of hot days in forests, while these effects were

dampened in urban broods. Thus, urban populations may become adapted to better tolerate

heat events.

Introduction

Current large-scale environmental changes are substantially influencing the ecological conditions for Earth's biota. Climate change is one of the dominant global processes that affect a wide range of organisms. Its effects include systematic changes in the long-term average meteorological conditions and also increases in the frequency of extreme weather events like heatwaves, droughts, and heavy storms [1–3]. These climatic changes are occurring in a world that is experiencing global-scale transformation of natural habitats into heavily modified anthropogenic habitats. For example, current estimates show that 77 % of the terrestrial habitats have already been modified by the direct effects of human activities such as agriculture and urbanization [4], and the remaining pristine habitats are also disappearing at a high rate [4–6] forcing an increasing number of wildlife populations to persist in anthropogenic environments [7]. Although the ecological and evolutionary consequences of different global processes such as climate change and anthropogenic land-use conversion are most often investigated separately, they obviously do not act independently. Thus to be able to better understand their current effects and to predict the changes likely induced in the future, we need to investigate their joint impacts in natural systems.

Hot days and heatwaves are among those extreme meteorological events that have become more frequent in the last few decades [2]. Extreme hot conditions can have strong biological impacts, affecting both survival and reproduction of organisms [8–10]. For example, the infamous heatwave that occurred in southern and western Europe in 2003 resulted in an estimated 70,000 heat-related human deaths [11] and was followed by detectable decreases in wild bird populations [12]. Similarly, extreme hot weather affects reproductive success negatively in a wide range of taxa, including both ectothermic [13, 14] and endothermic animals [10, 15–21]. These effects of extreme hot weather may be mediated by several, non-

exclusive mechanisms, including both direct effects of heat stress on physiological and

cognitive functioning [18, 20, 22] and indirect effects of heat *via* altered ecological conditions

like food or water availability [23, 24].

Anthropogenic habitat change, especially urbanization, is likely to interact with extreme

weather events because it fundamentally alters several basic environmental conditions,

including microclimate and food availability [7, 25]. Urban heat island (UHI) refers to the

generally higher ambient temperature in cities compared to the surrounding non-urban areas,

which is largely generated by heat storage in buildings and sealed roads. Its intensity can be as

high as + 5 °C (+ 9 °F) in some metropolitan areas [26–28], so organisms in cities experience

higher temperatures and more frequent heat events compared to organisms in non-urban areas

[28]. The UHI effect may interact with heat events in at least two ways. First, UHI can

exacerbate the biological effects of extreme heat events. For example, models show that

daytime temperature in temperate-zone cities during a heatwave can be higher by, on average,

2.8 °C as a result of the synergistic effect between UHI and heat events [29]. Moreover, heat

stress in cities might be even higher during a heat event than the sum of the background UHI

effect and the heatwave effect [30]. According to these model predictions, the number of hot

days is higher in urban than in non-urban areas with an additional increase in UHI resulting in

increased human mortality [30, 31]. Similarly, animals are more likely to reach their upper

limit of thermal tolerance in cities during heat events than in non-urban areas [32]. Thus,

urban populations may suffer a stronger reduction in survival and reproductive success than

non-urban populations due to hot days ('superimposed heat and UHI effects' hypothesis).

Second, it is also possible that during long-term exposure to UHI, urban populations

become adapted to higher temperatures ('adaptation to UHI' hypothesis). For example, heat

tolerance, expressed as the thermal maximum where animals become unable to coordinate

their motor performance, is hihger in urban compared to rural populations of acorn ants

(*Temnothorax curvispinosus*) [33]*,* water fleas (*Daphnia magna*)[34], and crested anoles

(*Anolis cristatellus*) [35]. Moreover, survival and production of winged individuals in acorn

ant colonies in an urban-rural reciprocal transplant experiment were higher in the home

environment for both urban and rural colonies, probably due to local thermal adaptations [33,

36]. The reproductive advantage of heat-tolerant or heat-adapted phenotypes in hot urban

conditions is also documented in the scale insect *Parthenolecanium quercifex*, as individuals

collected from warm urban environments reproduce at a higher rate under hot than under cool

laboratory conditions [37]. Although there is much less data on urban thermal adaptations in

endotherms, the smaller body size of the rodent *Peromyscus maniculatus* [38] and reduced

number of feathers in some birds [39] in urban populations might also reflect adaptations to

the UHI.

So far, only a few studies have tested the above hypotheses for joint effects of extreme hot

weather events and habitat urbanization. Their results show either additional mortality in

cities in humans [30] or adaptation to UHI effects in water fleas [34]. Importantly, there is no

field study for endotherm animal populations, despite that these are the most frequent targets

of conservation efforts and play crucial roles in urban ecosystems. To start to fill this

knowledge gap, here we tested whether heat events influence avian breeding success

differently in cities and forest habitats. We analyzed six years of breeding biology data of two

urban and two forest populations of great tits (*Parus major*), a small songbird that frequently

breeds in both urban and forest habitats. In this species, similarly to several other birds,

reproductive performance is typically lower in urban sites than in more natural areas [40, 41].

In the current study, we measured air temperature hourly at each site with own weather

stations to calculate the number of hot days for each brood's pre-fledging nestling period, and

tested whether nestlings' body mass, tarsus length, and survival are related to the number of

hot days differently in urban and forest sites. According to the 'superimposed heat and UHI

effects' hypothesis, we expect that heat events would result in stronger negative impacts on reproduction in urban than forest habitats. Alternatively, the 'adaptation to UHI' hypothesis predicts that heat events would have reduced impact in urban than forest populations.

Results

Habitat differences in heat events

Great tit offspring experienced markedly different temperature characteristics in urban and forest sites during the nestling season (Figure S1). Both the number of hot days (Figure S1A; urban-forest difference on log-odds scale: mean ± SE: 0.99 ± 0.13, t=7.39, df=750, p < 0.001) and the average temperatures within the nestling periods were significantly higher in the urban habitat (Figure S1B; urban-forest difference: 1.52 ± 0.27 °C, t=5.74, df=751, p < 0.001; model estimates controlled for year effects). That means the odds of hot days occurring during the nestling period was 2.7 times greater in the cities than in the forests: at least one hot day occurred in 179 out of 390 (45.9%) urban broods and in 77 out of 370 (20.8 %) forest broods. The highest maximum temperature was 40.1 °C in urban nestling periods (in June 2013) and 33.6 °C in forest nestling periods (in May 2014).

Effects of hot days on reproductive success

Nestling body mass

In the simple model where predictors were the number of hot days, study site and their two-way interaction, the effect of hot days on nestling mass differed between sites (as indicated by a significant interaction between sites and number of hot days, Table 1) and between urban and forest habitats (according to the linear contrast post-hoc test comparing the effects between the two urban and the two forest populations, Table 2): increase in the number of hot days negatively affected nestling mass in forest broods while no such trend was

present for the urban broods (Figure 1A). This result is consistent with the site-specific trends:

the two forest populations had the most negative slopes for the relationship of nestling mass

with the number of hot days, whereas the slope was flat or even positive in the two urban

populations (Table S4).

We also tested whether the effect of hot days persist when further potentially important

variables (i.e. hatching date, year, brood size an brood age) are taken into account. We built a

set of models containing various combinations of these variables in addition to the number of

hot days, study site and their interaction, and we compared these models using the

information-theoretic approach based on AICc. Model selection for these multi-predictor

models resulted in two supported models for nestling mass (ΔAICc < 2; the model with the

third best fit had ΔAICc = 4; Table S1). The first supported model included year, brood age,

and brood size besides the number of hot days, study sites, and their interaction; the second

supported model contained the same predictors except brood size. Linear contrasts calculated

from each of these supported multi-predictor models showed a significant overall habitat

difference (Table 2) with a more negative effect of hot days in forest compared to urban

populations (Table S4), corroborating the results of the simple model.

Nestling tarsus length

In the simple model, the interaction between site and number of hot days was significant

(Table 1), due to the difference between the two urban sites, with a positive slope in one site

and negative in the other (Table S4). The two forest sites both showed negative relationship

between the number of hot days and nestling tarsus length (Table S4). The linear contrast for

urban-forest difference suggested no systematic habitat difference in the effect of hot days on

the tarsus length of nestlings (Table 2, Figure 1B).

Model selection for the multi-predictor model set resulted in a single supported model,

which contained the number of hot days, study site, their interaction, the average temperature

of the nestling period, brood age, and brood size (for the other models ΔAICc ≥ 6.68, Table

S2). The result of linear contrast comparing urban and forest slopes from this model was

qualitatively identical to the result from the simple model, showing no consistent habitat

difference in the effect of hot days on nestling tarsus length (Table 2, Table S4).

Nestling mortality

The simple model showed that the effect of the number of hot days on nestling mortality

differed significantly between study sites (Table 1). The slope of the relationship between

nestling mortality and the number of hot days was positive in all populations, being highest in

one of the forest sites and lowest in one of the urban sites (Table S4). The slopes also differed

between urban and forest populations with a stronger effect in the latter (Table 2, Figure 1C),

i.e. nestling mortality increased significantly more with the number of hot days in the forests

than in the cities.

Multi-predictor model selection resulted in a single supported model, which was the full

model of our model set (for the other models ΔAICc ≥ 42.87, Table S3), containing the

number of hot days, study site and their interaction, the average temperature of the nestling

period, year, hatching date, and the quadratic term of hatching date. The linear contrast

calculated from this model showed a significantly steeper slope in forests compared to urban

sites (Table 2), corroborating again the result of the simple model.

Discussion

Our study found clear differences between urban and non-urban populations in the

effects of hot days on important fitness components. Although the influence of thermal

environment on fitness has been shown to vary with habitat urbanization in ectotherms [36],

our study is the first to demonstrate such habitat-dependent responses to heat stress in an

endothermic animal. While UHI caused a 1.19 °C higher temperature on average and there

were more hot days in the cities compared to forests (Figure S1), hot days affected nestling

mass and survival more negatively in the forest habitat type. These results are robust, as they

are based on >600 broods from 6 years and four sites, and were qualitatively unchanged by

taking into account several potentially confounding factors that may differ between urban and

forest habitats, such as the start of breeding and brood size. These findings highlight that both

lethal and sublethal effects of heat events (i.e. the proportion of surviving offspring and their

quality in terms of body mass) can differ between urban and non-urban habitats. Although

information about sublethal fitness costs of hot weather is scarce [9], developing under hot

weather conditions may have long-term consequences for fitness. Our results thus support the

'adaptation to UHI' hypothesis and suggest that non-urban populations of birds (and possibly

other vertebrates) are more vulnerable to extreme heat than urban populations.

Heat may affect the reproductive output of birds in at least two ways: directly through

offspring physiology and indirectly through food availability. Firstly, the negative effects of

hot days on offspring mass and survival may emerge due to the direct physiological

consequences of heat stress. Nestlings usually cannot maintain stable body temperature in the

first few days of their lives, and their metabolic processes also differ from those of adults [42]

leaving them potentially more vulnerable to extreme heat than adults. The heat responses of

dependent offspring are poorly known, although in birds the nestling phase seems to be the

most relevant period during ontogeny regarding heat effects compared to the incubation and

post-fledging periods [43]. In hot environments, individuals can less effectively dissipate the

excess heat as a consequence of increased metabolic rate, leading to hyperthermia. Increasing

metabolism causes decreased utilization of food and faster mobilization of energy reserves,

thus it may cause lower body mass. Additionally, during heat stress, evaporative water loss is

elevated as the organism tries to cool itself by evaporation to maintain body temperature in

the physiologically normal range, causing dehydration [44, 45]. Thus, increased metabolic

rate and water loss both can lead to decreased body mass, as it has been shown in several bird

species [16, 46–48].

These direct physiological effects of heat might be better tolerated by urban animals due

to changes in their physiological and morphological characteristics. For example, we found in

the same study populations that urban great tit nestlings have fewer feathers and increased

bare body surfaces than forest nestlings [39], which may facilitate heat dissipation. Reduced

body size may also help to cope with the heat in cities because animals with smaller size have

a higher surface-biomass ratio which facilitates heat loss. Bergmann's rule [49] predicts that

reduced size is beneficial against dehydration and overheating in warmer environments [44]

(but see [50]) such as cities. This idea is supported by urban-rural comparisons of invertebrate

communities [51], as well as by a study of water flea populations which found that urban

individuals were generally smaller than rural ones, and smaller individuals were more heat

tolerant [34]. In birds, a study of white-browed scrubwrens (*Sericornis frontalis*) also found

that smaller individuals survived better when more extreme hot and dry events occurred [52].

Urban birds are usually smaller than their non-urban conspecifics [53–55] and this is also the

case with the great tits in our study populations [40]. Although experimental evidence shows

that low availability of nestling food is a major cause of the reduced size of urban great tit

nestlings [56], selection for better heat tolerance might also contribute to it. Finally, life-

history changes may also help animals to cope with the direct effects of urban heat. For

example, urban birds usually have lower clutch size than forest birds [40, 53] which may also

be advantageous for heat dissipation when environmental temperature is high because fewer

offspring produce less heat and have more space to maintain distance in the nest, which helps

better heat conduction.

Secondly, indirect effects may also contribute to lower body mass and higher mortality in

response to heat events. For example, hot weather may reduce food availability for

insectivorous nestlings by affecting their parents and/or their prey. Caterpillars are the main

source of nestling diet in great tits and many other birds, and increasing temperature strongly

decreases the time to pupation in several lepidopteran species [57–59], so optimal caterpillar

food may be available for a shorter time in hot periods. Moreover, the growth of caterpillars

declines rapidly above a critical temperature [58] and their mortality increases when

temperature is constantly high [59, 60]. These heat effects may not only result in less food for

nestlings but may also contribute to dehydration because caterpillars are the most water-rich

items in the diet of nestlings [61]. These effects might be stronger in forests where the amount

of available caterpillar prey is much higher in general [40] and the vast majority of nestling

diet consists of caterpillars. In such environments, it might be more difficult for parent birds

to compensate for a reduction in caterpillar biomass compared to cities, where nestlings are

usually fed a greater proportion of other food types [62, 63]. Additionally, the foraging

activity of parents may also decrease when temperature is high [16, 64, 65], resulting in less

food delivered to the offspring, and this effect might also be diminished in urban populations

if parent birds have adapted to tolerate heat better (see above). Further studies are required to

test whether heat affects parental care differentially in urban and non-urban habitats.

Our study suggests that the increasing frequency of hot days expected as a consequence

of ongoing climate change would be more harmful for natural than urban populations, further

endangering biodiversity. For example, our results predict that the reproductive advantage of

non-urban populations (e.g. higher body mass and lower mortality of nestlings) over urban

populations can be reversed when the number of hot days experienced by nestlings becomes

high (exceeds 7-8 days in our study). This raises the intriguing possibility that urban

populations might serve as sources for the conservation of some species in the future.

Adaptation to constantly warmer urban environments may be achieved by microevolution,

epigenetic changes, and/or phenotypic plasticity [66]. In some ectotherm species, artificial

thermal selection and common garden experiments have revealed that one or more of these

mechanisms can play a role in the increased upper thermal tolerance of urban populations

[34–36, 67]. To our knowledge, no such experiments have yet been done to identify the

mechanisms of higher thermal tolerance in urbanized birds or other endothermic animals,

which is an important knowledge gap in our understanding of urban adaptations. Therefore,

there is a great need for further studies that explore the differences in thermal tolerance

between different habitats to understand how wild animals are affected by, and may adapt to,

the interacting challenges of climate change and urbanization in the Anthropocene.

Methods

Data collection

We installed nest-boxes for great tits in an urban site (city of Veszprém 47°05'17"N,

17°54'29"E), and in two sites in natural habitats (forests near Vilma-puszta 47°05'06.7"N,

17°51'51.4"E and Szentgál 47°06'39"N, 17°41'17"E) in 2012, and additionally in another

urban site (Balatonfüred 46°57'30"N, 17°53'34"E) in 2013. Urban nest-boxes are located

mostly in public parks, university campuses, and a cemetery, where vegetation contains both

native and introduced plant species. Forest study sites are located in deciduous woodlands,

characterized by beech *Fagus sylvatica* and hornbeam *Carpinus betulus* (in Szengtál) or

downy oak *Quercus cerris* and South European flowering ash *Fraxinus ornus* (in Vilma-

puszta). See detailed description of study sites in [40]. In March 2013, we installed a WH

2080 weather station (Ambient, LLC, AZ, USA) at each study site that recorded hourly

temperature (°C) data throughout the six years of the study.

Great tits usually raise one or two broods per breeding season. We collected data on all

breeding attempts at each site from 2013 to 2018, by recording the number of eggs and

nestlings in the nest-boxes every 3-4 days from March to the end of July. We captured parent

birds using a nest-box trap 6-15 days after their first nestling had hatched. We determined

parents' sex based on their plumage characteristics and banded each bird with a unique

combination of a numbered metal band and three plastic color bands. Breeding adult birds that

had been color-banded on previous occasions were identified from recordings made during

the nestling period by using a small, concealed video camera attached to the nest-boxes [68].

In these video samples, we considered a color-banded individual to be a parent bird if it was

recorded to enter the nest-box with food at least once. Close to fledging (at day 14-16 post-

hatch; day 1 being the hatching day of the first nestling in the brood), we measured nestlings'

body mass with a Pesola spring balance (± 0.1 g) and tarsus length with a Vernier caliper (±

0.1 mm).

All procedures applied during our study were in accordance with the guidelines for

animal handling outlined by ASAB/ABS (www.asab.org) and Hungarian laws. We have all

the required permissions for capturing, measuring of the birds and monitoring their breeding

from the Government Office of Veszprém County, Nature Conservation Division (former

Middle Transdanubian Inspectorate for Environmental Protection, Natural Protection and

Water Management; permission number: 24861/2014 and VE-09Z/03454-8/2018) and from

the National Scientific Ethical Committee on Animal Experimentation (permission number:

VE/21/00480-9/2019).

Data processing

We used both the first and second annual broods of each pair in the study. We omitted

broods where we could not precisely identify the nestling period due to unknown dates of

hatching and/or the death of the entire brood before hatching, and also those broods where we

had gaps in the temperature data for more than two days during the nestling period (n= 190

broods). For analyzing the effects of weather on nestling size, we used the average body mass

and the average tarsus length of each brood as response variables, and we included broods

where at least one offspring was alive at the time of nestlings' measuring and banding (i.e. the

age of 14-16 days). Because chick size varies with age, we omitted those few broods where

the nestlings were measured before or after 14-16 days of nestling age (n= 30); thus we had

n= 674 broods for nestling size analyzes.

For analyzing the effects of weather on nestling mortality, we used broods in which at

least one offspring was alive on the third day after the hatching of the first nestling. We

excluded a small number of broods (n= 7) that failed within the first three days after hatching

because the average interval between our nest monitoring visits was three days, so estimating

the number of hatched chicks and their survival before the first nest check after hatching

would be very uncertain. We also excluded those broods where chick mortality was likely

unrelated to weather conditions (n= 7), i.e. occurred due to predation or human disturbance

(i.e. recorded during the next observation after the occasion of capturing the parent bird on its

nest). This yielded n= 760 broods for nestling mortality analyzes, from which 385 had no

chick mortality, 319 had partial mortality, and 56 had complete brood loss during the nestling

period.

We calculated all weather variables (average temperature and number of hot days,

detailed below) for the nestling period of each brood individually. For investigating nestling

size, weather variables were calculated for days from the day of hatching of the first chick to

the day preceding the day of banding and measuring the nestlings. This sample included only

broods where banding occurred at age 14-16 days (mean ± SE: 14.04 ± 0.03). For analyzing

nestling mortality, we calculated weather variables for the period from the day of hatching of

the first offspring to either the day preceding the day of recording the last nestling's death or

the day preceding the day of banding and measuring the nestlings (mean ± SE: 13.72 ± 0.07,

range: 3 – 18 days). We calculated the average temperature as the mean of hourly

temperatures over the nestling period. There are several methods for quantifying extreme hot

weather [69], using various thresholds from 21 to 43 °C (70 to 110 °F) depending on the

geographic location. We calculated the number of hot days during each nestling period using

a threshold specific to our region. To calculate a threshold for hot days, first, we defined a

reference period from the earliest hatching date to the latest chick banding date recorded in

our six-year data set (pooling all sites and years), which ranged from 9th April to 15th July.

Then, to estimate the typical temperatures in our geographic area during this reference period,

we used a 26-years dataset from an external reference weather station located in

Szentkirályszabadja, a small village near Veszprém (47°57'06" É, 17°58'10" K, ca. 9.5 – 22

km from our study sites). This weather station is maintained by the Hungarian Meteorological

Service and its temperature data are available from the National Oceanic and Atmospheric

Administration database (www.ncdc.noaa.gov), with records of air temperature every three

hours a day (0, 3, 6, 12, 15, 18, 21 UTC) since 1993. Using this temperature dataset, we

calculated the 90 % percentile of all daily maximum temperatures for the reference period

from this 26-year long dataset, which was 28.7 °C. Finally, we calculated the number of hot

days as the number of days when the daily maximum temperature was higher than the 28.7 °C

threshold during the nestling period for each brood. Fixed temperature thresholds for

calculating hot days are widely used in ecological and evolutionary research [69], based on

the assumption of a fixed threshold for heat tolerance in adults and offspring [18, 19, 52, 70].

Analyzing the effects of extreme weather is challenging. Extreme events are rare by definition, so their distribution is strongly skewed. Furthermore, weather variables are often correlated with each other and with other variables influencing reproductive success (e.g. with calendar date), so multicollinearity can be a problem in models containing multiple predictors. Statistical methods that can handle multicollinearity, such as covariance-based structural equation modelling, are less well suited for handling non-normal residual distributions and the non-independence structure of ecological data (e.g. multiple broods per pair). Therefore, here we used general and generalized linear mixed-effects (LME and GLMM) models and Akaike's information-criterion (AIC) based model selection [71] to focus on the interaction between habitat type and the number of hot days in explaining reproductive variables. Mixed-effects models can be applied appropriately to non-normal and non-independent data, while model selection based on the models' total explanatory power (model fit) is not sensitive to multicollinearity [72].

All analyzes were run in R (version 4.0.0) [73]. First, to examine the temperature characteristics of our study sites, we compared the number of hot days and the average temperature in the nestling periods (calculated separately for each brood) between urban and forest habitats. We constructed a generalized linear model with generalized Poisson distribution for the number of hot days using "glmmTMB" package [74] to handle overdispersion, and a linear model for the average temperatures. In both models, study site (4 sites) and year (6 years) were the predictors. To statistically compare the mean of each meteorological variable between the two habitat types, we calculated marginal means from the models for each study site, then calculated a linear contrast for habitat comparison as the difference between the average of the two urban sites versus the average of the two forest sites, using the "emmeans" R package [75]. We used this approach rather than including

habitat type as a fixed effect and site as a random effect in the models because variance

estimations of random effects with few levels are unreliable [76, 77], whereas including both

habitat type and site as fixed effects would have resulted in strong collinearity between these

two factors [78]. Instead, we treated the four sites as if they were two control groups and two

treatment groups in an experiment, and we used a pre-planned comparison to test the

prediction that the two treatment (i.e. urban) groups would differ from the two control (i.e.

forest) groups. Such pre-planned comparisons are a powerful approach for testing *a priori*

hypotheses [79].

To analyze average nestling mass and average tarsus length as response variables, we

used the "lme" function of the "nlme" R package [80] which assumes Gaussian error. For

nestling mortality (proportion of hatched chicks that died by the age of 14-16 days) as the

response variable, we used GLMM models using the "glmer" function of the "lme4" R

package [81] with binomial error distribution and logit link function. All of our models

contained pair identity as a random factor to control for the non-independence of broods

produced by the same pair. Broods got the same pair ID if both the female and the male

parents were the same, and a new pair ID was assigned to the brood when it had at least one

different parent. There were 111 pairs of parents that had more than one brood in our dataset,

ranging 2-6 broods/pair. Additionally, to handle overdispersion in models of nestling

mortality, we used an observation-level random effect [82] and ran the models with the

BOBYQA optimizer [81].

First, we built a simple model for each of the three response variables (nestling mass,

tarsus length, and mortality) including only the number of hot days (used as a numeric

covariate) and study site (used as a factor with four levels), and their two-way interaction as

predictors. Then, to statistically compare the effect of hot days on nestling size and mortality

between the two habitat types, we calculated the slopes of the relationship between

reproductive success measurements and the number of hot days for each study population,

and we compared the average of the two urban slopes with the average of the two forest

slopes using a linear contrast.

To infer the robustness of the results from the simple models, we also built multi-

predictor models which included further potentially important predictor variables. In all

models of our three response variables, we included year as a categorical variable, the average

temperature during the nestling period and hatching date as numeric covariates, the latter

defined as the number of days elapsed from the 1st January annually to the hatching of the first

nestling in each brood. We also added the quadratic term of hatching date, because a

preliminary inspection of diagnostic graphs suggested the possibility of non-linear seasonal

changes in the response variables. For models of nestling size (i.e. average body mass and

average tarsus length of nestlings), we additionally included brood age (i.e. the number of

days elapsed from the hatching of the first chick to the day of nestlings' measuring, ranging

from 14 to 16 days), and brood size (i.e. the number of offspring alive at measuring). We

standardized all numeric predictors using z-transformation for our multi-predictor analyzes to

avoid model convergence problems due to different scales of variables. Variance inflation

factor (VIF), a measure of multi-collinearity, ranged from 1.04 to 6.24 for the predictors in

these multi-variate models. Because multi-collinearity with VIF values > ca. 2 can lead to

unreliable standard errors, non-significant parameter estimates from such models must be

treated with caution [72]. However, the set of predictors that yields the best explanatory

power can be correctly identified by model selection based on model fit statistics (like AIC),

even if multi-collinearity is present [71]. Therefore, we created a candidate model set for each

response variable using all combinations of predictors but always including the number of hot

days, study site, and their interaction (because this interaction was the focus of our study, and

our goal with the model-selection procedure was to see if this interaction yields the same

habitat contrasts in the multi-predictor models as alone in the simple models). The model set

contained 33 models for nestling size (Table S1-S2) and 9 models for nestling mortality

(Table S3). We compared the models in each model set based on their AICc (AIC corrected

for sample size) to identify the most supported model(s) for each response variable [71] using

the "model.avg" function of the "MuMIn" R package [83]. From the model(s) with superior

support based on ΔAICc and Akaike weight, we calculated the slope for each study site and

the linear contrast that compares the effect of hot days between urban and forest sites the

same way as from the simple models.

Acknowledgements

We thank several field assistants for helping during fieldwork. The project was primarily

financed by the National Research, Development and Innovation Office of Hungary (NKFIH,

grants K84132, K112838 & K132490 to AL), and by the Hungarian Academy of Sciences. It

was also supported by the TKP2020-IKA-07 project financed under the 2020-4.1.1-TKP2020

Thematic Excellence Programme by NKFIH, and by the European Union with the co-funding

of the European Social Fund (TÁMOP-4.2.2.A-11/1/KONV-2012-0064). I.P. was supported

by the ÚNKP-20-4 New National Excellence Programme of the Ministry of Innovation and

Technology from the Source of the National Research, Development and Innovation Fund.

V.B. was supported by the János Bolyai Scholarship of the Hungarian Academy of Sciences.

EV was supported by the PD-134985 grant by NKFIH and by the MSCA EF Seal of

Excellence IF-2019 grant by Vinnova, the Swedish Governmental Agency for Innovation

Systems (grant number: 2021-01102). K.S. was supported by MTA-ELTE Comparative

Ethology Research Group (F01/031).

References

1. Perkins-Kirkpatrick SE, Lewis SC (2020) Increasing trends in regional heatwaves. Nat Commun 11:1–8. doi: 10.1038/s41467-020-16970-7

2. Drumond A, Margarida LR, Reboita MS, Taschetto AS (2020) Weather and climate extremes: Current developments. Atmosphere (Basel) 11:11–14. doi: 10.3390/ATMOS11010024

3. Buckley LB, Huey RB (2016) Temperature extremes: geographic patterns, recent changes, and implications for organismal vulnerabilities. Glob Chang Biol 22:3829–3842. doi: 10.1111/gcb.13313

4. Watson JEM, Allan JR, Venter O, et al. (2018) Protect the last of the wild. Nature 563:27–30.

5. Watson JEM, Shanahan DF, Di Marco M, et al. (2016) Catastrophic Declines in Wilderness Areas Undermine Global Environment Targets. Curr Biol 26:2929–2934. doi: 10.1016/j.cub.2016.08.049

6. Di Marco M, Ferrier S, Harwood TD, et al. (2019) Wilderness areas halve the extinction risk of terrestrial biodiversity. Nature 573:582–585. doi: 10.1038/s41586-019-1567-7

7. Alberti M, Marzluff JM (2004) Ecological resilience in urban ecosystems: Linking urban patterns to human and ecological functions. Urban Ecosyst 7:241–265. doi: 10.1023/B

8. Mora C, Dousset B, Caldwell IR, et al. (2017) Global risk of deadly heat. Nat Clim Chang 7:501–506. doi: 10.1038/nclimate3322

9. Conradie SR, Woodborne SM, Cunningham SJ, McKechnie AE (2019) Chronic, sublethal effects of high temperatures will cause severe declines in southern African arid-zone birds during the 21st century. Proc Natl Acad Sci U S A 116:14065–14070. doi: 10.1073/pnas.1821312116

10. Woodroffe R, Groom R, McNutt JW (2017) Hot dogs: High ambient temperatures impact

reproductive success in a tropical carnivore. J Anim Ecol 86:1329–1338. doi:

10.1111/1365-2656.12719

11. Robine JM, Cheung SLK, Le Roy S, et al. (2008) Death toll exceeded 70,000 in Europe

during the summer of 2003. Comptes Rendus - Biol 331:171–178. doi:

10.1016/j.crvi.2007.12.001

12. Jiguet F, Julliard R, Thomas CD, et al. (2006) Thermal range predicts bird population

resilience to extreme high temperatures. Ecol Lett 9:1321–1330. doi: 10.1111/j.1461-

0248.2006.00986.x

13. Zizzari ZV, Ellers J (2011) Effects of exposure to short-term heat stress on male

reproductive fitness in a soil arthropod. J Insect Physiol 57:421–426. doi:

10.1016/j.jinsphys.2011.01.002

14. Dayananda B, Murray BR, Webb JK (2017) Hotter nests produce hatchling lizards with

lower thermal tolerance. J Exp Biol 220:2159–2165. doi: 10.1242/jeb.152272

15. Bourne AR, Cunningham SJ, Nupen LJ, et al. (2021) No sex-specific differences in the

influence of high air temperatures during early development on nestling mass and

fledgling survival in the Southern Pied Babbler (Turdoides bicolor). Ibis (Lond 1859).

doi: 10.1111/ibi.12990

16. Oswald KN, Smit B, Lee ATK, et al. (2021) Higher temperatures are associated with

reduced nestling body condition in a range-restricted mountain bird. J Avian Biol. doi:

10.1111/jav.02756

17. Conrey RY, Skagen SK, Yackel Adams AA, Panjabi AO (2016) Extremes of heat,

drought and precipitation depress reproductive performance in shortgrass prairie

passerines. Ibis (Lond 1859) 158:614–629. doi: 10.1111/ibi.12373

18. McCowan LSC, Griffith SC (2021) Baked eggs: catastrophic heatwave induced

reproductive failure in the desert-adapted Zebra Finch (Taeniopygia guttata). Ibis (Lond.

1859). 163:

19. Marrot P, Garant D, Charmantier A (2017) Multiple extreme climatic events strengthen

selection for earlier breeding in a wild passerine. Philos Trans R Soc L B Biol Sci. doi:

http://dx.doi.org/10.1098/rstb.2016.0372

20. Welbergen JA, Klose SM, Markus N, Eby P (2008) Climate change and the effects of

temperature extremes on Australian flying-foxes. Proc R Soc B Biol Sci 275:419–425.

doi: 10.1098/rspb.2007.1385

21. Van de Ven TMFN, Fuller A, Clutton-Brock TH (2020) Effects of climate change on pup

growth and survival in a cooperative mammal, the meerkat. Funct Ecol 34:194–202. doi:

10.1111/1365-2435.13468

22. Danner RM, Coomes CM, Derryberry EP (2021) Simulated heat waves reduce cognitive

and motor performance of an endotherm. Ecol Evol 11:2261–2272. doi:

10.1002/ece3.7194

23. du Plessis KL, Martin RO, Hockey PAR, et al. (2012) The costs of keeping cool in a

warming world: Implications of high temperatures for foraging, thermoregulation and

body condition of an arid-zone bird. Glob Chang Biol 18:3063–3070. doi:

10.1111/j.1365-2486.2012.02778.x

24. Fuller A, Mitchell D, Maloney SK, Hetem RS (2016) Towards a mechanistic

understanding of the responses of large terrestrial mammals to heat and aridity associated

with climate change. Clim Chang Responses 3:1–19. doi: 10.1186/s40665-016-0024-1

25. Seress G, Liker A (2015) Habitat urbanization and its effects on birds. Acta Zool Acad Sci

Hungaricae 61:373–408. doi: 10.17109/AZH.61.4.373.2015

26. Landsberg HE (1981) The Urban Climate. Acedemic Press, New York

27. Zhao L, Lee X, Smith RB, Oleson K (2014) Strong contributions of local background

climate to urban heat islands. Nature 511:216–219. doi: 10.1038/nature13462

28. Central, Climate (2021) Hot Zones: Urban heat islands. Res Br by Clim Cent 10. doi:

10.4324/9781315765853-11

29. Zhao L, Oppenheimer M, Zhu Q, et al. (2018) Interactions between urban heat islands and

heat waves. Environ Res Lett. doi: 10.1088/1748-9326/aa9f73

30. Li D, Bou-Zeid E (2013) Synergistic interactions between urban heat islands and heat

waves: The impact in cities is larger than the sum of its parts. J Appl Meteorol Climatol

52:2051–2064. doi: 10.1175/JAMC-D-13-02.1

31. Rizvi SH, Alam K, Iqbal MJ (2019) Spatio -temporal variations in urban heat island and

its interaction with heat wave. J Atmos Solar-Terrestrial Phys 185:50–57. doi:

10.1016/j.jastp.2019.02.001

32. Krause JS, Chmura HE, Wingfield JC, et al. (2017) How birds cope physiologically and

behaviourally with extreme climatic events. Philos Trans R Soc B Biol Sci

372:20160140. doi: 10.1098/rstb.2016.0140

33. Diamond SE, Chick L, Perez ABE, et al. (2017) Rapid evolution of ant thermal tolerance

across an urban-rural temperature cline. Biol J Linn Soc 121:248–257.

34. Brans K, Jansen M, Vanoverbeke J, et al. (2017) The heat is on: genetic adaptation to

urbanization mediated by thermal tolerance and body size. Glob Chang Biol 23:5218–

5227. doi: 10.1111/gcb.13784

35. Campbell-Staton SC, Winchell KM, Rochette NC, et al. (2020) Parallel selection on

thermal physiology facilitates repeated adaptation of city lizards to urban heat islands.

Nat Ecol Evol 4:652–658. doi: 10.1038/s41559-020-1131-8

36. Martin RA, Chick LD, Garvin ML, Diamond SE (2021) In a nutshell, a reciprocal

transplant experiment reveals local adaptation and fitness trade-offs in response to urban

evolution in an acorn-dwelling ant. Evolution (N Y) 75:876–887. doi:

10.1111/evo.14191

37. Meineke EK, Dunn RR, Sexton JO, Frank SD (2013) Urban Warming Drives Insect Pest
582   Abundance on Street Trees. PLoS One 8:2–8. doi: 10.1371/journal.pone.0059687

583   38. Guralnick R, Hantak MM, Li D, McLean BS (2020) Body size trends in response to
584   climate and urbanization in the widespread North American deer mouse, Peromyscus
585   maniculatus. Sci Rep 10:1–13. doi: 10.1038/s41598-020-65755-x

586   39. Sándor K, Liker A, Sinkovics C, et al. (2021) Urban nestlings have reduced number of
587   feathers in Great Tits (Parus major). Ibis (Lond 1859) 163:1369–1378. doi:
588   10.1111/ibi.12948

589   40. Seress G, Hammer T, Bókony V, et al. (2018) Impact of urbanization on abundance and
590   phenology of caterpillars and consequences for breeding in an insectivorous bird. Ecol
591   Appl 28:1143–1156.

592   41. Bailly J, Scheifler R, Berthe S, et al. (2016) From eggs to fledging: negative impact of
593   urban habitat on reproduction in two tit species. J Ornithol 157:377–392. doi:
594   10.1007/s10336-015-1293-3

595   42. Price ER, Dzialowski EM (2018) Development of endothermy in birds: patterns and
596   mechanisms. J Comp Physiol B Biochem Syst Environ Physiol 188:373–391. doi:
597   10.1007/s00360-017-1135-0

598   43. Marques-Santos F, Dingemanse NJ (2020) Weather effects on nestling survival of great
599   tits vary according to the developmental stage. J Avian Biol. doi: 10.1111/jav.02421

600   44. Weathers WW (1972) Physiological thermoregulation in heat-stressed birds:
601   consequences of body sizes. Physiol Zool 54:345–361.

602   45. Arad Z, Horowitz M, Eylath U, Marder J (1989) Osmoregulation and body fluid
603   compartmentalization in dehydrated heat-exposed pigeons. Am J Physiol - Regul Integr
604   Comp Physiol 257:377–382. doi: 10.1152/ajpregu.1989.257.4.R732

46. Quinteiro-Filho WM, Ribeiro A, Ferraz-de-Paula V, et al. (2010) Heat stress impairs

performance parameters , induces intestinal injury , and decreases macrophage activity in

broiler chickens. Poult Sci 89:1905–1914. doi: 10.3382/ps.2010-00812

47. Kruuk LEB, Osmond HL, Cockburn A (2015) Contrasting effects of climate on juvenile

body size in a Southern Hemisphere passerine bird. Glob Chang Biol 21:2929–2941. doi:

10.1111/gcb.12926

48. Corregidor-Castro A, Jones OR (2021) The effect of nest temperature on growth and

survival in juvenile Great Tits Parus major. Ecol Evol 11:7346–7353. doi:

10.1002/ece3.7565

49. Bergmann C (1847) Ueber die verhältnisse der wärmeökonomie der thiere zu ihrer grösse.

Gottinger Stud 3:595–708.

50. Whitfield MC, Smit B, Mckechnie AE, Wolf BO (2015) Avian thermoregulation in the

heat : scaling of heat tolerance and evaporative cooling capacity in three southern

African arid-zone passerines. J Exp Biol 218:1705–1714. doi: 10.1242/jeb.121749

51. Merckx T, Souffreau C, Kaiser A, et al. (2018) Body-size shifts in aquatic and terrestrial

urban communities. Nature 558:113–116. doi: 10.1038/s41586-018-0140-0

52. Gardner JL, Rowley E, De Rebeira P, et al. (2017) Effects of extreme weather on two

sympatric australian passerine bird species. Philos Trans R Soc B Biol Sci. doi:

10.1098/rstb.2016.0148

53. Chamberlain DE, Cannon AR, Toms MP, et al. (2009) Avian productivity in urban

landscapes: A review and meta-analysis. Ibis (Lond 1859) 151:1–18. doi:

10.1111/j.1474-919X.2008.00899.x

54. Liker A, Papp Z, Bókony V, Lendvai ÁZ (2008) Lean birds in the city: Body size and

condition of house sparrows along the urbanization gradient. J Anim Ecol 77:789–795.

doi: 10.1111/j.1365-2656.2008.01402.x

55. Caizergues AE, Charmantier A, Lambrechts MM, et al. (2021) An avian urban

morphotype: how the city environment shapes great tit morphology at different life
stages. Urban Ecosyst 1–15. doi: 10.1007/s11252-020-01077-0

56. Seress G, Sándor K, Evans KL, Liker A (2020) Food availability limits avian reproduction
in the city: An experimental study on great tits Parus major. J Anim Ecol 89:1570–1580.
doi: 10.1111/1365-2656.13211

57. Lemoine NP, Capdevielle JN, Parker JD (2015) Effects of in situ climate warming on
monarch caterpillar (Danaus plexippus) development. PeerJ 1–10. doi:
10.7717/peerj.1293

58. Kingsolver JG, Woods HA, Kingsolver JG, Woods HA (1997) Thermal Sensitivity of
Growth and Feeding in Manduca sexta Caterpillars. Physiol Zool 70:631–638.

59. Lee KP, Roh C (2010) Temperature-by-nutrient interactions affecting growth rate in an
insect ectotherm. Entomol Exp Appl 136:151–163. doi: 10.1111/j.1570-
7458.2010.01018.x

60. York A, Oberhauser K (2002) Effects of duration and timing of heat stress on Monarch
butterfly (Danaus plexippus) (Lepidoptera : Nymphalidae) development. J Kansas
Entomol Soc 75:290–298.

61. Zandt HS (1997) Water content of prey of nestling Blue tits in a Corsican habitat.
Netherlands J Zool 47:125–131.

62. Pollock CJ, Capilla-Lasheras P, McGill RAR, et al. (2017) Integrated behavioural and
stable isotope data reveal altered diet linked to low breeding success in urban-dwelling
blue tits (Cyanistes caeruleus). Sci Rep 7:5014. doi: 10.1038/s41598-017-04575-y

63. Jarrett C, Powell LL, McDevitt H, et al. (2020) Bitter fruits of hard labour: diet
metabarcoding and telemetry reveal that urban songbirds travel further for lower-quality
food. Oecologia 193:377–388. doi: 10.1007/s00442-020-04678-w

64. Nilsson J-Å, Nord A (2018) Testing the heat dissipation limit theory in a breeding

passerine. Proc R Soc B 285:1–8. doi: 10.1098/rspb.2018.0652

65. Wiley EM, Ridley AR (2016) The effects of temperature on offspring provisioning in a

cooperative breeder. Anim Behav 117:187–195. doi: 10.1016/j.anbehav.2016.05.009

66. Zhu L, Hoffmann AA, Li SM, Ma C Sen (2021) Extreme climate shifts pest dominance

hierarchy through thermal evolution and transgenerational plasticity. Funct Ecol

35:1524–1537. doi: 10.1111/1365-2435.13774

67. Calosi P, Bilton DT, Spicer JI (2008) Thermal tolerance, acclimatory capacity and

vulnerability to global climate change. Biol Lett 4:99–102. doi: 10.1098/rsbl.2007.0408

68. Seress G, Vincze E, Pipoly I, et al. (2017) Effects of capture and video-recording on the

behavior and breeding success of Great Tits in urban and forest habitats. J F Ornithol

88:299–312. doi: 10.1111/jofo.12205

69. Bailey LD, van de Pol M (2016) Tackling extremes: Challenges for ecological and

evolutionary research on extreme climatic events. J Anim Ecol 85:85–96. doi:

10.1111/1365-2656.12451

70. Anderson T (2006) Biology of the ubiquitous house sparrow: from genes to populations.

Oxford University Press

71. Burnham KP, Anderson DR (2002) Model Selection and Multi-Model Inference: A

Practical Information-Theoretic Approach, 2nd ed.

72. Graham MH (2003) Confronting multicollinearity in ecological multiple regression.

Ecology 84:2809–2815. doi: 10.1890/02-3114

73. R Core Team (2020) R: A Language and Environment for Statistical Computing.

74. Brooks ME, Kristensen K, Benthem KJ van, et al. (2017) glmmTMB Balances Speed and

Flexibility Among Packages for Zero-inflated Generalized Linear Mixed Modeling. R J

9:378–400.

75. Lenth R V (2018) emmeans: Estimated Marginal Means, aka Least-Squares Means. R

package version 1.2.3.

76. Bolker BM, Brooks ME, Clark CJ, et al. (2008) Generalized linear mixed models : a

practical guide for ecology and evolution. Trends Ecol Evol 24:127–135. doi:

10.1016/j.tree.2008.10.008

77. Piepho HP, Büchse A, Emrich K (2003) A Hitchhiker's Guide to Mixed Models for

Randomized Experiments. Agron Crop Sci 189:310–322.

78. Dormann CF, Elith J, Bacher S, et al. (2013) Collinearity : a review of methods to deal

with it and a simulation study evaluating their performance. Ecography (Cop) 36:27–46.

doi: 10.1111/j.1600-0587.2012.07348.x

79. Ruxton GD, Beauchamp G (2008) Time for some a priori thinking about post hoc testing.

Behav Ecol 19:690–693. doi: 10.1093/beheco/arn020

80. Pinheiro J, Bates D, DebRoy S, Sarkar D (2013) NLME: Linear and nonlinear mixed

effects models. Version 3.:1–336.

81. Bates D, Maechler M, Bolker B, Walker S (2015) Fitting linear mixed-effects models

using lme4. J Stat Softw 67:1–48.

82. Maindonald J, Braun J (2010) Data Analysis and Graphics Using R: An Example-Based

Approach, 3rd ed. Brief Bioinform. doi: 10.1093/bib/bbn012

83. Barton K (2009) Mu-MIn: Multi-model inference. R package.

 Tables and Figures

700 **Table 1. Relationship between the number of hot days during nestling development and**
701 **nestlings' body mass, tarsus length, and mortality, according to the simple models.** Type
702 3 ANOVA (analyzis of deviance) table is shown for each model. Significant effects (p < 0.05)
703 are highlighted in bold.
704

Model terms	Nestling mass			Nestling tarsus length			Nestling mortality		
	χ^2	df	P	χ^2	df	P	χ^2	df	P
Nr. hot days	3.119	1	0.077	**6.134**	**1**	**0.013**	**5.799**	**1**	**0.016**
Study site	**243.518**	**3**	**< 0.001**	**103.380**	**3**	**< 0.001**	**91.078**	**3**	**< 0.001**
Nr. hot days × study site	**29.289**	**3**	**< 0.001**	**9.555**	**3**	**0.023**	**11.844**	**3**	**0.008**

705 For average nestling mass and tarsus length, the number of pairs was 535 and the number of broods
706 was 674. For nestling mortality, the number of pairs was 600 and the number of broods was 760.
707

Table 2. Differences between urban and forest habitats in the effect of hot days on nestling's body mass, tarsus length, and mortality. The table shows linear contrasts comparing the average slope of two urban *versus* two forest sites, calculated from the estimates of the simple models (left) and the most supported multi-predictor models (right) for each response variable. For nestling mass, results of two supported multi-predictor models are separated by "/". Positive contrasts mean more positive slopes (i.e. less negative effects of hot days) in the urban habitat. Significant habitat differences (p < 0.05) are highlighted in bold.

Response	Simple model				Multi-predictor model			
	contrast ± SE	df	t	p	contrast ± SE	df	t	p
Nestling mass	**0.338 ± 0.085**	**135**	**3.978**	**< 0.001**	**0.656 ± 0.166 / 0.720 ± 0.165**	**127 / 129**	**3.941 / 4.355**	**< 0.001 / < 0.001**
Nestling tarsus length	0.046 ± 0.039	135	1.187	0.237	0.044 ± 0.075	127	0.590	0.556
Nestling mortality[*]	**-0.408 ± 0.159**	**156**	**-2.564**	**0.010**	**-0.643 ± 0.281**	**149**	**-2.288**	**0.022**

[*] Estimates for nestling mortality are on the logit scale

Figure 1. Relationships between the number of hot days during the nestling period and (A) body mass, (B) tarsus length, and (C) mortality of great tit nestlings. Circles represent brood means for nestling mass and tarsus length. Colored stripes show the 95% confidence band of the slope estimated from the simple models (Table S4).

      Supplementary Material for

724   **"**Extreme hot weather has stronger impacts on avian reproduction in forests than

725   in cities" by

726   Ivett Pipoly, Bálint Preiszner, Krisztina Sándor, Csenge Sinkovics, Gábor Seress, Ernő

727   Vincze, Veronika Bókony, András Liker

728
Figure S1. Distribution of the number of hot days and average temperature during the

nestling periods in our study sites. Boxes show the interquartile range, the thick line is the

median, and whiskers refer to the range of data distribution. Data points represent great tit

broods (green: forest, blue: urban).

Table S1. Result of AICc-based model selection for average nestling mass. Supported
models are highlighted in bold.

Model	df	logLik	AICc	ΔAICc	Akaike weight
456789	**17**	**-1220.19**	**2475.31**	**0**	**0.55**
45789	**16**	**-1221.97**	**2476.78**	**1.47**	**0.26**
23456789	19	-1220.17	2479.49	4.19	0.07
1456789	18	-1221.48	2480.00	4.69	0.05
145789	17	-1223.37	2481.67	6.36	0.02
2345789	18	-1222.4	2481.84	6.53	0.02
123456789	20	-1221.12	2483.52	8.22	0.01
46789	16	-1226.23	2485.29	9.99	0
12345789	19	-1223.17	2485.51	10.2	0
4789	15	-1227.84	2486.41	11.11	0
2346789	18	-1226.43	2489.91	14.61	0
146789	17	-1227.61	2490.15	14.84	0
234789	17	-1227.96	2490.85	15.54	0
14789	16	-1229.11	2491.05	15.75	0
2345679	14	-1232.31	2493.25	17.95	0
12346789	19	-1227.39	2493.94	18.63	0
45679	12	-1234.77	2494.02	18.71	0
1234789	18	-1228.84	2494.72	19.42	0
12345679	15	-1232.37	2495.48	20.17	0
145679	13	-1235.07	2496.68	21.38	0
4679	11	-1241.41	2505.22	29.91	0
234679	13	-1239.51	2505.57	30.26	0
1234679	14	-1239.99	2508.62	33.31	0
14679	12	-1242.58	2509.63	34.32	0
1234579	14	-1245.06	2518.77	43.46	0
4579	11	-1249.02	2520.44	45.13	0
14579	12	-1249.51	2523.49	48.18	0
234579	13	-1248.72	2524.00	48.69	0
123479	13	-1251.77	2530.09	54.78	0
479	10	-1255.36	2531.04	55.74	0
23479	12	-1254.36	2533.20	57.89	0
1479	11	-1256.61	2535.61	60.31	0
(Null)	3	-1369.13	2744.30	269	0

736
737 Model term codes: 1=average temperature of nestling periods, 2=(hatching date)^2, 3=
738 hatching date, 4=number of hot days, 5=brood age, 6=brood size, 7= study site, 8=years, 9=
739 number of hot days × study site. Null model contains only the random effect on nestlings'
740 body mass.
741

742 **Table S2. Result of AICc-based model selection for average nestling tarsus length.**
743 Supported model is highlighted in bold.

Model	df	logLik	AICc	ΔAICc	Akaike weight
145679	**13**	**-686.49**	**1399.52**	**0**	**0.94**
12345679	15	-687.74	1406.21	6.68	0.03
1456789	18	-686.07	1409.18	9.65	0.01
14679	12	-692.46	1409.39	9.86	0.01
2345679	14	-691.27	1411.17	11.65	0
123456789	20	-685.2	1411.68	12.16	0
45679	12	-693.68	1411.83	12.31	0
1234679	14	-693.67	1415.99	16.46	0
4679	11	-697.1	1416.59	17.07	0
456789	17	-690.98	1416.89	17.36	0
146789	17	-691.88	1418.69	19.17	0
234679	13	-696.21	1418.98	19.45	0
12346789	19	-690.66	1420.49	20.97	0
46789	16	-695.18	1423.18	23.66	0
23456789	19	-692.23	1423.63	24.1	0
2346789	18	-696.89	1430.82	31.3	0
12345789	19	-698.68	1436.53	37.01	0
1234789	18	-702.76	1442.56	43.03	0
45789	16	-708.91	1450.64	51.12	0
145789	17	-707.92	1450.76	51.24	0
4789	15	-712.47	1455.66	56.14	0
1234579	14	-713.53	1455.70	56.17	0
2345789	18	-709.65	1456.34	56.82	0
14789	16	-712.36	1457.55	58.03	0
234789	17	-712.68	1460.30	60.78	0
14579	12	-718.76	1461.98	62.46	0
123479	13	-718.12	1462.79	63.27	0
1479	11	-723.78	1469.97	70.44	0
4579	11	-724.26	1470.91	71.39	0
479	10	-727.16	1474.64	75.12	0
234579	13	-726.24	1479.04	79.52	0
23479	12	-729.11	1482.70	83.17	0
(Null)	3	-783.63	1573.30	173.78	0

744
745 Model term codes: 1=average temperature of nestling periods, 2=(hatching date)^2, 3=
746 hatching date, 4=number of hot days, 5=brood age, 6=brood size, 7= study site, 8=years, 9=
747 number of hot days × study site. Null model contains only the random effect on nestlings'
748 tarsus length.
749

750 **Table S3. Result of AICc-based model selection for nestling mortality.** Supported model is
751 highlighted in bold.

model	df	logLik	AICc	ΔAICc	Akaike weight
1234567	**18**	**-1032.7**	**2102.33**	**0**	**1**
234567	17	-1055.19	2145.20	42.87	0
123457	13	-1074.94	2176.37	74.04	0
4567	15	-1088.84	2208.32	105.99	0
14567	16	-1088.53	2209.79	107.46	0
23457	12	-1105.89	2236.20	133.87	0
1457	11	-1125.98	2274.31	171.98	0
457	10	-1130.95	2282.19	179.86	0
(Null)	3	-1203.8	2413.62	311.29	0

752
753 Model term codes: 1=average temperature of nestling periods, 2=(hatching date)^2, 3=
754 hatching date, 4=number of hot days, 5= study site, 6=years, 7= number of hot days × study
755 site. Null model contains only the random effect on nestling mortality.

Table S4. The effect of the number of hot days during the nestling period on reproductive success in great tits. The slope of the relationship with 95% confidence interval (CI) is shown for each site, as estimated by marginal means from A) the simple models (containing number of hot days, study site, and their two-way interaction) and B) the supported multi-predictor models (Table 1). For nestling mass, results of the two supported multi-predictor models are separated by "/". Slopes significantly different from zero (i.e. zero not included between the lower and upper limit of CI) are highlighted in bold.

Study sites	Average nestling mass				Average nestling tarsus length				Nestling mortality[*]			
	Slope	SE	lower CI	upper CI	Slope	SE	lower CI	upper CI	Slope	SE	lower CI	upper CI
A) simple model												
Veszprém city	-0.081	0.046	-0.172	0.010	**-0.052**	**0.021**	**-0.094**	**-0.011**	**0.198**	**0.082**	**0.037**	**0.360**
Balatonfüred city	**0.163**	**0.049**	**0.065**	**0.260**	0.036	0.023	-0.009	0.081	0.011	0.090	-0.165	0.187
Szentgál forest	-0.106	0.085	-0.273	0.061	-0.050	0.038	-0.126	0.026	0.161	0.176	-0.183	0.505
Vilma-puszta forest	**-0.489**	**0.131**	**-0.749**	**-0.230**	-0.059	0.060	-0.178	0.060	**0.864**	**0.236**	**0.401**	**1.327**
B) multi-predictor model												
Veszprém city	-0.201 / **-0.269**	0.098 / **0.095**	-0.394 / **-0.457**	-0.007 / **-0.082**	**-0.146**	**0.049**	**-0.243**	**-0.050**	0.246	0.201	-0.148	0.639
Balatonfüred city	**0.271** / **0.236**	**0.098** / **0.098**	**0.076** / **0.042**	**0.466** / **0.430**	0.002	0.049	-0.094	0.098	0.113	0.193	-0.266	0.492
Szentgál forest	**-0.473** / **-0.609**	**0.185** / **0.179**	**-0.840** / **-0.962**	**-0.107** / **-0.255**	-0.120	0.080	-0.279	0.038	0.527	0.362	-0.182	1.236
Vilma-puszta forest	**-0.768** / **-0.866**	**0.258** / **0.256**	**-1.278** / **-1.373**	**-0.258** / **-0.359**	-0.112	0.120	-0.349	0.125	**1.118**	**0.442**	**0.251**	**1.985**

For average nestling mass and tarsus length, number of pairs was 535 and number of broods was 674. Sample sizes: Veszprém n= 222, Balatonfüred n= 115, Szentgál forest n= 242, Vilma-puszta forest n= 95 broods.
For nestling mortality, number of pairs was 600 and number of broods was 760. Sample sizes: Veszprém n= 260, Balatonfüred n= 130, Szentgál forest n= 260, Vilma-puszta forest n= 110 broods.
[*]Estimates for nestling mortality are on the logit scale