# The proteasomal deubiquitinating enzyme PSMD14 regulates macroautophagy by controlling Golgi-to-ER retrograde transport

Bustamante HA[1,2], Cereceda K[3], González AE[1,4], Valenzuela GE[5,6], Cheuquemilla Y[6], Hernández S[3], Arias-Muñoz E[3], Cerda-Troncoso C[3], Bandau S[8], Soza A[3], Kausel G[5], Kerr B[3], Mardones GA[1,7], Cancino J[3], Hay RT[8], Rojas-Fernandez A[6,8¥], Burgos PV[3,9¥]

[1]Instituto de Fisiología, Facultad de Medicina, Universidad Austral de Chile, 5110566, Valdivia, Chile

[2]Instituto de Microbiología Clínica, Facultad de Medicina, Universidad Austral de Chile, 5110566, Valdivia, Chile

[3]Centro de Biología Celular y Biomedicina (CEBICEM), Facultad de Medicina y Ciencia, Universidad San Sebastián, Lota 2465, 7510157, Santiago, Chile

[4]Institute of Biochemistry II, School of Medicine, Goethe University Frankfurt, Theoder-Stern-Kai 7, 60590, Frankfurt am Main, Germany

[5]Instituto de Bioquímica y Microbiología, Facultad de Ciencias, Universidad Austral de Chile, 5110566, Valdivia, Chile.

[6]Instituto de Medicina & Centro Interdisciplinario de Estudios del Sistema Nervioso (CISNe), Universidad Austral de Chile, 5110566, Valdivia, Chile.

[7]Centro Interdisciplinario de Estudios del Sistema Nervioso (CISNe), Universidad Austral de Chile, 5110566, Valdivia, Chile.

[8]Centre for Gene Regulation and Expression, College of Life Sciences, University of Dundee, DD1 4HN, Dundee, United Kingdom.

[9]Centro de Envejecimiento y Regeneración (CARE-UC), Facultad de Ciencias Biológicas, Pontificia Universidad Católica de Chile, 83330023, Santiago, Chile

**Correspondence to:**
¥ Patricia Burgos, PhD. Centro de Biología Celular y Biomedicina, Facultad de Medicina y Ciencia, Universidad San Sebastián, Lota 2465, Santiago 7510157, Chile. Email: patricia.burgos@uss.cl

¥Alejandro Rojas-Fernández, PhD. Instituto de Medicina & Centro Interdisciplinario de Estudios del Sistema Nervioso (CISNe), Universidad Austral de Chile, Valdivia 5110566, Chile. Email: alejandro.rojas@uach.cl

**ABSTRACT**

Ubiquitination regulates several biological processes. Here, we search for ubiquitin-related genes implicated in protein membrane trafficking performing a High-Content siRNA Screening including 1,187 genes of the human "ubiquitinome" using Amyloid Precursor Protein (APP) as a reporter. We identified the deubiquitinating enzyme PSMD14, a subunit of the 19S regulatory particle of the proteasome, specific for K63-Ub chains in cells, as a novel key regulator of Golgi-to-endoplasmic reticulum (ER) retrograde transport. Silencing or pharmacological inhibition of PSMD14 caused a robust and rapid inhibition of Golgi-to-ER retrograde transport which leads to a potent blockage of macroautophagy by a mechanism associated with the retention of Atg9A and Rab1A at the Golgi apparatus. Because pharmacological inhibition of the proteolytic core of the 20S proteasome did not recapitulate these effects, we concluded that PSMD14, and their K-63-Ub chains, act as a crucial regulator factor for macroautophagy by controlling Golgi-to-ER retrograde transport.

Keywords: PSMD14, Ubiquitin, Retrograde, Trafficking, APP

## Introduction

Several post-translational modifications (PTMs) contribute to membrane transport [1,2]. Among these, ubiquitination acts as a relevant player in protein membrane trafficking [3–5] participating as an important protein localization signal [3,6–8]. However, to date most studies have focused on its role in endocytosis, particularly in the inclusion of cargoes in intraluminal vesicles (ILVs) of multivesicular bodies (MVBs) [9–11] such as cell-surface receptors [12]. Ubiquitination is a regulated post-translational modification that conjugates ubiquitin (Ub) to lysine (K) residues and is involved in many cellular pathways including the degradation of target proteins through the proteasomes and autophagy [13,14]. Ubiquitin modification is a cascade of reactions catalyzed by three classes of enzymes (E1, E2 and E3). An ATP dependent E1 activating enzymes first forms a covalent intermediate with ubiquitin, followed by the transfer of Ub to E2 conjugating enzymes and finally the covalent attachment of Ub to a K residue in the target protein by E3 Ub ligases [15]. Ubiquitination is a reversible reaction, with specific deubiquitinating enzymes (DUBs) that catalyze the removal of Ub-moieties for their recycling. DUBs serve to counterbalance ubiquitination reactions within the cell, thus dynamically contributing to the regulation of various cellular processes, such as endosomal sorting [11,16–18].

Ub increases proteome complexity, providing an additional surface for protein-protein interactions and functional regulation [19,20]. Ub-modified proteins and Ub chains are recognized by several Ub receptors including a family of specialized proteins carrying Ub binding domains (UBDs). To date, over 20 UBD families have been identified and characterized in mammals participating in recognition of Ub monomers or Ub chains on specific substrates [4,14,21]. Together, the Ub network offers a variety of choices to modulate cellular processes including protein membrane trafficking. However, the contribution of the ubiquitinome regarding their impact in intracellular membrane trafficking is not yet fully understood.

To investigate the contribution of most of the members of the ubiquitinome in protein membrane trafficking, we performed High-Content siRNA Screening (HCS) including 1,187 genes of the human "ubiquitinome". For this screening we used a stable cell line expressing Amyloid Precursor Protein (APP) fused to EGFP. APP was the protein selected as target because it traffics dynamically through the secretory and endocytic pathways

containing specific sorting signal motifs [22–24] and specific lysine residues targets of

ubiquitination [25–28]. We thus identified the deubiquitinating (DUB) enzyme PSMD14, a

subunit of the 19S regulatory particle (RP) of the proteasome, as a crucial player of Golgi-

to-Endoplasmic reticulum (ER) retrograde transport. The DUB enzyme PSMD14 has been

shown to be specific for K63-Ub chains in cells [29]. Here, we found that the inhibition of its

activity blocks Golgi-to-ER retrograde transport causing the swelling of the Golgi

apparatus. Also, we found that PSMD14 inhibition acts as a potent blocker of

macroautophagy as a result of its negative impact in Golgi-to-ER retrograde transport.

Inhibition of the proteolytic core of the 20S proteasome by MG132 did not recapitulate

these effects, indicating that the 19S RP plays a role on macroautophagy by controlling

membrane trafficking at the early secretory pathway.

**Results**

***High-Content siRNA Screening revealed PSMD14 deubiquitinating enzyme as a novel***
***regulator of protein trafficking.***

To investigate the contribution of the ubiquitinome in protein membrane trafficking, we
performed an automated HCS using a human small interfering RNA (siRNA)
"ubiquitinome" library, consisting of 1,187 siRNA duplex pools targeting all known and
predicted Ub-genes. This library, used previously in mammalian screening assays [30,31]
includes E1 activating enzymes (0.8%), E2 conjugating enzymes (4.7%), E3 Ub ligases
(61.2%), UBD-containing proteins (12.8%), DUBs (8.8%), SUMO-specific proteases
(SENPs) (0.2%) and others (11.4%) (Fig. 1A). We used this library to identify novel
regulators of protein membrane trafficking using an H4 neuroglioma cell line stably
expressing APP-EGFP, a cell line previously characterized and used in siRNAs knockdown
experiments [24,32,33]. We standardized the basal (background) and maximum fluorescence
intensity by analyzing total fluorescence of silenced and non-silenced APP-EGFP,
respectively. A reduction in total fluorescence intensity from 100% to 6% was observed
with the EGFP siRNA, compared to the Non-Target (NT) siRNA with a Z factor of 0.69
(Fig. 1B). Representative images of basal and maximum levels of total fluorescence
intensity in reporter cells are shown in Fig. 1C. Cells were further assessed with a primary
siRNA screening consisting of a pool of four-different siRNA duplexes for each of the
1,187 genes selected, each siRNA targeting a distinct region to minimize their off-targets
effects [34]. Fig. 1D shows a summary of the results with all siRNA pools tested and
normalized in respect to the expression of the NT siRNA. Among the 1,187 tested genes,
we found that the highest total fluorescence increase (4.15-fold increase) corresponded to
protein PSMD14 (POH1, also known as Rpn11/MPR1/SS13/CepP1) (Fig. 1D), a subunit of
the 19S regulatory particle (RP) of the proteasome, which has DUB activity [35,36].
Representative images of reporter cells knock-down (KD) of PSMD14 in comparison to
cells transfected with NT siRNA are shown in Fig. 1E. Further, we validated these results
analyzing the effect of PSMD14 KD on the levels of full-length endogenous APP in
parental H4 cells by western blot. The effect of all four PSMD14 siRNAs in KD cells tested
on endogenous APP levels are shown in Fig. 2A (first panel, lanes 3-6) and compared to
un-transfected cells (Mock) or cells transfected with NT siRNA (Fig. 2A, first panel, lanes

**Figure 1**

**Figure 2**

1 and 2, respectively). Quantification analysis showed a 3.13-fold increase of full-length
endogenous APP levels in PSMD14 KD cells, compared to Mock cells (Fig. 2B). Besides,
we found that silencing of PSMD14 caused a strong increase in high molecular weight Ub
conjugates, consistent with the role of PSMD14 as a proteasomal subunit with DUB
activity (Fig. 2A, second panel, lanes 3-6 compared to lanes 1 and 2). Confirmation of the
efficient KD of PSMD14 with all four siRNAs tested was demonstrated by western blot
(Fig. 2A, third panel, lanes 3-6 compared to lanes 1 and 2), observing a 5.22-fold decrease
on PSMD14 levels (Fig. 2C). Moreover, efficient KD of PSMD14 was also determined by
quantitative reverse transcription PCR (RT-qPCR) (Fig. 2D). Finally, to evaluate whether
the increase of APP protein levels in PSMD14 KD cells could be the result of an up-
regulation of APP transcription, we evaluated *APP* mRNA levels by RT-qPCR, observing
no significant changes (Fig. 2E). Altogether, these findings indicate that the silencing of
PSDM14 caused a robust increase of APP protein levels, a finding that could suggests
impairment in its turnover due to trafficking alterations.

***Acute inhibition of the deubiquitinating enzyme PSMD14 of the 19S RP accumulates***
***APP in a swollen Golgi apparatus.***

To determine whether PSMD14 is involved in the control of endogenous APP levels, we
investigated the effect of incubating cells with Capzimin (CZM), a potent and specific
inhibitor of the DUB activity of PSMD14[37], a subunit of the 19S RP of the proteasome
(Fig. 3A). We observed that CZM led to a significant increment in full-length endogenous
APP levels in a dose-dependent manner, compared to untreated cells (Fig. 3B and 3C).
Interestingly, compared to untreated cells, and in contrast to CZM, treatment with a
standard concentration of 10 μM MG132, a reversible inhibitor of the β1, β2 and β5
subunits of the 20S catalytic core of the proteasome[38,39] (Fig. 3A), caused a significant
decrease in full-length endogenous APP levels (Fig. 3B and 3C). To confirm the inhibition
of PSMD14 by CZM, we tested the effect of this inhibitor on the levels of high molecular
weight Ub conjugates, comparing it with MG132. In agreement with the effect of PSMD14
KD, we observed that CZM caused a robust increase in high molecular weight Ub
conjugates in a dose-dependent manner, finding a maximum effect with 10 μM CZM (Fig.
3D). However, we noticed that 10 μM MG132 caused a more powerful increase in Ub
conjugates (Fig. 3D). To confirm the accumulation of Ub conjugates by these treatments,

**Figure 3**

we studied the presence of Ub-inclusion bodies by immunofluorescence analysis. We found

that, compared to untreated cells (Fig. 3E, left panel), 10 µM CZM (Fig. 3E, middle panel)

caused a similar accumulation of cytoplasmic Ub-inclusion bodies than 10 µM MG132

(Fig. 3E, right panel). Together, our results confirm that acute inhibition of PSMD14 by

CZM replicates the phenotype obtained by PSMD14 KD regarding to the impact on the

endogenous APP levels. Thus, CZM offers a pharmacological tool to address whether the

activity of PSDM14 is required for APP membrane trafficking.

To unveil this possibility, we performed an immunofluorescence analysis of endogenous

APP in parental H4 cells. We observed that the treatment with 10 µM CZM resulted in a

perinuclear redistribution of APP, which is highly indicative of Golgi apparatus localization

(Fig. 4D, compared to 4A). Moreover, similar to the PSMD14 KD, we observed that the

CZM treatment caused a significant 2.50-fold increase in APP total fluorescence intensity

compared to untreated cells (Fig. 4G). We also observed a significant 1.92-fold increase in

the amount of APP in the area positive to the Golgi matrix protein GM130 (GM130),

compared to the total area (Fig. 4F, compared to 4C and Fig. 4H). Moreover, we observed

that the treatment with CZM caused a significant 1.30-fold increase in the total cell area

(Fig. 4I). In addition, we found that CZM caused the swelling of the Golgi apparatus (Fig.

4E compared to 4B), a phenotype that is shown in a zoom in Fig. 4J. To confirm this

phenotype, we performed measurements of GM130 images from confocal 3D

reconstructions from the Z-stacks of cells treated with CZM (Fig. 4K). We observed a 1.53-

fold increase in the Golgi apparatus volume in CZM treated cells, compared to control cells

(Fig. 4L). Similar results were observed in quantitative 2D image analysis, observing a

significant 2.53-fold increase in the Golgi apparatus area upon CZM treatment (Suppl. Fig.

1). Altogether, our results show that acute inhibition of PSMD14 by CZM accumulates

APP in a swollen Golgi apparatus. These findings strongly suggest that CZM impairs the

Golgi apparatus due to perturbations of the trafficking through this organelle.

***Acute inhibition of the deubiquitinating enzyme PSMD14 perturbs Golgi-to-ER***

***retrograde transport***

The Golgi apparatus is a highly dynamic organelle that requires fine regulation of

trafficking pathways in order to maintain its size, shape and composition. In particular, it

has been shown that Golgi-to-ER retrograde transport plays a crucial role in the

**Figure 4**

maintenance of Golgi morphology. Indeed, inhibition of this specific trafficking pathway

results in a significant swelling of this organelle [40–45]. To investigate whether the swelling

of the Golgi by the inhibition of PSMD14 with CZM could be the result of the inhibition of

the Golgi-to-ER retrograde transport, we first validated our findings in HeLa cells. We

measured Golgi apparatus volume in 3D reconstructions from the Z-stacks of untreated

HeLa cells or treated with CZM and MG132 by using Giantin as Golgi apparatus reporter.

Similar to our findings in H4 cells, we found that the inhibition of PSMD14 DUB activity

by CZM causes a 2.14-fold increase in Golgi apparatus volume (Fig. 5A, middle panel

compared to the left panel and 5B). Interestingly, in contrast to CZM, we observed that

MG132 caused no effect on Golgi apparatus volume (Fig. 5A, right panel compared to the

left panel and 5B), strongly suggesting that Golgi homeostasis is regulated by the PSMD14

DUB activity. Then, we investigated whether this phenotype was the result of Golgi-to-ER

retrograde transport inhibition. We assessed the distribution of stably overexpressed KDEL

(Lys-Asp-Glu-Leu) Receptor 1 (KDELR1) fused to GFP (KDELR1-GFP) in HeLa cells.

As it was previously reported [44,46], overexpressed KDELR1-GFP is mostly distributed to

the ER due to its efficient transport from the Golgi apparatus to the ER (Fig. 5C, left panel

and 5D). Thus, impairment of Golgi-to-ER retrograde transport causes the accumulation of

KDELR1-GFP in the Golgi apparatus working as an assay to identify novel regulators of

this pathway [44]. Interestingly, we found that CZM caused a rapid time-dependent

accumulation of KDELR1-GFP at the Golgi apparatus, observing a maximal effect after 90

min of treatment (Suppl. Fig. 2). At this time, we observed a significant 1.50-fold increase

in the amount of KDELR1-GFP within the Giantin-positive Golgi apparatus, compared to

the total area (Fig. 5C, middle panel compared to the left panel, and 5D). As before, we

tested the effect of MG132, observing no effect on the distribution of KDELR1-GFP (Fig

5C, right panel compared to left panel, and 5D), which strongly supports that MG132 has

no effect on Golgi-to-ER retrograde transport. Next, to confirm blockage of Golgi-to-ER

retrograde transport by the acute inhibition of the PSMD14, we tracked the redistribution of

the transiently overexpressed thermo-sensitive KDELR1-VSVG-YFP chimera (vesicular

stomatitis virus G protein fused to KDEL receptor 1 and YFP) in HeLa cells upon treatment

with CZM by live cell imaging at different temperatures. Briefly, at a permissive

temperature of 32°C, KDELR1-VSVG-YFP cycles between the Golgi apparatus and the

**Figure 5**

**KDELR1-GFP**

**Supplementary Figure 2**

ER, showing distribution mainly at the Golgi apparatus. Upon shifting to a restrictive

temperature of 40°C, KDELR1-VSVG-YFP is progressively accumulated at the ER due to

its specific retention in this compartment at this temperature. In addition, because the

Golgi-to-ER retrograde transport is not affected at 40ºC, the rapid decay of the fluorescence

of KDELR1-VSVG-YFP fluorescence at the Golgi apparatus, a measurement that is used to

evaluate inhibition of Golgi-to-ER retrograde transport pathway [47]. We found that in

control cells nearly 50% of the fluorescence of KDELR1-VSVG-YFP decayed after 15 min

of shifting the temperature to 40°C (Fig. 5E, upper panels and 5F), indicating that

retrograde transport is working normally [47]. In contrast, when cells were treated with CZM,

the decay of the fluorescence of KDELR1-VSVG-YFP was not apparent, confirming that

acute inhibition of the PSMD14 blocked Golgi-to-ER retrograde transport (Fig. 5E, lower

panels and 5F). In contrast, MG132 treated cells showed similar results as controls (data not

shown), confirming that the 20S proteasome has no effect on retrograde transport.

Importantly, we found that CZM is able to block retrograde transport for short times having

no effect on the catalytic activity of the 20S proteasome (Suppl. Fig. 3). In contrast, a rapid

and robust inhibition on the catalytic activity of the 20S proteasome is observed with

MG132 (Suppl. Fig. 3). These findings strongly indicate that acute inhibition of PSMD14

DUB activity by CZM acts as a powerful blocker of Golgi-to-ER retrograde transport,

explaining the swelling of the Golgi apparatus and the accumulation of protein cargoes

such as APP at this location.

***Inhibition of Golgi-to-ER retrograde transport by CZM has a negative impact on***

***macroautophagy.***

Several lines of evidence have shown that Golgi-to-ER retrograde transport plays a relevant

role in autophagosome biogenesis at the level of the ER [48–51]. In addition, it has been shown

that PSMD14 participates in the activation of the aggresome clearance by cleaving K63 Ub

chains of aggregate proteins[52,53]. Thus, we investigated the effect of acute inhibition of

PSMD14 DUB activity on the number of autophagosomes by testing the classical marker

microtubule-associated protein 1 light chain 3B (LC3B) [54]. Therefore, parental H4 cells were

treated under nutrient starvation with Earle's balanced salt solution (EBSS), an established

culture medium used for activation of autophagosomal formation [55]. As expected, we found

that starvation strongly increased the number of autophagosomes (Fig. 6B), compared to

**Figure 6**

**Supplementary Figure 3**

cells treated under normal nutrient conditions (Fig. 6A). In contrast, we observed that

treatment with CZM abolished the appearance of autophagosomes upon EBSS treatment

(Fig. 6C compared to 6B). We also tested the effect of CZM in cells under normal nutrients

conditions, but in the absence or presence of Torin-1, a potent and selective inhibitor of

mammalian target of rapamycin complexes (mTORC1/2), a trigger of autophagosomal

formation[56]. Similar to the results with EBSS, we observed that CZM prevented the

appearance of autophagosomes promoted by Torin-1 (Fig. 6F compared to 6E). In

agreement with these findings, we observed that CZM alone did not increase the number of

autophagosomes (Fig. 6D compared to 6A), in contrast to the effect of MG132 (Suppl. Fig.

4). To confirm these results, we quantified the number of LC3-positive structures under all

conditions tested, observing a significant decrease in the number of autophagosomes when

cells, treated with EBSS solution or Torin-1, where also treated with CZM (Fig. 6G). In

addition, we validated these results biochemically performing western blot analysis of

endogenous LC3B (Fig. 6H). We found that the levels of LC3B-II were strongly increased

with EBSS or Torin-1 treatment (Fig. 6H, lanes 2 and 4). In contrast, when cells were

treated with EBSS or Torin-1 in the presence of CZM, LC3B-II levels did not change (Fig.

6H, lanes 3 and 5) compared to control cells (Fig. 6H, lane 1), results that were quantified

and depicted in Fig. 6J. Interestingly, this biochemical analysis also showed that CZM

increased LC3B-I levels under all conditions tested (Fig. 6H, lanes 3, 5 and 6, and Fig. 6I)

compared to control cells (Fig. 6H, lane 1), suggesting that the reduction of the

autophagosomal structures caused by CZM was not due to a reduction in LC3B total levels.

Altogether, these results confirm that acute inhibition of PSMD14 DUB activity acts as a

potent blocker of autophagosome biogenesis induced by EBSS or Torin-1. In addition,

these findings suggest that blockage of autophagosomal biogenesis could be a consequence

of Golgi-to-ER retrograde transport inhibition.

***Inhibition of Golgi-to-ER retrograde transport by CZM accumulates RAB1A and***

***ATG9A at the Golgi apparatus***

To evaluate this hypothesis, we tested the effect of acute inhibition of the PSMD14 DUB

activity on the distribution of proteins implicated in the initial steps of autophagosome

formation, which traffics in early compartments of the secretory pathway. We first tested

RAB1A, a small GTPase with an essential role in the initiation of autophagy, facilitating

**A**

Control       10 μM CZM      10 μM MG132

LC3B

**B**

Control
10 μM CZM
10 μM MG132

kDa
20 — LC3B-I
— LC3B-II

50 — β-actin

1    2    3

**C**

4.03
***
n.s.

Ratio LC3B-II/LC3B-I

Control   10 μM CZM   10 μM MG132

# Supplementary Figure 4

the recruitment of the unc-51-like kinase 1 (ULK1) complex to subdomains of the ER, a

crucial early step during autophagosome formation [57,58]. We found that CZM treatment

caused a redistribution of RAB1A to the perinuclear zone (Fig. 7A). Measuring these

images confirmed a significant increase of RAB1A in this area (Fig. 7B), a result that was

accompanied by a decrease of RAB1A in the cell periphery defined as radial zone (Fig.

7B). Importantly, we found that RAB1B was not affected by the same condition (data not

shown). Moreover, we observed that RAB1A was distributed in the swollen Golgi

apparatus, similar to GM130, upon CZM treatment (Suppl. Fig. 5). With the same aim, we

studied the distribution of ATG9A, an essential transmembrane protein involved in

macroautophagy, which plays a crucial role in the early steps of autophagosome formation

[59,60]. ATG9A-containing vesicles are formed from the Golgi apparatus translocating to the

ER to form the initiation site at the ER for autophagosome formation [61]. Similar to RAB1A,

we found that acute inhibition of PSMD14 caused a redistribution of ATG9A to the

perinuclear zone, together with a decrease in its distribution in the radial zone (Fig. 7C).

The quantification analysis of these images is shown in Fig. 7C. Moreover, and similar to

RAB1A, we observed that ATG9A is distributed to the swollen GM130-Golgi apparatus

area upon CZM treatment (Suppl. Fig. 6). These results strongly indicate that blockage of

Golgi-to-ER retrograde transport by acute inhibition of PSMD14 DUB activity causes the

retention at the Golgi apparatus of key proteins implicated in early steps of autophagosome

formation. The PSDM14-dependent inhibition of autophagy explains the accumulation of

APP at the Golgi apparatus since macroautophagy has been recently demonstrated as a

positive regulator of protein secretion from the Golgi apparatus[62]. Collectively, these results

show the strong interplay between membrane transport and autophagy through a novel

mechanism involving the proteasome complex through the deubiquitinating activity of

PSMD14.

## Discussion

300  We report here that PSMD14 DUB activity, a subunit of the 19S RP of the proteasome,

301  functions as a novel regulator of autophagosome formation. To our knowledge, this is the

302  first report demonstrating that impairment of the proteasome can have a negative impact on

303  the initiation of autophagy. Several reports have shown that inhibition of the 20S catalytic

**Figure 7**

**Supplementary Figure 5**

**Supplementary Figure 6**

core of the proteasome by blockers of the β-subunits trigger the enhancement of the

biogenesis of LC3B-positive autophagosomes [63–68]. Contrary to these findings, we found

that the blockage of the PSMD14 DUB activity perturbs the biogenesis of LC3B-positive

autophagosomes. Because the inhibition of PSMD14 DUB activity but not the blockage of

the 20S catalytic core blocked Golgi-to-ER retrograde transport, a pathway implicated in

the initiation of autophagosomes [48–51], we postulated that PSMD14 DUB activity controls

autophagy by a process independent of bulk proteasomal degradation but dependent on

K63-Ub chains. Indeed, K63-Ub chains has been directly involved in the recycling of

transmembrane cargoes into the endosomal pathway in *C. elegans*[69]. In addition, K63-Ub

chains regulate positively autophagy by mTORC inactivation through K63-linked

polyubiquitination of RagA[70], and by promoting K63-chain autoubiquitination of the E3

ligase TRIM13 that increases the interaction with the autophagy receptor p62 promoting

initiation of ER-phagy[71]. Likewise, the deconjugation of K63-Ub chains has biological

relevance. Free unanchored K63-Ub chains release in a PSMD14 dependent manner are

crucial to coordinate the elimination of protein aggregates by autophagy[52]. Free unanchored

K63-Ub chains bind and activate the deacetylase HDAC6[52,53], which contributes to relocate

the aggregates to the aggresome for final autophagic clearence[72-83]. Here, we unveil the first

insight into the regulation of Golgi-to-ER retrograde transport by K63-Ub chains by the

functional role of the deubiquitinating enzyme PSMD14. We hypothesize that the function

of key cytosolic proteins implicated in Golgi-to-ER retrograde transport are under the

control of K63-Ub chains, confirming the crucial regulatory role of deubiquitination in

membrane protein trafficking [73,74].

In this regard, key regulatory proteins involved in Golgi-to-ER transport are regulated by

the state of ubiquitination/deubiquitination. In yeast, deletion of the DUB Ube3p and its co-

factor Bre5p accumulates ubiquitinated β′-COP facilitating its rapid degradation by the

proteasome [75,73]. Because β′-COP is a subunit of the COP-I coatomer complex, a key

machinery implicated in Golgi-to-ER retrograde transport, reduction in β′-COP levels

perturbs this trafficking pathway [73]. Two other proteins that participates in this retrograde

trafficking pathway, such as PKA and UVRAG [44,76], have been shown to be regulated by

ubiquitination. Blockers of PKA signaling cause inhibition of this trafficking pathway, a

process that is accompanied by the swelling of the Golgi apparatus [44,45]. It has been shown

that the catalytic PKA subunit (PKAc) is ubiquitinated by the CHIP E3 ligase, resulting in

proteasomal degradation of PKAc and signaling shutdown [77]. UVRAG mediates the

interaction of β′-COP with ER tethers and COP-I coatomer for efficient fusion of

retrograde vesicles to the ER, a crucial step during Golgi-to-ER retrograde transport [49].

UVRAG is ubiquitinated by SMURF1 [78] and its silencing causes the swelling of the Golgi

apparatus and the inhibition of the Golgi-to-ER retrograde transport [49]. Whether these

proteins are regulated by K63-Ub chains is unknown and should be further addressed.

In addition, we propose that Golgi-to ER retrograde transport inhibition by PSMD14

dysfunction might result on the accumulation of structural and/or autophagy regulatory

elements. In fact, the inhibition of PSMD14 caused the accumulation of ATG9A at the

Golgi. ATG9A is a multispanning membrane protein essential for autophagy [59,60]. ATG9A

is actively transported through Golgi-to-ER retrograde transport [61] and participates as a key

player in the biogenesis of autophagosomes [59,79]. Moreover, along with the accumulation of

ATG9A at the Golgi, we found increased levels of RAB1A at the Golgi apparatus. RAB1A

is an essential small GTPase that participates in the recruitment of the ULK1 complex to

subdomains of the ER for autophagy initiation [57,58]. Interestingly, and similar to PSMD14

inhibition, ATG9A deficiency causes an increase in LC3B-I levels, accompanied by a

reduction in the number of autophagosome structures [59,79,80]. Together, our results strongly

support that PSMD14 inhibition perturbs autophagosome biogenesis due to the

sequestration of key proteins of this process at the Golgi apparatus. In addition, it supports

the hypothesis of a closed intersection between Golgi-to-ER retrograde and autophagy

pathway [49,76] whereas PSMD14 DUB activity emerged as a new regulatory element of this

intersection. The PSDM14-dependent inhibition of macroautophagy might affect protein

secretion from the Golgi apparatus since autophagy has been recently demonstrated as a

positive regulator of this process[62] as observed by the strong accumulation of APP at the

Golgi apparatus. Collectively, the results demonstrate the strong functional interplay

between membrane transport and macroautophagy mediated by a novel mechanism

involving the proteasome complex through the deubiquitinating activity of PSMD14

(Model on Fig. 8). We propose that inhibition of PSDM14 DUB activity by CZM is a new

strategy to cause inhibition of the Golgi-to-ER retrograde pathway. Thus, CZM should now

be considered a new pharmacological tool to study the impact of Golgi-to-ER retrograde

**Figure 8**

transport inhibition in other cell biology processes, such as autophagy. Moreover, because CZM was discovered as a new drug for cancer treatment [37], it would be now interesting to investigate whether part of its anti-cancer effects could be the result of inhibiting Golgi-to-ER retrograde pathway. Interestingly, recent studies have shown that PSMD14 is upregulated (mRNA and protein levels) in different tumoral cell types [81,82]. Whether cancer cells are more dependent of Golgi-to-ER retrograde pathway than normal cells, as it happens regarding the mechanisms of protein quality control [83] is still unclear. CZM could offer an interesting tool to evaluate this hypothesis positioning PSMD14 as a promising target for therapeutic intervention.

**Materials and Methods**

***Chemical Reagents***

Earle's balanced salt solution (EBSS) and the cocktail of protease inhibitors were purchased from Sigma-Aldrich (St. Louis, MO, USA). MG132 was purchased from Millipore (Burlington, MA, USA). Torin-1 was purchased from Tocris Bioscience (Bristol, UK). Dr. Raymond Deshaies from California Institute of Technology (Caltech), CA, USA and Dr. Yuyong Ma from University of California, CA, USA kindly donated Capzimin (CZM).

***Antibodies***

We used the following monoclonal antibodies: mouse anti-ubiquitin clone P4D1 (Cytoskeleton, Inc, Denver, Co, USA), mouse anti-β-actin clone BA3R (Thermo Fisher Scientific, Waltham, MA, USA), rabbit anti-RAB1A clone D3X9S, mouse anti-GM130 clone 35/GM130, rabbit anti-PSMD14 clone D18C7 (Cell Signaling Technology, Danvers, MA, USA) and rabbit anti-ATG9A clone EPR2450(2) (Abcam, Cambridge, UK). We used the following polyclonal antibodies: rabbit anti-ubiquitin (cat: Z0458, Dako, Carpintería, CA, USA), rabbit anti-Giantin (cat: AB24586, Abcam, Cambridge, UK), rabbit anti-LC3 (cat: 2775S, Cell Signaling Technology, Danvers, MA, USA), rabbit anti-APP CT695 (cat: 51-2700, Thermo Fisher Scientific). Horseradish Peroxidase-conjugated secondary antibodies were purchased from Jackson ImmunoResearch Laboratories (West Grove, PA, USA), and DAPI probe, Alexa and Dylight fluorophore-conjugated secondary antibodies were purchased from Thermo Fisher Scientific.

 *Cell Culture*

 H4 ATCC ® HTB-148™ Homo sapiens brain neuroglioma cells, referred here as H4

 human neuroglioma cells and HeLa ATCC ® CCL-2™ cells were obtained from the

 American Type Culture Collection (Manassas, VA, USA). Cell lines were cultured in

 Dulbecco's modified Eagle's medium (DMEM; Thermo Fisher Scientific) supplemented

 with 10% (vol/vol) heat-inactivated fetal bovine serum (FBS; Thermo Fisher Scientific),

 and penicillin/streptomycin (Thermo Fisher Scientific), in a 5% $CO_2$ atmosphere at 37°C.

 The generation of the H4 stable cell line expressing hemagglutinin-tagged $APP_{695}$-F/P-

 D/A-EGFP (APP-EGFP) and the HeLa stable cell line expressing KDELR1-GFP were

 previously reported [32,44,84]. Stably transfected cells were maintained in culture medium

 supplemented with 100 μg/ml G418. Cells were grown to sub confluence and then treated

 with drugs or transfected with siRNAs for further western blot and immunofluorescence

 analyses. Nutrient starvation assays were performed in the presence of EBSS. Assays to

 detect *Mycoplasm* were performed periodically.

 *High Content siRNA Transfection and Imaging*

 The primary siRNA screen for levels of APP-EGFP were performed in duplicates with our

 custom-assembled "Ubiquitinome" siRNA library, which consists of 1,187 SMARTpools

 siRNAs targeting all known and assumed components of the ubiquitin and ubiquitin-like

 systems in 96-well format, previously reported [30,31]. The H4 cells stably expressing APP-

 EGFP were reverse transfected [30,31] in μClear bottom 96-well plates (Greiner Bio-One,

 Kremsmünster, Austria) with ON- TARGETplus SMARTpools siRNAs (GE Dharmacon,

 Lafayette, CO, USA). Additionally, ON-TARGETplus Non-targeting pool (NT siRNA)

 was used together with untransfected cells (Mock) as neutral control; siRNA against GFP

 (GFP siRNA) was used as positive control for APP downregulation. Briefly, 10 μl siRNA

 (200 nM) were stamped from 96-well plates of the library on to μClear bottom 96 well

 plates. Thereafter, 10 μl of Opti-MEM I Reduced Serum Medium (Thermo Fisher

 Scientific) containing Lipofectamine RNAiMAX transfection reagent (Thermo Fisher

 Scientific) was added to the siRNA (dilution 1:50). Plates were shaken at 900 rpm for 1

 min followed by incubation for 20 min at room temperature. During this incubation cells

 were resuspended in DMEM supplemented with 10% (vol/vol) FBS and 1x Normocin

 (InvivoGen, San Diego, CA, USA) and 6,000 cells were dispensed onto each well loaded

with medium for a final siRNA concentration of 20 nM. Plates were incubated for 72 h and

further prepared for high content measurement.

After transfection, cells were stained using 100 µl of reagent per step, dispensed by the

automated reagent dispenser XPP-721 (fluidX, Manchester, UK), according to the

following protocol: two washes in phosphate buffered saline (PBS); fixation in 3.7%

(vol/vol) paraformaldehyde for 10 min; one wash in PBS; incubation with 0.2% (vol/vol)

Triton X-100 in PBS for 10 min; two washes in PBS; incubation with 0.1 µg/ml DAPI for 5

min; two washes with PBS. Finally, 100 µl of PBS were left in each well. Images were

acquired and analyzed using the automated microscope IN Cell 2000 Analyzer (GE

Healthcare, Little Chalfont, UK). Images of six randomized fields per well having ~ 600

cells each were acquired, and further analyzed the total fluorescence intensity in n > 2,000

cells per condition. A secondary siRNA screening was performed in triplicate targeting the

35 most responsive hits, using each single siRNA duplex derived from the SMARTpools

used in the primary siRNA screening.

***siRNA Transfection for the siRNA Screening Validation Stage***

Four single siRNA sequences targeting human PSMD14 (Accession number:

NM_005805), derived from the ON-TARGETplus SMARTpool used in the siRNA

Screening (Suppl. Fig. 7) were purchased from GE Dharmacon (Lafayette, CO, USA).

siRNA transfections were carried out in 60 mm tissue culture plates using the

Lipofectamine RNAiMax transfection reagent (Thermo Fisher Scientific) according to the

manufacturer's protocol, and after 72 h cells were collected for further analysis.

***RNA isolation and RT-qPCR Analysis***

Total RNA extraction from H4 cells was carried out using the E.Z.N.A.® Total RNA Kit I

(Omega Biotek, Norcross, GA, USA), and either purity (260nm/280nm ratio and

260nm/230nm ratio) and quantity (260nm absorbance) were determined by

spectrophotometry using NanoVue Spectrophotometer (GE Healthcare). The cDNA

synthesis was performed from 2.5 µg of total RNA and oligo-dT and MMLV reverse

transcriptase (Promega, Madison, WI, USA) according to supplier instructions. Specific

primer pairs for *tbp* (NM_003194), *psmd14* (NM_005805) and *app* (NM_000484) human

genes were designed for quantitative reverse transcription PCR on cDNA template (RT-

qPCR) (Suppl. Fig. 8). First, the specificity of amplicons was verified by cloning and

| | Sequence |
|---|---|
| **siRNA #1** | (5´-GAACAAGUCUAUAUCUCUU-3´) |
| **siRNA #2** | (5´-GGCAUUAAUUCAUGGACUA-3´) |
| **siRNA #3** | (5´-AGAGUUGGAUGGAAGGUUU-3´) |
| **siRNA #4** | (5´-GAUGGUUGUUGGUUGGUAU-3´) |

**Supplementary Figure 7**

| Target | Sequence |
|---|---|
| **hTBP1** | f(5´-TAGTCCAATGATGCCTTACG-3´) <br> r(5´-TGGTCAGAGTTTGAGAATGG-3´) |
| **hPSMD14** | f(5´-ACCTTAAGAGTTGTAGTTACTGACC-3´) <br> r(5´-TTTAACAGTGCCAGGGAAGAG-3´) |
| **hAPP** | f(5´-CCTAAAGCATTTCGAGCATG-3´) <br> r(5´-GTTTCCGTAACTGATCCTTG-3´) |

# Supplementary Figure 8

sequencing, including *tbp* (223bp), *psmd14* (150bp) and *app* (247bp). mRNA levels were

quantified in cDNA by qPCR with GoTaq qPCR Master Mix (Promega) according to

supplier's instructions in a Mx3000 Real-Time Thermocycler (Stratagene, San Diego, CA,

USA). In a 40 cycles PCR reaction, each cycle consisted in 20 s at 94°C, 15 s at 55°C and

15 s at 72°C, followed by a final heating at 95°C, revealing melting curves that confirmed

single amplification products. All analyses were performed in triplicate. The expression

level of each gene was normalized to *tbp* expression as reference gene using exon-spanning

primers to control for genomic DNA contamination since no DNAse treatment of total

RNA was included. RT-qPCR assays were analyzed with 2(-ΔΔCt) method [85] via MxPro

software (Stratagene) and expressed as relative quantity to normalizer [86].

***Preparation of protein extracts, Electrophoresis, SDS-PAGE and Western Blot analysis***

Cells were washed in ice-cold Phosphate Buffered Saline (PBS) and lysed at 4°C in lysis

buffer (50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1 mM EDTA, 1% (vol/vol) Triton X-100)

supplemented with a cocktail of protease inhibitors (Sigma-Aldrich). All lysates were

cleared by centrifugation at 16,000 x *g* for 20 min at 4°C, and protein concentration was

determined with a protein assay dye reagent (Bio-Rad Laboratories, Hercules, CA, USA).

Samples with an equivalent amount of protein were boiled for 5 min with Laemmli SDS-

PAGE sample buffer, and then analyzed by SDS-PAGE. Proteins were electroblotted onto

nitrocellulose membranes, blocked by incubation for 30 min in PBS containing 5% (wt/vol)

free-fat dry milk, and incubated sequentially with primary and secondary antibodies, both

diluted in blocking solution, for 1 h at room temperature, or overnight at 4°C.

Chemiluminescence protein detection was performed using SuperSignal West Pico

(Thermo Fisher Scientific). β-Actin was used as an internal loading control.

***In vitro Proteasomal Activity Assay***

Proteasome activity was quantitatively assessed in H4 cell extracts using the β5-selective

fluorogenic substrate succinyl-leucine-leucine-valine-tyrosine-4-methyl-7-courmarylamide

(Suc-LLVY-AMC, Calbiochem, Burlington, MA, USA) using an adapted protocol [87].

Briefly, cells were left untreated or treated for 4 h with different amounts of CZM (between

2 and 10 μM) or with 10 μM MG132. Cells were lysed in lysis buffer (20 mM Tris-HCl,

pH 7.2, 1 mM EDTA, 1 mM dithiothreitol (DTT) and 0.1% (vol/vol) Nonidet P-40)

supplemented with a cocktail of protease inhibitors (Sigma-Aldrich). Lysates were cleared

489 by centrifugation at 16,000 x $g$ for 20 min at 4°C, and protein concentration was

490 determined with a protein assay dye reagent (Bio-Rad Laboratories, Hercules, CA, USA).

491 Soluble extracts (20 μg) were incubated for 60 min at 37°C in proteasome assay buffer (50

492 mM Tris-HCl, pH 7.2, 1 mM DTT, 0,5 mM EDTA and 100 μM Suc-LLVY-AMC) all

493 dispensed in triplicate into a 96-well black opaque plate (Nunc, Thermo Fisher Scientific).

494 Proteasome activity was measured by monitoring the production of free AMC using a

495 Synergy HT Multi-detection Microplate Reader (BioTek Instruments, VT, USA) with

496 excitation and emission wavelengths of 360 nm and 460 nm, respectively, and the data

497 were obtained by Gen5 Version 2.09.1 data analysis software. Statistical significance was

498 determined by One-Way ANOVA, followed by Dunnett's test. Value of P <0,01(**) and P

499 <0,001(***) were regarded as statistically significant and is indicated in the figure.

500 *Immunofluorescence*

501 Cells grown on glass coverslips were washed with PBS and fixed in 4% (vol/vol)

502 paraformaldehyde for 30 min at room temperature. After fixation, cells were washed in

503 PBS and permeabilized with 0.2% (vol/vol) Triton X-100 in PBS for 10 min at room

504 temperature. Cells were incubated with the indicated primary antibodies diluted in

505 immunofluorescence buffer (PBS containing 10% (vol/vol) FBS and 0.1% (wt/vol)

506 Saponin) for 30 min at 37°C. Coverslips were washed in PBS and incubated with the

507 corresponding Alexa-conjugated secondary antibody diluted in immunofluorescence buffer

508 for 30 min at 37°C. For nuclei staining, cells were washed with PBS and incubated for 10

509 min at room temperature with 0.1 mg/ml DAPI. After the final wash, coverslips were

510 mounted onto glass slides with Fluoromount-G (SouthernBiotech, Birmingham, AL, USA).

511 *Fluorescence Microscopy*

512 Images of fixed cells were acquired by using a TCS SP8 laser-scanning confocal

513 microscope (Leica Microsystems, Wetzlar, Germany) equipped with a 63X oil immersion

514 objective (1.4 NA), 405 nm, 488 nm and 561 nm laser lines, with Photomultiplier (PMT), a

515 hybrid detector system (Leica HyD) and the Leica Application Suite LAS X software. For

516 quantification of fluorescent signals, 8-bit images were acquired under identical settings

517 avoiding signal saturation and corrected for background signal on each image. The

518 corrected fluorescent signal in each cell of each image was used in Image J (version 1.44o;

519 Wayne Rasband, NIH, http://imagej.nih.gov) to determine the total integrated pixel

intensity per cell area. Colocalization analyses were performed with sets of

immunofluorescence images (Z-stack, with 0.2 μm intervals) of the same cells for each

marker. Quantification of the acquired images was performed with the ICY software

(Quantitative Image Analysis Unit, Institut Pasteur, http://icy.bioimageanalysis.org/) using

protocols plugin to create a pipeline to analyze the images in batch, active contours plugin

was used to perform the cell segmentation, hk-means plugin was used for threshold

detection, wavelet spot detector plugin was used for spot detection and colocalization

studio plugin for colocalization analysis.

For live cell imaging assays, H4 cells were grown in glass bottom culture dishes (MatTek

Corporation, Ashland, MA, USA) and transiently transfected for 48-h with KDELR-

VSVG-YFP, using TransIT-LT1 Transfection Reagent (Mirus Bio LLC, Madison, Wi,

USA) according to the manufacturer's protocol. Before the live cell imaging assay, the

culture medium was replaced with phenol red-free DMEM supplemented with HEPES

(10mM, pH 7.4), and the cells were treated with CZM (10μM) at 32°C in a controlled

temperature chamber in the TCS SP8 laser-scanning confocal microscope. Cells were kept

at 32°C to allow KDELR-VSVG-YFP localization on the Golgi, followed by a shift in

temperature to 40°C. Imaging was done with a 63X oil immersion objective (1.4 NA),

running the Leica Application Suite LAS X software, acquiring 8-bit images at 1-min

interval for 15 min at 40°C (488 laser for excitation; HyD: 510-550 nm; 1024×1024 pixels;

frame average 1). Quantification of the acquired images was performed with the

MetaMorph Software version 7.0.

***3D Golgi Reconstruction and Golgi Volume and Area Measurements***

H4 and HeLa cells stably expressing KDELR1-GFP, under the specified conditions, were

fixed and immunostained with anti-GM130 or anti-Giantin, respectively. The

immunofluorescence protocol was performed as described above to visualize the Golgi

structure. For Golgi volume and area measurements, Z-stack (250 nm) fluorescence images

were acquired by using a TCS SP8 laser-scanning confocal microscope (Leica

Microsystems, Wetzlar, Germany) equipped with a 63X oil immersion objective (1.4 NA)

running the Leica Application Suite LAS X software. Images were then processed with

ImageJ software version FIJI to remove the background by using threshold. The

thresholded images were then visualized using the "3D Viewer" plugin. The Golgi volume

was quantified using ImageJ software version FIJI by setting a thresholded region to select

only the Golgi fluorescence. Then, the Golgi structures were separated by ROI (Regions of

Interest) and the individual Golgi volume was measured with the plugin "Voxel Counting".

The Golgi volume in $\mu m^3$ was determined by the number of voxels contained in the stack of

images (voxel 0.2x0.2x0.25 nm). The Golgi Area was quantified using ICY software and

the Golgi marker GM130 was used to determine the ROI. To separate the specific signal

from background "k means threshold" plugin was used, using the same threshold level to

all images. Data analysis was performed using GraphPad Prism 6 (GraphPad Software, La

Jolla, CA, USA) and the results represented in graphs depicting the mean ± SEM of at least

20 cells. Statistical significance of the data was determined with Student's T-test. The value

of P <0,001(***) was regarded as statistically significant and is indicated in the respective

figures.

***Densitometric Quantification and Statistical Analysis***

The amount of immunoblot signal was estimated using Image J software version 1.48v

(Wayne Rasband, NIH, http://imagej.nih.gov). For each condition, protein bands were

quantified from at least three independent experiments in order to ensure adequate

statistical power. Data analysis was performed using Microsoft Excel 2013 for Windows

(Redmond, WA, USA) or GraphPad Prism 6. Results are represented in graphs depicting

the mean ± standard deviation. Statistical significance of data comparisons from two groups

comparisons was determined with Student's T-test for parametric data. Values of P <0.05

(*), P <0.01 (**), P <0.001 (***) were regarded as statistically significant and are indicated

in the figures. Statistical significance of data from many groups was analyzed using One-

Way ANOVA, followed by Tukey's test in order to evaluate pair-wise comparisons. The

value of P <0.05 was regarded as statistically significant and indicated in the figure in

different letters above bars mean.

**Author Contributions:** Conceptualization, B.H.A., R.F.A. and B.P.V.; Experimental

design, B.H.A., R.F.A., M.G.A., C.J., and B.P.V.; Execution of experiments, B.H.A.,

R.F.A., B.S., C.K., V.G.E., A.M.E., C.T.C., Data analysis, B.H.A., C.K., G.A.E., V.G.E.,

C.Y., H.S., A.M.E., C.T.C. Reagents, Materials, and Analysis tools, V.G.E., H.S., B.S.,

A.S., K.G., M.G.A., C.J., H.R.T., R.F.A. and B.P.V. writing—original draft preparation,

B.H.A., R.F.A. and B.P.V.; writing—review and editing, B.H.A., B.S., A.S., K.B., M.G.A.,

C.J., R.F.A., H.R.T and B.P.V.; funding acquisition, B.P.V., R.F.A and H.R.T.

**Funding Statement:** This work was funded by Fondo Nacional de Desarrollo Científico y

Tecnológico of Chile (FONDECYT; http://www.conicyt.cl/fondecyt) No. 1171649 to

B.P.V. & No. 11150532 to R.F.A.; Associative Investigation Program (PIA;

https://www.conicyt.cl/pia) including No. ACT-172066 to B.P.V. &No. AFB-170005 to

B.P.V.; Academy Insertion Program (PAI; https://www.conicyt.cl/pai) No. 79150075 to

R.F.A.; Fondo de Equipamiento Científico y Tecnológico of Chile (FONDEQUIP;

http://www.conicyt.cl/fondequip) No. EQM150118 to B.P.V.; Cooperation International

Programme (CONICYT-RCUK; https://www.conicyt.cl/pci) No. DPI20140068 to B.P.V.;

B.H.A, G.A.E., and V.G.E. were supported by National Ph.D Fellowships No. 21130315,

201110746 and 21130511 respectively (CONICYT; https://www.conicyt.cl/becasconicyt);

Programa de Mejoramiento de la Calidad y la Equidad de la Educación superior,

MECESUP AUS1203 and Vicerrectoría de Investigación de la Universidad Austral de

Chile No. D #2015-02 to B.H.A., D #2013-07 to A.E.G. and D #2015-05 to V.G.E. The

funders had no role in study design, data collection and analysis, decision to publish, or

preparation of the manuscript.

**Acknowledgments:** We thank Gonzalo Astroza (Universidad Austral de Chile) and Ellis

Jaffray (GRE, University of Dundee) for technical assistance. Dr. Raymond Deshaies

(Division of Biology & Biological Engineering, California Institute of Technology) & Dr.

Yuyong Ma (Department of Chemistry and Biochemistry, University of California) for

proving Capzimin for this study.

**Conflicts of Interest:** The authors declare no conflict of interest.

**Abbreviations**

APP: Amyloid Precursor Protein

CZM: Capzimin

DUB: Deubiquitinating enzyme

EBSS: Earle's balanced salt solution

EGFP: Enhanced Green Fluorescent Protein

ER: Endoplasmic Reticulum

FBS: Fetal bovine serum

GFP: Green Fluorescent Protein

GM130: Golgi matrix protein GM130

HCS: High-Content siRNA Screening

ILV: intraluminal vesicle

K: Lysine residue

KDELR1: KDEL (Lys-Asp-Glu-Leu) Receptor 1

KD: Knock-down

LC3B: Microtubule-associated protein 1 light chain 3B

mTORC: mammalian target of rapamycin complex

MVB: multivesicular body

NT: Non-Target

PBS: Phosphate Buffered Saline

PCR: Polymerase Chain Reaction

PTM: Post-translational Modification

RP:  19S regulatory particle

RT-qPCR: Quantitative reverse transcription PCR

SENP: SUMO-specific protease

siRNA: Small interfering RNA

Ub: Ubiquitin

UBD: Ub-binding domain

ULK1: unc-51-like kinase 1

VSVG: Vesicular Stomatitis Virus G

YFP: Yellow Fluorescent Protein

 **Figure Legends**

**Figure 1. High-Content siRNA Screening assay revealed PSMD14 as a novel regulator**
**of APP levels. (A)** Graphical distribution of the targets evaluated in the primary High
Content siRNA screening using the siRNA "ubiquitinome" library in H4 cells. **(B)**
Quantification of the total fluorescence intensity of reporter APP-EGFP cells transfected
for 72 h with NT siRNA and EGFP siRNA. Bars represent the mean ± SD with a statistical
Z factor = 0.69. **(C)** High content images (20X) captured in reporter APP-EGFP cells
transfected for 72 h with NT siRNA and EGFP siRNA. **(D)** Graphical representation of
total fluorescence intensity of all 1,187 genes analyzed in primary siRNA screening with
the reporter APP-EGFP cells. PSMD14 appears indicated as the top hit. **(E)** High content
images (20X) in pseudo color of reporter APP-EGFP cells transfected for 72 h with siRNA
SMARTpool targeted against PSMD14 (PSMD14 siRNA) in comparison to cells
transfected with NT siRNA. Scale Bar of the images indicates the scale of fluorescence
intensity.

**Figure 2. PSMD14 is validated as a regulator of the endogenous APP levels.**
**(A)** Protein extracts of parental H4 cells either untransfected (Mock), transfected with NT
siRNA, or transfected with four different PSMD14 siRNA sequences for 72 h were
analyzed by western blot. Polyclonal antibodies to endogenous APP (CT695) and to
Ubiquitin, and monoclonal antibodies to PSMD14 (clone D18C7) and to β-actin (clone
BA3R), were tested. The position of molecular mass markers are indicated on the left.
Densitometric quantification of the levels of endogenous APP protein levels **(B)** and
PSMD14 **(C)** in H4 cells transfected with PSMD14 siRNA#1, compared to untransfected
cells (Mock). Statistical significance was determined by Student's t-test. Bars represent the
mean ± SD of biological replicates (APP n =5; PSMD14 n=4). **P <0.01 and ***P <0.001.
**(D)** mRNA levels of *psmd14* and **(E)** mRNA levels of *app* were measured using RT-qPCR
from parental H4 cells transfected for 72 h. All data were normalized for TATA binding
protein expression in either untransfected cells (Mock), cells transfected with NT siRNA or
cells transfected with four different PSMD14 siRNAs duplexes. Statistical significance was
determined by One-Way ANOVA, followed by Tukey's test. Bars represent the mean ± SD

of biological replicates (*psmd14* n=3; *app* n=3). Different letters above the mean bars apply

to significant differences between groups P <0.01.

**Figure 3. Acute inhibition of PSMD14 by CZM shows a similar phenotype as PSMD14**

**KD on the levels of APP and high molecular weight Ub conjugates.**

**(A)** Schematic diagram of the molecular targets of Capzimin and MG132 in the 19S RP

and 20S catalytic core of the proteasome, respectively. **(B)** Parental H4 cells were treated

either with vehicle (DMSO; Control), or increasing doses of CZM for 4 h, or MG132 for 6

h. Protein extracts were analyzed by western blot with a polyclonal antibody to endogenous

APP. Monoclonal antibody to β-actin (clone BA3R) was used as a loading control. The

position of molecular mass markers is indicated on the left. **(C)** Densitometric

quantification of APP protein levels as shown in (D). Statistical significance was

determined by One-Way ANOVA, followed by Tukey's test. Bars represent the mean ± SD

of biological replicates (n=4). Different letters above the mean bars apply to significant

differences between groups P <0.05. **(D)** Parental H4 cells were treated as in (B), and the

protein extracts were analyzed by western blot with a polyclonal antibody to Ub.

Monoclonal antibody to β-actin (clone BA3R) was used as a loading control. The position

of molecular mass markers is indicated on the left. **(E)** Immunofluorescence microscopy

images of the cellular localization of Ub in parental H4 cells treated with either the vehicle

(DMSO; Control), CZM for 4 h or MG132 for 6 h. Cells were fixed, permeabilized and

stained with a mouse monoclonal antibody to Ub (clone P4D1) followed by Alexa-488-

conjugated donkey anti-mouse IgG. Scale bar, 10 μm. (n=3).

**Figure 4. Acute inhibition of PSMD14 by CZM triggers the accumulation of APP in a**

**swollen Golgi apparatus.**

Immunofluorescence analysis of endogenous APP in H4 parental cells treated either with

the vehicle (DMSO; Control) **(A-C)** or CZM **(D-F)** for 4 h. Cells were fixed,

permeabilized, and double stained with a rabbit polyclonal antibody to APP (CT695) (A

and D) and a mouse monoclonal antibody to GM130 (clone35/GM130) (B and E), followed

by Alexa-594-conjugated donkey anti-Rabbit IgG and Alexa-488-conjugated donkey anti-

Mouse IgG. Merging of the images generated the third picture (C and F). Scale bar, 10 μm.

**(G)** Quantitative analysis of the mean of total fluorescence intensity of APP upon treatment

with CZM, in comparison to control cells. The statistical significance was determined by

Student's t-test. Bars represent the mean ± SD of the fluorescent signal per cell area (n=43

cells). ***P <0.001. **(H)** Quantitative analysis of the fraction of APP colocalizing with

GM130 under CZM treatment and compared to control cells. Statistical significance was

determined by Student's t-test. Bars represent the mean ± SD of the fluorescent signal per cell

area (n=43 cells). ***P <0.001. **(I)** Quantitative analysis of the cell area. Statistical significance

was determined by Student's t-test. Bars represent the mean ± SD of the cell area (n=43 cells)

**P <0.001. **(J)** Immunofluorescence microscopy analysis of GM130 in parental H4 cells

treated either with the vehicle (DMSO; Control) or CZM for 4 h. Cells were fixed,

permeabilized and stained with mouse monoclonal antibody to GM130 (clone 35/GM130)

followed by Alexa-488-conjugated donkey anti-mouse IgG, and nuclei were stained with

DAPI. Scale bar, 10 μm. **(K)** 3D reconstructions of the Golgi apparatus using GM130 as

Golgi marker were generated from Z-stacks (250 nm). **(L)** Golgi Volume was measured

from 3D reconstructions as shown in (K). Statistical significance was determined by

Student's t-test. Bars represent the means ± SEM (n=20 cells). *** P <0.001.

**Figure 5. The PSMD14 DUB inhibitor CZM impairs Golgi-to-ER retrograde**
**transport.**

**(A)** 3D reconstructions of the Golgi apparatus using Giantin as Golgi marker were

generated from Z-stacks (250 nm) obtained from HeLa cells stably expressing KDELR1-

GFP treated for 90 min either with vehicle (DMSO; Control), CZM or MG132. **(B)** Golgi

volume was measured from 3D reconstructions as shown in (A). Statistical significance

was determined by Student's t-test. Bars represent the means ± SEM (n=30 cells). *** P

<0.001. **(C)** HeLa cells stably expressing KDELR1-GFP were treated for 90 min either

with vehicle (DMSO; Control), CZM or MG132. Cells were fixed and representative

confocal images were acquired. **(D)** Measurement of Giantin and total KDELR1-GFP total

fluorescent intensity. Statistical significance was determined by Student`s t-test. Bars

represent the means ± SEM (n=34 cells). *** P <0.001. **(E)** H4 cells were transiently

transfected to express the thermo-sensitive retrograde transport reporter KDELR-VSVG-

YFP. Cells were kept at 32°C to allow KDELR-VSVG-YFP localization at the Golgi. Cells

were then shifted to 40°C (restrictive temperature) and images acquired at 1 min interval

for 15 min. **(F)** Quantitative image analysis was performed to measure the integrated

fluorescence of KDELR-VSVG-YFP at the Golgi at 1 min interval for 15 min. Statistical
significance was determined by Student`s t-test. Bars represent the mean ± SEM (n=3
cells). *p<0.05

**Figure 6. Inhibition of autophagosome formation by CZM.**
Immunofluorescence microscopy analysis of the subcellular localization of endogenous
LC3B in parental H4 cells treated with either the vehicle (DMSO; Control) **(A)**, EBSS for 4
h **(B),** CZM for 6 h **(D)** or Torin-1 for 4 h **(E)**. EBSS **(C)** and Torin-1 **(F)** were tested using
a 2-h pretreatment with CZM followed by the treatment with EBSS or Torin-1 for 4 h in the
presence of CZM. Cells were fixed, permeabilized and stained with a rabbit polyclonal
antibody to LC3B followed by Alexa-594-conjugated donkey anti-Rabbit IgG, and nuclei
were stained with DAPI. Scale bar 10 μm. **(G)** Quantification of the puncta positive to
LC3B. Statistical significance was determined by One-Way ANOVA, followed by Tukey's
test. Bars represent the mean ± SEM (n=50 cells). Different letters above the mean bars
indicate the significant differences between groups P <0.05. **(H)** Protein extracts from
parental H4 cells treated as in (A-F) were analyzed by western blot with a rabbit polyclonal
antibody to LC3B. Monoclonal antibody to β-actin (clone BA3R) was used as a loading
control. The position of molecular mass markers is indicated on the left. **(I)** Densitometric
quantification of LC3B-I levels and **(J)** LC3B-II levels. Statistical significance was
determined by One-Way ANOVA, followed by Tukey's test. Bars represent the mean ±
SEM of biological replicates (LC3B-I n=3; LC3B-II n=3). Different letters above the mean
bars indicate the significant differences between groups P <0.05.

**Figure 7. Redistribution of RAB1A and ATG9A to the Golgi apparatus with CZM.**
**(A)** Immunofluorescence analysis of endogenous RAB1A and ATG9A in H4 parental cells
treated for 4 h either with the vehicle (DMSO; Control) (left panel) or CZM (right panel).
Cells were fixed, permeabilized, and stained with a rabbit monoclonal antibody to RAB1A
(clone D3X9S) (upper panel) and a rabbit monoclonal antibody to ATG9A (clone
EPR2450(2)) (lower panel), followed by Alexa-594-conjugated donkey anti-Rabbit IgG.
Scale Bar, 10 μm. (B) Quantitative analysis of the fluorescence intensity of RAB1A upon
treatment with CZM, in comparison to control cells. Statistical significance was determined
by Student's t-test. Bars represent the mean ± SEM of the fluorescent signal per cell area

(n=227 cells). **P <0.01; ***P < 0.001; n.s., not significant. (C) Quantitative analysis of

the fluorescence intensity of ATG9A upon treatment with CZM, in comparison to control

cells. Statistical significance was determined by Student's t-test. Bars represent the mean ±

SEM of the fluorescent signal per cell area (n=95 cells). *P< 0.05; ***P < 0.001; n.s., not

significant.

**Figure 8. Model of mechanism underlying regulation of protein membrane trafficking**

**and macroautophagy by the proteasome 19S RP PSMD14 DUB activity.** The model

depicts the closed interplay between membrane transport and macroautophagy by a novel

mechanism involving the proteasome complex through the deubiquitinating activity of

PSMD14. We propose that active PSMD14 and their K63-Ub chains (1, left panel)

positively regulate Golgi-to-ER retrograde transport (2, left panel), pathway implicated in

the retrieval of key proteins for autophagosome biogenesis and macroautophagy (3, left

panel). Reduction of free K63-Ub chains by inactive PSM14 (1, right panel) results on the

blockage of Golgi-to-ER retrograde transport (2, right panel) causing the accumulation of

ATG9A and RAB1A at the Golgi apparatus. Thus, blockage of Golgi-to-ER retrograde

transport inhibits the biogenesis of autophagosomes and macroautophagy (3, right panel).

Macroautophagy has been recently demonstrated to act as a potent positive regulator of

protein transport from the Golgi apparatus to the cell surface (Golgi-secretion ON; 4, left

panel). Thus, inhibition of macroautophagy upon inactive PSMD14 (siRNA/CZM) blocks

protein transport from the Golgi apparatus to the cell surface (Golgi-secretion OFF)

explaining the effect on APP transport (5 left and 5 right).

**Bibliography**

1.  Shinde, S. R. & Maddika, S. Post translational modifications of Rab GTPases. *Small*
*GTPases* 9, 49–56 (2018).
2.  Luo, P. M. & Boyce, M. Directing Traffic: Regulation of COPI Transport by Post-
translational Modifications. *Front. Cell Dev. Biol.* 7, 190 (2019).
3.  Li, M. *et al.* Mono- versus polyubiquitination: differential control of p53 fate by
Mdm2. *Science* 302, 1972–1975 (2003).
4.  Komander, D. & Rape, M. The ubiquitin code. *Annu. Rev. Biochem.* 81, 203–229
(2012).
5.  Foot, N., Henshall, T. & Kumar, S. Ubiquitination and the regulation of membrane
proteins. *Physiol. Rev.* 97, 253–281 (2017).
6.  Pelham, H. R. B. Membrane traffic: GGAs sort ubiquitin. *Curr. Biol.* 14, R357–9
(2004).
7.  Yang, B. & Kumar, S. Nedd4 and Nedd4-2: closely related ubiquitin-protein ligases
with distinct physiological functions. *Cell Death Differ.* 17, 68–77 (2010).
8.  Tan, J. & Evin, G. B-site APP-cleaving enzyme 1 trafficking and Alzheimer's disease
pathogenesis. *J. Neurochem.* 120, 869–880 (2012).
9.  Raiborg, C. & Stenmark, H. The ESCRT machinery in endosomal sorting of
ubiquitylated membrane proteins. *Nature* 458, 445–452 (2009).
10.  Ren, X. & Hurley, J. H. VHS domains of ESCRT-0 cooperate in high-avidity binding
to polyubiquitinated cargo. *EMBO J.* 29, 1045–1054 (2010).
11.  Clague, M. J., Liu, H. & Urbé, S. Governance of endocytic trafficking and signaling by
reversible ubiquitylation. *Dev. Cell* 23, 457–467 (2012).
12.  Haglund, K., Di Fiore, P. P. & Dikic, I. Distinct monoubiquitin signals in receptor
endocytosis. *Trends Biochem. Sci.* 28, 598–603 (2003).
13.  Clague, M. J. & Urbé, S. Ubiquitin: same molecule, different degradation pathways.
*Cell* 143, 682–685 (2010).
14.  Kwon, Y. T. & Ciechanover, A. The Ubiquitin Code in the Ubiquitin-Proteasome
System and Autophagy. *Trends Biochem. Sci.* 42, 873–886 (2017).
15.  Ciechanover, A. & Kwon, Y. T. Degradation of misfolded proteins in
neurodegenerative diseases: therapeutic targets and strategies. *Exp Mol Med* 47, e147
(2015).
16.  Clague, M. J. & Urbé, S. Endocytosis: the DUB version. *Trends Cell Biol.* 16, 551–
559 (2006).
17.  Komander, D., Clague, M. J. & Urbé, S. Breaking the chains: structure and function of
the deubiquitinases. *Nat. Rev. Mol. Cell Biol.* 10, 550–563 (2009).
18.  Mevissen, T. E. T. & Komander, D. Mechanisms of deubiquitinase specificity and
regulation. *Annu. Rev. Biochem.* 86, 159–192 (2017).
19.  Bienko, M. *et al.* Ubiquitin-binding domains in Y-family polymerases regulate
translesion synthesis. *Science* 310, 1821–1824 (2005).
20.  Bienko, M. *et al.* Regulation of translesion synthesis DNA polymerase eta by
monoubiquitination. *Mol. Cell* 37, 396–407 (2010).
21.  Husnjak, K. & Dikic, I. Ubiquitin-binding proteins: decoders of ubiquitin-mediated
cellular functions. *Annu. Rev. Biochem.* 81, 291–322 (2012).

22. Lai, A., Sisodia, S. S. & Trowbridge, I. S. Characterization of sorting signals in the
821           beta-amyloid precursor protein cytoplasmic domain. *J. Biol. Chem.* 270, 3565–3573
822           (1995).
823    23. Perez, R. G. *et al.* Mutagenesis identifies new signals for beta-amyloid precursor
824           protein endocytosis, turnover, and the generation of secreted fragments, including
825           Abeta42. *J. Biol. Chem.* 274, 18851–18856 (1999).
826    24. Burgos, P. V. *et al.* Sorting of the Alzheimer's disease amyloid precursor protein
827           mediated by the AP-4 complex. *Dev. Cell* 18, 425–436 (2010).
828    25. Watanabe, T., Hikichi, Y., Willuweit, A., Shintani, Y. & Horiguchi, T. FBL2 regulates
829           amyloid precursor protein (APP) metabolism by promoting ubiquitination-dependent
830           APP degradation and inhibition of APP endocytosis. *J. Neurosci.* 32, 3352–3365
831           (2012).
832    26. El Ayadi, A., Stieren, E. S., Barral, J. M. & Boehning, D. Ubiquilin-1 regulates
833           amyloid precursor protein maturation and degradation by stimulating K63-linked
834           polyubiquitination of lysine 688. *Proc. Natl. Acad. Sci. USA* 109, 13416–13421
835           (2012).
836    27. Morel, E. *et al.* Phosphatidylinositol-3-phosphate regulates sorting and processing of
837           amyloid precursor protein through the endosomal system. *Nat. Commun.* 4, 2250
838           (2013).
839    28. Williamson, R. L. *et al.* Disruption of amyloid precursor protein ubiquitination
840           selectively increases amyloid β (Aβ) 40 levels via presenilin 2-mediated cleavage. *J.*
841           *Biol. Chem.* 292, 19873–19889 (2017).
842    29. Cooper, E. M. *et al.* K63-specific deubiquitination by two JAMM/MPN+ complexes:
843           BRISC-associated Brcc36 and proteasomal Poh1. *EMBO J.* 28, 621–631 (2009).
844    30. Bett, J. S., Ibrahim, A. F. M., Garg, A. K., Rocha, S. & Hay, R. T. SiRNA screening to
845           identify ubiquitin and ubiquitin-like system regulators of biological pathways in
846           cultured mammalian cells. *J. Vis. Exp.* (2014). doi:10.3791/51572
847    31. MacKay, C. *et al.* E3 ubiquitin ligase HOIP attenuates apoptotic cell death induced by
848           cisplatin. *Cancer Res.* 74, 2246–2257 (2014).
849    32. Prabhu, Y. *et al.* Adaptor protein 2-mediated endocytosis of the β-secretase BACE1 is
850           dispensable for amyloid precursor protein processing. *Mol. Biol. Cell* 23, 2339–2351
851           (2012).
852    33. González, A. E. *et al.* Autophagosomes cooperate in the degradation of intracellular C-
853           terminal fragments of the amyloid precursor protein via the MVB/lysosomal pathway.
854           *FASEB J.* 31, 2446–2459 (2017).
855    34. Moffat, J., Reiling, J. H. & Sabatini, D. M. Off-target effects associated with long
856           dsRNAs in Drosophila RNAi screens. *Trends Pharmacol. Sci.* 28, 149–151 (2007).
857    35. Verma, R. *et al.* Role of Rpn11 metalloprotease in deubiquitination and degradation by
858           the 26S proteasome. *Science* 298, 611–615 (2002).
859    36. Yao, T. & Cohen, R. E. A cryptic protease couples deubiquitination and degradation
860           by the proteasome. *Nature* 419, 403–407 (2002).
861    37. Li, J. *et al.* Capzimin is a potent and specific inhibitor of proteasome isopeptidase
862           Rpn11. *Nat. Chem. Biol.* 13, 486–493 (2017).
863    38. Tsubuki, S., Saito, Y., Tomioka, M., Ito, H. & Kawashima, S. Differential inhibition of
864           calpain and proteasome activities by peptidyl aldehydes of di-leucine and tri-leucine. *J.*
865           *Biochem.* 119, 572–576 (1996).

39. Kisselev, A. F. & Goldberg, A. L. Proteasome inhibitors: from research tools to drug
candidates. *Chem. Biol.* 8, 739–758 (2001).
40. Wieland, F. T., Gleason, M. L., Serafini, T. A. & Rothman, J. E. The rate of bulk flow
from the endoplasmic reticulum to the cell surface. *Cell* 50, 289–300 (1987).
41. Martínez-Menárguez, J. A., Geuze, H. J., Slot, J. W. & Klumperman, J. Vesicular
tubular clusters between the ER and Golgi mediate concentration of soluble secretory
proteins by exclusion from COPI-coated vesicles. *Cell* 98, 81–90 (1999).
42. Klumperman, J. Transport between ER and Golgi. *Curr. Opin. Cell Biol.* 12, 445–449
(2000).
43. Thor, F., Gautschi, M., Geiger, R. & Helenius, A. Bulk flow revisited: transport of a
soluble protein in the secretory pathway. *Traffic* 10, 1819–1830 (2009).
44. Cancino, J. *et al.* Control systems of membrane transport at the interface between the
endoplasmic reticulum and the Golgi. *Dev. Cell* 30, 280–294 (2014).
45. Tenorio, M. J., Luchsinger, C. & Mardones, G. A. Protein kinase A activity is
necessary for fission and fusion of Golgi to endoplasmic reticulum retrograde tubules.
*PLoS One* 10, e0135260 (2015).
46. Hsu, V. W., Shah, N. & Klausner, R. D. A brefeldin A-like phenotype is induced by
the overexpression of a human ERD-2-like protein, ELP-1. *Cell* 69, 625–635 (1992).
47. Cole, N. B., Ellenberg, J., Song, J., DiEuliis, D. & Lippincott-Schwartz, J. Retrograde
transport of Golgi-localized proteins to the ER. *J. Cell Biol.* 140, 1–15 (1998).
48. Lewis, M. J., Rayner, J. C. & Pelham, H. R. A novel SNARE complex implicated in
vesicle fusion with the endoplasmic reticulum. *EMBO J.* 16, 3017–3024 (1997).
49. He, S. *et al.* PtdIns(3)P-bound UVRAG coordinates Golgi-ER retrograde and Atg9
transport by differential interactions with the ER tether and the beclin 1 complex. *Nat.*
*Cell Biol.* 15, 1206–1219 (2013).
50. Lemus, L., Ribas, J. L., Sikorska, N. & Goder, V. An ER-Localized SNARE Protein Is
Exported in Specific COPII Vesicles for Autophagosome Biogenesis. *Cell Rep.* 14,
1710–1722 (2016).
51. Chen, Q. *et al.* ATL3 Is a Tubular ER-Phagy Receptor for GABARAP-Mediated
Selective Autophagy. *Curr. Biol.* 29, 846–855.e6 (2019).
52. Hao, R. *et al.* Proteasomes activate aggresome disassembly and clearance by producing
unanchored ubiquitin chains. *Mol. Cell* 51, 819–828 (2013).
53. Nanduri, P., Hao, R., Fitzpatrick, T. & Yao, T.-P. Chaperone-mediated 26S
proteasome remodeling facilitates free K63 ubiquitin chain production and aggresome
clearance. *J. Biol. Chem.* 290, 9455–9464 (2015).
54. Kabeya, Y. *et al.* LC3, a mammalian homologue of yeast Apg8p, is localized in
autophagosome membranes after processing. *EMBO J.* 19, 5720–5728 (2000).
55. Munafó, D. B. & Colombo, M. I. A novel assay to study autophagy: regulation of
autophagosome vacuole size by amino acid deprivation. *J. Cell Sci.* 114, 3619–3629
(2001).
56. Thoreen, C. C. *et al.* An ATP-competitive mammalian target of rapamycin inhibitor
reveals rapamycin-resistant functions of mTORC1. *J. Biol. Chem.* 284, 8023–8032
(2009).
57. Winslow, A. R. *et al.* α-Synuclein impairs macroautophagy: implications for
Parkinson's disease. *J. Cell Biol.* 190, 1023–1037 (2010).
58. Webster, C. P. *et al.* The C9orf72 protein interacts with Rab1a and the ULK1 complex
to regulate initiation of autophagy. *EMBO J.* 35, 1656–1676 (2016).

59. Orsi, A. *et al.* Dynamic and transient interactions of Atg9 with autophagosomes, but
not membrane integration, are required for autophagy. *Mol. Biol. Cell* 23, 1860–1873
(2012).

60. Kishi-Itakura, C., Koyama-Honda, I., Itakura, E. & Mizushima, N. Ultrastructural
analysis of autophagosome organization using mammalian autophagy-deficient cells. *J.*
*Cell Sci.* 127, 4984 (2014).

61. Karanasios, E. *et al.* Autophagy initiation by ULK complex assembly on ER
tubulovesicular regions marked by ATG9 vesicles. *Nat. Commun.* 7, 12420 (2016).

62. Tapia, D. *et al.* KDEL receptor regulates secretion by lysosome relocation- and
autophagy-dependent modulation of lipid-droplet turnover. *Nat. Commun.* 10, 735
(2019).

63. Iwata, A., Riley, B. E., Johnston, J. A. & Kopito, R. R. HDAC6 and microtubules are
required for autophagic degradation of aggregated huntingtin. *J. Biol. Chem.* 280,
40282–40292 (2005).

64. Ding, W.-X. *et al.* Linking of autophagy to ubiquitin-proteasome system is important
for the regulation of endoplasmic reticulum stress and cell viability. *Am. J. Pathol.*
171, 513–524 (2007).

65. Lan, D., Wang, W., Zhuang, J. & Zhao, Z. Proteasome inhibitor-induced autophagy in
PC12 cells overexpressing A53T mutant α-synuclein. *Mol. Med. Rep.* 11, 1655–1660
(2015).

66. Peng, H. *et al.* Ubiquitylation of p62/sequestosome1 activates its autophagy receptor
function and controls selective autophagy upon ubiquitin stress. *Cell Res.* 27, 657–674
(2017).

67. Milani, M. *et al.* The role of ATF4 stabilization and autophagy in resistance of breast
cancer cells treated with Bortezomib. *Cancer Res.* 69, 4415–4423 (2009).

68. Albornoz, N., Bustamante, H., Soza, A. & Burgos, P. Cellular responses to proteasome
inhibition: molecular mechanisms and beyond. *Int. J. Mol. Sci.* 20, (2019).

69. Zhang, J., Liu, J., Norris, A., Grant, B. D. & Wang, X. A novel requirement for
ubiquitin-conjugating enzyme UBC-13 in retrograde recycling of MIG-14/Wntless and
Wnt signaling. *Mol. Biol. Cell* 29, 2098–2112 (2018).

70. Deng, L. *et al.* The ubiquitination of rag A GTPase by RNF152 negatively regulates
mTORC1 activation. *Mol. Cell* 58, 804–818 (2015).

71. Ji, C. H. *et al.* The N-Degron Pathway Mediates ER-phagy. *Mol. Cell* 75, 1058–
1072.e9 (2019).

72. Kawaguchi, Y. *et al.* The deacetylase HDAC6 regulates aggresome formation and cell
viability in response to misfolded protein stress. *Cell* 115, 727–738 (2003).

73. Cohen, M., Stutz, F. & Dargemont, C. Deubiquitination, a new player in Golgi to
endoplasmic reticulum retrograde transport. *J. Biol. Chem.* 278, 51989–51992 (2003).

74. Millard, S. M. & Wood, S. A. Riding the DUBway: regulation of protein trafficking by
deubiquitylating enzymes. *J. Cell Biol.* 173, 463–468 (2006).

75. Cohen, M., Stutz, F., Belgareh, N., Haguenauer-Tsapis, R. & Dargemont, C. Ubp3
requires a cofactor, Bre5, to specifically de-ubiquitinate the COPII protein, Sec23. *Nat.*
*Cell Biol.* 5, 661–667 (2003).

76. He, S., O'Connell, D., Zhang, X., Yang, Y. & Liang, C. The intersection of Golgi-ER
retrograde and autophagic trafficking. *Autophagy* 10, 180–181 (2014).

77. Rinaldi, L. *et al.* Feedback inhibition of cAMP effector signaling by a chaperone-
assisted ubiquitin system. *Nat. Commun.* 10, 2572 (2019).

78. Feng, X. *et al.* Ubiquitination of UVRAG by SMURF1 promotes autophagosome
maturation and inhibits hepatocellular carcinoma growth. *Autophagy* 15, 1130–1149
(2019).
79. Saitoh, T. *et al.* Atg9a controls dsDNA-driven dynamic translocation of STING and
the innate immune response. *Proc. Natl. Acad. Sci. USA* 106, 20842–20846 (2009).
80. Mattera, R., Park, S. Y., De Pace, R., Guardia, C. M. & Bonifacino, J. S. AP-4
mediates export of ATG9A from the trans-Golgi network to promote autophagosome
formation. *Proc. Natl. Acad. Sci. USA* 114, E10697–E10706 (2017).
81. Song, Y. *et al.* Blockade of deubiquitylating enzyme Rpn11 triggers apoptosis in
multiple myeloma cells and overcomes bortezomib resistance. *Oncogene* 36, 5631–
5638 (2017).
82. Wang, C.-H. *et al.* POH1 knockdown induces cancer cell apoptosis via p53 and bim.
*Neoplasia* 20, 411–424 (2018).
83. Deshaies, R. J. Proteotoxic crisis, the ubiquitin-proteasome system, and cancer therapy.
*BMC Biol.* 12, 94 (2014).
84. Bustamante, H. A. *et al.* Turnover of C99 is controlled by a crosstalk between ERAD
and ubiquitin-independent lysosomal degradation in human neuroglioma cells. *PLoS*
*One* 8, e83096 (2013).
85. Livak, K. J. & Schmittgen, T. D. Analysis of relative gene expression data using real-
time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* 25, 402–408
(2001).
86. Pfaffl, M. W. A new mathematical model for relative quantification in real-time RT-
PCR. *Nucleic Acids Res.* 29, e45 (2001).
87. Mlynarczuk-Bialy, I. *et al.* Biodistribution and efficacy studies of the proteasome
inhibitor bsc2118 in a mouse melanoma model. *Transl Oncol* 7, 570–579 (2014).
985
986

**Supplementary Figures Legends**

**Supplementary Figure 1. The PSMD14 DUB inhibitor CZM increases the Golgi**

**apparatus area**. Immunofluorescence microscopy analysis of the Golgi area in parental H4

cells treated for 4 h either with the vehicle (DMSO; Control) or CZM. The Golgi marker

GM130 was used to determine the region of interest in each condition. Statistical

significance was determined by Student's t-test. Bars represent the mean ± SEM (n =43

cells). ***P <0.001.

**Supplementary Figure 2. CZM causes the accumulation of KDELR1-GFP at the Golgi**

**apparatus.** HeLa cells expressing KDELR1-GFP were either left untreated or treated with

CZM for 30, 60 or 90 min. Cells were fixed and representative confocal images were

acquired.

**Supplementary Figure 3.  Effect of CZM on proteasome activity.** Parental H4 cells were

treated either with the vehicle (DMSO; Control), CZM or MG132, for 90 min. Protein

extracts were used to measure *in vitro* the Chymotrypsin-like peptidase activity of the

proteasome. The enzymatic activity was quantified according to the cleavage of the

fluorogenic substrate Suc-LLVY-AMC to AMC, and normalized to that of control cells.

The statistical significance was determined by One-Way ANOVA, followed by Tukey's

test. Bars represent the mean ± SD of biological replicates (n=3). **P <0.01; n.s., not

significant.

**Supplementary Figure 4. Effect of CZM and MG132 on basal macroautophagy. (A)**

Immunofluorescence microscopy analysis of the subcellular localization of LC3 in parental

H4 cells treated with either with the vehicle (DMSO; Control), CZM for 4 h or MG132 for

6 h. Cells were fixed, permeabilized and stained with a rabbit polyclonal antibody to LC3B

followed by Alexa-594-conjugated donkey anti-Rabbit IgG. Scale bar 10 μm. **(B)** Parental

H4 cells were treated as in (A) and protein extracts were analyzed by western blot with a

polyclonal antibody to LC3B. Monoclonal antibody to β-actin (clone BA3R) was used as a

loading control. The position of molecular mass markers is indicated on the left. **(C)** Ratio

LC3B-II/LC3B-I ratio as shown in (B). The statistical significance was determined by One-

Way ANOVA, followed by Tukey's test. Bars represent the mean ± SD of biological replicates (n=3). ***P <0.001; n.s., not significant.

**Supplementary Figure 5. Distribution of RAB1A upon CZM treatment.** Immunofluorescence analysis of endogenous RAB1A in H4 parental cells treated either with vehicle (DMSO; Control) **(A-C)** or CZM for 4 h **(D-F)**. Cells were fixed, permeabilized, and double stained with a rabbit monoclonal antibody to RAB1A (clone D3X9S) (A and D) and a mouse monoclonal antibody to GM130 (clone35/GM130) (B and E), followed by Alexa-594-conjugated donkey anti-Rabbit IgG and Alexa-488-conjugated donkey anti-Mouse IgG. Merging of the images generated the third picture (C and F). Scale bar, 10 μm. **(G)** Quantitative analysis of the fraction of RAB1A colocalizing with GM130 under CZM treatment and compared to control cells. The statistical significance was determined by Student's t-test. Bars represent the mean ± SEM of the fluorescent signal per cell area (n=173 cells). *P< 0.05.

**Supplementary Figure 6. ATG9A is distributed in the swollen Golgi apparatus upon CZM treatment.** Immunofluorescence analysis of endogenous ATG9A in H4 parental cells treated either with the vehicle (DMSO; Control) **(A-C)** or CZM for 4 h **(D-F)**. Cells were fixed, permeabilized, and double stained with a rabbit monoclonal antibody to ATG9A (clone EPR2450(2)) (A and D) and a mouse monoclonal antibody to GM130 (clone35/GM130) (B and E), followed by Alexa-594-conjugated donkey anti-Rabbit IgG and Alexa-488-conjugated donkey anti-Mouse IgG. Merging of the images generated the third picture (C and F). Scale bar, 10 μm. **(G)** Quantitative analysis of the fraction of ATG9A colocalizing with GM130 under CZM treatment and compared to control cells. The statistical significance was determined by Student's t-test. Bars represent the mean ± SEM of the fluorescent signal per cell area (n=93 cells). **P <0.01.

**Supplementary Figure 7. siRNA sequences directed against human PSMD14 used for Validation Stage.**

**Supplementary Figure 8. Primer pairs sequences used for RT-qPCR.**