The crystal structure of the TonB-dependent transporter YncD reveals

a positively charged substrate binding site.

Rhys Grinter[a,b]# and Trevor Lithgow[b]

#To whom correspondence should be addressed: rhys.grinter@monash.edu

[a]School of Biological Sciences, Monash University, Clayton 3800, Victoria, Australia

[b]Infection and Immunity Program, Biomedicine Discovery Institute and Department of Microbiology,

Monash University, Clayton 3800, Victoria, Australia

Keywords: YncD, Crystallography, Membrane Transport, Gram-Negative Bacteria, TonB-Dependent

Transporter

Abstract

The outer membrane of Gram-negative bacteria is highly impermeable to hydrophilic molecules larger than 600 Da, protecting these bacteria from toxins present in the environment. In order to transport nutrients across this impermeable membrane, Gram-negative bacteria utilise a diverse family of outer-membrane proteins called TonB-dependent transporters. The majority of this family transport iron-containing substrates. However, it is becoming increasingly clear that TonB-dependent transporters target chemically diverse substrates. In this work, we investigate the structure and phylogenetic distribution of the TonB-dependent transporter YncD. We show that while YncD is present in some enteropathogens including *E. coli* and *Salmonella* spp., it is also widespread in Gamma and Betaproteobacteria of environmental origin. We determine the structure of YncD, showing that despite a distant evolutionary relationship, it shares structural features with the ferric-citrate transporter FecA, including a compact positively-charged substrate-binding site. Despite these shared features, we show that YncD does not contribute to the growth of *E. coli* in pure culture under-iron limiting conditions or with ferric-citrate as an iron source. Previous studies on transcriptional regulation in *E. coli* show that YncD is not induced under iron-limiting conditions and is unresponsive to the Ferric uptake regulator (Fur). These observations combined with the data we present, suggest that YncD is not responsible for the transport of an iron-containing substrate.

Introduction

The outer membrane of Gram-negative bacteria represents a formidable permeability barrier to hydrophilic molecules larger than ~600 Da (1). Hydrophilic molecules smaller than this diffusion limit cross the outer membrane through proteins called porins, which can be either promiscuous or substrate-specific (2). To import molecules that are too large for diffusion through outer-membrane porins, bacteria employ a diverse family of membrane proteins termed TonB-dependent transporters (TBDTs) (3). These transporters in the outer membrane are so named because they engage in active transport, the energy for which is provided by the TonB-ExbBD complex in the inner membrane, which spans the periplasm to interact with TBDTs (4,5). TBDTs are specific for a particular substrate, which they bind with high affinity in an extracellular binding pocket (6,7). This high-affinity binding allows TBDTs to capture molecules that are of low abundance in the environment, enabling bacteria to scavenge scarce nutrients (6). One such scarce nutrient is iron, which is the substrate for the majority of characterised TBDTs (3). Most iron-transporting TBDTs bind Fe^{3+} ions in complex with siderophores, organic molecules that bind metal ions with high affinity and are secreted by the bacteria or by competitors from the microbial community (8). In addition to importing Fe-siderophore complexes, some TBDTs harvest iron directly from proteins (9,10). These protein-binding TBDTs often target host proteins and are employed by pathogens to obtain iron during infections (11,12).

While many of the most intensively studied TBDTs import iron-containing substrates, it is becoming increasingly clear that members of the TBDT protein family are responsible for the import of chemically diverse substrates (13). Non-iron TBDT substrates range in size from free metal ions, like Cu^{2+} and Zn^{2+}, to cobalamin, polysaccharides, polypeptides, and small globular proteins (9,14-22). As functionally characterised TBDTs represent a small segment of this protein family, the extent of the substrate diversity of this family remains to be discovered.

The model bacterium *Escherichia coli* K12 possesses 9 TBDTs which target different substrates for import (7). The TBDTs FepA, CirA and Fiu transport Fe^{3+} containing catecholate siderophores (7,23,24), FhuA and FhuE transport Fe^{3+} containing hydroxamate siderophores

of fungal origin (6,25), FecA transports iron citrate (26), while BtuB targets cobalamin (16).

Recent research suggests that the TBDT YddB from *E. coli* K12 functions in the import of a

small iron-containing protein, although the exact nature of this substrate remains to be

identified (27). The final TBDT possessed by *E. coli* K12 is YncD, and the substrate for this

transporter remains unknown. The homolog of YncD in *Salmonella enterica* ssp. *enterica*

serovar Typhi has been shown to be important for virulence in a mouse model of infection

(28). As such, characterisation of YncD is important for understanding infections caused by

Salmonella and other enterobacterial pathogens that possess it.

In this study, we investigated the structure and phylogenetic distribution of YncD. Using a

comparative genomics approach, we show that YncD is present in members of

Enterobacteriaceae that form commensal or pathogenic associations with the human

intestine, and is more broadly present in environmental Proteobacterial isolates, including

species which are opportunistic pathogens of humans. We determined the crystal structure

of YncD demonstrating that it possesses a compact positively-charged extracellular substrate-

binding site that is characteristic of TBDTs that import negatively charged small-molecule

substrates. Finally, using an *E. coli* strain deficient in all iron transporting TBDTs we show that

YncD does not play a role in iron acquisition under laboratory conditions, providing further

evidence that this transporter does not transport an iron-containing substrate.

Results

The TonB-dependent transporter YncD is widespread in environmental bacteria

To determine the evolutionary relationship between YncD and other functionally characterised TBDTs, we performed clustering analysis of these sequences based on pairwise sequence similarity scores, using the program CLANS (29). This analysis reveals that YncD is distantly related to other TBDTS of known function and does not form a cluster with any of these sequences (Figure 1A). This analysis shows that while YncD is most closely related to the TBDTs FecA and BtuB, this reflects only 21% and 18% sequence identity, respectively.

To gain insight into the distribution of YncD in Gram-negative bacteria, we interrogated the Uniprot reference proteomes for YncD homologues with the HMMER search algorithm (30,31). An initial sequence similarly E-value cut-off of $1x10^{-50}$ was used to capture all YncD sequences present in the database. To remove false-positive non-YncD sequences from this dataset, we subsequently clustered the sequences using CLANS (29) with an E-value cut-off of $1x10^{-130}$. The largest sequence cluster had 359 distinct sequences including YncD from *E. coli* and was extracted for curation and analysis (Table S1, Figure S1A). Partial sequences were removed from this dataset, yielding 331 sequences. These were reclustered with an E-value cut-off of 0, revealing several distinct sub-sets within the YncD protein sub-family (Figure S1B). A multiple sequence alignment of these diverse YncD sequences was then performed revealing that they share amino acid sequence identities ranging between 33 and 99 % (Table S2, Data S1), and highlighted a number of conserved regions. Most notably the region of the corresponding in amino acids 79-150 of YncD from *E. coli* is highly conserved (Data S1). This region encompasses the extracellular loops of the N-terminal plug domain of the transporter, known to mediate substrate binding, suggesting that the YncD homologues identified target a chemically analogous substrate.

To explore potential niche-specific functions associated with YncD, we interrogated genome metadata from those bacteria identified in the YncD HMMER search. The environment of isolation of 105 YncD-containing species was determined, with all bacteria belonging to either the gamma- or beta-proteobacteria (Table S1). While some of these bacterial species were isolated from a human host (largely Enterobacteriaceae associated with the gut or

Pseudomonadales associated with respiratory infection), the majority of YncD-containing

bacteria were environmental isolates (Figure 1B). The environmental isolates originated from

soil, sediment or water samples (Figure 1C). The widespread distribution of YncD in

environmental isolates, and its absence from most specialist bacterial pathogens, would be

most consistent with YncD playing a role in the acquisition of a molecule produced by the

microbial community.

The crystal structure of YncD from *E. coli* reveals a positively charged substrate-binding site

To gain insight into the evolutionary relationship between YncD and other TBDTs and to

provide insight into the nature of the YncD substrate, we solved the crystal structure of YncD

by X-ray crystallography. Crystals of YncD from *E. coli* were prepared and diffraction was

collected to 3.2 Å (Table 1). Molecular replacement using the closest YncD homologue FecA

(PDB ID: 1KMO, 1KMP) proved unsuccessful (26). To obtain experimental phases, YncD

crystals were soaked with potassium tetranitroplatinate and anomalous diffraction was

collected to 3.5 Å (Table 1). Phases were obtained for this dataset using single-wavelength

anomalous dispersion (SAD) phasing, with an anomalous substructure of 2 high occupancy

and 1 low occupancy platinum atoms located (Figure S3). An initial model of YncD was build

using maps from this dataset and utilised to obtain phases for the higher resolution dataset

by molecular replacement. The structure of YncD was then built and refined using the native

3.2 Å data (Table 1).

As expected, based on other members of the TBDT family, structurally YncD consists of at 22

stranded transmembrane β-barrel, which is occluded by a globular N-terminal plug domain

(Figure 2A, B). The Dali server was utilised to search the protein databank (PDB) for structural

homologues to YncD (32), revealing that consistent with our sequence analysis the closest

structural homologue to YncD is FecA (Table S3). Despite the low sequence identity of 21 %

between YncD and FecA, these proteins share a backbone atom root mean square deviation

(RMSD) of 2.2 Å. Consistent with this, the extracellular loop length, secondary and tertiary

structure is relatively well conserved between the two proteins, compared to the more

distantly related TBDT FhuE (Figure 2C, Figure S2).

Previous analysis of the structure of diverse TBDTs in complex with their substrates revealed

the location of a conserved substrate-binding site in these transporters (7). Based on this

analysis we identified a compact substrate-binding site formed by the extracellular loops of

YncD in our structure (Figure 3A). Analysis of sequence conservation among YncD

homologues using the program Consurf (33) and our alignment of YncD sequences revealed

that the residues that constitute this binding pocket are well conserved in the YncD

homologues, suggesting they play an important role in transporter function (Figure 1B, C,

Data S1). These conserved residues include a number of arginine, glutamine and lysine

residues that give the substrate-binding site of YncD a highly positive charged, based on a

calculated electrostatic surface (Figure 3B-D). Electrostatic analysis of the structure of FecA

shows that this distant YncD homologue also possesses a highly positively charged substrate-

binding site (Figure 3C). In FecA, several arginine and glutamine residues, which give the

substrate-binding site its positive character, are responsible for coordinating the carboxylic

acid groups of the Fe-citrate complex, and thus are important for substrate binding by FecA

(Figure 3C) (26). Superimposition of the FecA-Fe-Citrate complex with YncD places Fe-citrate

ligand centrally in the positively charged substrate binding site of YncD (Figure 3C). The

similarities between the character of the substrate bindings sites of YncD and FecA suggest

that despite their differences they may target chemically similar substrates.

YncD does not import ferric citrate or support growth of *E. coli* during iron limitation in pure

culture

To test the hypothesis that YncD could support growth under iron-limiting conditions and

specifically that it imports ferric citrate, we monitored the growth of an isogenic *E. coli* strain

with and without YncD. Because TBDT iron-uptake systems are highly redundant, the *yncD*

gene was deleted in a strain which is deficient in the 7 TBDTs involved in iron-uptake (Figure

S4) (7). This ΔTBDT strain grows poorly under iron-limiting conditions (6), making any

additional iron-acquisition defect easier to discern. In M9 minimal media, the growth of the

ΔTBDT +/- *yncD* was identical, demonstrating that YncD does not play a role in iron

acquisition under these conditions (Figure 4A). Since there is some residual free iron in M9

minimal media, the available iron was chelated with the addition of citrate or the monomeric

catecholate compound 2,3-dihydroxybenzoic acid (DHB). As the ΔTBDT strain lacks the outer

membrane transporters required to import these substrates, they limit the exogenous iron

available to the bacteria. Accordingly, the growth of *E. coli* ΔTBDT was somewhat reduced in

the presence of DHB and drastically reduced in the presence of citrate (Figure 4B). Deletion

of *yncD* in this strain had no further effect on growth, demonstrating that YncD does not play

a role in importing Fe-citrate or Fe-DHB under these conditions (Figure 4B). As expected

growth of wildtype *E. coli* BW25113, which possesses TBDTs for both ferric-citrate (FecA) and

Fe-DHB (Fiu, CirA, FepA) import (7,26) was identical in the presence or absence of these

compounds (Figure 4C).

Discussion

Determination of the function of proteins is one of the most challenging aspects of modern

protein biochemistry. The expansion of genomic and metagenomic sequencing data has

provided a wealth of proteins sequences, most of which are only partially characterised or of

completely unknown function. Despite its difficulty, determining the function of novel

proteins is invaluable, as without a robust understanding of the functional capacity of an

organism's proteome it is difficult to accurately predict its metabolic capabilities, physiology

or lifestyle.

In this work, we show that YncD is a TBDT which is only distantly related to previously

characterised members of this family and that YncD is widespread in gamma- and beta-

proteobacteria. The crystal structure of YncD shows common structural features with the

ferric-citrate transporter FecA, but from growth assays, YncD appears not to transport ferric-

citrate. The structure of YncD represents the final structure of the 9 TBDTs possessed by the

model bacterium *E. coli* K12 (6,7,23,26,34-36). This structure improves our understanding of

the structural variation and similarities that exist in the TBDT family and paves the way for

future studies addressing the function of YncD.

Seven of the nine TBDTs present in *E. coli* are involved in iron acquisition. All of these

transporters are regulated by the Ferric uptake regulator (Fur) and are induced under iron-

limiting conditions (37). BtuB, which imports the non-iron containing substrate cobalamin, is

neither Fur regulated or induced under iron limiting conditions (37,38). Interestingly, like

BtuB, YncD is neither Fur regulated or upregulated under iron-limiting conditions, suggesting
that it may not function in the import of an iron-containing substrate (37,38). While the
specific substrate transported by YncD remains to be determined, this work provides a
number of clues and avenues for further research towards its identification. Based on its
compact negatively-charged extracellular binding site, the substrate of YncD is likely to be
relatively small and negatively charged.

As YncD is prevalent in bacterial species from soil and aquatic communities, its substrate is
likely produced by members of these ecological communities. In addition, the impact of YncD
on virulence phenotypes in *Salmonella* Typhi suggest that it is also a molecule which is
relevant in a human host-context (28). Based on these considerations and structural
parameters in the ligand-binding site, possible substrates for YncD would include molecules
containing negatively charged phosphate groups. Other members of the TBDT family have
been previously been implicated in thiamine pyrophosphate import through comparative
genomics (39). Alternatively, YncD may import an organic acid containing compound in
complex with a non-ferrous metal ion (26).

The importance of YncD for virulence in *Salmonella* Typhi (28), makes understanding the
function of this transporter an important question for medical microbiology. Understanding
the substrate of YncD and the role of this compound in bacterial virulence will improve our
fundamental understanding of infection by this bacterium and provide a potential target for
therapeutic intervention.

Methods

YncD Homologue Identification and Analysis

To determine the evolutionary relationship between YncD and other TBDTs of known structure
and/or function these sequences were classified by an all-against-all BLAST clustering algorithm,
based on pairwise similarities. The resulting data set was visualized with CLANS with an E-value cut
off of $1{\times}10^{-10}$ (29).

To identify YncD sequences in available bacterial genomes, a HMMER search was performed on the Uniprot reference proteomes database using YncD as the search sequence (30,31). An E-value cut-off of 1×10^{-50} applied to hits. This search yielded a total of 568 YncD homologue sequences. These sequences were classified by an all-against-all BLAST clustering algorithm, based on pairwise similarities. The resulting data set was visualized with CLANS with an E value cut off of 1×10^{-130}. Sequence clusters were identified in CLANS using a network-based algorithm, with a minimum group size of 10 (29). The largest sequence cluster from this clustering contained 359 sequences, including YncD from *E. coli* and designated the YncD group. This group was isolated and subjected to further clustering with an E value cut off of 0. A multiple sequence alignment was performed on 331 full-length sequences from the YncD group using the clustal algorithm (40). Environment or host of isolation for a subset of YncD sequences (105 sequences) was determined from genome metadata from the Ensemble and Uniprot databases (41,42).

YncD Expression and Purification

The open reading frame encoding YncD was amplified by PCR from *E. coli* BW25113 using primers containing 5' BamHI and 3' XhoI restriction sites (Forward: CCATCGGATCCGGCTGA TGAACAGACTATGATTGTC, Reverse: CCATCCTCGAGTTACTCAAATCTCCACGCAATATTCAT). This YncD open reading frame was then cloned into a modified pET20b vector with a PelB signal sequence, followed by an N-terminal 10 × His-tag and TEV cleavage site, via restriction digestion and ligation. The resulting vector was transformed into *E. coli* BL21 (DE3) C41 cells (43). Protein expression was performed in terrific broth (12 g tryptone, 24 g yeast extract, 61.3 g K_2HPO_4, 11.55 g KH_2PO_4, 10 g glycerol) with 100 mg.ml^{-1} ampicillin for selection. Cells were grown at 37 °C until OD_{600} of 1.0, induced with 0.3 mM IPTG, and grown for a further 14 hours at 25 °C. Cells were harvested by centrifugation and lysed using a cell disruptor (Emulseflex) in Lysis Buffer (50 mM Tris, 200 mM NaCl [pH 7.9]) in the presence of 0.1 mg.ml^{-1} Lysozyme, 0.05 mg.ml^{-1} DNAse1 and complete protease cocktail inhibitor tablets (Roche).

The resulting lysate was clarified by centrifugation at 10,000 g for 10 min, the supernatant from this low-speed spin was then centrifuged to 1 h at 100,000 g to isolate a membrane fraction. The supernatant was decanted, and the membrane pellet was suspended in lysis buffer using a tight-fitting homogeniser. The resuspended membranes were solubilised by the addition of 10 % Elugent (Santa Cruz Biotechnology) and incubated with gentle stirring at room temperature for 20 min. The solubilised membrane protein fraction was clarified by centrifugation at 20,000 g for 10 min. The supernatant containing the solubilized proteins was applied to Ni-agarose resin equilibrated in Ni-

binding buffer DDM (50 mM Tris, 500 mM NaCl, 20 mM Imidazole, 0.03% Dodecylmaltoside (DDM)

[pH7.9]). The resin was washed with 10-20 column volumes of Ni binding buffer DDM before elution

of the protein with a step gradient of, 10, 25 and 50, 100 % Ni gradient buffer DDM (50 mM Tris, 500

mM NaCl, 1 M Imidazole, 0.03 % DDM [pH7.9]). YncD eluted at the 50 and 100 % gradient steps.

Eluted fractions containing YncD were pooled and applied to a 26/600 S200 Superdex size exclusion

column equilibrated in SEC buffer DDM (50 mM Tris, 200 mM NaCl, 0.03 % DDM [pH 7.9]). To

exchange YncD into the detergent Octyl β-D-glucopyranoside (βOG) for crystallographic and

biochemical analysis, SEC fractions containing YncD were pooled and applied to Ni-agarose resin,

equilibrated in βOG buffer (50 mM Tris, 200 mM NaCl, 0.8 % βOG [pH 7.9]). The resin was washed

with 10 column volumes of βOG buffer before elution with βOG buffer + 250 mM imidazole. Fractions

containing YncD were pooled, and 6 × His-tagged TEV protease (final concentration 2 mg.ml^{-1}) and DTT

(final concentration 1 mM) were added. This solution was then dialysed against of βOG buffer at 4-6h

at 20 °C to allow TEV cleavage of the 10 × his-tag from YncD and removal of excess imidazole. The

sample was then applied to Ni-agarose resin to remove TEV protease and the cleaved histidine-

containing peptide. The flow-through containing YncD from this step was collected, concentrated to

10 mg.ml^{-1} in a 30 kDa cut-off centrifugal concentrator, snap-frozen and stored at -80 °C.

YncD Crystallisation, Data Collection and Structure Solution

Purified YncD in βOG was screened for crystallisation conditions using commercially available screens

(approximately 600 conditions). Crystals grew in a number of conditions, with a condition containing

0.2 M Ca acetate, 0.10 M Tris pH 8.5 and 20 % PEG 3000 chosen for optimisation. Concentration and

pH grid screening was performed yielding an optimised crystallisation conditions containing 0.17 M Ca

acetate, 0.10 M Tris pH 7.0 and 16 % PEG 3000. Crystals from this condition were looped,

crystallisation solution was removed by wicking and crystals were flash-cooled in liquid N$_2$. Diffraction

data were collected at 100 K at the Australian synchrotron and processed in the space group P2$_1$22$_1$

to 3.2 Å. Molecular replacement of this dataset using the structures of various TBDTs as a starting

model failed and so soaking was performed to obtain a heavy atom derivative. Using a fine

acupuncture needle a small quantity of potassium tetranitroplatinate salt was added to the drop

containing YncD crystals, the well was resealed and crystal incubated for 60 minutes. These crystals

were then were looped, crystallisation solution was removed by wicking and crystals were flash-

frozen in liquid N$_2$. Data was collected from the potassium tetranitroplatinate soaked crystals at 100 K

at the Australian synchrotron using a wavelength of 0.987 Å, with crystals diffracting to 3.5 Å. A heavy

atom search was performed by SAD using Shelx within the CCP4 software package (44,45). Three

heavy atom sites were identified, which were provided to Autosol withing the Phenix software

package for phasing and density modification (46). An initial structure of YncD was constructed using

these experimentally-phased density modified maps using Coot (47). This model was used for

molecular replacement of the 3.2 Å native data using phaser (48). The model of YncD was then build

and refined using Coot, Phenix package and Buster (46,47,49).

E. coli ΔyncD Mutant Generation and Growth Analysis

The *E. coli* BW25113 *ΔyncD* mutant strain was created using the λ Red system (50). A Kanamycin-

resistance cassette flanked by 300 bp of genomic DNA either side of the chromosomal location of

yncD was amplified by PCR (primers: F: AAACAGGCTATTTCGCTTAGCGA, R:

GAACCTAACAGTAATGAACCACG using the specific mutant from the *E. coli* Keio collection (51) as a

template, generating the yncD-Kan KO cassette. *E. coli* ΔTBDT (6) was transformed with the λ Red

recombinase plasmid pKD46 (50) and grown at 30 °C (LB broth + 100 µg.ml^{-1} ampicillin + 100 uM

Fe(II)SO$_4$) to an OD600 nm of 0.1 before λ Red recombinase was induced by the addition of 0.2% L-

arabinose. The culture was then grown at 30 °C until an OD600 nm of 0.6–0.8 was attained and were

transformed with the yncD-Kan KO cassette using the room-temperature electroporation method

(52). Briefly, bacterial cells were isolated by centrifugation at 3000 g for 3 min and washed twice with

a volume of sterile 10% glycerol equal to the volume of culture used. The cells were then

resuspended in 10% glycerol to a volume of 1/15 of that of the culture. The yncD-Kan KO cassette

DNA (100–500 ng) was then added to 100 µl of the resuspended bacteria and the mixture was

electroporated. 1 ml of LB broth was added to the cells post-electroporation, and the culture was

recovered at 37°C for 1 h before plating onto LB agar + 30 µg ml^{-1} kanamycin + 100 uM Fe(II)SO$_4$. PCR

was used to validate that colonies did indeed have the KanR cassette in place of the gene of interest.

To remove the KanR gene and generate a "clean" *yncD* deletion, the mutant strain was transformed

with the plasmid pCP20 (53) containing the `flippase cassette'. Cells were grown at 30 °C under either

ampicillin (100 µg ml^{-1}) or chloramphenicol (30 µg ml^{-1}) selection to maintain the plasmid. A single

colony of the mutant pCP20-containing strain was used to inoculate 1 ml LB broth + 100 uM Fe(II)SO$_4$

(no selection). The culture was grown overnight at 43°C to activate expression of the flippase gene.

This culture was then subjected to tenfold serial dilution in sterile LB and plated onto LB agar + 100

uM Fe(II)SO$_4$ with no selection. The resulting colonies were patched onto LB agar containing

kanamycin, chloramphenicol or no selection. PCR was used to validate colonies that, while unable to

grow in the presence of kanamycin or chloramphenicol, grew in the absence of selection and had no

remnant of the KanR cassette in the deletion of *yncD*.

E. coli ΔTBDT, *E. coli* ΔTBDT/*ΔyncD* and wildtype *E. coli* BW25113 were grown in LB broth until

stationary phase. These cultures were used to inoculate 20 ml of M9 minimal media +/- 10 mM citrate

or +/- 100 µM 2,3-Dihydrobenzoic acid, to an OD600nm of 0.05. Cultures were grown at 37 °C with

shaking and the rate of growth was quantified by measuring OD600nm at hourly intervals.

References

1. van den Berg, B., Prathyusha Bhamidimarri, S., Dahyabhai Prajapati, J., Kleinekathöfer, U., and Winterhalter, M. (2015) Outer-membrane translocation of bulky small molecules by passive diffusion. *Proceedings of the National Academy of Sciences* **112**, E2991-E2999

2. Vergalli, J., Bodrenko, I. V., Masi, M., Moynié, L., Acosta-Gutiérrez, S., Naismith, J. H., Davin-Regli, A., Ceccarelli, M., van den Berg, B., and Winterhalter, M. (2019) Porins and small-molecule translocation across the outer membrane of Gram-negative bacteria. *Nat. Rev. Microbiol.*, 1-13

3. Noinaj, N., Guillier, M., Barnard, Travis J., and Buchanan, S. K. (2010) TonB-Dependent Transporters: Regulation, Structure, and Function. *Annual Review of Microbiology* **64**, 43-60

4. Maki-Yonekura, S., Matsuoka, R., Yamashita, Y., Shimizu, H., Tanaka, M., Iwabuki, F., and Yonekura, K. (2018) Hexameric and pentameric complexes of the ExbBD energizer in the Ton system. *Elife* **7**

5. Celia, H., Noinaj, N., Zakharov, S. D., Bordignon, E., Botos, I., Santamaria, M., Barnard, T. J., Cramer, W. A., Lloubes, R., and Buchanan, S. K. (2016) Structural insight into the role of the Ton complex in energy transduction. *Nature*

6. Grinter, R., and Lithgow, T. (2019) Determination of the molecular basis for coprogen import by Gram-negative bacteria. *IUCrJ* **6**

7. Grinter, R., and Lithgow, T. (2019) The structure of the bacterial iron–catecholate transporter Fiu suggests that it imports substrates via a two-step mechanism. *Journal of Biological Chemistry* **294**, 19523-19534

8. Chu, B., Garcia-Herrero, A., Johanson, T., Krewulak, K., Lau, C., Peacock, R., Slavinskaya, Z., and Vogel, H. (2010) Siderophore uptake in bacteria and the battle for iron with the host; a bird's eye view. *Biometals* **23**, 601-611

9. Grinter, R., Josts, I., Mosbahi, K., Roszak, A. W., Cogdell, R. J., Bonvin, A. M., Milner, J. J., Kelly, S. M., Byron, O., and Smith, B. O. (2016) Structure of the bacterial plant-ferredoxin receptor FusA. *Nature communications* **7**

10. Noinaj, N., Easley, N. C., Oke, M., Mizuno, N., Gumbart, J., Boura, E., Steere, A. N., Zak, O., Aisen, P., Tajkhorshid, E., Evans, R. W., Gorringe, A. R., Mason, A. B., Steven, A. C., and Buchanan, S. K. (2012) Structural basis for iron piracy by pathogenic Neisseria. *Nature* **483**, 53-58

11. Leduc, I., Banks, K. E., Fortney, K. R., Patterson, K. B., Billings, S. D., Katz, B. P., Spinola, S. M., and Elkins, C. (2008) Evaluation of the Repertoire of the TonB-Dependent Receptors of Haemophilus ducreyi for Their Role in Virulence in Humans. *Journal of Infectious Diseases* **197**, 1103-1109

12. Cornelissen, C. N., Kelley, M., Hobbs, M. M., Anderson, J. E., Cannon, J. G., Cohen, M. S., and Sparling, P. F. (1998) The transferrin receptor expressed by gonococcal strain FA1090 is required for the experimental infection of human male volunteers. *Molecular Microbiology* **27**, 611-616

13. Schauer, K., Rodionov, D. A., and de Reuse, H. (2008) New substrates for TonB-dependent transport: do we only see the 'tip of the iceberg'? *Trends in biochemical sciences* **33**, 330-338

14. Grinter, R., Milner, J., and Walker, D. (2012) Ferredoxin containing bacteriocins suggest a novel mechanism of iron uptake in *Pectobacterium* spp. *PLoS ONE* **7**, e33033

15. Grinter, R., Hay, I. D., Song, J., Wang, J., Teng, D., Dhanesakaran, V., Wilksch, J. J., Davies, M. R., Littler, D., and Beckham, S. A. (2018) FusC, a member of the M16 protease family acquired by bacteria for iron piracy against plants. *PLoS Biol.* **16**, e2006026

16. Chimento, D. P., Mohanty, A. K., Kadner, R. J., and Wiener, M. C. (2003) Substrate-induced transmembrane signaling in the cobalamin transporter BtuB. *Nature Structural & Molecular Biology* **10**, 394

17. Yoneyama, H., and Nakae, T. (1996) Protein C (OprC) of the outer membrane of Pseudomonas aeruginosa is a copper-regulated channel protein. *Microbiology* **142**, 2137-2144

18. Calmettes, C., Ing, C., Buckwalter, C. M., El Bakkouri, M., Lai, C. C.-L., Pogoutse, A., Gray-
Owen, S. D., Pomès, R., and Moraes, T. F. (2015) The molecular mechanism of Zinc acquisition
by the neisserial outer-membrane transporter ZnuD. *Nature communications* **6**, 7996
19. Bolam, D. N., and van den Berg, B. (2018) TonB-dependent transport by the gut microbiota:
novel aspects of an old problem. *Current opinion in structural biology* **51**, 35-43
20. Glenwright, A. J., Pothula, K. R., Bhamidimarri, S. P., Chorev, D. S., Baslé, A., Firbank, S. J.,
Zheng, H., Robinson, C. V., Winterhalter, M., Kleinekathöfer, U., Bolam, D. N., and van den
Berg, B. (2017) Structural basis for nutrient acquisition by dominant members of the human
gut microbiota. *Nature* **541**, 407-411
21. Madej, M., White, J. B., Nowakowska, Z., Rawson, S., Scavenius, C., Enghild, J. J., Bereta, G. P.,
Pothula, K., Kleinekathoefer, U., and Baslé, A. (2019) Dynamic oligopeptide acquisition by the
RagAB transporter from Porphyromonas gingivalis. *bioRxiv*, 755678
22. Grinter, R., Josts, I., Zeth, K., Roszak, A. W., McCaughey, L. C., Cogdell, R. J., Milner, J. J., Kelly,
S. M., Byron, O., and Walker, D. (2014) Structure of the atypical bacteriocin pectocin M2
implies a novel mechanism of protein uptake. *Molecular microbiology* **93**, 234-246
23. Buchanan, S. K., Smith, B. S., Venkatramani, L., Xia, D., Esser, L., Palnitkar, M., Chakraborty, R.,
Van Der Helm, D., and Deisenhofer, J. (1999) Crystal structure of the outer membrane active
transporter FepA from Escherichia coli. *Nature Structural & Molecular Biology* **6**, 56
24. Nikaido, H., and Rosenberg, E. Y. (1990) Cir and Fiu proteins in the outer membrane of
Escherichia coli catalyze transport of monomeric catechols: study with beta-lactam antibiotics
containing catechol and analogous groups. *Journal of bacteriology* **172**, 1361-1367
25. Pawelek, P. D., Croteau, N., Ng-Thow-Hing, C., Khursigara, C. M., Moiseeva, N., Allaire, M., and
Coulton, J. W. (2006) Structure of TonB in complex with FhuA, *E. coli* outer membrane
receptor. *Science* **312**, 1399-1402
26. Ferguson, A. D., Chakraborty, R., Smith, B. S., Esser, L., Van Der Helm, D., and Deisenhofer, J.
(2002) Structural basis of gating by the outer membrane transporter FecA. *Science* **295**, 1715-
1719
27. Grinter, R., Leung, P. M., Wijeyewickrema, L. C., Littler, D., Beckham, S., Pike, R. N., Walker, D.,
Greening, C., and Lithgow, T. (2019) Protease-associated import systems are widespread in
Gram-negative bacteria. *PLoS Genet.* **15**
28. Xiong, K., Chen, Z., Xiang, G., Wang, J., Rao, X., Hu, F., and Cong, Y. (2012) Deletion of yncD
gene in Salmonella enterica ssp. enterica serovar Typhi leads to attenuation in mouse model.
FEMS microbiology letters **328**, 70-77
29. Frickey, T., and Lupas, A. (2004) CLANS: a Java application for visualizing protein families
based on pairwise similarity. *Bioinformatics* **20**, 3702-3704
30. Chen, C., Natale, D. A., Finn, R. D., Huang, H., Zhang, J., Wu, C. H., and Mazumder, R. (2011)
Representative Proteomes: A Stable, Scalable and Unbiased Proteome Set for Sequence
Analysis and Functional Annotation. *PLoS ONE* **6**, e18910
31. Finn, R. D., Clements, J., and Eddy, S. R. (2011) HMMER web server: interactive sequence
similarity searching. *Nucleic Acids Res.* **39**, W29-W37
32. Holm, L., and Laakso, L. M. (2016) Dali server update. *Nucleic acids research* **44**, W351-W355
33. Ashkenazy, H., Abadi, S., Martz, E., Chay, O., Mayrose, I., Pupko, T., and Ben-Tal, N. (2016)
ConSurf 2016: an improved methodology to estimate and visualize evolutionary conservation
in macromolecules. *Nucleic acids research* **44**, W344-W350
34. Buchanan, S. K., Lukacik, P., Grizot, S., Ghirlando, R., Ali, M. M., Barnard, T. J., Jakes, K. S.,
Kienker, P. K., and Esser, L. (2007) Structure of colicin I receptor bound to the R-domain of
colicin Ia: implications for protein import. *The EMBO journal* **26**, 2594-2604
35. Locher, K. P., Rees, B., Koebnik, R., Mitschler, A., Moulinier, L., Rosenbusch, J. P., and Moras,
D. (1998) Transmembrane signaling across the ligand-gated FhuA receptor: crystal structures
of free and ferrichrome-bound states reveal allosteric changes. *Cell* **95**, 771-778

36. Grinter, R., Leung, P. M., Wijeyewickrema, L. C., Littler, D., Beckham, S., Pike, R. N., Walker, D.,
Greening, C., and Lithgow, T. (2019) Protease-associated import systems are widespread in
Gram-negative bacteria. *PLoS Genetics* **15**, e1008435

37. Seo, S. W., Kim, D., Latif, H., O'Brien, E. J., Szubin, R., and Palsson, B. O. (2014) Deciphering Fur
transcriptional regulatory network highlights its complex role beyond iron metabolism in
Escherichia coli. *Nat. Commun.* **5**, 4910

38. McHugh, J. P., Rodríguez-Quiñones, F., Abdul-Tehrani, H., Svistunenko, D. A., Poole, R. K.,
Cooper, C. E., and Andrews, S. C. (2003) Global iron-dependent gene regulation in Escherichia
coli A new mechanism for iron homeostasis. *J. Biol. Chem.* **278**, 29478-29486

39. Rodionov, D. A., Vitreschak, A. G., Mironov, A. A., and Gelfand, M. S. (2002) Comparative
genomics of thiamin biosynthesis in procaryotes New genes and regulatory mechanisms.
Journal of Biological chemistry **277**, 48949-48959

40. Larkin, M. A., Blackshields, G., Brown, N., Chenna, R., McGettigan, P. A., McWilliam, H.,
Valentin, F., Wallace, I. M., Wilm, A., and Lopez, R. (2007) Clustal W and Clustal X version 2.0.
bioinformatics **23**, 2947-2948

41. Hubbard, T., Barker, D., Birney, E., Cameron, G., Chen, Y., Clark, L., Cox, T., Cuff, J., Curwen, V.,
and Down, T. (2002) The Ensembl genome database project. *Nucleic Acids Res.* **30**, 38-41

42. Apweiler, R., Bairoch, A., Wu, C. H., Barker, W. C., Boeckmann, B., Ferro, S., Gasteiger, E.,
Huang, H., Lopez, R., and Magrane, M. (2004) UniProt: the universal protein knowledgebase.
Nucleic Acids Res. **32**, D115-D119

43. Miroux, B., and Walker, J. E. (1996) Over-production of proteins in Escherichia coli: mutant
hosts that allow synthesis of some membrane proteins and globular proteins at high levels.
Journal of molecular biology **260**, 289-298

44. Sheldrick, G. M. (2008) A short history of SHELX. *Acta Crystallographica Section A:*
Foundations of Crystallography **64**, 112-122

45. Winn, M. D., Ballard, C. C., Cowtan, K. D., Dodson, E. J., Emsley, P., Evans, P. R., Keegan, R. M.,
Krissinel, E. B., Leslie, A. G., and McCoy, A. (2011) Overview of the CCP4 suite and current
developments. *Acta Crystallographica Section D: Biological Crystallography* **67**, 235-242

46. Adams, P. D., Afonine, P. V., Bunkoczi, G., Chen, V. B., Davis, I. W., Echols, N., Headd, J. J.,
Hung, L.-W., Kapral, G. J., Grosse-Kunstleve, R. W., McCoy, A. J., Moriarty, N. W., Oeffner, R.,
Read, R. J., Richardson, D. C., Richardson, J. S., Terwilliger, T. C., and Zwart, P. H. (2010)
PHENIX: a comprehensive Python-based system for macromolecular structure solution. *Acta*
Crystallogr. Sect. D. **66**, 213-221

47. Emsley, P., Lohkamp, B., Scott, W. G., and Cowtan, K. (2010) Features and development of
Coot. *Acta Crystallogr. Sect. D.* **66**, 486-501

48. McCoy, A. J., Grosse-Kunstleve, R. W., Adams, P. D., Winn, M. D., Storoni, L. C., and Read, R. J.
(2007) Phaser crystallographic software. *Journal of applied crystallography* **40**, 658-674

49. Smart, O. S., Womack, T. O., Flensburg, C., Keller, P., Paciorek, W., Sharff, A., Vonrhein, C., and
Bricogne, G. (2012) Exploiting structure similarity in refinement: automated NCS and target-
structure restraints in BUSTER. *Acta Crystallographica Section D: Biological Crystallography*
68, 368-380

50. Datsenko, K. A., and Wanner, B. L. (2000) One-step inactivation of chromosomal genes in
Escherichia coli K-12 using PCR products. *Proceedings of the National Academy of Sciences* **97**,
6640-6645

51. Baba, T., Ara, T., Hasegawa, M., Takai, Y., Okumura, Y., Baba, M., Datsenko, K. A., Tomita, M.,
Wanner, B. L., and Mori, H. (2006) Construction of Escherichia coli K-12 in-frame, single-gene
knockout mutants: the Keio collection. *Molecular systems biology* **2**

52. Tu, Q., Yin, J., Fu, J., Herrmann, J., Li, Y., Yin, Y., Stewart, A. F., Müller, R., and Zhang, Y. (2016)
Room temperature electrocompetent bacterial cells improve DNA transformation and
recombineering efficiency. *Scientific reports* **6**, 24648

53.   Doublet, B., Douard, G., Targant, H., Meunier, D., Madec, J.-Y., and Cloeckaert, A. (2008)
Antibiotic marker modifications of λ Red and FLP helper plasmids, pKD46 and pCP20, for
inactivation of chromosomal genes using PCR products in multidrug-resistant strains. *Journal*
of microbiological methods **75**, 359-361

Figures

Figure 1 The relationship between YncD and characterised TBDTs and the origin of bacteria that possess YncD (A) Sequence clustering analysis (E-value cutoff = 1×10^{-10}) of YncD with TBDTs of known structure and/or function. Sequences are colored-coded according to the sequence cluster to which they belong. The circle representing the sequence of YncD is enlarged and colored red, showing that YncD does not cluster with other sequences. (B) The environment or host of isolation of a subset of the YncD containing bacteria summarised in Table S1. (C) The environment type of YncD containing bacterial isolates from environmental sources.

Figure 2 The crystal structure of YncD reveals structural homology to the distantly-related

transporter FecA. (A) The crystal structure of YncD shown as a cartoon representation with

rainbow coloring from the N-terminus (blue) to the C-terminus (red). Selected loops are

labelled for reference i.e. L3 = Loop 3. The outer membrane-embedded region is boxed and

with the label 'OM'. Ca^{2+} ions and βOG detergent molecules co-crystallised with YncD are

shown and sphere and stick representations respectively. (B) The crystal structure of YncD

with the barrel domain shown as a cutaway surface and stick representation and the plug-

domain shown as a cartoon representation. The labile section of the plug domain displaced

by TonB is shown in blue, including the TonB box, which is disordered in the structure. (C) The

extracellular loops of YncD, FecA (PDB ID = 1KMP) and FhuE (PDB ID = 6E4V) shown as a

cartoon representation, with rainbow coloring as in panel A.

Figure 3: YncD possesses a conserved positively charged substrate-binding pocket. (A) A

cartoon representation of YncD colored as in Figure 1A (left) and a surface representation in

the same orientation showing the substrate-binding pocket (right). (B) A stereoview of the

amino acids in the YncD substrate-binding pocket which are conserved in YncD homologues

according to the program Consurf (33). The region shown corresponds to the boxed region in

panel A, YncD is shown as a cartoon representation and conserved residues are shown as

sticks. (C) A stereoview of the region of YncD from panel B shown as an electrostatic surface

(top left), as surface colored according to Consurf conservation scores: dark purple = highly

conserved, lighter purple = somewhat conserved, light blue = less conserved, dark blue =

poorly conserved (top right). An electron static surface of FecA in complex with Fe-citrate

shown as a stick and sphere representation (bottom left) and the position of Fe-citrate in the

YncD binding pocket when the FecA-Fe-citrate complex with superimposed with YncD

(bottom right). (D) A stereoview of positively charged conserved residues in the YncD

substrate-binding pocket.

Figure 4 YncD does not import ferric citrate or assist in iron acquisition under laboratory

conditions. (A) A growth curve of E. *coli* ΔTBDT and E. *coli* ΔTBDT/*ΔyncD* grown in M9 minimal

media in the presence or absence of 10 mM citrate or 100 µM 2,3-dihydroxybenzoic acid. (B)

A zoomed view of the growth curve from panel A hours 1 to 7. (C) the growth of wildtype *E.*

coli BW25113 under the same conditions as the growth curve shown in panel A.

Table 1 YncD crystallography data collection and refinement statistics

	K$_2$Pt(NO$_2$)$_4$ Soak	Native
Data Collection[a]		
Space Group	*P2$_1$22$_1$*	*P2$_1$22$_1$*
Cell Dimensions		
a, b, c (Å)	90.61, 96.28, 127.39	87.85, 98.47, 124.70
α, β, γ (°)	90, 90, 90	90, 90, 90
Wavelength	0.9763	0.9763
Resolution (Å)	48.14-3.50 (3.83-3.50)	49.24-3.20 (3.42-3.20)
R$_{merge}$	38.6 (665.70)	28.1 (202.0)
R$_{pim}$	11.7 (201.2)	11.4 (79.7)
I/s(*I*)	7.6 (0.8)	7.9 (1.7)
CC(1/2)	0.999 (0.338)	0.998 (0.757)
Completeness (%)	99.8 (99.5)	100.0 (100.0)
Redundancy	22.1 (22.5)	13.2 (13.1)
No. unique reflections	14619 (3395)	18454 (3272)
Refinement statistics		
R$_{work}$/R$_{free}$		27.4/31.4
No. atoms		5127
Average B-factors		
Protein/Ligand		84.29, 77.10
R.m.s deviations		
Bond lengths (Å)		0.005
Bond angles (°)		0.735
Ramachandran Plot		
Favored/Allowed/Outliers (%)		94.1, 5.43, 0.47
PDB ID		6V81

[a] Values in parentheses are for the highest-resolution shell.