Degrees of compositional shift in tree communities vary along a gradient of

temperature change rates over one decade: Application of an individual-based

temporal beta-diversity concept

Running title: Individual-based temporal-beta diversity

Ryosuke Nakadai[*12]

ORCID: 0000-0002-9512-8511

[1] Department of Environmental and Biological Sciences, Faculty of Science and Forestry,

University of Eastern Finland, Yliopistokatu 7, 80101 Joensuu, Finland

[2] Department of Ecosystem Studies, Graduate School of Agricultural and Life Sciences,

The University of Tokyo,1-1-1 Yayoi, Bunkyo-ku, Tokyo 113-8657, Japan

* Correspondence: R. Nakadai

E-mail: **r.nakadai66@gmai.com**

Declarations

Author contribution–

R.N. conceived the study, designed the methods described herein, conducted the

analyses using empirical data, and wrote the manuscript.

Acknowledgments–

I thank Drs. Anu Valtonen and Shunsuke Matsuoka who made insightful comments

on an early version of the manuscript and Dr. Takaya Iwasaki who advised on the

data visualizations. I also thank my friend for inspiring me to reconsider the

importance and the preciousness of individual life in both society and ecology.

Funding–

Financial support was provided by the Japan Society for the Promotion of Science

(grant 18J00093).

Conflicts of interest–

I declare no conflicts of interest.

Abstract

Temporal patterns in communities have gained widespread attention recently, to the

extent that temporal changes in community composition are now termed "temporal

beta-diversity". Previous studies of beta-diversity have made use of two classes of

dissimilarity indices: incidence-based (e.g., Sørensen and Jaccard dissimilarity) and

abundance-based (e.g., Bray–Curtis and Ružička dissimilarity). However, in the

context of temporal beta-diversity, the persistence of identical individuals and

turnover among other individuals within the same species over time have not been

considered, despite the fact that both will affect compositional changes in

communities. To address this issue, I propose new index concepts for beta-diversity

and the relative speed of compositional shifts in relation to individual turnover based

on individual identity information. Individual-based beta-diversity indices are novel

dissimilarity indices that consider individual identity information to quantitatively

evaluate temporal change in individual turnover and community composition. I

applied these new indices to individually tracked tree monitoring data in deciduous

and evergreen broad-leaved forests across the Japanese archipelago with the objective

of quantifying the effect of climate change trends (i.e., rates of change of both annual

mean temperature and annual precipitation) on individual turnover and compositional

shifts at each site. A new index explored the relative contributions of mortality and

recruitment processes to temporal changes in community composition. Clear patterns

emerged showing that an increase in the temperature change rate facilitated the

relative contribution of mortality components. The relative speed of compositional

shift increased with increasing temperature change rates in deciduous forests but

decreased with increasing warming rates in evergreen forests. These new concepts

provide a way to identify novel and high-resolution temporal patterns in

communities.

Keywords: climate change, compositional equilibrium, compositional shift,

individual-based beta-diversity, individual-tracked monitoring data, mortality,

recruitment

Introduction

All living organisms have finite life spans. Each individual tries to leave offspring

before its death. The persistence of individuals and their reproduction in the past shape

the current diversity patterns through temporal dynamics of populations, communities,

and ecosystems. The effects of recent anthropogenically induced climate change on

local and global biodiversity have received considerable attention (Parmesan and Yohe

2003, Chen et al. 2011). Specifically, intensive efforts have focused on biodiversity

loss (i.e., alpha- and gamma-diversity) and compositional shifts (i.e., beta-diversity)

within the context of biodiversity conservation (e.g., Dornelas et al. 2014, Valtonen et

al. 2016, Hillebrand et al. 2017). Ideally, long-term biodiversity monitoring programs

from multiple countries employing consistent methodologies should be used to identify

the effects of global climate change. However, multi-decade monitoring programs

remain very rare (Magurran et al. 2010; but see Valtonen et al. 2016).

Whittaker (1960, 1972) developed the concept of beta-diversity, which he defined

as the variation in community composition among sites in a region. Since then, many

new beta-diversity indices have been developed, and attempts have been made to

relate the properties of beta-diversity and dissimilarity indices to background

ecological processes (Harrison et al. 1992, Williams 1996, Lennon et al. 2001,

Tuomisto and Ruokolainen 2006, Baselga 2010, Legendre 2010; reviewed by Koleff

et al. 2003, Anderson et al. 2011, Jost et al. 2011, Legendre and De Cáceres 2013).

More attention has been focused recently on temporal changes in community

composition in either single sites or in a series of sites surveyed repeatedly over time

(Magurran 2011, Bahram et al. 2015, Matsuoka et al. 2016, Legendre and Condit

2019). Temporal changes in community composition are now termed "temporal beta-

diversity" (Hatosy et al. 2013, Legendre and Gauthier 2014, Shimadzu et al. 2015).

Legendre (2019) elegantly partitioned temporal beta-diversity into loss and gain

components of the compositional changes in each community, thereby facilitating

mechanistic understanding of the temporal changes in community composition.

Furthermore, recent studies have started to show that the temporal beta-diversity

patterns differ among sites (Legendre and Condit 2019, Magurran et al. 2019).

Environmental changes, including anthropogenically induced shifts such as global

climate change, are likely factors that drive spatial differences in temporal beta-

diversity (Legendre 2019).

The persistence of individuals over time is an important component of the temporal

beta-diversity concept that has not been explicitly considered previously. The

persistence of individuals contributes to a static equilibrium in community

composition. By contrast, turnover of different individuals belonging to the same

species, even when the community composition is stable over time, contributes to a

dynamic equilibrium in community composition. Therefore, conventional temporal

beta-diversity is zero when the community composition is at equilibrium. However, it

can be difficult to detect differences in turnover within the same species and the

speed or frequency of compositional changes in communities differs between sites

with high and low individual turnover rates across time. Therefore, an appropriate

evaluation of temporal changes in community composition could benefit from

considering the fates of identifiable individuals in a community over time. Here, I

propose new indices of temporal beta-diversity that take into account individual

identity information. These new coefficients are individual-based in contrast to others

that are incidence-based (e.g., the Sørensen index; Sørensen 1948), or abundance-

based (e.g., the Bray–Curtis index; Bray and Curtis 1957). My new indices

quantitatively evaluate the temporal changes in community composition by

considering individual identity information and reveal the relative contribution of

different background processes to these temporal changes. I also propose a new index

to quantify the relative speed of change away from compositional equilibrium in a

community relative to the speed of individual turnover. First, I calculated these

indices for 16 forest inventory datasets compiled across the Japanese archipelago.

These datasets have individual identity information for two time periods. I then aimed

to empirically quantify the extent of the compositional shift response to climate

change by excluding the effect of differences in the speed of individual turnover

among site. The indices identified the effects of climate change on the composition of

tree communities on a decadal time scale.

Materials and Methods

New indices of temporal beta-diversity and new statistics to evaluate the relative speed

of compositional shift in relation to individual turnover

Two types of dissimilarity indices have been proposed in previous studies: incidence-

based and abundance-based (Baselga et al. 2013). Specifically, the widely used Bray–

Curtis dissimilarity index (Bray and Curtis 1957, which was proposed earlier as a

proportional difference index by Odum 1950), is an abundance-based extension of

the Sørensen index (Legendre and Legendre 1998, Baselga et al. 2013). I have

excluded details of incidence-based indices that are tangential to the main topic in my

study (see Anderson et al. 2011, Jost et al. 2011, Baselga et al. 2013). The conceptual

basis for conventional beta-diversity indices and my new procedures relies on the fact

that the intersection (*A* component) and the relative complements (*B* and *C*

components) for species abundance can be formulated as follows:

$A = \sum_i \min\left(x_{ij}, x_{ik}\right)$ (1)

$B = \sum_i x_{ij} - \min\left(x_{ij}, x_{ik}\right)$ (2)

$C = \sum_i x_{ik} - \min\left(x_{ij}, x_{ik}\right)$ (3)

where x_{ij} is the abundance of species i at site j, and x_{ik} is the abundance of species i at

site k. Therefore, A is the number of individuals of each species that occur at both

sites j and k, while B and C are the numbers of individuals that are unique to sites j

and k, respectively. This formulation of the Bray–Curtis (d_{BC}) index is expressed as

follows:

$$d_{BC} = \frac{B + C}{2A + B + C}.$$ (4)

In the temporal beta-diversity concept, the differences between data vectors

correspond to observations made at time 1 (abbreviated T1) and time 2 (T2), which are

referred to sites k and j in the spatial beta-diversity concept. The A, B, and C

components are described as follows:

$$A_t = \sum_i \min(y_{i1}, y_{i2})$$ (5)

$$B_t = \sum_i y_{i1} - \min(y_{i1}, y_{i2})$$ (6)

$$C_t = \sum_i y_{i2} - \min(y_{i1}, y_{i2})$$ (7)

where y_{i1} is the abundance of species i at time T1 at a site, and y_{i2} is the abundance of

species i at time T2 at the site. Therefore, A_t is the number of individuals of each

species that occur at both times T1 and T2, whereas B_t and C_t are the numbers of

individuals that are unique to times T1 and T2, respectively. The temporal beta-

diversity of the Bray–Curtis ($d_{t.BC}$) index is expressed as follows:

$$d_{t.BC} = \frac{B_t + C_t}{2A_t + B_t + C_t}.$$ (8)

Using Equation 8, Legendre (2019) partitioned temporal beta-diversity indices into loss

and gain components because a change through time is directional; species abundances

would identify losses and/or gains between T1 and T2 as follows:

$$d_{t.BC} = \frac{B_t + C_t}{2A_t + B_t + C_t}$$

$$= \frac{B_t}{2A_t + B_t + C_t} + \frac{C_t}{2A_t + B_t + C_t} = d_{t.loss} + d_{t.gain} \quad . \qquad (9)$$

Legendre and his collaborators applied the loss/gain partitioning procedure for

temporal beta-diversity to many empirical datasets (e.g., Kuczynski et al. 2018, Brice

et al. 2019), and constructed a new analytical trend in temporal beta-diversity.

Individual-based components and novel indices

In this study, I developed the concept of individual-based temporal beta-diversity

based on Bray–Curtis dissimilarity through extension of the abundance-based temporal

beta-diversity indices. Traditionally, the equilibrium and non-equilibrium elements in

community ecology indicate properties of the whole community that are directly

related to species coexistence (Pickett 1980). However, some individuals replace others,

and thus ecological communities are always dynamic and vary by at least some degree

in space and time (Mori et al. 2018, Ryo et al. 2019). In addition, some individuals of

long-lived species (e.g., trees and vertebrates) persist over the long term, whereas

individuals of short-lived species are replaced by conspecifics or different species.

Therefore, it should be possible to categorize the components of community dynamics

into persistence (P), and those that change over time, i.e., mortality and recruitment (M

and R, respectively). M refers to individuals occurring at T1 that die before T2, thereby

representing the death of individuals during the time period between T1 and T2 (Fig.

1). R identifies individuals that were not alive at T1 but were subsequently found at T2,

thereby representing recruitment during the period between T1 and T2. Component P

refers to the persistence of individuals from T1 to T2 (Fig. 1). Although P, M, and R

relate to components A, B, and C, respectively, here I have used different characters to

emphasize the processes represented by each component. The respective abundances

were calculated as follows:

$$P = \sum_l \min(z_{l1}, z_{l2}) \qquad (10)$$

$$M = \sum_l z_{l1} - \min(z_{l1}, z_{l2}) \qquad (11)$$

$$R = \sum_l z_{l2} - \min(z_{l1}, z_{l2}) \qquad (12)$$

where z_{l1} is individual l at time T1, and z_{l2} is individual l at time T2. Both z_{l1} and z_{l2}

take only 1 or 0, i.e., presence or absence. Therefore, P is the number of individuals

present at both T1 and T2, whereas M and R are the numbers of individuals that are

unique to T1 or T2, respectively. This formulation of the individual-based temporal

beta-diversity index (d_{MR}) can be expressed as follows:

$$d_{MR} = \frac{M + R}{2P + M + R}. \qquad (13)$$

Furthermore, this individual-based temporal beta-diversity can be partitioned into

the relative contributions of mortality (d_M) and recruitment (d_R) processes using a

similar procedure for the loss and gain component-partitioning of the temporal beta-

diversity in Equation 9 (Legendre 2019) as follows:

$$d_{MR} = \frac{M + R}{2P + M + R}$$

$$= \frac{M}{2P + M + R} + \frac{R}{2P + M + R} = d_M + d_R. \qquad (14)$$

In some cases, there is turnover of different individuals belonging to the same

species even when the community composition is stable over time, which contributes

to a dynamic compositional equilibrium across time periods. Therefore, the

components that change over time (M and R) are also separated into two components;

compositional equilibrium (E) and shift (B_t and C_t), respectively. Component M can be

partitioned into lost individuals that contribute to compositional change in a

community (B_t) and lost individuals that are replaced by conspecifics, contributing

substantially to equilibrium in community composition (E_{loss}). Similarly, component R

can be partitioned into gained individuals that contribute to compositional change in a

community (C_t), and gained individuals replacing conspecifics, thereby contributing

substantially to equilibrium in community composition (E_{gain}). By definition, the

numbers of E_{loss} and E_{gain} individuals are identical; hence, I replaced both E_{loss} and E_{gain}

with the coefficient E, the component that contributes to dynamic compositional

equilibrium. The formulation is as follows:

$$
\begin{aligned}
d_{MR} &= \frac{M + R}{2P + M + R} \\
&= \frac{B_t + E_{loss} + C_t + E_{gain}}{2P + M + R} = \frac{B_t + C_t + 2E}{2P + M + R} \\
&= \frac{2E}{2P + M + R} + \frac{B_t + C_t}{2P + M + R} = d_E + d_S
\end{aligned}
\tag{15}
$$

where d_E and d_S indicate the contributions of equilibrium and shift to individual-based

temporal beta-diversity. I note that E is equal to the A_t component minus the P

component.

Both d_M/d_R and d_E/d_S are indicators of the relative contribution to individual-

based temporal beta-diversity (d_{MR}); they represent rates of mortality vs. recruitment

processes, and compositional equilibrium to shift, respectively. Furthermore, the

components M and R can also be analyzed in the same manner as community data (Fig.

1). Therefore, the calculation of Bray–Curtis dissimilarity between communities using

mortality and recruitment is as follows:

$$
\begin{aligned}
v_s &= \frac{B_t + C_t}{2E + B + C_t} \\
&= 1 - \frac{2E}{2E + B_t + C_t} = 1 - \frac{2E}{M + R} = 1 - v_e .
\end{aligned}
\tag{16}
$$

255 v_s indicates the speed of compositional shifts in a community relative to the total speed

256 of individual turnover through mortality and recruitment processes. If v_s is 0, then

257 individual turnover would not contribute to compositional shifts in a community, i.e.,

258 the community composition of mortality and recruitment are identical and an

259 equilibrium state exists. If v_s is 1, all individual turnovers contribute to compositional

260 shifts, and the community composition of mortality and recruitment are completely

261 different. It is also possible to calculate v_s as $d_{t.BC}/d_{MR}$. Although I did not use v_e for

262 analyses in the present study, v_e should be an index of dynamic compositional

263 equilibrium. The relative contributions of compositional equilibrium and shift (d_E/d_S)

264 are formulated as follows:

266 $$d_E \div d_S = \frac{2E}{2P + M + R} \div \frac{B_t + C_t}{2P + M + R} = \frac{2E}{B_t + C_t} = \frac{2E}{M + R - 2E} \quad (17)$$

268 Because "$2E$" in the denominator is simply the difference between v_e (equation 16) and

269 d_E/d_S (equation 17), the two are strongly correlated (Table S1), but these two indices

270 have different meanings.

271   In my main analyses, I focused only on an extension of Bray–Curtis

272 dissimilarity. I also summarized explanations of individual-based beta-diversity using

273 the Ružička dissimilarity index (Ružička 1958), which was identified by Podani et al.

274 (2013) and Legendre (2014) as another appropriate index for beta-diversity studies.

275 The Ružička dissimilarity index is also an abundance-based extension of the Jaccard

276 dissimilarity index (Jaccard 1901). The explanations of individual-based beta-diversity

277 using the Ružička dissimilarity index are described Supplementary text 1.

279 *Applying Japanese forest monitoring datasets to testing patterns in temporal beta-*

280 *diversity indices along climate change gradients*

281 I predicted the indices of individual-based beta-diversity and the relative speed of

282 compositional shift variation along degrees of climate change rate per year, i.e.,

283 average speed of local climate change. I applied the new indices using individual-

tracked monitoring data from forests in Japan. Two types of temporal change in tree

communities may result from climate change. One is due to changes in the obligate

speed of individual turnover and the balance of mortality and recruitment, e.g.,

climate warming facilitates the faster growth and shorter life span of each individual

tree as a physiological process within individuals (Körner 2017, Munné-Bosch 2018).

The other is due to the relative speed of compositional change, e.g., simple

compositional shifts driven by climate change as an ecological process among

individuals (i.e., competitive interactions and adaptation to the environment). Those

can be referred to as d_{MR}, including the components (d_M, d_R) and v_s, respectively.

The Japanese archipelago comprises a long chain of continental islands lying off the

eastern coast of Asia; it is recognized as a hotspot of biodiversity (Mittermeier et al.

2011). The mean temperatures of the coldest months range from −18.4 to 22.3°C, mean

temperatures of the warmest months from 6.2 to 29.2°C, and the annual precipitation

ranges from 700.4 to 4,477.2 mm (Japan Meteorological Agency, 1981–2010;

http://www.jma.go.jp/jma/indexe.html). Typical forest biomes are classified into four

types: alpine meadow/creeping pine zone, subalpine coniferous forest zone, deciduous

broad-leaved forest zone, and evergreen broad-leaved forest zone. The empirical tree

census dataset used in this study was obtained from the forest plots used in the

Monitoring Sites 1000 Project; this project was launched by the Ministry of the

Environment, Japan (see Ishihara et al. 2011 for details). The monitoring sites include

five types of forest: deciduous broad-leaved, evergreen broad-leaved, mixed broad-

leaved and coniferous, evergreen coniferous, and artificial forest (Ishihara et al., 2011,

Suzuki et al. 2015). The full dataset is available on the website of the Biodiversity

Center of Japan (http://www.biodic.go.jp/moni1000/index.html, in Japanese: accessed

on November 11, 2019), and a subset of the data has been published as a data article

(Ishihara et al. 2011).

To apply the individual-based temporal beta-diversity approach, I selected 16 forest

plots. These included nine plots of deciduous broad-leaved forest and seven plots of

evergreen broad-leaved forest (Fig. 2, Table S2) that met the following criteria: species

level identifications had been performed in each plot, the plots had been surveyed one

decade after a previous survey (in the period 2005–2008 to 2015–2018), individual

identity information was recorded throughout the two time periods, plot size was 1.0

ha (100 × 100 m), and sufficient plot sample sizes in various environments within each

forest type were available for use in analyses. In this analysis, I selected tree individuals

with stem girths at breast height (GBH) ≥ 15.7 cm, which was approximately

equivalent to a diameter at breast height (DBH) ≥ 5 cm; these were considered target

individuals in the communities. I organized the data at the level of individual trees

before the analyses, the original forest datasets had been compiled at the stem level,

using taxonomic information provided in the Ylist (Yonekura and Kajita 2003). The

data I analyzed included 18,749 individuals belonging to 254 species found in the 16

plots (Fig. 2).

Climate variables

To obtain meteorological observation data and their changes over a decade, I searched

for the meteorological station closest to each of the forest plots in the Japanese

Automated Meteorological Data Acquisition System database (Japan Meteorological

Agency; see details in Table S2, Fig. S1). Specifically, I extracted the monthly mean

and total precipitation data (accessed December 12, 2019), and calculated the annual

mean temperature (°C) and annual precipitation (mm) for each year and site. I

estimated temperature differences from elevation differences between each of the

forest plots and its nearest meteorological station using 0.55°C/100 m in elevation as

the mean lapse rate; this relationship between temperature and elevation is commonly

used by Japanese forestry researchers (e.g., Oshima et al. 1961, Fukushima et al. 2009).

I regressed annual mean temperature and annual precipitation against year for each plot

using an ordinary least squares approach in R (R Development Core Team 2019). The

resulting regression coefficients indicated the average temperature change per year in

the targeted decade, and hence average change rate in annual mean temperature and

annual precipitation in each plot. The average rates of change in annual mean

temperature (°C) and annual precipitation (mm) were 0.004 ± 0.047 (−0.105 to 0.063)

and 5.714 ± 32.795 (−37.082 to 77.791), respectively, in deciduous broad-leaved

forests, and 0.003 ± 0.043 (−0.040 to 0.059) and 20.860 ± 18.737 (−0.176 to 47.261),

respectively, in evergreen broad-leaved forests [mean ± SD (range)].

Modeling patterns of existing and novel indices describing community changes across

time

I focused mainly on the following indices: $d_{t.BC}$, d_{MR}, d_M/d_R, d_E/d_S, and v_s. Formulas

and ecological interpretation are summarized in Table 1. First, I analyzed the

relationships between these indices and the average rates of change in annual mean

temperature and annual precipitation using both simple and polynomial regressions.

Then I compared the Akaike's information criterion (AIC) values (Akaike 1976)

among models, including the null model, and selected the best model for each. I did

not use multiple regressions for further analyses because the number of data points

available was insufficient. All statistical analyses were conducted in R. R scripts to

calculate the new indices are provided in Supplementary file 1.

Results

I detected no trends in $d_{t.BC}$ and d_{MR} in relation to average rate of change in annual mean

temperature in either deciduous or evergreen broad-leaved forests (Fig. 3a, b, c, d,

Table S3). In analyses of d_M/d_R, the polynomial model was chosen by the AIC

comparison as the best model for deciduous broad-leaved forests; the simple model

was the best for evergreen broad-leaved forests. Values of d_M/d_R tended to be greater at

sites that warmed the most in the targeted decades. This was true for both forest types.

Thus, the relative contribution of mortality to individual based beta-diversity increased

with warming rate over time (Fig. 3e, f, Table S3). The d_E/d_S index did not show any

statistical trends over time for deciduous forests (Fig. 3g, Table S3); the simple

regression model was chosen by AIC comparison as the best for evergreen forests (Fig.

3h, Table S3). In evergreen forests, d_E/d_S increased most over a decade in warming

sites. Thus, the relative contribution of individual turnover without compositional shift

in relation to that with compositional shift increased over time (Fig. 3h, Table S3). The

polynomial model of v_s was selected as the best for the deciduous forest type: v_s

increased with larger temperature changes (Fig. 3i, Table S3). The simple model was

selected for the evergreen forest type: v_s decreased linearly with increasing warming

over time (Fig. 3j, Table S3). The correlations of all calculated indices are provided in

Table S1.

The average rate of change in annual precipitation did not affect trends over time for

any of the calculated indices (Table S3).

Discussion

The effects of anthropogenic climate change have been intensively studied in many

disciplines, e.g., analyses of species interactions, range shifts, phenology, and

ecosystem functioning (Thomas et al. 2004, Kordas et al. 2011, Valtonen et al. 2014).

However, studies of temporal changes in diversity and composition in local

communities have often produced mixed and/or contradictory results among sites and

researchers (Vellend et al. 2013, Dornelas et al. 2014, Newbold et al. 2015, reviewed

in Hisano et al. 2018). Quantitative analyses of biological community change over time

are required to expand our understanding of global climate change and its relationship

to biodiversity change (Dornelas et al. 2014). The novel concepts and procedures that

I have developed quantitatively evaluated the relative speed of compositional shift and

the relative contribution of mortality and recruitment processes in tree communities

along a gradient in temperature change rates across a decade. These patterns had

previously been difficult to detect using existing abundance-based dissimilarity indices

(Fig. 3a, b). Compositional shifts in tree communities are generally recognized as long-

term phenomena in the absence of disturbance. Therefore, 10 years is a relatively short-

time period in which to analyze compositional changes in tree communities comprising

species with life spans extending from decades to centuries (Loehle 1988). I found that

community responses to local climate change trends differed between forest types, but

nevertheless, there were real changes in community composition over only a decade.

These changes were associated with change in annual mean temperatures, but not with

changes in annual precipitation across the Japanese archipelago.

One advantage of the method I have developed is that it dissects individual turnover

into mortality and recruitment components. For example, I detected no clear patterns

in the relationship between individual turnover (d_{MR}), which is the obligate speed of

temporal change in communities, and temperature change rate (Fig. 3c, d). However,

by splitting the index into two components, I was able to identify relative contributions

to individual turnover (d_M/d_R). When this was done, clear patterns emerged showing

that an increase in the temperature change rate facilitated the relative contribution of

mortality components (Fig. 3e, f). Furthermore, by assessing the trend of each

component separately, I was able to confirm that decreases in recruitment components

with increases in the temperature change rate drove the pattern in deciduous forests,

but I was not able to confirm this in evergreen forests (Table S4).

The relative contribution of the index d_E/d_S to individual-based temporal beta-

diversity (d_{MR}) increased positively in evergreen forests as the warming rate increased,

although no trends in the individual components (d_E, d_S) were discerned. Thus, the

balance between the components changed in a consistent manner with temperature

change, although no patterns of variation in the individual components were found.

There were no clear trends in d_E/d_S in relation to temperature change rates in deciduous

forests. However, d_E was affected by the temperature change rate in these forests: d_E

decreased with increased warming rates; thus, there was a clear contribution of

compositional equilibrium decrease along these gradients (Table S4).

I used a novel index to evaluate compositional changes, i.e., the relative speed of the

compositional shift in communities (v_s), which clearly changed along the gradient of

temperature change rates (Fig. 3i, j, Table S3). Furthermore, the responses were

different between deciduous and evergreen broad-leaved forests. Specifically, the

relative speed in evergreen forests was smaller at sites with greater temperature

warming. Furthermore, the range size of the values (v_s) in deciduous forests was about

7-fold greater than the range size in evergreen forests (Table S5). Thus, deciduous

forests are more affected by differences in temperature change rate. I did not detect

statistically significant trends along a gradient of precipitation change rate, although

climate change induced drought is a major factor in tree mortality (Allen et al. 2010,

Saiki et al. 2017). These contrasting results may be due to the humid climate of the

Japanese archipelago, but assessing this relationship requires the inclusion of many

other candidate explanatory variables (e.g., continuous dry days) which were not

available. To capture the effects of global climate change accurately, a broad multi-site

analysis across the globe using similar methods and other relevant environmental

variables will be required.

Predicting community persistence from snapshot data on community structure is a

complicated issue for community ecologists. For example, Ulrich et al. (2018) argued

that it may be possible to evaluate species persistence and their relative proportions

from the patterns of species abundance distribution. If this were the case, the relative

speed of compositional shift should provide a comparable estimate of community

persistence from both individual and species-based perspectives. Specifically, using

some time series datasets containing substantial annual individual information, it

should be technically possible to calculate the retention time for each community, i.e.,

the lifespan durations of all individuals replaced in a focal community. These

calculations could be cross-compared with snapshot community structure data (e.g.,

species abundance patterns). In the present study, I used data for only two time periods,

so this cross-comparison was not possible. Nevertheless, clarification of the

relationship between relative abundance and temporal changes in individual and

community composition should become possible in the near future. Underlying

mechanisms explaining these relationships may then be discernible.

Compilation of long-term temporal datasets for large macroscopic organisms (e.g.,

trees and mammals) requires considerable human effort and cost for monitoring and

research surveys because large organisms often have long life spans. Nevertheless,

the new indices I developed clearly highlighted the importance of monitoring datasets

assembled using traditional detailed surveys (e.g., individual tracking in forest plots)

in detecting the temporal trends over short time periods. Therefore, I re-emphasize

here the significance of classical monitoring surveys to collect biodiversity

information. I also developed an approach that introduced the concept of individual-

based beta-diversity within the context of temporal changes in community

composition. Theoretically, however, the procedure could be expanded to a spatial

context for use in studies of animal behavioral ranges and movements, including

long-distance migration, using monitoring and biologging datasets (Viana et al. 2016,

Börger et al. 2020) that include multiple time series datasets collected from the same

place within a shorter time frame than the life spans of the targeted organisms (e.g.,

bird ringing and classical mark–recapture procedures). Specifically, the approach

could detect detailed spatial turnover for individuals within same species in

communities.

Many ecology theories have been developed from the concept of species. The

original concept of beta-diversity was proposed to reflect species turnover (Whittaker

1960). By contrast, much recent research has pointed out the problems of species-based

studies and the importance of considering individuals in community ecology (Dupont

et al. 2011, Nakazawa 2017, Tatsumi et al. 2019). Species are the consequences of

ecological/evolutionary processes among individuals; to be accurate, the species

themselves do not drive these processes, i.e., individuals do by competing with other

individuals, and this does not involve the species directly. Although individuals were

considered in the unified neutral theory (Hubbel 2001) and subsequent related studies

(Hubert et al. 2015, Vanoverbeke et al. 2016, Tatsumi et al. 2019), the focus was

restricted to the patterns of species richness, abundance, and related diversity indices.

The effects of individual births and deaths on biodiversity patterns have not yet been

considered sufficiently. Procedures for incorporating individual identity information

into biodiversity evaluation, i.e., individual-based diversity indices, are required to

facilitate further conceptual advances in ecology. New ecological concepts will emerge

from combined considerations of species abundance and information relevant to

individual identity in the future.

Conclusions

Previous studies have often detected temporal changes in local communities, likely as

a result of human activities, but not at all times and places. In this study, I clearly

showed the responses of tree communities to local and short-term climate change

trends using datasets collected according to novel index concepts. Notably, the

relative speed of compositional shift in tree communities varied along a gradient of

temperature change rates over a single decade. This community level response

resulted from individual-level mortality and recruitment processes. Therefore,

accumulating more detailed long-term information at global scales, specifically

individual identity information at the local community level, is required for

understanding both actual biodiversity responses to global climate change and

background processes. Mortality and recruitment rates are generally difficult to

measure in empirical studies compared to loss and gain information based only on

snapshot data, especially in the case of plants or vertebrates. The individual-based

approach may not be tractable for all taxonomic groups. However, information on

individual organisms will help to bridge the disciplines of macroecology and local

studies focusing on within-individual phenomena, for example, physiological

ecology. The calculation of individual-based diversity indices from individual identity

monitoring data will support efforts to quantitatively evaluate the impacts of climate

change on future biodiversity. Increasing availability of monitoring data with

individual identification combined with individual-based diversity indices such as

presented here can thus be key in quantitatively evaluating the impacts of climate

change on biodiversity over time.

Data accessibility statement

All data were obtained from open sources. Forest monitoring data were obtained from

the Biodiversity Center of Japan (http://www.biodic.go.jp/moni1000/index.html, in

Japanese: accessed on November 11, 2019) published in part by Ishihara et al. (2011).

Climate data were obtained from the Japanese Automated Meteorological Data

Acquisition System (AMeDAS) database (http://www.jma.go.jp/jma/indexe.html,

Japan Meteorological Agency accessed December 12, 2019). Taxonomic information

was obtained from the Ylist (Yonekura and Kajita 2003:

http://ylist.info/ylist_simple_sea rch.html, in Japanese: accessed December 8, 2019).

References

Akaike, H. 1976. An information criterion (AIC). Math. Sci., 14, 5–7.

Allen, C.D. et al. 2010. A global overview of drought and heat-induced tree mortality

reveals emerging climate change risks for forests. For. Ecol. Manag., 259, 660–

684.

Anderson, M. J. et al. 2011. Navigating the multiple meanings of β diversity: a

roadmap for the practicing ecologist. Ecol. Lett. 14, 19–28.

Bahram, M. et al. 2015. Local-scale biogeography and spatiotemporal variability in

communities of mycorrhizal fungi. New. Phytol. 205, 1454–1463.

Baselga, A. 2010. Partitioning the turnover and nestedness components of beta

diversity. Glob. Ecol. Biogeogr. 19, 134–143.

Baselga, A. 2013. Separating the two components of abundance-based dissimilarity:

balanced changes in abundance vs. abundance gradients. Methods Ecol. Evol., 4,

552–557.

Börger, L. et al. 2020. Biologging special feature. J. Anim. Ecol. 13 January 2020

https://doi.org/10.1111/1365–2656.13163

Bray, J.R., and Curtis, J.T. 1957. An ordination of the upland forest communities of

southern Wisconsin. Ecol. Monogr. 27, 325–349.

Brice, M.H. et al. 2019. Disturbances amplify tree community responses to climate

change in the temperate–boreal ecotone. Glob. Ecol. Biogeogr., 28, 1668–1681.

Chen, I. C. et al. 2011. Rapid range shifts of species associated with high levels of

climate warming. Science 333, 1024–1026.

Dornelas, M. et al. 2014. Assemblage time series reveal biodiversity change but not

systematic loss. Science 344, 296–299.

Dupont, Y.L. et al. 2011. Scaling down from species to individuals: a flower-

visitation network between individual honeybees and thistle plants. Oikos 120,

170–177.

Fukushima, K. et al. 2009 Water yield and nitrogen loss during regrowth of Japanese

cedar forests after clearcutting. In: Taniguchi M, Fukushima Y, Burnett WC,

Haigh M, Umezawa Y (eds) From headwaters to the ocean: hydrological change

and watershed management. Taylor & Francis, London, pp. 97–104

Harrington, R. et al. 1999. Climate change and trophic interactions. Trends Ecol.

Evol. 14, 146–150.

Harrison, S. et al. 1992. Beta diversity on geographic gradients in Britain. J. Anim.

Ecol. 61, 151–158.

Hatosy, S.M. et al. 2013. Beta diversity of marine bacteria depends on temporal

scale. Ecology 94, 1898–1904.

Hillebrand, H. et al. 2018. Biodiversity change is uncoupled from species richness

trends: consequences for conservation and monitoring. J. Appl. Ecol. 55, 169–

184.

Hisano, M. et al. 2018. Biodiversity as a solution to mitigate climate change impacts

on the functioning of forest ecosystems. Biol. Rev. 93, 439–456.

Hubbell, S. P. 2001. The unified neutral theory of biodiversity and biogeography.

Princeton University Press.

Hubert, N. et al. 2015. Metacommunity speciation models and their implications for

diversification theory. Ecol. Lett. 18, 864–881.

Ishihara, M. I. et al. 2011. Forest stand structure, composition, and dynamics in 34

sites over Japan. Ecol. Res. 26, 1007.

Jaccard, P. 1901. Distribution de la flore alpine dans le bassin des Dranses et dans

quelques régions voisines. Bull. Soc. Vaudoise. Sci. Nat. 37, 241–272.

Jost, L. et al. 2011 Compositional similarity and beta diversity. *Biological Diversity:*

Frontiers in Measurement and Assessment (eds A. Magurran and B.J. McGill),

pp. 66–84. Oxford University Press, Oxford.

Koleff, P. et al. 2003. Measuring beta diversity for presence-absence data. J. Anim.

Ecol. 72, 367–382.

Kordas, R. L. et al. 2011. Community ecology in a warming world: the influence of

temperature on interspecific interactions in marine systems. J. Exp. Mar. Biol.

Ecol. 400, 218–226.

Körner, C. 2017. A matter of tree longevity. Science 355, 130–131.

Kuczynski, L. et al. 2018. Concomitant impacts of climate change, fragmentation and

non-native species have led to reorganization of fish communities since the

1980s. Glob. Ecol. Biogeogr. 27, 213–222.

Legendre, P., and Condit, R. 2019. Spatial and temporal analysis of beta diversity in

the Barro Colorado Island forest dynamics plot, Panama. For. Ecosyst. 6, 7.

Legendre, P., and De Cáceres, M. 2013. Beta diversity as the variance of community

data: dissimilarity coefficients and partitioning. Ecol. Lett 16, 951–963.

Legendre, P. and Gauthier, O. 2014. Statistical methods for temporal and space–time

analysis of community composition data. Proc. Roy. Soc. B. 281, 20132728.

Legendre, P. and Legendre, L. 1998. Numerical Ecology, 2nd edn. Elsevier,

Amsterdam.

Legendre, P. 2014. Interpreting the replacement and richness difference components

of beta diversity. Glob. Ecol. Biogeogr. 23, 1324–1334.

Legendre, P. 2019. A temporal beta-diversity index to identify sites that have changed

in exceptional ways in space‑time surveys. Ecol. Evol. 9, 3500–3514.

Lennon, J. J. et al. 2001. The geographical structure of British bird distributions:

diversity, spatial turnover and scale. J. Anim. Ecol. 70, 966–979.

Loehle, C. 1988. Tree life history strategies: the role of defenses. Can. J. For. Res. 18,

209–222.

Magurran, A. E. 2011. Measuring biological diversity in time (and space). *Biological*

Diversity: Frontiers in Measurement and Assessment (eds A. Magurran & B.J.

McGill), pp. 85–104. Oxford University Press, Oxford.

Magurran, A. E. et al. 2010. Long-term datasets in biodiversity research and

monitoring: assessing change in ecological communities through time. Trends

Ecol. Evol. 25, 574–582.

Magurran, A. E. et al. 2019. Temporal β diversity: A macroecological perspective.

608   Glob. Ecol. Biogeogr. 28, 1949–1960.

609 Matsuoka, S. et al. 2016. Temporal distance decay of similarity of ectomycorrhizal

610   fungal community composition in a subtropical evergreen forest in Japan. FEMS

611   Microbiol. Ecol. 92.

612 Mittermeier R.A. et al. 2011. Global biodiversity conservation: the critical role of

613   hotspots. In: Zachos FE, Habel JC (eds) Biodiversity hotspots. Springer, Berlin,

614   pp 3–22

615 Mori, A. S. et al. 2018. β-diversity, community assembly, and ecosystem functioning.

616   Trends Ecol. Evol. 33, 549–564.

617 Munné-Bosch, S. 2018. Limits to tree growth and longevity. Trends Plant Sci. 23,

618   985–993.

619 Nakazawa, T. 2017. Individual interaction data are required in community ecology: a

620   conceptual review of the predator-prey mass ratio and more. Ecol. Res. 32, 5–

621   12.

622 Newbold, T. et al. 2015. Global effects of land use on local terrestrial biodiversity.

623   Nature 520, 45.

624 Odum, E.P. 1950. Bird populations of the Highlands (North Carolina) Plateau in

625   relation to plant succession and avian invasion. Ecology 31, 587–605.

626 Oshima, Y. 1961. Ecological studies of *Sasa* communities. I. Productive structure of

627   some of the *Sasa* communities in Japan. Bot. Mag. Tokyo 74, 199–210.

628 Parmesan, C., and Yohe, G. 2003. A globally coherent fingerprint of climate change

629   impacts across natural systems. Nature 421, 37.

630 Pickett, S. T. 1980. Non-equilibrium coexistence of plants. Bull. Torrey. Bot. Club.

631   238–248.

632 Podani, J. et al. 2013. A general framework for analyzing beta diversity, nestedness

633   and related community-level phenomena based on abundance data. Ecol.

634   Complex. 15, 52–61.

R Development Core Team 2019. R: a language and environment for statistical

computing. R Foundation for Statistical Computing, Vienna, Austria.

http://www.R-project.org/.

Ružička, M. 1958. Anwendung mathematisch–statisticher Methoden in der

Geobotanik (synthetische Bearbeitung von Aufnahmen). Biologia, Bratislava 13,

647–661.

Ryo, M. et al. 2019. Basic principles of temporal dynamics. Trends Ecol. Evol. 34,

723–733

Saiki, S. T. et al. 2017. Physiological mechanisms of drought-induced tree die-off in

relation to carbon, hydraulic and respiratory stress in a drought-tolerant woody

plant. Sci. Rep. 7, 2995.

Shimadzu, H. et al. 2015. Measuring temporal turnover in ecological communities.

Methods Ecol. Evol. 6, 1384–1394.

Sørensen, T. A. 1948. A method of establishing groups of equal amplitude in plant

sociology based on similarity of species content and its application to analyses

of the vegetation on Danish commons. Biol. Skar. 5, 1–34.

Suzuki, S. N. et al. 2015. Regional-scale directional changes in abundance of tree

species along a temperature gradient in Japan. Glob. Change Biol. 21, 3436–

3444.

Tatsumi, S. et al. 2019. Individual-based models of community assembly:

Neighbourhood competition drives phylogenetic community structure. J. Ecol.

107, 735–746.

Thomas, C. D. et al. 2004. Extinction risk from climate change. Nature, 427, 145.

Tuomisto, H., and Ruokolainen, K. 2006. Analyzing or explaining beta diversity?

Understanding the targets of different methods of analysis. Ecology 87, 2697–

2708.

Ulrich, W. et al. 2018. The two-parameter Weibull distribution as a universal tool to

model the variation in species relative abundances. Ecol. Complex. 36, 110–116.

Valtonen, A. et al. 2017. Long-term species loss and homogenization of moth

communities in Central Europe. J. Anim. Ecol. 86, 730–738.

Valtonen, A. et al. 2014. Is climate warming more consequential towards poles? The

phenology of Lepidoptera in Finland. Glob. Change Biol. 20, 16–27.

Vanoverbeke, J. et al. 2016. Community assembly is a race between immigration and

adaptation: eco-evolutionary interactions across spatial scales. Ecography 39,

858–870.

Vellend, M. et al. 2013. Global meta-analysis reveals no net change in local-scale

plant biodiversity over time. PNAS 110, 19456–19459.

Viana, D. S. et al. 2016. Migratory birds as global dispersal vectors. Trends Ecol.

Evol. 31, 763–775.

Whittaker, R. H. 1960. Vegetation of the Siskiyou mountains, Oregon and California.

Ecol. Monogr. 30, 279–338.

Whittaker, R. H. 1972. Evolution and measurement of species diversity. Taxon 21,

213–251.

Williams, P. H. 1996. Mapping variations in the strength and breadth of

biogeographic transition zones using species turnover. Proc. Roy. Soc. B. 263,

579–588.

Yonekura, K. and Kajita, T. 2003. BG Plants Ylist: an online service of Japanese plant

names, including a nomenclature index (in Japanese).

(http://ylist.info/ylist_simple_search.html) accessed Dec 8 2019

Figure 1 Schematic representation of individual-based temporal beta-diversity. T1 and T2 refer to the communities at times 1 (abbreviated T1) and 2 (T2), respectively. Both communities comprise four species (sp. 1, 2, 3, and 4) and all individuals are identified separately (*Zl*). M refers to individuals in existence at T1 but dead before T2, thereby separately identifying the deaths of individuals during the period between T1 and T2. R identifies individuals that were not alive at T1, but found at T2, thereby identifying the recruitment of individuals during the period between T1 and T2. P refers to individuals that persisted from T1 to T2, thus both T1 and T2 are 1. The arrows indicate the temporal trajectory for each individual; specifically, when they are present or absent in communities between T1 and T2.

Figure 2 Locations of the study sites within the Japanese archipelago. The crosses and asterisks indicate the locations of targeted plots in deciduous broad-leaved forest (DB) and evergreen broad-leaved forest (EB) included in the present study. The lettering adjacent to each symbol is the plot identification code (ID). Detailed plot information is listed in Table S3. The colors indicate four major vegetation types: purple, Alpine meadow or creeping pine zone; blue, subalpine coniferous forest zone; red, deciduous broad-leaved forest zone; green, evergreen broad-leaved forest zone. The information on vegetation type at a 1.0-km grid scale was obtained from the results of the fifth censuses (1992–1996) of the National Survey of the Natural Environment in Japan (http://www.biodic.go.jp/dload/mesh_vg.html; in Japanese, accessed December 12, 2019).

Figure 3 Trends in tree community indices across a gradient of temperature change rates over a decade. The left column of panels refers to trends in deciduous broad-leaved forests; the right column of panels refers to trends in evergreen broad-leaved forests. Panels (a) and (b) show the trends in typical abundance-based temporal beta-diversity indices (Bray–Curtis dissimilarity: d_{BC}). Panels (c) and (d) show the trends

in individual-based temporal beta-diversity indices (d_{MR}). Panels (e) and (f) show

trends in the d_M/d_R index, which is an indicator of the relative contribution of

mortality and recruitment processes to individual-based temporal beta-diversity (e:

adjusted-R^2 = 0.523, P = 0.046; f: *adjusted-R^2* = 0.391, P = 0.079). Panels (g) and (h)

show trends in the d_E/d_S index, which is an indicator of the relative contribution of

equilibrium and shift processes to individual-based temporal beta-diversity (h:

adjusted-R^2 = 0.472, P = 0.053). Panels (i) and (j) show trends in the index v_s, which

is an indicator of the relative speed of compositional shift in relation to individual

turnover (i: *adjusted-R^2* = 0.413, P = 0.085; j: *adjusted-R^2* = 0.462, P = 0.056). The

solid black lines indicate statistically significant (P < 0.05) trends. The dashed black

line indicates a marginally significant (0.05 < P < 0.1) trend. The blue dashed lines

enclose the 95% confidence interval envelopes. Detailed results of statistical analyses

are presented in Table S3.

732   **Figure 2**

Figure 3

Average change rate of annual mean temperature (°C/year)

738 Table 1. Formulations and ecological interpretation of Bray–Curtis dissimilarity (top: Bray and Curtis 1957) and the novel indices

739 proposed in this study. y_{il} is the abundance of species i at time T1 at a site, and y_{i2} is the abundance of species i at time T2 at the site.

740 z_{ll} is individual l at time T1, and z_{l2} is individual l at time T2. Both z_{ll} and z_{l2} take only 1 or 0, i.e., presence or absence.

Index	Formula	Ecological interpretation
$d_{t.\,BC}$	$$\frac{B_t + C_t}{2A_t + B_t + C_t} = \frac{\sum_i y_{i1} - \min(y_{i1}, y_{i2}) + \sum_i y_{i2} - \min(y_{i1}, y_{i2})}{2[\sum_i \min(y_{i1}, y_{i2})] + \sum_i y_{i1} - \min(y_{i1}, y_{i2}) + \sum_i y_{i2} - \min(y_{i1}, y_{i2})}$$	The compositional change based on the change of species abundance in a target community across two time periods ($0 \le d_{t.\,BC} \le 1$).
d_{MR}	$$\frac{M + R}{2P + M + R} = \frac{\sum_l z_{l1} - \min(z_{l1}, z_{l2}) + \sum_l z_{l2} - \min(z_{l1}, z_{l2})}{2[\sum_l \min(z_{l1}, z_{l2})] + \sum_l z_{l1} - \min(z_{l1}, z_{l2}) + \sum_l z_{l2} - \min(z_{l1}, z_{l2})}$$	The turnover of individuals by mortality and recruitment in a target community across two time periods ($0 \le d_{MR} \le 1$).
d_M	$$\frac{M}{2P + M + R} = \frac{\sum_l z_{l1} - \min(z_{l1}, z_{l2})}{2[\sum_l \min(z_{l1}, z_{l2})] + \sum_l z_{l1} - \min(z_{l1}, z_{l2}) + \sum_l z_{l2} - \min(z_{l1}, z_{l2})}$$	The contribution of mortality to individual turnover in a target community across two time periods ($0 \le d_M \le 1$).

d_R

$$\frac{R}{2P + M + R} = \frac{\sum_l z_{l2} - \min(z_{l1}, z_{l2})}{2[\sum_l \min(z_{l1}, z_{l2})] + \sum_l z_{l1} - \min(z_{l1}, z_{l2}) + \sum_l z_{l2} - \min(z_{l1}, z_{l2})}$$

The contribution of recruitment to individual turnover in a target community across two time periods ($0 \leq d_R \leq 1$).

d_E

$$\frac{2E}{2P + M + R} = \frac{2[\sum_i \min(y_{i1}, y_{i2}) - \sum_l \min(z_{l1}, z_{l2})]}{2[\sum_l \min(z_{l1}, z_{l2})] + \sum_l z_{l1} - \min(z_{l1}, z_{l2}) + \sum_l z_{l2} - \min(z_{l1}, z_{l2})}$$

The contribution of compositional equilibrium (i.e., individual turnover within the same species) to individual turnover in a target community across two time periods ($0 \leq d_E \leq 1$).

d_S

$$\frac{B_t + C_t}{2P + M + R} = \frac{\sum_i y_{i1} - \min(y_{i1}, y_{i2}) + \sum_i y_{i2} - \min(y_{i1}, y_{i2})}{2[\sum_l \min(z_{l1}, z_{l2})] + \sum_l z_{l1} - \min(z_{l1}, z_{l2}) + \sum_l z_{l2} - \min(z_{l1}, z_{l2})}$$

The contribution of compositional shift (i.e., individual turnover among different species) to individual turnover in a target community across two time periods ($0 \leq d_S \leq 1$).

d_M/d_R	$$\dfrac{M}{R} = \dfrac{\sum_l z_{l1} - \min\,(z_{l1}, z_{l2})}{\sum_l z_{l2} - \min\,(z_{l1}, z_{l2})}$$	The balance between mortality and recruitment in a target community across two time periods, i.e., the abundance ratio between mortality and recruitment.
d_E/d_S	$$\dfrac{2E}{B_t + C_t} = \dfrac{2[\sum_i \min(y_{i1}, y_{i2}) - \sum_l \min(z_{l1}, z_{l2})]}{\sum_i y_{i1} - \min\,(y_{i1}, y_{i2}) + \sum_i y_{i2} - \min\,(y_{i1}, y_{i2})}$$	The balance between compositional equilibrium and shift in a target community across two time periods, i.e., the abundance ratio between individual turnover within same species and among different species.
v_s	$$\dfrac{\dfrac{B_t + C_t}{2E + B_t + C_t}}{} $$ $$= \dfrac{\sum_i y_{i1} - \min\,(y_{i1}, y_{i2}) + \sum_i y_{i2} - \min\,(y_{i1}, y_{i2})}{2[\sum_i \min(y_{i1}, y_{i2}) - \sum_l \min(z_{l1}, z_{l2})] + \sum_i y_{i1} - \min\,(y_{i1}, y_{i2}) + \sum_i y_{i2} - \min\,(y_{i1}, y_{i2})}$$	The relative speed of compositional shift in relation to individual turnover ($0 \leq v_s \leq 1$). v_s is also calculated as d_{BC}/d_{MR}.

Appendices

Table S1. Correlations among geographical and environmental variables and their respective calculated indices.

Table S2. Geographical and climate attributes of each plot and the nearest metrological station. See Fig. 2 for plot locations.

Table S3. Statistical analyses of the relationships between the indices d_{BC}, d_{MR}, d_M/d_R, d_E/d_S, and v_s and the change rate of annual mean temperature and annual precipitation across a decade for two different forest types. The four indices are defined in the legend of Fig. 3. The analyses were performed using three models (null, simple regression, polynomial regression). *adj. R^2*, adjusted coefficient of determination. AIC, Akaike's information criterion. Fig. 3 presents plots of the trends analyzed in part A of this table.

Table S4. Results of simple linear regression analyses of other calculated indices, which were not included in the main analyses, on the change rates of annual mean temperature and annual precipitation.

Table S5. The values (mean, standard deviation, and range) of calculated indices for deciduous broad-leaved forest and evergreen broad-leaved forest in the present study.

Table S6. Results of simple linear regression analyses of the calculated indices using latitude and the means and SDs of annual mean temperature and annual precipitation

Table S7. Indices calculated in this study

Figure S1. Annual mean temperature and annual precipitation in each year (2003–2018) at each site. The red circles and lines indicate the annual mean temperature; the blue bars indicate annual precipitation. The dashed line indicates the average annual mean temperature (1981–2010). The dot-dash line indicates the average of annual precipitation (1981–2010).

Supplementary text 1. Explanation of individual-based temporal beta-diversity in the Ružička dissimilarity index version.

Supplementary file 1. R software script of individual-based temporal beta-diversity calculations using sample data