Degrees of compositional equilibrium in tree communities vary along a gradient

of temperature change rates over one decade: Application of an individual-based

temporal beta-diversity concept

Running title: Individual-based temporal beta diversity

Ryosuke Nakadai*[1][2]

[1] Department of Environmental and Biological Sciences, Faculty of Science and

Forestry, University of Eastern Finland, Yliopistokatu 7, 80101 Joensuu, Finland

[2] Department of Ecosystem Studies, Graduate School of Agricultural and Life Sciences,

The University of Tokyo,1-1-1 Yayoi, Bunkyo-ku, Tokyo 113-8657, Japan

* Correspondence: R. Nakadai

E-mail: r.nakadai66@gmai.com

Type of article: Letters

Number of words in the abstract: 148

Number of words in the main text: 4952

Number of references: 69

Number of figures: 4

Number of tables: 1

Number of text boxes: 0

Keywords: climate change, compositional equilibrium vs. compositional shift,

individual-based beta-diversity, individual-tracked monitoring data, mortality,

recruitment

Author contribution

R.N. conceived the study, designed the methods described herein, conducted the

analyses using empirical data, and wrote the manuscript.

Acknowledgments

I thank Drs. Anu Valtonen and Shunsuke Matsuoka who made insightful comments

on an early version of the manuscript and Dr. Takaya Iwasaki who gave advice on the

dataset for visualization by map. I also thank my friend for inspiring me to reconsider

the importance and the preciousness of individual life in both society and ecology.

Financial support was provided by the Japan Society for the Promotion of Science

(grant 18J00093).

Data accessibility statement

All the data used in the present study were obtained from open sources. The forest

monitoring data were extracted from the website of the Biodiversity Center of Japan

(http://www.biodic.go.jp/moni1000/index.html, in Japanese: accessed on November

11, 2019) published in part by Ishihara *et al*. (2011). Climate data are from the

Japanese Automated Meteorological Data Acquisition System (AMeDAS) database

(http://www.jma.go.jp/jma/indexe.html, Japan Meteorological Agency accessed

December 12, 2019). Taxonomic information was obtained from the Ylist (Yonekura

& Kajita 2003: http://ylist.info/ylist_simple_sea rch.html, in Japanese: accessed

December 8, 2019).

Abstract

Temporal patterns in communities have gained much attention recently. However, the

persistence of identical individuals and turnover among other individuals within the

same species over time have not been considered. To deal with this issue, I propose

new index concepts for beta-diversity and the relative speed of community

equilibrium in relation to individual turnover based on individual identity information

within communities. I applied this new approach to individually tracked tree

monitoring data in deciduous and evergreen broad-leaved forests across the Japanese

archipelago. A new index explored the relative contributions of mortality and

recruitment processes. Clear patterns emerged showing that an increase in the

temperature change rate facilitated the relative contribution of mortality components.

The relative speed of community equilibrium decreased with increasing temperature

change rates in deciduous forests but increased in evergreen forests. The new

concepts provide a way to identify novel and high-resolution temporal patterns in

communities.

Introduction

All living organisms have finite life spans. Each individual tries to leave offspring

before its death. The persistence of individuals and their reproduction in the past

shape the current diversity patterns through temporal dynamics of populations,

communities, and ecosystems. The effects of recent anthropogenically induced

climate change on local and global biodiversity have received considerable attention

(Permesan & Yohe 2003, Chen *et al*. 2011). Specifically, intensive efforts have

focused on biodiversity loss (i.e., alpha- and gamma-diversity) and compositional

shifts (i.e., beta-diversity) within the context of biodiversity conservation (e.g.,

Dornelas *et al*. 2014, Valtonen *et al*. 2016, Hillebrand *et al*. 2017). Ideally, long-term

biodiversity monitoring programs that span multiple decades across replicated study

sites using consistent methodologies and sampling efforts should be used to identify

the effects of climate change. However, multi-decade monitoring programs are as yet

very rare (Magurran *et al*. 2010; but see Valtonen *et al*. 2016).

Whittaker (1960, 1972) developed the concept of beta-diversity, which he defined

as the variation in community composition among sites in a region. Since then, many

new beta-diversity indices have been developed, and attempts have been made to

relate the properties of beta-diversity and dissimilarity indices to background

ecological processes (Harrison *et al*. 1992, Williams 1996, Lennon *et al*. 2001,

Tuomisto & Ruokolainen 2006, Baselga 2010, Legendre 2010; reviewed by Koleff *et*

al. 2003, Anderson *et al*. 2011, Jost *et al*. 2011, Legendre & De Cáceres 2013). More

attention has been focused recently on temporal changes in community composition

in either single sites or in a series of sites surveyed repeatedly over time (Magurran

2011, Matsuoka *et al*. 2016, Legendre & Condit 2019). Temporal changes in

community composition are now termed "temporal beta-diversity" (Hatosy *et al*.

2013, Legendre & Gauthier 2014, Shimadzu *et al*. 2015). The patterns of decay in

temporal distance, which involve increasing community dissimilarity from one time

point to the next in the same location, have been considered within the context of

patterns of decay in geographical distance (Hatosy *et al*. 2013, Bahram *et al*. 2015,

Matsuoka *et al*. 2016). Legendre (2019) elegantly partitioned temporal beta-diversity

into loss and gain components of the compositional changes in each community,

thereby facilitating mechanistic understanding of the temporal changes in community

composition. Furthermore, recent studies have started to show that the temporal

beta-diversity patterns differ among sites (Legendre & Condit 2019, Magurran *et al*.

2019). Environmental changes, including anthropogenically induced shifts, such as

the present worldwide climate warming, are likely factors that drive spatial

differences in temporal beta-diversity (Legendre 2019).

The persistence of individuals over time is an important component of the temporal

beta-diversity concept, which has not been explicitly considered before. The

persistence of individuals contributes to a static equilibrium in community

composition. Nevertheless, there is turnover of different individuals belonging to the

same species even when the community composition is stable over time, i.e.,

conventional temporal beta-diversity is zero when the community composition is at

equilibrium, which contributes to a dynamic equilibrium in community composition.

However, it is difficult to detect differences in turnover within the same species. Thus,

the meaning of compositional changes in communities differs between sites with high

and low individual turnover rates across time. Therefore, an appropriate evaluation of

temporal changes in community composition could benefit from considering the fates

of identifiable individuals in a community over time. Here, I propose new indices of

temporal beta-diversity that take into account individual identity information. These

new coefficients are individual-based in contrast to others that are incidence-based

(e.g., the Sørensen index; Sørensen 1948), or abundance-based (e.g., the Bray–Curtis

index; Bray & Curtis 1957). My new indices quantitatively evaluate the temporal

changes in community composition by considering individual identity information

and reveal the relative contribution of different background processes to these

temporal changes. I also propose a new index to quantify the relative speed of change

away from compositional equilibrium in a community relative to the speed of

individual turnover. I calculated the indices for 16 forest inventory datasets compiled

across the Japanese archipelago. These datasets have individual identity information

for two time periods. The indices identified the effects of climate changes on the

composition of tree communities on a decadal time scale.

Materials and Methods

New indices of temporal beta-diversity and new statistics to evaluate the relative

speed compositional equilibrium in relation to individual turnover

Two types of dissimilarity indices have been proposed in previous studies:

incidence-based and abundance-based (Baselga *et al*. 2013) types. Specifically, the

widely used Bray–Curtis dissimilarity index (Bray & Curtis 1957, which was

proposed earlier as a proportional difference index by Odum 1950), is an

abundance-based extension of the Sørensen index (Legendre & Legendre 1998,

Baselga *et al*. 2013). I have excluded details of incidence-based indices that are

tangential to the main topic in my study (see Anderson *et al*. 2011, Jost *et al*. 2011,

Baselga *et al*. 2013). The conceptual basis for conventional beta-diversity indices and

my new procedures relies on the fact that the intersection (A component) and the

relative complement (B and C components) for species abundance can be formulated

as follows:

$$A = \sum_i \min\left(x_{ij}, x_{ik}\right) \qquad\qquad (1)$$

$$B = \sum_i x_{ij} - \min\left(x_{ij}, x_{ik}\right) \qquad\qquad (2)$$

$$C = \sum_i x_{ik} - \min\left(x_{ij}, x_{ik}\right) \qquad\qquad (3)$$

where x_{ij} is the abundance of species i at site j, and x_{ik} is the abundance of species i at

site k. Therefore, A is the number of individuals of each species that occur at both

sites j and k, while B and C are the numbers of individuals that are unique to sites j

and k, respectively. This formulation of the Bray–Curtis (d_{BC}) index is expressed as

follows:

$$d_{BC} = \frac{B + C}{2A + B + C} .$$ (4)

In the temporal beta-diversity concept, the differences between data vectors

correspond to observations made at time 1 (abbreviated T1) and time 2 (T2), which

are referred to sites k and j in the spatial beta-diversity concept. The A, B, and C

components are described as follows:

$$A_t = \sum_i \min(y_{i1}, y_{i2})$$ (5)

$$B_t = \sum_i y_{i1} - \min(y_{i1}, y_{i2})$$ (6)

$$C_t = \sum_i y_{i2} - \min(y_{i1}, y_{i2})$$ (7)

where y_{i1} is the abundance of species i at time T1 at a site, and y_{i2} is the abundance of

species i at time T2 at the site. Therefore, A_t is the number of individuals of each

species that occur at both times T1 and T2, whereas B_t and C_t are the numbers of

individuals that are unique to times T1 and T2, respectively. The temporal

beta-diversity of the Bray–Curtis ($d_{t.BC}$) index is expressed as follows:

$$d_{t.BC} = \frac{B_t + C_t}{2A_t + B_t + C_t} .$$ (8)

Using Equation 8, Legendre (2019) partitioned the temporal beta-diversity indices

into loss and gain components because a change through time is directional; species

abundances would identify losses and/or gains between T1 and T2 as follows:

$$d_{t.BC} = \frac{B_t + C_t}{2A_t + B_t + C_t}$$

$$= \frac{B_t}{2A_t + B_t + C_t} + \frac{C_t}{2A_t + B_t + C_t} = d_{t.loss} + d_{t.gain} \ . \qquad (9)$$

Legendre and his collaborators applied the loss/gain partitioning procedure for

temporal beta-diversity to many empirical datasets (e.g., Kuczynski *et al.* 2018, Brice

et al. 2019), and constructed a new analytical trend in temporal beta-diversity.

In this study, I developed the concept of abundance-based temporal beta-diversity

based on Bray–Curtis dissimilarity through extension of the individual-based

temporal beta-diversity indices. The following P, M, and R components correspond to

components A, B, and C, respectively. M refers to individuals occurring at T1, but

dead before T2, thereby separately identifying the deaths of individuals during the

time period between T1 and T2 (Fig. 1). R identifies individuals that were not alive at

T1, but subsequently found at T2, thereby identifying the recruitment of individuals

during the period between T1 and T2. Component P refers to the persistence of

individuals from T1 to T2 (Fig. 1). The respective abundances were calculated as

follows:

$$P = \sum_l \min(z_{l1}, z_{l2}) \qquad\qquad (10)$$

$$M = \sum_l z_{l1} - \min(z_{l1}, z_{l2}) \qquad\qquad (11)$$

$$R = \sum_l z_{l2} - \min(z_{l1}, z_{l2}) \qquad\qquad (12)$$

where z_{l1} is individual l at time T1, and z_{l2} is individual l at time T2. Both z_{l1} and z_{l2}

take only 1 or 0, i.e., presence or absence. Therefore, *P* is the number of individuals

present at both T1 and T2, whereas M and R are the numbers of individuals that are

unique to T1 or T2, respectively. Although P, M, and R refer to components A, B, and

C, respectively, here I have used different characters to emphasize the meaning of the

processes of each component. This formulation of the individual-based temporal

beta-diversity index (d_{MR}) can be expressed as follows:

$$d_{MR} = \frac{M + R}{2P + M + R} \, . \tag{13}$$

Furthermore, this individual-based temporal beta-diversity can be partitioned into

the relative contributions of mortality (d_M) and recruitment (d_R) processes using a

similar procedure for the loss and gain component-partitioning of the temporal

beta-diversity in Equation 9 (Legendre 2019) as follows:

$$d_{MR} = \frac{M + R}{2P + M + R}$$

$$= \frac{M}{2P + M + R} + \frac{R}{2P + M + R} = d_M + d_R \, . \tag{14}$$

Traditionally, the equilibrium and non-equilibrium elements in community ecology

indicate properties of the whole community that are directly related to species

coexistence (Pickett 1980). However, some individuals replace others, and thus

ecological communities are always dynamic and vary at least by some degree in

space and time (Mori *et al.* 2018, Ryo *et al.* 2019). In addition, some individuals of

long-lived species (e.g., trees and vertebrates) are persistent over the term, while

individuals of short-lived species are replaced by conspecifics or different species.

Therefore, it is possible to categorize the components of community dynamics into

persistence (P), and components that change over time, i.e., mortality and recruitment

part (M and R), as explained above. The elements of change are also separated into

two component parts of compositional equilibrium (E) and shift (B_t and C_t).

Specifically, component M can be partitioned into lost individuals that contribute to

compositional change in a community (B_t) and lost individuals that are replaced by

conspecifics, contributing substantially to equilibrium in community composition

(E_{loss}). Similarly, component R can be partitioned into gained individuals that

contribute to compositional change in a community (C_t), and gained individuals

replacing conspecifics, thereby contributing substantially to equilibrium in

community composition (E_{gain}). By definition, the numbers of E_{loss} and E_{gain}

individuals are identical; hence, I replaced both E_{loss} and E_{gain} with the coefficient E,

the component that contributes to dynamic compositional equilibrium. The

formulation is as follows:

$$d_{MR} = \frac{M + R}{2P + M + R}$$

$$= \frac{B_t + E_{loss} + C_t + E_{gain}}{2P + M + R} = \frac{B_t + C_t + 2E}{2P + M + R}$$

$$= \frac{2E}{2P + M + R} + \frac{B_t + C_t}{2P + M + R} = d_E + d_S \qquad (15)$$

where d_E and d_S indicate the contributions of equilibrium and shift to

individual-based temporal beta-diversity.

Both M and R in equation 14 refer to total numbers of individuals, but original data

for mortality and recruitment processes of individual species can also be analyzed in

the same manner as community data (Fig. 1). Therefore, the calculation of

Bray–Curtis dissimilarity between communities using mortality and recruitment is

possible, as follows:

$$v_s = \frac{B_t + C_t}{2E + B + C_t}$$

$$= 1 - \frac{2E}{2E + B_t + C_t} = 1 - \frac{2E}{M + R} = 1 - v_e \, . \qquad (16)$$

v_s indicates the speed of compositional shifts in a community relative to the total

speed of individual turnover through mortality and recruitment processes. I used v_e

$= 2E/(M+R)$ for analyses of an empirical dataset because the meaning is intuitively

clearer. If v_e is 1, the individual turnover would not contribute to compositional shifts

in a community, i.e., the community composition of mortality and recruitment are

identical and an equilibrium state exists. If v_e is 0, all individual turnovers contribute

to compositional shifts, and the community composition of mortality and recruitment

are completely different. In my main analyses, I focused only on an extension of

Bray–Curtis dissimilarity. I also summarized explanations of individual-based

beta-diversity using the Ružička dissimilarity index (Ružička 1958), which was

identified by Podani *et al*. (2013) and Legendre (2014) as another appropriate index

for beta-diversity studies. The Ružička dissimilarity index is also an abundance-based

extension of the Jaccard dissimilarity index (Jaccard 1901).

Individual-tracked monitoring data for forests

I predicted the indices of individual-based beta-diversity and the relative speed of

compositional equilibrium variation along degrees of climate change rate per year,

i.e., average speed of local climate change. I tested the performance of the new

indices using individual-tracked monitoring data for forests in Japan. Two types of

temporal change in tree communities may result from climate change—the change

due to changes in the obligate speed of individual turnover, e.g., climate warming

facilitates the faster growth and shorter life span of each individual tree as a

physiological process within individuals (Körner 2017, Munné-Bosch 2018), and the

change due to the relative speed of compositional change, e.g., simple compositional

shifts caused by climate change as an ecological process among individuals. Those

can be referred to as d_{MR} and v_e, respectively.

The Japanese archipelago comprises a long chain of continental islands lying off

the eastern coast of Asia; it is recognized as a hotspot of biodiversity (Mittermeier *et*

al. 2011). The mean temperatures of the coldest months range from −18.4 to 22.3°C,

mean temperatures of the warmest months from 6.2 to 29.2°C, and the annual

precipitation ranges from 700.4 to 4477.2 mm (Japan Meteorological Agency,

1981–2010; http://www.jma.go.jp/jma/indexe.html). Typical forest biomes are

classified into four types: alpine meadow/creeping pine zone, subalpine coniferous

forest zone, deciduous broad-leaved forest zone, and evergreen broad-leaved forest

zone. The empirical tree census dataset used in this study was obtained from the

forest plots used in the Monitoring Sites 1000 Project; this project was launched by

the Ministry of the Environment, Japan (see Ishihara *et al*. 2011 for details). The

monitoring sites include five types of forest: deciduous broad-leaved, evergreen

broad-leaved, mixed broad-leaved and coniferous, evergreen coniferous, and artificial

forest (Ishihara *et al*., 2011, Suzuki *et al*. 2015). The full dataset is available on the

website   of   the   Biodiversity   Center   of   Japan

(http://www.biodic.go.jp/moni1000/index.html, in Japanese: accessed on November

11, 2019), and a subset of the data has been published as a data article (Ishihara *et al*.

2011).

To test the performance of the individual-based temporal beta-diversity approach, I

chose 16 forest plots: nine plots of deciduous broad-leaved forest and seven plots of

evergreen broad-leaved forest (Fig. 2, Table S1) using the following criteria: species

level identifications had been performed in each plot, the plots had been surveyed one

decade after a previous survey (in the period 2005–2008 to 2015–2018), individual

identity information was recorded throughout the two time periods, plot size was 1.0

ha square (100 × 100 m), and sufficient numbers of plots in various environments

within the forest types were available for use in analyses. In this analysis (which was

based on the contents of the original datasets) I chose tree individuals with stem

girths at breast height (GBH) ≥ 15.7 cm, which was approximately equivalent to a

diameter at breast height (DBH) ≥ 5 cm; these were considered target individuals in

the communities. I re-compiled the data at the level of individual trees before the

analyses; the original forest datasets had been compiled at the stem level. I used

taxonomic information provided in the Ylist (Yonekura and Kajita 2003). The data I

analyzed included 18,749 individuals belonging to 254 species found in the 16 plots

(Fig. 2).

Climate variables

To obtain meteorological observation data and their changes over a decade, I

searched for the meteorological station closest to each of the forest plots in the

Japanese Automated Meteorological Data Acquisition System database (Japan

Meteorological Agency; see details in Table S1, Fig. S1). Specifically, I extracted the

monthly mean and total precipitation data (accessed December 12, 2019), and

calculated the annual mean temperature (°C) and annual precipitation (mm) for each

year and site. I estimated temperature differences from elevation differences between

each of the forest plots and its nearest meteorological station using 0.55°C/100 m in

elevation as the mean lapse rate; this relationship between temperature and elevation

is commonly used by Japanese forestry researchers (e.g., Oshima *et al.* 1961,

Fukushima *et al.* 2009). I used simple linear model in R software (R Development

Core Team 2019) to calculate regression coefficients between years and annual

climate (annual mean temperature and annual precipitation) for the targeted years (i.e.,

from the year of the first survey to the year of the second survey) for each plot. The

correlation coefficients indicate the average temperature change per year in the

targeted decade, and hence average change rate in annual mean temperature and

annual precipitation in each plot. The average rates of change in annual mean

temperature (°C) and annual precipitation (mm) were 0.004±0.047 (−0.105 to 0.063)

and 5.714±32.795 (−37.082 to 77.791), respectively, in deciduous broad-leaved

forests, and 0.003±0.043 (−0.040 to 0.059) and 20.860±18.737 (−0.176 to 47.261),

respectively, in evergreen broad-leaved forests (mean ± SD [range]).

Macroecological patterns of indices describing individual-based temporal

beta-diversity and the relative speed of compositional equilibrium

I focused mainly on the following indices: d_{MR}, d_M/d_R, d_E/d_S, and v_e. d_{MR} indicates

total individual turnover between two time periods (i.e., individual-based temporal

beta-diversity). Both d_M/d_R and d_E/d_S are indicators of the relative contribution to

individual-based temporal beta-diversity (d_{MR}); they are rates of mortality to

recruitment processes, and compositional equilibrium to shift, respectively. v_e is the

relative speed of compositional equilibrium relative to total individual turnover. The

relative contributions of compositional equilibrium and shift are formulated as

follows:

$$d_E \div d_S = \frac{2E}{2P + M + R} \div \frac{B_t + C_t}{2P + M + R} = \frac{2E}{B_t + C_t} = \frac{2E}{M + R - 2E} \quad (17)$$

Because "$2E$" in the denominator is simply the difference between v_e (equation 16)

and d_E/d_S (equation 17), the two are strongly correlated (Table S6), but the two

indices have different meaning.

First, I analyzed the relationships between these indices and the average rates of

change in annual mean temperature and annual precipitation using both simple and

polynomial regressions. Then I compared the Akaike's information criterion (AIC)

values (Akaike 1976) among models, including the null model, and selected the best

model for each. I did not use multiple regressions for further analyses because the

number of data points available was insufficient.

To test the relationship between compositional equilibrium and structural change in

communities over time, I also calculated the change in community evenness for each

plot over a decade using Pielou's equation $J = H/Ln(S)$, where H is the Shannon

diversity and S is total species richness in a community; the analyses were executed

with the "vegan" package in R software. I checked the trends of change in

community evenness using the procedures applied in the analyses of temporal trends

in indices d_{MR}, d_M/d_R, d_E/d_S, and v_e (described above). I examined the relationships

between compositional equilibrium (v_e) and structural change (J_{change}). All statistical

analyses were conducted in R.

Results

The values of d_{MR}, d_M/d_R, d_E/d_S, and v_e were 0.154±0.056 (0.084–0.242), 1.211±0.749

(0.415–2.460), 1.025±0.779 (0.134–2.909), and 0.455 ±0.167 (0.119–0.7449),

respectively, in deciduous broad-leaved forests, and 0.137±0.037 (0.102–0.198),

1.211±0.352 (0.541–1.733), 1.153±0.168 (0.978–1.412), and 0.533±0.036

(0.495–0.585), respectively, in evergreen broad-leaved forests.

I detected no trends in d_{MR} in relation to average rate of change in annual mean

temperature in either deciduous or evergreen broad-leaved forests (Fig. 3a, b, Table 1).

In analyses of d_M/d_R, the polynomial model was chosen by the AIC comparison as the

best model for deciduous broad-leaved forests; the simple model was the best for

evergreen broad-leaved forests. The increase in d_M/d_R was greatest at sites that

warmed most in the targeted decades. This was true for both types of forest. Thus, the

relative contribution of mortality process to individual based beta-diversity increased

with warming rate over time (Fig. 3c, d). The d_E/d_S index did not show any statistical

trends over time for deciduous forests (Fig. 3e); the simple regression model was

chosen by AIC comparison as the best for evergreen forests (Table 1). In evergreen

forests, d_E/d_S increased most over a decade in warming sites. Thus, the relative

contribution to individual based beta-diversity of individual turnover increased over

time in relation to the contribution with compositional shift (Fig. 3f). The polynomial

model of v_e was selected as the best for the deciduous forest type: v_e decreased with

larger temperature changes (Fig. 3g, Table 1). The simple model was selected for the

evergreen forest type: v_e increased linearly with increasing warming over time (Fig.

3h, Table 1).

The values of the J_{change} were −0.008±0.033 (−0.091 to 0.019) in deciduous

broad-leaved forests, and 0.010±0.022 (−0.025 to 0.041) in evergreen broad-leaved

forests. J_{change} decreased most at sites with larger temperature change rates in both

forest types (Fig. 3i, j). The statistical analyses of J_{change} changes in relation to the

average change rate in annual mean temperature are summarized in Table 1. I

detected a significant positive correlation for the relationship between these variables

in the deciduous forest type, (Fig. 4a), but not in the evergreen forest type (Fig. 4b).

The average rate of change in annual precipitation did not affect trends over time

for any of the calculated indices (Table S3).

Discussion

Quantitative analyses of biological community change over time are required to

expand our understanding of global climate change and its relationship to rapid losses

in biodiversity. The novel concepts and procedures that I have developed

quantitatively evaluated the relative speed of compositional equilibrium and the

relative contribution of mortality and recruitment processes in tree communities

along a gradient in temperature change rates across a decade. Compositional shifts in

tree communities are generally recognized as long-term phenomena in the absence of

disturbance. Therefore, 10 years is a relatively short-time period in which to analyze

compositional changes in tree communities comprising species with life spans

extending from decades to centuries (Loehle 1988). I found that community

responses to local climate change trends differed between forest types, but

nevertheless, there were real changes in community composition over only a decade.

The changes were caused by change in annual mean temperatures, and not by

changes in annual precipitation across the Japanese archipelago.

The effects of climate changes caused by human activity have been intensively

studied in many disciplines, e.g., analyses of species interactions, range shifts,

phenology, and ecosystem functioning (Thomas *et al*. 2004, Kordas *et al*. 2011,

Valtonen *et al*. 2014). However, studies of temporal changes in diversity and

composition in local communities have often produced mixed and/or different results

among sites and researchers (Vellend *et al*. 2013, Dornelas *et al*. 2014, Newbold *et al*.

2015, reviewed in Hisano *et al*. 2018). I used a novel index to evaluate compositional

changes, i.e., the relative speed of the compositional equilibrium in communities (v_e),

which clearly changed along the gradient of temperature change rates (Fig. 4, Table

1). Furthermore, the responses were different between deciduous and evergreen

broad-leaved forests. Specifically, the relative speed in evergreen forests was greater

at sites with most temperature warming. Furthermore, the range size of the values (v_e)

in deciduous forests was about 7-fold greater than the range size in evergreen forests.

Thus, deciduous forests are more affected by the differences in temperature change

rate. I did not detect statistically significant trends along a gradient of precipitation

change rate, although drought caused by climate change is a major factor in tree

mortality (Allen *et al*. 2010, Saiki *et al*. 2017). These contrasting results may be due

to the humid climate of the Japanese archipelago. To capture the effects of global

climate change accurately, a broad multi-site analysis across the globe using similar

methods will be required.

The association between the relative speed of compositional equilibrium and

structural changes in communities over time is an important ecological issue. I found

that in deciduous forests the relative speed was significantly and positively correlated

with changes in community evenness, but this was not the case for evergreen forests

(Fig. 4). Therefore, compositional changes were not always accompanied by

structural change (Fig. 4). I examined the patterns of community evenness in

mortality and recruitment communities (J_M and J_R respectively) to develop a

mechanistic interpretation of the temporal change in community evenness in

deciduous forests (Table S5). In detail, the evenness of the mortality communities

increased with an increasing rate of temperature change; thus, mortality processes

were a major contributor of the change in community evenness, but recruitment

communities did not track a significant trend (Table S4). Only d_R responded

positively to the warming rate, but no clear patterns were apparent for this index

(Table S4). The major response components differed between the changes in

composition and structures of communities. Although more supporting evidence is

needed, I therefore suggest that there might not be a direct relationship between

changes in community structure and composition over time.

Predicting community persistence from snapshot data on community structure is a

complicated issue for community ecologists. For example, Ulrich *et al.* (2018) argued

that it may be possible to evaluate species persistence and their relative proportions

from the patterns of species abundance distribution. If this were the case, the relative

speed of compositional equilibrium should provide a comparable estimate of

community persistence from both individual and species-based perspectives.

Specifically, using some time series datasets containing substantial annual individual

information, it should be technically possible to calculate the retention time for each

community, i.e., the lifespan durations of all individuals replaced in a focal

community. These calculations could be cross-compared to with snapshot community

structure data (e.g., species abundance patterns). In the present study, I used data for

only two time periods, so this cross-comparison was not possible. Nevertheless,

clarification of the relationship between changes in community structure and

composition should become possible in the near future. Underlying mechanisms

explaining the relationship may then be discernable.

One advantage of the methodology I have developed is that it dissects individual

turnover into mortality and recruitment components. For example, I detected no clear

patterns in the relationship between individual turnover (d_{MR}), which is the obligate

speed of temporal change in communities, and temperature change rate (Fig. 3).

However, by splitting the index into two components, I was able to identify relative

contributions to individual turnover (d_M/d_R). When this was done, clear patterns

emerged showing that an increase in the temperature change rate facilitated the

relative contribution of mortality components (Fig. 3, Table 1). Furthermore, by

checking the trend of each component separately, I was able to confirm that decreases

in recruitment components with increases in the temperature change rate drove the

pattern in deciduous forests, but I was not able to confirm this in evergreen forests

(Table S5).

The relative contribution of the index d_E/d_S to individual-based temporal

beta-diversity (d_{MR}) increased positively in evergreen forests as the warming rate

increased, although no trends in the individual components (d_E, d_S) were discerned.

Thus, the balance between the components changed in a consistent manner with

temperature change, although no patterns of variation in the individual components

were found. There were no clear trends in d_E/d_S in relation to temperature change

rates in deciduous forests. However, d_E was affected by the temperature change rate

in these forests: d_E decreased with increased warming rates; thus, there was a clear

contribution of compositional equilibrium decrease along these gradients (Table S5).

Compilation of long-term temporal datasets for macro species requires

considerable human effort and cost for monitoring and research surveys because large

organisms often have long life spans. Nevertheless, the new indices I developed

clearly enhanced the importance of the monitoring datasets assembled using

traditional detailed surveys (e.g., individual tracking in forest plots) in detecting the

temporal trends that occurred over short time periods. Therefore, I re-emphasize here

the significance of classical monitoring surveys to collect biodiversity information. I

also developed an approach that introduced the concept of individual-based

beta-diversity within the context of temporal changes in community composition.

Theoretically, however, the procedure could be expanded to a spatial context for use

in studies of animal behavioral ranges and movements, including long-distance

migration, using monitoring and biologging datasets (Viana *et al.* 2016, Börger *et al.*

2020) that include multiple time series datasets collected from the same place within

a shorter time frame than the life spans of the targeted organisms (e.g., bird ringing

and classical mark–recapture procedures). Specifically, the approach could detect

detailed spatial turnover for individuals within same species in communities.

Many ecology theories have been developed from the concept of species. The

original concept of beta-diversity was proposed to reflect species turnover (Whittaker

1960). By contrast, much recent research has pointed out the problems of

species-based studies and the importance of considering individuals in community

ecology (Dupont *et al.* 2011, Nakazawa 2017, Tatsumi *et al.* 2019). Species are the

consequences of ecological/evolutionary processes among individuals; to be accurate,

the species themselves do not drive these processes, i.e., individuals do by competing

with other individuals, and this does not involve the species directly. Although

individuals were considered in the unified neutral theory (Hubbel 2001) and

subsequent related studies (Hubert *et al.* 2015, Vanoverbeke *et al.* 2016, Tatsumi *et al.*

2019), the focus was restricted to the patterns of species richness, abundance, and

related diversity indices. The effects of individual births and deaths on biodiversity

patterns have not yet been considered sufficiently. Procedures for incorporating

individual identity information into biodiversity evaluation, i.e., individual-based

diversity indices, are required to facilitate further conceptual advances in ecology.

New ecological concepts will emerge from combined considerations of species

abundance and information relevant to individual identity in the future.

Conclusions

Previous studies have often detected temporal changes in local communities, likely as

a result of human activities, but not at all times and places. In this study, I clearly

showed the responses of tree communities to local and short-term climate change

trends using datasets collected according to novel index concepts. Notably, the

relative speed of compositional equilibrium in tree communities varied along a

gradient of temperature change rates in this decade. This community level response

resulted from individual sorting in mortality and recruitment processes. Therefore,

accumulating more detailed information across the globe, specifically individual

identity information at the local community level collected over time, is required for

understanding both the actual changes due to global climate change in biodiversity

and the background processes mechanistically. Mortality and recruitment rates are

generally difficult to measure in empirical studies compared to loss and gain

information based only on snapshot data, especially except in the case of plants or

vertebrates. The individual-based approach may not be tractable for all taxonomic

groups. However, information on individual organisms will help to bridge the

disciplines of macroecology and study fields focusing on within-individual

phenomena, for example, physiological ecology. The calculation of individual-based

diversity indices from individual identity monitoring data will support efforts to

quantitatively evaluate the impacts of climate change on future biodiversity. The

further development of novel diversity index concepts that make use of

individual-based biodiversity information to provide a more detailed and mechanistic

understanding of biodiversity in general is needed.

References

Akaike, H. (1976). An information criteron (AIC). *Math. Sci.*, 14, 5-7.

Allen, C.D., Macalady, A.K., Chenchouni, H., Bachelet, D., McDowell, N., Vennetier,

M., *et al.* (2010). A global overview of drought and heat-induced tree mortality

reveals emerging climate change risks for forests. *For. Ecol. Manag.*, 259,

660–684.

Anderson, M. J., Crist, T. O., Chase, J. M., Vellend, M., Inouye, B. D., Freestone, A.

L., *et al*. (2011). Navigating the multiple meanings of β diversity: a roadmap for

the practicing ecologist. *Ecol. Lett.,* 14, 19-28.

Bahram, M., Peay, K. G., & Tedersoo, L. (2015). Local–scale biogeography and

spatiotemporal variability in communities of mycorrhizal fungi. *New. Phytol.*,

205, 1454-1463.

Baselga, A. (2010). Partitioning the turnover and nestedness components of beta

diversity. *Glob. Ecol. Biogeogr.*, 19, 134-143.

Baselga, A. (2013). Separating the two components of abundance–based

dissimilarity: balanced changes in abundance vs. abundance gradients. *Methods*

Ecol. Evol., 4, 552-557.

Börger, L., Bijleveld, A., Fayet, A., Machovsky-Capuska, G., Patrick, S., Street, G.

M., & Vander Wal, E. (2020). Biologging special feature. *J. Anim. Ecol.,* 13

January 2020 https://doi.org/10.1111/1365-2656.13163

Bray, J.R., & Curtis, J.T. (1957). An ordination of the upland forest communities of

southern Wisconsin. *Ecol. Monogr.*, 27, 325-349.

Brice, M.H., Cazelles, K., Legendre, P., & Fortin, M.J. (2019). Disturbances amplify

tree community responses to climate change in the temperate–boreal ecotone.

Glob. Ecol. Biogeogr., 28, 1668-1681.

Chen, I. C., Hill, J. K., Ohlemüller, R., Roy, D. B., & Thomas, C. D. (2011). Rapid

range shifts of species associated with high levels of climate warming. *Science*,

333, 1024-1026.

Dornelas, M., Gotelli, N. J., McGill, B., Shimadzu, H., Moyes, F., Sievers, C., &

Magurran, A. E. (2014). Assemblage time series reveal biodiversity change but

not systematic loss. *Science*, 344, 296-299.

Dupont, Y.L., Trøjelsgaard, K., & Olesen, J.M. (2011). Scaling down from species to

individuals: a flower–visitation network between individual honeybees and

thistle plants. *Oikos*, 120, 170-177.

Fukushima K, Tokuchi N, Tateno R, Katsuyama M (2009) Water yield and nitrogen

loss during regrowth of Japanese cedar forests after clearcutting. In: Taniguchi

M, Fukushima Y, Burnett WC, Haigh M, Umezawa Y (eds) From headwaters to

the ocean: hydrological change and watershed management. Taylor & Francis,

London, pp. 97–104

Harrington, R., Woiwod, I., & Sparks, T. (1999). Climate change and trophic

interactions. *Trends Ecol. Evol.*, 14, 146-150.

Harrison, S., Ross, S.J., & Lawton, J.H. (1992). Beta diversity on geographic

gradients in Britain. *J. Anim. Ecol.*, 61, 151-158.

Hatosy, S.M., Martiny, J.B., Sachdeva, R., Steele, J., Fuhrman, J.A., & Martiny, A. C.

(2013). Beta diversity of marine bacteria depends on temporal scale. *Ecology*, 94,

1898-1904.

Hillebrand, H., Blasius, B., Borer, E.T., Chase, J.M., Downing, J.A., Eriksson,

B.K.,*et al*. (2018). Biodiversity change is uncoupled from species richness

trends: consequences for conservation and monitoring. *J. Appl. Ecol.*, 55,

169-184.

Hisano, M., Searle, E. B., & Chen, H. Y. (2018). Biodiversity as a solution to mitigate

climate change impacts on the functioning of forest ecosystems. *Biol. Rev.*, 93,

439-456.

Hubbell, S. P. (2001). The unified neutral theory of biodiversity and biogeography.

Princeton University Press.

Hubert, N., Calcagno, V., Etienne, R. S., & Mouquet, N. (2015). Metacommunity

speciation models and their implications for diversification theory. *Ecol. lett.*, 18,

864-881.

611 Ishihara, M. I., Suzuki, S. N., Nakamura, M., Enoki, T., Fujiwara, A., Hiura, T., *et al*.

612   (2011). Forest stand structure, composition, and dynamics in 34 sites over Japan.

613   *Ecol. Res*., 26, 1007.

614 Jaccard, P. (1901). Distribution de la flore alpine dans le bassin des Dranses et dans

615   quelques régions voisines. *Bull. Soc. Vaudoise. Sci. Nat.*, 37, 241-272.

616 Jost, L., Chao, A. & Chazdon, R. (2011) Compositional similarity and beta diversity.

617   *Biological Diversity: Frontiers in Measurement and Assessment* (eds A.

618   Magurran & B.J. McGill), pp. 66–84. Oxford University Press, Oxford.

619 Koleff, P., Gaston, K. J., & Lennon, J. J. (2003). Measuring beta diversity for

620   presence–absence data. *J. Anim. Ecol.*, 72, 367-382.

621 Kordas, R. L., Harley, C. D., & O'Connor, M. I. (2011). Community ecology in a

622   warming world: the influence of temperature on interspecific interactions in

623   marine systems. *J. Exp. Mar. Biol. Ecol.*, 400, 218-226.

624 Körner, C. (2017). A matter of tree longevity. *Science,* 355, 130-131.

625 Kuczynski, L., Legendre, P., & Grenouillet, G. (2018). Concomitant impacts of

626   climate change, fragmentation and non‐native species have led to

627   reorganization of fish communities since the 1980s. *Glob. Ecol. Biogeogr.*, 27,

628   213-222.

629 Legendre, P., & Condit, R. (2019). Spatial and temporal analysis of beta diversity in

630   the Barro Colorado Island forest dynamics plot, Panama. *For. Ecosyst.*, 6, 7.

631 Legendre, P., & De Cáceres, M. (2013). Beta diversity as the variance of community

632   data: dissimilarity coefficients and partitioning. *Ecol. lett*, 16, 951-963.

633 Legendre, P., & Gauthier, O. (2014). Statistical methods for temporal and space–time

634   analysis of community composition data. *Proc. Roy. Soc. B.*, 281, 20132728.

635 Legendre, P. & Legendre, L. (1998) Numerical Ecology, 2ndedn. Elsevier,

636   Amsterdam.

637 Legendre, P. (2014). Interpreting the replacement and richness difference components

of beta diversity. *Glob. Ecol. Biogeogr.*, 23, 1324-1334.

Legendre, P. (2019). A temporal beta-diversity index to identify sites that have

changed in exceptional ways in space‑time surveys. *Ecol. Evol.*, 9, 3500-3514.

Lennon, J. J., Koleff, P., GreenwooD, J. J., & Gaston, K. J. (2001). The geographical

structure of British bird distributions: diversity, spatial turnover and scale. *J.*

Anim. Ecol., 70, 966-979.

Loehle, C. (1988). Tree life history strategies: the role of defenses. *Can. J. For. Res.*,

18, 209-222.

Magurran, A. E. (2011). Measuring biological diversity in time (and space).

Biological Diversity: Frontiers in Measurement and Assessment (eds A.

Magurran & B.J. McGill), pp. 85-104. Oxford University Press, Oxford.

Magurran, A. E., Baillie, S. R., Buckland, S. T., Dick, J. M., Elston, D. A., Scott, E.

M., *.et al*. (2010). Long-term datasets in biodiversity research and monitoring:

assessing change in ecological communities through time. *Trends Ecol. Evol.*, 25,

574-582.

Magurran, A. E., Dornelas, M., Moyes, F., & Henderson, P. A. (2019). Temporal β

diversity–A macroecological perspective. *Glob. Ecol. Biogeogr.*, 28, 1949-1960.

Matsuoka, S., Kawaguchi, E., & Osono, T. (2016). Temporal distance decay of

similarity of ectomycorrhizal fungal community composition in a subtropical

evergreen forest in Japan. *FEMS Microbiol. Ecol.*, 92.

Mittermeier R.A., Turner W.R., Larsen F.W., Brooks T.M., & Gascon C. (2011).

Global biodiversity conservation: the critical role of hotspots. In: Zachos FE,

Habel JC (eds) Biodiversity hotspots. Springer, Berlin, pp 3–22

Mori, A. S., Isbell, F., & Seidl, R. (2018). β-diversity, community assembly, and

ecosystem functioning. *Trends Ecol. Evol.*, 33, 549-564.

Munné-Bosch, S. (2018). Limits to tree growth and longevity. *Trends Plant Sci.*, 23,

985-993.

Nakazawa, T. (2017). Individual interaction data are required in community ecology:

a conceptual review of the predator–prey mass ratio and more. *Ecol. Res.*, 32,

5-12.

Newbold, T., Hudson, L.N., Hill, S.L., Contu, S., Lysenko, I., Senior, R.A., *et al*.

(2015). Global effects of land use on local terrestrial biodiversity. *Nature*, 520,

45.

Odum, E.P. (1950). Bird populations of the Highlands (North Carolina) Plateau in

relation to plant succession and avian invasion. *Ecology*, 31, 587–605.

Oshima, Y. (1961). Ecological studies of *Sasa* communities. I. Productive structure of

some of the *Sasa* communities in Japan. *Bot. Mag. Tokyo*, 74, 199-210.

Parmesan, C., & Yohe, G. (2003). A globally coherent fingerprint of climate change

impacts across natural systems. *Nature*, 421, 37.

Pickett, S. T. (1980). Non-equilibrium coexistence of plants. *Bull. Torrey. Bot. Club.*,

238-248.

Podani, J., Ricotta, C., & Schmera, D. (2013). A general framework for analyzing

beta diversity, nestedness and related community–level phenomena based on

abundance data. *Ecol. Complex.*, 15, 52–61.

R Development Core Team (2019). R: a language and environment for statistical

computing. R Foundation for Statistical Computing, Vienna, Austria.

http://www.R-project.org/.

Ružička, M. (1958). Anwendung mathematisch–statisticher Methoden in der

Geobotanik (synthetische Bearbeitung von Aufnahmen). *Biologia, Bratislava*,

13, 647–661.

Ryo, M., Aguilar-Trigueros, C. A., Pinek, L., Muller, L. A., & Rillig, M. C. (2019).

Basic principles of temporal dynamics. *Trends Ecol. Evol.*, 34, 723-733

Saiki, S. T., Ishida, A., Yoshimura, K., & Yazaki, K. (2017). Physiological

mechanisms of drought-induced tree die-off in relation to carbon, hydraulic and

692   respiratory stress in a drought–tolerant woody plant. *Sci. Rep.*, 7, 2995.

693 Shimadzu, H., Dornelas, M., & Magurran, A. E. (2015). Measuring temporal turnover

694   in ecological communities. *Methods Ecol. Evol.*, 6, 1384-1394.

695 Sørensen, T. A. (1948). A method of establishing groups of equal amplitude in plant

696   sociology based on similarity of species content and its application to analyses

697   of the vegetation on Danish commons. *Biol. Skar.*, 5, 1-34.

698 Suzuki, S. N., Ishihara, M. I., & Hidaka, A. (2015). Regional‐scale directional

699   changes in abundance of tree species along a temperature gradient in Japan.

700   *Glob. Change Biol.*, 21(9), 3436-3444.

701 Tatsumi, S., Cadotte, M.W., & Mori, A.S. (2019). Individual–based models of

702   community assembly: Neighbourhood competition drives phylogenetic

703   community structure. *J. Ecol.*, 107, 735-746.

704 Thomas, C. D., Cameron, A., Green, R. E., Bakkenes, M., Beaumont, L. J.,

705   Collingham, Y. C., *et al*. (2004). Extinction risk from climate change. *Nature*,

706   427, 145.

707 Tuomisto, H., & Ruokolainen, K. (2006). Analyzing or explaining beta diversity?

708   Understanding the targets of different methods of analysis. *Ecology*, 87,

709   2697–2708.

710 Ulrich, W., Nakadai, R., Matthews, T. J., & Kubota, Y. (2018). The two-parameter

711   Weibull distribution as a universal tool to model the variation in species relative

712   abundances. *Ecol. Complex.*, 36, 110-116.

713 Valtonen, A., Hirka, A., Szőcs, L., Ayres, M.P., Roininen, H., & Csóka, G. (2017).

714   Long–term species loss and homogenization of moth communities in Central

715   Europe. *J. Anim. Ecol.*, 86, 730-738.

716 Valtonen, A., Leinonen, R., Pöyry, J., Roininen, H., Tuomela, J., & Ayres, M. P.

717   (2014). Is climate warming more consequential towards poles? The phenology

718   of Lepidoptera in Finland. *Glob. Change Biol.*, 20, 16-27.

Vanoverbeke, J., Urban, M. C., & De Meester, L. (2016). Community assembly is a

race between immigration and adaptation: eco‐evolutionary interactions across

spatial scales. *Ecography*, 39, 858-870.

Vellend, M., Baeten, L., Myers-Smith, I. H., Elmendorf, S. C., Beauséjour, R., Brown,

C. D., *et al*. (2013). Global meta-analysis reveals no net change in local-scale

plant biodiversity over time. *PNAS*, 110, 19456-19459.

Viana, D. S., Santamaría, L., & Figuerola, J. (2016). Migratory birds as global

dispersal vectors. *Trends Ecol. Evol*, 31, 763-775.

Whittaker, R. H. (1960). Vegetation of the Siskiyou mountains, Oregon and

California. *Ecol. Monogr.*, 30(3), 279-338.

Whittaker, R. H. (1972). Evolution and measurement of species diversity. *Taxon*, 21,

213-251.

Williams, P. H. (1996). Mapping variations in the strength and breadth of

biogeographic transition zones using species turnover. *Proc. Roy. Soc. B.*,

263(1370), 579-588.

Yonekura K, Kajita T (2003) BG Plants Ylist: an online service of Japanese plant

names, including a nomenclature index (in Japanese).

(http://ylist.info/ylist_simple_search.html) accessed on 8[th] Dec 2019

Legends

Figure 1 Schematic representation of individual-based temporal beta-diversity. T1

and T2 refer to the communities at times 1 (abbreviated T1) and 2 (T2), respectively.

Both communities comprise four species (sp. 1, 2, 3, and 4) and all individuals are

identified separately (Zi). M refers to individuals in existence at T1 but dead before

T2, thereby separately identifying the deaths of individuals during the period between

T1 and T2. R identifies individuals that were not alive at T1, but found at T2, thereby

identifying the recruitment of individuals during the period between T1 and T2. P

refers to individuals that persisted from T1 to T2, thus both T1 and T2 are 1. The

arrows indicate the temporal trajectory for each individual; specifically, when they

are present or absent in communities between T1 and T2.

Figure 2 Locations of the study sites within the Japanese archipelago. The crosses

and asterisks indicate the locations of targeted plots in deciduous broad-leaved forest

(DB) and evergreen broad-leaved forest (EB) included in the present study. The

lettering adjacent to each symbol is the plot identification code (ID). Detailed plot

information is listed in Table S2. The colors indicate four major vegetation types:

purple, Alpine meadow or creeping pine zone; blue, subalpine coniferous forest zone;

red, deciduous broad-leaved forest zone; green, evergreen broad-leaved forest zone.

The information on vegetation type at a 1.0 km grid scale was obtained from the

results of the fifth censuses (1992–1996) of the National Survey of the Natural

Environment in Japan (http://www.biodic.go.jp/dload/mesh_vg.html; in Japanese,

accessed December 12, 2019).

Figure 3 Trends in tree community indices across a gradient of temperature change

rates over a decade. The left column of panels refers to trends in deciduous

broad-leaved forests; the right column of panels refers to trends in evergreen

broad-leaved forests. Panels (a) and (b) show the trends in individual-based temporal

beta-diversity indices (d_{MR}). Panels (c) and (d) show trends in the d_M/d_R index, which

is an indicator of the relative contribution of mortality and recruitment processes to

individual-based temporal beta-diversity (d_{MR}). Panels (e) and (f) show trends in the

d_E/d_S index, which is an indicator of the relative contribution of equilibrium and shift

processes to individual-based temporal beta-diversity (d_{MR}). Panels (g) and (h) show

trends in the index v_e, which is an indicator of the relative speed of compositional

equilibrium. Panels (i) and (j) show trends in the index J_{change}, which is an indicator

of community evenness. The solid black lines indicate statistically significant ($P <$

0.05) trends. The dashed and the dot-dashed black lines indicate marginally

significant ($0.05 < P < 0.1$, $0.1 < P < 0.15$) trends respectively. The blue dashed lines

enclose the 95% confidence interval envelopes. Detailed results of statistical analyses

are presented in Table 1.

Figure 4 Relationships between the change in community evenness (J_{change}) and the

relative speed of compositional equilibrium (v_e) in deciduous forests (a) (coefficients

= 3.190, $adj.R^2 = 0.314$, $P = 0.067$), and in evergreen forests (b) (coefficients = 0.716,

$adj.R^2 = 0.037$, $P = 0.318$).

 Figure 1

787   **Figure 2**

Figure 3

Average change rate of annual mean temperture (°C/year)

792   Figure 4

Table 1 Statistical analyses of the relationships between the indices d_{MR}, d_M/d_R, d_E/d_S, v_e, and J_{change} and the temperature change rate across a decade for two different types of forest. The four indices are defined in the legend to Fig. 3. The analyses were performed using three models (null, simple regression, polynomial regression). *adj. R^2*, adjusted coefficient of determination. AIC, Akaike's information criterion. Fig. 3 presents plots of the trends analyzed in this table.

		Temperature change rate		Squared temperature change rate					
		Coefficient	*P*	Coefficient	*P*	*adj. R^2*	*P*		AIC
(a) Deciduous broad-leaved forest									
d_{MR}	Null	–	–	–	–	–	–		–23.503 †
	Simple	–0.408	0.361	–	–	–0.006	0.361		–22.654
	Polynomial	–0.620	0.344	–4.119	0.625	–0.124	0.598		–21.044
d_M/d_R	Null	–	–	–	–	–	–		23.267
	Simple	10.577	0.049 *	–	–	0.366	0.049 *	19.958
	Polynomial	17.099	0.018 *	127.068	0.119	0.523	0.046 *	18.018 †
d_E/d_S	Null	–	–	–	–	–	–		23.990 †
	Simple	–2.601	0.685	–	–	–0.114	0.685		25.763
	Polynomial	–8.456	0.351	–114.075	0.343	–0.105	0.570		26.302

v_e	Null	–	–		–	–		–	–		–3.700	
	Simple	–1.461	0.270		–	–		0.052	0.270		–3.378	
	Polynomial	–3.513	0.036	*	–39.994	0.061	.	0.413	0.085	.	–7.083	†
J_{change}	Null	–	–		–	–		–	–		–32.832	
	Simple	–0.197	0.464		–	–		–0.053	0.464		–31.572	
	Polynomial	–0.729	0.010	*	–10.354	0.008	**	0.658	0.017	**	–41.082	†

(b) Evergreen broad–leaved forest

d_{MR}	Null	–	–		–	–		–	–		–23.200	†
	Simple	0.024	0.953		–	–		–0.199	0.953		–21.206	
	Polynomial	0.081	0.884		–3.666	0.855		–0.485	0.980		–19.272	
d_M/d_R	Null	–	–		–	–		–	–		8.171	
	Simple	5.717	0.079	.	–	–		0.391	0.079	.	5.419	†
	Polynomial	4.762	0.233		61.633	0.642		0.284	0.228		6.990	
d_E/d_S	Null	–	–		–	–		–	–		–2.211	

	Simple	2.902	0.053	.	–	–	0.472	0.053	.	−5.952	†
	Polynomial	2.529	0.171	.	24.065	0.683	0.370	0.177		−4.281	
v_e	Null	–	–	–	–	–	–	–		−23.911	
	Simple	0.612	0.056	.	–	–	0.462	0.056	.	−27.531	†
	Polynomial	0.536	0.175	.	4.918	0.697	0.356	0.184		−25.830	
J_{change}	Null	–	–	–	–	–	–	–		−30.588	
	Simple	0.352	0.087	.	–	–	0.369	0.087	.	−33.085	
	Polynomial	0.0500	0.050	.	−9.567	0.216	0.487	0.117		−34.104	†

***$P < 0.001$; **$P < 0.01$; *$P < 0.05$; '.,' $P < 0.1$

† The best of three models judged by the AIC

Appendices

Table S1 Data on the geography and climate of each plot and its nearest metrological station. See Fig. 2 for locations of plots in the Japanese archipelago.

Table S2 Indices calculated in this study

Table S3 Changes in the calculated individual-based temporal beta-diversity indices and the relative speed of compositional equilibrium in relation to temporal changes in annual precipitation.

Table S4 Results of simple linear regression analyses of the calculated indices on latitude, and the means and SDs of annual mean temperature and annual precipitation

Table S5 Results of simple linear regression analyses of other calculated indices, which were not included in the main analyses, on the change rates of annual mean temperature and annual precipitation

Table S6 Correlations among the geographical and environmental variables and their respective calculated indices.

Figure S1 Annual mean temperature and annual precipitation in each year (2003–2018) at each site. The red circles and lines indicate the annual mean temperature; the blue bars indicate annual precipitation. The dashed line indicates the average annual mean temperature (1981–2010). The dot-dash line indicates the average of annual precipitation (1981–2010).

Supplementary text 1 Explanation of individual-based temporal beta-diversity in the Ružička dissimilarity index version

Supplementary file 1 R software script of individual-based temporal beta-diversity calculations using sample data

	T1	M	R	T2	P
Z_1	1	0	0	1	1
Z_2	1	0	0	1	1
Z_3	1	1	0	0	0
Z_4	0	0	1	1	0
Z_5	0	0	1	1	0
Z_6	1	0	0	1	1
Z_7	1	0	0	1	1
Z_8	1	1	0	0	0
Z_9	1	1	0	0	0
Z_{10}	1	1	0	0	0
Z_{11}	1	1	0	0	0
Z_{12}	1	1	0	0	0
Z_{13}	0	0	1	1	0
Z_{14}	0	0	1	1	0
Z_{15}	0	0	1	1	0
Z_{16}	1	1	0	0	0
Z_{17}	1	1	0	0	0
Z_{18}	1	1	0	0	0
Z_{19}	1	0	0	1	1
Z_{20}	0	0	1	1	0

Groups: sp.1 (Z_1–Z_5), sp.2 (Z_6–Z_{10}), sp.3 (Z_{11}–Z_{15}), sp.4 (Z_{16}–Z_{20})

Plot type

× Deciduous broad−leaved forest (DB)
✳ Evergreen broad−leaved forest (EB)

Vegetation type

■ Alpine meadow/Creeping pine zone
■ Subalpine coniferous forest zone
■ Deciduous broad−leaved forest zone
■ Evergreen broad−leaved forest zone

Average change rate of annual mean temperture (°C/year)