aiMeRA: A generic modular response analysis R package and its application to

estrogen and retinoic acid receptors crosstalk

Gabriel Jimenez-Dominguez[1,2,3], Patrice Ravel[1,2,3], Stéphan Jalaguier[1,2,3], Vincent Cavaillès[1,2,3,*] and

Jacques Colinge[1,2,3,*]

[1]Inserm U1194, Institut de Recherche en Cancérologie de Montpellier, Montpellier, France

[2]University of Montpellier, Montpellier, France

[3]ICM, Institut régional du Cancer de Montpellier, Montpellier, France

*Corresponding authors

E-mails: vincent.cavailles@inserm.fr, jacques.colinge@inserm.fr

Abstract

Modular response analysis (MRA) is a widely used modeling technique to uncover coupling strengths in molecular networks under a steady-state condition by means of perturbation experiments. We propose an extension of this methodology to search genomic data for new associations with a network modeled by MRA and to improve the predictive accuracy of MRA models. These extensions are illustrated by exploring the cross talk between estrogen and retinoic acid receptors, two nuclear receptors implicated in several hormone-driven cancers such as breast. We also present a novel, rigorous and elegant mathematical derivation of MRA equations, which is the foundation of this work and of an R package that is freely available at https://github.com/bioinfo-ircm/aiMeRA/. This mathematical analysis should facilitate MRA understanding by newcomers.

Author summary

Estrogen and retinoic acid receptors play an important role in several hormone-driven cancers and share co-regulators and co-repressors that modulate their transcription factor activity. The literature shows evidence for crosstalk between these two receptors and suggests that spatial competition on the promoters could be a mechanism. We used MRA to explore the possibility that key co-repressors, i.e., NRIP1 (RIP140) and LCoR could also mediate crosstalk by exploiting new quantitative (qPCR) and RNA sequencing data. The transcription factor role of the receptors and the availability of genome-wide data enabled us to explore extensions of the MRA methodology to explore genome-wide data sets *a posteriori*, searching for genes associated with a molecular network that was sampled by perturbation experiments. Despite nearly two decades of use, we felt that MRA lacked a systematic mathematical derivation. We present here an elegant and rather simple analysis that should greatly facilitate

42   newcomers' understanding of MRA details. Moreover, an easy-to-use R package is released that should

43   make MRA accessible to biology labs without mathematical expertise. Quantitative data are embedded

44   in the R package and RNA sequencing data are available from GEO.

Introduction

Modular response analysis (MRA) was introduced to infer the coupling between components of a

biological system in a steady-state [1]. It can be applied to components at different levels of details, e.g.,

individual genes or subsystems such as pathways or processes. It relies on the perturbation of individual

components, the so-called modules. Various developments of MRA and related methods were recently

reviewed [2] but, despite its success, MRA mathematical derivation was not provided in a systematic

and rigorous manner. We thus reasoned that such an analysis was needed and it would facilitate the

understanding of the methodology for newcomers. It is presented as a result and is the basis of the

development of an open source R library (aiMeRA) that should make MRA accessible to biology labs

without mathematical expertise. We illustrate the use of the aiMeRA package by investigating the

crosstalk between nuclear receptors (NRs) in a breast cancer (BC) cell line. A new extension of the

method is also introduced to perform inferences at the genomic scale. The Blüthgen Lab recently

released another R package to perform MRA computations [3], although with a specific focus on their

particular edge-pruning and associated maximum likelihood extension of MRA [4] that is not our interest

in this study.

Estrogen receptors (ERs) belong to the NR superfamily, which act as transcription factors activated upon

ligand binding. The two isoforms of ERs (ERα and ERβ) are involved in the control of cell proliferation

and exhibit essential functions in tissue development and homeostasis, in particular in organs related to

reproduction [5]. ERα overexpression is frequently observed in breast, ovarian, endometrial, and other

hormone-driven tumors. The transcriptional activity of ERs is modulated by several coregulatory

complexes including coactivators and corepressors [5]. In the presence of estrogens or any agonist

ligand, ERs interact preferentially with coactivators, or with a specific subclass of corepressors including

nuclear receptor-interacting protein 1 (NRIP1 or RIP140) and Ligand-dependent corepressor (LCoR).

NRIP1 is a corepressor of particular interest because its expression is directly induced by estrogen, i.e.,

NRIP1 installs a negative feedback loop to keep ER signaling under control [6]. NRIP1 abnormal

expression is indeed observed in ER-driven tumors [7,8]. LCoR represses transcription of estrogen-

induced gene expression [9], and NRIP1 expression was shown to be necessary for LCoR inhibitory

activity in BC cells [10].

Interestingly, NRIP1 and LCoR function as corepressors for several liganded NRs. For instance, LCoR can

repress vitamin D receptor (VDR), retinoic acid receptors (RARs), and RXR ligand-dependent

transcription [9] in addition to ERs. Moreover, NRIP1 is a known direct target and negative regulator of

RAR transcription [11].

There is experimental evidence of crosstalk between ER and RAR signaling [12]. For instance, ERα can

suppress the basal expression of retinoic acid (RA)-responsive gene RARβ2, but also turns out to be

necessary for its RA induction [13]. It was also found that ERα activates RARα1 expression in BC cells

[14]. Other authors intersected RAR targets identified by ChIP-seq with ER binding sites to discover a

significant overlap [15]. This work suggested a space competition mechanism for estrogen and RA

signaling in BC. A potential cooperative interaction between RARα and ER was also shown in BC [16].

Since NRIP1 and LCoR expression can be both regulated by RAR and ER transcription, we can further

hypothesize that these molecules mediate part of ER-RAR crosstalk. The induced expression of NRIP1

and LCoR by one receptor produces molecules able to repress signaling of both receptors subsequently.

We aimed at characterizing the ER-RAR-NRIP1-LCoR network, at the transcriptional level, utilizing

transcript abundance measurements and MRA. Accordingly, we considered a steady-state situation in a

BC (MCF7-derived) cell line that would model BC cells with or without constant estrogenic stimulation.

Perturbation experiments were realized to generate quantitative PCR data unraveling interaction

strengths in the network, i.e., coupling according to MRA principles.

Given the nature of ER and RAR, i.e., transcription factors, and the general ability of MRA to perform

predictions [4], we introduced an extension of the method to perform genome-wide inferences

exploiting mRNA sequencing (RNA-seq) data. For this purpose, we acquired whole transcriptomes under

perturbed conditions identical to those used for qPCR. We first established that MRA could produce

accurate results from RNA-seq data. Next, we asked whether the ER-RAR-NRIP1-LCoR network inferred

by MRA could predict the mRNA abundance of estrogen-targeted genes better than a trivial model. This

extension of MRA, where one or several modules do not experience perturbations, was called

unidirectional to underline the implied absence of potential influence on the other modules.

Results

Mathematical derivation

MRA original paper [1] introduced the concept of modeling interdependencies (coupling) within a

biological system modularly. That is, subsystems involving molecules and their relationship at a detailed

level, which would not be the interest of the study, could be captured as a single module with one

measurable quantity defining the overall module activity. For instance, in the case of ERα signaling, it is

possible to represent the complex process of ligand-binding and transcriptional activity by a single

module (Fig. 1). The activity of this module is measured by a reporter gene, which is the luciferase in

MELN cells. NRIP1 and LCoR activities were determined by their respective mRNA abundances. We ask

the question of the direct dependence of each module activity with respect to the other modules

activity. That is, we want to compute (signed) weights to put on the directed edges of Fig. 1. The answer

is searched in a steady-state through successive elementary perturbations of each module activity.

Depending on the application, this framework can be applied to different molecular species and

processes, e.g., protein or metabolite concentrations, protein phosphorylation levels, etc. [2].

Fig. 1. MRA general principle. Illustrated with the ERα-NRIP1-LCoR transcriptional network. Each

module's activity level is given by a measured reporter. Coupling (edge weights) is determined from

perturbation experiments.

Now, in full generality, we assume that there are n modules whose activities are given by $x \in \mathbb{R}^n$. We

further admit the existence of n intrinsic parameters, $p \in \mathbb{R}^n$, one *per* module, which are perturbed by

the elementary perturbations. One can imagine mRNA abundance parameters for perturbations such as

siNRIP1 or siLCoR, and numbers of available ERα-E2 bound complexes for the E2 perturbation. In other

circumstances, perturbations could change affinity constants or other physical characteristics. Finally,

we assume that there exist $f: \mathbb{R}^n \times \mathbb{R}^n \longrightarrow \mathbb{R}^n$ of class $\mathcal{C}^1$ (continuously differentiable) such that

$$\dot{x} = f(x, p). \tag{1}$$

We do not need to know $f(x,p) = (f_1(x,p), \cdots, f_n(x,p))^t$ explicitly, but we need one more hypothesis

that is the existence of a time $T > 0$ such that all the solutions we consider for any p and initial

conditions of x, have reached a steady-state, i.e.,

$$\dot{x} = 0, \forall t > T.$$

The unperturbed, basal state of the modules is denoted $x^0 \in \mathbb{R}^n$ and it has corresponding parameters

$p^0 \in \mathbb{R}^n$. According to our hypotheses, $f(x^0, p^0) = 0 \Leftrightarrow f_i(x^0, p^0) = 0, \forall i \in \{1, \cdots, n\}$. By the implicit

function theorem, $\forall i$, there exists open neighborhoods $V_i \times W_i \subset \mathbb{R}^{n-1} \times \mathbb{R}^n$ of

$(x_1^0, \cdots, x_{i-1}^0, x_{i+1}^0, \cdots, x_n^0, p_1^0, \cdots, p_n^0)$, $U_i \subset \mathbb{R}$ of x_i^0, and $g_i : V_i \times W_i \rightarrow U_i$ (also $\mathcal{C}^1$) with

$$f_i\left(x_1^0, \cdots, x_{i-1}^0, g_i(\cdots), x_{i+1}^0, \cdots, x_n^0, p_1^0, \cdots, p_n^0\right) = 0. \tag{2}$$

We denote $x(p^0 + \Delta p)$, the steady-state corresponding to the changed parameters $p^0 + \Delta p$. We

introduce the notation $x_{j \neq i}$ to denote all the x_j but x_i. Now, if we assume that $(x_{j \neq i}(p^0 + \Delta p), p^0 +$

$\Delta p)$ belong to $V_i \times W_i$ for all the perturbations considered experimentally, then by Taylor's Formula

$$x_i(p^0 + \Delta p) = x_i(p^0) + g_i'(p^0) \begin{pmatrix} x_1(p^0 + \Delta p) - x_1(p^0) \\ \vdots \\ x_{i-1}(p^0 + \Delta p) - x_{i-1}(p^0) \\ x_{i+1}(p^0 + \Delta p) - x_{i+1}(p^0) \\ \vdots \\ x_n(p^0 + \Delta p) - x_n(p^0) \\ \Delta p_1 \\ \vdots \\ \Delta p_n \end{pmatrix} + o(\|\Delta p\|). \tag{3}$$

Dividing each side by $x_i(p^0)$, Eq. (3) can be rewritten

$$\frac{x_i(p^0 + \Delta p) - x_i(p^0)}{x_i(p^0)} =$$

$$\sum_{j=1, j \neq i}^{n} \frac{x_j(p^0)}{x_i(p^0)} \frac{\partial g_i}{\partial x_j}(p^0) \left(\frac{x_j(p^0 + \Delta p) - x_j(p^0)}{x_j(p^0)}\right)$$

147
$$+ \frac{1}{x_i(p^0)} \sum_{j=1}^{n} \frac{\partial g_i}{\partial p_j}(p^0)(\Delta p_j) + \mathcal{O}(\|\Delta p\|). \tag{4}$$

149   Since parameter p_j influences module j only, $\frac{\partial g_i}{\partial p_j} = 0$ if $j \neq i$. Moreover, $g_i(x_{j\neq i}, p) = x_i(x_{j\neq i}, p)$ in

150    $V_i \times W_i$, and if we denote

151
$$\frac{\Delta x_i}{x_i} = \frac{x_i(p^0 + \Delta p) - x_i(p^0)}{x_i(p^0)},$$

152   and

153
$$r_{i,j} = \frac{x_j(p^0)}{x_i(p^0)} \frac{\partial x_i}{\partial x_j}(p^0), j \neq i, \tag{5}$$

154   then

155
$$\frac{\Delta x_i}{x_i} = \sum_{j\neq i} r_{i,j}\left(\frac{\Delta x_j}{x_j}\right) + \frac{1}{x_i(p^0)} \frac{\partial x_i}{\partial p_i}(p^0)(\Delta p_i) + \mathcal{O}(\|\Delta p\|). \tag{6}$$

156   We next consider elementary perturbations q_k, $k \in \{1, \cdots, n\}$, which only perturb module k, i.e., the

157   parameter p_k. Neglecting the second-order term $\mathcal{O}(\|\Delta p\|)$ and writing

158
$$\left(\frac{\Delta x_i}{x_i}\right)_{q_k}$$

159   the relative difference in module i activity upon Δp_k change induced by perturbation q_k, we find

160
$$\left(\frac{\Delta x_i}{x_i}\right)_{q_k} = \sum_{j\neq i} r_{i,j}\left(\frac{\Delta x_j}{x_j}\right)_{q_k}, k \neq i, \tag{7}$$

161
$$\left(\frac{\Delta x_i}{x_i}\right)_{q_i} = \sum_{j\neq i} r_{i,j}\left(\frac{\Delta x_j}{x_j}\right)_{q_k} + \frac{\partial x_i}{\partial p_i}(p^0)\left(\frac{\Delta p_i}{x_i}\right). \tag{8}$$

162   By defining $r_{i,i} = -1$, we can write Eqs (7) and (8) in matrix form:

163
$$rR = -P, \tag{9}$$

164   where R is the matrix that contains the experimentally observed relative activity changes $R_{j,k} =$

165      $\left(\frac{\Delta x_j}{x_j}\right)_{q_k}$, $j,k \in \{1,\cdots,n\}$. P is a diagonal matrix with $P_{i,i} = \frac{\partial x_i}{\partial p_i}(p^0)\left(\frac{\Delta p_i}{x_i}\right)$, $i \in \{1,\cdots,n\}$. The system (9)

166   can be solved in two steps [1]. Firstly, $r = -PR^{-1}$ and because $r_{i,i} = -1$, we have $P_{i,i}(R^{-1})_{i,i} = 1$ and

167   thus $P_{i,i} = \frac{1}{(R^{-1})_{i,i}}$. Secondly,

168
$$r = -[\text{diag}(R^{-1})]^{-1}R^{-1}.$$

169   The elements of R are defined by $\left(\frac{x_i(p^0+\Delta p_k)-x_i(p^0)}{x_i(p^0)}\right)_{q_k}$ but as previously suggested [1], we preferred to

170   estimate this quantity by

171
$$R_{j,k} = 2\left(\frac{x_i(p^0+\Delta p_k)-x_i(p^0)}{x_i(p^0+\Delta p_k)+x_i(p^0)}\right)_{q_k}, \tag{10}$$

which avoids divisions by 0 and is numerically more stable.

Finally, from Eq. (5), we see that $r_{i,j}$ contains the searched coupling between MRA modules: direct

action of j on i normalized by the ratio x_j/x_i. Similarly, $P_{i,i}$ measures the relative effect of q_i on x_i. We

call it q_i magnitude. The implicit function theorem provides analytical expressions for $g_i{'}$ in terms of f

partial derivatives, but since f is generally unknown, we did not use them. In his seminal work,

Kholodenko made additional hypotheses to show that r contains coupling information, which is not

necessary with our derivation. To be rigorous, one should ultimately restrict the model to

neighborhoods included in all the V_i's, W_i's, and U_i's.

MRA models have been largely used for their inference capabilities [4]. Let us define a multiple

perturbation q to be the linear combination of elementary perturbations q_k. For instance, a

perturbation on modules i and j with the same individual magnitudes would be coded by a column

vector c with 1's at positions i and j and 0's elsewhere. From Eq. (9), we compute

184
$$\left(\frac{\Delta x}{x}\right)_q = -r^{-1}Pc,$$
(11)

185 with $\left(\frac{\Delta x}{x}\right)_q$ the column vector containing the inferred relative changes on each module activity.

186 Denoting Δp the parameter changes induced by q, individual module activities are given by

187
$$\left(\frac{\Delta x_i}{x_i}\right)_q = -(r^{-1}Pc)_i = 2\frac{x_i(p^0 + \Delta p) - x_i(p^0)}{x_i(p^0 + \Delta p) + x_i(p^0)}$$

188
$$\Longleftrightarrow \quad x_i(p^0 + \Delta p) = \frac{x_i(p^0)}{1+\frac{2}{(r^{-1}Pc)_i}}\left(\frac{2}{(r^{-1}Pc)_i} - 1\right).$$
(12)

In case elementary perturbations contribute for different amounts to q, the vector c contains q_k's

relative weights. In every case, linearity between the perturbation strength and its impact on p is

assumed.

Confidence intervals (CI) around model parameters are estimated by a bootstrap procedure [17]. We

considered experimental designs with biological and technical replicates. We average the technical

replicates to obtain one activity value *per* biological replicate. The latter are again averaged to compute

the R matrix according to Eq. (10). From the biological replicates, we estimate the variance of each x_i

employing an estimator optimized for a small sample size from Statistical Process Control theory [18,19].

Finally, a Gaussian distribution is assumed and 10^6 R matrices are generated, which are submitted to

MRA computations. The 95% CI is obtained from the 2.5[th] and 97.5[th] percentiles. In case 0 is not included

in the CI, the MRA parameter is deemed significant and marked by an asterisk in the figures.

Inferences obtained from Eq. (12) were also complemented by the estimation of CIs following the

principles above.

Transcriptional data

ERβ and RARβ expression could not be quantified in MELN cells. We hence learned networks involving

an ERα module, which transcriptional activity was reported by the ERE/luciferase construct. That is,

luciferase mRNA abundance measured ERα activity. ERα mRNA abundance would combine ligand-bound

and free amounts of the receptor, but only the ligand-bound ones matter in the model. We did not try

to distinguish between RARα and RARγ. We estimated their combined transcriptional activity by the

mRNA abundance of the *HOXA5* gene and the corresponding MRA module was named RARs. NRIP1 and

LCoR activity were determined by their gene mRNA abundance. Since MELN cells are BC cells, we

considered the E2-, RA-, or E2+RA-stimulated conditions as basal. That is, perturbations at ERα and RARs

were negative (switch to ethanol). Perturbations at NRIP1 and LCoR were achieved by siRNAs, i.e., they

were also negative.

The ERα-NRIP1-LCoR network

In an unstimulated condition (no E2), it is known that NRIP1 expression induces LCoR expression [10].

We started by assessing this coupling under the E2 basal condition and found similar coupling (Fig. 2A).

We also observed negative coupling from LCoR to NRIP1, which is logical since NRIP1 is a direct target of

E2-bound ERα and LCoR one of its corepressor. We next inferred the ERα-NRIP1-LCoR network under E2

(Fig. 2B). We could observe the known induction of NRIP1 by ERα with negative feedback [6]. We also

reconstituted the known inhibition of ERα by LCoR [9]. Interestingly, the induction of LCoR upon NRIP1

expression observed in Fig. 2A became a double inhibition via ERα in Fig. 2B. This makes sense since

there is no transcriptional control by NRIP1 alone, it can only modulate ERα activity. Perturbation

magnitudes are in Fig. 2C. Finally, we assessed the validity of the inferred network by checking its

predictive power. From Fig. 2D, we note a reasonable fidelity of the model and that the relative errors

are commensurate with the CI sizes, i.e., with data variability.

Fig. 2. ER and RAR separated networks. A. Coupling between the two corepressors under E2 condition.

A 95% CI for each model parameter was estimated and the parameter marked by an asterisk provided 0

was not included (nonzero with 5% significance). **B.** The ERα-NRIP1-LCoR transcriptional network. **C.**

ERα, NRIP1, and LCoR perturbation magnitudes. **D.** Inference of gene expression under the dual siNRIP1

and siLCoR perturbation. **E.** Similar to A but under RA condition. **F.** The RARs-NRIP1-LCoR network. **G.**

Similar to C. **H.** Similar to D.

The RARs-NRIP1-LCoR network

Our next endeavor was to build a RARs-NRIP1-LCoR network before switching to the full network with

both NRs. NRIP1 and LCoR coupling under RA stimulation (Fig. 2E) remained similar to its state under E2.

That was expected since these two corepressors are used by several NRs. In Fig. 2F, we reconstituted the

induction of NRIP1 expression by RAR as well as the inhibition of RAR expression by NRIP1 [11]. The

inhibition of RAR by LCoR was also known [9]. Coupling between LCoR and NRIP1 is essentially similar to

Fig. 2B since the NRIP1-to-LCoR arrows featured weak coupling. NRIP1 perturbation magnitude

remained close, but LCoR perturbation changed 2-fold (Fig. 2G) although the same siRNAs were used.

Inferences (Fig.2H) also supported the accuracy of the model.

The full ERα-RARs-NRIP1-LCoR network

We followed the same approach as above to construct a full model of ER-RAR crosstalk (Fig. 3A).

Perturbation magnitudes (Fig. 3B) were in the same range as before under the new dual E2 and RA basal

condition. Values for NRIP1 and LCoR were closer to the RAR-NRIP1-LCoR network. No literature reports

coupling with the corepressors NRIP1 and LCoR under this particular condition. Only the crosstalk

between RAR and ER mentioned in the introduction is known [15,16]. We hence first challenged the

model by testing its predictive accuracy (Fig. 3C), which was again satisfying.

Fig. 3. The ERα-RARs-NRIP1-LCoR network. A. MRA network model. **B.** Perturbation magnitudes under

the dual E2 and RA stimulation. **C.** Inferred activity of the modules upon double siRNA inhibition of

NRIP1 and LCoR.

Interestingly, cross inhibition of ER and RAR signaling acted along two paths. The model shows direct

inhibition of ER transcriptional activity by the RAR module, which was described in the literature [15,16].

Reciprocal inhibition was suggested but not significant in our data. In agreement with our hypothesis,

we found a parallel crosstalk mechanism through the induction of NRIP1 expression, which could

subsequently repress both RARs and ERα. MRA modeling thus supported the coexistence of the two

phenomena. Although LCoR reversed action on NRIP1 compared to the E2 and RA independent

conditions might counterbalance cross inhibition of the two NRs, the strengths of coupling on the model

edges and the much-attenuated induction of LCoR by NRIP1 suggested that it was not the case.

MRA models from RNA-seq data

Since MRA relies on module activity relative change (Eqs. (7-8)), absolute quantitation is not necessary.

We thus computed an RARs-NRIP1-LCoR MRA models as HOXA5, NRIP1, and LCoR mRNA abundances

were available in our RNA-seq data (Fig. 4A). Comparing with the qPCR-based model in Fig. 2F, we notice

that all the significant edges of Fig. 2F plus NRIP1-to-LCoR were recovered. The only change is very weak

coupling LCoR-to-RARs (-0.03) that became slightly positive (0.13) with RNA-seq data. CIs were not

determined for RNA-seq data since only two replicates were available.

Fig. 4. Genomic-scale inferences. A. RARs-NRIP1-LCoR model trained from RNA-seq data. **B.** Eight

closest replacement genes for ERE-Luc in the ERα-NRIP1-LCoR model. **C.** ERα-NRIP1-LCoR models with

ERE-Luc replaced by PGR, trained from qPCR and RNA-seq data. **D.** Principle of unidirectional MRA. **E.**

Accuracy of unidirectional MRA inference (udMRA & udMRA.ab) under the E2 condition with double

siNRIP1/siLCoR perturbation versus simple predictors (mean, geometric mean (gMean), and maximum

of the two siRNAs).

The results above indicated that MRA could be applied to RNA-seq data. We, therefore, decided to

exploit this opportunity by performing a new type of investigation. We used MRA to find a gene that

would function as ERα transcription reporter, and would thus replace the ERE-Luc construct. Existing

ChIP-seq data [20] intersected with our RNA-seq data allowed us to identify 884 genes targeted by ERα

and E2-regulated (edgeR analysis, P-value<0.01, fold-change>2). We hence computed 884 MRA models

with siNRIP1 and siLCoR RNA-seq data, replacing ERE-Luc by each of those genes successively. The genes

with closest Euclidean distances between their model coupling parameters (the $r_{i,j}$ matrix) and those of

the original Fig. 2B qPCR model are listed in Fig. 4B. We decided to test PGR and measured its expression

by qPCR. The qPCR- and RNA-seq-based models are featured in Fig. 4C. They accurately reproduced the

original model of Fig. 2B and were very similar to each other, thereby further validating the use of RNA-

seq data for MRA model training.

Unidirectional MRA on a genome-scale

We reasoned that the ERα-NRIP1-LCoR MRA model might provide means of predicting E2-regulated

gene expression. We hence introduced a modified ERα-NRIP1-LCoR MRA model with one additional

module that cannot influence the other modules (Fig. 4D). The gray unidirectional arrows in Fig. 4D

represent the coupling between NRIP1, LCoR, the ERα module, and the added gene denoted by X. This

coupling can be learned in the E2 basal condition by applying elementary perturbations as above. This

must be repeated for each gene X considered. Gene X mRNA abundance is an $n + 1^{\text{th}}$ module and, by

hypothesis, $r_{i,n+1} = 0, \forall i \in \{1, \cdots, n\}$, since no return arrows exist. From Eq. (7), we can compute

$r_{n+1,j}, \forall j \in \{1, \cdots, n\}$, by solving the system

$$\left(\frac{\Delta x_{n+1}}{x_{n+1}}\right)_{q_k} = \sum_{j=1}^{n} r_{n+1,j} \left(\frac{\Delta x_j}{x_j}\right)_{q_k}, \; k \in \{1, \cdots, n\}.$$

The performance of this new type of MRA model (udMRA) was assessed by its ability to predict module

$n + 1$ activity under the dual siNRIP1/siLCoR condition, as we did above for the other models. To avoid

trivially successful predictions on genes that would not vary, we limited the benchmark to the 884 genes

above that were also significantly regulated upon siNRIP1 or siLCoR in the E2-stimulated condition. That

left us with 60 genes. In Fig. 4E, we report the relative errors observed applying udMRA and comparing

with naïve predictions. udMRA yielded significantly better estimates of the added module activity.

One could wonder whether perturbation magnitudes during double siRNA interference on the same

biological system remain identical. That is, whether filling the vector c in Eq. (11) with 1's at the

perturbed module indices (what we did so far) is the best option. Eq. (11) is written such that we can try

different values. We searched for optimal coefficients a and b applied to siNRIP1 and siLCoR

perturbations (at the corresponding indices in vector c), such that prediction errors of Luciferase, NRIP1,

and LCoR expression (as in Fig. 2D) would be minimal. We found $a = 1$, and $b = 0.4$. Then, we used

those coefficients in the udMRA model to try to predict the expression of the 61 benchmark genes more

accurately. Indeed, we see in Fig. 4E that this new model called udMRA.ab achieved much better

accuracy.

aiMeRA usage

The R package was designed to be generally applicable; it relies on the formulae presented here. It is

able to work with any quantitative input, including biological and technical replicates. We included

functionality to facilitate the definition of MRA model topologies (Fig. 5A). Model construction only

involves the execution of a few generic R functions and network plots can be generated within R directly

(Fig. 5B). It is also possible to export such graphs in the graphML format for loading into Cytoscape [21].

More details are provided in the package documentation. aiMeRA is available from GitHub; submission

to Bioconductor is pending.

Fig. 5. The aiMeRA R package. A. Example R code to load data, prepare them and compute the model.

Note that NRIP1 was called after its common alternative name RIP140. Basal condition is E2 plus RA and

we see that LCoR perturbation is defined as E2+RA+siLCoR. Same logic for RIP140 (=NRIP1). Perturbation

on the HOXA5 module reporting RARs activity is defined as E2, i.e., loss of RA stimulation. Etc. **B.** Plot of

an MRA model in R using the igraph library.

Discussion

MRA modeling [1,2] is a widely used technique to learn the coupling between the modules of a

biological system from perturbation data sets (Fig. 1). We introduced a new mathematical derivation of

the method and implemented a generic R package called aiMeRA (Fig. 5). Application thereof was

illustrated on an unpublished data set combining specific qPCR and broad RNA-seq data to explore

crosstalk between ER and RAR, two important NRs involved in several tumors such as BC. Data analysis

reconstituted some known interactions (Fig. 2) and supported a novel hypothesis that reciprocal

negative coupling could be mediated by shared corepressors (Fig. 3).

We showed that MRA transcriptional models trained from RNA-seq data are close to those trained from

qPCR (Fig. 4A-C). Which lead us to introduce an innovative application of MRA to probe genes genome-

wide, searching for replacement reporters of module activity. That is, new genes that could be

functionally related to an MRA module. In particular, we found that the progesterone receptor gene

(*PGR*) reported on ligand-bound ERα transcriptional activity accurately. That observation indicates the

potential value of this new use of MRA models since PGR is a widely used reporter of estrogen activity in

BC in the clinic.

We additionally investigated the possibility to build hybrid MRA models (udMRA & udMRA.ab) including

unidirectional coupling to add modules that were not perturbed in the training data set (Fig. 4D). In the

context of the application presented in this report, i.e., the transcriptional activity of a system of

transcription factors, we showed that the ERα-RARs-NRIP1-LCoR udMRA and udMRA.ab models could

outperform naïve predictors (Fig. 4E). Other biological systems might be amenable to such modified

models in the absence of strong back-coupling. The aiMeRA package methods support both RNA-seq

data and udMRA models.

Methods

Cell culture and perturbation experiments

We used MELN cells, an MCF7-derived cell line stably transfected with the estrogen-responsive

luciferase reporter gene ERE-βGlob-Luc-SV-Neo [22]. The cell line was authenticated by short tandem

repeat profiling and tested for mycoplasma contamination.

MELN cells were cultured in phenol red-free Dulbecco's modified Eagle medium (Gibco) containing 5%

dextran-charcoal treated FCS (Invitrogen) and antibiotics (Gibco). Perturbations at NRIP1 and LCoR were

obtained by siRNAs that were transfected using Interferrin (Polyplus). Perturbations at ERα and RARs

were induced by their respective natural ligands: the hormones estrogen (17β-estradiol or E2 for short)

and all-trans retinoic acid (RA).

MELN cells were obtained in the following conditions: basal (untreated), E2, RA, E2+RA, siNRIP1, siLCOR,

siNRIP1+siLCOR, E2+siNRIP1, E2+siLCOR, E2+siNRIP1+siLCOR, RA+siNRIP1, RA+siLCOR,

RA+siNRIP1+siLCOR, E2+RA+siNRIP1, E2+RA+siLCOR, and E2+RA+siNRIP1+siLCOR. These experiments

were realized in triplicates. Cells were harvested after 18 hours of culture. E2-treated cells received

100nM E2, RA-treated cells 10 uM RA, and untreated cells ethanol. Validations of the response to E2 and

siRNA interference are in Suppl. Fig. 1.

mRNA quantification

RNA was isolated using the Zymo Research kit (Zymo Research) and reverse transcription (RT)-qPCR

assays were done using qScript (VWR) according to the manufacturer's protocol. Transcripts were

quantified using SensiFAST SYBR (BioLine) on an LC480 instrument. The nucleotide sequences of the

primers used for real-time PCR were:

RIP140-f (5'- AATGTGCACTTGAGCCATGATG -3'),

RIP140-r (5'- TCGGACACTGGTAAGGCAGG -3'),

LCoR-f (5'- GAACCTAGCGAACAAGACGGTG -3'),

LCoR-r (5'- TGGAGAGTGGCTCAGGGAAGT -3'),

Luciferase-f (5'- CTCACTGAGACTACATCAGC -3'),

Luciferase-r (5'- TCCAGATCCACAACCTTCGC -3'),

HOXA5-f (5'- GCGCAAGCTGCACATAAGTC -3'),

HOXA5-r (5'- GAACTCCTTCTCCAGCTCCA -3'),

ERα-f (5'- TGGAGATCTTCGACATGCTG -3'),

ERα-r (5'- TCCAGAGACTTCAGGGTGCT -3'),

RARα-f (5'- GGATATAGCACACCATCCCC -3'),

RARα-r (5'- TTGTAGATGCGGGGTAGAGG -3'),

PGR-f (5'- CGCGCTCTACCCTGCACTC-3'),

PGR-r (5'-TGAATCCGGCCTCAGGTAGTT-3').

(RT)-qPCR data are available from the R package.

mRNA sequencing

For two of the triplicates, in each condition, RNA was extracted as above described. Libraries were

prepared with Illumina TruSeq kit and submitted to NextSeq500 sequencing (1x75bp/40M reads). The

first 13 and last 7 bps were cut by an in-house Perl script to eliminate compositional bias. Cut reads were

submitted to sickle to eliminate remaining low-quality regions. Alignments were performed against the

human genome (hg38) with TopHat v2.10 [23] and read counts extracted with HTSeq-Count [24]. The

read count matrix was normalized with edgeR [25] TMM algorithm. Data are available from GEO under

GSE143956.

aiMeRA library implementation

We implemented the MRA method according to the mathematical formulation above as an R library.

(RT)-qPCR data of this project were embedded in the R library for convenience and to provide an

example. We also included the data used in the MRA original paper [1] such that users can check that

our code gives the same results as those reported in the latter publication.

Acknowledgments

412   We thank Simon Cabello-Aguilar and Meriem Mekedem for useful discussions during the development of

413   the project. JC was supported by the Agence Nationale de la Recherche (grant

414   ANR-16-CE17-0002-02), the Groupement des entreprises françaises dans la lutte contre le cancer

415   (GEFLUC), and the Fondation ARC (PJA 20151203332).

References

1.   Kholodenko BN, Kiyatkin A, Bruggeman FJ, Sontag E, Westerhoff HV, Hoek JB. Untangling the
wires: a strategy to trace functional interactions in signaling and gene networks. Proc Natl Acad Sci
U S A. 2002;99: 12841–6. doi:10.1073/pnas.192442699

2.   Santra T, Rukhlenko O, Zhernovkov V, Kholodenko BN. Reconstructing static and dynamic models
of signaling pathways using Modular Response Analysis. Current Opinion in Systems Biology.
2018;9: 11–21. doi:10.1016/j.coisb.2018.02.003

3.   Dorel M, Klinger B, Gross T, Sieber A, Prahallad A, Bosdriesz E, et al. Modelling signalling networks
from perturbation data. Bioinformatics. 2018;34: 4079–4086. doi:10.1093/bioinformatics/bty473

4.   Klinger B, Sieber A, Fritsche-Guenther R, Witzel F, Berry L, Schumacher D, et al. Network
quantification of EGFR signaling unveils potential for targeted combination therapy. Mol Syst Biol.
2013;9: 673. doi:10.1038/msb.2013.29

5.   Heldring N, Pike A, Andersson S, Matthews J, Cheng G, Hartman J, et al. Estrogen receptors: how
do they signal and what are their targets. Physiol Rev. 2007;87: 905–931.
doi:10.1152/physrev.00026.2006

6.   Cavaillès V, Dauvois S, L'Horset F, Lopez G, Hoare S, Kushner PJ, et al. Nuclear factor RIP140
modulates transcriptional activation by the estrogen receptor. EMBO J. 1995;14: 3741–3751.

7.   Docquier A, Garcia A, Savatier J, Boulahtouf A, Bonnet S, Bellet V, et al. Negative regulation of
estrogen signaling by ERβ and RIP140 in ovarian cancer cells. Mol Endocrinol. 2013;27: 1429–1441.
doi:10.1210/me.2012-1351

8.   Sixou S, Müller K, Jalaguier S, Kuhn C, Harbeck N, Mayr D, et al. Importance of RIP140 and LCoR
Sub-Cellular Localization for Their Association With Breast Cancer Aggressiveness and Patient
Survival. Transl Oncol. 2018;11: 1090–1096. doi:10.1016/j.tranon.2018.06.006

9.   Fernandes I, Bastien Y, Wai T, Nygard K, Lin R, Cormier O, et al. Ligand-Dependent Nuclear
Receptor Corepressor LCoR Functions by Histone Deacetylase-Dependent and -Independent
Mechanisms. Molecular Cell. 2003;11: 139–150. doi:10.1016/S1097-2765(03)00014-5

10. Jalaguier S, Teyssier C, Nait Achour T, Lucas A, Bonnet S, Rodriguez C, et al. Complex regulation of
LCoR signaling in breast cancer cells. Oncogene. 2017;36: 4790–4801. doi:10.1038/onc.2017.97

11. White KA, Yore MM, Warburton SL, Vaseva AV, Rieder E, Freemantle SJ, et al. Negative Feedback
at the Level of Nuclear Receptor Coregulation SELF-LIMITATION OF RETINOID SIGNALING BY
RIP140. J Biol Chem. 2003;278: 43889–43892. doi:10.1074/jbc.C300374200

12. White KA, Yore MM, Deng D, Spinella MJ. Limiting effects of RIP140 in estrogen signaling: potential
mediation of anti-estrogenic effects of retinoic acid. J Biol Chem. 2005;280: 7829–7835.
doi:10.1074/jbc.M412707200

13. Rousseau C, Pettersson F, Couture MC, Paquin A, Galipeau J, Mader S, et al. The N-terminal of the
estrogen receptor (ERalpha) mediates transcriptional cross-talk with the retinoic acid receptor in
human breast cancer cells. J Steroid Biochem Mol Biol. 2003;86: 1–14. doi:10.1016/s0960-
0760(03)00255-3

14. Laganière J, Deblois G, Giguère V. Functional genomics identifies a mechanism for estrogen
activation of the retinoic acid receptor alpha1 gene in breast cancer cells. Mol Endocrinol. 2005;19:
1584–1592. doi:10.1210/me.2005-0040

15. Hua S, Kittler R, White KP. Genomic antagonism between retinoic acid and estrogen signaling in
breast cancer. Cell. 2009;137: 1259–1271. doi:10.1016/j.cell.2009.04.043

16. Ross-Innes CS, Stark R, Holmes KA, Schmidt D, Spyrou C, Russell R, et al. Cooperative interaction
between retinoic acid receptor-alpha and estrogen receptor in breast cancer. Genes Dev. 2010;24:
171–182. doi:10.1101/gad.552910

17.   Davison AC, Hinkley DV. Bootstrap Methods and their Applications. Cambridge University Press;
464   1997. Available: https://doi.org/10.1017/CBO9780511802843

465   18.   Wheeler DJ, Chambers DS. Understanding Statistical Process Control. 2nd ed. SPC Press, Inc.; 1992.

466   19.   Harter HL. Tables of Range and Studentized Range. Ann Math Statist. 1960;31: 1122–1147.
467   doi:10.1214/aoms/1177705684

468   20.   Madak-Erdogan Z, Charn T-H, Jiang Y, Liu ET, Katzenellenbogen JA, Katzenellenbogen BS.
469   Integrative genomics of gene and metabolic regulation by estrogen receptors α and β, and their
470   coregulators. Mol Syst Biol. 2013;9: 676. doi:10.1038/msb.2013.28

471   21.   Smoot ME, Ono K, Ruscheinski J, Wang PL, Ideker T. Cytoscape 2.8: new features for data
472   integration and network visualization. Bioinformatics. 2011;27: 431–2.
473   doi:10.1093/bioinformatics/btq675

474   22.   Balaguer P, Boussioux AM, Demirpence E, Nicolas JC. Reporter cell lines are useful tools for
475   monitoring biological activity of nuclear receptor ligands. Luminescence. 2001;16: 153–158.
476   doi:10.1002/bio.630

477   23.   Kim D, Pertea G, Trapnell C, Pimentel H, Kelley R, Salzberg SL. TopHat2: accurate alignment of
478   transcriptomes in the presence of insertions, deletions and gene fusions. Genome Biology.
479   2013;14: R36. doi:10.1186/gb-2013-14-4-r36

480   24.   Anders S, Pyl PT, Huber W. HTSeq--a Python framework to work with high-throughput sequencing
481   data. Bioinformatics. 2015;31: 166–169. doi:10.1093/bioinformatics/btu638

482   25.   McCarthy DJ, Chen Y, Smyth GK. Differential expression analysis of multifactor RNA-Seq
483   experiments with respect to biological variation. Nucleic Acids Res. 2012;40: 4288–4297.
484   doi:10.1093/nar/gks042

Supporting information captions

Suppl. Fig. 1. A. MELN cells were transfected with either a siRIP140 (siNRIP1), a siLCoR or a combination

of the two siRNAs. RIP140 and LCoR mRNA levels were quantified by real time PCR. Results are corrected

to 28S mRNA and normalized to cells transfected with the control siRNA. **B.** MELN cells were transfected

as described in A and treated with estradiol (10^{-7} M) when indicated. Luciferase mRNA expression is

quantified as in A.

Figures

Figure 1

D
Basal condition E2, double siNRIP1/siLCOR perturbation

Module	Exper.	Inferred	95% CI	Rel. error (%)
LCoR	0.70	0.58	[0.47; 0.73]	13.9
RIP140	0.40	0.41	[0.33; 0.50]	7.5
Luciferase	2.45	3.44	[2.41; 5.43]	40.4

1 = expression under basal condition

H
Basal condition RA, double siNRIP1/siLCOR perturbation

Module	Exper.	Inferred	95% CI	Rel. error (%)
LCoR	0.70	0.55	[0.32; 0.84]	20.8
RIP140	0.40	0.37	[0.32; 0.42]	7.6
Hoxa5	2.14	1.96	[1.17; 3.39]	8.3

497
1 = expression under basal condition

498 Figure 2

C

Basal condition E2 & RA, double siNRIP1/siLCOR perturbation

Module	Exper.	Inferred	95% CI	Rel. error (%)
LCoR	0.55	0.52	[0.29; 0.83]	5.0
RIP140	0.36	0.38	[0.31; 0.47]	6.5
Luciferase	3.05	1.99	[1.61; 2.61]	30.9
Hoxa5	2.45	1.70	[0.83; 3.30]	34.7

1 = expression under basal condition

500   Figure 3

Gene	Eclidean distance
PGR	0.94
CCD88A	0.94
SYTL5	1.06
LOXL4	1.07
PNPLA3	1.13
RBBP8	1.14
RHOBTB1	1.22
XRCC2	1.23
...	...

504   Figure 4

A

```
library(aiMeRA)

# load data and define the model topology
data=data.setup(list(estr1_A,estr1_B,estr2_A,
                      estr2_B,estr3_A,estr3_B))
data=data2sdmean(data)
matp=read.rules(c("E2+RA+siLCoR->LCoR",
                  "E2+RA+siRIP140->RIP140",
                  "E2->Hoxa5","RA->Luciferase",
                  "E2+RA->0"))

# compute the model, CIs, and plot
res=mra(data$mean,matp)
inter=interval(data$mean,sd.ex,rules,nrep=2)
netgraph(res,inter = inter)
```

B

507   Figure 5