aiMeRA: A generic modular response analysis R package and its application to estrogen and retinoic acid receptors crosstalk

Gabriel Jimenez-Dominguez[1,2,3], Patrice Ravel[1,2,3], Stéphan Jalaguier[1,2,3], Vincent Cavaillès[1,2,3,*] and Jacques Colinge[1,2,3,*]

[1]Inserm U1194, Institut de Recherche en Cancérologie de Montpellier, 34298 Montpellier, France, [2]University of Montpellier, 34070 Montpellier, France, [3]ICM, Institut régional du Cancer de Montpellier, 34298 Montpellier, France

*To whom correspondence should be addressed.

Abstract

Motivation: Modular response analysis (MRA) is a widely used modeling technique to uncover coupling in biological networks under steady-state conditions. We wanted to explore the crosstalk between two nuclear receptors involved in several solid tumors. No open-source software package exists to apply MRA without advanced mathematical skills.

Results: We present a novel, synthetic mathematical derivation of MRA equations and make an R package available. Using this tool, we show the potential existence of different forms of crosstalk between estrogen and retinoic acid receptors. We also show that MRA models can be trained from RNA sequencing (RNA-seq) data. We introduce the concept of employing MRA models to search for new genes associated with a module activity. Finally, we propose an extension to MRA to add modules *a posteriori* and obtain activity inferences that outperform naïve predictions.

Availability: The aiMeRA R package is available at https://github.com/bioinfo-ircm/aiMeRA/, including qPCR data.

Contact: vincent.cavailles@inserm.fr, jacques.colinge@inserm.fr

1 Introduction

Modular response analysis (MRA) was introduced to infer the coupling between components of a biological system in a steady-state (Kholodenko *et al.*, 2002). It can be applied to components at different levels of details, e.g., individual genes or subsystems such as pathways or processes. It relies on the perturbation of individual components, the so-called modules. Various developments of MRA and related methods were recently reviewed (Santra *et al.*, 2018) but, despite its success, MRA is not accessible through a standard software package to the best of our knowledge. We thus reasoned that it would help the community to implement an R library (aiMeRA) that everyone could use without specific skills. Moreover, we propose a novel, elegant, and rather straightforward mathematical derivation of MRA basic equations, which should greatly facilitate understanding by new users. We showcase the use of the aiMeRA package by investigating the crosstalk between nuclear receptors (NRs) in a breast cancer (BC) cell line. A new extension of the method is also introduced to perform inferences at the genomic scale.

Estrogen receptors (ERs) belong to the NR superfamily, which act as transcription factors activated upon ligand binding. The two isoforms of ERs (ERα and ERβ) are involved in the control of cell proliferation and exhibit essential functions in tissue development and homeostasis, in particular in organs related to reproduction (Heldring *et al.*, 2007). ERα over-expression is frequently observed in breast, ovarian, endometrial, and other hormone-driven tumors. The transcriptional activity of ERs is modulated by several coregulatory complexes including coactivators and corepressors (Heldring *et al.*, 2007). In the presence of estrogens or any agonist ligand, ERs interact preferentially with coactivators, or with a specific subclass of corepressors including nuclear receptor-interacting protein 1 (NRIP1 or RIP140) and Ligand-dependent corepressor (LCoR). NRIP1 is a corepressor of particular interest because its expression is directly induced by estrogen, i.e., NRIP1 installs a negative feedback loop to keep ER signaling under control (Cavaillès *et al.*, 1995). NRIP1 abnormal expression is indeed observed in ER-driven tumors (Docquier *et al.*, 2013; Sixou *et al.*, 2018). LCoR represses transcription of estrogen-induced gene expression (Fernandes *et al.*, 2003), and NRIP1 expression was shown to be necessary for LCoR inhibitory activity in BC cells (Jalaguier *et al.*, 2017).

Interestingly, NRIP1 and LCoR function as corepressors for several liganded NRs. For instance, LCoR can repress vitamin D receptor (VDR), retinoic acid receptors (RARs), and RXR ligand-dependent transcription (Fernandes *et al.*, 2003) in addition to ERs. Moreover, NRIP1 is a known direct target and negative regulator of RAR transcription (White *et al.*, 2003).

There is experimental evidence of crosstalk between ER and RAR signaling (White *et al.*, 2005). For instance, ERα can suppress the basal expression of retinoic acid (RA)-responsive gene RARβ2, but also turns out to be necessary for its RA induction (Rousseau *et al.*, 2003). It was also found that ERα activates RARα1 expression in BC cells (Laganière *et al.*, 2005). Other authors intersected RAR targets identified by ChIP-seq with

ER binding sites to discover a significant overlap (Hua *et al.*, 2009). This work suggested a space competition mechanism for estrogen and RA signaling in BC. A potential cooperative interaction between RARα and ER was also shown in BC (Ross-Innes *et al.*, 2010). Since NRIP1 and LCoR expression can be both regulated by RAR and ER transcription, we can further hypothesize that these molecules mediate part of ER-RAR crosstalk. The induced expression of NRIP1 and LCoR by one receptor produces molecules able to repress signaling of both receptors subsequently.

We aimed at characterizing the ER-RAR-NRIP1-LCoR network, at the transcriptional level, utilizing transcript abundance measurements and MRA. Accordingly, we considered a steady-state situation in a BC (MCF7-derived) cell line that would model BC cells with or without constant estrogenic stimulation. Perturbation experiments were realized to generate quantitative PCR data unraveling interaction strengths in the network, i.e., coupling according to MRA principles.

Given the nature of ER and RAR, i.e., transcription factors, and the general ability of MRA to perform predictions (Klinger *et al.*, 2013), we introduced an extension of the method to perform genome-wide inferences exploiting mRNA sequencing (RNA-seq) data. For this purpose, we acquired whole transcriptomes under perturbed conditions identical to those used for qPCR. We first established that MRA could produce accurate results from RNA-seq data. Next, we asked whether the ER-RAR-NRIP1-LCoR network inferred by MRA could predict the mRNA abundance of estrogen-targeted genes better than a trivial model. This extension of MRA, where one or several modules do not experience perturbations, was called *unidirectional* to underline the implied absence of potential influence on the other modules.

2 Methods

Cell culture and perturbation experiments

We used MELN cells, an MCF7-derived cell line stably transfected with the estrogen-responsive luciferase reporter gene ERE-βGlob-Luc-SV-Neo (Balaguer *et al.*, 2001). The cell line was authenticated by short tandem repeat profiling and tested for mycoplasma contamination.

MELN cells were cultured in phenol red-free Dulbecco's modified Eagle medium (Gibco) containing 5% dextran-charcoal treated FCS (Invitrogen) and antibiotics (Gibco). Perturbations at NRIP1 and LCoR were obtained by siRNAs that were transfected using Interferrin (Polyplus). Perturbations at ERα and RARs were induced by their respective natural ligands: the hormones estrogen (17β-estradiol or E2 for short) and all-trans retinoic acid (RA).

MELN cells were obtained in the following conditions: basal (untreated), E2, RA, E2+RA, siNRIP1, siLCOR, siNRIP1+siLCOR, E2+siNRIP1, E2+siLCOR, E2+siNRIP1+siLCOR, RA+siNRIP1, RA+siLCOR, RA+siNRIP1+siLCOR, E2+RA+siNRIP1, E2+RA+siLCOR, and E2+RA+siNRIP1+siLCOR. These experiments were realized in triplicates. Cells were harvested after 18 hours of culture. E2-treated cells received 100nM E2, RA-treated cells 10 uM RA, and untreated cells ethanol. Validations of the response to E2 and siRNA interference are in Suppl. Fig. 1.

mRNA quantification

RNA was isolated using the Zymo Research kit (Zymo Research) and reverse transcription (RT)-qPCR assays were done using qScript (VWR) according to the manufacturer's protocol. Transcripts were quantified using SensiFAST SYBR (BioLine) on an LC480 instrument. The nucleotide sequences of the primers used for real-time PCR were:

RIP140-f (5'- AATGTGCACTTGAGCCATGATG -3'),
RIP140-r (5'- TCGGACACTGGTAAGGCAGG -3'),
LCoR-f (5'- GAACCTAGCGAACAAGACGGTG -3'),
LCoR-r (5'- TGGAGAGTGGCTCAGGGAAGT -3'),
Luciferase-f (5'- CTCACTGAGACTACATCAGC -3'),
Luciferase-r (5'- TCCAGATCCACAACCTTCGC -3'),
HOXA5-f (5'- GCGCAAGCTGCACATAAGTC -3'),
HOXA5-r (5'- GAACTCCTTCTCCAGCTCCA -3'),
ERα-f (5'- TGGAGATCTTCGACATGCTG -3'),
ERα-r (5'- TCCAGAGACTTCAGGGTGCT -3'),
RARα-f (5'- GGATATAGCACACCATCCCC -3'),
RARα-r (5'- TTGTAGATGCGGGGTAGAGG -3'),
PGR-f (5'- CGCGCTCTACCCTGCACTC-3'),
PGR-r (5'-TGAATCCGGCCTCAGGTAGTT-3').
(RT)-qPCR data are available from the R package.

mRNA sequencing

For two of the triplicates, in each condition, RNA was extracted as above described. Libraries were prepared with Illumina TruSeq kit and submitted to NextSeq500 sequencing (1x75bp/40M reads). The first 13 and last 7 bps were cut by an in-house Perl script to eliminate compositional bias. Cut reads were submitted to sickle to eliminate remaining low-quality regions. Alignments were performed against the human genome (hg38) with TopHat v2.10 (Kim *et al.*, 2013) and read counts extracted with HTSeq-Count (Anders *et al.*, 2015). The read count matrix was normalized with edgeR (McCarthy *et al.*, 2012) TMM algorithm. RNA-seq data will be available upon publication.

aiMeRA library implementation

We implemented the MRA method according to the mathematical formulation above as an R library. (RT)-qPCR data of this project were embedded in the R library for convenience and to provide an example. We also included the data used in the MRA original paper (Kholodenko *et al.*, 2002) such that users can check that our code gives the same results as those reported in the latter publication.

3 Results

3.1 Mathematical derivation

MRA original paper (Kholodenko *et al.*, 2002) introduced the concept of modeling interdependencies (coupling) within a biological system modularly. That is, subsystems involving molecules and their relationship at a detailed level, which would not be the interest of the study, could be captured as a single module with one measurable quantity defining the overall module activity. For instance, in the case of ERα signaling, it is possible to represent the complex process of ligand-binding and transcriptional activity by a single module (Fig. 1). The activity of this module is measured by a reporter gene, which is the luciferase in MELN cells. NRIP1 and LCoR activities were determined by their respective mRNA abundances. We ask the question of the direct dependence of each module activity with respect to the other modules activity. That is, we want to compute (signed) weights to put on the directed edges of Fig. 1. The answer is searched in a steady-state through successive elementary perturbations of each module activity. Depending on the application, this framework can be applied to different molecular species and processes, e.g., protein or metabolite concentrations, protein phosphorylation levels, etc. (Santra *et al.*, 2018).

Fig. 1. MRA general principle. Illustrated with the ERα-NRIP1-LCoR transcriptional network. Each module's activity level is given by a measured reporter. Coupling (edge weights) is determined from perturbation experiments.

Now, in full generality, we assume that there are n modules whose activities are given by $x \in \mathbb{R}^n$. We further admit the existence of n intrinsic parameters, $p \in \mathbb{R}^n$, one *per* module, which are perturbed by the elementary perturbations. One can imagine mRNA abundance parameters for perturbations such as siNRIP1 or siLCoR, and numbers of available ERα-E2 bound complexes for the E2 perturbation. In other circumstances, perturbations could change affinity constants or other physical characteristics. Finally, we assume that there exist $f: \mathbb{R}^n \times \mathbb{R}^n \longrightarrow \mathbb{R}^n$ of class $\mathcal{C}^1$ (continuously differentiable) such that

$$\dot{x} = f(x, p). \tag{1}$$

We do not need to know $f(x,p) = (f_1(x,p), \cdots, f_n(x,p))^t$ explicitly, but we need one more hypothesis that is the existence of a time $T > 0$ such that all the solutions we consider for any p and initial conditions of x, have reached a steady-state, i.e.,

$$\dot{x} = 0, \forall t > T.$$

The unperturbed, basal state of the modules is denoted $x^0 \in \mathbb{R}^n$ and it has corresponding parameters $p^0 \in \mathbb{R}^n$. According to our hypotheses, $f(x^0, p^0) = 0 \Leftrightarrow f_i(x^0, p^0) = 0, \forall i \in \{1, \cdots, n\}$. By the implicit function theorem, $\forall i$, there exists open neighborhoods $V_i \times W_i \subset \mathbb{R}^{n-1} \times \mathbb{R}^n$ of $(x_1^0, \cdots, x_{i-1}^0, x_{i+1}^0, \cdots, x_n^0, p_1^0, \cdots, p_n^0)$, $U_i \subset \mathbb{R}$ of x_i^0, and $g_i: V_i \times W_i \longrightarrow U_i$ (also $\mathcal{C}^1$) with

$$f_i(x_1^0, \cdots, x_{i-1}^0, g_i(\cdots), x_{i+1}^0, \cdots, x_n^0, p_1^0, \cdots, p_n^0) = 0. \tag{2}$$

We denote $x(p^0 + \Delta p)$, the steady-state corresponding to the changed parameters $p^0 + \Delta p$. We introduce the notation $x_{j \neq i}$ to denote all the x_j but x_i. Now, if we assume that $(x_{j \neq i}(p^0 + \Delta p), p^0 + \Delta p)$ belong to $V_i \times W_i$ for all the perturbations considered experimentally, then by Taylor's Formula

$$x_i(p^0 + \Delta p) = x_i(p^0) + g_i'(p^0) \begin{pmatrix} x_1(p^0 + \Delta p) - x_1(p^0) \\ \vdots \\ x_{i-1}(p^0 + \Delta p) - x_{i-1}(p^0) \\ x_{i+1}(p^0 + \Delta p) - x_{i+1}(p^0) \\ \vdots \\ x_n(p^0 + \Delta p) - x_n(p^0) \\ \Delta p_1 \\ \vdots \\ \Delta p_n \end{pmatrix}$$

$$+ o(\|\Delta p\|). \tag{3}$$

Dividing each side by $x_i(p^0)$, Eq. (3) can be rewritten

$$\frac{x_i(p^0 + \Delta p) - x_i(p^0)}{x_i(p^0)} =$$

$$\sum_{j=1, j \neq i}^n \frac{x_j(p^0)}{x_i(p^0)} \frac{\partial g_i}{\partial x_j}(p^0) \left(\frac{x_j(p^0 + \Delta p) - x_j(p^0)}{x_j(p^0)} \right)$$

$$+ \frac{1}{x_i(p^0)} \sum_{j=1}^n \frac{\partial g_i}{\partial p_j}(p^0)(\Delta p_j) + o(\|\Delta p\|). \tag{4}$$

Since parameter p_j influences module j only, $\frac{\partial g_i}{\partial p_j} = 0$ if $j \neq i$. Moreover, $g_i(x_{j \neq i}, p) = x_i(x_{j \neq i}, p)$ in $V_i \times W_i$, and if we denote

$$\frac{\Delta x_i}{x_i} = \frac{x_i(p^0 + \Delta p) - x_i(p^0)}{x_i(p^0)},$$

and

$$r_{i,j} = \frac{x_j(p^0)}{x_i(p^0)} \frac{\partial x_i}{\partial x_j}(p^0), j \neq i, \tag{5}$$

then

$$\frac{\Delta x_i}{x_i} = \sum_{j \neq i} r_{i,j} \left(\frac{\Delta x_j}{x_j} \right) + \frac{1}{x_i(p^0)} \frac{\partial x_i}{\partial p_i}(p^0)(\Delta p_i) + o(\|\Delta p\|). \tag{6}$$

We next consider elementary perturbations q_k, $k \in \{1, \cdots, n\}$, which only perturb module k, i.e., the parameter p_k. Neglecting the second-order term $o(\|\Delta p\|)$ and writing

$$\left(\frac{\Delta x_i}{x_i} \right)_{q_k}$$

the relative difference in module i activity upon Δp_k change induced by perturbation q_k, we find

$$\left(\frac{\Delta x_i}{x_i} \right)_{q_k} = \sum_{j \neq i} r_{i,j} \left(\frac{\Delta x_j}{x_j} \right)_{q_k}, k \neq i, \tag{7}$$

$$\left(\frac{\Delta x_i}{x_i} \right)_{q_i} = \sum_{j \neq i} r_{i,j} \left(\frac{\Delta x_j}{x_j} \right)_{q_k} + \frac{\partial x_i}{\partial p_i}(p^0) \left(\frac{\Delta p_i}{x_i} \right). \tag{8}$$

By defining $r_{i,i} = -1$, we can write Eqs (7) and (8) in matrix form:

$$rR = -P, \tag{9}$$

where R is the matrix that contains the experimentally observed relative activity changes $R_{j,k} = \left(\frac{\Delta x_j}{x_j} \right)_{q_k}$, $j, k \in \{1, \cdots, n\}$. P is a diagonal matrix with $P_{i,i} = \frac{\partial x_i}{\partial p_i}(p^0) \left(\frac{\Delta p_i}{x_i} \right)$, $i \in \{1, \cdots, n\}$. The system (9) can be solved in two steps (Kholodenko *et al.*, 2002). Firstly, $r = -PR^{-1}$ and because $r_{i,i} = -1$, we have $P_{i,i}(R^{-1})_{i,i} = 1$ and thus $P_{i,i} = \frac{1}{(R^{-1})_{i,i}}$. Secondly,

$$r = -[\text{diag}(R^{-1})]^{-1}R^{-1}.$$

The elements of R are defined by $\left(\frac{x_i(p^0 + \Delta p_k) - x_i(p^0)}{x_i(p^0)} \right)_{q_k}$ but as previously suggested (Kholodenko *et al.*, 2002), we preferred to estimate this quantity by

$$R_{j,k} = 2 \left(\frac{x_i(p^0 + \Delta p_k) - x_i(p^0)}{x_i(p^0 + \Delta p_k) + x_i(p^0)} \right)_{q_k}, \tag{10}$$

which avoids divisions by 0 and is numerically more stable.

Finally, from Eq. (5), we see that $r_{i,j}$ contains the searched coupling between MRA modules: direct action of j on i normalized by the ratio x_j/x_i. Similarly, $P_{i,i}$ measures the relative effect of q_i on x_i. We call it q_i magnitude. The implicit function theorem provides analytical expressions for g_i' in terms of f partial derivatives, but since f is generally unknown, we did not use them. In his seminal work, Kholodenko made additional hypotheses to show that r contains coupling information, which is not necessary with our derivation. To be rigorous, one should ultimately restrict the model to neighborhoods included in all the V_i's, W_i's, and U_i's.

MRA models have been largely used for their inference capabilities (Klinger *et al.*, 2013). Let us define a multiple perturbation q to be the linear combination of elementary perturbations q_k. For instance, a perturbation on modules i and j with the same individual magnitudes would be coded by a column vector c with 1's at positions i and j and 0's elsewhere. From Eq. (9), we compute

$$\left(\frac{\Delta x}{x}\right)_q = -r^{-1}Pc, \qquad (11)$$

with $\left(\frac{\Delta x}{x}\right)_q$ the column vector containing the inferred relative changes on each module activity. Denoting Δp the parameter changes induced by q, individual module activities are given by

$$\left(\frac{\Delta x_i}{x_i}\right)_q = -(r^{-1}Pc)_i = 2\frac{x_i(p^0 + \Delta p) - x_i(p^0)}{x_i(p^0 + \Delta p) + x_i(p^0)}$$

$$\Leftrightarrow \quad x_i(p^0 + \Delta p) = \frac{x_i(p^0)}{1+\frac{2}{(r^{-1}Pc)_i}}\left(\frac{2}{(r^{-1}Pc)_i} - 1\right). \qquad (12)$$

In case elementary perturbations contribute for different amounts to q, the vector c contains q_k's relative weights. In every case, linearity between the perturbation strength and its impact on p is assumed.

Confidence intervals (CI) around model parameters are estimated by a bootstrap procedure (Davison and Hinkley, 1997). We considered experimental designs with biological and technical replicates. We average the technical replicates to obtain one activity value *per* biological replicate. The latter are again averaged to compute the R matrix according to Eq. (10). From the biological replicates, we estimate the variance of each x_i employing an estimator optimized for a small sample size from Statistical Process Control theory (Wheeler and Chambers, 1992; Harter, 1960). Finally, a Gaussian distribution is assumed and 10^6 R matrices are generated, which are submitted to MRA computations. The 95% CI is obtained from the 2.5th and 97.5th percentiles. In case 0 is not included in the CI, the MRA parameter is deemed significant and marked by an asterisk in the figures.

Inferences obtained from Eq. (12) were also complemented by the estimation of CIs following the principles above.

3.2 Transcriptional data

ERβ and RARβ expression could not be quantified in MELN cells. We hence learned networks involving an ERα module, which transcriptional activity was reported by the ERE/luciferase construct. That is, luciferase mRNA abundance measured ERα activity. ERα mRNA abundance would combine ligand-bound and free amounts of the receptor, but only the ligand-bound ones matter in the model. We did not try to distinguish between RARα and RARγ. We estimated their combined transcriptional activity by the mRNA abundance of the *HOXA5* gene and the corresponding MRA module was named RARs. NRIP1 and LCoR activity were determined by their gene mRNA abundance. Since MELN cells are BC cells, we considered the E2-, RA-, or E2+RA-stimulated conditions as basal. That is, perturbations at ERα and RARs were negative (switch to ethanol). Perturbations at NRIP1 and LCoR were achieved by siRNAs, i.e., they were also negative.

3.3 The ERα-NRIP1-LCoR network

In an unstimulated condition (no E2), it is known that NRIP1 expression induces LCoR expression (Jalaguier *et al.*, 2017). We started by assessing this coupling under the E2 basal condition and found similar coupling (Fig. 2A). We also observed negative coupling from LCoR to NRIP1, which is logical since NRIP1 is a direct target of E2-bound ERα and LCoR

one of its corepressor. We next inferred the ERα-NRIP1-LCoR network under E2 (Fig. 2B). We could observe the known induction of NRIP1 by ERα with negative feedback (Cavaillès *et al.*, 1995). We also reconstituted the known inhibition of ERα by LCoR (Fernandes *et al.*, 2003). Interestingly, the induction of LCoR upon NRIP1 expression observed in Fig. 2A became a double inhibition via ERα in Fig. 2B. This makes sense since there is no transcriptional control by NRIP1 alone, it can only modulate ERα activity. Perturbation magnitudes are in Fig. 2C. Finally, we assessed the validity of the inferred network by checking its predictive power. From Fig. 2D, we note a reasonable fidelity of the model and that the relative errors are commensurate with the CI sizes, i.e., with data variability.

Basal condition E2, double siNRIP1/siLCOR perturbation

Module	Exper.	Inferred	95% CI	Rel. error (%)
LCoR	0.70	0.58	[0.47; 0.73]	13.9
RIP140	0.40	0.41	[0.33; 0.50]	7.5
Luciferase	2.45	3.44	[2.41; 5.43]	40.4

1 = expression under basal condition

Basal condition RA, double siNRIP1/siLCOR perturbation

Module	Exper.	Inferred	95% CI	Rel. error (%)
LCoR	0.70	0.55	[0.32; 0.84]	20.8
RIP140	0.40	0.37	[0.32; 0.42]	7.6
Hoxa5	2.14	1.96	[1.17; 3.39]	8.3

1 = expression under basal condition

Fig. 2. ER and RAR separated networks. A. Coupling between the two corepressors under E2 condition. A 95% CI for each model parameter was estimated and the parameter marked by an asterisk provided 0 was not included (nonzero with 5% significance). **B.** The ERα-NRIP1-LCoR transcriptional network. **C.** ERα, NRIP1, and LCoR perturbation magnitudes. **D.** Inference of gene expression under the dual siNRIP1 and siLCoR perturbation. **E.** Similar to A but under RA condition. **F.** The RARs-NRIP1-LCoR network. **G.** Similar to C. **H.** Similar to D.

Fig. 3. The ERα-RARs-NRIP1-LCoR network. A. MRA network model. **B.** Perturbation magnitudes under the dual E2 and RA stimulation. **C.** Inferred activity of the modules upon double siRNA inhibition of NRIP1 and LCoR.

Module	Exper.	Inferred	95% CI	Rel. error (%)
LCoR	0.55	0.52	[0.29; 0.83]	5.0
RIP140	0.36	0.38	[0.31; 0.47]	6.5
Luciferase	3.05	1.99	[1.61; 2.61]	30.9
Hoxa5	2.45	1.70	[0.83; 3.30]	34.7

Basal condition E2 & RA, double siNRIP1/siLCOR perturbation
1 = expression under basal condition

3.4 The RARs-NRIP1-LCoR network

Our next endeavor was to build a RARs-NRIP1-LCoR network before switching to the full network with both NRs. NRIP1 and LCoR coupling under RA stimulation (Fig. 2E) remained similar to its state under E2. That was expected since these two corepressors are used by several NRs. In Fig. 2F, we reconstituted the induction of NRIP1 expression by RAR as well as the inhibition of RAR expression by NRIP1 (White *et al.*, 2003). The inhibition of RAR by LCoR was also known (Fernandes *et al.*, 2003). Coupling between LCoR and NRIP1 is essentially similar to Fig. 2B since the NRIP1-to-LCoR arrows featured weak coupling. NRIP1 perturbation magnitude remained close, but LCoR perturbation changed 2-fold (Fig. 2G) although the same siRNAs were used. Inferences (Fig.2H) also supported the accuracy of the model.

3.5 The full ERα-RARs-NRIP1-LCoR network

We followed the same approach as above to construct a full model of ER-RAR crosstalk (Fig. 3A). Perturbation magnitudes (Fig. 3B) were in the same range as before under the new dual E2 and RA basal condition. Values for NRIP1 and LCoR were closer to the RAR-NRIP1-LCoR network. No literature reports coupling with the corepressors NRIP1 and LCoR under this particular condition. Only the crosstalk between RAR and ER mentioned in the introduction is known (Hua *et al.*, 2009; Ross-Innes *et al.*, 2010). We hence first challenged the model by testing its predictive accuracy (Fig. 3C), which was again satisfying.

Interestingly, cross inhibition of ER and RAR signaling acted along two paths. The model shows direct inhibition of ER transcriptional activity by the RAR module, which was described in the literature (Hua *et al.*, 2009; Ross-Innes *et al.*, 2010). Reciprocal inhibition was suggested but not significant in our data. In agreement with our hypothesis, we found a parallel crosstalk mechanism through the induction of NRIP1 expression, which

could subsequently repress both RARs and ERα. MRA modeling thus supported the coexistence of the two phenomena. Although LCoR reversed action on NRIP1 compared to the E2 and RA independent conditions might counterbalance cross inhibition of the two NRs, the strengths of coupling on the model edges and the much-attenuated induction of LCoR by NRIP1 suggested that it was not the case.

3.6 MRA models from RNA-seq data

Since MRA relies on module activity relative change (Eqs. (7-8)), absolute quantitation is not necessary. We thus computed an RARs-NRIP1-LCoR MRA models as HOXA5, NRIP1, and LCoR mRNA abundances were available in our RNA-seq data (Fig. 4A). Comparing with the qPCR-based model in Fig. 2F, we notice that all the significant edges of Fig. 2F plus NRIP1-to-LCoR were recovered. The only change is very weak coupling LCoR-to-RARs (-0.03) that became slightly positive (0.13) with RNA-seq

Fig. 4. Genomic-scale inferences. A. RARs-NRIP1-LCoR model trained from RNA-seq data. **B.** Eight closest replacement genes for ERE-Luc in the ERα-NRIP1-LCoR model. **C.** ERα-NRIP1-LCoR models with ERE-Luc replaced by PGR, trained from qPCR and RNA-seq data. **D.** Principle of unidirectional MRA. **E.** Accuracy of unidirectional MRA inference (udMRA & udMRA.ab) under the E2 condition with double siNRIP1/siLCoR perturbation versus simple predictors (mean, geometric mean (gMean), and maximum of the two siRNAs).

data. CIs were not determined for RNA-seq data since only two replicates were available.

The results above indicated that MRA could be applied to RNA-seq data. We, therefore, decided to exploit this opportunity by performing a new type of investigation. We used MRA to find a gene that would function as ERα transcription reporter, and would thus replace the ERE-Luc construct. Existing ChIP-seq data (Madak-Erdogan *et al.*, 2013) intersected with our RNA-seq data allowed us to identify 884 genes targeted by ERα and E2-regulated (edgeR analysis, P-value<0.01, fold-change>2). We hence computed 884 MRA models with siNRIP1 and siLCoR RNA-seq data, replacing ERE-Luc by each of those genes successively. The genes with closest Euclidean distances between their model coupling parameters (the $r_{i,j}$ matrix) and those of the original Fig. 2B qPCR model are listed in Fig. 4B. We decided to test PGR and measured its expression by qPCR. The qPCR- and RNA-seq-based models are featured in Fig. 4C. They accurately reproduced the original model of Fig. 2B and were very similar to each other, thereby further validating the use of RNA-seq data for MRA model training.

3.7 Unidirectional MRA on a genome-scale

We reasoned that the ERα-NRIP1-LCoR MRA model might provide means of predicting E2-regulated gene expression. We hence introduced a modified ERα-NRIP1-LCoR MRA model with one additional module that cannot influence the other modules (Fig. 4D). The gray unidirectional arrows in Fig. 4D represent the coupling between NRIP1, LCoR, the ERα module, and the added gene denoted by X. This coupling can be learned in the E2 basal condition by applying elementary perturbations as above. This must be repeated for each gene X considered. Gene X mRNA abundance is an $n + 1^{\text{th}}$ module and, by hypothesis, $r_{i,n+1} = 0, \forall i \in \{1, \cdots, n\}$, since no return arrows exist. From Eq. (7), we can compute $r_{n+1,j}, \forall j \in \{1, \cdots, n\}$, by solving the system

$$\left(\frac{\Delta x_{n+1}}{x_{n+1}}\right)_{q_k} = \sum_{j=1}^n r_{n+1,j}\left(\frac{\Delta x_j}{x_j}\right)_{q_k}, k \in \{1, \cdots, n\}.$$

The performance of this new type of MRA model (udMRA) was assessed by its ability to predict module $n + 1$ activity under the dual siN-RIP1/siLCoR condition, as we did above for the other models. To avoid trivially successful predictions on genes that would not vary, we limited the benchmark to the 884 genes above that were also significantly regulated upon siNRIP1 or siLCoR in the E2-stimulated condition. That left us with 60 genes. In Fig. 4E, we report the relative errors observed applying udMRA and comparing with naïve predictions. udMRA yielded significantly better estimates of the added module activity.

One could wonder whether perturbation magnitudes during double siRNA interference on the same biological system remain identical. That is, whether filling the vector c in Eq. (11) with 1's at the perturbed module indices (what we did so far) is the best option. Eq. (11) is written such that we can try different values. We searched for optimal coefficients a and b applied to siNRIP1 and siLCoR perturbations (at the corresponding indices in vector c), such that prediction errors of Luciferase, NRIP1, and LCoR expression (as in Fig. 2D) would be minimal. We found $a = 1$, and $b = 0.4$. Then, we used those coefficients in the udMRA model to try to predict the expression of the 61 benchmark genes more accurately. Indeed, we see in Fig. 4E that this new model called udMRA.ab achieved much better accuracy.

3.8 aiMeRA usage

The R package was designed to be generally applicable; it relies on the formulae presented here. It is able to work with any quantitative input, including biological and technical replicates. We included functionality to facilitate the definition of MRA model topologies (Fig. 5A). Model construction only involves the execution of a few generic R functions and network plots can be generated within R directly (Fig. 5B). It is also possible to export such graphs in the graphML format for loading into Cytoscape (Smoot *et al.*, 2011). More details are provided in the package documentation. aiMeRA is available from GitHub; submission to Bioconductor is pending.

4 Discussion

MRA modeling (Kholodenko *et al.*, 2002; Santra *et al.*, 2018) is a widely used technique to learn the coupling between the modules of a biological system from perturbation data sets (Fig. 1). We introduced a new mathematical derivation of the method and implemented a generic R package

A
```
library(aiMeRA)

# load data and define the model topology
data=data.setup(list(estr1_A,estr1_B,estr2_A,
                 estr2_B,estr3_A,estr3_B))
data=data2sdmean(data)
matp=read.rules(c("E2+RA+siLCoR->LCoR",
              "E2+RA+siRIP140->RIP140",
              "E2->Hoxa5","RA->Luciferase"
              "E2+RA->0"))

# compute the model, CIs, and plot
res=mra(data$mean,matp)
inter=interval(data$mean,sd.ex,rules,nrep=2)
netgraph(res,inter = inter)
```

B

Fig. 5. The aiMeRA R package. A. Example R code to load data, prepare them and compute the model. Note that NRIP1 was called after its common alternative name RIP140. Basal condition is E2 plus RA and we see that LCoR perturbation is defined as E2+RA+ siLCoR. Same logic for RIP140 (=NRIP1). Perturbation on the HOXA5 module reporting RARs activity is defined as E2, i.e., loss of RA stimulation. Etc. **B.** Plot in R using the igraph library (converted in gray-scale for the figure).

called aiMeRA (Fig. 5). Application thereof was illustrated on an un-published data set combining specific qPCR and broad RNA-seq data to explore crosstalk between ER and RAR, two important NRs involved in several tumors such as BC. Data analysis reconstituted some known inter-actions (Fig. 2) and supported a novel hypothesis that reciprocal negative coupling could be mediated by shared corepressors (Fig. 3).

We showed that MRA transcriptional models trained from RNA-seq data are close to those trained from qPCR (Fig. 4A-C). Which lead us to introduce an innovative application of MRA to probe genes genome-wide, searching for replacement reporters of module activity. That is, new genes that could be functionally related to an MRA module. In particular, we found that the progesterone receptor gene (*PGR*) reported on ligand-bound ERα transcriptional activity accurately. That observation indicates the po-tential value of this new use of MRA models since PGR is a widely used reporter of estrogen activity in BC in the clinic.

We additionally investigated the possibility to build hybrid MRA mod-els (udMRA & udMRA.ab) including unidirectional coupling to add mod-ules that were not perturbed in the training data set (Fig. 4D). In the con-text of the application presented in this report, i.e., the transcriptional ac-tivity of a system of transcription factors, we showed that the ERα-RARs-NRIP1-LCoR udMRA and udMRA.ab models could outperform naïve predictors (Fig. 4E). Other biological systems might be amenable to such modified models in the absence of strong back-coupling. The aiMeRA package methods support both RNA-seq data and udMRA models.

Acknowledgments

We thank Simon Cabello-Aguilar and Meriem Mekedem for useful discussions dur-ing the development of the project.

Funding

JC was supported by the Agence Nationale de la Recherche (grant ANR-16-CE17-0002-02), the Groupement des entreprises françaises dans la lutte contre le cancer (GEFLUC), and the Fondation ARC (PJA 20151203332).

References

Anders,S. *et al.* (2015) HTSeq--a Python framework to work with high-throughput sequencing data. *Bioinformatics*, **31**, 166–169.

Balaguer,P. *et al.* (2001) Reporter cell lines are useful tools for monitoring biologi-cal activity of nuclear receptor ligands. *Luminescence*, **16**, 153–158.

Cavaillès,V. *et al.* (1995) Nuclear factor RIP140 modulates transcriptional activa-tion by the estrogen receptor. *EMBO J*, **14**, 3741–3751.

Davison,A.C. and Hinkley,D.V. (1997) Bootstrap Methods and their Applications Cambridge University Press.

Docquier,A. *et al.* (2013) Negative regulation of estrogen signaling by ERβ and RIP140 in ovarian cancer cells. *Mol. Endocrinol.*, **27**, 1429–1441.

Fernandes,I. *et al.* (2003) Ligand-Dependent Nuclear Receptor Corepressor LCoR Functions by Histone Deacetylase-Dependent and -Independent Mech-anisms. *Molecular Cell*, **11**, 139–150.

Harter,H.L. (1960) Tables of Range and Studentized Range. *Ann. Math. Statist.*, **31**, 1122–1147.

Heldring,N. *et al.* (2007) Estrogen receptors: how do they signal and what are their targets. *Physiol. Rev.*, **87**, 905–931.

Hua,S. *et al.* (2009) Genomic antagonism between retinoic acid and estrogen sig-naling in breast cancer. *Cell*, **137**, 1259–1271.

Jalaguier,S. *et al.* (2017) Complex regulation of LCoR signaling in breast cancer cells. *Oncogene*, **36**, 4790–4801.

Kholodenko,B.N. *et al.* (2002) Untangling the wires: a strategy to trace functional interactions in signaling and gene networks. *Proc Natl Acad Sci U S A*, **99**, 12841–6.

Kim,D. *et al.* (2013) TopHat2: accurate alignment of transcriptomes in the presence of insertions, deletions and gene fusions. *Genome Biology*, **14**, R36.

Klinger,B. *et al.* (2013) Network quantification of EGFR signaling unveils poten-tial for targeted combination therapy. *Mol Syst Biol*, **9**, 673.

Laganière,J. *et al.* (2005) Functional genomics identifies a mechanism for estrogen activation of the retinoic acid receptor alpha1 gene in breast cancer cells. *Mol. Endocrinol.*, **19**, 1584–1592.

Madak-Erdogan,Z. *et al.* (2013) Integrative genomics of gene and metabolic regu-lation by estrogen receptors α and β, and their coregulators. *Mol. Syst. Biol.*, **9**, 676.

McCarthy,D.J. *et al.* (2012) Differential expression analysis of multifactor RNA-Seq experiments with respect to biological variation. *Nucleic Acids Res.*, **40**, 4288–4297.

Ross-Innes,C.S. *et al.* (2010) Cooperative interaction between retinoic acid recep-tor-alpha and estrogen receptor in breast cancer. *Genes Dev.*, **24**, 171–182.

Rousseau,C. *et al.* (2003) The N-terminal of the estrogen receptor (ERalpha) medi-ates transcriptional cross-talk with the retinoic acid receptor in human breast cancer cells. *J. Steroid Biochem. Mol. Biol.*, **86**, 1–14.

Santra,T. *et al.* (2018) Reconstructing static and dynamic models of signaling path-ways using Modular Response Analysis. *Current Opinion in Systems Biology*, **9**, 11–21.

Sixou,S. *et al.* (2018) Importance of RIP140 and LCoR Sub-Cellular Localization for Their Association With Breast Cancer Aggressiveness and Patient Survival. *Transl Oncol*, **11**, 1090–1096.

Smoot,M.E. *et al.* (2011) Cytoscape 2.8: new features for data integration and net-work visualization. *Bioinformatics*, **27**, 431–2.

Wheeler,D.J. and Chambers,D.S. (1992) Understanding Statistical Process Control 2nd ed. SPC Press, Inc.

White,K.A. *et al.* (2005) Limiting effects of RIP140 in estrogen signaling: potential mediation of anti-estrogenic effects of retinoic acid. *J. Biol. Chem.*, **280**, 7829–7835.

White,K.A. *et al.* (2003) Negative Feedback at the Level of Nuclear Receptor Coregulation SELF-LIMITATION OF RETINOID SIGNALING BY RIP140. *J. Biol. Chem.*, **278**, 43889–43892.