A Prokaryotic Membrane Sculpting BAR Domain Protein

Daniel A. Phillips[1*], Lori A. Zacharoff[2*], Cheri M. Hampton[3], Grace W. Chong[4], Anthony P. Malanoski[5], Lauren Ann Metskas[6], Shuai Xu[2], Lina J. Bird[5], Brian J. Eddie[5], Grant J. Jensen[6,7], Lawrence F. Drummy[3], Mohamed Y. El-Naggar[2,4,8], and Sarah M. Glaven[5]

[1]American Society for Engineering Education, Washington, DC, USA.
[2]University of Southern California, Department of Physics and Astronomy, Los Angeles, CA 90089
[3]Materials and Manufacturing Directorate, Air Force Research Laboratory, Wright-Patterson Air Force Base, WPAFB, OH, USA.
[4]University of Southern California, Department of Biological Sciences, Los Angeles, CA 90089
[5]Center for Bio/Molecular Science and Engineering, Naval Research Laboratory, Washington, DC, USA.
[6]California Institute of Technology, Division of Biology and Biological Engineering, Pasadena, CA 91125
[7]California Institute of Technology, Howard Hughes Medical Institute, Pasadena, CA 91125
[8]University of Southern California, Department of Chemistry, Los Angeles, CA 90089

[*]These authors contributed equally to this work

Classification: Biological sciences

Keywords: BAR domain | extracellular electron transport | membrane shaping | outer membrane vesicles | outer membrane extensions | *Shewanella*

Send all correspondence to:
Daniel A. Phillips
Daniel.phillips.ctr@nrl.navy.mil
(423) 215-2256
4555 Overlook Ave.
Washington, DC 20375

Lori A. Zacharoff
zacharof@usc.edu
(763) 221-0960
920 Bloom Walk
SSC, 215C
Los Angeles, CA 90089

Abstract

Bin/Amphiphysin/RVS (BAR) domain proteins belong to a superfamily of coiled-coil proteins that influence membrane curvature in eukaryotes and are associated with vesicle biogenesis, vesicle-mediated protein trafficking, and intracellular signaling[1-6]. BAR domain proteins have not been identified in bacteria, despite certain organisms displaying an array of membrane curvature phenotypes[7-16]. Here we identify a prokaryotic BAR domain protein, BdpA, from *Shewanella oneidensis* MR-1, an iron reducing bacterium known to produce redox active membrane vesicles and micrometer-scale membrane extensions. BdpA is required for uniform size distribution of outer membrane vesicles and is responsible for scaffolding outer membrane extensions (OMEs) into membrane structures with consistent diameter and curvature. While a strain lacking BdpA produces OMEs, cryogenic transmission electron microscopy reveals more lobed, disordered OMEs rather than the membrane tubes produced by the wild type strain. Overexpression of BdpA promotes OME formation even during planktonic conditions where *S. oneidensis* OMEs are less common. Heterologous expression also results in OME production in *Marinobacter atlanticus* CP1 and *Escherichia coli*. Based on the ability of BdpA to alter membrane curvature *in vivo*, we propose that BdpA and its homologs comprise a newly identified class of prokaryotic BAR (P-BAR) domains that will aid in identification of putative P-BAR proteins in other bacterial species.

Significance Statement:

BAR domain proteins are a crucial component of membrane architecture and vesicle trafficking in eukaryotic cells. Despite the recognized importance of BAR domain proteins in eukaryotic cell biology, this family of proteins has never before been reported for prokaryotes. Previously, outer membrane extensions of *Shewanella oneidensis* were shown to transition between chains of vesicles and tubules. We identified a component critical to structural transition as the first BAR domain protein in prokaryotes, which we term BdpA (BAR domain-like protein A). We show that BdpA (i) is enriched in outer membrane structures; (ii) is critical to regulate size of outer membrane vesicles; (iii) controls the shape of outer membrane extensions; (iv) confers membrane extension formation capabilities when expressed in other bacteria.

Main

Eukaryotic Bin/Amphiphysin/Rvs (BAR) domain-containing proteins generate membrane curvature through electrostatic interactions between positively charged amino acids and negatively charged lipids, scaffolding the membrane along the intrinsically curved surface of the antiparallel coiled-coil protein dimers[17-20]. Some BAR domain-containing proteins, such as the N-BAR protein BIN1, contain amphipathic helical wedges that insert into the outer membrane leaflet and can assist in membrane binding[21]. Other BAR domains can be accompanied by a membrane targeting domain, such as PX for phosphoinositide binding[22, 23], in order to direct membrane curvature formation at specific sites, as is the case with sorting nexin BAR proteins[4]. The extent of accumulation of BAR domain proteins at a specific site can influence the degree of the

resultant membrane curvature(24), and tubulation events arise as a consequence of BAR

domain multimerization in conjunction with lipid binding(25). Interactions between BAR

domain proteins and membranes resolve membrane tension, promote membrane

stability, and aid in localizing cellular processes, such as actin binding, signaling through

small GTPases, membrane vesicle scission, and vesicular transport of proteins(1, 26,

27). Despite our knowledge of numerous eukaryotic BAR proteins spanning a variety of

modes of curvature formation, membrane localizations, and subtypes (N-BAR, F-BAR,

and I-BAR), characterization of a functional prokaryotic BAR domain protein has yet to be

reported.

Bacterial cell membrane curvature can be observed during the formation of outer

membrane vesicles (OMV) and outer membrane extensions (OME). OMV formation is

ubiquitous and has many documented functions(9). OMEs are less commonly observed,

remain attached to the cell, and various morphologies can be seen extending from single

cells including *Myxococcus xanthus*(14, 15), flavobacterium strain Hel3_A1_48(8), *Vibrio*

vulnificus(10), *Francisella novicida*(28), *Shewanella oneidensis*(7, 29-31), and as cell-cell

connections in *Bacillus subtilis*(32-34) and *Eschericia coli*(35). Several bacterial proteins

have demonstrated membrane tubule formation capabilities *in vitro*(16, 36-40), but

despite the growing number of reports, proteins involved in shaping bacterial membranes

into OMV/Es have yet to be identified. Recently, researchers have begun to suspect that

OMV and OME formation has some pathway overlap(8), and it is proposed that proteins

are necessary to stabilize these structures(13).

S. oneidensis OMVs are redox-active and enriched with BdpA

Shewanella oneidensis is known to produce redox-active OMVs(41) and OMEs

coated with mulitheme cytochromes, particularly upon surface attachment(7, 30, 41).

However, little is known about their formation mechanism, control of shape or curvature,

and electrochemical properties that influence EET function.   OMVs were purified from

cells grown in batch cultures to characterize the redox features and unique proteome of

S. oneidensis OMVs, as well as to identify putative membrane shaping proteins.

Cryogenic transmission electron microscopy (cryo-TEM) tomographic reconstruction

slices of the purified samples showed uniform OMVs with the characteristic single

membrane phenotype and an approximate diameter of 200 nm (Fig. 1a).   Previous

measurements showed OMVs can reduce extracellular electron acceptors(41) and that

vesicles from *G. sulfurreducens* can mediate electron transfer(42). Electrochemical

activity of multiheme cytochrome complex MtrCAB and their ability to mediate

micrometer-scale electron transport has been characterized in whole cells(43), but no

electrochemical characterization of OME/Vs has been reported that link activity to

multiheme cytochromes. Here, electrochemical measurements of isolated OMVs were

performed to determine if purified OMVs maintain the characteristic redox features of the

MtrCAB complex when detached from cells. Cyclic voltammetry (CV) demonstrated redox

activity of isolated membrane vesicles adhered to a gold electrode via self-assembled

monolayers (Fig. 1b). The first derivative (Fig. 1b inset) revealed a prominent peak with

a midpoint potential of 66 mV and a smaller peak at -25 mV versus a standard hydrogen

reference electrode (SHE). This midpoint potential is consistent with the characteristics

of multiheme cytochromes such as MtrC/OmcA from previous microbial electrochemical

studies(43, 44), suggesting that the extracellular redox molecules of the cellular outer

membrane extends to OMVs.

The proteome of the OMVs was compared to the proteome of purified outer

membranes extracted from whole cells. Using a label-free quantification method(45),

significant differences in the ratio of individual proteins in the vesicle to the outer

membrane could be computed (log fold change) (Fig 1c). The proteome of the purified

OMVs showed 328 proteins were significantly enriched in the vesicles as compared to

the outer membrane, and 314 proteins were significantly excluded from the vesicles (Fig.

1c, Supplemental table 1). MtrCAB cytochromes were present in the OMVs as well as

the outer membrane, supporting redox activity of OMVs observed by CV. Active protein

sorting into eukaryotic vesicles is a coordinated process involving protein sorting signal

recognition, localized membrane protein recruitment, initiation of membrane curvature

induction, and coating nascent vesicles with protein scaffolds(46). Several proteins

significantly enriched in the vesicles were identified that could contribute to OMV

formation, including the murein transglycosylase, the peptidoglycan degradation enzyme

holin, cell division coordinator CpoB, and a highly enriched putative BAR domain-

containing protein encoded by the gene at open reading frame SO_1507, hereafter

named BAR domain-like protein A (BdpA) (Fig. 1d).

Vesicle enrichment of BdpA led us to the hypothesis that BdpA could be involved

in membrane shaping of OMVs based on the role of BAR domain proteins in eukaryotes.

The C-terminal BAR domain of BdpA is predicted to span an alpha-helical region from AA

276-451 (E-value = 2.96e-03); however, since the identification of the protein is based on

homology to the eukaryotic BAR domain consensus sequence (cd07307), it is possible

that the BAR domain region extends beyond these bounds (Fig. 1d). Coiled coil

prediction(47) suggests BdpA exists in an oligomeric state of antiparallel alpha-helical

dimers, as is the case for all known BAR domain proteins(18, 48-50). The predicted

structure of the BAR domain-containing region generated in i-TASSER(51) shows a 3

helix bundle (Supplemental Fig. 1a). When the BdpA BAR domain monomers were

aligned to the dimeric structure of the F-BAR protein Hof1p(52), the predicted dimer

interface residues of BdpA came into proximity, revealing an intrinsically curved dimer

with positively charged residues along the concave surface (Supplemental Fig. 1b). BdpA

has a N-terminal signal peptide with predicted cleavage sites between amino acids 22-

23, suggesting non-cytoplasmic localization (Fig. 1d). A galactose-binding domain-like

region positioned immediately downstream of the signal peptide supports lipid targeting

activity seen in other BAR domain proteins, such as the eukaryotic sorting nexins(3) which

have phox (PX) domains that bind phosphoinositides(53). The *S. oneidensis* rough-type

lipopolysaccharide (LPS) contains 2-acetamido-2-deoxy-D-galactose(54), which

suggests possible localization of the protein to the outer leaflet of the outer membrane.

BdpA controls size distribution of vesicles

To determine whether BdpA influences vesicle morphology, OMVs were harvested

from wild type (WT) cells and cells in which the gene for BdpA had been deleted (Δ*bdpA*),

and their diameters were measured by dynamic light scattering (DLS). WT OMVs had a

median diameter of 190 nm with little variability in the population (standard deviation (s.d.)

= ±21 nm), while the diameters of Δ*bdpA* OMVs were distributed over a wider range with

a median value of 280 nm (s.d. = ± 131 nm (Fig 2a). The data suggest BdpA controls

vesicle diameter in membrane structures *ex vivo*, potentially acting by stabilizing OMVs.

Cell-associated OMV frequency and size distribution was also measured in live

cultures using a perfusion flow imaging platform and the membrane stain FM 4-64, as

described previously(29). *S. oneidensis* strains were monitored for OME/V production

and progression over the course of 5 hours (>5 fields of view per replicate, n=3) using

time-lapse imaging. Spherical membrane stained extracellular structures were classified

as OMVs, while larger aspect ratio (i.e. length greater than the width) structures were

classified as OMEs. The proportion of cells producing 'large' vesicles, defined as having

a membrane clearly delineated from the interior of the vesicles and typically >300 nm,

was quantified, and Δ*bdpA* cells produced significantly more large vesicles compared to

WT cells (Fig. 2b) even though both the overall frequency of vesiculation and extensions

were the same (Fig. 2c). Previous studies showed that OMEs transition between large

vesicles and OMEs over time(29). BdpA appears to be involved in this transition in

Shewanella due to the increased frequency of large vesicles from Δ*bdpA* cells. The

median diameter of the OMVs is also the apparent maximum diameter observed in outer

membrane extensions(29) suggesting BdpA influences membrane morphologies of both

structures.

BdpA constrains membrane extension morphology

The resolution of fluorescence microscopy was insufficient to identify

morphological differences between OMEs. Cells were also visualized after deposition

onto a glass coverslip instead of a perfusion flow chamber as previously reported(7).

BdpA was expressed from a 2,4-diacetylphloroglucinol (DAPG)-inducible promoter(55)

(P$_{PhlF}$-BdpA) in the Δ*bdpA* strain containing the plasmid p452-*bdpA* with 12.5 μM DAPG.

After 3 hours post deposition on cover glass, OMEs can be seen extending from WT,

Δ*bdpA*, and Δ*bdpA* p452-*bdpA* cells (Fig. 3a, Supplemental Fig. 2, Supplemental videos

1-3, 5 fields of view, n=3). Similar to perfusion flow experiments, no statistically significant

difference in the overall frequency of OME production was observed between the cells in

static cultures.

Cryo-TEM was used to assess morphological differences between the OMEs in

each of the strains at the ultrastructural level. *S. oneidensis* OMEs from unfixed WT,

Δ*bdpA,* and Δ*bdpA* p452-*bdpA* strains were visualized at 90 minutes (Supplemental Fig.

3) and 3 hours (Fig. 3a) following cell deposition onto EM grids. By 3 hours post

inoculation, images of WT cells consistently show narrow, tubule-like OMEs (p = 0.01,

student's t-test) or vesicle chains of symmetric curvature (Fig. 3b,c n=31).   The Δ*bdpA*

OMEs generally appear as lobed, disordered vesicle chains with irregular curvature (p =

0.01, student's t-test), and vesicles can be observed branching laterally from lobes on the

extensions (Fig. 3b,c n=13).   Nascent WT OMEs from previous studies also exhibited

lateral branching of vesicles and lobes, but they exhibited uniform curvature and diameter

between lobes and were observed immediately following OME formation(29).   Tubules

were not observed in any Δ*bdpA* OMEs at 3 hours (Fig. 3d, p = 0.01, student's t-test).

OMEs from Δ*bdpA* p452-*bdpA* cells appear as a narrow tubules of a uniform curvature or

as ordered vesicle chains (Fig. 3b,c n=3), showing that expression of BdpA from a

plasmid rescues the mutant phenotype by constricting and ordering OMEs into narrow

tubules and chains (Fig. 3d, p = 0.01, student's t-test). Earlier in OME progression at 90

minutes, WT OME phenotypes appeared narrow, tubule-like, and seldom interspersed

with lobed regions (Supplemental figure 3a).   In Δ*bdpA* OMEs, lobed regions are

prevalent with irregular curvature (Supplemental figure 3b). Several narrow Δ*bdpA* p452-

bdpA OMEs evenly interspersed with slight constriction points or "junction densities" were

observed extending from a single cell (Supplemental Fig. 3c).

Expression of BdpA results in OMEs during planktonic growth

S. oneidensis OMEs are more commonly observed in surface attached cells than

planktonic cells(7, 29). BAR domain proteins can directly promote tubule formation from

liposomes *in vitro*(24), so inducing expression of an additional copy of the *bdpA* gene

prior to attachment could result in OME formation even during planktonic growth. Growth

curves were similar in cultures with the pBBR1-mcs2 empty vector in either of the WT or

Δ*bdpA* background strains, but induction of *bdpA* in Δ*bdpA* p452-*bdpA* cells at higher

concentrations of 1.25 and 12.5 µM 2,4-diacetylphloroglucinol affected the growth rate

(Supplemental figure 4).   Planktonic cultures inoculated from overnight cultures were

induced with 12.5 µM DAPG for 1 hour, labeled with FM 4-64, and imaged by confocal

microscopy. Neither WT (Fig. 4) nor WT with the empty plasmid exposed to 12.5 µM

DAPG (not shown) displayed OMEs immediately following deposition onto cover glass.

However, 12.5 µM DAPG-induced bdpA expression from p452-*bdpA* in the WT backround

strain (WT p452-*bdpA*) displayed OMEs immediately, ranging between 1-7 extensions

per cell (Figure 4, Supplemental video 4). The abundance of OMEs suggest that

increased *bdpA* expression in planktonic cultures can initiate membrane sculpting into

OMEs at the expense of cell division.

The ultrastructure of OMEs resulting from expression of *bpdA* from WT p452-*bdpA*

cells was examined by cryo-TEM, but in this case samples from planktonic cultures were

vitrified on EM grids after induction rather than incubation during induction on the EM

grids. By expressing extra copies of *bdpA* in the WT strain during planktonic induction,

we predicted that OME morphology would be predominately reliant on curvature

formation by BdpA rather than other unidentified structural proteins involved in intrinsic

OME formation. OMEs appear as tubule-like segments interspersed with pearled regions

proximal to the main cell body (Fig. 4b). OMEs from the MR-1 p452-*bdpA* strain are

observed as thin, tubule-like outer membrane vesicle chains, suggesting BdpA

involvement in the constriction of the larger outer membrane vesicle chains into longer,

tubule-like extensions with more evenly interspersed junction densities. The BdpA OME

phenotype more closely resembles membrane tubules formed by the F-BAR protein

Pacsin1 from eukaryotic cells, showing a mixture of tubule regions interspersed with

pearled segments(56, 57). Orientation and association of Pacsin1 dimers on the

membrane surface with one another impacted tubule morphology, and BdpA membrane

sculpting could be similar mechanistically.

BdpA-mediated membrane extensions in *Marinobacter atlanticus* CP1 and *E. coli*.

To test the effect of expressing BdpA in an organism with no predicted BAR

domain-containing proteins and no apparent OME production, BdpA was expressed in

Marinobacter atlanticus CP1(58). *Marinobacter* and *Shewanella* are of the same

phylogenetic order (*Alteromonadales*) and have been used for heterologous expression

of other *S. oneidensis* proteins, such as MtrCAB(59, 60). Upon exposure to DAPG, *M.*

atlanticus containing the p452-*bdpA* construct (CP1 p452-*bdpA*) forms membrane

extensions (Figure 4). OMEs ranged from small membrane blebs to OME tubules

extending up to greater than 10 μm in length from the surface of the cell (Supplemental

Fig. 5). As noted previously, variation in the tubule phenotypes are commonly seen in

tubules from eukaryotic F-BAR proteins(56, 57), showing possible mechanistic overlap of

mutable membrane curvature functionalities between these two separate BAR domain

proteins.

In previous membrane curvature formation experiments with eukaryotic BAR

domain proteins, localized BAR domain protein concentrations affected the resultant

shape of the membranes, ranging from bulges to tubules and branched, reticular tubule

networks at the highest protein densities(61-63). We predicted that expression of BdpA

in cells optimized for protein overexpression, such *E. coli* BL21(DE3), would show OMEs

resembling BAR protein concentration-dependent structures previously observed from

eukaryotic BAR protein experiments *in vitro*. While the uninduced *E. coli* BL21(DE3)

p452-*bdpA* cells had uniform, continuous cell membranes similar to those of plasmid-free

BL21(DE3) cells under the conditions tested, *E. coli* BL21(DE3) cells containing the p452-

bdpA vector induced with 12.5 µM DAPG had outer membrane extensions and vesicles

(Figure 4). When visualized over time, OMEs progressed towards a network of reticular

membrane structures extending from the cell (Fig. 4c). After 30 minutes, additional

membrane blebs were observed that developed into elongated OMEs by 60 minutes.

Growth of *E. coli* OMEs was coincident with shrinking of the cell body (from initial cell

length = 4.5 µm to 3.5 µm at 60 minutes), supporting direct membrane sculpting activity

of BdpA.

P-BAR: a new BAR domain subtype

The discovery of a novel, functional BAR domain protein in prokaryotes provokes

questions into the evolutionary origin of BAR domains, such as whether the BdpA BAR

domain in *S. oneidensis* arose as a result of convergent evolution, a horizontal gene

transfer event, or has a last common ancestor across all domains of life. BdpA homologs

were identified by PSI-BLAST in other *Gammaproteobacteria*, including most but not all

species of *Shewanella*, as well as *Alishewanella*, *Rheinheimera*, and *Cellvibrio*

(Supplemental Fig. 6).   The current BAR domain Pfam Hidden Markov Model (HMM)

prediction analysis identified BAR domain features in only 5 of the 52 prokaryotic

homologs despite greater than 90% homology to *S. oneidensis* BdpA. An amino acid

alignment of the 52 BdpA homologs was used to generate a maximum likelihood

phylogenetic tree showing evolutionary relatedness of BdpA orthologs to the BAR domain

prediction sequences (Supplemental Fig. 6). The 5 BdpA orthologs predicted to contain

a BAR domain based on the current model were subsequently aligned with representative

known BAR proteins from the various BAR domain subtypes (N-BAR, F-BAR, and I-

BAR)(64). BdpA and its prokaryotic orthologs cluster separately from the eukaryotic BAR

proteins in their own distinct clade (Fig. 5), suggesting that while BdpA contains a

functional BAR domain, it represents its own class of BAR domain, hereafter named P-

BAR (P̲rokaryotic BAR). It is possible that the P-BAR domain arose as a result of

horizontal gene transfer from a eukaryote due to the prevalence of eukaryotic coiled-coil

proteins with predicted homology to BdpA after 2 iterations of PSI-BLAST. However, the

branch lengths and low bootstrap values supporting the placement of P-BAR relative to

other BAR domain subtypes make it challenging to directly infer the evolutionary history

of P-BAR domains. Discovery of other putative P-BAR proteins would help to build this

analysis, and if future comparative proteomics analysis of OME/Vs demonstrates

overlapping activity of BdpA with preferential cargo loading into OME/Vs, it could hint at

the evolutionary origins of vesicle-based protein trafficking. Conservation of BAR domain

proteins supports the notion that three-dimensional organization of proteins in lipid

structures is as important to prokaryotes as it is eukaryotes, and suggests additional novel

P-BAR proteins are waiting to be discovered.

Methods

Bacterial strains, plasmids, and medium The bacterial strains used in this study can

be found in Supplemental Table 1. *S. oneidensis* strains were grown aerobically in Luria

Bertani (LB) media at 30°C with 50 µg/mL kanamycin when maintaining the plasmid. To

observe membrane extensions, cells were centrifuged and resuspended in a defined

media comprised of 30 mM Pipes, 60 mM sodium DL-lactate as an electron donor, 28mM

NH_4Cl, 1.34 mM KCl, 4.35 mM NaH_2PO_4, 7.5 mM NaOH, 30 mM NaCl, 1mM $MgCl_2$, 1

mM $CaCl_2$, and 0.05 mM ferric nitrilotriacetic acid(30). *Marinobacter atlanticus* CP1

strains were grown in BB media (50% LB media, 50% Marine broth) at 30°C with 100

µg/mL kanamycin to maintain the plasmids as described previously(58).

Inducible BdpA expression plasmids were constructed for use in *S. oneidensis*

MR-1, *M. atlanticus* CP1, and *E. coli* BL21(DE3) using the pBBR1-mcs2 backbone

described previously(58). The Marionette sensor components (*phlF* promoter,

consitutively expressed PhlF repressor, and yellow fluorescence protein (YFP)) cassette

from pAJM.452(55) was cloned into the pBBR1-mcs2 backbone, and the YFP cassette

was replaced with the gene encoding BdpA by Gibson assembly (primers in

Supplemental Table 2). The resulting plasmid was given the name p452-*bdpA*. The

Gibson assembly reactions were electroporated into *E. coli* Top10 DH5α cells

(Invitrogen), and the sequences were confirmed through Sanger sequencing (Eurofins

genomics). Plasmid constructs were chemically transformed into conjugation-competent

E. coli WM3064 cells for conjugative transfer into the recipient bacterial strains of *S.*

oneidensis MR-1 and *M. atlanticus* CP1. The same BdpA expression vector was

transformed into *E. coli* BL21(DE3) cells (Invitrogen) by chemical transformation.

Generation of a scarless *ΔbdpA* knockout mutant of *S. oneidensis* was performed

by combining 1 kilobase fragments flanking upstream and downstream from *bdpA* by

Gibson assembly into the pSMV3 suicide vector. The resultant plasmid pSMV3_1507KO

was transformed into *E. coli* DH5α *λpir* strain UQ950 cells for propagation. Plasmid

sequences were confirmed by Sanger sequencing before chemical transformation into *E.*

coli WM3064 for conjugation into *S. oneidensis*. Conjugation of pSMV3_1507KO into *S.*

oneidensis MR-1 was performed as described previously[31]. Optical densities at 600nm

were measured to determine growth curves for each strain in either LB or SDM with 50

µg/mL kanamycin and 2,4-diacetylphloroglucinol (DAPG) as indicated. Cultures of 500

µL samples diluted to an initial OD_{600} of 0.1 were grown in Costar® polystyrene 48 well

plates (Corning Incorporated) within a Tecan Infinite M1000 Pro (Grödig, Austria) plate

reader at 30°C with shaking agitation at 258 rpm. Optical densities were recorded every

15 minutes with i-Control software (2.7). All measurements were performed in triplicate.

Purification of Outer Membrane Vesicles *S. oneidensis* MR-1 cells were grown in LB

in 1L non-baffled flasks at 30° C at 200 RPM. When an OD_{600} of 3.0 was reached, cells

were pelleted by centrifugation at 5000 x g for 20 min at 4°C, resulting supernatant was

filtered through a 0.45 µm filter to remove remaining bacterial cells. Vesicles were

obtained by centrifugation at 38,400 x g for 1 h at 4°C in an Avanti J-20XP centrifuge

(Beckman Coulter, Inc). Pelleted vesicles were resuspended in 20 ml of 50 mM HEPES

(pH 6.8) and filtered through 0.22 µm pore size filters. Vesicles were again pelleted as

described above and finally resuspended in 50 mM HEPES, pH 6.8, except for vesicle

preparations used for electrochemistry which were suspended in 100 mM MES, 100 mM

KCl, pH 6.8. Extracellular DNA, flagella, and pili can all be co-purified. Protocol was

adapted from Perez-Cruz et al(65).

Cryoelectron tomography Vesicle samples were diluted to a protein concentration of

0.4 mg/mL and applied to glow-discharged, X-thick carbon-coated, R2/2, 200 mesh

copper Quantifoil grid (Quantifoil Micro Tools) using a Vitrobot chamber (FEI). Grids were

automatically plunge frozen and saved for subsequent imaging. No fixative was used.

Images were collected on an FEI Krios transmission electron microscope equipped with

a K2 Summit counting electron-detector camera (Gatan). Data were collected using

customized scripts in SerialEM(66), with each tilt series ranging from −60° to 60° in 3°

increments, an underfocus of ~1-5 μm, and a cumulative electron dose of 121 e/A^2 for

each individual tilt series. Tomograms were reconstructed using a combination of

ctffind4(67) and the IMOD software package(68).

332
Dynamic Light Scattering Distributions of vesicle diameters were measured with Wyatt

Technology's Möbiuζ dynamic light scattering instrument with DYNAMICS software for

data collection and analysis. Data was collected using a 0-50 mW laser at 830 nm. The

scattered photons were detected at 90˚. Measurements were recorded from 11 biological

replicates for WT OMVs, and 9 replicates for Δ*bdpA* OMVs. Mobius software analyzed

the population of particles to generate a table with binned diameters and the percentage

of particles at each diameter. In order to compute the average vesicle size from each

sample, a weighted average was computed so that diameter bins that had the most

number of vesicles would be accurately represented in the final average weight of the

population. The product of each diameter is multiplied by its percentage in the population, these products are added together for each sample, divided by the sum of weights. The weighted diameters per replicate were then averaged for each genotype. Statistical significance was determined by Student's *t*-test, and error bars represent standard deviation.

Electrochemistry CHA Industries Mark 40 e-beam and thermal evaporator was used to deposit a 5 nm Ti adhesion layer and then a 100 nm Au layer onto cleaned glass coverslips (43mm x 50mm #1 Thermo Scientific Gold Seal Cover Glass, Portsmouth NH, USA). Self-assembled monolayers were formed by incubated the gold coverslip in a solution of 1mM 6-mercaptohexanoic acid in 200 proof ethanol for at least 2 hours. Electrode was then rinsed several time in ethanol followed by several rinses in milliQ water. The SAMs layer was then activated by incubation in 100 mM N-(3-Dimethylaminopropyl)-N′-ethylcarbodiimide hydrochloride and 25 mM N-hydroxysuccinimide, pH 4, for 30 minutes. A sample of outer membrane vesicles was deposited on the surface of the electrode and incubated at room temperature overnight in a humid environment. Cyclic voltammetry was performed in a 50 mL three-electrode half-cell completed with a platinum counter electrode, and a 1 M KCl Ag/AgCl reference electrode electrical controlled by a Gamry 600 potentiostat (Gamry, Warminster, PA). The whole experiment was completed in an anaerobic chamber with 95% nitrogen, 5% hydrogen atmosphere.

Proteomics Vesicle samples were prepared as described above. *S. oneidensis* outer membrane (OM) was purified via the Sarkosyl method described by Brown et al.(69). A 50 mL overnight culture of cells was harvested by centrifugation at 10,000 × g for 10 min. The cell pellet suspended in 20 mL of 20 mM ice-cold sodium phosphate (pH 7.5) and passed four times through a French Press (12000 lb/in^2). The lysate was centrifuged at 5,000 × g for 30 min to remove unbroken cells. The remaining supernatant was centrifuged at 45,000 × g for 1 h to pellet membranes. Crude membranes were suspended in 20 mL 0.5% Sarkosyl in 20 mM sodium phosphate and shaken horizontally at 200 rpm for 30 min at room temperature. The crude membrane sample was centrifuged at 45,000 × g for 1 h to pellet the OM. The pellet of OM was washed in ice-cold sodium phosphate and recentrifuged.

To prepare for mass spectrometry samples were treated sequentially with urea, TCEP, iodoactinamide, lysl endopeptidase, trypsin, and formic acid. Peptides were then desalted by HPLC with a Microm Bioresources C8 peptide macrotrap (3x8mm). The digested samples were subjected to LC-MS/MS analysis on a nanoflow LC system, EASY-nLC 1200, (Thermo Fisher Scientific) coupled to a QExactive HF Orbitrap mass spectrometer (Thermo Fisher Scientific, Bremen, Germany) equipped with a Nanospray Flex ion source. Samples were directly loaded onto a PicoFrit column (New Objective, Woburn, MA) packed in house with ReproSil-Pur C18AQ 1.9 um resin (120A° pore size, Dr. Maisch, Ammerbuch, Germany). The 20 cm x 50 μm ID column was heated to 60° C. The peptides were separated with a 120 min gradient at a flow rate of 220 nL/min. The gradient was as follows: 2–6% Solvent B (7.5 min), 6-25% B (82.5 min), and 25-40% B (30 min) and to 100% B (9min). Solvent A consisted of 97.8% H2O, 2% acetonitrile,

and 0.2% formic acid and solvent B consisted of 19.8% H2O, 80% ACN, and 0.2% formic

acid. The QExactive HF Orbitrap was operated in data dependent mode with the Tune

(version 2.7 SP1build 2659) instrument control software. Spray voltage was set to 2.5

kV, S-lens RF level at 50, and heated capillary at 275 °C. Full scan resolution was set to

60,000 at m/z 200. Full scan target was 3×106 with a maximum injection time of 15 ms.

Mass range was set to 300−1650 m/z. For data dependent MS2 scans the loop count

was 12, target value was set at 1×105, and intensity threshold was kept at 1×105.

Isolation width was set at 1.2 m/z and a fixed first mass of 100 was used. Normalized

collision energy was set at 28. Peptide match was set to off, and isotope exclusion was

on. Data acquisition was controlled by Xcalibur (4.0.27.13) and all data was acquired in

profile mode.

Bioinformatics Putative BAR domain SO_1507 (BdpA) was identified in search of

annotation terms of *S. oneidensis* MR-1. The conserved domain database (CDD-

search)(NCBI) was accessed to identify the position-specific scoring matrix (PSSM) of

the specific region of SO_1507 that represented the BAR domain (amino acid residues

at positions 276-421). The domain prediction matched to BAR superfamily cl12013 and

specifically to the family member BAR cd07307. LOGICOIL multi-state coiled-coil

oligomeric state prediction was used to predict the presence of coiled-coils within

BdpA(47). SignalP 6.1 was used to detect the presence of the signal peptide and cellular

localization of BdpA(70). I-TASSER protein structure prediction(51) was used to generate

a predicted model of the BAR domain region of BdpA. Alignment of the predicted BdpA

BAR domain structure model to the structure of Hof1p was performed in the PyMOL

Molecular Graphics System, Version 2.0 (Schrödinger, LLC).

A PSI-BLAST(71) search against the NCBI nr database was performed using the

BdpA BAR sequence as the initial search seed to determine how prevalent the BdpA BAR

domain is in related species. Conserved BdpA orthologs were annotated as hypothetical

proteins in all of the species identified. In the initial round, 24 proteins were found from

other organisms identified as *Shewanella* with a high conservation among the proteins

and another 28 proteins were found in more distant bacteria species that had similarity of

65% to 44%. A second iteration identified a few proteins much more distantly related from

bacterial species and then proteins from eukaryote phylum *Arthropoda* that were

annotated as being centrosomal proteins. All of the found proteins from bacterial species

were hypothetical proteins with no known function. Only five of the proteins from the

search returned hits to the PSSM of the BAR cd07307. The identity among the proteins

was very high and examination of the proteins suggests that a functional form similar to

the BAR domain would result for all the found proteins. Overall, this places BdpA as a

protein that just barely meets criteria via PSSM models to be assigned as matching the

BAR domain while the rest of the proteins found have enough differences to fail to match

the BAR model while still being very similar to BdpA. An attempt was made to build up a

HMM (Hidden Markov Model) using hmmer(72) to use for searching for other proteins

that might match, but as with the PSI-BLAST search, only the proteins that formed the

model returned as good matches. There appears to be a tight clade of very similar

proteins with very little differentiation in the sequence. This indicates that while sequence

homology between BdpA and the existing BAR domain consensus sequence predicted

the BAR domain region in BdpA using hmmer or NCBI tools, the sequence conservation

is at the cusp of a positive hit by the HMM since other closely related (>90% homology)

BdpA orthologs were not predicted to contain a BAR domain by this method. The most

homologous eukaryotic protein to BdpA (27%) is a putative centrosomal protein in

Vollenhovia emeryi (accession #: XP_011868153) that is predicted to contain an amino

terminal C2 membrane binding domain and a carboxy-terminal SMC domain within a

coiled-coil region. Despite CDD search failing to predict the presence of a BAR domain

in this protein, it does not preclude the presence of one, pending an updated BAR Pfam

HMM. Alignments and phylogenies were constructed in MEGA 7. MUSCLE was used to

align protein sequences, and Maximum Likelihood phylogenies were inferred using the

Le-Gascuel (LG+G) substitution matrix(73). Initial trees for the heuristic search were

obtained automatically by applying Neighbor-Join and BioNJ algorithms to a matrix of

pairwise distances estimated using a JTT model, and then selecting the topology with

superior log likelihood value. A discrete Gamma distribution was used to model

evolutionary rate differences among sites as indicated in figure legends. All positions with

less than 85% site coverage were eliminated.

Confocal microscopy For *in vivo* imaging of intrinsic outer membrane extension

production, *S. oneidensis* MR-1 strains were grown in LB media overnight, washed twice

with SDM, and diluted to an OD_{600} of 0.05 in 1 mL of SDM with appropriate antibiotics.

Prior to pipetting, ~1cm of the pipette tip was trimmed to minimize shear forces during

transfer. 100 µL of each culture was labeled with 1 µL 1M FM 4-64 to visualize the cell

membranes. After staining, 10 µL of the labeled cell suspension was gently pipetted onto

22 x 22 mm No.1 cover glass (VWR) and sealed onto glass slides with clear acrylic nail

polish for confocal imaging or onto Lab-Tek chambered #1 cover glass (Thermo Fischer

Scientific)(for widefield fluorescence).   On average, intrinsic membrane extension

formation could be observed starting after 45 minutes sealed onto cover glass. For

planktonic OME production from BdpA induction in trans, diluted cells (in either SDM for

S. oneidensis, BB for *M. atlanticus*, or LB for *E. coli*) were induced with 12.5 µM DAPG

for 1 hour at 30˚C with 200 RPM shaking agitation. Cells were labeled with FM 4-64 and

sealed onto glass slides as before. Induced OMEs were imaged immediately after

mounting onto slides.

Confocal images were taken by a Zeiss LSM 800 confocal microscope with a Plan-

Apochromat 63x/1.4 numerical aperture oil immersion M27 objective. FM 4-64

fluorescence was excited at 506 nm: 0.20% laser power. Emission spectra was detected

from 592-700 nm using the LSM 800 GaAsP-Pmt2 detector. To capture the dynamics of

the OMEs, images were collected over the designated length of time between 0.27 – 0.63

seconds per frame. Single frame time series images were collected of either a 50.71 µm

by 50.71 µm (2x zoom) or a 20.28 µm by 20.28 µm (5x zoom) field of view. Widefield

fluorescence images were taken using a LED-Module 511 nm light source at 74.2%

intensity with 583-600 nm filters and a 91 He CFP/YFP/mCherry reflector. Excitation and

emission spectra were 506 nm and 751 nm, respectively. Images were collected using a

Hamamatsu camera with a 250 ms exposure time. Images were recorded using the Zeiss

Zen software (Carl Zeiss Microscopy, LLC, Thornwood, NY, USA). Frequency of OMEs

was calculated from 3 biological replicates of each strain from at least 5 fields of view

selected at random (2444 cells from WT, 4378 from Δ*bdpA*, and 3354 from Δ*bdpA* p452-

bdpA). OMEs were counted per cell per field of view from a 20 second video if a

membrane extension was observed extending from the cell over that timeframe.

Statistical significance was determined by two-tailed Student's *t*-test.

Perfusion flow microscopy For OME statistics comparing *S. oneidensis* strains MR-1 and Δ*bdpA,* cells were pre-grown aerobically from frozen (-80°C) stock in 10 mL of Luria-Bertani (LB) broth (supplemented with 50 µg/mL Kanamycin for strains with plasmid) in a 125-mL flask overnight at 30°C and 225 rpm. The next day, the stationary phase (OD_{600} 3.0 – 3.3) preculture was used to inoculate 1:100 into 10 mL of fresh LB medium in a 125-mL flask. After ~6 hours at 30°C and 225 rpm, when the OD_{600} was 2.4 (late log phase), 5 mL of cells were collected by centrifugation at 4226 x *g* for 5 min and washed twice in defined medium. The perfusion chamber, microscope, and flow medium described previously[7, 29, 30] were used for all perfusion flow OME statistics experiments. During each 5 hour imaging experiment, the perfusion chamber was first filled with this flow medium, then <1 mL of washed cells were slowly injected for a surface density of ~100-300 cells per 112 x 112 µm field of view on a Nikon Eclipse Ti-E inverted microscope with the NIS-Elements AR software. Cells were allowed to attach for 5-15 minutes on the coverslip before perfusion flow was resumed at a volumetric flow rate of 6.25 ± 0.1 µL/s. Cells and OMEs were visualized with the red membrane stain FM 4-64FX in the flow medium (0.25 µg/mL of flow medium). A total of 1,831 wild type and 2,265 Δ*bdpA* cells were used for extension and vesicle quantification. Experiments were performed from three individual biological replicates for each strain, and statistical significance was determined by two-tailed χ^2 test.

Cryo transmission electron microscopy

Shewanella strains were streaked onto LB plates with or without kanamycin and allowed to incubate 3 days on a benchtop. The night before freezing, individual colonies were inoculated into 3 ml LB +/- kanamycin and incubated at 30 °C overnight with 200 rpm

shaking. The following morning optical densities of the cultures were measured at

600nm and adjusted to a final OD_{600} of 1. Cells were pelleted at 8,000 rpm for three

minutes for buffer exchange/washes. For the Δ*bdpA* p452-*bdpA* transformed cells, 12.5

μM DAPG was added. A freshly glow discharged 200 mesh copper grid with R2/1

Quantifoil carbon film was placed into a concavity slide. Approximately 150 μl of a 1:10

dilution of the cell suspensions, with or without the inducer, was added to cover the grid.

A glass coverslip was then lowered onto the concavity to exclude air bubbles. The

edges of the coverslip were then sealed with nail polish to prevent media evaporation.

The slide assembly was then incubated in a 30 °C incubator for 1.5 to 3 hours.

Immediately prior to plunge freezing, the top coverslip was removed by scoring the nail

polish with a razor blade. TEM grids with cells were gently retrieved with forceps and

loaded into a Leica grid plunge for automated blotting and plunging into LN_2-cooled

liquid ethane. Vitrified grids were transferred to a LN_2 storage dewar. Imaging of frozen

samples was performed on either a Titan (ThermoFisher Scientific) microscope

equipped with a Gatan Ultrascan camera and operating at 300 kV or a Talos

(ThermoFisher Scientific) equipped with a Ceta camera and operating at 200 kV.

Images were acquired at 10,000x to 20,000x magnification and were adjusted by

bandpass filtering. Unfixed OMEs were sorted based on appearance into 4 categories.

Tubules were narrow OMEs with relatively uniform or slight symmetric curvature.

Narrow chains were recorded as OMEs with a narrow, consistent diameter and

symmetric curvature at constriction points. Irregular chains were classified as OMEs

without a consistent diameter throughout the length of the OME and asymmetric

curvature on either side of the extension. Blebs \ bulges were outer membrane

structures that did not resemble OMEs but still extended from the cell membrane

surface. Phenotypes were documented from observations of 31 WT, 13 Δ*bdpA*, and 3

Δ*bdpA* p452-*bdpA* OMEs over three separate biological replicates, with two technical

replicates of each strain per biological replicate.

Acknowledgements

We thank Dr. Jeffery Gralnick for helpful discussions and advice; Dr. Adam Meyer

and Dr. Chris Voigt for the DAPG-inducible Marionette promoter; Dr. Annie Moradian and

Dr. Mike Sweredoski and the California Institute of Technology Proteome Exploration Lab

for useful discussions on the preparation and analysis of proteomics data. Some of the

cryo-TEM work was done in the Beckman Institute Resource Center for Transmission

Electron Microscopy at Caltech. This work was supported by the United States

Department of Defense Synthetic Biology for Military Environments (SBME) Applied

Research for the Advancement of Science and Technology Priorities (ARAP) program.

Work in ME-N's lab was supported by the U.S. Office of Naval Research Multidisciplinary

University Research Initiative Grant No. N00014-18-1-2632. LAZ was partially supported

by the National Science Foundation grant DEB-1542527. SX was supported by the

Division of Chemical Sciences, Geosciences, and Biosciences, Office of Basic Energy

Sciences of the U.S. Department of Energy through grant DE-FG02-13ER16415. Work

in GJJ's lab was supported by the National Institute of Health (GM122588 to GJJ).

Author Contributions

DP and LZ conceived the study independently then combined projects when

complementary data on BdpA was discovered.　LZ purified OMVs, prepared samples for

LC MS-MS, and performed DLS measurements.　LZ and SX made electrochemical

measurements and analysis.　DP conducted BdpA domain prediction and validation

analysis, generated the p452-*bdpA* plasmid, Δ*bdpA* and p452-*bdpA* strains. DP and GC

conducted fluorescence imaging experiments, and DP, LZ, and GC analyzed the data.

LB adapted the Marionette sensor (P_{phlF}-YFP) into pBBR1-mcs2.　LZ and LAM performed

cryo-TEM of OMVs.　GC and LZ analyzed perfusion flow system data.　CH, DP, and LD

performed cryo-TEM experiments of OMEs and image processing / analysis.　DP and AM

generated phylogenetic data, and DP, AM, and BE analyzed the data.　DP, LZ, CM, GC,

AM, LAM, BE, GJJ, LD, MEN, and SG provided data interpretation.　DP, LZ, MEN, and

SG wrote the manuscript, with input from all coauthors.

References

1.   Carman PJ & Dominguez R (2018) BAR domain proteins-a linkage between cellular membranes,
signaling pathways, and the actin cytoskeleton. *Biophys Rev* 10(6):1587-1604.
2.   Schoneberg J*, et al.* (2017) Lipid-mediated PX-BAR domain recruitment couples local membrane
constriction to endocytic vesicle fission. *Nat Commun* 8:15873.
3.   van Weering JR & Cullen PJ (2014) Membrane-associated cargo recycling by tubule-based
endosomal sorting. *Semin Cell Dev Biol* 31:40-47.
4.   Knaevelsrud H*, et al.* (2013) Membrane remodeling by the PX-BAR protein SNX18 promotes
autophagosome formation. *J Cell Biol* 202(2):331-349.
5.   Mim C & Unger VM (2012) Membrane curvature and its generation by BAR proteins. *Trends
Biochem Sci* 37(12):526-533.
6.   Rao Y & Haucke V (2011) Membrane shaping by the Bin/amphiphysin/Rvs (BAR) domain protein
superfamily. *Cell Mol Life Sci* 68(24):3983-3993.
7.   Chong GW, Pirbadian S, & El-Naggar MY (2019) Surface-Induced Formation and Redox-
Dependent Staining of Outer Membrane Extensions in Shewanella oneidensis MR-1. *Frontiers in
Energy Research* 7.
8.   Fischer T*, et al.* (2019) Biopearling of Interconnected Outer Membrane Vesicle Chains by a
Marine Flavobacterium. *Appl Environ Microbiol* 85(19).
9.   Toyofuku M, Nomura N, & Eberl L (2018) Types and origins of bacterial membrane vesicles.
Nature Reviews Microbiology.
10.   Hampton CM*, et al.* (2017) The Opportunistic Pathogen Vibrio vulnificus Produces Outer
Membrane Vesicles in a Spatially Distinct Manner Related to Capsular Polysaccharide. *Front
Microbiol* 8:2177.
11.   Kim EY, Tyndall ER, Huang KC, Tian F, & Ramamurthi KS (2017) Dash-and-Recruit Mechanism
Drives Membrane Curvature Recognition by the Small Bacterial Protein SpoVM. *Cell Syst
5(5):518-526 e513.
12.   Cornejo E, Subramanian P, Li Z, Jensen GJ, & Komeili A (2016) Dynamic Remodeling of the
Magnetosome Membrane Is Triggered by the Initiation of Biomineralization. *MBio* 7(1):e01898-
01815.
13.   Bohuszewicz O, Liu J, & Low HH (2016) Membrane remodelling in bacteria. *J Struct Biol* 196(1):3-
14.
14.   Remis JP*, et al.* (2014) Bacterial social networks: structure and composition of Myxococcus
xanthus outer membrane vesicle chains. *Environ Microbiol* 16(2):598-610.
15.   Wei X, Vassallo CN, Pathak DT, & Wall D (2014) Myxobacteria produce outer membrane-
enclosed tubes in unstructured environments. *J Bacteriol* 196(10):1807-1814.
16.   Low HH, Sachse C, Amos LA, & Lowe J (2009) Structure of a bacterial dynamin-like protein lipid
tube provides a mechanism for assembly and membrane curving. *Cell* 139(7):1342-1352.
17.   Peter BJ*, et al.* (2004) BAR domains as sensors of membrane curvature: the amphiphysin BAR
structure. *Science* 303(5657):495-499.
18.   Frost A, De Camilli P, & Unger VM (2007) F-BAR proteins join the BAR family fold. *Structure
15(7):751-753.
19.   Shimada A*, et al.* (2007) Curved EFC/F-BAR-domain dimers are joined end to end into a filament
for membrane invagination in endocytosis. *Cell* 129(4):761-772.

20.   Weissenhorn W (2005) Crystal structure of the endophilin-A1 BAR domain. *J Mol Biol*
351(3):653-661.
21.   Drin G & Antonny B (2010) Amphipathic helices and membrane curvature. *FEBS Lett*
584(9):1840-1847.
22.   Seet LF & Hong W (2006) The Phox (PX) domain proteins and membrane traffic. *Biochim Biophys*
Acta 1761(8):878-896.
23.   Itoh T & De Camilli P (2006) BAR, F-BAR (EFC) and ENTH/ANTH domains in the regulation of
membrane-cytosol interfaces and membrane curvature. *Biochim Biophys Acta* 1761(8):897-912.
24.   Simunovic M, Voth GA, Callan-Jones A, & Bassereau P (2015) When Physics Takes Over: BAR
Proteins and Membrane Curvature. *Trends Cell Biol* 25(12):780-792.
25.   Mim C*, et al.* (2012) Structural basis of membrane bending by the N-BAR protein endophilin. *Cell*
149(1):137-145.
26.   Habermann B (2004) The BAR-domain family of proteins: a case of bending and binding? *EMBO*
reports 5(3):250-255.
27.   Miki H, Yamaguchi H, Suetsugu S, & Takenawa T (2000) IRSp53 is an essential intermediate
between Rac and WAVE in the regulation of membrane ruffling. *Nature* 408(6813):732-735.
28.   Sampath V, McCaig WD, & Thanassi DG (2018) Amino acid deprivation and central carbon
metabolism regulate the production of outer membrane vesicles and tubes by Francisella. *Mol*
Microbiol 107(4):523-541.
29.   Subramanian P, Pirbadian S, El-Naggar MY, & Jensen GJ (2018) Ultrastructure of Shewanella
oneidensis MR-1 nanowires revealed by electron cryotomography. *Proc Natl Acad Sci U S A*
115(14):E3246-E3255.
30.   Pirbadian S*, et al.* (2014) Shewanella oneidensis MR-1 nanowires are outer membrane and
periplasmic extensions of the extracellular electron transport components. *Proc Natl Acad Sci U*
S A 111(35):12883-12888.
31.   Gorby YA*, et al.* (2006) Electrically conductive bacterial nanowires produced by Shewanella
oneidensis strain MR-1 and other microorganisms. *Proc Natl Acad Sci U S A* 103(30):11358-
11363.
32.   Bhattacharya S*, et al.* (2019) A Ubiquitous Platform for Bacterial Nanotube Biogenesis. *Cell Rep*.
33.   Dubey GP*, et al.* (2016) Architecture and Characteristics of Bacterial Nanotubes. *Dev Cell*
36(4):453-461.
34.   Dubey GP & Ben-Yehuda S (2011) Intercellular nanotubes mediate bacterial communication. *Cell*
144(4):590-600.
35.   Pande S*, et al.* (2015) Metabolic cross-feeding via intercellular nanotubes among bacteria. *Nat*
Commun 6:6238.
36.   Tanaka M, Arakaki A, & Matsunaga T (2010) Identification and functional characterization of
liposome tubulation protein from magnetotactic bacteria. *Mol Microbiol* 76(2):480-488.
37.   Danne L*, et al.* (2017) Membrane Remodeling by a Bacterial Phospholipid-Methylating Enzyme.
MBio 8(1).
38.   Danne L*, et al.* (2017) Dissection of membrane-binding and -remodeling regions in two classes of
bacterial phospholipid N-methyltransferases. *Biochim Biophys Acta* 1859(12):2279-2288.
39.   Danne L*, et al.* (2015) Membrane-binding mechanism of a bacterial phospholipid N-
methyltransferase. *Mol Microbiol* 95(2):313-331.
40.   Low HH & Lowe J (2006) A bacterial dynamin-like protein. *Nature* 444(7120):766-769.
41.   Gorby Y*, et al.* (2008) Redox-reactive membrane vesicles produced by Shewanella. *Geobiology*
6(3):232-241.
42.   Liu X*, et al.* (2019) Bacterial Vesicles Mediate Extracellular Electron Transfer. *Environmental*
Science & Technology Letters.

43. Xu S, Barrozo A, Tender LM, Krylov AI, & El-Naggar MY (2018) Multiheme Cytochrome Mediated
Redox Conduction through Shewanella oneidensis MR-1 Cells. *J Am Chem Soc* 140(32):10085-
10089.

44. Okamoto A, Nakamura R, & Hashimoto K (2011) In-vivo identification of direct electron transfer
from Shewanella oneidensis MR-1 to electrodes via outer-membrane OmcA–MtrCAB protein
complexes. *Electrochimica Acta* 56(16):5526-5531.

45. Cox J*, et al.* (2014) Accurate proteome-wide label-free quantification by delayed normalization
and maximal peptide ratio extraction, termed MaxLFQ. *Mol Cell Proteomics* 13(9):2513-2526.

46. Hsu VW, Bai M, & Li J (2012) Getting active: protein sorting in endocytic recycling. *Nat Rev Mol*
Cell Biol 13(5):323-328.

47. Vincent TL, Green PJ, & Woolfson DN (2013) LOGICOIL--multi-state prediction of coiled-coil
oligomeric state. *Bioinformatics* 29(1):69-76.

48. Linkner J*, et al.* (2014) The inverse BAR domain protein IBARa drives membrane remodeling to
control osmoregulation, phagocytosis and cytokinesis. *J Cell Sci* 127(Pt 6):1279-1292.

49. Cui H*, et al.* (2013) Understanding the role of amphipathic helices in N-BAR domain driven
membrane remodeling. *Biophys J* 104(2):404-411.

50. Henne WM*, et al.* (2007) Structure and analysis of FCHo2 F-BAR domain: a dimerizing and
membrane recruitment module that effects membrane curvature. *Structure* 15(7):839-852.

51. Roy A, Kucukural A, & Zhang Y (2010) I-TASSER: a unified platform for automated protein
structure and function prediction. *Nat Protoc* 5(4):725-738.

52. Moravcevic K*, et al.* (2015) Comparison of Saccharomyces cerevisiae F-BAR domain structures
reveals a conserved inositol phosphate binding site. *Structure* 23(2):352-363.

53. Pylypenko O, Lundmark R, Rasmuson E, Carlsson SR, & Rak A (2007) The PX-BAR membrane-
remodeling unit of sorting nexin 9. *EMBO J* 26(22):4788-4800.

54. Vinogradov E, Korenevsky A, & Beveridge TJ (2003) The structure of the rough-type
lipopolysaccharide from Shewanella oneidensis MR-1, containing 8-amino-8-deoxy-Kdo and an
open-chain form of 2-acetamido-2-deoxy-d-galactose. *Carbohydrate Research* 338(19):1991-
1997.

55. Meyer AJ, Segall-Shapiro TH, Glassey E, Zhang J, & Voigt CA (2018) Escherichia coli "Marionette"
strains with 12 highly optimized small-molecule sensors. *Nat Chem Biol*.

56. Mahmood MI, Noguchi H, & Okazaki KI (2019) Curvature induction and sensing of the F-BAR
protein Pacsin1 on lipid membranes via molecular dynamics simulations. *Sci Rep* 9(1):14557.

57. Wang Q*, et al.* (2009) Molecular mechanism of membrane constriction and tubulation mediated
by the F-BAR protein Pacsin/Syndapin. *Proc Natl Acad Sci U S A* 106(31):12700-12705.

58. Bird LJ*, et al.* (2018) Development of a Genetic System for Marinobacter atlanticus CP1 (sp.
nov.), a Wax Ester Producing Strain Isolated From an Autotrophic Biocathode. *Front Microbiol*
9:3176.

59. Coursolle D & Gralnick JA (2010) Modularity of the Mtr respiratory pathway of Shewanella
oneidensis strain MR-1. *Mol Microbiol* 77(4):995-1008.

60. Bird LJ*, et al.* (2019) Engineered living conductive biofilms as functional materials. *MRS*
Communications:1-13.

61. Ayton GS*, et al.* (2009) New insights into BAR domain-induced membrane remodeling. *Biophys J*
97(6):1616-1625.

62. Simunovic M*, et al.* (2013) Protein-mediated transformation of lipid vesicles into tubular
networks. *Biophys J* 105(3):711-719.

63. Noguchi H (2016) Membrane tubule formation by banana-shaped proteins with or without
transient network structure. *Sci Rep* 6:20935.

64.   Salzer U, Kostan J, & Djinovic-Carugo K (2017) Deciphering the BAR code of membrane
696   modulators. *Cell Mol Life Sci* 74(13):2413-2438.

697   65.   Perez-Cruz C*, et al.* (2013) New type of outer membrane vesicle produced by the Gram-negative
698   bacterium Shewanella vesiculosa M7T: implications for DNA content. *Appl Environ Microbiol*
699   79(6):1874-1881.

700   66.   Hagen WJH, Wan W, & Briggs JAG (2017) Implementation of a cryo-electron tomography tilt-
701   scheme optimized for high resolution subtomogram averaging. *J Struct Biol* 197(2):191-198.

702   67.   Rohou A & Grigorieff N (2015) CTFFIND4: Fast and accurate defocus estimation from electron
703   micrographs. *J Struct Biol* 192(2):216-221.

704   68.   Kremer JR, Mastronarde DN, & McIntosh JR (1996) Computer visualization of three-dimensional
705   image data using IMOD. *J Struct Biol* 116(1):71-76.

706   69.   Brown RN, Romine MF, Schepmoes AA, Smith RD, & Lipton MS (2010) Mapping the subcellular
707   proteome of Shewanella oneidensis MR-1 using sarkosyl-based fractionation and LC-MS/MS
708   protein identification. *J Proteome Res* 9(9):4454-4463.

709   70.   Nielsen H (2017) Predicting Secretory Proteins with SignalP. *Protein Function Prediction:*
710   *Methods and Protocols*, ed Kihara D (Springer New York, New York, NY), pp 59-73.

711   71.   Altschul SF*, et al.* (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database
712   search programs. *Nucleic Acids Res* 25(17):3389-3402.

713   72.   Finn RD, Clements J, & Eddy SR (2011) HMMER web server: interactive sequence similarity
714   searching. *Nucleic Acids Res* 39(Web Server issue):W29-37.

715   73.   Le SQ & Gascuel O (2008) An improved general amino acid replacement matrix. *Mol Biol Evol*
716   25(7):1307-1320.

717   74.   Myers CR & Nealson KH (1988) Bacterial manganese reduction and growth with manganese
718   oxide as the sole electron acceptor. *Science* 240(4857):1319-1321.

719   75.   Saltikov CW & Newman DK (2003) Genetic identification of a respiratory arsenate reductase.
720   *Proc Natl Acad Sci U S A* 100(19):10983-10988.

721   76.   Kovach ME*, et al.* (1995) Four new derivatives of the broad-host-range cloning vector
722   pBBR1MCS, carrying different antibiotic-resistance cassettes. *Gene* 166(1):175-176.

723   77.   Simon R, Priefer U, & Pühler A (1983) A Broad Host Range Mobilization System for In Vivo
724   Genetic Engineering: Transposon Mutagenesis in Gram Negative Bacteria. *Bio/Technology*
725   1(9):784-791.

Fig. 1| Redox active vesicles are enriched with BAR domain protein BdpA. a.
Cryoelectron tomography image of *S. oneidensis* MR-1 outer membrane vesicles (OMVs)
(scale = 200 nm). **b**. Cyclic voltammetry of vesicles adhered to gold electrode via small
self-assembled monolayers, as diagramed. Inset shows first derivative of anodic scan. **c**.
Volcano plot of vesicle proteome compared to cell-associated outer membrane (red =
enriched in vesicles, blue = enriched in cell-associated outer membrane). **d**. Schematic
of BdpA domains.

Fig. 2| BdpA is responsible for maintaining vesicle size but does not alter the frequency of OMV or OME formation. a. Dynamic light scattering of OMV size distributions from the deletion strain (red, 9 biological replicates) compared to wild type (blue, 11 biological replicates) and Δ*bdpA* cells expressing *bdpA* from a plasmid (yellow, (left), with weighted averages of OMV diameters (right). Color opacity denotes overlapping data points ($p < 0.05$, *t*-test). **b**. Proportion of cells forming large vesicles (>300 nm diameter) during perfusion flow monitored by time lapse fluorescence imaging ($p < 0.00459$, χ^2 test). **c**. Proportion of cells forming vesicles and extensions relative to the total number of cells observed by fluorescence microscopy during perfusion flow ($p > 0.05$, χ^2 test). Time lapse images were recorded from 3 biological replicates per strain with 5-13 fields of view per replicate. Error bars represent standard deviation.

Fig. 3| BdpA promotes OME maturation into ordered tubules. a. Fluorescence images of *S. oneidensis* WT (top), Δ*bdpA* (middle), and Δ*bdpA* p452-*bdpA* with 12.5 µM DAPG (bottom) OMEs. Scale = 2 µm. **b.** Cryo-TEM images of WT (top), Δ*bdpA* (middle), and Δ*bdpA* p452-*bdpA* with 12.5 µM DAPG (bottom) OMEs. Scale = 100 nm. **c.** Representative cartoon of OME phenotypes. **d**. Pie charts show relative frequency of OME phenotypes observed from each strain. Membrane blebs/bulges were defined as non-structured membrane protrusions that did not resemble either of the other OME categories.

Fig. 4| Heterologous expression of BdpA promotes OME formation. **a.** Induction of BdpA expression with 12.5 µM DAPG during planktonic, non-attached growth results in OME formation in *S. oneidensis* (left), *M. atlanticus* CP1 (middle), and *E. coli* BL21(DE3) (right). Scale = 2 µm. **b.** Cryo-TEM image of OMEs following planktonic induction of BdpA expression in *S. oneidensis* WT p452-*bdpA* cells. Scale = 200 nm. Insets enlarged to show detail of regularly ordered electron densities at the surface of OME junctions (blue) and tubule regions (yellow). Scale = 50 nm. **c.** OME growth over time at 30 minute intervals of *E. coli* BL21(DE3) expressing BdpA while attached to a glass surface. Scale = 2 µm.

Fig. 5| Comparative phylogenetic analysis of BdpA with prokaryotic homologs and eukaryotic BAR domains. Maximum Likelihood evolutionary histories were inferred from 1000 bootstrap replicates, and the percentage of trees in which the taxa clustered together is shown next to the branches. Arrows indicate multiple branches collapsed to a single node. *S. oneidensis* BdpA and 5 prokaryotic orthologs (WP_011623497 – unclassified *Shewanella*, ESE40074 – *S. decolorationis* S12, WP_039978560 - *S. decolorationis*, KEK29176 – *S. xiamenensis*, and WP_055648003 – *Shewanella sp.* Sh95) predicted by the current BAR domain Pfam HMM to contain a BAR domain aligned with representative BAR domains from various BAR domain subtypes (N-BAR, F-BAR, SNX-BAR, I-BAR) at a total of 196 positions. The Gamma distribution used to model evolutionary rate differences among sites was 12.9598.

787   **Supplemental Table 2**

Strain, Plasmid, or Primer	Description or relevant genotype	Source or reference
Strains		
Shewanella oneidensis strains		
MR-1	Wild type	[(74)]
Δ*bdpA*	*bdpA* scarless deletion	This study
Marinobacter atlanticus CP1	Wild type	[(58)]
Eschericia coli strains		
BL21(DE3)	OneShot *E. coli* BL21(DE3)	Invitrogen
UQ950	Cloning strain	[(75)]
BW29427 (WM3064)	Conjugation strain	[(75)]
Plasmids		
pBBR1-mcs2	empty vector	[(76)]
pBBJM	Cloning backbone	This study
pSMV3	Suicide vector	[(77)]
pSMV3_1507KO	Contains up and downstream regions of SO_1507	This study
p452-*bdpA*	DAPG inducible *bdpA* vector	This study
Primers		
pAJMF2	TTAACGCGAATTTTAACAAAATATTAACGCcccgcttaacgatcgttggctg	
pAJMR3	AGCGGATAACAATTTCACACAGGAAACAGCTacctcagataaaatatttgc	
pBBRF3	gggctcatgagcaaatattttatctgaggtAGCTGTTTCCTGTGTGAAATTG	
pBBRR2	acccgcgctcagccaacgatcgttaagcgggGCGTTAATATTTTGTTAAAATTCGC	
1507F_insert	ttaatactagagaaagaggggaaatactagATGCGCACCGCTGC	
1507R_insert	gaggcctcttttctggaatttggtaccgagCTACATAAAGGCTTTAGTAAAGGCTT	
BBJMV_reverse	CAGCATTGAGATGACTGCAGCGGTGCGCATctagtatttcccctctttctctagtat	
BBJMV_forward	AAGGAAGCCTTTACTAAAGCCTTTATGTAGctcggtaccaaattccagaaaag	

1   Myers, C. R. & Nealson, K. H. Bacterial manganese reduction and growth with manganese oxide as the sole
electron acceptor. *Science* **240**, 1319-1321, (1988).
2   Bird, L. J. *et al.* Development of a Genetic System for *Marinobacter atlanticus* CP1 (sp. nov.), a Wax Ester
Producing Strain Isolated From an Autotrophic Biocathode. *Front Microbiol* **9**, 3176, (2018).
3   Saltikov, C. W. & Newman, D. K. Genetic identification of a respiratory arsenate reductase. *Proc Natl Acad*
Sci U S A **100**, 10983-10988, (2003).
4   Kovach, M. E. *et al.* Four new derivatives of the broad-host-range cloning vector pBBR1MCS, carrying
different antibiotic-resistance cassettes. *Gene* **166**, 175-176, (1995).
5   Simon, R., Priefer, U. & Pühler, A. A Broad Host Range Mobilization System for In Vivo Genetic
Engineering: Transposon Mutagenesis in Gram Negative Bacteria. *Bio/Technology* **1**, 784-791, (1983).

    Supplemental Figure 1

803 **Predicted structure of the BAR domain region of BdpA.**
(A) Ribbon diagram of i-TASSER prediction of BdpA structure as a dimeric molecule, residues 175-
502. Dimer was modeled from an alignment of BdpA monomers to homodimeric Hof1p (Protein
Data Bank [PDB] ID code 4WPE) structures that resulted in closest proximity of putative dimer
interface (purple) residues from the initial BAR domain prediction. (B) Surface representation of
predicted BdpA homodimer colored according to electrostatic potential. The bottom concave face
has an accumulation of distributed positively charged residues (Lys, Arg, His - blue), while the top
face has clusters of positive and negatively charged (Glu, Asp - red) residues, within a range of -3.0
to 3.0 V.

804 **Supplemental Figure 2**

806 Proportion of cells making OMEs without perfusion flow at three hours post deposition onto
807 chambered cover glass, measured from five random fields of view from each of three
808 independent cultures per strain.

Supplemental Figure 3

Cryo-TEM of *S. oneidensis* WT (A), Δ*bdpA* (B), and Δ*bdpA* p452-*bdpA* (C) OMEs at 90 minutes post-surface attachment. Scale = 100 nm.

816   Supplemental Figure 4

Growth of *S. oneidensis* strains in LB (top) or *Shewanella* Defined Medium (SDM) (bottom) in
response to DAPG exposure and BdpA induction. Error bars are standard deviation of three
biological replicates.

Supplemental Figure 5

Variability in OME phenotypes following BdpA induction in *M. atlanticus* CP1 p452-*bdpA*
cells. Cells displayed an array of membrane curvature phenotypes, ranging from short OMEs
(<2µm, A-C), long OMEs (2-10+ µm, D-F), membrane blebbing (C,F), and branched OME/OMV
chains (F). Scale = 2 µm.

    Supplemental Figure 6

BdpA has homologs in other bacterial species. A phylogenetic tree of the 23 BAR domain sequences that seed the BAR domain HMM predictions, *S. oneidensis* BdpA, and conserved BdpA orthologs in other prokaryotes. The protein regions corresponding to the BdpA BAR domain sequence from the 52 prokaryotic BdpA orthologs were aligned with the 23 representative eukaryotic BAR domain-containing proteins used to generate the BAR domain consensus sequence (cd07307) at a total of 222 positions. Maximum Likelihood evolutionary histories were inferred from 1000 bootstrap replicates, and the percentage of trees in which the taxa clustered together is shown next to the branches. The Gamma distribution used to model evolutionary rate differences among sites was 11.9548.

Supplementary Video 1 Widefield imaging of *S. oneidensis* WT cells 3h post-
deposition onto the surface of a chambered cover glass. Scale = 5 µm.

Supplementary Video 2 Widefield imaging of *S. oneidensis* Δ*bdpA* cells 3h post-
deposition onto the surface of a chambered cover glass. Scale = 5 µm.

Supplementary Video 3 Widefield imaging of *S. oneidensis* Δ*bdpA* p452-*bdpA* cells 3h
post-deposition onto the surface of a chambered cover glass. Scale = 5 µm.

Supplementary Video 4 Confocal imaging of *S. oneidensis* MR-1 p452-*bdpA* cells after
1h planktonic induction of BdpA with 12.5 µM DAPG. Scale = 5 µm.