1   **Understanding population structure**

2   **in an evolutionary context:**

3   **population-specific F_{ST} and pairwise F_{ST}**

5   Shuichi Kitada*, Reiichiro Nakamichi[†], and Hirohisa Kishino[‡]

*Tokyo University of Marine Science and Technology, Tokyo 108-8477, Japan

[†]Japan Fisheries Research and Education Agency, Yokohama 236-8648, Japan

[‡]Graduate School of Agriculture and Life Sciences, The University of Tokyo, Tokyo

113-8657, Japan

Short running title:

Population-specific and pairwise F_{ST}

Keywords:

Adaptation, demographic history, genetic diversity, migration, population structure

Corresponding:

Shuichi Kitada,

Tokyo University of Marine Science and Technology, Minato-ku, Tokyo 108-8477,

Japan

+81-297-45-5267

kitada@kaiyodai.ac.jp

Abstract

Populations are shaped by their history. Therefore, it is crucial to interpret population

structure in an evolutionary context. Pairwise F_{ST} measures population structure, whereas

population-specific F_{ST} measures deviation from the ancestral population. To understand

current population structure and a population's history of range expansion, we propose a

representation method that overlays population-specific F_{ST} estimates on an unrooted

neighbor-joining tree inferred from a pairwise F_{ST} distance matrix and on a map of sampling

locations. We examined the usefulness of our procedure by conducting simulations that

mimicked population colonization from an ancestral population and analyzing published

human, Atlantic cod, and wild poplar genotype data sets. Our results demonstrated that

population-specific F_{ST} values identify the source population and trace the evolutionary

history of its derived populations based on genetic diversity. In contrast, pairwise F_{ST} values

represent the current population structure. By integrating results of both estimators, we

obtained a new picture of current population structure that incorporates evolutionary history.

The generalized least squares of genome-wide population-specific F_{ST} indicated that the wild

poplar population expanded its distribution to the north where it adapted to longer day lengths,

to seashores where it adapted to abundant rainfall, and to the south where it adapted to dry

summers. Genomic data highlight the power of the bias-corrected moment estimators of F_{ST}.

All F_{ST} moment estimators described in this paper have reasonable CPU times and are useful

in population genomics studies. The R codes for our representation method and simulations

are available in the Supporting Information.

Quantifying genetic relationships among populations is of substantial interest in population

biology, ecology, and human genetics (Weir and Hill 2002). Appropriate estimates of

population structure are the basis of our understanding of biology and biological applications,

which vary from evolutionary and conservation studies to association mapping and forensic

identification (Weir and Hill 2002; Weir and Goudet 2017). For such objectives, Wright's F_{ST}

(Wright 1951) is commonly used to quantify the genetic divergence of populations and there

are many informative reviews on traditional and population-specific F_{ST} estimators (*e.g.*,

Excoffier 2001; Rousset 2001, 2004; Balloux and Lugon-Moulin 2002; Weir and Hill 2002;

Beaumont 2005; Holsinger and Weir 2009; Gaggiotti and Foll 2010; Bhatia et al. 2013; Weir

and Goudet 2017). Because inferences include methods of moment, maximum likelihood and

Bayesian estimation, those reviews tended to focus on theoretical perspectives. Therefore,

these issues are well understood, particularly among statistical and theoretical population

geneticists.

The traditional F_{ST} estimators were originally developed to estimate F_{ST} over a

metapopulation based on a set of population samples. In practice, this is often called global

F_{ST} (*e.g.*, Pérez-Lezaun et al. 1997). A moment estimator of F_{ST} (Weir & Cockerham in 1984)

and Analysis of molecular variance (Excoffier et al. 1992) have been widely used in various

fields. Additionally, F_{ST} values between pairs of population samples (pairwise F_{ST}) are

routinely used to estimate population structure in human genetics (Pérez-Lezaun et al. 1997;

Ramachandran et al., 2005), conservation biology (Palsbøll et al. 2007; Schwartz et al. 2007),

and evolutionary biology (*e.g.*, Hemmer-Hansen et al. 2013a; Therkildsen et al. 2013a;

Geraldes et al. 2014; McKown et al. 2014a; Jorde et al. 2015; Rougemont et al. 2020).

Divergent selection in an environmental gradient can impact population structure (Nosil et al.

2009; Orsini et al. 2013), and prior studies examined geographic distance and habitat

differences between populations as explanatory variables that impact population structure

estimated based on genome-wide (average over loci) pairwise F_{ST} values (*e.g.*, Rousset 1997;

Bradbury and Bentzen 2007; Petrou et al. 2014; Jorde et al. 2015; Kitada et al. 2017).

An alternative approach for estimating population differentiation is the use of

population-specific F_{ST} estimators (reviewed by Gaggiotti and Foll 2010). Model-based

Bayesian approaches, based on beta and/or Dirichlet distributions, for estimating

population-specific F_{ST} have been proposed (Balding and Nichols 1995; Nicholson *et al.*

2002; Falush *et al.* 2003; Beaumont and Balding 2004). Locus-population-specific F_{ST} was

developed to identify adaptive divergence of a gene among populations using empirical Bayes

(Beaumont and Balding 2004) and full Bayesian methods (BayeScan) (Foll and Gaggiotti

2006, 2008). These methods have been applied in many cases of various species (*e.g.*,

Therkildsen et al. 2013a; Geraldes et al. 2014; Limborg et al. 2012). In addition to the

model-based methods, bias-corrected moment estimators of population-specific F_{ST} have been

proposed (Weir and Hill 2002; Weir and Goudet 2017), which is expected to accurately reflect

population history and have a wide range of applications (Weir and Goudet 2017).

Despite the expected usefulness of population-specific F_{ST}, particularly in evolutionary

biology, applications have been sparce to date (*e.g.*, Nicholson *et al.* 2002; Weir *et al.* 2005;

Foll and Gaggiotti 2006; Buckleton *et al.* 2013; Rougemont et al. 2020). The main goal of our

paper was to demonstrate to empirical population geneticists how the two types of F_{ST}

(population-specific and pairwise) can be used in combination to help elucidate population

structure in the evolutionary context. In our approach, the current population structure is

represented by an unrooted neighbor-joining (NJ) tree (Saitou and Nei 1987) based on

pairwise F_{ST}. The colors on the terminal nodes based on the genome-wide population-specific

F_{ST} values enable inference of the historical order of population colonization. Moreover, we

present a representation on the geographical map. We first examined the usefulness of our

procedure by simulations that mimicked population colonization from a single ancestral

population with three scenarios of population range expansion. The three scenarios, one-, two-,

and three-directional population range expansion, reflect the empirical datasets of the human,

Atlantic cod (*Gadus morhua*), and wild poplar (*Populus trichocarpa*), respectively, and well

supported the results. Because the populations close to the ancestral population had more

opportunities for mutations than recently founded populations, they are likely to have the

highest heterozygosity and low values of population-specific F_{ST}. Consistently, our

simulations demonstrated that Weir and Goudet's (2017) (WG) population-specific F_{ST}

moment estimator correctly identified the source population and traced the evolutionary

history of its derived populations based on genetic diversity. The population structure

predicted by the pairwise F_{ST} moment estimator (Nei and Chesser 1983, NC83) was

consistent with the current population structure.

All F_{ST} estimators were computed using the R package FinePop2_ver.0.2, which is available

at CRAN. The R codes for our representation method exemplified by the human data and

simulations of population colonization used in this study are available in the Supporting

Information. They can be used for microsatellite and single-nucleotide polymorphism (SNP)

genotype data with Genepop format (Raymond and Rousset 1995; Rousset 2008), which has

been particularly widely used among biologists.

MATERIALS AND METHODS

Visual representation of population structure and demographic history

We integrated genome-wide population-specific and pairwise F_{ST} estimates on a NJ tree and

on a map of sampling locations. We drew the NJ tree based on the distance matrix of

genome-wide pairwise F_{ST} value using the `nj` function in the R package ape and

superimposed the magnitude of genome-wide population-specific F_{ST} values using a color

gradient on the NJ tree based on rgb $(1 - F_{ST,0}, 0, F_{ST,0})$, where $F_{ST,0} = (F_{ST} - \min F_{ST})/$

$(\max F_{ST} - \min F_{ST})$. This conversion represents the standardized magnitude of a

population-specific F_{ST} value at the sampling point, with colors between blue (for the largest

F_{ST}) and red (smallest F_{ST}). The F_{ST} maps were drawn using the `sf` package in R, where

sampling locations were plotted based on the longitudes and latitudes; they were visualized by

the population-specific F_{ST} color gradient, and the size of each sampling point was

proportional to the expected heterozygosity (H_e). Genome-wide population-specific F_{ST}

values were sorted by decreasing values and plotted with standard errors. Sampling points

with genome-wide pairwise F_{ST} values smaller than a given threshold were connected by

yellow lines to visualize the connectivity between populations. Under the assumption of

Wright's island model at equilibrium between drift, mutation, and migration (Wright 1931),

$F_{ST} \approx \frac{1}{4N_e m + 1}$, where N_e is the effective population size and m is the average rate of

migration between populations (see, Slatkin 1987). For a threshold of $F_{ST} = 0.02$,

$4N_e m \approx 49$ migrants per generation (also see, Whitlock and Mccauley 1999; Waples and

Gaggiotti 2006).

Population colonization simulations

To test the performance of our visual representation, we conducted simulations that mimicked

colonization of populations from a single ancestral population (population 1). We modeled

three types of colonization, one-, two-, and three-directional population expansion (Figure

1A–C), with 24 demes (populations 2–25). We set the effective population size of the

ancestral population to $N_e = 10^4$ (twice the number of individuals in diploid organisms in a

random mating population). At the beginning of colonization, 1% of N_e migrated into the

adjacent vacant habitat once every 10 generations. We considered one simulation cycle to be

one generation for convenience. The effective population size of the newly derived population

increased to the same size of the ancestral population ($N_e = 10^4$) after one generation, and

the population exchanged 1% of N_e genes with adjacent population(s) in every generation.

Like the ancestral population, 1% of N_e individuals migrated into the adjacent vacant habitat

once every 10 generations. We simulated the allele frequencies of SNPs in the ancestral and

24 derived populations.

The initial allele frequencies in the ancestral population, q, at 100,000 neutral SNP loci were

generated from the predictive equilibrium distribution, $f(q) \propto q^{-1}(1 - q)^{-1}$ (Wright 1931).

Additionally, 10 newly derived SNPs were introduced to each existing population in each

generation. Therefore, in total, 35,100 SNPs were generated. When a new SNP emerged in a

population, we set the initial allele frequency of the newly derived SNP to 0.01 in the

population and 0 in the other populations. This mimicked new mutations that survived at the

initial phase after their birth. These 100,000 ancestral SNPs and 35,100 newly derived SNPs

were considered "unobserved." The allele frequencies of these SNPs were changed by random

drift under a binomial distribution in every generation. Many of the SNPs had reduced

frequencies of the derived allele over simulation generations and lost their polymorphism.

After 260 simulation generations, SNPs that retained their polymorphism were randomly

selected as "observed" SNPs; in this simulation, we selected 10,000 ancestral SNPs and 500

newly derived SNPs. Then, we generated 50 individuals for each population. Genotypes of

these 10,500 SNPs were randomly generated for each individual following the allele

frequencies in the population to which each individual belonged. Thus, we obtained

"observed" genotypes of 1,250 individuals (= 50 individuals $\times$ 25 populations) at 10,500 SNP

loci (10,000 ancestral SNPs + 500 newly derived SNPs by mutation). We converted the

simulated genotypes into Genepop format (Raymond & Rousset, 1995; Rousset, 2008). We

computed genome-wide population-specific and pairwise F_{ST} values between 25 populations

and overlaid genome-wide population-specific F_{ST} values on the unrooted NJ tree.

Inferring environmental selection from observed population structure

Hereafter, notations consistent with those of Weir and Hill (2002) were used: i for populations

($i = 1, ..., r$), u for alleles ($u = 1, ..., m$), and l for loci ($l = 1, ..., L$). To infer the geography

and environment that were experienced by the population range expansion, we regressed the

genome-wide population-specific F_{ST} values on the geographical and environmental variables.

Residuals are correlated because of population structure; therefore, the effective sample size is

lower than the actual sample size. In such circumstances, ordinary least squares overestimate

the precision. To take the correlation into account, we derived the components of the

variance–covariance matrix of the population-specific F_{ST} estimator. It can be written as

$\mathrm{ps}\hat{F}_{ST}^i = \frac{\bar{y}^i}{\bar{x}}$. Using the Taylor series expansion for the first term, we inferred the asymptotic

variance as

$$V\left[\mathrm{ps}\hat{F}_{ST}^i\right] \simeq \frac{\bar{y}^2}{\bar{x}^2}\left\{\frac{V[\bar{x}]}{\bar{x}^2} + \frac{V[\bar{y}^i]}{\bar{y}^{i2}} - \frac{2Cov[\bar{x}, \bar{y}^i]}{\bar{x}\bar{y}^i}\right\} \quad (1)$$

194 Similarly, the asymptotic covariances between population-specific F_{ST} values of i, j

195   populations were obtained by

$$Cov\left[ps\hat{F}_{ST}^i, ps\hat{F}_{ST}^j\right] \simeq \frac{\bar{y}^i \bar{y}^j}{\bar{x}^4} V[\bar{x}] - \frac{\bar{y}^i}{\bar{x}^3} Cov\left[\bar{x}, \bar{y}^j\right] - \frac{\bar{y}^j}{\bar{x}^3} Cov\left[\bar{x}, \bar{y}^i\right], \quad (2)$$

where the variance and covariance components were calculated by

$V[\bar{x}] = \frac{1}{L(L-1)}\sum_{l=1}^{L}(x_l - \bar{x})^2$, $V[\bar{y}^i] = \frac{1}{L(L-1)}\sum_{l=1}^{L}(y_l^i - \bar{y}^i)^2$,

$Cov[\bar{x}, \bar{y}^i] = \frac{1}{L(L-1)}\sum_{l=1}^{L}(x_l - \bar{x})(y_l^i - \bar{y}^i)$ and

$Cov[\bar{x}, \bar{y}^j] = \frac{1}{L(L-1)}\sum_{l=1}^{L}(x_l - \bar{x})(y_l^j - \bar{y}^j).$

This analysis was performed on the wild poplar data set, for which 11

environmental/geographical parameters were available for each sampling location. We used

Equations 1 and 2 for the components of the variance–covariance matrix Ω of the GLS

function in FinePop2_ver.0.2. Once the key factors that are associated with the population

range expansion are identified, it would be interesting to search for the crucial genes that

enabled adaptation to the local environment by determining outliers of locus-specific,

population-specific F_{ST} values (Foll and Gaggiotti 2008; Coop et al. 2010).

F_{ST} estimators applied in this study

210   We used WG population-specific F_{ST} moment estimators, because we expected that

211   population-specific F_{ST} values accurately reflect population history (Weir and Goudet 2017).

212   In our analyses, we extended WG population-specific F_{ST} estimator to overall loci

213   (genome-wide population-specific F_{ST}), and the combined ratio estimator (Cochran 1977) for

214   overall loci (Buckleton et al. 2016) was:

$$ps\hat{F}_{ST}^i = \frac{\sum_{l=1}^{L}(\widetilde{M}_{W,l}^i - \widetilde{M}_l^B)}{\sum_{l=1}^{L}(1 - \widetilde{M}_l^B)}, \quad (3)$$

where $\widetilde{M}_W^i$ is the unbiased within-population matching of two distinct alleles of population i,

and $\widetilde{M}^B$ is the between-population-pair matching average over pairs of populations i, i'

(Supplemental Note). This is called the "ratio of averages" F_{ST} estimator, which

asymptotically converges to unbiased estimates of F_{ST} as the number of independent SNPs

increases (Weir and Cockerham 1984; Weir and Hill 2002; Bhatia et al. 2013).

We also applied empirical (Beaumont and Balding, 2004) and full Bayesian (Foll and

Gaggiotti, 2006) population-specific F_{ST} estimators. Beaumont and Balding (2004)

maximized the Dirichlet-multinomial marginal likelihood in their Equation 1 and estimated

θ_{li}:

$$L_{li}\left(\theta_{li} \middle| n_{li1}, \ldots, n_{lim_l}\right) = \frac{\Gamma(\theta_{li})}{\Gamma(N_{li} + \theta_{li})} \prod_{u=1}^{m_l} \frac{\Gamma(n_{liu} + \theta_{li}\bar{p}_{lu})}{\Gamma(\theta_{li}\bar{p}_{lu})} . \qquad (4)$$

Here, θ_{li} is the scale parameter of the Dirichlet prior distribution for locus l and population i,

$\tilde{p}_{lu}$ is the observed frequency of allele u at locus l, n_{liu} is the observed allele count in

population i, and N_{li} is the total number of alleles. Importantly, $\bar{p}_{lu}$ is the mean allele

frequency over all populations, whereas $\theta_{li}\bar{p}_{lu} = \alpha_{liu}$, where $\theta_{li} = \sum_{u=1}^{m_l} \alpha_{liu}$. The

parametrization reduces the number of parameters to be estimated. Based on a Dirichlet

(multi-allelic) and/or a beta (bi-allelic) scale parameter, population-specific F_{ST} values were

estimated for each locus using the following function of $\hat{\theta}_{li}$ (Beaumont and Balding, 2004):

$$ps\tilde{F}_{ST,l}^i = \frac{1}{\hat{\theta}_{li} + 1} . \qquad (5)$$

233   To estimate genome-wide pairwise F_{ST}, we used the NC83 bias-corrected G_{ST} moment

234   estimator:

$$\mathrm{pw}\hat{F}_{\mathrm{ST}} = \frac{\sum_{l=1}^{L}(\hat{H}_{T,l} - \hat{H}_{S,l})}{\sum_{l=1}^{L}\hat{H}_{T,l}}, \qquad (6)$$

where $\hat{H}_T$ and $\hat{H}_S$ are the unbiased estimators of total and within-population heterozygosity,

respectively (Supplemental Note). G_{ST} (Nei 1973) is defined "by using the gene frequencies at

the present population, so that no assumption is required about the pedigrees of individuals,

selection, and migration in the past" (Nei 1977). G_{ST} assumes no evolutionary history

(Holsinger and Weir 2009), whereas NC83 does not consider any population replicates (Weir

and Cockerham 1984; Excoffier 2001). The pairwise F_{ST} values obtained from NC83

therefore measured genetic differentiation among current populations based on a fixed set of

samples of populations, which were derived from an ancestral population. Our previous

coalescent simulations demonstrated that NC83 performs the best among F_{ST} estimators when

estimating pairwise F_{ST} values, particularly for larger numbers of loci (Kitada et al. 2017).

Computing F_{ST} values

We converted the genotype data into Genepop format (Raymond and Rousset 1995; Rousset

2008) for implementation in the R package FinePop2_ver.0.2. Genome-wide WG

population-specific F_{ST} (Equation 3) values were computed using the `pop_specificFST`

function. In addition to the "ratio of averages" (Weir and Cockerham 1984; Weir and Hill

2002) used for the F_{ST} functions in FinePop2, we computed the "average of ratios" (Bhatia et

al. 2013) of the WG population-specific F_{ST} for human data for comparison by averaging

locus-specific, population-specific F_{ST} values over loci. We maximized Equation 4 and

estimated the empirical Bayesian population-specific F_{ST} (Beaumont and Balding 2004) at

each locus according to Equation 5. We then averaged these values over all loci. For the full

Bayesian model, GESTE_ver. 2.0 (Foll and Gaggiotti 2006) was used to compute

genome-wide population-specific F_{ST} values. The Bayesian genome-wide population-specific

F_{ST} values were computed for human data to compare the results with that from the WG

population-specific F_{ST} moment estimator. F_{ST} is equal to G_{ST} for diploid random mating

populations (Excoffier 2001; also see Supplemental Note); therefore, NC83 pairwise F_{ST}

values (Equation 6) were computed using the `pop_pairwiseFST` function in FinePop2.

Expected heterozygosity was calculated for each population with the `read.GENEPOP`

function.

Three empirical data sets

As the first empirical data set, we chose microsatellite genotypes collected from human

populations worldwide, because human evolutionary history, migration, and population

structure have been particularly well studied (*e.g*., Diamond 1997; Rosenberg et al. 2002;

Ramachandran et al. 2005; Liu et al. 2006; Rutherford 2016; Nielsen et al. 2017). These

patterns are well known by statistical/theoretical population geneticists and biologists;

therefore, testing our new integrative F_{ST} analysis on this data set could provide a good

example of the usefulness of our practical approach. The human microsatellite data in

Rosenberg et al. (2002) were retrieved from

https://web.stanford.edu/group/rosenberglab/index.html. We converted the genotype data to

Genepop format (Raymond and Rousset 1995; Rousset 2008). We removed the Surui sample

(Brazil) from the data because that population was reduced to 34 individuals in 1961 as a

result of introduced diseases (Liu et al. 2006). We retained genotype data ($n = 1,035$) of 377

microsatellite loci from 51 populations categorized into six groups as in the original study: 6

populations from Africa, 12 from the Middle East and Europe, 9 from Central/South Asia, 18

from East Asia, 2 from Oceania, and 4 from America. Longitudes and latitudes of the

sampling sites were obtained from Cann et al. (2002) (Table S1).

An SNP data set was obtained from a commercially important fish, Atlantic cod, in the North

Atlantic. We combined the Atlantic cod genotype data of 924 SNPs common to 29

populations reported in Therkildsen et al. (2013a, b) and 12 populations in Hemmer-Hansen et

al. (2013a, b) in Genepop format. We compared genotypes associated with each marker in

samples that were identical between the two studies, namely, CAN08 and

Western_Atlantic_2008, ISO02 and Iceland_migratory_2002, and ISC02 and

Iceland_stationary_2002, and standardized the gene codes. We removed cgpGmo.S1035,

whose genotypes were inconsistent between the two studies. We also removed

cgpGmo.S1408 and cgpGmo.S893, for which genotypes were missing in several population

samples in Therkildsen et al. (2013b). Temporal replicates in Norway migratory, Norway

stationary, North Sea, and Baltic Sea samples were removed for simplicity. The final data set

consisted of genotype data (n = 1,065) at 921 SNPs from 34 populations: 3 from Iceland, 25

from Greenland, 3 from Norway, and 1 each from Canada, the North Sea, and the Baltic Sea.

The genotype data included historical samples collected 50–80 years ago and contemporary

samples from the northern range margin of the species in Greenland, Norway, and the Baltic

Sea. The inclusion of both types of data might facilitate detection of migration history of this

highly migratory marine fish. Two ecotypes (migratory and stationary) that were able to

interbreed but were genetically differentiated (Hemmer-Hansen et al., 2013a; Berg et al.,

2016) were included in the Norway and Iceland samples. All individuals in the samples were

adults, and most were mature (Therkildsen et al., 2013a). The longitudes and latitudes of the

sampling sites in Hemmer-Hansen et al. (2013a) were used. For the data from Therkildsen et

al. (2013a), approximate sampling points were estimated from the map of the original study,

and longitudes and latitudes were recorded (Table S2).

The other SNP data set was from a tree, wild poplar, in the American Pacific Northwest. The

SNP genotype data and environmental/geographical data were retrieved from the original

studies of McKown et al. (2014a, b). The genotype data contained 29,355 SNPs of 3,518

genes of wild poplar ($n = 441$) collected from 25 drainage areas (McKown et al., 2014c).

Details of array development and selection of SNPs are provided in Geraldes et al. (2011,

2013). We converted the data to Genepop format. The samples covered various regions over a

range of 2,500 km near the Canadian–US border at altitudes between 0 and 800 m. A

breakdown of the 25 drainages (hereafter, populations) is as follows: 9 in northern British

Colombia (NBC), 2 in inland British Colombia (IBC), 12 in southern British Colombia (SBC),

and 2 in Oregon (ORE) (Geraldes et al., 2014). The original names of clusters and population

numbers were combined and used for our population labels (NBC1, NBC3,…, ORE30). Each

sampling location was associated with 11 environmental and geographical parameters:

latitude (lat), longitude (lon), altitude (alt), longest day length (DAY), frost-free days (FFD),

mean annual temperature (MAT), mean warmest month temperature (MWMT), mean annual

precipitation (MAP), mean summer precipitation (MSP), annual heat-moisture index (AHM),

and summer heat-moisture index (SHM) (Table S3). The AHM was calculated in the original

study as (MAT+10)/(MAP/1000); a large AHM indicates extremely dry conditions.

RESULTS

Simulations of population colonization

In the one-directional simulation, our method correctly identified the ancestral population

with the highest genetic diversity, and populations were located in order from 1 to 25 on the

NJ tree (Figure 1D). In the two-directional simulation, our method correctly identified the

ancestral population and detected that populations were split at population 9 and expanded in

two directions, which was consistent with the simulation scenario (Figure 1E). In the

three-directional simulation, the ancestral population was closely located to the adjacent

populations 2, 9, and 17, but correctly detected three directions, as in the other simulations

(Figure 1F).

Our simulation results demonstrated that our method represents insight into current

population structure that incorporates evolutionary history. Our results revealed that WG

population-specific F_{ST} values (standardized by a color gradient) identified the source

population and traced the evolutionary history of its derived populations based on genetic

diversity. The NC83 pairwise F_{ST} estimator correctly estimated the current population

structure. Genome-wide WG population-specific F_{ST} values were negative in the ancestral

population and adjacent populations, and the phenomenon was particularly significant in the

one- and two-directional models, whereas no negative population-specific F_{ST} values were

obtained in the ancestral and adjacent populations in the three-directional model (Figures

1G-I). In contrast, H_e values were larger in the ancestral population in the one- and

two-directional models than in the three-directional model, though the variation in H_e values

was not substantial because of the relatively few generations (260) in the simulation (Figures

1D-F). Our simulation results indicated that, when gene flow from other populations into the

source population was limited, relatively large H_e could be maintained, which resulted in

substantial negative population-specific F_{ST} values. Very similar results were obtained from

different data with more than 20 simulations (Figures not shown).

Humans

The map (Figure 2A) shows integrated information from genome-wide population-specific

and pairwise F_{ST} and H_e, which revealed population structure with migration and genetic

diversity of human populations. Interestingly, Bantu Kenyans had the smallest F_{ST} value (as

shown in red). Figure 2B ordered population-specific F_{ST} values from Africa to Central/South

Asia, the Middle East, Europe, East Asia, Oceania, and America (Table S4). As indicated by

sampling points connected by yellow lines with pairwise F_{ST} values below the 0.02 threshold

(Figure 2A), genetic connectivity from Africa was low. In contrast, migration was substantial

within Eurasia but was much smaller than that inferred from Eurasia to Oceania and America.

As illustrated by sampling point radii (Figure 2A), H_e was the highest in Africa, followed by

the Middle East, Central/South Asia, Europe, and East Asia, but relatively small in Oceania

and lowest in South America. The Karitiana in Brazil had the lowest H_e. The NJ tree (Figure

2C) integrated with population-specific F_{ST} values revealed that human populations originated

from Bantu Kenyans and expanded to Europe through Mozabite, the Middle East,

Central/South Asia, and East Asia. The Kalash was isolated from Europe/Middle East and

Central/South Asia populations. Papuans/Melanesians and American populations diverged

from between Central/South Asian and East Asian populations. The ordinal NJ tree of

pairwise F_{ST} values divided the populations into five clusters: 1) Africa, 2) the Middle East,

Europe, and Central/South Asia, 3) East Asia, 4) Oceania, and 5) America (Figure S1).

Bayesian population-specific F_{ST} values estimated using the methods of Beaumont and

Balding (2004) (empirical Bayes) and Foll and Gaggiotti (2006) (full Bayes) were nearly

identical; however, in African populations, they were higher than WG population-specific F_{ST}

values (Figure S2). The distributions of F_{ST} values obtained from the two Bayesian methods

were very similar, with the smallest F_{ST} values observed in the Middle East, Europe, and

Central/South Asia (Figure S2, Table S4). The "ratio of averages" and "average of ratios" of

the WG population-specific F_{ST} estimator were almost identical in all populations for this data

set (Figure S3).

Atlantic cod

The F_{ST} map (Figure 3A) visualized population structure, migration, and genetic diversity of

the Atlantic cod populations. The Canadian population had the smallest population-specific

F_{ST} value (as shown in red) and the greatest H_e (indicated by circle radii). As shown by

sampling point radii (Figure 3A), H_e was greatest in Canada. It was also high in Greenland,

low in other areas, and lowest in the Baltic Sea. Figure 3B ordered population-specific F_{ST}

values from Canada to the Baltic sea (Table S5). Greenland west coast populations (green in

Figure S4) generally had small population-specific F_{ST} values, whereas fjord populations had

relatively higher population-specific F_{ST} values. The F_{ST} values were much higher for

populations in Iceland, Norway, and the North Sea, and highest in the Baltic Sea. Based on

pairwise F_{ST} values between sampling points (< 0.02 threshold) (Figure 3A), substantial

migration was detected between Greenland, Iceland, and Norway. In contrast, migration was

low from Canada to Greenland and from Iceland and Norway to the North and Baltic Seas.

Our integrated NJ tree (Figure 3C) with population-specific F_{ST} values indicated that Atlantic

cod originated from Canada, migrated to the west coast of Greenland, and then expanded their

distribution to Iceland, Norway, the North Sea, and the Baltic Sea.

Atlantic cod might migrate to find new habitat when they reach carrying capacity, and

individuals with genomic variation were able to adapt to changing environments and formed

the current local populations. According to the ordinal NJ tree of the pairwise F_{ST} distance

matrix (Figure S4), the populations were divided into four large clusters: 1) Canada, 2)

Greenland west coast, 3) Greenland east coast, Iceland, Norway, and 4) North and Baltic Seas.

Fjord populations (in purple) formed a sub-cluster within the Greenland west coast, and

migratory (orange) and stationary (magenta) ecotypes also formed a sub-cluster.

Wild poplar

The F_{ST} map (Figure 4A) indicated that population-specific F_{ST} values were lowest in SBC27,

IBC15, and IBC16 (as shown in red). Sampling points connected by yellow lines (< 0.02

threshold pairwise F_{ST} values) indicated migration between all populations. As shown by

sampling point radii (Figure 4A), H_e was highest in SBC27, IBC15, and IBC16, and lowest

in NBC5. Figure 4B showed that samples collected from areas close to the SBC coast had

higher population-specific F_{ST} values than other SBC samples (Figure S5, Table S6). NBC

samples had population-specific F_{ST} values similar to those of SBC. Among NBC samples,

NBC8 had the smallest population-specific F_{ST}, and NBC5 had the highest value, followed by

NBC6 and NBC7. The pairwise F_{ST} NJ tree integrated with population-specific F_{ST} values

(Figure 4C) indicated that wild poplar originated from Inner BC and SBC27, and expanded in

three directions, namely, to the BC southern coast, northern BC and south-western Alaska,

and Oregon. The ordinal NJ tree based on the pairwise F_{ST} distant matrix divided populations

into three large clusters: 1) IBC, 2) SBC, 3) NBC, and 4) ORE (Figure S7). The population

represented by sample ORE30 was isolated from ORE29.

To avoid multicollinearity, we excluded seven out of 11 environmental variables that were

significantly correlated with each other, namely, lat, lon, alt, FFD, MWMT, MSP, and AHM.

Our GLS of genome-wide population-specific F_{ST} values on the four environmental variables

(DAY, MAT, MAP, and SHM) indicated that DAY, MAP, and SHM were significant (Table 1).

All estimates were positive, which indicated that higher population-specific F_{ST} values were

expected for longer DAY (longer daylight time), higher MAP (abundant rain), and higher

SHM (dry summers), and these values reflected directions of population expansion. The

scatter plot of DAY and SHM (each population colored by the population-specific F_{ST} value)

(Figure 5A) suggested three directions of population range expansion; the wild popular that

originated from IBC15 expanded its distribution to NBC, where it adapted to longer DAY;

that which originated from SBC27 expanded to SBC seashores, where it adapted to lower

SHM, and to ORE29 and ORE30, where it adapted to higher SHM. This was consistent in the

scatter plot of DAY and MAP (Figure 5B), which suggested that the expansion in SBC could

have been facilitated by adaptation to higher MAP.

CPU times

Using a laptop computer with an Intel Core i7-8650U CPU, only 89.8 s of CPU time was

required to compute WG population-specific F_{ST} estimates and SEs of wild poplar (29,355

SNPs; 25 populations, $n = 441$). Alternatively, 120.7 s was required to obtain pairwise F_{ST}

(NC83) between all population pairs. Based on the results, we may need 50 min to compute

WG population-specific F_{ST} and 70 min to compute pairwise NC83 F_{ST} estimates for 1

million SNPs using this laptop. This computation could be much faster if we used a

workstation.

Data availability

The authors affirm that all data necessary for confirming the conclusions of the article are

present within the article, figures, a table, and supplemental information.

DISCUSSION

Genome-wide population-specific F_{ST} traced population history as reflected by genetic diversity

Our simulations demonstrated that the WG population-specific F_{ST} estimator was able to identify the source population and trace the evolutionary history of its derived populations based on genetic diversity. We also found that the NC83 pairwise F_{ST} estimator correctly estimated current population structure. Our one-, two-, and three-directional simulations corresponded to Atlantic cod, human, and wild poplar data sets, respectively, which further supported our results. As discussed below, the WG population-specific F_{ST} estimator is a rescaling of expected heterozygosity, and a linear relationship between expected heterozygosity and population F_{ST} is expected. This shows that the population-specific F_{ST} estimator implicitly assumes that populations closest to the ancestral population have the highest heterozygosity. Our simulations evaluated the performance of the WG population-specific F_{ST} estimator for such cases.

In our analysis, genome-wide WG population-specific F_{ST} values successfully illustrated human evolutionary history, and indicated that humans originated in Kenya, expanded from the Middle East into Europe and from Central/South Asia into East Asia, and then possibly migrated to Oceania and America (Figure 2). Kenya is located just below Ethiopia, where the earliest anatomically modern humans were found from fossils (Nielsen et al. 2017). Our results are also in good agreement with the highest levels of genetic diversity being detected in Africa (Rosenberg et al., 2002), the relationship uncovered between genetic and geographic distance (Ramachandran et al., 2005), the shortest colonization route from East Africa (Liu et

al., 2006), and major migrations inferred from genomic data (Nielsen et al., 2017). The

genome-wide WG population-specific F_{ST} values are consistent with results obtained from 24

forensic STR markers (Buckleton et al., 2016).

Our analysis indicated that Atlantic cod originated in Canada (CAN08). Figure 3 suggested

that the population expansion of Atlantic cod began by minimal gene flow from Canada. They

might have first expanded to the west coast of Greenland before spreading to Iceland, the

North Sea, Norway, and the Baltic Sea. This result was consistent with genomic evidence that

Atlantic cod inhabit both sides of the Atlantic Ocean and evolved from a common

evolutionary origin (Berg et al. 2017). The migratory ecotypes characterized by deeper and

more offshore habitats and long-distance migrations (Hemmer-Hansen et al., 2013a) may

have played an important role in this expansion, as suggested in Figure 3C. In the original

Atlantic cod study (Therkildsen et al. 2013a), strong differentiation of CAN08 was found at

neutral markers, which prompted the authors to suggest that Greenland populations were the

result of colonization from Iceland rather than from refugial populations in southern North

America. In our study, CAN08 had the highest H_e, which was lower in Iceland than in

Greenland (Figure 3A); this result implies that Icelandic populations were the descendants of

colonists from Greenland, which in turn originated in Canada. The BAS0607 sample from the

Baltic Sea had the highest population-specific F_{ST} and the lowest H_e, which suggests that

Baltic cod is the newest population. This result agrees with the findings of a previous study,

which identified Baltic cod as an example of a species subject to ongoing selection for

reproductive success in a low salinity environment (Berg et al. 2015).

Although the wild poplar samples used in this study might not cover the whole distribution

range of the species, which extends from southern California to northern Alaska, Montana,

and Idaho (Geraldes et al. 2013), genome-wide population-specific F_{ST} values suggested that

wild poplar trees in southern British Colombia (SBC27) and inland British Colombia (IBC15,

IBC16) are the closest to the ancestral population. Our results showed that the wild popular

expanded in three directions: coastal British Colombia, southern Oregon, and northern British

Colombia (Figure 5). The largest population-specific F_{ST} value was found in the population

with the smallest heterozygosity, SBC22, which may have resulted from a bottleneck

(Geraldes et al., 2014).

Genome-wide population-specific F_{ST} detects key environments that promote

adaptation

Our GLS of genome-wide population-specific F_{ST} values revealed that long daylight hours,

abundant rainfall, and dry summer conditions are the key environmental factors that

influenced the evolution of wild poplar (Table 1). Wild poplar could have expanded their

distribution by their fluffy seeds being blown away by wind, and individuals that had genomic

variation were able to adapt to local environments. The results suggested that wild popular

originated from southern British Colombia (SBC27) and inland British Colombia (IBC15, 16)

and expanded its distribution to NBC by adapting to longer day lengths, to SBC seashores

adapting to the rainy environment, and to southern Oregon (ORE30) adapting to dry summer

conditions (Figure 5). A previous study on wild poplar revealed that genes involved in

drought response were identified as F_{ST} outliers along with other genes related to

transcriptional regulation and nutrient uptake (Geraldes et al. 2014), which is a finding

consistent with our GLS results. Our results were also consistent with the F_{ST} outlier test of

the original study (Geraldes et al., 2014), in which Bayescan (Foll and Gaggiotti 2008)

revealed that genes involved in circadian rhythm and response to red/far-red light had high

locus-specific global F_{ST} values. Moreover, the first principal component of SNP allele

frequencies was significantly correlated with day length, and a previous enrichment analysis

for population structuring uncovered genes related to circadian rhythm and photoperiod

(McKown et al. 2014a). Our results were in agreement with the previous findings, which

show the usefulness of using GLS of genome-wide population-specific F_{ST} to infer

environmental adaptation and population expansion of species.

Properties of F_{ST} moment estimators

Previous studies have suggested or indicated that the "ratio of averages" works better than the

"average of ratios" as the number of independent SNPs increases (Weir and Cockerham 1984;

Weir and Hill 2002; Bhatia et al. 2013). In regard to the WG population-specific F_{ST} estimator,

similar results were obtained for the 377 human microsatellite loci using either the "ratio of

averages" or the "average of ratios" (Figure S3). This similar outcome may have been due to

the relatively small variation in the locus-specific global F_{ST} values (Figure not shown) and

the relatively large number of alleles (12 ± 4) of human microsatellites.

To explicitly show the underlying mechanism, we used the observed heterozygosity of

population i ($\hat{H}_{Si}$) as derived in Nei and Chesser (1983) (Supplemental Note). When the

number of loci (L) increases, the average observed heterozygosity over all loci converges to

its expected value according to the law of large numbers as

$$\frac{1}{L}\sum_{l=1}^{L}\left(1-\sum_{u=1}^{m}\tilde{p}_{iu}^2\right) \to \frac{1}{L}\sum_{l=1}^{L}\left(1-E\left[\sum_{u=1}^{m}\tilde{p}_{iu}^2\right]\right).$$

543 The observed heterozygosity thus converges to the expected value:

$$\widehat{H}_{Si} = \widehat{H}_{Si}\left(1 - \frac{1}{n_i}\right) + \frac{\widehat{H}_{0i}}{2n_i} \quad \rightarrow H_{Si}\left(1 - \frac{1}{n_i}\right) + \frac{H_{0i}}{2n_i}.$$

Similarly, $\widehat{H}_S$ and $\widehat{H}_T$ converge to their expected values. This example indicates that the

numerators and denominators of bias-corrected F_{ST} moment estimators, whether global,

pairwise, or population-specific, converge to their true means and provide unbiased estimates

of F_{ST} in population genomics analyses with large numbers of SNPs. Our analyses show that

genomic data highlight the usefulness of the bias-corrected moment estimators of traditional

F_{ST} developed in the early 1980s (Nei and Chesser 1983; Weir and Cockerham 1984) and

population-specific F_{ST} (Weir and Goudet 2017).

To estimate pairwise F_{ST}, our previous coalescent simulations based on ms (Hudson 2002)

showed that NC83 performed best among the present F_{ST} estimators for cases with 10,000

SNPs (Kitada et al. 2017). Other F_{ST} moment estimators within an ANOVA framework

produce values approximately double those of true values when used to estimate pairwise F_{ST}.

NC83 considers a fixed set of population samples; in contrast, the other F_{ST} moment

estimators consider replicates of a set of populations (Weir and Cockerham 1984; Holsinger

and Weir 2009). The models for replicates of population samples were considered to

appropriately estimate global F_{ST} and/or mean ancestral coefficient, but cause over-estimation

when used to estimate pairwise F_{ST}.

The WG population-specific F_{ST} moment estimator measures population genetic diversity

under the framework of relatedness of individuals and identifies the population with the

largest genetic diversity as the ancestral population. This estimator thus works to infer

evolutionary history through genetic diversity. The WG population-specific F_{ST} estimator is

based on allele matching probabilities, where within-population observed heterozygosity can

be written as $1 - \widetilde{M}_W^i$. When Hardy–Weinberg equilibrium is assumed ($\widehat{H}_{0i} = \widehat{H}_{Si}$), the

preceding formula is equivalent to the NC83 unbiased estimator of the gene diversity of

population i $\left(\widehat{H}_{Si}\right)$ (Supplemental Note):

$$1 - \widetilde{M}_W^i = \frac{2n_i}{2n_i - 1}\left(1 - \sum_{u=1}^m \tilde{p}_{iu}^2\right) = \widehat{H}_{Si} \, .$$

570   Another variable, $\widetilde{M}^B$, is "average over pairs of populations of between-population-pair

571   matching" (Weir and Goudet 2017). $\widetilde{M}^B$ is the homozygosity over pairs of populations, and

572   we can write observed heterozygosity over pairs of populations as $1 - \widetilde{M}^B = \widehat{H}_B$. $\widehat{H}_B$ is an

573   estimator for the denominator of Hudson et al. (1992). When using only allele frequencies, the

574   population-specific F_{ST} estimator can be written in terms of gene diversity as

$$\text{ps}\widehat{F}_{ST}^i = \frac{\widetilde{M}_w^i - \widetilde{M}^B}{1 - \widetilde{M}^B} = \frac{\widehat{H}_B - \widehat{H}_{Si}}{\widehat{H}_B} = 1 - \frac{\widehat{H}_{Si}}{\widehat{H}_B} \, . \quad (7)$$

This formulation is reasonable, because WG population-specific F_{ST} uses "allele matching,

equivalent to homozygosity and complementary to heterozygosity as used by Nei (1973),

rather than components of variance" (Weir and Goudet 2017). In our three case studies, a

linear relationship between H_e of each population (= H_{Si}) and $\text{ps}\widehat{F}_{ST}^i$ was evident (Figure

S6). The coefficient of determination, R^2, was 0.91 for 51 human populations ($n = 1,035$),

0.99 for 34 Atlantic cod populations ($n = 1,065$), and 0.82 for 25 wild poplar populations ($n =$

441). The goodness of fit to the linear function should depend on population sample size

(number of individuals).

In the Atlantic cod case study, CAN08 had the highest H_e (Figure 4b) and a very large

negative population-specific F_{ST} value of −0.21 ± 0.019 compared with the maximum value

of 0.22 ± 0.014 in BAS0607 (Figure S6, Table S5). The Atlantic cod data corresponded to the

one-directional model of our simulations, where the WG population-specific F_{ST} value was

significantly negative in the ancestral population, whereas H_e was the largest (Figure 1D,G).

Our consistent results between the simulations and Atlantic cod case study indicate that, when

gene flow from other populations into the source population is limited, relatively large H_e

($\widehat{H}_{Si}$) is maintained in the source population. In such cases with $\widehat{H}_{Si} > \widehat{H}_B$, Equation 7

produces negative values of population-specific F_{ST}.

Bayesian F_{ST} estimators measure the deviation from the average of the sampled

populations

Our F_{ST} map based on the empirical Bayesian population-specific F_{ST} values (Beaumont and

Balding 2004) indicated that the Middle East, Europe, and Central/South Asia were centers of

human origin (Figure 6A), which was consistent with that from the full Bayesian

population-specific F_{ST} estimator (Foll and Gaggiotti 2006) (figure not shown). Our

integrated NJ tree showed that the Hazara, Pakistan population was genetically closest to the

human ancestors (Figure 6B). The results obtained with Bayesian estimators were a

consequence of Equation 4, which uses the mean allele frequency over populations ($\bar{p}_{lu}$) to

reduce the number of parameters to be estimated. The numbers of sampling locations of the

51 human populations were as follows: 6 from Africa, 12 from the Middle East/Europe, 9

from Central/South Asia, 18 from East Asia, 2 from Oceania, and 4 from America. The mean

allele frequency ($\bar{p}_{lu}$) reflected the weight of samples from the Middle East, Europe, and

Central/South Asia, thereby resulting in these areas being identified as centers of origin.

Instead of $\bar{p}_{lu}$, the full Bayesian method uses allele frequencies in the ancestral population,

p_{lu}, which are generated from a noninformative Dirichlet prior, $p_{lu} \sim Dir$ (1, ...,1). Our

results indicate that not enough information is available to estimate allele frequencies in the

ancestral population assumed in the models. The shrinkage effect on allele frequencies in

Bayesian inference (Stein, 1956) may shift population-specific F_{ST} values toward the average

of the whole population. Indeed, when we used a subsample of the 37 human populations (3

from Africa, 6 from the Middle East/Europe, 4 from Central/South Asia, 18 from East Asia, 2

from Oceania, and 4 from America; Table S7), the area with the highest population-specific

F_{ST} values shifted toward Central/South Asia and East Asia, whereas Bantu Kenyans had the

smallest WG population-specific F_{ST} value; this was consistent with the results from the full

data set (Figure 7).

In contrast, because of shrinkage toward mean allele frequencies, maximum likelihood and

Bayesian estimators of locus-specific global F_{ST} improve the power to detect genes under

environmental selection (Beaumont and Balding 2004). Our empirical Bayes pairwise F_{ST}

estimator (EBF_{ST}; Kitada et al. 2007), which is based on Equation 4, is also useful in cases

involving a relatively small number of polymorphic marker loci, such as microsatellites; it

performs best by averaging large sampling variation of allele frequencies in populations with

small sample sizes, particularly in high gene flow scenarios (Kitada et al., 2017). However,

EBF_{ST} suffers from a shrinkage effect similar to that of Bayesian population-specific F_{ST}

estimators. We note that the shrinkage effect on allele frequencies can enhance the bias of

Bayesian F_{ST} estimators, particularly in genome analyses where large numbers of SNPs are

used.

Conclusions

The genome-wide WG population-specific F_{ST} moment estimator was able to identify the

source population and trace the evolutionary history of the derived population's history based

on genetic diversity under the assumption that populations closest to the ancestral population

have the highest heterozygosity. In contrast, NC83 genome-wide pairwise F_{ST} moment

estimator represents the current population structure. By integrating estimates from both F_{ST}

estimators on NJ trees and maps of sampling locations, we obtained a picture of population

structure by incorporating evolutionary history. Our GLS analysis of genome-wide

population-specific F_{ST}, which takes the correlation between population-specific F_{ST} values

into account, provides insights into how a species has adapted to key environments and

expanded its distribution. Given a large number of loci, bias-corrected F_{ST} moment estimators,

whether global, pairwise, or population-specific, provide unbiased estimates of F_{ST} supported

by the law of large numbers. Genomic data highlight the usefulness of the bias-corrected

moment estimators of F_{ST}.

ACKNOWLEDGEMENTS

We express our appreciation to editors Mark Beaumont and Nicholas Barton for reviewing

this manuscript and providing the opportunity for revision. We thank the reviewers for

providing essential comments that significantly improved the manuscript. This study was

supported by Japan Society for the Promotion of Science Grants-in-Aid for Scientific

Research KAKENHI nos. 16H02788 and 19H04070 to HK and 18K0578116 to SK.

LITERATURE CITED

Balloux, F., and N. Lugon-Moulin, 2002 The estimation of population differentiation with

microsatellite markers. Mol. Ecol. 11: 155–165.

https://doi.org/10.1046/j.0962-1083.2001.01436.x

Balding, D. J., and R. A. Nichols, 1995 A method for quantifying differentiation between

populations at multi-allelic loci and its implications for investigating identity and

paternity. Genetica 96: 3–12. https://doi.org/10.1007/BF01441146

Beaumont, M. A., and D. J. Balding, 2004 Identifying adaptive genetic divergence among populations from genome scans. Molec. Ecol. 13: 969–980. https://doi.org/10.1111/j.1365-294X.2004.02125.x

Beaumont, M. A., 2005 Adaptation and speciation: what can F_{ST} tell us? Trends Ecol. Evol. 20: 435–440. https://doi.org/10.1016/j.tree.2005.05.017

Berg, P. R., S. Jentoft, B. Star, K. H. Ring, H. Knutsen *et al.* 2015 Adaptation to low salinity promotes genomic divergence in Atlantic cod (*Gadus morhua* L.). Genome Biol. Evol. 7: 1644–1663. https://doi.org/10.1093/gbe/evv093

Berg, P. R., B. Star, C. Pampoulie, M. Sodeland, J. M. Barth *et al.* 2016 Three chromosomal rearrangements promote genomic divergence between migratory and stationary ecotypes of Atlantic cod. Sci. Rep. 6: 23246. https://doi.org/10.1038/srep23246

Berg, P. R., B. Star, C. Pampoulie, I. R. Bradbury, P. Bentzen *et al.* 2017 Trans-oceanic genomic divergence of Atlantic cod ecotypes is associated with large inversions. *Heredity*, *119*, 418–428. https://doi.org/10.1038/hdy.2017.54

Bhatia, G., N. Patterson, S. Sankararaman, and A. L. Price, 2013 Estimating and interpreting F_{ST}: the impact of rare variants. Genome Res. 23: 1514–1521. http://www.genome.org/cgi/doi/10.1101/gr.154831.113

Bradbury, I. R., and P. Bentzen, 2007 Non-linear genetic isolation by distance: implications for dispersal estimation in anadromous and marine fish populations. Mar. Ecol. Prog. Ser. 340: 245–257. https://doi:10.3354/meps340245

Buckleton, J., J. Curran, J. Goudet, D. Taylor, A. Thiery *et al.* 2016 Population-specific F_{ST} values for forensic STR markers: A worldwide survey. Foren. Sci. Intnatl. Genetics 23: 91–100. https://doi.org/10.1016/j.fsigen.2016.03.004

Cann, H. M., C. De Toma, L. Cazes, M. F. Legrand, V. Morel, *et al.*, 2002 A human genome

diversity cell line panel. Science 296: 261–262.

https://doi:10.1126/science.296.5566.261b

Cockran, W. G., 1977 *Sampling Techniques*. Wiley, New York.

Coop, G., D. Witonsky, A. Di Rienzo, and J. K. Pritchard, 2010 Using environmental

correlations to identify loci underlying local adaptation. Genetics 185: 1411–1423.

https://doi.org/10.1534/genetics.110.114819

Diamond, J. 1997 *Guns, Germs and Steel: The Fates of Human Societies.* Random House,

London.

Excoffier, L., 2001 Analysis of population subdivision, pp. 271–307 in Handbook of

Statistical Genetics, edited by D. J. Balding, M. Bishop and C. Cannings. Wiley,

Chichester, UK.

Excoffier, L., P. E. Smouse, & J. M. Quattro, 1992. Analysis of molecular variance inferred

from metric distances among DNA haplotypes: application to human mitochondrial

DNA restriction data. Genetics 131: 479-491. PubMed 1644282

Falush, D., M. Stephens, and J. K. Pritchard, 2003 Inference of population structure using

multilocus genotype data: linked loci and correlated allele frequencies. Genetics 164:

1567–1587.

Foll, M., and O. Gaggiotti, 2006 Identifying the environmental factors that determine the

genetic structure of populations. Genetics 174: 875–891.

https://doi.org/10.1534/genetics.106.059451

Foll, M., and O. E. Gaggiotti, 2008 A genome-scan method to identify selected loci

appropriate for both dominant and codominant markers: a Bayesian perspective.

Genetics 180: 977–993. https://doi.org/10.1534/genetics.108.092221

Gaggiotti, O. E., and M. Foll, 2010 Quantifying population structure using the F-model.

Molec. Ecol. Resour. 10: 821–830. https://doi.org/10.1111/j.1755-0998.2010.02873.x

Geraldes, A., J. Pang, N. Thiessen, T. Cezard, R. Moore *et al.* 2011 SNP discovery in black

cottonwood (*Populus trichocarpa*) by population transcriptome resequencing. Molec.

Ecol. Resour. 11: 81–92. https://doi.org/10.1111/j.1755-0998.2010.02960.x

Geraldes, A., S. P. Difazio, G. T. Slavov, P. Ranjan, W. Muchero *et al.* 2013 A 34K SNP

genotyping array for *Populus trichocarpa*: design, application to the study of natural

populations and transferability to other *Populus* species. Molec. Ecol. Resour. 13:

306–323. https://doi.org/10.1111/1755-0998.12056

Geraldes, A., N. Farzaneh, C. J. Grassa, A. D. McKown, R. D. Guy *et al.* 2014 Landscape

genomics of *Populus trichocarpa* the role of hybridization limited gene flow and

natural selection in shaping patterns of population structure. Evolution 68: 3260–3280.

https://doi.org/10.1111/evo.12497

Hemmer-Hansen, J., E. E. Nielsen, N. O. Therkildsen, M. I. Taylor, R. Ogden *et al.* 2013a A

genomic island linked to ecotype divergence in Atlantic cod. Molec. Ecol. 22:

2653–2667. https://doi.org/10.1111/mec.12284

Hemmer-Hansen, J., E. E. Nielsen, N. O. Therkildsen, M. I. Taylor, R. Ogden *et al.* 2013b

Data from: A genomic island linked to ecotype divergence in Atlantic cod, Dryad,

Dataset, https://doi.org/10.5061/dryad.9gf10

Holsinger, K.E., and B. S. Weir, 2009 Genetics in geographically structured populations:

defining, estimating and interpreting F_{ST}. Nat. Rev. Genet. 9: 639–650.

https://doi.org/10.1038/nrg2611

Hudson, R. R. 2002 Generating samples under a Wright–Fisher neutral model of genetic

variation. Bioinformatics 18: 337–338. https://doi.org/10.1093/bioinformatics/18.2.337

Hudson, R. R., M. Slatkin, and W. P. Maddison, 1992 Estimation of levels of gene flow from

DNA sequence data. Genetics 132: 583–589. PubMed 1427045

Jorde, P. E., G. Søvik, J. I. Westgaard, J. Albretsen, C. André *et al.* 2015 Genetically distinct

populations of northern shrimp, *Pandalus borealis*, in the North Atlantic: adaptation to

different temperatures as an isolation factor. Molec. Ecol. 24: 1742–1757.

https://doi.org/10.1111/mec.13158

Kitada, S., T. Kitakado, and H. Kishino, 2007 Empirical Bayes inference of pairwise F_{ST} and

its distribution in the genome. Genetics 177: 861–873.

https://doi.org/10.1534/genetics.107.077263

Kitada, S., R., Nakamichi, and H. Kishino, 2017 The empirical Bayes estimators of fine-scale

population structure in high gene flow species. Molec. Ecol. Resour. 17: 1210–1222.

https://doi.org/10.1111/1755-0998.12663

Liu, H., F. Prugnolle, A. Manica, and F. Balloux, 2006 A geographically explicit genetic

model of worldwide human-settlement history. Am. Hum. Genet. 79: 230–237.

https://doi.org/10.1086/505436

Limborg, M. T., S. J. Helyar, M. De Bruyn, M. I. Taylor, E. E. Nielsen *et al.* 2012

Environmental selection on transcriptome☐derived SNPs in a high gene flow marine

fish, the Atlantic herring (*Clupea harengus*). Molec, Ecol. 21: 3686–3703.

https://doi.org/10.1111/j.1365-294X.2012.05639.x

McKown, A. D., R. D. Guy, J. Klápště, A. Geraldes, M. Friedmann *et al.*, 2014a Geographical

and environmental gradients shape phenotypic trait variation and genetic structure in

Populus trichocarpa. New Phytol. 201, 1263–1276. https://doi.org/10.1111/nph.12601

McKown, A. D., J. Klápště, R. D. Guy, A. Geraldes, I. Porth *et al.*, 2014b Genome -wide

association implicates numerous genes underlying ecological trait variation in natural

populations of *Populus trichocarpa*. New Phytol. 203: 535–553.

https://doi.org/10.1111/nph.12815

McKown, A. D., R. D. Guy, L. Quamme, J. Klápště, J. La Mantia *et al.*, 2014c Association

genetics, geography and ecophysiology link stomatal patterning in *Populus trichocarpa*

with carbon gain and disease resistance trade-offs. Molec, Ecol. 23: 5771–5790.

https://doi.org/10.1111/mec.12969

Nei, M., 1973 Analysis of gene diversity in subdivided populations. Proc. Natl. Acad. Sci.

USA 70: 3321–3323. https://doi.org/10.1073/pnas.70.12.3321

Nei, M., 1977 F‑statistics and analysis of gene diversity in subdivided populations. Ann.

Hum. Genet. 41: 225–233. https://doi.org/10.1111/j.1469-1809.1977.tb01918.x

Nei, M., and R. K. Chesser, 1983 Estimation of fixation indices and gene diversities. Ann.

Hum. Genet. 47: 253–259. https://doi.org/10.1111/j.1469-1809.1983.tb00993.x

Nicholson, G., A. V. Smith, F. Jónsson, Ó. Gústafsson, K. Stefánsson *et al.*, 2002 Assessing

population differentiation and isolation from single‑nucleotide polymorphism data. J.

Roy. Stat. Soc. B. Stat. Method. 64: 695–715.

https://doi.org/10.1111/1467-9868.00357

Nielsen, R., J. M. Akey, M. Jakobsson, J. K. Pritchard, S. Tishkoff, and E. Willerslev, 2017

Tracing the peopling of the world through genomics. Nature 541: 302–310.

https://doi.org/10.1038/nature21347

Nosil, P., D. J. Funk, and D. Ortiz-Barrientos, 2009 Divergent selection and heterogeneous

genomic divergence. Molec, Ecol. 18: 375–402.

https://doi.org/10.1111/j.1365-294X.2008.03946.x

Orsini, L., J. Vanoverbeke, I. Swillen, J. Mergeay, and L. De Meester, 2013 Drivers of

population genetic differentiation in the wild: isolation by dispersal limitation,

isolation by adaptation and isolation by colonization. Molec, Ecol. 22, 5983–5999.

https://doi.org/10.1111/mec.12561

Palsbøll, P. J., M. Berube, and F. W. Allendorf, 2007 Identification of management units using population genetic data. Trends Ecol. Evol. 22: 11–16.

https://doi.org/10.1016/j.tree.2006.09.003

Pérez-Lezaun, A., F. Calafell, E. Mateu, D. Comas, R. Ruiz-Pacheco *et al.* 1997 Microsatellite variation and the differentiation of modern humans. Human Genetics 99: 1–7.

https://doi.org/10.1007/s004390050299

Petrou, E. L., J. E. Seeb, L. Hauser, M. J. Witteveen, W. D. Templin *et al.* 2014 Fine-scale sampling reveals distinct isolation by distance patterns in chum salmon (*Oncorhynchus keta*) populations occupying a glacially dynamic environment. Cons. Genet. 15: 229–243. https://doi.org/10.1007/s10592-013-0534-3

Ramachandran, S., O. Deshpande, C. C. Roseman, N. A. Rosenberg, M. W. Feldman *et al.* 2005 Support from the relationship of genetic and geographic distance in human populations for a serial founder effect originating in Africa. Proc. Natl. Acad. Sci. 102: 15942–15947. https://doi.org/10.1073/pnas.0507611102

Raymond, M., and F., Rousset,1995 GENEPOP (version 1.2): population genetics software for exact tests and ecumenicism. J. Hered. 86: 248–249.

Rosenberg, N. A., J. K. Pritchard, J. L. Weber, H. M. Cann, K. K. Kidd *et al.* 2002 Genetic structure of human populations. Science 298: 2381–2385.

https://doi.org/10.1126/science.1078311

Rougemont, Q., J. S. Moore, T. Leroy, E. Normandeau, E. B. Rondeau *et al.* 2020 Demographic history shaped geographical patterns of deleterious mutation load in a broadly distributed Pacific Salmon. PLoS Genetics 16: e1008348.

https://doi.org/10.1371/journal.pgen.1008348

Rousset, F., 1997 Genetic differentiation and estimation of gene flow from F-statistics under

isolation by distance. Genetics 145: 12193–1228.

Rousset, F., 2001 Inferences from spatial population genetics, pp. 239–269 in Handbook of

Statistical Genetics, edited by D. J. Balding, M. Bishop and C. Cannings. Wiley,

Chichester, UK.

Rousset, F., 2004 *Genetic Structure and Selection in Subdivided Populations*. Princeton

University Press, Princeton, NJ.

Rousset, F., 2008 Genepop'007: a complete reimplementation of the Genepop software for

Windows and Linux. Mol. Ecol. Resour. 8: 103–106.

https://doi.org/10.1111/j.1471-8286.2007.01931.x

Rutherford, A. 2016 *A Brief History of Everyone Who Ever Lived: The Human Story Retold*

Through Our Genes. The Experiment, NY.

Saitou, N., and M. Nei, 1987 The neighbor-joining method: a new method for reconstructing

phylogenetic trees. Molec. Biol. Evol. 4: 406–425.

https://doi.org/10.1093/oxfordjournals.molbev.a040454

Schwartz, M. K., G. Luikart, and R. S. Waples, 2007 Genetic monitoring as a promising tool

for conservation and management. Trends Ecol. Evol. 22: 25–33.

https://doi.org/10.1016/j.tree.2006.08.009

Slatkin, M. 1987 Gene flow and the geographic structure of natural populations. Science, 236:

787–792. https://doi.org/10.1126/science.3576198

Stein, C. 1956 Inadmissibility of the usual estimator for the mean of a multivariate

distribution, pp. 197–206 in Proceedings of the Third Berkeley Symposium on

Mathematical Statistics and Probability Vol. 1, University of California Press, Berkeley,

CA.

Therkildsen, N. O., J. Hemmer-Hansen, R. B. Hedeholm, M. S. Wisz, C. Pampoulie *et al.*

2013a Spatiotemporal SNP analysis reveals pronounced biocomplexity at the northern

range margin of Atlantic cod *Gadus morhua*. Evol. Appl. 6: 690–705.

https://doi.org/10.1111/eva.12055

Therkildsen, N. O., J. Hemmer-Hansen, R. B. Hedeholm, M. S. Wisz, C. Pampoulie *et al.*

2013b Data from: Spatiotemporal SNP analysis reveals pronounced biocomplexity at

the northern range margin of Atlantic cod *Gadus morhua*, v2, Dryad, Dataset,

https://doi.org/10.5061/dryad.rd250

Waples, R. S., and O. Gaggiotti, 2006 What is a population? An empirical evaluation of some

genetic methods for identifying the number of gene pools and their degree of

connectivity. Mol. Ecol. 15: 1419–1439.

https://doi.org/10.1111/j.1365-294X.2006.02890.x

Weir, B. S., and C. C. Cockerham, 1984 Estimating *F*-statistics for the analysis of population

structure. Evolution 38: 1358–1370.

Weir, B. S., and W. G. Hill, 2002 Estimating *F*-statistics. Annu. Rev. Genet. 36: 721–750.

https://doi.org/10.1146/annurev.genet.36.050802.093940

Weir, B. S., L. R. Cardon, A. D. Anderson, D. M. Nielsen, and W. G. Hill, 2005 Measures of

human population structure show heterogeneity among genomic regions. Genome Res.

15: 1468–1476. https://doi.org/10.1101/gr.4398405

Weir, B. S., and J. Goudet, 2017 A unified characterization of population structure and

relatedness. Genetics 206: 2085–2103. https://doi.org/10.1534/genetics.116.198424

Whitlock, M. C., and D. E. Mccauley, 1999 Indirect measures of gene flow and migration:

$F_{\mathrm{ST}} \neq 1(4Nm + 1)$. Heredity 82: 117–125.

https://doi.org/10.1046/j.1365-2540.1999.00496.x

Wright, S., 1931 Evolution in mendelian populations. Genetics 16: 97–158. PMID: 17246615

Wright, S., 1951 The genetical structure of populations. Ann. Eugen. 15: 323–354.

https://doi.org/10.1111/j.1469-1809.1949.tb02451.x

Table 1 Regression of genome-wide population-specific F_{ST} of 25 wild poplar populations on environmental variables

Variables	Estimate	SE	Z	p
DAY	0.0489	0.0164	2.99	0.003**
MAT	-0.0088	0.0086	-1.03	0.305
MAP	0.0001	0.0000	2.79	0.005**
SHM	0.0022	0.0009	2.38	0.018*

858 DAY; longest day length (hours),
859 MAT; mean annual temperature (°C),
860 MAP; mean annual precipitation (mm),
861 SHM; summer heat-moisture index,
862 *$p < 0.05$ and **$p < 0.01$

Figure 1 Results from population colonization simulations. Schematic diagrams of the models: (A) one-, (B) two-, and (C) three-directional colonization. Population 1 in red is ancestral, and arrows show the direction of colonization. Each population exchanged 1% of N_e genes with adjacent population(s) in every generation, as shown by arrows (see text). Neighbor-joining unrooted trees based on pairwise F_{ST} distance matrix overlaid with population-specific F_{ST} values for each model (D–F). The color of each population shows the magnitude of population-specific F_{ST} values. The radius of each population is proportional to the level of He, as visualized by $(H_e \times 10)^{2.5}$. Population-specific F_{ST} values for 25 simulated populations are presented for each model (G–I).

874
875
Figure 2 Population structure of 51 human populations (n = 1,035; 377
microsatellites). (A) Map showing genetic diversity and population connectivity with
the magnitude of population-specific F_{ST} values. Populations connected by yellow
lines are those with pairwise F_{ST} < 0.02. The radius of each sampling point is
proportional to the level of He, as visualized by H_e^{12}. (B) Distribution of
population-specific F_{ST} values ± 2×SE. (C) Unrooted NJ tree based on pairwise F_{ST}
overlaid with population-specific F_{ST} values on population labels. The color of each
population shows the magnitude of population-specific F_{ST} values.

884
885
Figure 3 Population structure of 34 geographical samples of wild Atlantic cod (n =
1,065; 921 SNPs). (A) Map showing genetic diversity and population connectivity with
the magnitude of population-specific F_{ST} values. Populations connected by yellow
lines are those with pairwise F_{ST} < 0.02. The radius of each sampling point is
proportional to the level of He, as visualized by H_e^{50}. (B) Distribution of
population-specific F_{ST} values ± 2×SE. (C) Unrooted NJ tree based on pairwise F_{ST}
overlaid with population-specific F_{ST} values on population labels. The color of each
population shows the magnitude of population-specific F_{ST} values.

894
895
Figure 4 Population structure for 25 geographical samples of wild poplar (n = 441;
29,355 SNPs). (A) Map showing genetic diversity and population connectivity with the
magnitude of population-specific F_{ST} values. Populations connected by yellow lines
are those with pairwise F_{ST} < 0.02. The radius of each sampling point is proportional to
the level of *He*, as visualized by H_e^{100}. (B) Distribution of population-specific F_{ST}
values $\pm$ 2×SE. (C) Unrooted NJ tree based on pairwise F_{ST} overlaid with
population-specific F_{ST} values on population labels. The color of each population
shows the magnitude of population-specific F_{ST} values.

904
905
Figure 5 Population range expansion and environmental adaptation. Longest day
length vs. (A) summer heat-moisture index and (B) mean annual precipitation for 25
geographical samples of wild poplar. The color of each population shows the
magnitude of population-specific F_{ST} values. The circles show inferred population
expansion from IBC15, IBC16, and SBC27. The color of the circles refers to the
population clusters (see, Figure S5).

912
913
Figure 6 Population structure of 51 human populations inferred based on Bayesian
population-specific F_{ST} values. (A) Map showing genetic diversity and population
connectivity with the magnitude of population-specific F_{ST} values. Populations
connected by yellow lines are those with pairwise $F_{ST} < 0.02$. The radius of each
sampling point is proportional to the level of *He*, as visualized by H_e^{12}. (b) Unrooted NJ
tree based on pairwise F_{ST} overlaid with population-specific F_{ST} values on population
labels. The color of each population shows the magnitude of population-specific F_{ST}
values.

Figure 7 Population structure of 37 subsamples of human populations inferred based on Bayesian and WG population-specific F_{ST} values. The numbers of sampling locations of the subsamples were as follows: 3 from Africa, 6 from the Middle East/Europe, 9 from Central/South Asia, 18 from East Asia, 2 from Oceania, and 4 from America. Maps show genetic diversity and population connectivity with the magnitude of (A) Bayesian and (C) WG population-specific F_{ST} values. Populations connected by yellow lines are those with pairwise $F_{ST} < 0.02$. The radius of each sampling point is proportional to the level of *He*, as visualized by H_e^{12}. Unrooted NJ tree based on pairwise F_{ST} overlaid with (B) Bayesian and (D) WG population-specific F_{ST} values on population labels. The color of each population shows the magnitude of population-specific F_{ST} values.