A Novel Multiplex qPCR Assay for Detection of *Plasmodium falciparum* with *Histidine-rich*

***Protein 2 and 3 (pfhrp2 and pfhrp3)* Deletions in Polyclonal Infections**

Lynn Grignard[1], Debbie Nolder[1,2], Nuno Sepúlveda[1,3], Araia Berhane[4], Selam Mihreteab[4], Robert
Kaaya[5], Jody Phelan[1], Kara Moser[6], Donelly A. van Schalkwyk[1], Susana Campino[1], Jonathan B.
Parr[6], Jonathan J. Juliano[2,7], Peter Chiodini[2,7], Jane Cunningham[8], Colin J. Sutherland[1], Chris
Drakeley[1] and Khalid B. Beshir[1]*

1. Faculty of Infectious Diseases, London School of Hygiene and Tropical Medicine, United Kingdom
2. PHE Malaria Reference Laboratory, London School of Hygiene & Tropical Medicine, United
Kingdom
3. Centre of Statistics and Applications of University of Lisbon, Portugal
4. Communicable Diseases Control Division, Ministry of Health, Eritrea
5. Kilimanjaro Christian Medical University College, Tanzania
6. University of North Carolina at Chapel Hill, United States
7. UCL Hospital for Tropical Diseases, United Kingdom
8. World Health Organization, Geneva, Switzerland

Abstract

Background: Rapid diagnostic tests (RDTs) that detect the malaria antigen histidine-rich protein

2 (HRP2) are widely used in endemic areas globally to confirm *Plasmodium falciparum* infection

in febrile patients. The emergence of parasites lacking the gene encoding HRP2 and escaping

RDT detection threatens progress in malaria control and elimination. Many health facilities in

malaria endemic countries are dependent on RDTs for diagnosis and some National Health

Service hospitals without expert microscopists rely on them for diagnosis out of hours. It is vital to

study the emergence and the extent of such parasites globally to guide diagnostic policy.

Currently, verification of the presence of such parasites in a blood sample requires a series of

PCR assays to confirm the presence of *P. falciparum* and in the absence of amplicons from

pfhrp2 and/or *pfhrp3*, which encodes a cross-reactive protein isoform. These tests have different

limits of detection and many laboratories have reported difficulty in confirming the absence of

pfhrp2 and *pfhrp3* with certainty.

Methods: We developed and validated a novel and rapid multiplex real time quantitative (qPCR)

assay to detect *pfhrp2, pfhrp3,* confirmatory parasite and human reference genes

simultaneously. We also applied the assay to detect *pfhrp2* and *pfhrp3* deletion in 462 field

samples from different endemic countries and UK travellers.

Results: The qPCR assay showed limit of detection and quantification of 0.76-1.5 parasites per

µl. The amplification efficiency, coefficient of determination (R^2) and slope for the genes were 96-

1.07%, 0.96-0.98 and -3.375 2 to - 3.416 respectively. The assay demonstrated diagnostic

sensitivity of 100% (n=19, 95% CI= (82.3%; 100%)) and diagnostic specificity of 100% (n=31;

95% CI= (88.8%; 100%)) in detecting *pfhrp2* and *pfhrp3* in. In addition, the qPCR assay

estimates *P. falciparum* parasite density and can detect *pfhrp2* and *pfhrp3* deletions masked in

polyclonal infections. We report *pfhrp2* and *pfhrp3* deletions in parasite isolates from Kenya,

Tanzania and in UK travellers.

Conclusion: The new qPCR assay is simple to use and offers significant advantages in speed

and ease of interpretation. It is easily scalable to routine surveillance studies in countries where

P. falciparum parasites lacking *pfhrp2* and *pfhrp3* are a threat to malaria control.

Key words: *pfhrp2, pfhrp3, pfldh,* qPCR, RDT and malaria

*Corresponding author: Khalid.Beshir@lshtm.ac.uk

Background

Malaria is caused by infecting protozoan parasites of the genus *Plasmodium. P. falciparum*

continues to be the predominant species with an estimated global incidence of more than 2228

million cases and about 405,000 deaths reported in 2018 (1). Immunochromatographic rapid

diagnostic tests (RDTs), which use membrane-bound antibodies to detect parasite proteins in

finger-prick blood samples, play a crucial role in malaria control successes in disease endemic

countries. Early diagnosis is critical to malaria elimination and eradication programs and RDT

deployment is an important component of the strategy. As a result, the global availability and

scale of use of RDTs has increased dramatically over the last 10 years (2). Most RDTs used

worldwide detect *P. falciparum* histidine-rich protein 2 (pfHRP2) and/or *Plasmodium* lactate

dehydrogenase (pLDH) antigens. Some studies have shown that at least some pfHRP2-based

RDTs also detect *P. falciparum* histidine-rich protein 3 (pfHRP3) due to a shared antigenic

epitope (2-5). In sub-Saharan Africa, which bears 90% of the global malaria burden, RDTs

accounted for 74% of diagnostic testing among suspected malaria cases in 2015, and pfHRP2-

based tests were the most widely used (2).

Parasites with *pfhrp2* and/or *pfhrp3* genes (*pfhrp2/3*) deletions were first observed in South

America and increasing reports of false-negative RDT results due to these parasites have now

emerged from selected region of Africa and Asia (6-8). In some countries, a high proportion of

RDT false-negative results due to these gene deletions has led to the changes in national

diagnostic guidelines (9, 10). However, before undertaking any drastic changes in diagnostic

testing policies or deploying less sensitive, less heat stable RDTs that detect alternative antigens,

malaria programs need, robust epidemiological data about local *pfhrp2/3* deletion prevalence.

The World Health Organization has prioritized studies of these parasites and developed a

protocol for *pfhrp2/3* deletion surveillance (11). However, confirmation of pfhrp2/3 deletions using

current techniques is challenging and time consuming. Most studies of *pfhrp2* and *pfhrp3*

deletions deploy conventional nested PCR (nPCR) amplification of several genes followed by

gel-electrophoresis (12). In this genotyping approach, at least three independent genes are used

to ascertain the quality of DNA and the presence of *P. falciparum* parasites to avoid unintentional

misclassification of pfhrp2/3 deletions in samples with low-concentration or degraded DNA (13,

14). The nPCR approach requires several rounds of PCR for each gene and running the gel-

electrophoresis for each PCR product. The nested-PCR genotyping approach is labour-intensive,

time consuming and is prone to contamination, particularly when deployed in large-scale

surveillance studies. The various nPCR methods used differ in limit of detection, and this can

cause type I and type II errors. Further, performance of reported the *pfhrp2* and *pfhrp3* PCR is

variable, with wide ranging limits of detection and the risk cross-reactivity in some assays. In

addition, gel-electrophoresis approaches do not detect deletions masked in polyclonal infections

(6).

In this study, we report the development of a multiplex qPCR assay which simultaneously detects

DNA from the human host, a single-copy parasite house-keeping gene, and the *pfhrp2* and

pfhrp3 genes, including in polyclonal *P. falciparum* infections, in a single reaction. We report the

validation and application of this novel method using DNA samples derived from DBS and whole

blood of field isolates and clinical samples. We also deploy the assay to estimate *P. falciparum*

parasite density to rule out low parasite density as a factor for false RDT negative results (5, 13)

when microscopic data is unavailable or if it is not reliable.

Methods

***Plasmodium falciparum* laboratory strains**

Initial validation of the qPCR assay was performed using culture-adapted laboratory isolates with different *pfhrp2* and *pfhrp3* status, 3D7 (wildtype, West Africa origin), Dd2 (*pfhrp2 deletion, Indochina origin*), HB3 (*pfhrp3* deletion, Honduras origin) were obtained from the Malaria Research Reference Reagent Repository (http://MR4.org). A culture-adapted isolate lacking both genes (3BD5, double deletion) was also obtained from Thomas Wellems (NIAID, US). Parasite cultures of 3D7, Dd2, HB3 and 3BD5 were tightly synchronized as ring stage trophozoites *in vivo* to simulate infected peripheral blood similar to previously used methods (15, 16). The WHO P*. falciparum* International standard (Pf INT), a reagent comprising lyophilised whole blood from a single hyperparasitaemic individual was obtained from NIBSC UK. Undiluted, this reagent represents a parasitaemia of 9.8% which is equivalent to 4.9 x 10^5 parasites per µl (15, 17).

Clinical and field DNA samples

Clinical validation and application of the qPCR assay was performed using a convenience of 462 DNA samples derived from both DBS and whole-blood samples. Study site, sample collection and other details including IRB approvals have already been published for the study in which Eritrean samples were collected (9, 10). Parasite DNA was isolated from fifty dried bloodspot (DBS) samples from suspected malaria patients in Eritrean; 299 samples from whole-blood collected in EDTA from symptomatic Tanzanian and Kenyan patients and anonymised 113 samples from UK malaria patients. Whole blood collected in EDTA from UK travellers with confirmed *P. falciparum* infections in 2018 was obtained from the Public Health England Malaria Reference Laboratory (MRL), London, UK. Samples from Kenya (Ahero) and Tanzania (Bagamoyo) were collected between Oct 2016 and Dec 2018 as part of a study of parasite clearance after treatment with artemisinin combination therapy. These samples from Kenya and Tanzania have aliquots of cryopreserved blood samples and were selected for *pfhrp2/3* deletion study to identify *pfhrp2/3*-deleted parasites for culture-adaptation.

DNA Extraction

DNA was extracted from DBS from Eritrea and from whole blood from the MRL and from cultured laboratory isolates using a robotic DNA extraction system (Qiasymphony, QIAGEN, Germany),

as previously described (17). For the clinical samples collected in Kenya and Tanzania, 200μl of whole-blood was extracted using the QIAamp Blood Mini Kit (Qiagen) into 200μl of Buffer EB as per the manufacturer's instructions.

Multiplex qPCR development

Gene target selection and primer design

To design highly specific amplification primers that are conserved across global *P. falciparum* isolates we carried out multiple alignment of *pfhrp2* gene sequences from 1581 published *P. falciparum* genomes (MalariaGEN) from Africa, SE Asia and South America, and a similar alignment was also carried out for *pfhrp3* (Figure S1). The DNA sequence of the genes were obtained from publicly available genomic data and the processing of the data has been described in our previous report (6). We have also aligned the 3D7 DNA sequence of *pfhrp2* (PF3D7_0831800) and *pfhrp3* (PF3D7_1372200) to ensure that the conserved primers of the two genes do not cross-bind and are specific to *pfhrp2* and *pfhrp3* respectively (Figure S2). The DNA sequences of *pfhrp2* and *pfhrp3* were aligned using Geneious v. 10 (Biomatters, USA)

We used *Plasmodium falciparum* lactate dehydrogenase *(pfldh,* PF3D7_1324900), coded by a single-copy gene on chromosome 13, as a confirmatory gene for the presence and quality of parasite DNA as well as a target for measuring parasite density. We used previously published qPCR methods, with some modification of reaction conditions to amplify *pfldh* (13) and the human beta tubulin gene (*HumTuBB*) (Table S1)(18). The latter was used both as an internal control and as a normalizer for measurement of parasite density and for detection of *pfhrp2/3* deletion in polyclonal infections. All primers and probes were ordered from Eurofins Scientific (Germany).

Modification of pfhrp2 primer at the 3' end

Due to limited availability of suitable conserved target sequences region in *pfhrp2* and *pfhrp3* that are dissimilar between the isoform genes and to prevent non-specific cross-binding of the *pfhrp2* primers to *pfhrp3*, we used a strategy altering nucleotides located at the 3' end region (within the last 5 nucleotides) of both *pfhrp2* primers (Table S1). In total, we designed six primers with different modification at the 3' end of the forward and reverse *pfhrp2* primers, which were then

tested empirically to identify primer pairs that delivered the best specificity while maintaining

product yield (sensitivity).

Assay optimization

The multiplex qPCR assay designed in this study was optimized for primer and probe

hybridization temperature; different primer and probe concentrations and different $MgCl_2$

concentrations. All the optimization analyses were performed in triplicates in a RGQ rotor-gene

(Qiagen, Germany).

The optimized final reaction conditions were performed in a final volume of 25 µl containing 1.6X

NH_4 buffer (Bioline); 4 mM $MgCl_2$ (Bioline); 800nM dNTPs (Bioline), 200nM of *pfhrp2* primers,

200nM *pfhrp3* primers, 120 nM of *pfldh*, 120 nM *HumTuBB* primers, 120 nM of *pfhrp2* probe, 120

nM *pfhrp3* probe, 80 nM of *pfldh* probe and 80 nM *HumTuBB* probe; 2 units of biotaq polymerase

and 5 µl of extracted DNA. The optimal thermocycling conditions selected were 3 mins at 95^0C,

followed by 45 cycles of 15 sec at 95^0C; 30 sec at 54^0C and 30 sec at 72^0C.

PCR efficiency, linear dynamic range and limit of detection

We conducted assay performance analysis based on the MIQE guidelines[19]. The efficiency of

primer and probe combinations for each gene and linear dynamics of the qPCR assays were

evaluated using seven 4-fold dilutions of Pf INT (from 12500 to 3 parasites per µl). The last

dilution series (3 parasites per µl) was then further diluted 2-fold (from 3 to 0.38 parasites per µl)

to determine the measured limit of detection (LOD). The three laboratory strains (Dd2, HB3 and

3BD5) with known *pfhrp2/3* status were included in the evaluation linear dynamic of the linear

range to examine the effect of DNA concentration on the cross reactivity of the primers.

Assay precision, analytical sensitivity and specificity

Performance of the multiplex qPCR *pfhrp2/3* assay was evaluated by measuring coefficient of

variation across the seven four-fold dilution series of Pf INT. The specificity and sensitivity as well

as the robustness of the assay was evaluated by testing the lowest two concentrations of Pf INT

replicates of eight and 20 *P. falciparum*-negative whole blood samples in three different

experiments.

Detection of parasites with pfhrp2 and pfhrp3 deletion hidden in polyclonal infections

To investigate whether the qPCR assay could detect *pfhrp2/3*-deleted parasites hidden in

polyclonal infections robustly and accurately, we generated pairwise mixtures of known *P.*

falciparum genotypes (Dd2 and Pf INT, HB3 and Pf INT, and 3BD5 and Pf INT) at different ratios

i.e. 1:1, 5:1, 10:1, 100:1, 1000:1, 10,000:1, 1:100,000, 1:10,000, 1:1000, 1:100, 1:10, 1:5, 1:1.

We estimated the abundance of *pfhrp2/3* deletion genotypes in the mix relative to the whole

parasite biomass as measured by relative quantification normalized to *pfldh* (measures the DNA

of all strains) and *humTuBB* genes. The qPCR assay was also used to detect *pfhrp2/3* deletions

in patient samples with known polyclonal infection in patient samples, as determined by a

peviously published high resolution melting qPCR assay (20).

Relative quantification using the pfldh gene

We used *pfldh* as a parasite target and *HumTuBB* as a normalizer to estimate relative parasite

density of each sample. The lowest parasite density (parasite per µl) with a coefficient of

variation of less than 35% was considered limit of quantification for the assay(21). We

determined the performance of the *pfldh* qPCR in the multiplex assay by comparing parasite

densities determined by a published *pgmet* duplex qPCR assay using samples from Eritrea (18).

Application to DNA from diverse field samples

The validated qPCR assay was then applied to DNA extracted from diverse field samples.

Samples with Cq values lower than the Cq value of the limit of detection of individual gene are

determined to be negative for the gene. Samples with *HumTuBB* positive and *pfldh* positive but

negative for *pfhrp2* or *pfhrp3* are determined to be *pfhrp2*-deleted and *pfhrp3*-deleted

respectively. Samples with *HumTuBB* positive but pfldh negative are determined to be parasite

negative. Finally, samples with *HumTuBB* negative are considered to be invalid and DNA

extraction and/or PCR experiment should be repeated.

Data Analysis

For all amplification curve analyses, the quantification cycle (Cq) threshold was placed above the

amplification curve of No Template Control (NTC) and any crossing point between the Cq

threshold and the amplification curve was considered positive (Cq value) for the specific sample.

To evaluate assay precision, we calculated the coefficient of variation (CV) of parasite density

(parasite per µl) as follows: CV% = (standard deviation/mean) × 100. To determine the limit of

detection (LOD), we calculated the percentage of positive samples and the lowest sample with

more than 3 parasites per PCR and with ≥ 95% of replicate samples detected was considered

LOD (19). Similarly, to determine the lower limit of quantification (LOQ), we calculated the

parasite density using delta delta method and the sample with lowest parasite density (parasite

per µl) with a CV ≤35% was considered LOQ for the assay (21). Throughout the manuscript CV

refers to variation in parasite density (parasite per µl).

For estimation of relative abundance of *pfhrp2/3* deletion in a mixed-strain infection we used

delta delta relative quantification method (18) as follows:

$$\Delta C_q = C_q \text{ of } \textit{pfhrp2/3} - C_q \text{ of human}$$

$$\Delta\Delta C_q = \Delta C_q \text{ samples} - \Delta C_q \text{ of calibrator (Pf INT)}$$

$$\textit{pfhrp2/3} \text{ positive clone abundance} = 2^{-\Delta\Delta C_q}$$

$$\Delta C_q = C_q \text{ of } \textit{pfldh} - C_q \text{ of human}$$

$$\Delta\Delta C_q = \Delta C_q \text{ samples} - \Delta C_q \text{ of calibrator (Pf INT)}$$

$$\text{Total parasite abundance} = 2^{-\Delta\Delta C_q}$$

$$\textit{pfhrp2/3} \text{ deletion \%} = \textit{pfhrp2/3} \text{ clone abundance / total parasite abundance x 100}$$

Before calculating the relative abundance of *pfhrp2/3* strains, the C_q threshold of the positive

control (calibrator, Pf INT, 0.98% parasitaemia) was adjusted in each channel in such a way

that the C_q value is similar in all parasite target genes.

For comparison of parasite density estimated by two different $_q$PCR assays, we used STATA

(v 15, USA) software to perform linear regression.

Ethics approval and consent to participate

Ethical approval for collection of samples was obtained from each local ethical committee in

Eritrea (Eritrean MOH Research and Ethical Committees), Kenya (KEMRI IRB, 3293) and

Tanzania (MUHAS IRB, DA.282/298/01). Ethical approval for the samples from MRL

patients was obtained from NHS England Research Ethics Committee (18/LO/0738). The

ethical approval for the laboratory work for the Eritrean and MRL samples was obtained from

LSHTM Ethical Review Committee (#11979 and #14710 respectively).

Results

Primer and probe selection, and *in silico* analysis.

Initial assessment of *pfhrp3* primers across exons 1 and 2 showed cross reactivity with *pfhrp2* target (Figure S3) and a new set of primers within a specific conserved region of exon2 of *pfhrp3* was designed. After initial assessment *pfhrp2 and pfhrp3* primers for specificity and length of the probe, two sets of *pfhrp3* primers and three sets of *pfhrp2* primers, including primers with modifications at the 3' end, were selected for testing (Table S1). For *pfhrp2* primers, of the nine combinations used, the lowest C_q value was obtained when *pfhrp2*_F1 and *pfhrp2*_R2 (modification at 3' end) were combined and were selected for further optimization (Table S2). Since one single mutation in *pfhrp2* forward primer was found in one sample in The Gambia and three samples in Ghana we have nucleotide redundancy in the synthesis of *pfhrp2*_F1 primer to reflect these mutations (Table S2). The other *pfhrp2* primer combinations either produced fluorescence signal in Dd2 (*pfhrp2*-deleted laboratory strain) due to cross binding to *pfhrp3* or generated relatively higher (more unfavourable) C_q values in *pfhrp2*-positive lab strains (HB3 and Pf INT) compared to the selected primer combinations (Figure S4). Interestingly, a single nucleotide change (T to G) decreased the C_q value by 7 (30 to 23) while two nucleotide changes (T to G and T to G) decreased the C_q value by 2 (30 to 28) (Table S2).

Detection of *pfhrp2* and *pfhrp3* in laboratory strains

We followed the MIQE guidelines for optimization of the qPCR assay; for analysis and reporting of the data (19). The analytical specificity and sensitivity of each newly designed primer and probe combination was validated in monoplex and multiplex qPCR assays on Dd2, HB3, 3BD5 and 3D7, laboratory strains and Pf INT. The assay correctly determined the *pfhrp2* and *pfhrp3* status of the four laboratory strains and the Pf INT. No amplification of either *pfhrp2* or *pfhrp3* was observed in Dd2 and HB3 respectively, while 3BD5 produced no fluorescence signal in either the *pfhrp2* or *pfhrp3* channels (Figure 1). 3D7 and Pf INT were *pfhrp2* and *pfhrp3* positive.

Figure 1. Amplification of four laboratory clones (3D7, Dd2, HB3 and 3BD5) and Pf INT in three parasite targets (*pfhrp2*, *pfhrp3* and *pfldh*) and a human beta tubulin gene (HumTuBB). Laboratory clones 3D7 (wild type), Dd2 (*pfhrp2* deletion), HB3 (*hrp3* deletion), 3BD5 (both *pfhrp2/3* deletion) and Pf INT (*Plasmodium falciparum* WHO International Standard, both *pfhrp2/3* present) were amplified in triplicate targeting four different genes; *pfhrp2*, *pfhrp3*, *pfldh* and *HumTuBB*. The red horizontal line marks the threshold, the normalised fluorescence is measured on the y-axis and the number of cycles on the x-axis. Orange arrows point to different clones and the negative no template control (NTC).

Analytical sensitivity of the qPCR assay

The *pfhrp2*, *pfhrp3* and *pfldh* qPCR assays allowed detection of three parasites per μl with a C_q standard deviation (SD) of 0.58, 0.47, 0.41 respectively (Table 1A), which corresponds to parasite density CVs of 9.4%, 9.3% and 8.9% of respectively (Table 1B). The sample with dilution of the 1.5 parasite per μl showed a C_q SD of 0.80, 0.71 and 0.74, which corresponds to parasite density CVs of 32.6%, 26.1% and 28.1% respectively (Table S4). Though the assay also detected as low as 0.76 parasites per μl the SD value was very high (1.49, 1.37

and 1.48 respectively), and this corresponds to parasite density of CVs of 109%, 99% and

108% (Table S4). Therefore, the lowest parasite density that can be quantified with CV of

35% lies between 1.5 and 0.76 parasites per µl.

Table 1A: Precision of the parasite target genes: Mean, standard deviation (SD) of

quantification cycle (C_q) were calculated from amplifications of seven Pf INT 4-fold

dilutions (in triplicate) with *pfhrp2*, *pfhrp3* and *pfldh* assays.

Mean C_q values						
Parasite density (p/µl)	*pfhrp2*		*pfhrp3*		*pfldh*	
	Mean	SD	Mean	SD	Mean	SD
12500	23.18	0.06	23.46	0.02	22.44	0.04
3125	25.49	0.02	25.82	0.19	24.77	0.06
781	27.11	0.07	27.55	0.07	26.48	0.08
195	29.02	0.07	29.54	0.12	28.33	0.16
49	31.28	0.59	32.16	0.73	30.81	0.77
12	33.17	0.66	33.73	0.54	32.40	0.57
3	35.36	0.58	34.76	0.47	35.12	0.41
Total variance	28.95	0.15	29.31	0.18	28.32	0.17

286 **Table 1B: Precision of the parasite target genes: Mean, standard deviation (SD) of**

287 **coefficient of variation (CV) parasite density were calculated from amplifications of**

288 **seven Pf INT 4-fold dilutions (in triplicate) with *pfhrp2*, *pfhrp3* and *pfldh* assays.**

Estimated parasite density (parasite per µl)									
Parasite density (p/µl)	*pfhrp2*			*pfhrp3*			*pfldh*		
	Mean	SD	CV	Mean	SD	CV	Mean	SD	CV
12500	11220.33	544.65	4.85	12137.31	776.08	6.39	13405.10	658.97	4.92
3125	2965.02	215.30	7.26	3098.08	196.15	6.33	3493.59	315.34	9.03
781	902.11	71.29	7.90	879.47	68.65	7.81	1003.27	80.72	8.05
195	197.70	16.65	8.42	181.92	13.88	7.63	229.74	22.57	9.82
49	47.96	5.81	8.43	34.23	2.81	8.20	47.41	3.52	7.42
12	14.23	1.28	8.99	12.76	0.93	7.29	17.35	1.30	7.48
3	2.24	0.21	9.40	4.49	0.42	9.29	1.90	0.17	8.88
Total variance	190.40	15.53	7.74	197.70	14.83	7.50	208.08	16.19	7.78

PCR efficiency and dynamic range

In order to measure abundance of *pfhrp2* and *pfhrp3* clones relative to *pfldh* and estimate

parasite density relative to human DNA, the four targets should have similar amplification

efficiency. The efficiency of each qPCR was evaluated on 4-fold serial dilutions of Pf INT

(from 12500 to 3 parasite per µl) and each primer and probe combination resulted in similar

PCR efficiency (Figure 2) and this was evident by the C_q value generated by each

combination. A 4-fold dilution of the Pf INT produced linear standard plots with 98%, 96%,

98% and 1.07% PCR amplification efficiency for *pfhrp2*, *pfhrp3*, *pfldh* and *HumTub* primer

pairs respectively (Figure 2). The coefficient of determination (R^2) of the standard curve was

0.96-0.98 with a slope value of -3.375 to -3.416 for each assay.

Figure 2. Standard curve of 7 Pf INT samples diluted 4-fold starting from 12500
parasites per µl. The three parasite assays (*pfhrp2*, *pfhrp3* and *pfldh*) detected as low
as 3 parasites per microliter with amplification efficiency of 98%, 97% and 96% and
coefficient of determination (R2) -3.37, -3.39 and -3.43 respectively.

Performance of the qPCR assays

We assessed the precision of the assay by testing seven 4-fold dilution series of the Pf INT
(starting concentration, 12500 parasites per µl) in triplicate on three separate occasions. The
mean SD for cycle quantification across the seven samples was 0.15, 0.18 and 0.17 for
pfhrp2, *pfhrp3* and *pfldh* target genes respectively (Table 1A), which corresponds to a
calculated parasite density of CVs of 7.4%, 7.5% and 7.8% respectively (Table 1B). The
sensitivity and specificity as well as the robustness of the assay was assessed using *P.
falciparum*-negative blood samples and Pf INT (3 and 1.5 parasites per µl) in eight replicates
on three different occasions. All the control samples were negative, while the Pf INT was
positive indicating the absence of amplification inhibition and non-specific amplification
(Table S4).

Detecting parasites with *pfhrp2/3* deletions hidden in polyclonal infections

The ability of the qPCR assay to detect minor or major laboratory clones with *pfhrp2/3*
deletion was assessed using artificially mixed laboratory strains Dd2, HB3, 3BD5 and Pf INT.
Both *pfhrp2* and *pfhrp3* qPCR assays showed good performance in detecting minor (as low
as 20%; CI, 7.94-27.05) and major (as high as 99.99%; CI, 99.99-100) *pfhrp2/3*-deleted
clones in the artificially mixed laboratory clones (Table S4). The *pfhrp2/3* assays can also
detect as low as 10% *pfhrp2/3* deleted clones but the value lies within the confidence interval
of replicate experiment of a single clone sample and therefore lacks confidence.

Workflow and throughput time

Performing the multiplex qPCR requires three steps: approximately 20 minutes for reaction
setup takes ~ 20 minutes, two hours for runtime 30 minutes for analysis. Over all the
estimated time was three hours for 72 reactions on the Rotorgene Q platform. In
comparison, the estimated time for the conventional method, deploying several nested PCR
assays in a 96-well plate format followed by electrophoresis is approximately 30 hours. Time
for DNA extraction is the same for both.

Validation and application of the qPCR assay on field samples

To determine the diagnostic sensitivity and specificity of the qPCR assay for detecting

pfhrp2/3 deletions in field samples and to provide population estimates of deletion

prevalence, we first investigated 50 DNA samples obtained from confirmed *P. falciparum*

patients from Eritrea whose *pfhrp2/3* deletions were previously determined using the

conventional nPCR method (10). Results obtained from the qPCR assay were fully

concordant with previously reported results from the same samples using the conventional

nPCR method (10). The qPCR assay correctly detected all the *pfhrp2/3* positives (100%

sensitivity for *pfhrp2*, n=19, 95% CI = (82.4%, 100%); 100% sensitivity for *pfhrp3*, 95% CI=

(66.4%; 100%), n=9) and accurately determined the absence of *pfhrp2/3* in the remaining

samples (100% specificity for *pfhrp2,* n=31, 95% CI=(88.8%,100%); 100% specificity for

pfhrp3, n=41, 95% CI=(91.4%,100%)) (Table 2). We then assessed samples obtained from

clinical malaria patients from the MRL (n=113), Kenya (n=150) and Tanzania (n=149). For

the MRL samples, we selected 113 samples from countries with reported *pfhrp*2/3 deletion

or high risk of emergence of *pfhrp*2/3 deletion (22). The countries include Eritrea (n=1),

Ethiopia (n=2), Kenya (n=20), Tanzania (n=16), Uganda (n=27), Sudan (n=29), South Sudan

(n=10), Djibouti (n=1), Somalia (3) and east Africa (unknown country, n=2). Analysis of the

pfhrp2/3 status of the MRL samples by qPCR showed 1.8%, 1.8% and 6.3% *pfhrp2-/3+*,

pfhrp2+/3- and *pfhrp2-/3-* deletions respectively. One *pfhrp2* deletion occurred in a UK

traveller from Sudan and five *pfhrp3* deletions occurred in UK travellers from Ethiopia (1),

Sudan (1), South Sudan (2) and Uganda (1). There was evidence of *pfhrp2/3* deletions in

polyclonal infections in 3.5% of MRL samples in total (Table 4). Of the 149 whole blood

samples collected from Tanzania, 1 sample (0.7%) carried a *pfhrp2* deletions and another

sample (0.7%) carried a *pfhrp3* deletion while no deletions were observed in the 150 Kenyan

samples. However, there were 5 samples (3.4%) carrying *pfhrp2*-deleted strains hidden in

polyclonal infections in the Kenyan samples and one (0.7%) in the Tanzanian samples.

Details of the deletions in each country are shown in Table 2.

Table 2: Prevalence of *pfhrp2* and *pfhrp3* deletions including in polyclonal infections in clinical samples.

DNA source	Country	*pfhrp2* and/or *pfhrp3* deletions in samples			*pfhrp2/3* deletions in polyclonal infections		
		pfhrp2 n(%)	*pfhrp3* n(%)	*pfhp2/3* n%)	*pfhrp2* n(%)	*pfhrp3* n(%)	*pfhp2/3* n(%)
Eritrea	Eritrea (n=50)	0	7 (14)	32 (64)	3(0.5)	4 (8)	0
HTD	Djibouti (n=1)	0	0	0	0	0	0
	Kenya (n=20)	0	0	0	1 (5)	1 (5)	0
	Eritrea (n=1)	0	0	0	0	0	1 (100)
	Ethiopia (n=2)	0	1 (50)	0	0	0	0
	Somalia (n=3)	0	0	0	0	(2)66.7	0
	South Sudan (n=10)	0	3(30)	0	1 (10)	0	0
	Sudan (n=29)	1 (3.5)	1 (3.5)	1 (3.5)	0	3 (10.3)	0
	Tanzania (n=16)	0	0	0	0	1 (6.3)	0
	Uganda (n=27)	0	1 (3.7)	0	2 (7.4)	3 (11.2)	0
	East Africa (n=2)	0	0	0	0	0	0
Kenya/ Tanzania	Kenya (n=150)	0	0	0	5 (3.4)	0	0
	Tanzania (n=149)	1 (0.7)	1 (0.7)	0	1 (0.7)	0	0

Estimation of parasite density using *pfldh*

As well as a positive control for parasite DNA quality, the *pfldh* gene was simultaneously

used as a parasite target for estimation of parasite density using relative quantification. After

assessing its sensitivity, specificity and amplification efficiency the usefulness of the qPCR

as an estimator of parasite density was evaluated by comparison to the previously published

duplex *pgmet* qPCR assay (18). We performed this comparison using 50 DNA samples from

Eritrea and the *pfldh* qPCR showed high degree of correlation (R^2=0.95) with the duplex

pgmet qPCR (Figure 3). The *pgmet* qPCR generated relatively higher parasite density

compared to the *pfldh* qPCR and this is expected as the former targets multi-copy genes in

the apicoplast genome, whereas the latter targets is a single-copy gene.

Figure 3: Comparing parasite density estimates produced by *pfldh* and *pgmet* qPCR methods using 50 Eritrean samples. The estimation was done using delta delta relative quantification methods in the presence of *HumTuBB* gene as a normalizer and Pf INT as a calibrator. The two qPCR methods showed strong agreement (R^2 = 0.95)

Discussion

In this study, we describe the development, validation and application of a high-throughput multiplex qPCR assay for simultaneous determination of *P. falciparum* with *pfhrp2* and *pfhrp3* deletion genotypes in mono- and polyclonal infections and estimation of parasite density. This highly sensitive and specific method enables accurate and robust assessment of parasite density between 1.5 and 0.76 parasites per µl with a CV of 35% (21). The qPCR detection of each of the targets described here was achieved with high efficiency, R^2 values greater than 0.96 and very low standard deviation among replicates of each dilution. An additional strength of the assay is the detection of *pfhrp2/3* deletions at relative abundance as low as 20% and as high as 99% in mixed infections, simulated by mixture of cultured parasites. This is particularly relevant in moderate to high transmission settings, where such parasites have already emerged but can be masked in polyclonal infections. The assay can be used as a tool to monitor changes in frequency of deletions providing potential early

warning signs of emergence of *P. falciparum* with *pfhrp2/3* deletions. This would allow

appropriate measures to be taken to identify, respond and contain the spread of such

parasites before they become sufficiently abundant to impact on case management and

malaria control programs as has already occurred independently in South America and

Eritrea (12, 23).

The design of the multiplex qPCR assay provides several advantages over existing detection

methods (5, 14, 24). Firstly, the qPCR assay uses only one parasite target to confirm the

presence of DNA and to assess its quality while the gel-electrophoresis based nested PCR

(nPCR) methods use three target genes (5, 25, 26). This reduces the cost and throughput

time and simplifies the algorithm for interpreting results. Secondly, the parasite target gene

(*pfldh*) used for DNA confirmation in the qPCR assay has a single copy, generating

equivalent sensitivity to the *pfhrp2/3* targets. When multi-copy genes (e.g., *18SrDNA* and

cytb) are used in the nPCR and other qPCR assays there is increased risk of false-*pfhrp2/3*

deletion calls due to sensitivity differences with the single copy *pfhrp2/3* genes (14, 27). It is

difficult to evaluate the accuracy of *pfhrp2/3* deletion calls in the literature, as most

laboratories do not report the limit of detection of the nPCR methods used. Thirdly, the

qPCR assay uses a human house-keeping gene as internal control. Variability in DNA yield,

or loss, introduced during sample collection or parasite DNA extraction or qPCR

amplification can thus be corrected for, reducing the risk of false-*pfhrp2/3* deletion calls.

Using conventional methods, those samples negative by 18SrDNA may be excluded from

further *pfhrp2/3* deletion analysis as they are presumed parasite negatives, potentially

underestimating the prevalence of *pfhrp2/3* deletions by missing those samples where

technical failure has caused this outcome. Fourthly, the multiplex qPCR assay accurately

detects laboratory *P. falciparum* strains and clinical samples with *pfhrp2/3* deletions when

mixed as minor or major clones. The ability of the multiplex qPCR assay to detect *pfhrp2/3*

deletions in samples with low parasitaemia and in polyclonal infections is made possible due

to three unique features: the choice of a single copy parasite gene (*pfldh*) for DNA quality

confirmation; the inclusion of a human gene for normalisation and the modification of the

primers. Finally, the qPCR assay can also estimate relative parasite density using the

human gene as a normalizer and Pf INT as a calibrator. This characteristic of the assay is

useful because one of the criteria for confirming deletion of *pfhrp2/3* is estimation of parasite

density using microscopy to rule out low parasite density as a factor for lack of parasite

target amplification (14). However, microscopy is not always performed during community

surveys, and quantification of parasite density using qPCR is required (5, 13). The qPCR

assay not only detects *P. falciparum parasites* with *pfhrp2/3* deletions but also

simultaneously estimates parasite density in the same experiments, hence reducing time

and cost.

Our study shows the presence of *pfhrp2/3* deletions in infected UK travellers for the first

time. While *pfhrp2/3* deletions were previously reported in Eritrea (10), Ethiopia (28) and

Uganda (29) this is the first time such deletions are reported in Sudan and South Sudan,

though a negative *pfhrp2* result was reported in Sudan (30). Interestingly, *pfhrp2* and *pfhrp3*

deletions were also detected in polyclonal infections in Kenya, Eritrea, Somalia, South

Sudan, Sudan and Uganda. This suggests the circulation of low frequency *pfhrp2*-deleted

parasites in Somalia as minor strains in mixed infections. WHO recommends surveillance to

determine the prevalence of *pfhrp2/3* deletions occur and in neighbouring areas if the

prevalence of *pfhrp2* gene deletions that cause false-negative HRP2-based RDT results in a

representative sample is higher than 5%, HRP2-based RDTs should be replaced with

alternative *P. falciparum* diagnostic tool that is not exclusively reliant on detection of HRP2

(11). If the prevalence is below 5% a repeat of the survey is recommended in 1-2 years and

the detection of *pfhrp2/3* deletions in polyclonal infections by the qPCR assay could be used

to inform decisions about how soon to repeat the survey. For example, if the *pfhrp2/3*

deletions in polyclonal infections occur in medium to high transmission endemic settings, it

may be preferable to survey during the dry season when the multiplicity of infection is lower

and would allow accurate estimation of the *pfhrp2/3* deletions.

Limitations

The challenges of confirming the absence of a gene target require careful attention to lab workflow and DNA quality. First, the sensitivity of the qPCR assay demands careful laboratory workflows that prevent contamination. This is true of all qPCR assays but particularly important for discrimination of low-concentration deleted strains in polyclonal infections. Second, while the qPCR assay was carefully designed and optimized to avoid cross-binding, use of appropriate DNA controls is needed to monitor for unintentional amplification of *pfhrp3* by *pfhrp2* primers and vice versa. We recommend including at least two parasite negative controls (preferably Dd2 and HB3) for *pfhrp2* and *pfhrp3,* respectively, in each experiment. The use of only a double-deleted strain (such as 3BD5) is not recommended. In addition, the qPCR assay targets only one additional single-copy parasite gene while the conventional methods for *pfhrp2* and *pfhrp3* genotyping have employed three independent parasite genes to ensure DNA quality and rule out DNA degradation (14). Because the *pfhrp2* and *pfhrp3* qPCR amplicon lengths are shorter (98bp and 84bp respectively) than typical amplicons generated by the conventional method (~ 300 - 800bp), detection of these targets is expected to be more reliable. If the frequency of deletions is observed to be markedly increased near the limits of detection of the assay in a particular study, then confirmatory testing using a second gene target could be considered. Finally, due to limited conserved regions, the *pfhrp2*-specific primer covered known variant positions present in minority of field samples in the MalariaGEN genome data. The qPCR assay was optimized taking into account the known sequence variants, and the use of nucleotide redundancies in the primer synthesis did not affect the yield in florescence signal (Cq value). The variants within our primer sequence are relatively fewer compared to the primer and probe sequences of other recently published qPCR assays (27, 31), which we predict may be challenged by sequence variation (mutations, insertions and deletion) in the primer and probe sequences of *pfhrp2* and *pfhrp3* found in field samples from seven and nine countries respectively (Table S6).

Conclusion

Our qPCR method for detection of *pfhrp2/3* deletions is a robust alternative with several advantages over existing approaches. The qPCR assay has superior performance to existing methods in speed, cost and ease of interpretation in detecting *pfhrp2/3*-deleted *P. falciparum* parasites from DNA derived from whole blood or filter-paper bloodspots. Data from screening endemic country samples in returning travellers to the United Kingdom suggest systematic surveillance of *pfhrp2/3* deletions in Ethiopia, Sudan and South Sudan is warranted. Careful monitoring of *pfhrp2/3* deletions in Somalia will also be required as the emergence of *pfhrp2/pfhrp3*-deleted parasites as a single-clone infection may soon occur. Based on the findings in this report and elsewhere in the literature, *pfhrp2/3* deletions are present in 31 countries but the scale and scope is still not well elucidated and efforts to dramatically scale up surveillance are needed (32, 33). This qPCR assay can accurately and efficiently support surveillance efforts so that endemic countries have the data required to guide policy on RDT procurement and avert a serious public health threat.

Declarations

Author contributions

LG, NS and KBB conceived/designed the study. DN, AB, SM, KM, JJJ, JBP, PC, JC and CJS conceived/designed the individual studies. DAS performed the parasite culturing. JP and SC carried out the genome analysis. LG and RK conducted the initial experiments of the study. KBB conducted the final primer and probe design and analysis, wrote the study protocol and established the assay. LG and KBB conducted the experiments, analysed the results and wrote the first draft. LG, DN, NS, DAS, JBP, PC, JC, CJS, CD and KBB contributed to the reviewing of the manuscript. LG and KB wrote the final manuscript. All authors read and approved the manuscript.

Ethical approval and consent to participate

All data included in this analysis were obtained in accordance with ethical approvals from country of origin. The data are fully anonymised and cannot be traced back to identifiable individuals. A separate ethical approval was given by London School of Hygiene & Tropical Medicine ethical review committee to conduct this study on the Eritrean samples.

Acknowledgement

We thank all the patients and staff who participated in this study. We thank Helena Stone for her support with the Malaria Reference Laboratory samples. We also thank Oksana Kharabora for assistance in the lab with the Kenyan and Tanzanian samples. We thank Qin Cheng (ADFMIDI/QIMR Berghofer) sharing with us the *pfhrp2/3* data generated by conventional PCR to validate our qPCR assay.

Funding

This study was partly supported by Department of Infection Biology Rosemary Weir Research Prize to LG, NS and KBB. KBB is funded by LSHTM ISSF3 Pump-Priming. DN is funded by the PHE Malaria Reference Laboratory. RDK is supported by the DELTAS Africa Initiative grant # DEL-15-011 to THRiVE-2. Collection of samples from Kenya and Tanzania was supported by the National Institutes of Health (R01AI121558) to JJJ. The Malaria Reference Laboratory is funded by Public Health England.

Availability of data and materials

Data are available from the corresponding author with reasonable request.

Consent for publication

Not applicable

Competing interest

JBP reports non-financial support in the form of in-kind donation of laboratory testing and reagents from Abbott Laboratories for studies of viral hepatitis and financial support from the World Health Organization. Other authors declare no conflict of interest.

Contributor information

Khalid B Beshir, email: khalid.beshir@lshtm.ac.uk

References

1.	WHO. *World Malaria Report 2019.* Geneva: World Health Organization. 2019.
2.	WHO. Malaria rapid diagnostic test performance. Results of WHO product testing of malaria RDTs: round 8 (2016-2018). 2018.
3.	Baker J, Gatton ML, Peters J, Ho MF, McCarthy JS, Cheng Q. Transcription and expression of Plasmodium falciparum histidine-rich proteins in different stages and strains: implications for rapid diagnostic tests. PLoS One. 2011;6(7):e22593.
4.	Lee N, Baker J, Andrews KT, Gatton ML, Bell D, Cheng Q, et al. Effect of sequence variation in Plasmodium falciparum histidine- rich protein 2 on binding of specific monoclonal antibodies: Implications for rapid diagnostic tests for malaria. J Clin Microbiol. 2006;44(8):2773-8.
5.	Beshir KB, Sepulveda N, Bharmal J, Robinson A, Mwanguzi J, Busula AO, et al. Plasmodium falciparum parasites with histidine-rich protein 2 (pfhrp2) and pfhrp3 gene deletions in two endemic regions of Kenya. Sci Rep. 2017;7(1):14718.
6.	Sepulveda N, Phelan J, Diez-Benavente E, Campino S, Clark TG, Hopkins H, et al. Global analysis of Plasmodium falciparum histidine-rich protein-2 (pfhrp2) and pfhrp3 gene deletions using whole-genome sequencing data and meta-analysis. Infect Genet Evol. 2018;62:211-9.
7.	Thomson R, Beshir KB, Cunningham J, Baiden F, Bharmal J, Bruxvoort KJ, et al. pfhrp2 and pfhrp3 Gene Deletions That Affect Malaria Rapid Diagnostic Tests for Plasmodium falciparum: Analysis of Archived Blood Samples From 3 African Countries. J Infect Dis. 2019;220(9):1444-52.
8.	Dorado EJ, Okoth SA, Montenegro LM, Diaz G, Barnwell JW, Udhayakumar V, et al. Genetic Characterisation of Plasmodium falciparum Isolates with Deletion of the pfhrp2 and/or pfhrp3 Genes in Colombia: The Amazon Region, a Challenge for Malaria Diagnosis and Control. PLoS One. 2016;11(9):e0163137.
9.	Berhane A, Russom M, Bahta I, Hagos F, Ghirmai M, Uqubay S. Rapid diagnostic tests failing to detect Plasmodium falciparum infections in Eritrea: an investigation of reported false negative RDT results. Malar J. 2017;16(1):105.
10.	Berhane A, Anderson K, Mihreteab S, Gresty K, Rogier E, Mohamed S, et al. Major Threat to Malaria Control Programs by Plasmodium falciparum Lacking Histidine-Rich Protein 2, Eritrea. Emerg Infect Dis. 2018;24(3):462-70.
11.	WHO. Protocol for estimating the prevalence of pfhrp2/pfhrp3 gene deletions among symptomatic falciparum patients with false-negative RDT results. https://www.who.int/malaria/publications/atoz/hrp2-deletion-protocol/en/ Accessed 21 January 2020. 2018(WHO/CDS/GMP/2018.03).
12.	Gamboa D, Ho MF, Bendezu J, Torres K, Chiodini PL, Barnwell JW, et al. A large proportion of P. falciparum isolates in the Amazon region of Peru lack pfhrp2 and pfhrp3: implications for malaria rapid diagnostic tests. PLoS One. 2010;5(1):e8091.
13.	Parr JB, Verity R, Doctor SM, Janko M, Carey-Ewend K, Turman BJ, et al. Pfhrp2-Deleted Plasmodium falciparum Parasites in the Democratic Republic of the Congo: A National Cross-sectional Survey. J Infect Dis. 2017;216(1):36-44.
14.	Cheng Q, Gatton ML, Barnwell J, Chiodini P, McCarthy J, Bell D, et al. Plasmodium falciparum parasites lacking histidine-rich protein 2 and 3: a review and recommendations for accurate reporting. Malar J. 2014;13:283.
15.	Beshir KB, Diallo N, Sutherland CJ. Identifying Recrudescent Plasmodium falciparum in Treated Malaria Patients by Real-time PCR and High Resolution Melt Analysis of Genetic Diversity. Sci Rep. 2018;8(1):10097.
16.	van Schalkwyk DA, Burrow R, Henriques G, Gadalla NB, Beshir KB, Hasford C, et al. Culture-adapted Plasmodium falciparum isolates from UK travellers: in vitro drug sensitivity, clonality and drug resistance markers. Malar J. 2013;12:320.

17. Robinson A, Busula AO, Muwanguzi JK, Powers SJ, Masiga DK, Bousema T, et al.
Molecular quantification of Plasmodium parasite density from the blood retained in used
RDTs. Sci Rep. 2019;9(1):5107.
18. Beshir KB, Hallett RL, Eziefula AC, Bailey R, Watson J, Wright SG, et al. Measuring
the efficacy of anti-malarial drugs in vivo: quantitative PCR measurement of parasite
clearance. Malar J. 2010;9:312.
19. Bustin SA, Benes V, Garson JA, Hellemans J, Huggett J, Kubista M, et al. The MIQE
guidelines: minimum information for publication of quantitative real-time PCR experiments.
Clin Chem. 2009;55(4):611-22.
20. Beshir KB, Diallo N, Sutherland CJ. Identifying Recrudescent Plasmodium falciparum
in Treated Malaria Patients by Real-time PCR and High Resolution Melt Analysis of Genetic
Diversity. Scientific Reports. 2018;8(1):10097.
21. Forootan A, Sjoback R, Bjorkman J, Sjogreen B, Linz L, Kubista M. Methods to
determine limit of detection and limit of quantification in quantitative real-time PCR (qPCR).
Biomol Detect Quantif. 2017;12:1-6.
22. Gatton ML, Dunn J, Chaudhry A, Ciketic S, Cunningham J, Cheng Q. Use of
PfHRP2-only RDTs rapidly select for PfHRP2-negative parasites with serious implications for
malaria case management and control. J Infect Dis. 2017.
23. Berhane A MS, Mohammed S,, Embaye G HF, Zehaie A, , Chinorumba A CJ. Pfhrp2
detecting malaria rdts: Alarming false negative results in eritrea,. In: Program and Absracts
of the 65th Conference on American Soceity for Tropical Medicine and Hygiene, Atlanta.
2016;879.
24. Parr JB, Anderson O, Juliano JJ, Meshnick SR. Streamlined, PCR-based testing for
pfhrp2- and pfhrp3-negative Plasmodium falciparum. Malar J. 2018;17(1):137.
25. Parr JB, Verity R, Doctor SM, Janko M, Carey-Ewend K, Turman BJ, et al. Pfhrp2-
Deleted Plasmodium falciparum Parasites in the Democratic Republic of the Congo: A
National Cross-sectional Survey. J Infect Dis. 2016.
26. Kumar N, Pande V, Bhatt RM, Shah NK, Mishra N, Srivastava B, et al. Genetic
deletion of HRP2 and HRP3 in Indian Plasmodium falciparum population and false negative
malaria rapid diagnostic test. Acta Trop. 2013;125(1):119-21.
27. Kreidenweiss A, Trauner F, Rodi M, Koehne E, Held J, Wyndorps L, et al. Monitoring
the threatened utility of malaria rapid diagnostic tests by novel high-throughput detection of
Plasmodium falciparum hrp2 and hrp3 deletions: A cross-sectional, diagnostic accuracy
study. EBioMedicine. 2019;50:14-22.
28. Girma S, Cheaveau J, Mohon AN, Marasinghe D, Legese R, Balasingam N, et al.
Prevalence and Epidemiological Characteristics of Asymptomatic Malaria Based on
Ultrasensitive Diagnostics: A Cross-sectional Study. Clin Infect Dis. 2019;69(6):1003-10.
29. Thomson R, Beshir KB, Cunningham J, Baiden F, Bharmal J, Bruxvoort KJ, et al.
pfhrp2 and pfhrp3 gene deletions that affect malaria rapid diagnostic tests for Plasmodium
falciparum: analysis of archived blood samples from three African countries. J Infect Dis.
2019.
30. Mussa A, Talib M, Mohamed Z, Hajissa K. Genetic diversity of Plasmodium
falciparum histidine-rich protein 2 (PfHRP2) and its effect on the performance of PfHRP2-
based rapid diagnostic tests. BMC Res Notes. 2019;12(1):334.
31. Schindler T, Deal AC, Fink M, Guirou E, Moser KA, Mwakasungula SM, et al. A
multiplex qPCR approach for detection of pfhrp2 and pfhrp3 gene deletions in multiple strain
infections of Plasmodium falciparum. Sci Rep. 2019;9(1):13107.
32. WHO. Malaria Threats Map: visualizing biological challenges to malaria control and
elimination https://www.who.int/malaria/maps/threats-about/en/ Accessed on 31 January
2020. 2020.
33. WHO. Response plan to pfhrp2 gene deletions.
https://apps.who.int/iris/bitstream/handle/10665/325528/WHO-CDS-GMP-2019.02-eng.pdf.
Accessed on 31 January, 2020. 2019.
645