Axonal ER Ca²⁺ Release Enhances Miniature, but Reduces Activity-Dependent Glutamate Release in a Huntington Disease Model

James P. Mackay[1], Amy I. Smith-Dijak[1,2], Ellen T. Koch[1,2], Peng Zhang[1], Evan Fung, Wissam B. Nassrallah[1,3], Caodu Buren[1,2], Mandi Schmidt[2,4], Michael R. Hayden[4] and *Lynn A. Raymond[1]

[1] Department of Psychiatry, Djavad Mowafaghian Centre for Brain Health,

[2] Graduate Program in Neuroscience

[3] MD/PhD Program

[4] Centre for Molecular Medicine and Therapeutics, BC Children's Hospital Research Institute

University of British Columbia, Vancouver, British Columbia, Canada

*Correspondence addressed to:

Lynn A. Raymond, MD, PhD

Department of Psychiatry and Djavad Mowafaghian Centre for Brain Health

University of British Columbia

4834-2255 Wesbrook Mall

Vancouver, BC Canada V6T 1Z3

Lynn.raymond@ubc.ca

Abstract

Action potential-independent (miniature) neurotransmission occurs at all chemical synapses, but

remains poorly understood, particularly in pathological contexts. Spontaneous release of Ca^{2+} from

the axonal endoplasmic reticulum (ER) is thought to facilitated miniature neurotransmission, and

aberrant ER Ca^{2+} handling is notably implicated in the progression of Huntington's disease (HD)

and other neurodegenerative diseases. Here, we report elevated glutamate-mediated miniature

synaptic event frequencies in YAC128 (HD-model) cortical neurons, which pharmacological

experiments suggest is mediated by enhanced spontaneous ER Ca^{2+} release. Calcium imaging

using an axon-localized sensor revealed slow action potential (AP)-independent axonal Ca^{2+}

waves, which were more common in YAC128 cortical neurons. Conversely, spontaneous axonal

ER Ca^{2+} release was associated with reduced AP-dependent axonal Ca^{2+} events and consequent

glutamate release. Together, our results suggest spontaneous release of axonal ER Ca^{2+} stores

oppositely regulates activity-dependent and -independent neurotransmitter release in HD, with

potential implications for the fidelity and plasticity of cortical excitatory signaling.

Introduction

Huntington's disease (HD) is a fatal, autosomal dominantly-inherited neurodegenerative disorder

caused by a polyglutamine-encoding CAG repeat-expansion (>35 repeats) in exon-1 of the

huntingtin gene ("A novel gene containing a trinucleotide repeat that is expanded and unstable on

Huntington's disease chromosomes. The Huntington's Disease Collaborative Research Group.,"

1993). Disease onset is typically in middle age, characterized by progressively disordered

movement and declining cognition (Bachoud-Lévi et al., 2019). Although the mutant huntingtin

protein (mHTT) is widely expressed, GABAergic spiny projection neurons (SPN)s of the striatum

and pyramidal neurons of the cerebral cortex show the most severe degeneration in HD

(Graveland, Williams, & DiFiglia, 1985; Vonsattel et al., 1985). Cortical glutamatergic afferents

extensively innervate striatal SPNs and dysfunction at these synapses is thought to precede overt

neuron loss in HD (Raymond et al., 2011).

Glutamate-mediated toxicity was initially proposed to contribute to pathogenesis in HD based on

studies showing that striatal injections of glutamate receptor agonists largely recapitulate HD

pathology in animals (Beal et al., 1986; Hantraye, Riche, Maziere, & Isacson, 1990). More

recently, studies in SPNs from transgenic and knock-in mouse models of HD demonstrate

enhanced extrasynaptic N-Methyl-D-Aspartate receptor (NMDAR) surface expression and

function that may explain, in part, increased susceptibility to excitotoxic challenges (Botelho et

al., 2014; Fan, Fernandes, Zhang, Hayden, & Raymond, 2007; Kovalenko et al., 2018; Milnerwood

et al., 2010; Plotkin et al., 2014; Zeron et al., 2002). Cell stress/death signaling mediated by

elevated levels of extrasynaptic NMDARs in HD SPNs may be further exacerbated by reduced

glutamate transporter function (Miller et al., 2008), although recent studies show glutamate

clearance after synaptic release is normal or accelerated in striatal and cortical brain slice (Parsons

et al., 2016). Additionally, some studies show synaptic glutamate release from cortical afferents is

altered in HD mouse models, possibly contributing to excitotoxicity and synaptic dysfunction

(Cepeda et al., 2003; Joshi et al., 2009; Raymond et al., 2011). However, the direction of this effect

appears to be model and disease-stage dependent; relative to wildtype (WT), glutamate release is

typically enhanced early, but reduced with disease progression in HD mice (Joshi et al., 2009).

Mutant HTT (mHTT) directly interacts with endoplasmic reticulum (ER) type-1 inositol (1,4,5)-

triphosphate receptors (IP3Rs) sensitizing their Ca^{2+} release in response to IP3 (Tang et al., 2003);

blocking this interaction normalizes IP3-induced ER Ca^{2+} release *in vitro* and improves behavioral

outcomes in HD-model mice (Tang, Guo, Wang, Chen, & Bezprozvanny, 2009). Furthermore,

evidence suggests Ryanodine receptors, which are ER-localized Ca^{2+} channels that mediate Ca^{2+}-

induced Ca^{2+} release, are constitutively leaky in HD mouse models (Suzuki, Nagai, Wada, &

Koike, 2012). Although these studies focused on the soma and dendrites of SPNs at the

postsynaptic side of cortical-striatal synapses, ER is found in all neuronal compartments, including

presynaptic terminals, where its Ca^{2+} stores, when released, modulate neurotransmission

(Emptage, Reid, & Fine, 2001; Llano et al., 2000). However, it is unknown whether presynaptic

ER Ca^{2+} handling is also dysfunctional in HD and if such a process contributes to altered glutamate

release from cortical synaptic terminals.

Here, we have investigated whether glutamate release is altered, and if ER Ca^{2+} dysregulation

contributes to aberrant function, in presynaptic terminals of cortical pyramidal neurons from

premanifest HD-model mice expressing full-length mHTT with 128 CAG-repeats in a yeast

artificial chromosome (YAC128). Our results from neuronal cultures and acute cortical-striatal

brain slices demonstrate a shift in balance favoring increased action potential (AP)-independent

(miniature) vs. AP-dependent glutamate release, and that altered ER Ca^{2+} release contributes to

this change. This may have important previously unidentified disease implications, given the

distinct physiologically relevant signaling potentially mediated by miniature neurotransmission

(Fishbein & Segal, 2007; Kavalali, 2015). Ultimately our data contribute to growing evidence for

cortex and striatal synaptic dysfunction in HD, and support a key role for ER Ca^{2+} dysregulation

in this pathophysiology.

Results

Mini EPSC frequencies are elevated in YAC128 cortical cultures at early time points

Our group previously recorded miniature excitatory post synaptic currents (mEPSCs) from striatal

spiny projection neurons (SPNs) co-cultured with cortical neurons from prenatal WT or yeast

artificial chromosome (YAC128) mice. YAC128 SPNs showed elevated mEPSC frequencies,

compared to WT, at day in vitro (DIV) 14 – a time point when SPN dendritic arborization patterns

and spine numbers were similar between genotypes (Buren, Parsons, Smith-Dijak, & Raymond,

2016). At DIV21, YAC128 SPNs showed a significant reduction in total dendritic length, and

therefore reduced total excitatory synapse numbers, compared to WT. Despite this, mEPSC

frequencies onto WT and YAC128 SPNs were comparable at DIV21.

The above results suggested a higher rate of action potential-independent glutamate release from

YAC128 presynaptic cortical pyramidal neuron (CPN) terminals onto individual SPN synapses.

Here, we capitalize on a relatively simpler culture preparation containing only cortex-derived

neurons and YAC128 mouse-derived brain slices to mechanistically dissect how mutant huntingtin

protein expression affects action potential-independent and -dependent glutamate release from

CPN terminals.

To establish whether miniature glutamate release is enhanced at YAC128 CPN terminals targeting

other CPNs, we first established WT and YAC128 CPN mEPSC parameters in cortical cultures at

various DIV ages (7, 14, 18 and 21). In general, both genotypes showed increased mEPSC

frequencies with culture time. However, YAC128 CPN mEPSC frequencies were consistently

higher than age-matched WT controls after DIV7 and up until DIV21, at which point mEPSC

frequencies became comparable between genotypes **(Figure 1),** as was also the case in the co-

culture model (Buren et al., 2016). The greatest genotype mEPSC frequency difference occurred

at DIV14, when mean YAC128 CPN frequencies were more than double that of WT: 10.83 ± 1.80

Hz vs 4.52 ± 0.60 Hz, respectively **(Figure 1D, E)**. YAC128 mEPSC frequencies remained

significantly higher than WT at DIV18: 14.07 ± 1.54 Hz vs 9.92 ± 1.54 Hz, respectively **(Figure**

1G, H). Age matched YAC128 and WT CPNs showed similar mEPSC amplitudes across all DIV

time points examined **(Figure 1 C, F, I and L)**. We chose to perform subsequent experiments

probing the mechanistic details underlying increased cortical mini glutamate release in DIV18

cortical monocultures, since this was the most mature culture stage at which mEPSC frequencies

were elevated in YAC128 CPNs.

Synapse numbers and dendritic complexity are similar in WT and YAC128 cortical pyramidal neurons at DIV 18

The higher mEPSC frequencies seen in DIV18 YAC128 CPNs suggest an increased presynaptic glutamate release probability. However, relative differences in synapse numbers could also account for this finding. To estimate numbers of synapses, we first expressed cytosolic GFP in a small proportion of neurons in WT and YAC128 cortical cultures and imaged full CPN dendritic arbors at DIV18. Sholl analysis revealed similar arborization patterns and total dendritic length in WT and YAC128 CPNs **(Supplemental Figure 1A, C)**. In separate cultures, we expressed an internal GFP-tagged anti-PSD95 antibody (Gross et al., 2013) in a subset of neurons and immuno-stained for VGlut1, to identify glutamatergic terminals, and the GluA2 AMPA receptor subunit, to identify functional synapses. Functional synapse numbers, defined as GFP-labeled PSD95 puncta colocalized with VGlut1 and GluA2 immunofluorescent-labeled puncta, were not significantly different between DIV18 WT and YAC128 CPNs, although there was a trend towards lower synapse density in YAC128 CPNs **(Supplemental Figure 1E, F)**. Together with the above data showing increased mEPSC frequency in YAC128 CPNs at DIV18 **(Figure 1)**, these results point to an increase in miniature vesicular glutamate release from cortical terminals in YAC128 cultures.

Releasing ER calcium with low dose ryanodine or caffeine increases the miniature EPSC frequency in WT, but not YAC128 cultures

Studies using mouse models suggest Ca^{2+} release from ER stores is aberrant in HD due to increased IP3 and ryanodine receptor activity (Suzuki et al., 2012; Tang et al., 2003). Although effects of mHTT on the presynaptic ER have not been specifically studied, we hypothesized a presynaptic ER leak elevates cytosolic Ca^{2+}, thereby mediating the increased miniature glutamate release seen in our YAC128 cultures **(Figure 2A, B)**. To test this hypothesis, we first recorded mEPSCs before and during local application of low dose (5 μM) ryanodine to cultured CPNs; 5 μM ryanodine releases ER Ca^{2+} by opening ryanodine receptors (Meissner, 2017). In WT cultures, 5 μM ryanodine nearly doubled the CPN mEPSC frequency: from 6.74 ± 1.17 Hz to 11.91 ± 2.36 Hz [Students' paired T-test; p=0.0280; n=13 cells] **(Figure 2C - E)**. In YAC128 cultures, however, ryanodine (5 μM) did not significantly alter the CPN mEPSC frequency: 11.98 ± 1.66 Hz vs 13.89 ± 2.01 Hz - under control conditions and in ryanodine (5 μM) respectively **(Figure 2F - H)**. A Comparison of the percent change in mEPSC frequency following ryanodine (5 μM) revealed a significantly greater response in WT than in YAC128 CPNs [31.7 ± 7.5 % (n=13) vs 5.2 ± 9.5 % (n=15) in WT and YAC128 CPNs respectively (Students' unpaired T-test; p=0.0418)]. We next used caffeine (1 mM) as an alternative means of agonizing ryanodine receptors; this more prominently increased the mEPSC frequency in WT cultures, but otherwise produced similar results **(Supplemental Figure 2)**.

Removing extracellular Ca^{2+} and blocking the ER SERCA pump substantially reduces the mEPSC frequency in YAC128 but not WT cultures

If an ongoing release of presynaptic ER Ca^{2+} in YAC128 cultures occludes potentiation of synaptic glutamate release by low dose ryanodine and caffeine, blocking ER Ca^{2+} release should reduce synaptic glutamate release in YAC128 cultures. To test this, we pre-incubated cultures with extracellular fluid (ECF) containing the sarco/endoplasmic reticulum Ca^{2+}-ATPase (SERCA) pump inhibitor cyclopiazonic acid (CPA) (30 μM) to deplete ER Ca^{2+} stores. Since ER Ca^{2+} depletion can increase cytosolic Ca^{2+} by engaging the store operated Ca^{2+} response, and this effect can increase mini neurotransmitter release (Emptage et al., 2001), we performed these experiments in the absence of extracellular Ca^{2+}. Under these conditions, YAC128 CPNs showed a mean mEPSC frequency of 3.33 ± 0.57 Hz (n=14), significantly lower than seen in WT CPNs under identical conditions 8.41 ± 1.50 Hz (n=17) (Student's unpaired t-test; p=0.0062) **(Figure 2I-K)**. These results suggest that an ER calcium leak into the cytoplasm elevates mini vesicular glutamate release from YAC128 CPN terminals, and that in the absence of this effect, the intrinsic vesicular release probability is actually reduced in YAC128 compared to WT cortical terminals.

Presynaptic Ca^{2+} sparks and waves are more frequent in YAC128 cultures

To directly monitor presynaptic Ca^{2+} dynamics, we next fused GCaMP6-M (T.-W. Chen et al., 2013) with the rat synaptophysin protein via a small glycine-serine linker to generate a genetically-encoded Ca^{2+} sensor that preferentially localizes to presynaptic terminals. We expressed this rat synaptophysin-tagged GCaMP6-M construct (rSyph-GCaMP6m) in neurons in cortical cultures

and performed Ca^{2+} imaging experiments, first in the presence of TTX (500 nM) to relate

presynaptic Ca^{2+} signaling to our mEPSC findings (above). Remarkably, both YAC128 and WT

cultures showed spontaneous axonal Ca^{2+} sparks, often beginning in single boutons, then initiating

slow Ca^{2+} waves spreading to neighboring boutons **(Figure 3A, B)**. In some cases, the above waves

slowly traversed axons encompassing entire 63X (178.6 μm x 113.1 μm) imaging fields over

multiple seconds of imaging. Conversely, axonal Ca^{2+} events in the absence of TTX could not be

temporally resolved (owing to relatively slow GCaMP kinetics) and appeared simultaneously

across all the boutons of a given imaged axon. These mini presynaptic Ca^{2+} events in TTX were

also strikingly long lasting at individual boutons - on average 5 - 10 times longer than that of

typical events seen in the absence of TTX **(Figure 3C, D)**. These miniature presynaptic Ca^{2+} events

in TTX were more than three times as frequent in YAC128 cultures than in WT cultures **(Figure**

3E), but event DF/F amplitudes did not significantly differ between genotypes **(Figure 3F)**. The

detection algorithm used here (see methods) considered Ca^{2+} waves involving multiple axonal

boutons as single events (as in **Figure 3A**). However, the mean event area was similar in both

genotypes (not illustrated), suggesting similar numbers of axonal boutons were recruited on

average by such events in YAC128 and WT cultures. In subsets of WT and YAC128 cultures we

also expressed an mCherry-tagged PSD95 construct. In these cultures, we repeatedly observed

clear colocalizations between rSyph-GCaMP6m-labeled boutons, spontaneously active in TTX,

and mCherry-labeled dendritic spines (not illustrated). Thus although we cannot be certain that all

boutons showing spontaneous Ca^{2+} events in TTX participate in synaptic connections, it is clear

that such events are not restricted to ectopic boutons.

Basal cytosolic Ca^{2+} is higher in YAC128 presynaptic boutons

In a subset of TTX experiments, we applied the Ca^{2+} ionophore ionomycin (10 μM) to cultures following GCaMP imaging. By equilibrating cytosolic Ca^{2+} to extracellular levels, this approach allowed quantification of GCaMP fluorescence in the presence of the known extracellular Ca^{2+} concentration (here 2 mM). Thus, smaller ionomycin-mediated increases in GCaMP fluorescence indicate relatively higher basal (pre-ionomycin) cytosolic Ca^{2+} concentrations (Lindhout et al., 2019). In both genotypes, ionomycin-mediated changes in presynaptic GCaMP fluorescence (DF/F) were significantly smaller in boutons that had shown at least one spontaneous Ca^{2+} event in the previous 3 min recording **(Figure 4A, D)**, indicating higher basal cytosolic Ca^{2+} concentrations in these spontaneously active boutons. Furthermore, overall ionomycin responses were significantly smaller in YAC128 cultures than in WT cultures when comparing the entire population of boutons imaged in a culture (both spontaneously active and inactive) **(Figure 4E)**.

Caffeine increases basal Ca^{2+} and miniature events in WT, but not YAC128 cortical boutons

The slow kinetics of the above miniature presynaptic Ca^{2+} events are consistent with ER-mediated Ca^{2+} waves reported in postsynaptic neuronal compartments (Ross, 2012). We next tested whether these presynaptic Ca^{2+} events were affected by caffeine (1 mM), an ER ryanodine receptor agonist that substantially increased mEPSC frequencies in WT, but not YAC128 cultures (above). In the presence of TTX, caffeine (1 mM) significantly increased the presynaptic Ca^{2+} event frequency in WT cultures **(Figure 5A)**, but did not significantly alter event frequency in YAC128 cultures **(Figure 5B)**.

Significant photobleaching of the rSyph-GCaMP6m construct occurred with prolonged imaging; this combined with the relative infrequency of mini presynaptic Ca^{2+} events necessitated imaging of separate fields for the above control and caffeine comparisons. We next performed within-bouton measurements of basal rSyph-GCaMP6m fluorescence before and immediately following application of caffeine; the relatively short duration of these experiments made before and after measurements feasible. A clear slow increase in basal rSyph-GCaMP6m fluorescence was seen in most WT boutons following caffeine (1 mM) application **(Figure 5C)**, which was generally absent or reduced in YAC128 boutons **(Figure 5D)**. Overall a significantly greater caffeine (1 mM)-mediated increase in rSyph-GCaMP6m fluorescence was seen in WT boutons compared to YAC128 **(Figure 5E)**.

Action potential-dependent Ca^{2+} signals in presynaptic cortical terminals Are less frequent in YAC128 cultures

We next used the rSyph-GCaMP6m construct in the absence of TTX to examine presynaptic action potential-dependent Ca^{2+} signals in WT and YAC128 cortical cultures. When neuronal action potential firing was intact, rSyph-GCaMP6m -expressing boutons in both WT and YAC128 cultures were dominated by presumably voltage-gated Ca^{2+} channel-mediated signals **(Figure 6A, B)**. These signals were more frequent than those seen in the presence of TTX, but of far shorter duration. Interestingly, these activity-dependent Ca^{2+} events were nearly twice as frequent in WT axonal boutons, compared to those in YAC128 cultures **(Figure 6A, C and E).** Ryanodine (5 μM)

modestly but significantly reduced the frequency of these events in WT cultures (by 17 %) **(Figure**

6A, B and E). Conversely, 5 μM ryanodine elicited a modest, but significant increase in the

frequency of these events in YAC128 cultures **(Figure 6C- E)**.

As discussed above, rSyph-GCaMP6m events in the presence of TTX were typically much longer

in duration than action potential-dependent events in both genotypes and appeared to be restricted

to a subset of boutons with higher resting Ca^{2+} concentrations. Interestingly, similar strikingly

long-lasting events continued to occur in subsets of boutons in the absence of TTX. In an attempt

to identify boutons exhibiting these slow Ca^{2+} events, we next categorized boutons imaged in the

absence of TTX based on whether one or more Ca^{2+} events with a duration greater than 6 s

(measured at half peak amplitude) occurred during a 3 min recording. 6 s was chosen as the cutoff

value, because it was more than 5 standard deviations greater than the mean event duration seen

in WT and YAC128 boutons in the absence of TTX, but comparable to average event durations in

TTX in both genotypes: [8.60 ± 0.92 s (n=164) and 6.23 ± 0.35 s (n=440) in WT and YAC128

boutons respectively]. Without TTX, 7.9 % (23/292) of WT boutons showed one or more events

lasting longer than 6 s, while under identical conditions 10.4 % (44/423) of YAC128 boutons

showed at least one such event. Interestingly in both genotypes, overall event frequencies were

significantly lower in boutons showing one or more slow events (>6 s) than in boutons showing

only short duration events **(Figure 6F- I)**. Assuming boutons displaying these long lasting Ca^{2+}

events correspond to the same population that is spontaneously active in TTX, these results suggest

action potential-dependent Ca^{2+} events are reduced in the bouton population with higher resting

cytosolic Ca^{2+} levels and spontaneous ER release events. Taken together, experiments up to this

point suggest that releasing presynaptic ER Ca^{2+} elevates miniature vesicular glutamate release,

but can also reduce activity-dependent presynaptic Ca^{2+} influx, and imply that the spontaneous or

pharmacological release of presynaptic ER Ca^{2+} oppositely regulates miniature and action

potential-dependent glutamate release.

Low dose ryanodine reduces evoked glutamate release in WT- but not YAC128-derived

brain slices

We next performed experiments in cortical-striatal *ex-vivo* brain slices prepared from 2 – 3 month

old WT and YAC128 mice expressing the fluorescent glutamate sensor iGluSnFR in striatal

neurons (Parsons et al. 2016; Koch et al. 2018). This preparation allowed direct optical

measurement of glutamate release in the striatum, independent of postsynaptic neuronal properties.

Striatal iGluSnFR signals evoked by stimulating cortical axons of the corpus callosum were

significantly decreased by ryanodine (5 µM) in WT, but not YAC128 slices **(Figure 7)**. These data

are consistent with results in culture (Figs. 6), suggesting that AP-dependent glutamate release is

reduced in YAC128 as a consequence of a tonic Ca^{2+} leak from ER stores

Discussion

Presynaptic neurotransmitter release and consequent postsynaptic signaling largely underlies

communication between neurons, thus forming the basis of brain circuitry. Neurotransmitter

release in turn, can be broadly divided into action potential-dependent or independent forms. The

former requires presynaptic voltage-gated Ca^{2+} channel activation coordinated by sodium action

potentials; while the latter persists in the absence of neuronal activity. Although miniature

neurotransmission is poorly understood compared to its activity-dependent counterpart, it is

increasingly accepted to subserve clear physiological functions (Frank, Kennedy, Goold, Marek,

& Davis, 2006; McKinney, Capogna, Dürr, Gähwiler, & Thompson, 1999; Sutton et al., 2006).

Moreover, miniature release can be regulated relatively independently of action potential-

dependent release and may even be mediated by distinct vesicular pools (Fredj & Burrone, 2009;

Sara, Virmani, Deák, Liu, & Kavalali, 2005).

Alterations in synaptic signaling processes, particularly relating to the excitatory transmitter

glutamate, have been repeatedly reported in models of HD (Raymond et al., 2011; Sepers et al.,

2017; Tyebji & Hannan, 2017) and other neurodegenerative diseases (R. Wang & Reddy, 2017).

Increased cell-surface expression of extra-synaptic NMDA receptors by SPNs favors excitotoxic

postsynaptic glutamate-mediated signaling in HD models (Milnerwood et al., 2010). Mounting

evidence also suggests glutamate release from cortical afferents is altered in HD; however, the

direction of this alteration appears to be disease-stage-dependent (Cepeda et al., 2003; Joshi et al.,

2009).

The endoplasmic reticulum (ER), a continuous intracellular membrane-system involved in Ca^{2+}

storage and protein synthesis, is expressed in all neuronal processes including axons and

presynaptic boutons. Increased cytosolic Ca^{2+}-release from the ER has been shown in HD models,

as a result of a mutant huntingtin protein (mHTT)-mediated increase in IP3 receptor

responsiveness (Tang et al., 2003) and a constituent ryanodine receptor Ca^{2+} leak (Suzuki et al.,

2012). Previous HD studies have focused on the postsynaptic ER, however Ca^{2+} release from the

presynaptic ER is known to modulate neurotransmitter release (Emptage et al., 2001; Llano et al.,

2000). Results of our study suggest mHTT expression alters ER Ca^{2+} handling at stores in close

proximity to cortical presynaptic terminals, thereby altering glutamate release. This mechanism

clearly increased the frequency of Ca^{2+}-dependent, action potential-independent (mini) synaptic

glutamate events in our YAC128 model. Conversely, our findings suggest reduced action

potential-dependent glutamate release from YAC128 cortical terminals. As we have focused

exclusively on glutamatergic synapses, future experiments will be required to determine if this

mechanism similarly affects the release of other neurotransmitters and neuromodulators.

Spontaneous axonal Ca^{2+} signaling increases the YAC128 miniature glutamate release

frequency

CPNs in YAC128 cortical cultures at DIV14 and DIV18 showed higher mEPSCs frequencies

compared to CPNs in age-matched WT cultures. However by DIV 21, mEPSC frequencies were

similar between WT and YAC128 CPNs. This pattern is reminiscent of SPN mEPSC frequencies

in YAC128 cortical-striatal co-cultures, which were also higher relative to WT at early DIV ages,

but comparable to WT by DIV21 (Buren et al., 2016). In the co-culture model, the total dendritic

length of YAC128 SPNs was reduced relative to WT at DIV21, indicating reduced total SPN

synapse numbers may have masked ongoing elevated glutamate release rates from individual

YAC128 cortical terminals. We suspect that in our cortical cultures, degenerative changes at

DIV21 likewise obscured the elevated YAC128 CPN mEPSC frequencies clearly seen at earlier

DIV ages. Notably in DIV18-aged cortical cultures, the time point at which we performed

mechanistic experiments, no genotype differences in CPN synapse density and dendritic

morphology were evident, indicating that differences in mEPSC frequencies likely reflected

altered rates of release at individual presynaptic sites. In any case, increased rates of miniature

glutamate release appear to be an early and perhaps enduring phenotype of cultured YAC128

CPNs.

Caffeine or low dose ryanodine, which release ER Ca^{2+} by opening ryanodine receptors, failed to

increase the mEPSC frequency in YAC128 CPNs, despite robustly increasing the WT mEPSC

frequency. This result, in concert with the increased basal YAC128 mEPSC frequency, suggests

an ongoing spontaneous Ca^{2+} store release in YAC128 cultures occludes facilitation of the mEPSC

frequency by drug-mediated Ca^{2+} store release. The substantially lower CPN mEPSC frequency

seen in YAC128 cultures incubated with the ER SERCA pump inhibitor CPA in nominally zero

extracellular Ca^{2+}, is also consistent with a strong ER-derived Ca^{2+}-dependence of YAC128 mini

glutamate release. Conversely, mEPSC frequencies in WT cultures were not substantially reduced

under identical conditions; this resulted in significantly higher WT than YAC128 mEPSC

frequencies in CPA and 0 mM extracellular Ca^{2+}, notably opposite the relationship seen without

CPA in standard (2 mM) extracellular Ca^{2+}. This was unexpected given overall WT and YAC128

CPN synapse numbers were similar in these cultures, but may indicate a reduction in functional

YAC128 synapses, not apparent in fixed culture images, or deficits in the YAC128 synaptic release

machinery, as suggested in previous studies (Morton & Edwardson, 2001; Morton, Faull, &

Edwardson, 2001). In either case, enhanced Ca^{2+}-dependent mini glutamate release in YAC128

cultures clearly surmounts any such deficits. These results also suggest mini glutamate release in

WT cultures is largely Ca^{2+}-independent. In contrast, Xu et. al (2009) reported mini glutamate

release in similar cortical cultures persisted in the absence of extracellular Ca^{2+}, but was nearly

completely dependent on internal Ca^{2+}stores and blocked by preincubation with the membrane-

permeant Ca^{2+} chelator BAPTA-AM (Xu, Pang, Shin, & Südhof, 2009). If this were true in our

cultures, we would expect Ca^{2+} store depletion (with CPA), in concert with removal of extracellular

Ca^{2+}, to similarly abolish mini release. However CPA is reported to only partially deplete

presynaptic ER Ca^{2+} (de Juan-Sanz et al., 2017), therefore a role of residual ER Ca^{2+} in maintaining

mini release in WT cultures under these conditions cannot be excluded.

Despite a clear role of ER Ca^{2+} in mini release in a variety of neuronal preparations, the temporal

and spatial dynamics of action potential-independent presynaptic ER Ca^{2+} signals are largely

unclear. Imaging the rSyph-GCaMP6m probe in TTX revealed spontaneous, presynaptic Ca^{2+}

events that often initiated Ca^{2+} waves traversing cortical axons. These events were more frequent

in YAC128 cultures and showed strikingly slow kinetics in both genotypes, consistent with ER

Ca^{2+} waves reported in dendrites and non-neuronal cell-types (Ross, 2012). Caffeine substantially

increased the mini axonal Ca^{2+} event frequency in WT cultures to levels comparable to that seen

in YAC128 cultures, but failed to significantly change the event frequency in YAC128 cultures.

The congruence of these results with our mEPSC findings (above) suggests these axonal Ca^{2+}

events underlie the increased YAC128 mini glutamate release frequencies.

Basal cytosolic Ca^{2+} concentrations were higher in axonal boutons showing spontaneous Ca^{2+}

activity in TTX, compared to inactive boutons, in both genotypes. Elevated cytosolic Ca^{2+} levels

may therefore be required to precipitate the initial ryanodine or IP3 receptor opening needed to

incite the regenerative ER Ca^{2+}-induced Ca^{2+} release we presume underlie these axonal Ca^{2+}

events. Interestingly, the restriction of spontaneous Ca^{2+} signals to a subset of cortical boutons

with higher basal Ca^{2+} concentrations may mean mini glutamate release is not uniform across

cortical boutons, but favored from this population of spontaneously active boutons. Indeed,

subpopulations of synapses favoring mini release have been described in other neuronal systems

(Atasoy et al., 2008; Peled, Newman, & Isacoff, 2014; Reese & Kavalali, 2016). We suspect the

higher basal cytosolic Ca^{2+} concentrations in the overall YAC128 axonal bouton population meant

a greater proportion were capable of sustaining spontaneous Ca^{2+} activity. Caffeine application

increased the baseline cytosolic Ca^{2+} concentration and frequency of spontaneous events in WT,

but not YAC128 boutons, consistent with this idea.

Action potential-dependent glutamate release is suppressed due to presynaptic ER depletion

Regardless of genotype, striking differences were seen in the rSyph-GCaMP6m signals that

predominated in the presence versus absence of TTX. Most rSyph-GCaMP6m-expressing boutons

in the absence of TTX showed numerous spontaneous events during standard 3 min imaging

sessions. However, only subsets of boutons were spontaneously active in TTX, with active boutons

characterized by higher resting cytosolic Ca^{2+} concentrations. Furthermore, although Ca^{2+} events

at individual boutons in TTX were less frequent, they lasted many times longer than corresponding

action potential-dependent events. Interestingly in the absence of TTX, subsets of boutons of both

genotypes showed longer-lasting Ca^{2+} events, like those that persisted in TTX, typically alongside

faster action potential-dependent events. We suspect these boutons correspond to the population

spontaneously active in TTX. Event frequencies in such boutons, in the absence of TTX, were

significantly lower than in boutons lacking these longer-lasting events, further supporting the idea

that subsets of cortical boutons favor either miniature or action potential-dependent glutamate

release.

Although spontaneous axonal Ca^{2+} events were more common in YAC128 cultures in the presence

of TTX, the opposite was true when experiments were performed without TTX, suggesting

reduced activity-dependent glutamate release in YAC128 cultures. Reduced action potential-

dependent axonal Ca^{2+} event frequencies in YAC128 cultures could reflect decreased CPN firing

rates, altered transduction of sodium action potentials to bouton Ca^{2+} signals or some combination

of these factors. 5 μM ryanodine significantly reduced the frequency of these action potential-

dependent bouton Ca^{2+} events in WT cultures, though not to YAC128 levels, consistent with

reduced action potential-dependent presynaptic Ca^{2+} signaling in YAC128 cultures being partly,

but not entirely mediated, by spontaneous release of ER Ca^{2+}. However, nuances of ryanodine's

pharmacology may have confounded this conclusion. Low dose ryanodine opens ryanodine

receptors, but to a sub-maximal conductance state. Furthermore, ryanodine binding to a second

low affinity receptor site mediates channel closure, an action that predominates at doses greater

than 10 μM. Caffeine more effectively releases ER Ca^{2+}, exemplified here in patch clamp mini

EPSC experiments, and may therefore have been capable of lowering action potential-dependent

event frequencies in WT to YAC128 levels. Unfortunately, caffeine was undesirable for these

experiments because, unlike ryanodine, it is an antagonist at adenosine receptors, which are often

expressed at presynaptic sites and inhibit VGCCs (Dunwiddie & Masino, 2001). We speculate that

the ryanodine receptor population in YAC128 boutons favors the open conformation under basal

conditions, consistent with the lack of caffeine responses in YAC128 boutons, and that under such

conditions, ryanodine-mediated channel closure would be exaggerated. If so, this could account

for the modest, but significant ryanodine-mediated increase in event frequencies in YAC128

boutons.

Precise mechanism(s) by which pharmacological or disease-mediated ER Ca^{2+} release might

reduce activity-dependent presynaptic Ca^{2+} signaling remain uncertain. Indeed axonal or cell body

ER stores could equally underlie this effect given the dependence of these signals on sodium action

potentials. Dendritic ryanodine receptor activation has been shown to reduce neuronal firing rates

via activation of a Ca^{2+}-dependent K^+ conductance (van de Vrede, Fossier, Baux, Joels, &

Chameau, 2007). Alternatively, elevated cytosolic Ca^{2+} levels in YAC128 boutons (mediated by

the release of axonal stores) might contribute to Ca^{2+}-dependent VGCC inactivation, thereby

reducing the coupling of cortical action potentials to presynaptic VGCC-mediated Ca^{2+} influx.

Indeed, more generalized increases in cytosolic Ca^{2+} mediated by ER store release, have been

shown to facilitate Ca^{2+}-dependent inactivation of voltage-gated Ca^{2+} channels including the N

and P/Q-types commonly expressed in presynaptic terminals and implicated in vesicular release

(Budde, Meuth, & Pape, 2002; Cens, Rousset, Leyris, Fesquet, & Charnet, 2006). If this were the

case in our system, the higher resting Ca^{2+} concentrations evident in YAC128 boutons would be

expected to maintain local voltage-gated Ca^{2+} channels in a higher resting state of inactivation.

Low-dose ryanodine decreased the amplitude of striatum, glutamate-mediated iGluSnFR signals

in WT brain slices, an effect absent or diminished in YAC128 slices, consistent with our findings

in culture. There is a lack of consensus as to whether single action potentials can trigger Ca^{2+}-

induced Ca^{2+} release from the presynaptic ER (Emptage et al., 2001), or whether prolonged,

repetitive action potential firing is required (de Juan-Sanz et al., 2017). It seems unlikely that the

stimulation protocol used here appreciably contributed to the action potential-dependent discharge

of the presynaptic ER, given the relatively low intensity stimulation used (repeated only twice at

100 Hz). Rather, we suspect a ryanodine-mediated increase in presynaptic Ca^{2+} decreased evoked

glutamate release by inactivating presynaptic voltage-gated Ca^{2+} channels in WT slices, and that

this process was occluded in YAC128 slices. However, future studies will be required to clarify

mechanistic details of this process.

Taken together, the evidence presented here strongly suggests enhanced spontaneous release of

presynaptic ER Ca^{2+} in YAC128 mice favors miniature glutamate release at the expense of evoked

release.

Implications for postsynaptic signaling

Mini glutamate release elicits postsynaptic NMDA receptor-mediated Ca^{2+} influx under

physiologically relevant conditions (Beaulieu-Laroche & Harnett, 2018; Espinosa & Kavalali,

2009) and can mediate postsynaptic signaling distinct from that of action potential-dependent

release (Sutton, Taylor, Ito, Pham, & Schuman, 2007). Indeed mini glutamate release may activate

distinct populations of NMDAR receptors (Atasoy et al., 2008). In cultures, it was reported that

miniature glutamate-mediated events become toxic to CPNs following prolonged silencing of

neuronal activity with TTX (Fishbein & Segal, 2007). Future studies will be necessary to

determine if and how the increased mini glutamate release shown here interacts with the well

described alterations in postsynaptic NMDA receptor expression in the YAC128 and other HD-

models, and whether a shift towards activity-independent glutamate release contributes to

neurodegeneration in HD.

Materials and Methods

Culture preparation:

All animal-related procedures were approved by and adhered to the guidelines of the University

of British Columbia Committee on Animal Care and the Canadian Council on Animal Care

(protocols A17-0295, A15-0069 and A19-0076). Cultures were prepared from both male and

female embryonic day 17-18 pups from either wild-type (WT) FVB/N or transgenic yeast artificial

chromosome-containing mice expressing the full-length human huntingtin genomic DNA with 128

CAG repeats (YAC128). YAC128 mice were maintained on the FVB/N background (homozygous

line 55). Wildtype and YAC128 mice used for ex vivo slice experiments (below) and bred for

culture preparation (above) were group housed under controlled conditions, free of know

pathogens, at room temperature (22 – 24 °C), under a 12 hr light/dark cycle. Cortical cultures used

in patch clamp electrophysiology and Ca^{2+}-imaging experiments were prepared as previously

described (Milnerwood et al., 2012; Smith-Dijak et al., 2019) and plated at a density of 225, 000

neurons/ml. In a subsets of experiments, a portion of the total 2.7 million cortical neurons (plated

per 24-well culture) were transfected with transgenic reporters including: GFP; a synaptophysin-

tagged GCaMP6-M construct (a generous gift from Dr. Anne Marie Craig, UBC); a postsynaptic

density 95 (PSD-95)–tagged M-cherry construct; or a GFP-tagged internally expressed anti-

PSD95 antibody (a generous gift from Dr. D.B. Arnold, University of Southern California; (Gross

et al., 2013)).

Electrophysiology:

An Axopatch 200B amplifier and pClamp 9.2 software (Molecular Devices, Sunnyvale, CA) were used to acquire whole-cell patch clamp electrophysiology recordings. Data was digitized at 20 kHz and low-pass filtered at 1 kHz. For electrophysiology experiments, cultures were perfused with extracellular fluid (ECF) containing (in mM): 167 NaCl, 2.4 KCl, 10 glucose, 10 HEPES, 2 $CaCl_2$ and 1 $MgCl_2$; NaOH (1 mM) was used to adjust the pH to 7.30, and the osmolarity was adjusted to 305 – 310 mOsm. Tetrodotoxin (TTX) (500 nM) and picrotoxin (PTX) (50 mM) were added to this ECF to block sodium channel-mediated action potentials and $GABA_A$ receptor-mediated currents respectively. Neurons were patched with borosilicate glass pipettes pulled to a tip resistance of 3-6 $M\Omega$ when back-filled with intracellular solution containing in (mM): 130 Cs-methanesulfonate, 5 CsCl, 4 NaCl, 1 MgCl, 10 HEPES, 5 EGTA, 5 QX-314 Cl, 0.5 Na-GTP, 10 Na-phosphocreatine, and 5 Mg-ATP (~286 mOsm). During experiments, neurons were held at – 70 mV in voltage-clamp, with hyperpolarizing voltage steps (-10 mV) performed periodically to measure intrinsic membrane properties. Under these conditions AMPA receptor-mediated miniature excitatory postsynaptic currents (mEPSC)s appeared as transient inward current deflections. Recordings with a series resistance greater than 25 $M\Omega$ were excluded from analysis; typical values were between 15 – 20 $M\Omega$. A minimum of 2 minutes following establishment of the whole cell configuration was allowed before experimental measurements, so neurons could fully dialyze with intracellular solution and achieve a stable membrane resistance and holding current. For experiments involving within-cell drug applications, a maximum 20 % change in series resistance between control and drug measurements was tolerated. In these cases, drugs were

applied locally to the neuron with a fast perfusion system. Mini Analysis software (Synaptosoft)

was used to detect mEPSCs and extract relevant parameters. A minimum of 100 and no more than

1000 mEPSC's were analyzed per neuron per experimental condition.

Cortical pyramidal neuron morphology:

Cortical pyramidal neurons (CPN)s in WT and YAC128 cultures were labeled by transfecting a

subset of neurons (1 of 2.7 million) with a cytoplasmic green fluorescent protein (GFP)

(Addgene plasmid 37825) at the time of platting. Cultures were subsequently fixed at DIV17 - 19

and GFP-labeled CPN dendritic arbors imaged on a Zeiss Axiovert 200 M fluorescence

microscope (20 x magnification, 0.8 NA), with a Zeiss 702 monochrome camera, using Zen

software. Multiple image Z-stacks were acquired and X,Y-tiling was used to ensure entire dendritic

arbors were visualized. Images were exported to Fiji-ImageJ for analysis by a blinded observer.

Images were flattened using the maximum Z-projection function. Background subtraction was

performed, and neuronal processes were thresholded following adjustment of brightness and

contrast. Automated Sholl analysis was performed using the ImageJ sholl analysis plugin.

Excitatory cortical synapse staining:

A subset of neurons (2 of 2.7 million) in WT and YAC128 cortical cultures were transfected with

a GFP-tagged internally-expressed anti-PSD95 antibody (intrabody) (Gross et al., 2013) at the time

of platting. At DIV17 – 19, cells were fixed and stained for VGlut1 and the GluA2 AMPA receptor

subunit as previously described (Buren et al., 2016). Briefly, cultures were first live-stained with

a primary mouse anti-GluA2 antibody (Millipore), then fixed and stained with a secondary Alexa

Fluor 568-conjugated donkey anti-mouse antibody (Invitrogen). Subsequently, cultures were

incubated with a primary guinea pig anti-VGlut1 antibody (Millipore), then stained with a

secondary AMCA-conjugated donkey anti-guinea pig antibody (Jackson Immuno Research

Laboratories). To amplify the GFP fluorescence of the anti-PSD95 intrabody, cultures were also

incubated with a primary chicken anti-GFP antibody (1:1000) (Millipore), followed by an

secondary Alexa Fluor 488-conjugated antibody (1:500) (Invitrogen). Cultures were imaged on a

Zeiss Axiovert 200 M fluorescence microscope (63 x magnification, 1.4 NA), using a Zeiss 702

monochrome camera and Zen software. CPNs expressing the anti-PSD95 intrabody were

identified based on their diffuse cytoplasmic GFP fill, with bright GFP-labelled puncta expressed

at dendritic spines. A portion of each CPNs arbor, containing multiple secondary and tertiary

dendrites, was selected for imaging and sufficient image Z-stacks were acquired to adequately

capture all dendritic processes present in a given 63x field. Images were exported to Fiji-ImageJ

for analysis by a blinded observer and flattened using the maximum Z-projection function. For

each CPN image, the GFP channel was used to identify 3 secondary or tertiary dendritic segments,

at least 40 μm away from the CPN soma, over which ROIs were drawn. Following background

subtraction, fluorescent puncta in the green (GFP), red (Alexa Fluor 568) and blue (AMCA)

channels, visible within dendritic ROIs, were manually thresholded and detected with the analyze

particles function. The ImageJ colocalization plugin was used to identify triple-colocalized puncta

(PSD95, GluA2 and VGlut1), which we interpreted as functional CPN glutamatergic synapses.

Synapse density was defined as the number of triple-colocalized puncta present within a dendritic

segment divided by the area of the segment and averaged across all 3 dendrites analyzed in a given

CPN.

Synaptophysin-GCaMP imaging:

To directly image cytosolic Ca^{2+} in axonal boutons of neurons in our cortical mono cultures we

transfected 1 million cells (of a total 2.7 million) at time of plating with a rat synaptophysin-tagged

GCaMP6-M construct (rSyph-GCaMP6m). The rSyph-GCaMP6m construct was created by fusing

GCaMP6-M (T.-W. Chen et al., 2013) with the full length rat synaptophysin protein ($1 - 307$

amino acids) via a small glycine-serine linker and inserting the fused rSyph-GCaMP6-M construct

into a pLL3.7-hSyn vector to achieve neuron-selective expression. For some experiments, the same

1 million cells were also transfected at time of platting with an M-cherry-tagged PSD95 construct;

in these cases, rSyt-GCaMP6m-expressing presynaptic boutons that colocalized with M-Cherry

labeled postsynaptic spines were presumed to be functional synapses.

For all Ca^{2+}-imaging experiments, cultures were plated on 8-well cover-glass chambers (Thermo

Scientific ™, Nunc ™, Lab-Tek ™) and imaged at DIV (17 - 19) with a Zeiss Axiovert 200 M

fluorescence microscope (63 x magnification, 1.4 NA), using a Zeiss 702 monochrome camera

and Zen software. Movies were acquired at 10 Hz (100 ms exposure per frame) using the Zen

time-series mode with camera-steaming enabled. These experiments were performed in standard

ECF (as above) with or without TTX (500 nM) present, but in the absence of PTX.

Spontaneous Ca^{2+} waves were commonly observed in rSyph-GCaMP6m-labelled axons, evident

particularly when action potential-dependent Ca^{2+} events were blocked with TTX. Automatically

detecting such Ca^{2+} waves proved difficult with algorithms commonly used to quantify activity-

619 related neuronal GCaMP signals. We therefore used the Astrocyte Quantitative Analysis (AQuA)

620 software (running in MATLAB) (Y. Wang et al., 2019) to quantify these waves in the presence of

621 TTX. AQua does not rely on spatially segmenting a Ca^{2+} movie into ROIs based on neuronal

622 morphology, but rather defines events as spatially and temporally connected Ca^{2+} signals

623 surpassing user-defined thresholds. This means spontaneous axonal waves spreading across

624 multiple boutons are typically classified as a single event, because such Ca^{2+}-activity is grouped

625 in time and space. Conversely, a given bouton can be involved in multiple axonal waves and thus

626 detected as part of multiple events, as long as the pertinent signals are sufficiently temporally

627 separated. To compare the frequency of aQua-detected spontaneous axonal events between

628 genotypes and under pharmacological manipulations, we counted numbers of events occurring

629 within a standardized area [178.6 μm x 113.1 μm (a maximal 63 x field of view)], during a

630 standardized (3 min) consecutive imaging interval. The same aQua detection parameters,

631 empirically determined to best match a small number of manually analyzed experiments, were

632 applied across all cultures and conditions analyzed to facilitate meaningful comparisons.

634 In a subset of experiments conducted in the presence of TTX, responses of individual WT or

635 YAC128 rSyph-GCaMP6m-labelled synaptic boutons to caffeine or ionomycin were measured. In

636 these experiments culture fields were imaged for 3 min (as above), after which caffeine (1 mM) or

637 ionomycin (10 μM) was applied and imaging continued for an additional 1-2 min. aQua was not

638 used to analyze these experiments, as an ROI-based algorithm was desired. For caffeine

639 applications, an analyzer (blinded to culture genotype) used Fiji-ImageJ (NIH) to manually assign

640 elliptical ROIs to 10 boutons per movie and exported each ROI's fluorescence-intensity time-

641 course. Time-courses were subsequently imported to MATLAB, where the curve fitting tool was

used to model each time-course's photobleaching profile based on the initial 3 min recording; this

curve was then extrapolated to the entire time-course including caffeine treatment. The resultant

"bleaching curve" was subtracted from the raw rSyph-GCaMP6m fluorescence curve and this

time-course subsequently divided by the "bleaching curve" to yield a final curve reflecting the

caffeine response in DF/F units. Ionomycin experiments were analyzed similarly, except in this

case, the blinded analyzer selected 10 active-boutons (showing at least 1 clear Ca^{2+} event during

the initial 3 min recording) and 10 inactive-boutons (with no Ca^{2+} events present during the initial

3 min recording). Time-courses derived from inactive-boutons were exported to MATLAB, where

the ionomycin-mediated DF/F responses were calculated as above. In the case of active-boutons,

spontaneous events occurring during the first 3 min were detected with MATLAB's findpeaks

function and excised from the raw fluorescence time-courses before curve fitting, but otherwise

processed as above.

As expected, spontaneous events were far more frequent when rSyph-GCaMP6m -expressing

cultures where imaged in the absence of TTX. Most boutons showed many events during a 3 min

imagining session. These events, which were presumably driven by the action potential-dependent

opening of voltage-gated Ca^{2+} channels, were far shorter in duration than typical TTX-resistant

events. To assess for genotype differences in these signals, a blinded analyzer used Fiji-ImageJ to

randomly assign elliptical ROIs to 20 boutons per movie and exported each ROI's fluorescence-

intensity time-course to MATLAB. The high frequency of action potential-dependent events often

precluded extracting the photobleaching profile of a bouton's time-course, necessitating an

alternative means of calculating the DF/F values of spontaneous events. We therefore averaged

the grey-value fluorescence intensity across an entire-time course, subtracted this average

fluorescence pointwise from the time-course, then divided pointwise by the average fluorescence

to convert to DF/F units. Any nonstationary present in these DF/F time-series due to

photobleaching was removed with the MATLAB detrend function. The MATLAB findpeaks

function was subsequently used to detect events in these DF/F traces and extract relevant

parameters including peak event DF/F amplitudes and event half-amplitude widths. Bouton event

frequencies were calculated by dividing numbers of detected events in a DF/F trace by the imaging

interval. Bouton event frequencies and event parameters in the absence of TTX were compared

between WT and YAC128 cultures under baseline conditions and following application of

ryanodine (5 μM).

iGluSnFR imaging in acute brain slices:

Expression of the genetically-encoded intensity-based glutamate-sensing fluorescence reporter

(iGluSnFR) (Marvin et al. 2013) in WT and YAC128 mice was achieved with stereotaxic injection

of a viral construct as described previously (Parsons et al., 2016). Briefly, under isoflurane

anesthesia, 1 – 1.4 μl of the AAV1.hSyn.iGluSnFr.WPRE.SV40 construct (Penn Vector Core; Dr.

Loren Looger, Janelia Farm Research Campus of the Howard Hughes Medical Institute) was

directly injected into the dorsal striatum of 4 – 6 week old mice. Following surgery, mice were

closely monitored for a week to ensure adequate recovery.

After waiting 3 – 6 weeks, to ensure optimal iGluSnFR expression, acute brain slices from 2-3

month-old YAC128 mice and age-matched WT controls were prepared as described previously

(Parsons et al., 2016; Koch et al., 2018). Briefly, mice were decapitated following deep

isoflurane anesthesia and their brains rapidly removed and placed in an ice-cold slicing solution,

bubbled with carbogen (95% O_2, 5% CO_2) gas, containing (in mM): 125 NaCl, 2.5 KCl, 25

$NaHCO_3$, 1.25 NaH_2PO_4, 2.5 $MgCl_2$, 0.5 $CaCl_2$, and 10 glucose. 300 μm thick striatum-containing

sagittal brain slices were cut with a Leica VT1200S vibratome. Slices were subsequently incubated

for 30 min in warmed artificial cerebral spinal fluid ACSF containing 2 mM $CaCl_2$ and 1 mM

$MgCl_2$; ACSF constituents and concentrations were otherwise identical to the slicing solution

(above).

Slices were transferred to a submerged recording chamber for experiments and perfused with

carbogen-bubbled ACSF at a rate of 2 – 3 ml/min at room temperature. Cortical release of

glutamate into the striatum was evoked by delivering paired 0.1 ms electrical pulses at 100 Hz

with an A-M Systems isolated pulse stimulator (Model 2100) and tungsten monopolar stimulating

electrode (tip resistance - 0.1 MΩ). The electrode was placed into a corpus callosum segment

adjacent to the dorsal striatum at an approximate 50 – 100 μm depth. During and immediately prior

to the electrical stimulation, iGluSnFR fluorescence was excited with a 470 nm LED; slices were

not illuminated between experimental measurements to minimize phototoxicity and bleaching.

Stimulation and LED activation were triggered by Clampex software (Molecular Devices,

Sunnyvale, CA). iGluSnFR fluorescence was isolated with a 530 nm bandpass filter and imaged

with a CCD camera (1 M60, Pantera, Dalsa) and XCAP software (Epix, inc.) at 150 Hz with 8 x

8-pixel binning. Experimental measurements encompassing four stimulation trials and two blank

trial were performed at 3-minute intervals. The four stimulation trials were averaged and the blank

trials, in which slice-fluorescence was imaged without electrical stimulation, were averaged and

used to account for photobleaching and to calculate the stimulation mediated changes in iGluSnFR

fluorescence over basal fluorescence (ΔF/F) as described previously (Parsons et al., 2016).

Videos were analyzed offline with ImageJ software. The average iGluSnFR signal was measured

over 10 x 10-pixel (93.8 x 93.8 μM) region of interested (ROI) placed over the maximal area of

evoked iGSnFR activity within the striatum adjacent to the stimulating electrode.

Experimental design and statistical analysis:

Statistical analysis and creation of figures was performed using GraphPad Prism (version 7). All

data distributions were tested for normality with the D'Agostino-Pearson omibus normality test.

The Student's unpaired t-test was used for unpaired comparisons between two data groups, such

as when mean mEPSC frequencies were compared between WT and YAC128 neurons, as long

both data groups were normally distributed. When one or both groups failed the D'Agostino-

Pearson omibus normality test, the non-parametric Mann Whitney test was used instead.

When parameters of the same group of neurons or axonal boutons were compared before and after

a drug treatment, statistical significance was assessed with the Student's paired t-test, unless data

points in the control or drug-treatment group failed the D'Agostino-Pearson omibus normality test;

in which case, the non-parametric Wilcoxon matched-pairs signed rank test was used instead.

A two-way ANOVA with the Bonferroni post-test was used when testing for genotype differences

in a dependent variable measured at different time points, as was the case for our brain-slice

iGluSnFR experiments.

Depending on the experimental design, "n" numbers in figures refer either to number of neurons, numbers of imaged culture fields, numbers of individual axonal boutons, or numbers of brain slices. Clarifying details are present within individual figure legends, as are the numbers of culture batches or mice used.

Differences in mean values were considered significant at $p<0.05$, and significance levels are indicated in figures as follows: $*p<0.05$, $**p<0.01$, $***p<0.001$, $***p<0.0001$. Comprehensive descriptions of statistical analysis are included in figure legends.

Acknowledgments

We thank Dr. Anne Marie Craig (UBC) for expert advice on experiments using rSyph-GCaMP6m, and Dr. Lily Zhang and Dr. Rujun Kang for technical support and assistance. The work was supported by funding from the Canadian Institutes of Health Research (CIHR) Foundation grants FDN-143210 to LAR and FDN-154278 to MRH. CB was supported by a University of British Columbia 4-year Graduate Fellowship (UBC 4-YF). ASD was supported by a CIHR Canada Graduate Scholarship Doctoral award and a UBC 4-YF. EK was supported by a CIHR Canada Graduate Scholarship Master's award and UBC 4-YF. MES was supported by a Vanier Canada Graduate Scholarship and a UBC 4-YF. WN was supported by a UBC-CIHR-MD/PhD studentship and a Vanier Canada Graduate Scholarship. LAR holds the UBC Department of Psychiatry Louise A. Brown Chair in Neuroscience. MRH holds a Canada Research Chair.

Declaration of Interests

760   The authors declare no competing interests.

 Figure legends:

Fig 1: YAC128 cortical cultures show elevated miniature EPSC frequencies at early DIV

time points.

Representative traces (**A, D, G, J**) and population data for mEPSCs (frequency in **B, E, H, K**;

amplitude in **C, F, I, L**) recorded from WT and YAC128 CPNs. All recordings were made in

voltage clamp with a holding potential of -70mV in solution containing TTX (500 nM) and PTX

(50 μM).

A – C: Recordings made from CPNs in DIV-7 cultures**.** Mean mEPSC frequency (**B**) was 1.8 ±

0.3 Hz (n=18; 3 cultures) in WT CPNs and 2.8 ± 0.7 Hz (n=17; 4 cultures) in YAC CPNs; this

difference was not statistically significantly [t(33)=1.449; p=0.1567; Student's unpaired-t test].

Mean mEPSC amplitudes (**C**) were similar in WT and YAC128 cultures: 17.2 ± 1.5 pA (n=18; 3

cultures) and 17.0 ± 1.7 pA (n=17; 4 cultures) respectively [t(33)=0.06936; p=0.9452; Student's

unpaired-t test].

D - F: Recordings were made from CPNs in DIV-14 cortical cultures. Mean mEPSC frequency

(**E**) was significantly higher in YAC128 CPNs [10.8 ± 1.8 Hz (n=20; 6 cultures)] compared to WT

CPNs [4.4 ± 2.6 Hz (n=22; 4 cultures)], [t(40)=3.555; p=0.0010; Student's unpaired-t test]. Mean

CPN mEPSC amplitudes (**F**) were not significantly different in WT and YAC128 cultures: 12.1 ±

0.8 pA (n=22; 4 cultures) and 15.0 ± 1.4 pA (n=20; 6 cultures), respectively [t(40)=1.807;

p=0.0786; Student's unpaired-t test].

G – I: Recordings were made from CPNs in DIV-18 cortical cultures. Mean CPN mEPSC

frequency (**H**) was significantly higher in YAC128 cortical cultures [14.1 ± 1.2 Hz (n=38; 10

cultures)] compared to WT cultures [9.9 ± 1.5 Hz (n=30; 15 cultures)], [p<0.0059 (exact); Mann

Whitney test]. Mean CPN mEPSC amplitudes (**I**) were similar in WT and YAC128 cultures: 17.0

± 1.8 pA (n=30; 15 cultures) and 17.4 ± 1.8 pA (n=38; 10 cultures) respectively [p=0.8643 (exact);

Mann Whitney test].

J – L: Recordings were made from CPNs in DIV-21 cortical cultures. Mean CPN mEPSC

frequency (**K**) was similar in WT and YAC128 cultures: 14.2 ± 1.2 Hz (n=62; 17 cultures) and

14.0 ± 0.9 Hz (n=71, 23 cultures), respectively [p=0.5626 (exact); Mann Whitney test]. Mean CPN

mEPSC amplitudes (**L**) were similar in WT and YAC128 cultures: 22.3 ± 1.3 pA (n=62; 17

cultures) and 24.3 ± 1.2 pA (n=71; 23 cultures), respectively [p=0.2143 (exact); Mann Whitney

test].

Supplemental Fig 1: Dendritic complexity and excitatory synapse numbers are similar in

WT and YAC128 cortical cultures.

A, B. Representative images generated by thresholding a merged Z-stack of a green fluorescent

protein (GFP)-expressing WT CPN (**A**) and YAC128 CPN (**B**).

C. Number of dendritic intersections through concentric sholl circles centered on the soma of GFP-

filled WT and YAC128 CPNs of radii between 10 μm and 450 μm. WT and YAC128 CPNs have

nearly identical dendritic branching distributions [2-way ANOVA; Genotype: $F_{(1, 6165)}= 1.168$,

p<0.2799; Distance from soma: $F_{(44, 6165)}=381.6$, p<0.0001; Interaction: $F_{(44, 6165)}=0.9868$,

p<0.4966].

D. Total numbers of sholl intersections, which reflect general neuronal complexity, were similar

in WT and YAC128 CPNs: 254.0 ± 8.8 (n=61; 4 cultures) and 249.5 ± 8.7 (n=78; 4 cultures)

respectively [p=0.7263 (exact); Mann Whitney test].

E. A representative merged Z-stack from a WT CPN (left panels) and YAC128 CPN (right panels)

with merged green, red and blue channels showing staining of a genetically-encoded internally-

expressed GFP-tagged anti-PSD-95 antibody (green), anti AMPA receptor GluA2 subunit

immuno-staining (red) and anti VGlut1 immuno-staining (blue) respectively. The bottom panel

shows an expanded view of a segment of dendrite with arrows pointing to a subset of co-localized

PSD-95, GluA2 and VGlut1 puncta, which were counted as presumed functional synapses. Note,

for illustrative purposes the brightness and contrast of the individual channels were adjusted to

best illustrate the punctate fluorescence.

F. Numbers of presumed functional synapses (identified as above) per 100 μm^2 of secondary and

tertiary dendrites of WT and YAC128 CPNs: 8.8 ± 1.1 (n=37; 2 cultures) and 6.8 ± 0.7 (n=57; 3

cultures), respectively. Although synapse numbers were lower in YAC128 cultures this did not

reach statistical significance [p=0.1039 (exact); Mann Whitney test].

Fig 2: Spontaneous release of ER Ca^{2+} elevates YAC128 pyramidal neuron miniature EPSC

frequencies.

A and B - Hypothetical model (tested below) A. We hypothesize, in WT cultures, spontaneous

Ca^{2+} release from ER stores in presynaptic cortical terminals only minimally augments action

potential-independent (mini) glutamate release. **B.** Conversely, in YAC128 cultures, we

hypothesize spontaneous Ca^{2+} release from presynaptic ER stores is increased, and that this

elevates Ca^{2+}-dependent, mini glutamate release.

C – H: Voltage clamp recordings were made at -70 mV from DIV-18 cultured WT (**C – E**) and

YAC128 CPNs (**F – H**) in the presence of TTX (500 nM) and PTX (50 μM), under control

conditions and during subsequent local ryanodine (5 μM) application. Note that in the

representative traces from the WT CPN (**C**), ryanodine (5 μM) substantially increased the mEPSC

frequency, whereas the drug had little effect on the mEPSC frequency in the representative

YAC128 CPN (**F**). Quantifying the population data revealed that ryanodine (5 μM) significantly

increased the mean mEPSC frequency in WT cultured CPNs (**D**) from 6.7 ± 1.2 Hz to 11.9 ± 2.4

Hz, [t(12)=2.50; p=0.0280; n=13; 8 cultures; Student's paired-t test]. Ryanodine (5 μM) did not

significantly affect the mean WT CPN mEPSC amplitude (**E**) (control: 21.0 ± 3.2 pA, ryanodine:

20.2 ± 2.3 pA), [t(12)=0.65; p=0.5278; n=13; 8 cultures; Student's paired-t test]. In YAC128

cultures, ryanodine (5 μM) did not significantly affect the mean CPN mEPSC frequency (**G**)

(control: 12.0 ± 1.7 Hz, ryanodine: 13.9 ± 2.0 Hz), [t(14)=1.65; p=0.1206; n=15; 4 cultures;

Student's paired-t test]. Ryanodine (5 μM) also did not significantly affect the mean CPN mEPSC

amplitude (**H**) (control: 20.5 ± 4.4 pA, ryanodine: 19.6 ± 3.4 pA), [t(14)=0.71; p=0.4904; n=15; 4

cultures; Student's paired-t test].

I. Voltage clamp recordings from representative WT (top) and YAC128 (bottom) CPNs at -70 mV

in the presence of TTX (500 nM) and PTX (50 μM) (as before), with the addition of CPA (30 μM)

and in the absence of extracellular Ca^{2+}. Note in all such recording, CPNs were incubated in this

CPA-containing, 0 mM Ca^{2+} ECSF for a minimum of 10 min.

J. In CPA (30 μM) and in the absence of extracellular Ca^{2+}, mEPSCs were significantly less

frequent in YAC128 CPNs [3.3 ± 0.6 Hz (n=14; 4 cultures)] compared WT CPNs [8.4 ± 1.5 Hz

(n=17; 5 cultures)] under like conditions [t(29)=2.952; p=0.0062; Student's unpaired-t test].

K. In CPA (30 μM) and in the absence of extracellular Ca^{2+}, YAC128 CPNs showed a trend

towards a lower mean mEPSC amplitude than in WT (WT: 26.7 ± 5.2 pA, YAC128: 16.1 ± 3.3

pA), but this did not reach statistical significance [t(27)=1.706; p=0.0995; Student's unpaired-t

test].

Supplemental Fig 2: Releasing ER Ca^{2+} with caffeine increases the CPN mEPSC frequency

in WT, but not YAC128, CPNs.

Voltage clamp recordings were made from cultured WT and YAC128 CPNs at -70 mV in the

presence of TTX (500 nM) and PTX (50 μM), under control conditions and during subsequent

local caffeine (1 mM) application.

A, D: Representative traces from WT (A) and YAC128 (D) CPNs, under control (top traces) and

caffeine (bottom traces) conditions. In this CPN, caffeine (1 mM) substantially increased the

mEPSC frequency.

B. Caffeine (1 mM) significantly increased the mean mEPSC frequency in WT cultured CPNs

from 12.3 ± 2.5 Hz to 22.8 ± 8.0 Hz, [p=0.0076 (exact); n=16; 8 cultures; Wilcoxon matched-pairs

signed rank test].

C. Caffeine (1 mM) did not significantly affect the mean WT CPN mEPSC amplitude (control:

13.7 ± 1.9 pA, caffeine: 17.2 ± 3.4 pA), [p=0.8603 (exact); n=16; 8 cultures; Wilcoxon matched-

pairs signed rank test].

E. In YAC128 CPNs, caffeine (1 mM) did not significantly affect the mean CPN mEPSC

frequency (control: 12.0 ± 2.1 Hz, caffeine: 11.5 ± 1.9 Hz), [t(12)=0.4808; p=0.6393; n=13; 5

cultures; Student's paired-t test].

F. In YAC128 CPNs, caffeine (1 mM) also did not significantly affect the mean CPN mEPSC

amplitude (control: 14.5 ± 1.4 pA, caffeine: 14.1 ± 1.2 pA), [t(12)=0.8086; p=0.4345; n=13; 5

cultures; Student's paired-t test].

G. Percent change in mEPSC frequency mediated by Caffeine (1 mM) application to WT and YAC128 CPNs (calculated from the responses in panels **B** and **E**). Caffeine (1 mM) elicited a significantly greater increase in the mEPSC frequency in WT, compared to YAC128, CPNs [WT: 99.2 ± 34.1 % (n=16; 8 cultures), YAC128: 1.2 ± 7.6 % (n=13; 5 cultures)], [p=0.0025 (exact); Mann Whitney test].

Fig 3: Axonal Ca^{2+} waves are more frequent in YAC128 Cortical Cultures.

A. Portion of a YAC128 rat synaptophysin-tagged GCaMP6-M (rSyph-GCaMP6m)-expressing axon during a spontaneous Ca^{2+} wave in the presence of ECF containing TTX (500 nM). Images reflect raw GCaMP fluorescence values (arbitrary units) and comprise 15 s of a 4 min video. For illustrative purposes, images were binned temporally from the original 10 Hz movie, such that each 2 s frame reflects the average fluorescence of 20 successively acquired frames.

B. Time course of DF/F (change in fluorescence over basal fluorescence) values extracted from 3 of the YAC128 rSyph-GCaMP6m-expressing axonal boutons (shown in panel **A** with color-coded arrows, panel labelled 15s). Signals were spatially averaged across elliptical regions of interest (ROI)s encompassing the indicated boutons and represent the entire 4 min imaging session. Note the multi-second delay in the propagation of signals between boutons and the long time course of individual events.

C. Representative 3 min DF/F time course from a WT rSyph-GCaMP6m-expressing axonal bouton in TTX (500 nM) (top). Note the substantially longer duration of this event compared to representative events recorded without TTX from a separate WT rSyph-GCaMP6m-expressing bouton (bottom).

D. Average half amplitude width of spontaneous Ca^{2+} events occurring in rSyph-GCaMP6m-expressing boutons in WT and YAC128 cortical cultures in the presence and absence of TTX (500 nM). Select comparisons were made with the Mann Whitney test because event widths in all groups failed the D'Agostino-Pearson omibus normality test. WT events were significantly longer with TTX [8.60 ± 0.92 s (n=164 events; 85 boutons; 4 cultures)] than without [1.06 ± 0.97 s (n=3725 events; 175 boutons; 4 cultures)] [p<0.0001 (approximate); Mann Whitney test]. Likewise, YAC128 half widths were significantly longer with TTX [6.23 ± 0.35 s (n=440 events; 153 boutons; 4 cultures)] than without [1.17 ± 1.46 s (n=5159 events; 422 boutons; 6 cultures)] [p<0.0001 (approximate); Mann Whitney test].

E. Numbers of detected spontaneous axonal events (as in Panels **A** and **B**) during 3 min recordings of 63 x objective (178.6 μm x 113.1 μm) fields imaged in the presence of TTX (500 nM). Action potential-independent events were significantly more frequent in YAC128 cortical cultures [115.9 ± 28.1 events/3 min (n=15 fields; 4 cultures)], compared to WT [36.3 ± 5.5 events/3 min (n=12 fields; 4 cultures)], [p=0.0016 (exact); Mann Whitney test].

F. Average peak DF/F amplitudes of all the spontaneous, action potential-independent axonal Ca^{2+} events imaged in a given 63 x objective field over 3 min from WT and YAC128 cortical cultures. Average event amplitudes were not significantly different between WT [0.240 ± 0.046 DF/F (n=10 fields; 4 cultures)] and YAC128 cultures [0.216 ± 0.027 DF/F (n=12 fields; 4 cultures)], [t(20)=0.465; p=0.647; Student's unpaired-t test].

Fig 4: Boutons spontaneously active in TTX show relatively elevated cytosolic Ca^{2+} in both genotypes. Resting cytosolic Ca^{2+} concentrations are higher in the overall YAC128 bouton population.

A. DF/F time courses from two rSyph-GCaMP6m-expressing axonal boutons from a

representative YAC128 culture in the presence of TTX (500 nM). The first bouton (top trace) is

inactive during the 3 min imaging session, while the second bouton (bottom trace) shows a clear

spontaneous event (black arrow). Subsequent application of the Ca^{2+} ionophore ionomycin (10

μM) elicits a smaller Ca^{2+} response in the spontaneously active bouton, suggesting a higher basal

cytosolic Ca^{2+} concentration (pre-ionomycin) in the active bouton.

B,C: Ionomycin responses of rSyph-GCaMP6m-expressing boutons in the presence of TTX

following a 3 min imaging session in WT (**B**) and YAC128 (**C**) cultures. Boutons were classified

as active or inactive based on the presence, or lack thereof, of at least one spontaneous Ca^{2+} event

during the initial 3 minute recording. Ionomycin responses were significantly larger in the

population of inactive boutons [WT: 0.633 ± 0.072 DF/F (n=67 boutons); YAC128: 0.410 ± 0.040

DF/F (n=51 boutons)], vs active boutons [WT: 0.191 ± 0.026 DF/F (n=41 boutons; 4 cultures),

p<0.0001 (exact) by Mann Whitney test; YAC128: 0.149 ± 0.016 DF/F (n=46 boutons), p<0.0001

(exact) by Mann Whitney test]. 4 cultures for WT and 4 cultures for YAC128.

D, E: Responses of rSyph-GCaMP6m-expressing axons from a representative WT (**D**) and

YAC128 (**E**) cortical culture to ionomycin (10 μM). Images are from a 4 min imaging session,

with ionomycin application at 3 min and reflect raw GCaMP fluorescence values (arbitrary units)

(as above). To minimize impacts of photobleaching, the top (before) image is the frame

immediately prior ionomycin. The bottom image is a maximum projection of all frames acquired

30s prior and following ionomycin treatment and reflect each pixel's maximum ionomycin

response. The visible grey-value dynamic range is the same in panels **D** and **E**, (150 increments of

the total unprocessed 14 bit-depth), however the 0 value in **E** has been adjusted so that brightness

of the before image is comparable to that of its WT counterpart in **D**; this is so the reader can better

appreciate the YAC128 culture's smaller relative change in fluorescence.

F. Ionomycin responses of the all WT and YAC128 rSyph-GCaMP6m-expressing axons

automatically detected with the aQua software suite (see methods for reference). The WT bouton

population showed a significantly greater ionomycin-mediated change in fluorescence [0.272 ±

0.006 DF/F (n=1681 boutons; 4 cultures)] compared to the YAC128 population [0.190 ± 0.002

DF/F (n=1462 boutons; 4 cultures)], [$p<0.0001$ (exact); Mann Whitney test].

Fig 5: Caffeine increases the frequency of spontaneous Ca^{2+} events and the resting Ca^{2+}

concentration in WT, but not YAC128, cortical axonal boutons.

A, B: Numbers of detected spontaneous rSyph-GCaMP6m axonal events in the presence TTX

(500 nM) during 3 min recordings of 63 x objective fields of WT (**A**) and YAC128 (**B**) cortical

cultures (as above). The control group consists of the same culture fields shown in **Figure 6: E**,

and is compared to separate WT (**A**) or YAC128 (**B**) culture fields imaged in the presence of

caffeine (1 mM) under otherwise like conditions. WT axons showed significantly more action

potential-independent events in the presence of caffeine, than in its absence [203.8 ± 55.2 events/3

min (n=10 fields; 5 cultures)] vs [36.3 ± 5.5 events/3 min (n=12 fields; 4 cultures)] respectively

[$t(20)=3.316$; $p=0.0034$; Student's unpaired-t test]. Numbers of detected YAC128 axonal events

were not significantly different in the presence or absence of caffeine [115.1 ± 28.21 events/3 min

(n=15 fields; 4 cultures)] vs [150.0 ± 28.55 events/3 min (n=8 fields; 4 cultures)] respectively

[$p=0.2188$ (exact); Mann Whitney test].

C. DF/F time courses from representative WT (top) and YAC128 (bottom) rSyph-GCaMP6m-

expressing axonal boutons in the presence of TTX (500 nM) before and immediately following

application of caffeine (1 mM). For this illustration, boutons were selected which did not show

spontaneous events. Rather, note the gradual sustained increase in rSyph-GCaMP6m fluorescence,

which is of larger magnitude in the WT bouton.

D. Caffeine responses of WT and YAC128 rSyph-GCaMP6m-expressing boutons in the presence

of TTX as illustrated in panel **C.** Boutons showing spontaneous events **(as in Figure 6)** were

excluded as the long time course of such events complicated quantification of steady-state

increases in DF/F. Caffeine application elicited a significantly greater increase in rSyph-

GCaMP6m fluorescence in WT boutons [0.342 ± 0.030 DF/F (n=87 boutons)] compared to

YAC128 [0.229 ± 0.039 DF/F (n=54 boutons)] [p=0.0004 (exact); Mann Whitney test]. For A, B

and D: 5 cultures for WT and 5 cultures for YAC128.

Fig 6: In the absence of TTX, presynaptic calcium events are more frequent in WT,

compared to YAC128, cultures.

A, B: DF/F time course from representative WT rSyph-GCaMP6m-expressing axonal boutons in

the absence of TTX, under control conditions (A) and in presences of 5 μM ryanodine (B). Note

these are not the same boutons, as experiments illustrated in this figure were not paired.

C, D: DF/F time course from two different representative YAC128 rSyph-GCaMP6m-expressing

axonal boutons in the absence of TTX, one under control conditions (C) and the other in the

presence of 5 μM ryanodine. (D)

E: Mean Ca^{2+} event frequencies of rSyph-GCaMP6m-expressing boutons in the absence of TTX

in WT and YAC128 cortical cultures in the presence and absence of ryanodine (5 μM). Select

comparisons were made with the Mann Whitney test because event widths in all groups failed the

D'Agostino-Pearson omibus normality test. Events were significantly more frequent in WT

boutons [0.1424 ± 0.0061 Hz (n=272 boutons; 5 cultures)] than in YAC128 boutons [0.0761 ±

0.0044 Hz (n=423 boutons; 6 cultures)] [p<0.0001 (approximate); Mann Whitney test]. Events

were significantly less frequent in WT boutons with ryanodine (5 μM) [0.1179 ± 0.0033 Hz (n=120

boutons, 3 cultures)] than without (above) [p=0.0038 (approximate); Mann Whitney test].

Conversely, event frequencies were significantly more frequent in YAC128 boutons with

ryanodine (5 μM) [0.0886 ± 0.0044 Hz (n=189 boutons, 4 cultures)] than without (above)

[p=0.0003 (approximate); Mann Whitney test].

F, G: DF/F time course from two representative rSyph-GCaMP6m-expressing axonal boutons

from the same WT culture field in the absence of TTX. Bouton (F) shows only faster events (lasting

< 6 s), while in bouton (G) a clear slow event (lasting >6 s) is present. Note the substantially lower

overall event frequency in (G).

H, I: Ca^{2+} event frequencies of WT (H) and YAC128 (I) rSyph-GCaMP6m-expressing boutons

(as above), grouped based on the presence or absence of one or more slow events (> 6 s at half

peak amp). (H): 23/292 (7.9 %) WT boutons showed at least on such slow event with an overall

mean bouton frequency of 0.107 ± 0.016 Hz (n=23), significantly lower than WT boutons lacking

slower events [0.143 ± 0.006 Hz (n=269)] [p=0.0058 (exact); Mann Whitney test] (n=292 boutons,

5 cultures). (I) 44/423 (10.4 %) YAC128 boutons showed at least one such slow event with an

overall mean bouton frequency of 0.052 ± 0.005 Hz (n=44), significantly lower than YAC128

boutons lacking slower events [0.076 ± 0.004 Hz (n=379)] [p=0.0058 (exact); Mann Whitney test]

(n=423 boutons, 6 cultures).

1025

Fig 7: Ryanodine reduces evoked striatal iGluSnR responses in WT, but not YAC128, brain

slices.

A. Image of raw iGluSnFR fluorescence from a representative WT mouse brain slice with viral-

mediated expression of iGluSnFR in its dorsal striatum. Note the square ROI over which evoked

spatially averaged DF/F signals were measured, and the tungsten monopolar stimulating electrode

placed in the corpus callosum.

B. A DF/F image montage from the WT brain slice (shown in A), illustrating a striatum iGluSnFR

response evoked by electrically stimulating cortical axons in the adjacent corpus callosum.

C. Striatum evoked DF/F iGluSnFR responses from a representative WT (top) and YAC128

(bottom) brain slice before and in the presence of ryanodine (5 μM). Ryanodine elicits a substantial

reduction in the peak WT, but not YAC128, iGluSnFR response.

D. Peak evoked striatum iGluSnFR responses from WT and YAC128 slices before and during

treatment of slices with ryanodine (5 μM). Responses were measured repeatedly at 3 min intervals

and were normalized to the average of each slice's 3 baseline (before drug) measurements.

Ryanodine application decreased the amplitude of the evoked iGluSnFR response in WT (n=9

slices; 9 mice), but not YAC128 (n=8 slices; 8 mice). The effect of ryanodine on WT slices was

significantly different than in YAC128 slices, based on a significant time-genotype interaction

(p=0.0094, repeated measure (time) two way ANOVA).

1044

1045

1046

1047

1048   **References:**
1049

A novel gene containing a trinucleotide repeat that is expanded and unstable on Huntington"s
disease chromosomes. The Huntington"s Disease Collaborative Research Group. (1993). A
novel gene containing a trinucleotide repeat that is expanded and unstable on Huntington"s
disease chromosomes. The Huntington"s Disease Collaborative Research Group. *Cell*, *72*(6),
971–983.
Atasoy, D., Ertunc, M., Moulder, K. L., Blackwell, J., Chung, C., Su, J., & Kavalali, E. T.
(2008). Spontaneous and evoked glutamate release activates two populations of NMDA
receptors with limited overlap. *The Journal of Neuroscience : the Official Journal of the*
Society for Neuroscience, *28*(40), 10151–10166. http://doi.org/10.1523/JNEUROSCI.2432-
08.2008
Bachoud-Lévi, A.-C., Ferreira, J., Massart, R., Youssov, K., Rosser, A., Busse, M., et al. (2019).
International Guidelines for the Treatment of Huntington's Disease. *Frontiers in Neurology*,
10, 710. http://doi.org/10.3389/fneur.2019.00710
Beal, M. F., Kowall, N. W., Ellison, D. W., Mazurek, M. F., Swartz, K. J., & Martin, J. B.
(1986). Replication of the neurochemical characteristics of Huntington's disease by
quinolinic acid. *Nature*, *321*(6066), 168–171. http://doi.org/10.1038/321168a0
Beaulieu-Laroche, L., & Harnett, M. T. (2018). Dendritic Spines Prevent Synaptic Voltage
Clamp. *Neuron*, *97*(1), 75–82.e3. http://doi.org/10.1016/j.neuron.2017.11.016
Botelho, E. P., Wang, E., Chen, J. Y., Holley, S., Andre, V., Cepeda, C., & Levine, M. S. (2014).
Differential Synaptic and Extrasynaptic Glutamate-Receptor Alterations in Striatal Medium-
Sized Spiny Neurons of Aged YAC128 Huntington's Disease Mice. *PLoS Currents*, *6*.
http://doi.org/10.1371/currents.hd.34957c4f8bd7cb1f5ec47381dfc811c3
Budde, T., Meuth, S., & Pape, H.-C. (2002). Calcium-dependent inactivation of neuronal
calcium channels. *Nature Reviews. Neuroscience*, *3*(11), 873–883.
http://doi.org/10.1038/nrn959
Buren, C., Parsons, M. P., Smith-Dijak, A., & Raymond, L. A. (2016). Impaired development of
cortico-striatal synaptic connectivity in a cell culture model of Huntington's disease.
Neurobiology of Disease, *87*, 80–90. http://doi.org/10.1016/j.nbd.2015.12.009
Cens, T., Rousset, M., Leyris, J.-P., Fesquet, P., & Charnet, P. (2006). Voltage- and calcium-
dependent inactivation in high voltage-gated Ca(2+) channels. *Progress in Biophysics and*
Molecular Biology, *90*(1-3), 104–117. http://doi.org/10.1016/j.pbiomolbio.2005.05.013
Cepeda, C., Hurst, R. S., Calvert, C. R., Hernández-Echeagaray, E., Nguyen, O. K., Jocoy, E., et
al. (2003). Transient and progressive electrophysiological alterations in the corticostriatal
pathway in a mouse model of Huntington's disease. *The Journal of Neuroscience : the*
Official Journal of the Society for Neuroscience, *23*(3), 961–969.
Chen, T.-W., Wardill, T. J., Sun, Y., Pulver, S. R., Renninger, S. L., Baohan, A., et al. (2013).
Ultrasensitive fluorescent proteins for imaging neuronal activity. *Nature*, *499*(7458), 295–
300. http://doi.org/10.1038/nature12354
de Juan-Sanz, J., Holt, G. T., Schreiter, E. R., de Juan, F., Kim, D. S., & Ryan, T. A. (2017).
Axonal Endoplasmic Reticulum Ca 2+ Content Controls Release Probability in CNS Nerve
Terminals. *Neuron*, *93*(4), 867–881.e6. http://doi.org/10.1016/j.neuron.2017.01.010

Dunwiddie, T. V., & Masino, S. A. (2001). The role and regulation of adenosine in the central
nervous system. *Annual Review of Neuroscience*, *24*(1), 31–55.
http://doi.org/10.1146/annurev.neuro.24.1.31
Emptage, N. J., Reid, C. A., & Fine, A. (2001). Calcium Stores in Hippocampal Synaptic
Boutons Mediate Short-Term Plasticity, Store-Operated Ca2+ Entry, and Spontaneous
Transmitter Release. *Neuron*, *29*(1), 197–208. http://doi.org/10.1016/S0896-6273(01)00190-
8
Espinosa, F., & Kavalali, E. T. (2009). NMDA Receptor Activation by Spontaneous
Glutamatergic Neurotransmission. *Journal of Neurophysiology*, *101*(5), 2290–2296.
http://doi.org/10.1152/jn.90754.2008
Fan, M. M. Y., Fernandes, H. B., Zhang, L. Y. J., Hayden, M. R., & Raymond, L. A. (2007).
Altered NMDA receptor trafficking in a yeast artificial chromosome transgenic mouse
model of Huntington's disease. *The Journal of Neuroscience : the Official Journal of the*
Society for Neuroscience, *27*(14), 3768–3779. http://doi.org/10.1523/JNEUROSCI.4356-
06.2007
Fishbein, I., & Segal, M. (2007). Miniature synaptic currents become neurotoxic to chronically
silenced neurons. *Cerebral Cortex (New York, N.Y. : 1991)*, *17*(6), 1292–1306.
http://doi.org/10.1093/cercor/bhl037
Frank, C. A., Kennedy, M. J., Goold, C. P., Marek, K. W., & Davis, G. W. (2006). Mechanisms
underlying the rapid induction and sustained expression of synaptic homeostasis. *Neuron*,
52(4), 663–677. http://doi.org/10.1016/j.neuron.2006.09.029
Fredj, N. B., & Burrone, J. (2009). A resting pool of vesicles is responsible for spontaneous
vesicle fusion at the synapse. *Nature Neuroscience*, *12*(6), 751–758.
http://doi.org/10.1038/nn.2317
Graveland, G. A., Williams, R. S., & DiFiglia, M. (1985). Evidence for degenerative and
regenerative changes in neostriatal spiny neurons in Huntington's disease. *Science (New*
York, N.Y.), *227*(4688), 770–773.
Gross, G. G., Junge, J. A., Mora, R. J., Kwon, H.-B., Olson, C. A., Takahashi, T. T., et al.
(2013). Recombinant probes for visualizing endogenous synaptic proteins in living neurons.
Neuron, *78*(6), 971–985. http://doi.org/10.1016/j.neuron.2013.04.017
Hantraye, P., Riche, D., Maziere, M., & Isacson, O. (1990). A primate model of Huntington's
disease: behavioral and anatomical studies of unilateral excitotoxic lesions of the caudate-
putamen in the baboon. *Experimental Neurology*, *108*(2), 91–104.
Joshi, P. R., Wu, N.-P., André, V. M., Cummings, D. M., Cepeda, C., Joyce, J. A., et al. (2009).
Age-dependent alterations of corticostriatal activity in the YAC128 mouse model of
Huntington disease. *The Journal of Neuroscience : the Official Journal of the Society for*
Neuroscience, *29*(8), 2414–2427. http://doi.org/10.1523/JNEUROSCI.5687-08.2009
Kavalali, E. T. (2015). The mechanisms and functions of spontaneous neurotransmitter release.
Nature Reviews. Neuroscience, *16*(1), 5–16. http://doi.org/10.1038/nrn3875
Kovalenko, M., Milnerwood, A., Giordano, J., St Claire, J., Guide, J. R., Stromberg, M., et al.
(2018). HttQ111/+ Huntington's Disease Knock-in Mice Exhibit Brain Region-Specific
Morphological Changes and Synaptic Dysfunction. *Journal of Huntington's Disease*, *7*(1),
17–33. http://doi.org/10.3233/JHD-170282
Lindhout, F. W., Cao, Y., Kevenaar, J. T., Bodzęta, A., Stucchi, R., Boumpoutsari, M. M., et al.
(2019). VAP-SCRN1 interaction regulates dynamic endoplasmic reticulum remodeling and

presynaptic function. *The EMBO Journal*, *38*(20), e101345.
http://doi.org/10.15252/embj.2018101345

Llano, I., González, J., Caputo, C., Lai, F. A., Blayney, L. M., Tan, Y. P., & Marty, A. (2000). Presynaptic calcium stores underlie large-amplitude miniature IPSCs and spontaneous calcium transients. *Nature Neuroscience*, *3*(12), 1256–1265. http://doi.org/10.1038/81781

McKinney, R. A., Capogna, M., Dürr, R., Gähwiler, B. H., & Thompson, S. M. (1999). Miniature synaptic events maintain dendritic spines via AMPA receptor activation. *Nature Neuroscience*, *2*(1), 44–49. http://doi.org/10.1038/4548

Meissner, G. (2017). The structural basis of ryanodine receptor ion channel function. *The Journal of General Physiology*, *149*(12), 1065–1089. http://doi.org/10.1085/jgp.201711878

Miller, B. R., Dorner, J. L., Shou, M., Sari, Y., Barton, S. J., Sengelaub, D. R., et al. (2008). Up-regulation of GLT1 expression increases glutamate uptake and attenuates the Huntington's disease phenotype in the R6/2 mouse. *Neuroscience*, *153*(1), 329–337. http://doi.org/10.1016/j.neuroscience.2008.02.004

Milnerwood, A. J., Gladding, C. M., Pouladi, M. A., Kaufman, A. M., Hines, R. M., Boyd, J. D., et al. (2010). Early Increase in Extrasynaptic NMDA Receptor Signaling and Expression Contributes to Phenotype Onset in Huntington's Disease Mice. *Neuron*, *65*(2), 178–190. http://doi.org/10.1016/j.neuron.2010.01.008

Milnerwood, A. J., Kaufman, A. M., Sepers, M. D., Gladding, C. M., Zhang, L., Wang, L., et al. (2012). Mitigation of augmented extrasynaptic NMDAR signaling and apoptosis in cortico-striatal co-cultures from Huntington's disease mice. *Neurobiology of Disease*, *48*(1), 40–51. http://doi.org/10.1016/j.nbd.2012.05.013

Morton, A. J., & Edwardson, J. M. (2001). Progressive depletion of complexin II in a transgenic mouse model of Huntington's disease. *Journal of Neurochemistry*, *76*(1), 166–172. http://doi.org/10.1046/j.1471-4159.2001.00059.x

Morton, A. J., Faull, R. L., & Edwardson, J. M. (2001). Abnormalities in the synaptic vesicle fusion machinery in Huntington's disease. *Brain Research Bulletin*, *56*(2), 111–117. http://doi.org/10.1016/s0361-9230(01)00611-6

Parsons, M. P., Vanni, M. P., Woodard, C. L., Kang, R., Murphy, T. H., & Raymond, L. A. (2016). Real-time imaging of glutamate clearance reveals normal striatal uptake in Huntington disease mouse models. *Nature Communications*, *7*, 11251. http://doi.org/10.1038/ncomms11251

Peled, E. S., Newman, Z. L., & Isacoff, E. Y. (2014). Evoked and spontaneous transmission favored by distinct sets of synapses. *Current Biology : CB*, *24*(5), 484–493. http://doi.org/10.1016/j.cub.2014.01.022

Plotkin, J. L., Day, M., Peterson, J. D., Xie, Z., Kress, G. J., Rafalovich, I., et al. (2014). Impaired TrkB Receptor Signaling Underlies Corticostriatal Dysfunction in Huntington's Disease. *Neuron*, *83*(1), 178–188. http://doi.org/10.1016/j.neuron.2014.05.032

Raymond, L. A., André, V. M., Cepeda, C., Gladding, C. M., Milnerwood, A. J., & Levine, M. S. (2011). Pathophysiology of Huntington's disease: time-dependent alterations in synaptic and receptor function. *Neuroscience*, *198*, 252–273. http://doi.org/10.1016/j.neuroscience.2011.08.052

Reese, A. L., & Kavalali, E. T. (2016). Single synapse evaluation of the postsynaptic NMDA receptors targeted by evoked and spontaneous neurotransmission. *eLife*, *5*, 6336. http://doi.org/10.7554/eLife.21170

Ross, W. N. (2012). Understanding calcium waves and sparks in central neurons. *Nature*
Reviews. Neuroscience, *13*(3), 157–168. http://doi.org/10.1038/nrn3168
Sara, Y., Virmani, T., Deák, F., Liu, X., & Kavalali, E. T. (2005). An isolated pool of vesicles
recycles at rest and drives spontaneous neurotransmission. *Neuron*, *45*(4), 563–573.
http://doi.org/10.1016/j.neuron.2004.12.056
Sepers, M. D., Smith-Dijak, A., LeDue, J., Kolodziejczyk, K., Mackie, K., & Raymond, L. A.
(2017). Endocannabinoid-specific impairment in synaptic plasticity in striatum of
Huntington's disease mouse model. *The Journal of Neuroscience : the Official Journal of the*
Society for Neuroscience, 1739–17. http://doi.org/10.1523/JNEUROSCI.1739-17.2017
Smith-Dijak, A. I., Nassrallah, W. B., Zhang, L. Y. J., Geva, M., Hayden, M. R., & Raymond, L.
A. (2019). Impairment and Restoration of Homeostatic Plasticity in Cultured Cortical
Neurons From a Mouse Model of Huntington Disease. *Frontiers in Cellular Neuroscience*,
13, 209. http://doi.org/10.3389/fncel.2019.00209
Sutton, M. A., Ito, H. T., Cressy, P., Kempf, C., Woo, J. C., & Schuman, E. M. (2006). Miniature
neurotransmission stabilizes synaptic function via tonic suppression of local dendritic protein
synthesis. *Cell*, *125*(4), 785–799. http://doi.org/10.1016/j.cell.2006.03.040
Sutton, M. A., Taylor, A. M., Ito, H. T., Pham, A., & Schuman, E. M. (2007). Postsynaptic
decoding of neural activity: eEF2 as a biochemical sensor coupling miniature synaptic
transmission to local protein synthesis. *Neuron*, *55*(4), 648–661.
http://doi.org/10.1016/j.neuron.2007.07.030
Suzuki, M., Nagai, Y., Wada, K., & Koike, T. (2012). Calcium leak through ryanodine receptor
is involved in neuronal death induced by mutant huntingtin. *Biochemical and Biophysical*
Research Communications, *429*(1-2), 18–23. http://doi.org/10.1016/j.bbrc.2012.10.107
Tang, T. S., Guo, C., Wang, H., Chen, X., & Bezprozvanny, I. (2009). Neuroprotective Effects
of Inositol 1,4,5-Trisphosphate Receptor C-Terminal Fragment in a Huntington's Disease
Mouse Model. *Journal of Neuroscience*, *29*(5), 1257–1266.
http://doi.org/10.1523/JNEUROSCI.4411-08.2009
Tang, T.-S., Tu, H., Chan, E. Y. W., Maximov, A., Wang, Z., Wellington, C. L., et al. (2003).
Huntingtin and huntingtin-associated protein 1 influence neuronal calcium signaling
mediated by inositol-(1,4,5) triphosphate receptor type 1. *Neuron*, *39*(2), 227–239.
Tyebji, S., & Hannan, A. J. (2017). Synaptopathic mechanisms of neurodegeneration and
dementia: Insights from Huntington's disease. *Progress in Neurobiology*, *153*, 18–45.
http://doi.org/10.1016/j.pneurobio.2017.03.008
van de Vrede, Y., Fossier, P., Baux, G., Joels, M., & Chameau, P. (2007). Control of IsAHP in
mouse hippocampus CA1 pyramidal neurons by RyR3-mediated calcium-induced calcium
release. *Pflugers Archiv : European Journal of Physiology*, *455*(2), 297–308.
http://doi.org/10.1007/s00424-007-0277-4
Vonsattel, J. P., Myers, R. H., Stevens, T. J., Ferrante, R. J., Bird, E. D., & Richardson, E. P.
(1985). Neuropathological classification of Huntington's disease. *Journal of Neuropathology*
and Experimental Neurology, *44*(6), 559–577.
Wang, R., & Reddy, P. H. (2017). Role of Glutamate and NMDA Receptors in Alzheimer's
Disease. *Journal of Alzheimer's Disease : JAD*, *57*(4), 1041–1048.
http://doi.org/10.3233/JAD-160763
Wang, Y., DelRosso, N. V., Vaidyanathan, T. V., Cahill, M. K., Reitman, M. E., Pittolo, S., et al.
(2019). Accurate quantification of astrocyte and neurotransmitter fluorescence dynamics for

single-cell and population-level physiology. *Nature Neuroscience*, *22*(11), 1936–1944.
http://doi.org/10.1038/s41593-019-0492-2
Xu, J., Pang, Z. P., Shin, O.-H., & Südhof, T. C. (2009). Synaptotagmin-1 functions as a Ca2+
sensor for spontaneous release. *Nature Neuroscience*, *12*(6), 759–766.
http://doi.org/10.1038/nn.2320
Zeron, M. M., Hansson, O., Chen, N., Wellington, C. L., Leavitt, B. R., Brundin, P., et al.
(2002). Increased Sensitivity to N-Methyl-D-Aspartate Receptor-Mediated Excitotoxicity in
a Mouse Model of Huntington's Disease. *Neuron*, *33*(6), 849–860.
http://doi.org/10.1016/S0896-6273(02)00615-3
1235

Figure 1:

Supplemental Figure 1:

Figure 2:

Supplemental Figure 2:

Figure 3:

Figure 4:

Figure 5:

Figure 6:

Figure 7: