Evolution of sperm morphology in a crustacean genus

with fertilization inside an open brood pouch

Duneau David[1,3,4], Markus Möst[2], Dieter Ebert[3]

[1] Université Toulouse 3 Paul Sabatier, CNRS, IRD; UMR5174, EDB (Laboratoire Évolution & Diversité Biologique); Toulouse, France

[2] University of Innsbruck, Department of Ecology, Technikerstrasse 25, 6020 Innsbruck, Austria

[3] University of Basel, Department of Environmental Sciences, Zoology, Vesalgasse 1, 4051 Basel, Switzerland

[4] Corresponding author: david.duneau@gmail.com

Abstract

Sperm is the most fundamental male reproductive feature. It serves the fertilization of eggs and evolves under sexual selection. Two components of sperm are of particular interest, their number and their morphology. Mode of fertilization is believed to be a key determinant of sperm length across the animal kingdom. External fertilization, unlike internal, favors small and numerous sperm, since sperm density is thinned out in the environment. Here, we study the evolution of sperm morphology in the genus *Daphnia*, where fertilization occurs in a receptacle, the brood pouch, where sperm can constantly be flushed out by a water current. Based on microscopic observations of sperm morphologies mapped on a phylogeny with 15 *Daphnia* and 2 outgroup species, we found that despite the internal fertilization mode, *Daphnia* have among the smallest sperm recorded, as would be expected with external fertilization. Despite being all relatively small compared to other arthropods, sperm length diverged at least twice, once within each of the *Daphnia* subgenera *Ctenodaphnia* and *Daphnia*. Furthermore, species in the latter subgenus also lost the ability of cell compaction by extracellular encapsulation and have very polymorphic sperm with long, and often numerous, filopodia. We discuss the different strategies that *Daphnia* evolved to achieve fertilization success in the females' brood pouch.

Introduction

Sexual selection is a form of natural selection acting on mating and fertilization success. Hence, sperm, the most fundamental male reproductive feature allowing egg fertilization, is expected to adapt to maximize fertilization success. In particular, two components of sperm are under selection, their number and their morphology, the latter including the associated quality. The number of sperm cells per ejaculate is known to evolve in response to several factor: risk of sperm competition for egg fertilization, cryptic female choice and, female receptacle size or the absence of receptacle (Roldan, 2019). While sperm are considered as one of the taxonomically most diverse and rapidly evolving cell types (Birkhead *et al.*, 2009; Ramm *et al.*, 2014), the understanding of the adaptive value of sperm morphology, such as length and shape, remains largely incomplete (Lüpold & Pitnick, 2018). Based on phylogenetic analyses across the animal kingdom, the general rule seems to be that fertilization mode (i.e. whether eggs are fertilized within or outside the female) is a key predictor of sperm length (Kahrl *et al.*, 2021). There is a trade-off between sperm number and length (Immler *et al.*, 2011), and therefore, the production of more sperm comes at the cost of sperm size (Parker, 1982). Sperm are generally shorter and more numerous, in species in which ejaculates are thinned out in aquatic environments, while they are longer and less numerous in species where fertilization occurs within the female (Kahrl *et al.*, 2021; Fitzpatrick *et al.*, 2022). This is because ejaculates are more thinned in the environment than they are within the female's receptacle. Hence, external fertilization mode selects for higher sperm number and associated with it, smaller sperm size, to increase the chance of egg fertilization. On the contrary, internal fertilization mode reduces the thinning effect and allows for the evolution of less sperm per ejaculate and investment in higher sperm quality, e.g. longer sperm cells.

The superorder *Peracarida* is a large clade of crustaceans (e.g. *Amphipoda* and *Isopoda*) having the particularity that females have a chamber where the eggs are brooded: the brood pouch. As fertilization does occur before the females expel their eggs in the environment, such fertilization mode may be considered as internal. However, a water current generated by the filtering apparatus oxygenates the eggs in the brood pouch (Seidl *et al.*, 2002), and it may flush out sperm. In this context, even if the fertilization is internal, we expect that the thinning effect, imposed by the water current, has

an impact on the evolution of sperm length (i.e. reduces their size). Furthermore, males are likely to

evolve persistence traits that allow them to avoid being flushed away. Out of the 4705 species from

which we have a sperm morphology description in the SpermTree database, only 9 species belong to

the superorder of the *Peracarida* (Fitzpatrick *et al.*, 2022). Three of them have sperm described as

aflagellate and the others are described as non-standard, as they have tail-like morphology which are

not a flagellum. Despite being a very recent and rich database, little is known about sperm in this

species rich clade with a particular fertilization mode that may affect the evolution of sperm

morphology.

The water fleas or *Daphnia* are not part of the *Peracarida*, they are *Cladocera*, but they

possess a brood chamber, where fertilization of sexual eggs takes place. They reproduce mostly by

cyclical parthenogenesis, with sexual reproduction and thus egg fertilization being sporadic, but

essential for diapause in freezing and drying habitats and for dispersal. Usually triggered by a change

in environmental conditions and following periods of clonal reproduction during which females only

produce genetically identical daughters, some females produce sexual (haploid) eggs while others

produce males (Deng & Lynch, 1996; DeMeester & Vanoverbeke, 1999). During mating, one or two

males attach to the female to fertilize eggs, which will be released by the female in her brood pouch

after the male(s) departed. The brood pouch is a receptacle formed by the carapace on the dorsal side

of all *Daphnia* species, where early development of clonal and sexual eggs takes place. For the latter,

the cuticular structure of the brood pouch changes to form a protective case, which will be released

upon molting. These resting stages contribute to genetically diverse egg-banks from which future

populations can be established. There is evidence for males competing for fertilization in *Daphnia*

magna (Duneau *et al.*, bioRxiv 2020), but the extend of sperm competition in this receptacle is

unknown. It is not entirely clear where fertilization exactly occurs, whether in the brood pouch or

inside the oviduct exiting in the brood pouch. On a one hand, sperm are not flagellated (Wingstrand,

1978; Wuerz *et al.*, 2017), and it is difficult to see how sperm could enter the oviduct as it is closed

and eggs have to push their way out (Lee *et al.*, 2019). On the other hand, *Daphnia* males have more

or less exaggerated genital papilla, the organ used to deposit sperm in the brood chamber. When

exaggerated, they could help to bring sperm close to the oviduct, which was used to argue that

fertilization occurs in the oviduct (Baldass, 1941) or at its entrance, and reduces the thinning effect of

the water current in the brood pouch.

Although pioneer studies have given key general descriptions to identify the main sperm

structures (Delavault & Berard, 1974; Wingstrand, 1978; Zaffagnini, 1987; Wuerz *et al.*, 2017), only

little is known about morphology of aflagellated sperm in *Daphnia*. All *Anomopoda*, an infraorder

including *Daphnia*, have a vacuolar type of spermatogenesis (Wingstrand, 1978), *i.e.* in the testes, the

spermatids are enclosed in "private" vacuoles in the nutritive cells and are exocytosed into the

testicular lumen after they have decreased strongly in size and matured. When compacted during

maturation, they are generally small, about a few microns in length and width. Sperm of *D. magna* has

been more thoroughly studied with recent technology. This representative of the subgenus

Ctenodaphnia has longer sperm (~10 µm) encapsulated by an acellular capsule, probably to pack more

sperm in the testes, and very short filopodia extending from the cell membrane (Wuerz *et al.*, 2017).

The roles of this capsule and of the filopodia are unclear (Wuerz *et al.*, 2017). Here, we reconstructed

a robust phylogeny of the *Daphnidae* using COI, 12S and 16S rRNA genes based on (Adamowicz *et*

al., 2009; Cornetti *et al.*, 2019) and assessed 15 species representing major clades within *Daphnia* to

better describe the variation in sperm morphology in this genus and to gain insight into its evolution.

Materials and methods

Male *Daphnia* were either sampled from female mass cultures in the laboratory in which male

production was naturally occurring as a consequence of high population density, or from females

exposed to the hormone methyl farnesoate (MF, 400nM final concentration) which is known to induce

male production (Olmstead and Leblanc 2002). We assessed samples covering major clades in the

genus *Daphnia*. For the subgenus *Ctenodaphnia* we included *D. similis* Claus, 1876, *D. sinensis* Gu,

Xu, Li, Dumont et Han, 2013, *D. lumholtzi* Sars, 1885, *D. carinata* King, 1853, *D. magna* Straus,

1820, *D. hispanica* Glagolev and Alonso, 1990, *D. dolichocephala* Sars, 1895, and *D. barbata*

Weltner, 1897. For the subgenus *Daphnia*, we sampled *D. pulex* Leydig, 1860 and the European clade

of *D. pulicaria* Forbes, 1893 as representatives of the *D. pulex* group *sensu lato*. From the same

subgenus we also included samples from the *D. longispina* group *sensu lato*, sometimes also referred

to as 'Hyalodaphnia' (Petrusek *et al.,* 2008). This group was represented by members of the *D.*

longispina complex (Petrusek *et al.,* 2008) - namely *D. longispina* O.F. Müller, 1785*, D. mendotae*

Birge, 1918, *D. galeata* Sars, 1864, and *D. dentifera* Forbes, 1893 – as well as *D. curvirostris*

Eylmann, 1887. For *D. longispina* our sampling also covered the different morphotypes "zschokkei",

"hyalina" and "longispina" (Petrusek *et al.,* 2008).

We induced male production for the *D. longispina* morphotypes "hyalina" and "zschokkei"*, D.*

dentifera, D. mendotea, *D. galeata*, and *D. curvirostris* and collected naturally produced males for *D.*

similis, *D. sinensis*, *D. lumholtzi*, *D. carinata*, *D. magna*, *D. hispanica*, *D. dolichocephala*, *D. barbata*,

D. longispina ("longispina" morphotype), *D. pulex* and *D. pulicaria*.

To collect sperm, we exposed mature males to a 1 % nicotine solution ((-)-Nicotin 162.23

g/mol, supplier: Carl Roth, Germany) to induce ejaculation as in Duneau *et al.* (2012). As only mature

spermatozoa are present in the testicular lumen (Wingstrand, 1978:11; Zaffagnini, 1987:277), this

method allowed us to describe and measure mature sperm and avoid immature ones. Presence of

filopodia on the sperm was recorded, but we did not measure filopodia length. Measurements of the

greatest length of the freshly released sperm were performed with ImageJ (Open source, v. 1.5i) using

photographs taken under phase contrast light at a magnification of 40x. In species with very small

sperm (*D. pulex*, *D. pulicaria*, *D. dolichocephala* and *D. barbata*) we paid particularly attention that

the sperm were photographed just after release from the spermiduct to reduce the possibility of

degradation or to confuse them with other particles. However, it was challenging to take good

photographs of them, and the measurement may be less accurate than for the other species. For

example, *D. pulex* sperm is only around 2 µm in length (Xu *et al.*, 2015). All sperm were also

observed at the moment of release from the ejaculatory opening to verify that their shape corresponds

to what was observed later when they settled and were photographed. We also observed sperm

morphology in sea water to test that osmolarity was not affecting our results. We did not see any effect

and do not report further on these observations. Drawings of male genital papilla were redrawn from

published identification keys (Benzie, 2005; Popova *et al.*, 2016).

For an ancestral trait reconstruction, we reconstructed a phylogeny based on the mitochondrial

COI and 12S and, where available, 16S rRNA genes published in Adamowicz *et al.* (2009) and

Cornetti *et al.* (2019) (for sample information see Supplementary Table 1). Sequences were aligned

with MUSCLE (Edgar, 2004) and a maximum-likelihood tree partially constrained for the topology of

the full mitochondrial genome phylogeny from Cornetti *et al.* (2019) was constructed with iqtree2

(Nguyen *et al.*, 2015; Minh *et al.*, 2020). First, the best model and best-fit partitioning scheme were

inferred with ModelFinder (Chernomor *et al.*, 2016; Kalyaanamoorthy *et al.*, 2017) (-m

MFP+MERGE; for full commands see Supplementary Text 1). We then used the best model and

scheme to calculate a constrained and an unconstrained tree and compared the two trees using tree

topology test with the RELL approximation using 10,000 replicates (Kishino *et al.*, 1990), i.e. the

bootstrap proportion, Kishino-Hasegawa test (Kishino & Hasegawa, 1989), Shimodaira-Hasegawa test

(Shimodaira & Hasegawa, 1999), and expected likelihood weights tests (Strimmer & Rambaut,

2002)), and the approximately unbiased test (Shimodaira, 2002) (-zb 10000 –au). Since there was no

significant difference between constrained and unconstrained trees, we continued with the constrained

tree and calculated SH-aLRT (Guindon *et al.*, 2010) and UFBoot (Hoang *et al.*, 2018) support values

with 10,000 replicates (-bb 10000 -alrt 10000). The constrained tree was then also used for ancestral

state reconstruction using the packages phytools 0.7-70 (Revell, 2012), ape 5.4-1 (Paradis & Schliep,

2019), and geiger (Harmon *et al.*, 2008; Pennell *et al.*, 2014) in R version 4.0.3 (2020-10-10) (R Core

Team, 2020). We obtained an ultrametric dated tree using our ML tree and multiple node calibration

using the mtDNA - fossil calibration - Late Jurassic age bounds from Cornetti *et al.* (2019) and a strict

clock model in *chronos()*. We then calculated mean sperm length values for each measured *Daphnia*

species and added mean values for *Ceriodaphnia reticulata* and *Simocephalus* sp. from (Wingstrand,

1978). For ancestral state reconstruction, we used the sperm length data and the dated tree and fitted a

Brownian motion model of evolutionary change with phytools (code, alignments and input files for

tree reconstruction and ancestral state analysis are deposited on GitHub -

).

Results

Among the species we studied, the genital papilla was only exaggerated in *Daphnia magna*, but

inconspicuous in the other species. Hence, we could not correlate this trait with sperm morphology.

Using microscopic observations of sperm morphologies allowing to measure the maximum length of

rod-shape aflagellated sperm cells, we found that sperm length varied greatly among *Daphnia* species,

ranging from about 2 µm to at least 20 µm (Figure 1 and supplementary figures 1 and 2). Based on

recent *Daphnia* phylogenies (Adamowicz *et al.*, 2009; Cornetti *et al.*, 2019), we found that there was a

clear phylogenetic signal in sperm length across *Daphnia*, but length clusters are polyphyletic. Sperm

length diverged at least twice in the genus *Daphnia* (Figure 1), once in the subgenus *Daphnia* and

once in the subgenus *Ctenodaphnia*. We also found that the assessed species in the *D. longispina*

clade, i.e. *D. longispina*, *D. dentifera*, *D. mendotae* and *D. galeata*, have entirely lost the capsule

compacting sperm and had very long filopodia (Figure 2 C and D, Supplementary figures 1, 2 and 3).

Ceriodaphnia and *Simocephalus*, both members of the family Daphniidae and used here as outgroups,

have sperm of the vacuolar spermatogenesis type, like *Daphnia* species and all other clades in the

infraorder *Anomopoda*. Their sperm have been described, based on electron microscopy, as small (2 to

6 µm), more or less rod-shaped and strongly compacted in their capsule (e.g. Figure 2A and

Wingstrand, 1978:25-26). This information based on several species allowed to assume, although with

caution, that the most parsimonious ancestral sperm type in the genus *Daphnia* was rather short. The

subgenus *Ctenodaphnia*, except for *D. dolichocephala* and *D. barbata* who had compacted and small

sperm, evolved non-compacted and elongated sperm, several times larger than the putative ancestral

morphology (Figure 1 and 2B). A similar adaptation occurred in *Daphnia s. l.*. In accordance with

reports from Xu et al. (2015) on *Daphnia pulex* sperm length, we observed that members of the *D.*

pulex group *sensu lato* conserved the small and compacted sperm morphology, whereas members of

the *D. longispina* group *sensu lato* evolved larger elongated sperm (Figure 1). Furthermore, in the

same ejaculate from a single *D. longispina* male, sperm can have a strong polymorphism in the

number and length of filopodia per sperm cell. The cell can have, on each side, one long filopodium or

many shorter ones (Figure 2C and 2D, and Supplementary figure 3). These filopodia can be several

times the length of the sperm (not measured here) (Figure 2 and Supplementary figure 1 and 3).

Discussion

Since *Daphnia* males ejaculate in the female brood pouch, the fertilization mode is here

considered to be internal. However, as a water current generated by the filtering apparatus constantly

circulates through the brood pouch (Seidl *et al.*, 2002), male *Daphnia* face the challenge to have their

sperm flushed out. This phenomenon could be seen as a form of cryptic female choice, as it increases

the thinning effect. When two males copulate at the same time with a female, the brood chamber is a

place for direct sperm competition to occur. Interspecies variations in such sexual selection may be

present and shape sperm evolution. Males of different species may have different features to increase

their chance to successfully sire offspring. We speculate that when species have a brood pouch, males

may be selected to: 1- produce more, but smaller sperm, 2- deposit their sperm the closest to the

oviduct, 3- develop structures to reduce the chance that sperm is flushed out.

Our assessment of sperm morphology over 15 *Daphnia* species uncovered clearly structured

phylogenetic variation in sperm length. It diverged at least twice, once in the subgenus *Daphnia* and

once in the subgenus *Ctenodaphnia*. Our study reports that *Daphnia* have small rod-shape aflagellated

sperm with the species averages ranging from 2.6 to 13 µm. Considering that sperm in arthropods

measure on average 1034.4 µm (±81.7 SE and ±3051.5 SD) (based on 1394 arthropod species in

SpermTree database, (Fitzpatrick *et al.*, 2022)), our work allows to conclude that sperm are relatively

small in *Daphnia*. This supports the hypothesis that the particular internal fertilization of species with

brood pouch do not necessarily favor the evolution of larger sperm, most probably due to the thinning

of sperm density imposed by the constant water flow.

Variation in aflagellated sperm length within each *Daphnia* subgenus was ranging from 2.6 to

13 µm in the subgenus *Ctenodaphnia* and from 3.8 to 9.9 µm in the subgenus *Daphnia*. From the 185

species with aflagellated sperm of which the length is available in the SpermTree database, the median

of the averaged total length is 5 µm, with a range from 1 to 500 µm (Calculated from SpermTree

database, Fitzpatrick *et al.*, 2022). Two *Daphnia* species from each subgenus had mean sperm length

below this median length of 5 µm, while all other species were above this. Part of the size of sperm is

due to an extracellular compaction by an extracellular vacuole which takes place before the mature

sperm is released into the spermiduct (Wingstrand, 1978). As most sperm production happens when

males are juveniles, the total amount of sperm material is limited by the size of the spermiduct (Wuerz

et al., 2017). By limiting the total amount of sperm material stored, this constraint may put selection

on the degree of compaction allowing to store more cells in the duct. In such a case, *Daphnia* species

with "large" sperm may had relatively relaxed selection on sperm compaction, in comparison to

species with "small" and very compacted sperm. We propose that although, fertilization inside a brood

pouch favors smaller sperm probably to favor numbers, the evolutionary changes in sperm length we

observed may in part be underlined by a change in the mechanism of cell compaction before

maturation due to relaxed selection for extreme compaction in some species.

Another way to increase the chance of fertilization in the brood pouch is to deposit the sperm

closer to the oviduct. Few *Daphnia* species are known to have exaggerated genital papilla (Flössner,

2000; Benzie, 2005). We wondered if the penis-like apparatus, which could help to position the sperm

closer to the oviduct and thus reduce the thinning effect, could correlate with sperm size. However, we

found that despite the phylogenetic signal in sperm length, only one species included in our analysis,

D. magna, has an exaggerated genital papilla. Hence, genital papilla morphology cannot be used to

explain the phylogenetic structure we observed. However, strength of sexual selection is a function of

how often male ejaculates compete for fertilization, in particular via direct sperm competition and

sperm competition through cryptic female choice. If exaggerated genital papilla evolved under such

male-male competition and reduce the thinning effect by improving the chance that sperm are

deposited close to the oviduct, one can speculate that this trait evolves when strong sexual selection

occurs in the system. It is difficult to estimate the intensity and frequency of sexual selection in the

system, especially for each species. The induction and frequency of sexual reproduction in *Daphnia*

depends on the species and the environment and thus constitutes a locally adapted trait (Roulin *et al.*,

2013). In unstable and short-lived habitats, such as small rockpools or ponds in strongly seasonal

environments like deserts and arctic sites, few asexual generations occur before diapause

recommences. In stable environments, such as large lakes and ponds in temperate mild climatic

regions, many asexual generations may occur before the next sexual generation and most individuals

will never go through a sexual cycle. Traditionally, the latter case received more attention by *Daphnia*

researchers, leading to the biased impression that sexual reproduction, and thus the occurrence of

males, is generally rare. However, in *Daphnia magna*, there is a large heterogeneity among

populations. In unstable environments, males can be periodically abundant, and several males can be

found copulating at the same time with a female, suggesting that sexual selection can be strong, maybe

more than in most other species (Duneau *et al.*, bioRxiv 2020). Noticeably, *D. magna* is the only

species here that have exaggerated genital papilla and the only species for which matings with multiple

males have been reported (Duneau *et al.*, 2020). One could wonder if the evolution of exaggerated

genital papilla could have evolved in *D. magna* due to selection to improve sperm deposition close to

the oviduct and reduce the thinning effect in a species under strong sexual selection. More work is

needed to strengthen this hypothesis. One way would be to compare the variation in genital papilla

size between stable and unstable *D. magna* populations, as those populations should differ in the

intensity of sexual selection. We would expect that individuals from stable populations would have

less exaggerated genital papilla than those from unstable populations.

In an open brood pouch, sperm would gain an advantage if they could attach to the inner lining

of the pouch, so the water current will not flush them out. We did not observe a structure supporting

the idea that encapsulated sperm can attach to the inner lining. However, it is possible that, like in

crayfish (Niksirat *et al.*, 2014), the extracellular capsule might break post-copulation and reveal

filopodia which could attach to surfaces. In *Daphnia magna*, such filopodia have been reported inside

the extracellular vacuole (Wuerz *et al.*, 2017), but they are very small and difficult to resolve, even

with electron microscopy. Our phylogenetic analysis of sperm morphology revealed a monophyletic

clade (i.e. *D. longispina* species complex) which evolved an apparent loss of the capsule and the gain

of long, sometimes numerous, filopodia of diverse shapes. The lengths of these filopodia may be a

multiple of the length of the sperm cell (Fig. 2). We did not observe any movement of the filopodia

and therefore consider it is unlikely that the flagella are used to move towards the eggs. The filopodia

are very flexible and can be forked. But, as sperm features are expected to be adaptations to their

specific fertilization environment (Pitnick *et al.*, 2009), it is reasonable to assume that the filopodia

play a role in fertilization. This role may be related to the fusion with the oocyte, but it may also

reduce the chances to become flushed out from the brood pouch. Hence, we suggest that the loss of the

capsule may have evolved from the reduced selection for compaction and the advantage to improve

fusion with the oocyte and attach to the brood pouch.

Acknowledgements

We thank Jürgen Hottinger for help in the laboratory, Luca Cornetti for help with the alignment files

from his study on *Daphnia* phylogeny, and Jeremias Brand, Vitor Faria, Scott Pitnick, Lukas Schärer

and Axel Wiberg for discussions about the study. We thank Birgit Schlick-Steiner and the Molecular

Ecology group for hosting DD during his research stay. DD was supported by the French Laboratory

of Excellence project 'TULIP' (ANR-10-LABX-41; ANR-11-IDEX-0002–02). MM was supported by

an FWF Austrian Science Fund stand-alone project (P29667-B25), DE was supported by a grant from

the Swiss National Science Foundation.

References

Adamowicz, S.J., Petrusek, A., Colbourne, J.K., Hebert, P.D.N. & Witt, J.D.S. 2009. The scale of

divergence: A phylogenetic appraisal of intercontinental allopatric speciation in a passively

dispersed freshwater zooplankton genus. *Molecular Phylogenetics and Evolution* **50**: 423–436.

Baldass, F. 1941. Entwicklung von Daphnia pulex. In: *Abteilung Fur Anatomie und Ontogenie der*

Tiere.

Benzie, J.A.H. 2005. *Cladocera: The genus Daphnia (including Daphniopsis)*. Backhuys Publishers,

292   Leiden, The Netherlands.

293   Birkhead, T.R., Hosken, D.J. & Pitnick, S. 2009. *Sperm biology: an evolutionary perspective*,

294   Academic P. Elsevier.

295   Chernomor, O., Von Haeseler, A. & Minh, B.Q. 2016. Terrace Aware Data Structure for

296   Phylogenomic Inference from Supermatrices. *Systematic Biology* **65**: 997–1008.

297   Cornetti, L., Fields, P.D., Van Damme, K. & Ebert, D. 2019. A fossil-calibrated phylogenomic

298   analysis of *Daphnia* and the Daphniidae. *Molecular Phylogenetics and Evolution* **137**: 250–262.

299   Delavault, R. & Berard, J.J. 1974. Ultrastructural study of spermatogenesis in *Daphnia magna* Straus

300   (Entomostraca, Branchiopoda, Cladocerae).

301   DeMeester, L. & Vanoverbeke, J. 1999. An uncoupling of male and sexual egg production leads to

302   reduced inbreeding in the cyclical parthenogen Daphnia. *Proceedings of the Royal Society of*

303   *London. Series B: Biological Sciences* **266**: 2471–2477.

304   Deng, H.W. & Lynch, M. 1996. Change of Genetic Architecture in Response to Sex. *Genetics* **143**:

305   203–212.

306   Duneau, D., Altermatt, F., Ferdy, J.-B.F., Ben-Ami, F. & Ebert, D. 2020. Estimation of the propensity

307   for sexual selection in a cyclical parthenogen. *bioRxiv*, doi:

308   https://doi.org/10.1101/2020.02.05.935148.

309   Duneau, D., Luijckx, P., Ruder, L.F. & Ebert, D. 2012. Sex-specific effects of a parasite evolving in a

310   female-biased host population. *BMC Biology* **10**: 104.

311   Edgar, R.C. 2004. MUSCLE: Multiple sequence alignment with high accuracy and high throughput.

312   *Nucleic Acids Research* **32**: 1792–1797.

313   Fitzpatrick, J.L., Kahrl, A.F. & Snook, R.R. 2022. SpermTree, a species-level database of sperm

314 morphology spanning the animal tree of life. *Scientific Data* **9**: 1–6. Springer US.

315 Flössner, D. 2000. *Die Haplopoda und Cladocera (ohne Bosminidae) Mitteleuropas.* Backhuys

316  Publishers, Leiden, The Netherlands.

317 Guindon, S., Dufayard, J.F., Lefort, V., Anisimova, M., Hordijk, W. & Gascuel, O. 2010. New

318  algorithms and methods to estimate maximum-likelihood phylogenies: Assessing the

319  performance of PhyML 3.0. *Systematic Biology* **59**: 307–321.

320 Harmon, L.J., Weir, J.T., Brock, C.D., Glor, R.E. & Challenger, W. 2008. GEIGER: Investigating

321  evolutionary radiations. *Bioinformatics* **24**: 129–131.

322 Hoang, D.T., Chernomor, O., von Haeseler, A., Minh, B.Q. & Vinh, L.S. 2018. UFBoot2: Improving

323  the Ultrafast Bootstrap Approximation. *Molecular Biology and Evolution* **35**: 518–522.

324 Immler, S., Pitnick, S., Parker, G.A., Durrant, K.L., Lüpold, S., Calhim, S., *et al.* 2011. Resolving

325  variation in the reproductive tradeoff between sperm size and number. *Proceedings of the*

326  *National Academy of Sciences of the United States of America* **108**: 5325–5330.

327 Kahrl, A.F., Snook, R.R. & Fitzpatrick, J.L. 2021. Fertilization mode drives sperm length evolution

328  across the animal tree of life. *Nature Ecology and Evolution* **5**: 1153–1164. Springer US.

329 Kalyaanamoorthy, S., Minh, B.Q., Wong, T.K.F., Von Haeseler, A. & Jermiin, L.S. 2017.

330  ModelFinder: Fast model selection for accurate phylogenetic estimates. *Nature Methods* **14**:

331  587–589.

332 Kishino, H. & Hasegawa, M. 1989. Evaluation of the maximum likelihood estimate of the

333  evolutionary tree topologies from DNA sequence …. *Journal of Molecular Evolution* **29**: 170–

334  179.

335 Kishino, H., Miyata, T. & Hasegawa, M. 1990. Maximum likelihood inference of protein phylogeny

336  and the origin of chloroplasts. *Journal of Molecular Evolution* **31**: 151–160.

337 Lee, D., Nah, J.S., Yoon, J., Kim, W. & Rhee, K. 2019. Live observation of the oviposition process in

338  *Daphnia magna*. *PLoS ONE* **14**: 1–9.

Lüpold, S. & Pitnick, S. 2018. Sperm form and function: What do we know about the role of sexual

selection? *Reproduction* **155**: R229–R243.

Minh, B.Q., Schmidt, H.A., Chernomor, O., Schrempf, D., Woodhams, M.D., Von Haeseler, A., *et al.*

2020. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the

Genomic Era. *Molecular Biology and Evolution* **37**: 1530–1534.

Nguyen, L.T., Schmidt, H.A., Von Haeseler, A. & Minh, B.Q. 2015. IQ-TREE: A fast and effective

stochastic algorithm for estimating maximum-likelihood phylogenies. *Molecular Biology and*

Evolution **32**: 268–274.

Niksirat, H., Kouba, A. & Kozák, P. 2014. Post-mating morphological changes in the spermatozoon

and spermatophore wall of the crayfish Astacus leptodactylus: Insight into a non-motile

spermatozoon. *Animal Reproduction Science* **149**: 325–334. Elsevier B.V.

Paradis, E. & Schliep, K. 2019. Ape 5.0: An environment for modern phylogenetics and evolutionary

analyses in R. *Bioinformatics* **35**: 526–528.

Parker, G.A. 1982. Why are there so many tiny sperm? sperm competition and the maintenance of two

sexes. *Journal of Theoretical Biology* **96**: 281–294.

Pennell, M.W., Eastman, J.M., Slater, G.J., Brown, J.W., Uyeda, J.C., Fitzjohn, R.G., *et al.* 2014.

Geiger v2.0: An expanded suite of methods for fitting macroevolutionary models to phylogenetic

trees. *Bioinformatics* **30**: 2216–2218.

Pitnick, S., Hosken, D.J. & Birkhead, T.R. 2009. Sperm morphological diversity. In: *Sperm Biology*,

pp. 69–149.

Popova, E. V., Petrusek, A., Kořínek, V., Mergeay, J., Bekker, E.I., Karabanov, D.P., *et al.* 2016.

Revision of the old world *Daphnia* (Ctenodaphnia) *similis* group (Cladocera: Daphniidae).

Zootaxa **4161**: 1–40.

R Core Team. 2020. R: A language and environment for statistical computing. Vienna, Austria.

Ramm, S.A., Schärer, L., Ehmcke, J. & Wistuba, J. 2014. Sperm competition and the evolution of

spermatogenesis. *Molecular Human Reproduction* **20**: 1169–1179.

Revell, L.J. 2012. phytools: An R package for phylogenetic comparative biology (and other things).

Methods in Ecology and Evolution **3**: 217–223.

Roldan, E.R.S. 2019. Sperm competition and the evolution of sperm form and function in mammals.

Reproduction in Domestic Animals **54**: 14–21.

Roulin, A.C., Routtu, J., Hall, M.D., Janicke, T., Colson, I., Haag, C.R., *et al.* 2013. Local adaptation

of sex induction in a facultative sexual crustacean: insights from QTL mapping and natural

populations of *Daphnia magna*. *Molecular Ecology* **22**: 3567–3579.

Seidl, M.D., Pirow, R. & Paul, R.J. 2002. Water fleas (*Daphnia magna*) provide a separate ventilatory

mechanism for their brood. *Zoology* **105**: 15–23.

Shimodaira, H. 2002. An approximately unbiased test of phylogenetic tree selection. *Systematic*

Biology **51**: 492–508.

Shimodaira, H. & Hasegawa, M. 1999. Multiple comparisons of log-likelihoods with applications to

phylogenetic inference. *Molecular Biology and Evolution* **16**: 1114–1116.

Strimmer, K. & Rambaut, A. 2002. Inferring confidence sets of possibly misspecified gene trees.

Proceedings of the Royal Society B: Biological Sciences **269**: 137–142.

Wingstrand, K.G. 1978. Comparative spermatology of the Crustacea Entomostraca; 1, Subclass

Branchiopoda. *Biologiske Skrifter* **22**: 1–67.

Wuerz, M., Huebner, E. & Huebner, J. 2017. The morphology of the male reproductive system,

spermatogenesis and the spermatozoon of *Daphnia magna* (Crustacea: Branchiopoda). *Journal*

of Morphology **278**: 1536–1550.

Xu, S., Ackerman, M.S., Long, H., Bright, L., Spitze, K., Ramsdell, J.S., *et al.* 2015. A male-specific

genetic map of the microcrustacean *Daphnia pulex* based on single-sperm whole-genome

sequencing. *Genetics* **201**: 31–38.

Zaffagnini, F. 1987. Reproduction in *Daphnia*. In: *Daphnia* (R. H. Peters & R. De Bernadi, eds), p.

280. Istituto Italiano di Idrobiologia, Pallanza.

Figure legends

Figure 1: Evolution of sperm length and genital papilla morphology in *Daphnia*. Phylogenetic tree is modified from Cornetti et al. (2019) and Adamowicz et al. (2009). Color gradient represents sperm length and length of scale bar indicates to branch length in million years. Drawings represent the genital papilla of the males (redrawn from Benzie (2005) and Popova et al. (2016)). The red arrow indicates the atypical exaggerated genital papilla of *D. magna*. Photographs show an example of sperm for each species. The graph on the right represents the difference in sperm length among males. The mean sperm length was calculated from 2 to 3 individuals per species. Numbers under the mean represents the number of measured sperm. Letters on the right illustrates whether median sperm lengths are significantly different (different letters) or not (same letter) among species based on pairwise Kruskal-Wallis comparisons corrected for multiple testing using the FDR method. Numbers in phylogeny represent the node of the two main *Daphnia* clades: 1: subgenus *Ctenodaphnia*; 2: subgenus *Daphnia.*

Figure 2: Examples of sperm morphologies in the Daphniidae. A- *Ceriodaphnia laticauda*, small and encapsulated elongated sperm (blue color in figure 1) representative of one of the species used as outgroup. Photo from Wingstrand (1978). B- *D. magna*, large and encapsulated elongated sperm (blue color in figure 1); C and D - *D. longispina*, two examples of typical non-encapsulated and elongated sperm with filopodia (green color in figure 1).

Figure 1: Evolution of sperm length and genital papilla morphology in *Daphnia*. Phylogenetic tree is modified from Cornetti et al. (2019) and Adamowicz et al. (2009). Color gradient represents sperm length and length of scale bar indicates to branch length in million years. Drawings represent the genital papilla of the males (redrawn from Benzie (2005) and Popova et al. (2016)). The red arrow indicates the atypical exaggerated genital papilla of *D. magna*. Photographs show an example of sperm for each species. The graph on the right represents the difference in sperm length among males. The mean sperm length was calculated from 2 to 3 individuals per species. Numbers under the mean represents the number of measured sperm. Letters on the right illustrates whether median sperm lengths are significantly different (different letters) or not (same letter) among species based on pairwise Kruskal-Wallis comparisons corrected for multiple testing using the FDR method. Numbers in phylogeny represent the node of the two main *Daphnia* clades: 1: subgenus *Ctenodaphnia*; 2: subgenus *Daphnia*.

Figure 2: Examples of sperm morphologies in the Daphniidae. A- *Ceriodaphnia laticauda*, small and encapsulated elongated sperm (blue color in figure 1) representative of one of the species used as outgroup. Photo from Wingstrand (1978). B- *D. magna*, large and encapsulated elongated sperm (blue color in figure 1); C and D - *D. longispina*, two examples of typical non-encapsulated and elongated sperm with filopodia (green color in figure 1).