# Metagenomic Insights into Microbial Metabolisms of a Sulfur-Influenced Glacial Ecosystem

**Running Title:**

**Metagenomics of a Sulfur-Influenced Glacial Ecosystem**

Christopher B. Trivedi[a], Blake W. Stamps[a], Graham E. Lau[b], Stephen E. Grasby[c], Alexis S. Templeton[b], John R. Spear[a,#]

[a]Department of Civil and Environmental Engineering, Colorado School of Mines, Golden, CO, 80401 USA

[b]Department of Geological Sciences, University of Colorado Boulder, Boulder, CO, 80309 USA

[c]Geological Survey of Canada-Calgary, Calgary, AB, T2L2A7 Canada

[#]Corresponding author: John R. Spear, jspear@mines.edu

Current addresses:

Christopher B Trivedi – Interface Geochemistry, GFZ German Research Centre for Geosciences, Helmholtz Centre Potsdam, Potsdam, Brandenburg 14473 Germany

Blake W Stamps – UES, Inc., Dayton, OH, 45432 USA

Graham E Lau – Blue Marble Space Institute of Science, Seattle, WA, 98154 USA

**Abstract**

Biological sulfur cycling in polar, low-temperature ecosystems is an understudied phenomenon in part due to difficulty of access and the ephemeral nature of such environments. One such environment where sulfur cycling plays an important role in microbial metabolisms is located at Borup Fiord Pass (BFP) in the Canadian High Arctic. Here, transient springs emerge from the toe of a glacier creating a large proglacial aufeis (spring-derived ices) that are often covered in bright yellow/white sulfur, sulfate, and carbonate mineral precipitates that are accompanied by a strong odor of hydrogen sulfide. Metagenomic sequencing from multiple sample types at sites across the BFP glacial system produced 31 highly complete metagenome assembled genomes (MAGs) that were queried for sulfur-, nitrogen- and carbon-cycling/metabolism genes. Sulfur cycling, especially within the Sox complex of enzymes, was widespread across the isolated MAGs and taxonomically associated with the bacterial classes *Alpha-*, *Beta-*, *Gamma-*, and *Epsilon- Proteobacteria*. While this does agree with previous research from BFP implicating organisms within the *Gamma-* and *Epsilon- Proteobacteria* as the primary classes responsible for sulfur oxidation, our new data suggests putative sulfur oxidation by organisms within *Alpha-* and *Beta- Proteobacterial* classes which was not predicted. These findings indicate that in a low-temperature, ephemeral sulfur-based environment such as this, functional redundancy may be a key mechanism that microorganisms use to co-exist whenever energy is limited and/or focused by redox chemistry.

**Importance**

Borup Fiord Pass is a unique environment characterized by a sulfur-enriched glacial ecosystem, in the low-temperature environment of the Canadian High Arctic. This unique

combination makes BFP one of the best analog sites for studying icy, sulfur-rich worlds outside

of our own, such as Europa and Mars. The site also allows investigation of sulfur-based

microbial metabolisms in cold environments here on Earth. Herein, we report whole genome

sequencing data that suggests sulfur cycling metabolisms at BFP are more widely used across

bacterial taxa than predicted. From our data, the metabolic capability of sulfur oxidation among

multiple community members appears likely due to functional redundancy within their genomes.

Functional redundancy, with respect to sulfur-oxidation at BFP, may indicate that this dynamic

ecosystem hosts microorganisms that are able to use multiple sulfur electron donors alongside

other important metabolic pathways, including those for carbon and nitrogen.

## 58  1    Introduction

Arctic and Antarctic ecosystems such as lakes (Jungblut et al., 2016), streams

(Niederberger et al., 2009; Andriuzzi et al., 2018), groundwater-derived ice (aufeis; Grasby et al.,

2014; Hodgkins et al., 2004), and saturated sediments (Lamarche-Gagnon et al., 2015) provide

insight into geochemical cycling and microbial community dynamics in low temperature

environments. One such ecosystem in the Canadian High Arctic, Borup Fiord Pass (BFP), is

located on northern Ellesmere Island, Nunavut, Canada and contains a transient sulfidic spring

system discharging through glacial ice and forming supra and proglacial ice deposits (aufeis) as

well as cryogenic mineral precipitates (Figure 1; Grasby et al., 2003).

Borup Fiord Pass is a north-south trending valley through the Krieger mountains at

81°01' N, 81°38' W (Figure 1, inset). The spring system of interest is approximately 210-240 m

above sea level  and occurs at the toe of two coalesced glaciers (Grasby et al., 2003). The spring

system is thought to be perennial in nature but discharges from different locations along the

glacier from year to year (Grasby et al., 2003; Gleeson et al., 2010; Wright et al., 2013; Lau et

al., 2017). However, given lack of winter observations it is not confirmed that spring flow is

year-round or limited to the warmer spring and summer months. However, growth of large aufeis

formations through the dark season support continued discharge through the winter. A lateral

fault approximately 100 m south of the toe of the glacier has been implicated in playing a role in

the subsurface hydrology, in that this may be one of the areas where subsurface fluid flow is

focused to the surface (Grasby et al., 2003; Scheidegger, et al. 2012). Furthermore, organic

matter in shales along this fault may be a source of carbon for subsurface microbial processes

such as biological sulfate reduction (BSR; Lau et al., 2017).

Previous research at BFP has surveyed microbial activity, redox bioenergetics (Wright et

al., 2013), the biomineralization of elemental sulfur ($S^0$; (Grasby et al., 2003; Gleeson et al.,

2011; Lau et al., 2017), and the cryogenic carbonate vaterite (Grasby, 2003). Wright et al. (2013)

produced a metagenome in the context of geochemical data to constrain the bioenergetics of

microbial metabolism from a mound of elemental sulfur sampled at the site in 2012, and found

that at least in surface mineral deposits, sulfur oxidation was likely the dominant metabolism

present among *Epsilonproteobacteria*. An exhaustive 16S rRNA gene sequencing study

conducted on samples collected in 2014-2017 revealed an active and diverse assemblage of

autotrophic and heterotrophic microorganisms in melt pools, aufeis, spring fluid, and surface

mineral deposits that persist over multiple years, contributing to a basal community present in

the system regardless of site or material type (Trivedi et al., 2018). However, to date no work has

been attempted to fully understand and characterize the metabolic capability of the microbial

communities adapted to this glacial ecosystem, including sulfur metabolism beyond the surface

precipitates at the site.

Microbial sulfur oxidation and reduction, which spans an eight electron transfer between

the most oxidized and reduced states, is important in a multitude of diverse extreme

environments, such as hydrothermal vents & deep subsurface sediments (Dahl and Friedrich,

2008), microbial mats (Ley et al., 2006; Feazel et al., 2008; Robertson et al., 2009), sea ice

(Boetius et al., 2015), glacial environments (Mikucki and Priscu, 2007; Purcell et al., 2014), and

Arctic hypersaline springs (Perreault et al., 2008; Niederberger et al., 2009). Two predominant

forms of sulfur-utilizing microorganisms exist: sulfur-oxidizing microorganisms (SOMs), which

use reduced compounds (e.g. $H_2S$ and $S^0$) as electron donors, and sulfate-reducing

microorganisms (SRMs), which use oxidized forms of sulfur (e.g. $SO_4^{2-}$ and $SO_3^{2-}$) as electron

acceptors. SOMs are metabolically and phylogenetically diverse (Friedrich et al., 2001, 2005;

Dahl and Friedrich, 2008), and can often autotrophically fix carbon dioxide ($CO_2$) while using a

variety of electron acceptors (Mattes et al., 2013). Likewise, SRMs can utilize a range of electron

donors, including organic carbon, inorganic carbon, methane, hydrogen, metallic iron (Enning et

al., 2012), and even heavier metals such as uranium, where the soluble U(VI) is converted to the

insoluble U(IV) under anoxic conditions (Spear et al., 1999, 2000). SRMs are particularly

important in sulfate-rich marine environments where they contribute as much as 50% of total

global organic carbon oxidation (Thullner et al., 2009; Bowles et al., 2014).

Some research has been conducted on microbial sulfur-cycling in low-temperature polar

environments in both Antarctica (Mikucki et al., 2009, 2016) and the Arctic (Niederberger et al.,

2009; Purcell et al., 2014), including BFP (Gleeson et al., 2011; Grasby et al., 2012; Wright et

al., 2013). However, the organisms that participate in sulfur cycling and which metabolic

pathways they utilize in these low-temperature environments remain understudied. It is important

that we better classify these systems as they are key to our understanding of how life can adapt to

potentially adverse conditions (e.g., low-temperature and sulfidic conditions). Furthermore, these

adaptations may also be able to inform us about what to search for on worlds outside of Earth,

for instance Europa, where we know that low-temperature, sulfur-rich conditions exist (Carlson

et al., 1999), or Mars, where gypsum was discovered in the North Polar Cap (Massé et al., 2012).

We collected samples over multiple years (2014, 2016, and 2017; Figure 2) and from

varied sample types (e.g., aufeis, melt pools, spring fluids, and surface mineral precipitates;

Figure S2) for metagenomic whole genome sequencing (WGS). These data were assembled and

binned into Metagenome Assembled Genomes (MAGs), which were queried to identify which

metabolic processes are present and dominant across multiple sample types at BFP. Analysis of

these MAGs revealed the presence of sulfur oxidation genes (namely those part of the Sox

operon) across multiple taxonomic genomes, including in genomes related to organisms where it

is not predicted to be metabolically viable. This may indicate a form of functional redundancy

present in the BFP system where organisms from other Phyla are able to take advantage of the

abundance of reduced sulfur for metabolic processes. While sulfur metabolism was the focus of

this study, genes for carbon, nitrogen, oxygen, hydrogen, and methane metabolisms were queried

as this can provide information about energy utilization and potential electron acceptors.

**2    Results**

*Assembly and Identification of Metagenome Assembled Genomes*

High throughput sequencing of total environmental genomic DNA (gDNA) extracted

from nine BFP biomass samples produced paired-end reads that were trimmed and co-assembled

*de novo*. The co-assembly produced 1.16 Gbp in 477,394 contigs, with the largest being 406,785

bp long and the average being 2,440 bp long. A summary of all assembly statistics is available

within supplementary Table S1. After assembly, a total of 166 bins were identified after

CONCOCT binning (Alneberg et al., 2014) and further refinement within Anvi'o  (Eren et al.,

2015). Of these, 31 were classified as MAGs (defined as bins with greater than 50% estimated

genome completeness and less than 10% redundancy) after quality control within CheckM

(Parks et al., 2015). Of the 31 MAGs, nine were considered highly complete with low

redundancy (90% and 10%, respectively). Further summary statistics of sequencing data are

available in Table 1. MAG distribution was not even between samples. One broadly distributed

MAG, MAG 31 (most closely related to the genus *Flavobacterium*) had abundant contribution

from samples A6 (aufeis), 3B, 4E, and 6B (all mineral precipitate samples), and moderate

contributions from A4b and M4b (aufeis and melt pool samples, respectively). However, MAG

31 had the lowest completion of reported MAGs.

All MAGs were identified within the domain Bacteria. The majority of MAGs (22 of 31)

classified within the phylum *Proteobacteria* (Table 1). Of the *Proteobacteria*, the

*Betaproteobacteria* and *Gammaproteobacteria* were the most represented, with six (MAG 29,

19, 14, 5, 20, & 27) and six (MAG 11, 23, 2, 4, 15, & 16) MAGs, respectively. The class

*Alphaproteobacteria* contained five MAGs (MAG 10, 3, 6, 25, & 25), with the

*Deltaproteobacteria* (three; MAG 7, 12, & 21) and *Epsilonproteobacteria* (two; MAG 1 and 9)

having the fewest. Two MAGs (7 and 12) were putatively identified by average nucleotide

identity (ANI) within the *Deltaproteobacterial* genera *Desulfocapsa* (Table S2). Each class

contained at least one highly complete, low redundancy MAG (Table 1).

*Non-Sulfur Associated Metabolic Potential Identified Across BFP*

Carbon fixation potential was inferred by looking at three pathways: Wood-Ljundahl

(reductive acetyl-coenzyme A); Reverse tricarboxylic acid cycle (rTCA); and ribulose-1,5-

bisphosphate carboxylase pathway (RuBisCO). ATP citrate lyase (*acl*), a key enzyme in the

rTCA cycle (Perner et al., 2007), was identified in two MAGs (1 and 9 respectively), while genes

for other rTCA enzymes were found within 17 of 31 of MAGs (Figure 3b). ATP citrate lyase

was found in both *Epsilonproteobacterial* MAGs (*Sulfurimonas* (MAG 1) and *Sulfurovum*

(MAG 9)). The most commonly identified carbon fixation pathway within the queried genomes

from BFP was the Wood–Ljungdahl pathway found in all but three MAGs (14, 22, and 31).

RuBisCo was also identified in four MAGs with both *cbbL* or *cbbM* genes present (10, 19, 23,

and 27).

Genes associated with denitrification, specifically those associated with nitrate (*nap* and

*nar*) and nitrite (*nrf* and *nir*) reduction were considered in MAG analysis as the reduction of

oxidized nitrogen species can be utilized in sulfur oxidation reactions. Genes identified as *nap*

were more abundant than those putatively identified with homology to *nar* (Figure 4). No MAGs

had both genes, and all *nap* genes were found in higher quality MAGs (1, 2, 7, and 9), with the

two instances of *nar* genes identified in lower quality MAGs (21 and 29). Nitrite reductase genes

were more abundant across MAGs (*nrf* - 1, 2, 7, 12, 21, and 22; *nir* - 1, 3, 5, 8, 20, 24, 25, and

28), however, were not especially abundant. Only two MAGs appear to have the potential to

completely reduce nitrate ($NO_3^-$) to ammonium ($NH_4^+$; MAG 1 - *nap, nrf,* and *nir*; MAG 7 - *nap*

and *nrf*).

Cytochrome *c* oxidase (*cbb3*-type) was searched as a marker for aerobic respiration

(Pitcher and Watmough, 2004), and hydrogenases *hydAB* (Takai et al., 2005) and *hoxU*

(Roalkvam et al., 2015) genes were also investigated due to their importance within

*Epsilonproteobacterial* metabolisms. Cytochrome *c* oxidase *cbb3* was found in the majority (18

of 31) MAGs. Only one instance of *hydA* was found in a contig associated with MAG 8. *HoxU*

gene abundance was found in 9 out of 31 MAGs (Figure 3b). The gene *hpnp,* a hopanoid

biosynthesis gene that codes for an enzyme that methylates hopanoids at the C2 position, is often

used as a diagnostic for biomarker potential (Garby et al., 2013). From an astrobiological

perspective identifying biomarkers such as hopanoids in extreme environments can be useful in

conjunction with potential metabolisms. No *hpnp* (hopanoid synthesis) genes were identified

within the BFP MAGs.

*Sulfur Oxidation is Abundant Across BFG Metagenome Assembled Genomes*

Of greatest interest for this study were sulfur cycling associated genes (including both the

oxidation and reduction of sulfur species). MAGs were searched for well-known genes involved

in sulfide oxidation (e.g., *fcc, sqr*), sulfite oxidation (*sorAB*), and thiosulfate oxidation (*sox*). A

gene critical to sulfide oxidation, *fcc*, was found in MAGs 1, 5, 9, 19, 20, and 27; two of these (1

and 9) correspond to nearest neighbors in the *Epilonproteobacteria* class (*Sulfurimonas* and

*Sulfurovum*). The other five *fcc*-containing MAGs correspond to organisms within the

*Betaproteobacteria* (Figure 4). Sulfide-quinone reductase (*sqr*) was only observed in four MAGs

(3, 5, 6, and 25), three of which (3, 6, and 25) are *Alphaproteobacteria* and the other most

closely related to the genus *Limnobacter* (also containing the aforementioned *fcc* gene). No

MAGs showed the presence of genes necessary for sulfite dehydrogenase (*sorAB*).

The Sox system is a widely-studied pathway for the complete oxidation of thiosulfate

($S_2O_3^{2-}$) to sulfate ($SO_4^{2-}$). This metabolism is typically broken up into four different multi-gene

proteins, which make up the larger Sox enzyme complex (*soxAX*, *soxYZ*, *soxB*, and *soxCD*). Sox

genes were identified in MAGs 1 (*soxXA*, *soxYZ*, and *soxB*) and 9 (*soxXA* and *soxYZ*), which

were identified as most closely related to the genera *Sulfurimonas* and *Sulfurovum* within the

*Epsilonproteobacteria*, respectively. Sox genes were also identified in MAGs (5, 10, 14, 15, 19,

20, 22, 23, 27, and 29), with complete operons observed in MAGs 5, 10, 15, 20, and 23 (Figure

3b). All but one of these MAGs are associated with *Proteobacterial* classes, including

*Betaproteobacteria* (5, 14, 19, 20, 27, and 29), *Alphaproteobacteria* (10), and

*Gammaproteobacteria* (15 and 23), with MAG 22 being nearest neighbors to the candidate class

*Endomicrobium* (Phylum *Elusimicrobia*). Not all MAGs had full sox gene complexes (Figure

3b) but all had more than one copy of Sox- associated genes (e.g. multiple copies of *soxA;* Figure

S3). Five MAGs (5, 10, 15, 20, and 23) have genes present for all four of the sox-related proteins

(Figure 3b).

Three other sulfur-oxidizing genes that co-associate are *apr*, *sat*, and *qmo,* which carry

out the oxidation of sulfite to sulfate (Frigaard and Dahl, 2008). In the green sulfur bacteria these

genes are seen as part of the *sat-aprBA-qmoABC* gene operon; however, a similar *aprAB-*

*qmoABC* operon is also present in sulfate-reducing bacteria (Dahl and Friedrich, 2008). The *sat-*

*aprBA-qmoABC* operon was identified in MAG 19, (genus *Thiobacillus*). MAG 7, putatively

identified as *Desulfocapsa*, contains both *apr* and *qmo* genes; however, this is not the case in

MAG 12, also identified as *Desulfocapsa*, where *apr* is absent, but *sat* is present alongside *qmo.*

Overall, only three MAGs had *qmo* genes present (MAGs 7, 12, and 19) out of the 31 queried.

In only one instance in the BFP MAGs were *apr* and *sat* found together (MAG 5), which is most

closely related to the *Betaproteobacterial* genus *Limnobacter*. Otherwise, *apr* (found in four

MAGs) was detected less often than *sat* (present in 15 MAGs), which can be seen in Figure 3b.

However, gene count totals were comparable with *apr* having 16 and *sat* having 26 instances,

respectively (Figure 5), most of which can be attributed to MAG 19, where 9 instances of *apr*

were present. The 3'-phosphoadenosine 5'-phosphosulfate (PAPS) gene, often found as an

intermediate in dissimilatory sulfide oxidation (Han and Perner, 2015), was also identified in the

majority of MAGs (18 of 31).

Dissimilatory sulfite-reductase (*dsrAB*) was also queried to identify the respiratory sulfate

reduction potential of the system. *dsrAB* was present in three MAGs (7, 12, and 19) two of which

correspond to the *Deltaproteobacterial* genus *Desulfocapsa*, while the other appears to be

nearest to the *Betaproteobacterial* genus *Thiobacillus*. All three MAGs also have the DsrMKJOP

transmembrane gene complex present which often interacts with *dsrAB*, as well as the

dissimilatory sulfite-reductase gamma subunit and some form of the assimilatory gene pathway.

Two copies of *phs* (thiosulfate reductase)*,* a gene associated with sulfur disproportionation, were

identified in MAG 2 (Figures 3 & 5).

**3      Discussion**

Arctic and Antarctic freshwater ecosystems provide insight into low-temperature

geochemical sulfur cycling and microbial community dynamics. One such low-temperature

environment in the Arctic is Borup Fiord Pass (BFP), a glacial ecosystem dominated by sulfur-

rich ice and mineral precipitates. The bioenergetic capability of microbes living in a low

temperature Arctic sediment mound at BFP was previously identified through metagenomic

sequencing (Wright et al., 2013) ; however, the work presented increases our understanding of

low-temperature microbial metabolisms by expanding the number of sites and sample types

beyond surface precipitates by including aufeis samples and fluid from an active spring in 2016.

Through the use of metagenomic sequencing we have identified microbial community members

from a variety of sample types at BFP that have an abundance of genes related to the oxidation of

reduced sulfur species via the *sox* operon including taxa known to perform this process (e.g.

*Epsilonproteobacteria – Sulfurimonas* and *Sulfurovum*), as well as others not previously known

to carry out this function in these environments (e.g. *Limnobacter* sp.). We conclude that sulfur

oxidation may be functionally redundant and more widespread phylogenetically in sulfur-rich

polar environments.

In addition to the cycling of sulfur identified at BFP we also searched for core metabolic

processes including carbon and nitrogen metabolism. Low levels of nitrate were previously

reported at BFP (0.0030 mM in 2009; Wright et al., 2013, and 0.0908 mM in 2014; Lau et al.,

2017) within both spring and melt pool fluids, respectively, possibly due to microbial nitrate

reduction. Based on gene presence/absence we infer that some MAGs from BFP may be capable

of nitrate/nitrite reduction. The *nap* gene, or periplasmic nitrate reductase, is the most abundant

identified nitrate reductase (Figure 5), with multiple copies identified in MAGs 1, 2, and 9

(*Sulfurimonas, Shewanella,* and *Sulfurovum*, respectively). This suggests that sulfur oxidizing

microorganisms (SOMs) such as *Sulfurimonas* and *Sulfurovum* found in abundance at some

sample sites (e.g., M2 and M4b, respectively; Figure 3a) are capable of oxidizing reduced sulfur

compounds and utilizing nitrate as an electron acceptor. The *nir* gene, a nitrite reductase, which

reduces nitrite to nitric oxide (NO) is slightly less abundant than *nrf*, which reduces nitrite to

ammonium ($NH_4^+$). It should be noted that only MAGs which show the presence of *nap* also

have co-occurrence of the *nir* gene, which suggests that denitrification in the system may occur

via nitric oxide and/or nitrite. While the majority organisms at BFP are most likely using oxygen

as an electron acceptor, metagenomic evidence supports the idea that anaerobic denitrification is

a viable mechanism of low-temperature respiration.

The utilization of carbon and how it plays a part in microbial growth dynamics at BFP is

important to determine, particularly how organic (or fixed) carbon might enter the system.

Organic carbon is available within the BFP system (Lau et al., 2017) and has a potential source

from nearby shale units (Grasby et al., 2003), however, previous research by Wright, et al.

(2013) indicated that the most likely form of energy production in the system is via aerobic

oxidation of $S_0$ rather than that of organic carbon. Because of this, sulfide oxidation coupled with

$CO_2$ fixation is potentially one of the most important metabolisms present at BFP. One such

phylum containing organisms capable of autotrophy and  sulfur oxidation are the

*Epsilonproteobacteria* (Campbell et al., 2006). Genes related to one such carbon fixation

pathway, the Wood-Ljungdahl pathway (reductive acetyl-coenzyme A pathway, rCoA), appear

across all but three of the MAGs (14, 22, and 31). This is in contrast to genes related to the

reductive TCA (rTCA) cycle that were present in 17 out of 31 MAGs (including *acl* which was

found in two MAGs). The rTCA cycle has previously been implicated as the primary means of

converting inorganic carbon into biomass for organisms within the *Epsilonproteobacteria* (Dahl

and Friedrich, 2008; Hügler et al., 2010).  However, the reductive acetyl-CoA cycle is the

predominant carbon fixation pathway found throughout all of our MAG data. Similarly, Momper

et al. (2017) found the reductive acetyl-CoA pathway to be the preferred pathway most likely

linked to energy limitations within deep subsurface sediments. However, while Borup Fiord Pass

(BFP) is an extreme surface environment, organic carbon data and bioenergetics calculations

suggest that BFP microbes are not at the lower thermodynamic limit of life (Wright et al., 2013;

Lau et al., 2017). Additionally, the reductive acetyl-CoA pathway requires anoxic conditions,

which is contrary to the oxygen-rich surface sampling conditions at BFP. A possible explanation

for reductive acetyl-CoA carbon fixation at BFP might be the establishment of anoxic zones just

beneath the surface of any mineral precipitate or melt pool sediment materials. Within glacial

cryoconite sediments similar to those sampled, an anoxic zone can re-establish at 2 to 4 mm from

the surface of the sediment within an hour after disturbance (Poniecka et al. 2017).

It should be noted that aerobic metabolism genes are also present in abundance across

sample types at BFP as evidenced by the *cbb3* cytochrome *c* oxidase gene. Cytochrome *c*

oxidases have been shown to operate under low oxygen tension (Koh et al., 2017) and could still

be useful in hypoxic zones near surface sediments; however, this has not yet been established for

the *cbb3* cytochrome oxidase like those identified here. Sulfur oxidation and carbon fixation may

occur under micro-oxic conditions where the reductive acetyl-CoA pathway is still effective as

the dominant carbon fixation process.

The research presented here identified sulfur-oxidizing genes in MAGs from different

sites and site types at BFP, and across multiple different taxonomic lineages. Previous research at

BFP identified (from both spring fluid and mineral deposits) a number of the same organisms

found within our identified MAGs, including: *Marinobacter*, *Loktanella* (Grasby et al., 2003;

Gleeson et al., 2011), and the sulfur oxidizers *Sulfurimonas, Sulfurovum,* and *Sulfuricurvum*

(Gleeson et al., 2011, 2012; Wright et al., 2013; Trivedi et al., 2018). Our work has revealed that

MAGs belonging to previously identified genera likely oxidize reduced sulfur compounds in

addition to organisms that were not previously predicted to be taking part in such metabolic

processes. One MAG (MAG 5), putatively identified within the genus *Limnobacter*, contains

genes associated with the oxidation of hydrogen sulfide (*fcc*, *sqr),* as well as a complete pathway

(sox) needed to oxidize hydrogen sulfide to sulfate. Chen et al. (2016) reported the full sox

pathway within a *Limnobacter* sp. in an anaerobic methane oxidizing microbial community,

while other studies have reported its ability to oxidize thiosulfate to sulfate (Kämpfer et al.,

2001; Lu et al., 2011). However, we are unaware of any previous research linking *Limnobacter*

sp. to environmental sulfide oxidation.

*Loktanella* sp. were also previously identified at BFP in 16S rRNA gene clone libraries

(Gleeson et al., 2011) as well as in other low-temperature environments throughout the Canadian

Arctic (Perreault et al., 2007; Niederberger et al., 2009; Battler et al., 2013). *Loktanella* was

previously reported as containing *soxB* from samples obtained at Gypsum Hill, a perennial Arctic

spring on Axel Heiberg Island (Perreault et al., 2008). The Perreault, et al. study also found that

the *Loktanella* sp. was part of a consortium along with a *Marinobacter* sp. similar to what was

found in previous samples at BFP (Gleeson et al., 2011; Trivedi et al., 2018).

In contrast to previous 16S rRNA gene sequencing analyses, we putatively identified the

genera *Herminiimonas* and *Rhodoferax* within the class *Betaproteobacteria* through whole

genome sequencing and binning. Despite its identification within a 120,000-year-old Greenland

ice core (Loveland-Curtze et al., 2009), *Herminiimonas* is rarely identified within polar

environments. Moreover, the data on the capability of *Herminiimonas* to oxidize sulfur is equally

sparse. In a recent study, Koh et al. (2017) found that *Herminiimonas arsenitoxidans*, which was

found to oxidize arsenite, was able to oxidize sulfur when grown on trypticase soy agar.

Conversely, *Rhodoferax* (also known as purple non-sulfur bacteria)*, another

*Betaproteobacterium* that is commonly found in polar environments (Madigan et al., 2000; Lutz

et al., 2015), are facultatively photoheterotrophic; however, some are capable of photoautotrophy

when sulfide is used as an electron donor (Ghosh and Dam, 2009). A BFP MAG putatively

identified as *Rhodoferax* (MAG 20) also shows the presence of genes related to the reductive

acetyl-CoA pathway, which could confirm that this particular organism is capable of autotrophy

and may serve as a primary producer in this sulfur-dominated polar ecosystem.

The soxCD complex has been shown to be responsible for the oxidation of sulfur to

thiosulfate, and in organisms that lack this complex the sulfur is either stored inside the cell or

excreted (Hensen et al., 2006). The lack of soxCD in BFP MAGs (e.g. MAGs 1, 9, and 19) from

known sulfur-oxidizing microorganisms (*Sulfurimonas, Sulfurovum,* and *Thiobacillus*) is an

interesting finding. In fact, this may be one possible biological explanation for the abundance of

elemental sulfur precipitated across the surface of BFP. Some organisms may employ other

pathways like reverse dissimilatory sulfite reduction (rDsr) to oxidize accumulations of $S^0$

(Hamilton et al., 2015); however, the low prevalence of any sulfite reductase genes in the BFP

system could point to an accumulation of elemental sulfur rather than additional biogenic

oxidation. Thermodynamically sulfur oxidation should occur abiotically, however, at such low

temperatures, this abiotic process may be kinetically slow without biological catalysis.

In addition to sulfur oxidation, dissimilatory sulfate reduction is also a key component in

the sulfur cycle and potentially in energy generation at BFP. Out of the 31 identified MAGs,

three (MAG 7, 12, and 19) contain dissimilatory sulfite reductase (*dsrAB*) genes, a key step in

the complete reduction of sulfate to sulfide. Two MAGs, MAG 7 and 12, are most closely related

to the genus *Desulfocapsa*, a known sulfur disproportionator within the *Deltaproteobacteria*

(Finster et al., 1998, 2013). This genus was previously reported at BFP via 16S rRNA gene

sequencing (Trivedi et al., 2018). It should also be noted that while genes specific to

disproportionation (*phs*) were searched for, only one instance was found, within MAG 2, which

is most closely related to the genus *Shewanella*. Another MAG most closely related to the

*Betaproteobacterial* genus *Thiobacillus* (MAG 19) also contains a complete *dsr* operon.

Interestingly, MAG 19 is the only MAG that appears capable of both sulfur-oxidation via the sox

pathway and sulfate-reduction via dissimilatory sulfite reductase. *Thiobacillus* sp. are

traditionally known as sulfur oxidizing microorganisms (SOMs) and have been found in polar

environments previously (Boyd et al., 2014; Purcell et al., 2014; Harrold et al., 2016); however,

it has also been suggested that certain *Thiobacillus* species carry out reverse dissimilatory sulfite

reduction (rDsr, or reverse dsr) as an alternative means to oxidize reduced sulfur compounds

(Loy et al., 2009; Purcell et al., 2014). This could well be an alternative pathway that is present

in the BFP system; however, it would require transcriptomic evidence to confirm its activity. It

should be noted that MAG 19 appears to lack soxCD; however, the MAG is only ~73%

complete, which may indicate that its full metabolic potential is not represented.

The abundance of sox genes present in BFP across sample types and multiple bacterial

lineages suggests that sulfur cycling is functionally redundant in the BFP ecosystem.

Microorganisms in cold, deep subsea floor aquifers have also been identified with high degrees

of functional redundancy (Tully et al., 2018). Functional redundancy was recently identified as a

commonplace process in environmental systems by Louca et al. (2018). The authors speculated

that the degree of functional redundancy in an ecosystem is largely determined by the type of

environment, in contrast to the belief that species should inhabit distinct microbial niches

independent of environment. Moreover, Louca et al. call into question a previous idea that

functional redundancy in an ecosystem might infer a form of 'neutral co-existence' between

competing microorganisms (Loreau, 2004). However, their findings challenge this, as coexisting

microorganisms will often differ in ways that will affect how they grow under environmental

conditions. Based on these ideas it is highly probable that functional redundancy may be an

important survival mechanism in use by microorganisms at BFP.

At BFP a core microbial community not predicted to participate in sulfur cycling exists as

identified by a 16S rRNA gene sequencing (Trivedi et al., 2018). Instead our metagenomic data

indicate that multiple phylogenetically disparate core communities may have the potential to

utilize sulfur compounds for growth. As suggested by Louca et. al (2018), coexisting organisms

may not be in equal abundance (such as those identified within the core community) as there will

always be differences in functional ability based on enzyme efficiencies and growth kinetics.

This also supports previous research of the BFP system, where "blooms" of SOMs over short

time periods occurred, followed by a return to a basal microbial community (Trivedi et al.,

2018). Such blooms of SOMs could be facilitated from above by the input of new microbiota via

aeolian transport in meteorologic storm events whereby snow storms contain significant amounts

of microbiota and genetic potential (Honeyman et al., 2018). If there is a large influx of reduced

sulfur compounds from beneath the ice, some phyla, like the occasionally (e.g. conditionally)

rare *Epsilonproteobacteria* (Trivedi et al., 2018), may be preferentially able to use these

compounds, much like in deep sea vent fluids (Akerman et al., 2013) leading to an increase in

their relative abundance until settling to previously identified levels. Undoubtedly, sulfur

oxidation is the predominant and preferred metabolism in these sulfur-rich surface sites at BFP.

The response to ecological change by conditionally rare taxa (i.e., the low representation

of microbiota in any environment that can become dominant upon environmental condition

change) thus reflects a biological manifestation of functional redundancy. The presence of

functional redundancy found at BFP is due to the multiple environmental pressures at the site

such as extreme cold, and high concentrations of multiple sulfur species (e.g., $H_2S$, $HS^-$, $S^0$,

$S_2O_3^{2-}$, and $SO_4^{2-}$) geochemically produced in the subsurface (Wright et al., 2013). These

compounds then emanate to the surface where microorganisms then take advantage of the

abundance of reduced sulfur for growth. This may be driven by the unique convergence of lower

levels of organic carbon, low temperatures, and an abundance of reduced sulfur compounds at

BFP, which may not necessarily be the case in all highly sulfidic systems.

These results show that in low-temperature environments, functional redundancy of key

metabolisms is an active strategy for multiple bacterial lineages to coexist. The results presented

here reflect the potential metabolisms present at BFP generated from whole genome sequencing

and the production of MAGs. The next step in truly defining how microorganisms are able to

survive and thrive in the BFP system will be to undertake metatranscriptomic sequencing. This

next step in the genomic characterization of this ecosystem will allow us to differentiate between

the metabolic potential and active processes within the system. Our data show that sulfur

oxidation is functionally redundant at BFP and may be utilized by more microorganisms across

the bacterial domain than would be expected.

**4      Materials and Methods**

*Sample Collection*

BFP sample material was collected during multiple days between 21 June and 2 July

2014, 4 July 2016, and 7 July 2017. Samples included: 1) sedimented sulfur-like cryoconite

material and fluid found in glacial surface melt pools (labelled M2 & M4b); 2) filtered fluid from

thawed aufeis (labelled A4b & A6); 3) aufeis surface precipitate samples (material scraped from

on top of the aufeis) from 2017 (labelled 14C, 3B, 4E, & 6B); and 4) spring fluid from 2016

(labelled BFP16 Spring); see Table S1 and Figures 3a & S2). Cryoconite sediment, aufeis, and

spring fluid samples for DNA extraction were filtered through 0.22 µm Luer-lok Sterivex™

filters (EMD Millipore; Darmstadt, Germany), which were then capped and kept on ice in the

field until returning to the laboratory where they were stored at -20 °C until extraction. Aufeis

surface scrapings (mineral precipitate samples) from the 2017 field campaign were collected

using sterile and field-washed transfer pipettes and up to 0.5 g of material were mixed in ZR

BashingBead™ lysis tubes containing 750 ml of DNA/RNA Shield™ (Zymo Research Corp.).

Samples were shaken and kept at 4 °C until returned to the laboratory, where they were then

stored at -80 °C until DNA/RNA extraction could be performed.

*DNA Extraction and Metagenomic Library Preparation*

DNA extraction of 2014 melt pool , aufeis, and 2016 spring fluid samples was performed

using the PowerWater® Sterivex™ DNA Isolation Kit (MO BIO Laboratories, USA), and DNA

extraction of 2017 surface precipitate samples was extracted using the ZymoBIOMICS™

DNA/RNA Mini Kit (Zymo Research Corp.). All extractions were performed according to the

manufacturer's instructions. Extracted DNA concentrations were determined using a QuBit 2.0

fluorometer (Thermo Fisher Scientific, Chino, CA, USA) and the QuBit dsDNA HS assay (Life

Technologies, Carlsbad, CA, USA).

Genomic DNA was first cleaned using KAPA Pure Beads (KAPA Biosystems Inc.,

Wilmington, MA, USA) at a final concentration of 0.8X v/v prior to library preparation. Two

metagenomic sequencing runs were completed for the included sample set. The first (including

samples BFP14 A4b, A6, and M4b) were normalized to 1 ng of DNA in 35 µl of molecular

biology grade water, and prepared using the KAPA HyperPlus Kit (KAPA Biosystems Inc.,

Wilmington, MA, USA) and KAPA WGS adapters. Fragmentation and adapter ligation was

performed according to the manufacturer's instructions. Prepped samples were quality checked

on an Agilent 2100 Bioanalyzer (Agilent Genomics, Santa Clara, CA, USA) and submitted to the

Duke Center for Genomic and Computational Biology (Duke University, Durham, NC, USA).

Final pooled libraries were run on an Illumina HiSeq 4000 (Illumina, San Diego, CA, USA)

using PE150 chemistry. The second sequencing run (which included samples BFP14 M2, BFP16

Spring, and BFP17 3B, 4E, 6B, and 14C) was prepped using the Nextera XT DNA Library

Preparation Kit (Illumina, San Diego, CA, USA) in conjunction with Nextera XT V2 Indexes.

Libraries were hand normalized and quality checked on an Agilent 2100 Bioanalyzer (Agilent

Genomics, Santa Clara, CA, USA). Samples were submitted to the Duke Center for Genomic

and Computational Biology (Duke University, Durham, NC, USA). These libraries were

sequenced on an Illumina HiSeq 2500 Rapid Run using PE250 chemistry.

*Metagenomic Sequencing Assembly and Analysis*

Processing of metagenomic sequencing was performed on the Summit High Performance

Cluster (HPC) located at the University of Colorado, Boulder (Boulder, CO, USA). Assembly of

whole-genome sequencing reads used a modified Joint Genome Institute (JGI; Walnut Creek,

CA, USA) metagenomics workflow. We began by using BBDuk of the BBMap/BBTools

(v37.56; Bushnell, 2014) suite to trim out Illumina adapters using the built-in BBMap adapters

database. Trimmed sequences were checked for quality using FastQC v11.5 (Andrews, 2014) to

ensure that adapters were removed successfully. Trimmed paired-end sequences were then joined

using the PEAR package (Zhang et al., 2014) v0.9.10 using a p-value of 0.001. Adapter trimmed

files, both forward (R1) and reverse (R2) were then processed using BBTools 'repair.sh' to

ensure read pairs were in the correct order. This script is designed to reorder paired reads that

have become disorganized, which can often occur during trimming. The BFC (Bloom Filter

Correction) software tool was then used to error correct paired sequence reads (Li, 2015). Reads

were again processed with BBTools 'repair.sh' to ensure proper read pairing and order. Next,

Megahit v1.1.2 (Li et al., 2015) was used to generate both a co-assembly, as well as individual

sample assemblies from the quality controlled reads. Post assembly, the co-assembly was

processed with 'anvi-script-reformat-fasta' for input into Anvi'o, and contigs below 2500 bp

were removed. Bowtie2 v2.3.0 (Langmead and Salzberg, 2012) was then used to map individual

sample sequence reads onto the co-assembly, needed for the Anvi'o pipeline. Anvi'o v4 (Eren et

al., 2015) was then used to characterize, and then bin co-assembled contigs into metagenome

assembled genomes (MAGs). The taxonomic classifier Centrifuge v1.0.2-beta (Kim et al., 2016)

was used to provide a preliminary taxonomy of the co-assembly contigs, useful for Anvi'o

binning. The NCBI Clusters of Orthologous Groups (COGs; (Tatusov et al., 2000; Galperin et

al., 2015)) database was used within Anvi'o to predict COGs present within the co-assembly,

and CONCOCT v1.0 (Alneberg et al., 2014) was employed to create genome bins using a

combination of sequence coverage and composition of our assembled contigs. Anvi'o was then

used to visualize the putative bins in conjunction with other metadata to refine and identify

MAGs with 50% completeness and 10% redundancy. The scripts used in the processing of these

data (up to the point of Megahit-generated contigs) are publicly available at

https://doi.org/10.5281/zenodo.1302787.

Anvi'o refined bins were then processed using CheckM v1.0.11 (Parks et al., 2015)

where a total of 31 bins were selected as MAGs (Metagenome Assembled Genomes) that

followed the > 50% completeness and < 10% redundancy criteria. MAGs that met these criteria

were then renamed; from 1-31 in the order of highest completeness to lowest (Table 1). CheckM

relies on other software including pplacer v1.1.alpha17 (Matsen et al., 2010) for phylogenetic

tree placement, prodigal v2.6.3 (Hyatt et al., 2012) for gene translation and initiation site

prediction, and HMMER v3.1b2 (Eddy, 2011) which analyzes sequence data using hidden

Markov models. The pplacer generated tree was then visualized using the interactive Tree of Life

(iTOL; (Letunic and Bork, 2016)) to search for nearest neighbors of identified MAGs (see Figure

4). Finally, to confirm general taxonomic classification and annotate genes within our MAGs,

samples were processed using the Rapid Annotations using Subsystem Technologies (RAST)

server (Aziz et al., 2008). Genes of interest to this study were identified within the RAST web

interface. Further taxonomic classification was done for MAGs 7 and 12 using the online tool

JSpeciesWS (Richter et al., 2016), which uses pairwise genome comparisons to measure the

probability of two or more genomes. Tests within JSpeciesWS include average nucleotide

identity (ANI) via BLAST and MUMmer and correlation between Tetra-nucleotide signatures.

Average nucleotide identity (ANI) was used to determine if MAGs 7 and 12 (putatively

classified as *Desulfotalea*) were more closely related to the genus *Desulfocapsa* as was

previously identified within BFP samples (Trivedi et al., 2018).

*Data Accessibility*

Raw metagenomic sequencing data are available at the NCBI sequencing read archive

(SRA) under the accession numbers SRR10066347-SRR10066355

(https://www.ncbi.nlm.nih.gov/sra). Assembled MAGs are provided via FigShare under the

following DOI: 10.6084/m9.figshare.9767564. Additionally, the bioinformatics workflow up to

the point of assembled contigs can be found under DOI: https://doi.org/10.5281/zenodo.1302787.

**5     Conflict of Interest**

The authors declare that the research was conducted in the absence of any commercial or

financial relationships that could be construed as a potential conflict of interest.

**6     Author Contributions**

AST, JRS, and SEG led the design of the study. Fieldwork was conducted by CBT, GEL, SEG,

AST, and JRS. Laboratory work was performed by CBT and GEL, and bioinformatic analysis

was done by CBT and BWS. All authors interpreted results. CBT is the primary author of the

manuscript with contributions and guidance from all other authors.

**7      Funding**

This research was funded by a grant (NNX13AJ32G) from the NASA Exobiology and

Evolutionary Biology Program to AST and JRS. This work was also supported by NASA

Astrobiology Institute Rock Powered Life Grant NNA15BB02A (CBT, AST, and JRS). BWS

was supported under the Sloan Foundation grant G-2017-9853. CBT is currently supported under

the Helmholtz Recruiting Initiative (award number I-044-16-01) to Liane G. Benning.

**8      Acknowledgements**

Field logistics and travel to/from Borup Fiord Pass were supported through Natural Resources

Canada's (NRCan) Polar Continental Shelf Program (PCSP). JRS and SEG thank Dr. Karsten

Piepjohn and the Federal Institute for Geosciences and Natural Resources at BGR, Hanover,

Germany for support during the 2017 field campaign of the Circum-Arctic Structural Events-19

(CASE-19) expedition.

**9      References**

Akerman, N. H., Butterfield, D. A., and Huber, J. A. (2013). Phylogenetic diversity and
functional gene patterns of sulfur-oxidizing subseafloor Epsilonproteobacteria in diffuse
hydrothermal vent fluids. *Front. Microbiol.* 4, 1–14. doi:10.3389/fmicb.2013.00185.

Alneberg, J., Bjarnason, B. S., de Bruijn, I., Schirmer, M., Quick, J., Ijaz, U. Z., et al. (2014).

Binning metagenomic contigs by coverage and composition. *Nat. Methods* 11, 1144–1146.
doi:10.1038/nmeth.3103.

Andrews, S. (2014). FastQC software package. Available at:
https://www.bioinformatics.babraham.ac.uk/projects/fastqc/ [Accessed March 13, 2018].

Andriuzzi, W. S., Stanish, L. F., Simmons, B. L., Jaros, C., Adams, B. J., Wall, D. H., et al.
(2018). Spatial and temporal patterns of microbial mats and associated invertebrates along
an Antarctic stream. *Polar Biol.*, 1–11. doi:10.1007/s00300-018-2331-4.

Aziz, R. K., Bartels, D., Best, A. A., DeJongh, M., Disz, T., Edwards, R. A., et al. (2008). The
RAST Server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9, 75.
doi:10.1186/1471-2164-9-75.

Battler, M. M., Osinski, G. R., and Banerjee, N. R. (2013). Mineralogy of saline perennial cold
springs on Axel Heiberg Island, Nunavut, Canada and implications for spring deposits on
Mars. *Icarus* 224, 364–381. doi:10.1016/j.icarus.2012.08.031.

Boetius, A., Anesio, A. M., Deming, J. W., Mikucki, J. A., and Rapp, J. Z. (2015). Microbial
ecology of the cryosphere: sea ice and glacial habitats. *Nat. Rev. Microbiol.* 13, 677–690.
doi:10.1038/nrmicro3522.

Bowles, M. W., Mogollón, J. M., Kasten, S., Zabel, M., and Hinrichs, K.-U. (2014). Global rates
of marine sulfate reduction and implications for sub-sea-floor metabolic activities. *Science*
344, 889–91. doi:10.1126/science.1249213.

Boyd, E. S., Hamilton, T. L., Havig, J. R., Skidmore, M. L., and Shock, E. L. (2014).
Chemolithotrophic primary production in a subglacial ecosystem. *Appl. Environ. Microbiol.*
80, 6146–6153. doi:10.1128/AEM.01956-14.

Bushnell, B. (2014). BBMap software package. Available at:
https://sourceforge.net/projects/bbmap/ [Accessed March 13, 2018].

Campbell, B. J., Engel, A. S., Porter, M. L., and Takai, K. (2006). The versatile ε-proteobacteria:
Key players in sulphidic habitats. *Nat. Rev. Microbiol.* 4, 458–468.
doi:10.1038/nrmicro1414.

Carlson, R. W., Johnson, R. E., and Anderson, M. S. (1999). Sulfuric acid on Europa and the
radiolytic sulfur cycle. *Science* 286, 97–99. doi:10.1126/science.286.5437.97.

Carr, S. A., Orcutt, B. N., Mandernack, K. W., and Spear, J. R. (2015). Abundant Atribacteria in
deep marine sediment from the Adélie Basin, Antarctica. *Front. Microbiol.* 6, 1–12.
doi:10.3389/fmicb.2015.00872.

Chen, Y., Feng, X., He, Y., and Wang, F. (2016). Genome Analysis of a Limnobacter sp.
Identified in an Anaerobic Methane-Consuming Cell Consortium. *Front. Mar. Sci.* 3, 257.
doi:10.3389/fmars.2016.00257.

Dahl, C., and Friedrich, C. G. (2008). *Microbial Sulfur Metabolism.* , eds. C. Dahl and C. G.
Friedrich Berlin, Heidelberg: Springer Berlin Heidelberg doi:10.1007/978-3-540-72682-1.

Eddy, S. R. (2011). Accelerated Profile HMM Searches. *PLoS Comput. Biol.* 7, e1002195.

doi:10.1371/journal.pcbi.1002195.

Enning, D., Venzlaff, H., Garrelfs, J., Dinh, H. T., Meyer, V., Mayrhofer, K., et al. (2012).
Marine sulfate-reducing bacteria cause serious corrosion of iron under electroconductive
biogenic mineral crust. *Environ. Microbiol.* 14, 1772–1787. doi:10.1111/j.1462-
2920.2012.02778.x.

Eren, A. M., Esen, Ö. C., Quince, C., Vineis, J. H., Morrison, H. G., Sogin, M. L., et al. (2015).
Anvi'o: an advanced analysis and visualization platform for 'omics data. *PeerJ* 3, e1319.
doi:10.7717/peerj.1319.

Feazel, L. M., Spear, J. R., Berger, A. B., Harris, J. K., Frank, D. N., Ley, R. E., et al. (2008).
Eucaryotic diversity in a hypersaline microbial mat. *Appl. Environ. Microbiol.* 74, 329–32.
doi:10.1128/AEM.01448-07.

Finster, K., Liesack, W., and Thamdrup, B. (1998). Elemental sulfur and thiosulfate
disproportionation by Desulfocapsa sulfoexigens sp. nov., a new anaerobic bacterium
isolated from marine surface sediment. *Appl. Environ. Microbiol.* 64, 119–125.

Finster, K. W., Kjeldsen, K. U., Kube, M., Reinhardt, R., Mussmann, M., Amann, R., et al.
(2013). Complete genome sequence of Desulfocapsa sulfexigens, a marine
deltaproteobacterium specialized in disproportionating inorganic sulfur compounds. *Stand.*
*Genomic Sci.* 8, 58–68. doi:10.4056/sigs.3777412.

Friedrich, C. G., Bardischewsky, F., Rother, D., Quentmeier, A., and Fischer, J. (2005).
Prokaryotic sulfur oxidation. *Curr. Opin. Microbiol.* 8, 253–259.
doi:10.1016/j.mib.2005.04.005.

Friedrich, C. G., Rother, D., Bardischewsky, F., Quentmeier, A., and Fischer, J. (2001).
Oxidation of reduced inorganic sulfur compounds by bacteria: emergence of a common
mechanism? *Appl. Environ. Microbiol.* 67, 2873–82. doi:10.1128/AEM.67.7.2873-
2882.2001.

Frigaard, N. U., and Dahl, C. (2008). *Sulfur Metabolism in Phototrophic Sulfur Bacteria*.
doi:10.1016/S0065-2911(08)00002-7.

Galperin, M. Y., Makarova, K. S., Wolf, Y. I., and Koonin, E. V. (2015). Expanded microbial
genome coverage and improved protein family annotation in the COG database. *Nucleic*
*Acids Res.* 43, D261–D269. doi:10.1093/nar/gku1223.

Garby, T. J., Walter, M. R., Larkum, A. W. D., and Neilan, B. A. (2013). Diversity of
cyanobacterial biomarker genes from the stromatolites of Shark Bay, Western Australia.
*Environ. Microbiol.* 15, 1464–1475. doi:10.1111/j.1462-2920.2012.02809.x.

Ghosh, W., and Dam, B. (2009). Biochemistry and molecular biology of lithotrophic sulfur
oxidation by taxonomically and ecologically diverse bacteria and archaea. *FEMS Microbiol.*
*Rev.* 33, 999–1043. doi:10.1111/j.1574-6976.2009.00187.x.

Gleeson, D. F., Pappalardo, R. T., Anderson, M. S., Grasby, S. E., Mielke, R. E., Wright, K. E.,
et al. (2012). Biosignature detection at an Arctic analog to Europa. *Astrobiology* 12, 135–
150. doi:10.1089/ast.2010.0579.

Gleeson, D. F., Pappalardo, R. T., Grasby, S. E., Anderson, M. S., Beauchamp, B., Castaño, R.,
et al. (2010). Characterization of a sulfur-rich Arctic spring site and field analog to Europa
using hyperspectral data. *Remote Sens. Environ.* 114, 1297–1311.
doi:10.1016/j.rse.2010.01.011.

Gleeson, D. F., Williamson, C., Grasby, S. E., Pappalardo, R. T., Spear, J. R., and Templeton, A.
S. (2011). Low temperature S0 biomineralization at a supraglacial spring system in the
Canadian high Arctic. *Geobiology* 9, 360–375. doi:10.1111/j.1472-4669.2011.00283.x.

Grasby, S. E. (2003). Naturally precipitating vaterite (μ-CaCO3) spheres: Unusual carbonates
formed in an extreme environment. *Geochim. Cosmochim. Acta* 67, 1659–1666.
doi:10.1016/S0016-7037(00)01304-2.

Grasby, S. E., Allen, C. C., Longazo, T. G., Lisle, J. T., Griffin, D. W., and Beauchamp, B.
(2003). Supraglacial sulfur springs and associated biological activity in the Canadian high
Arctic—signs of life beneath the ice. *Astrobiology* 3, 583–596.
doi:10.1089/153110703322610672.

Grasby, S. E., Beauchamp, B., and Bense, V. (2012). Sulfuric acid speleogenesis associated with
a glacially driven groundwater system - Paleo-spring "pipes" at Borup Fiord Pass, Nunavut.
*Astrobiology* 12, 19–28. doi:10.1089/ast.2011.0700.

Grasby, S. E., Proemse, B. C., and Beauchamp, B. (2014). Deep groundwater circulation through
the High Arctic cryosphere forms Mars-like gullies. *Geology* 42, 651–654.
doi:10.1130/G35599.1.

Hamilton, T. L., Jones, D. S., Schaperdoth, I., and Macalady, J. L. (2015). Metagenomic insights
into S(0) precipitation in a terrestrial subsurface lithoautotrophic ecosystem. *Front.
Microbiol.* 5, 756. doi:10.3389/fmicb.2014.00756.

Han, Y., and Perner, M. (2015). The globally widespread genus Sulfurimonas: versatile energy
metabolisms and adaptations to redox clines. *Front. Microbiol.* 6, 989.
doi:10.3389/fmicb.2015.00989.

Harrold, Z. R., Skidmore, M. L., Hamilton, T. L., Desch, L., Amada, K., Van Gelder, W., et al.
(2016). Aerobic and anaerobic thiosulfate oxidation by a cold-adapted, subglacial
chemoautotroph. *Appl. Environ. Microbiol.* 82, 1486–1495. doi:10.1128/AEM.03398-15.

Hensen, D., Sperling, D., Trüper, H. G., Brune, D. C., and Dahl, C. (2006). Thiosulphate
oxidation in the phototrophic sulphur bacterium Allochromatium vinosum. *Mol. Microbiol.*
62, 794–810. doi:10.1111/j.1365-2958.2006.05408.x.

Hodgkins, R., Tranter, M., and Dowdeswell, J. A. (2004). The characteristics and formation of a
high-arctic proglacial icing. *Geogr. Ann. Ser. A, Phys. Geogr.* 86, 265–275.
doi:10.1111/j.0435-3676.2004.00230.x.

Honeyman, A. S., Day, M. L., and Spear, J. R. (2018). Fresh snowfall microbiology and
chemistry are driven by geography in storm-tracked events. *bioRxiv*.
doi:https://doi.org/10.1101/300772.

Hügler, M., Gärtner, A., and Imhoff, J. F. (2010). Functional genes as markers for sulfur cycling
and CO2 fixation in microbial communities of hydrothermal vents of the Logatchev field.

*FEMS Microbiol. Ecol.* 73, no-no. doi:10.1111/j.1574-6941.2010.00919.x.

Hyatt, D., LoCascio, P. F., Hauser, L. J., and Uberbacher, E. C. (2012). Gene and translation
initiation site prediction in metagenomic sequences. *Bioinformatics* 28, 2223–2230.
doi:10.1093/bioinformatics/bts429.

Jungblut, A. D., Hawes, I., Mackey, T. J., Krusor, M., Doran, P. T., Sumner, D. Y., et al. (2016).
Microbial Mat Communities along an Oxygen Gradient in a Perennially Ice-Covered
Antarctic Lake. *Appl. Environ. Microbiol.* 82, 620–30. doi:10.1128/AEM.02699-15.

Kämpfer, P., Spring, S., and Schleifer, K. H. (2001). Limnobacter thiooxidans gen. nov., sp.
nov., a novel thiosulfate-oxidizing bacterium isolated from freshwater lake sediment. *Int. J.*
*Syst. Evol. Microbiol.* 51, 1463–1470. doi:10.1099/00207713-51-4-1463.

Kim, D., Song, L., Breitwieser, F. P., and Salzberg, S. L. (2016). Centrifuge: rapid and sensitive
classification of metagenomic sequences. *Genome Res.* 26, 1721–1729.
doi:10.1101/gr.210641.116.

Koh, H.-W., Hur, M., Kang, M.-S., Ku, Y.-B., Ghai, R., and Park, S.-J. (2017). Physiological
and genomic insights into the lifestyle of arsenite-oxidizing Herminiimonas arsenitoxidans.
*Sci. Rep.* 7, 15007. doi:10.1038/s41598-017-15164-4.

Lamarche-Gagnon, G., Comery, R., Greer, C. W., and Whyte, L. G. (2015). Evidence of in situ
microbial activity and sulphidogenesis in perennially sub-0 °C and hypersaline sediments of
a high Arctic permafrost spring. *Extremophiles* 19, 1–15. doi:10.1007/s00792-014-0703-4.

Langmead, B., and Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nat.*
*Methods* 9, 357–359. doi:10.1038/nmeth.1923.

Lau, G. E., Cosmidis, J., Grasby, S. E., Trivedi, C. B., Spear, J. R., and Templeton, A. S. (2017).
Low-temperature formation and stabilization of rare allotropes of cyclooctasulfur (β-S 8 and
γ-S 8 ) in the presence of organic carbon at a sulfur-rich glacial site in the Canadian High
Arctic. *Geochim. Cosmochim. Acta* 200, 218–231. doi:10.1016/j.gca.2016.11.036.

Letunic, I., and Bork, P. (2016). Interactive tree of life (iTOL) v3: an online tool for the display
and annotation of phylogenetic and other trees. *Nucleic Acids Res.* 44, W242–W245.
doi:10.1093/nar/gkw290.

Ley, R. E., Harris, J. K., Wilcox, J., Spear, J. R., Miller, S. R., Bebout, B. M., et al. (2006).
Unexpected diversity and complexity of the Guerrero Negro hypersaline microbial mat.
*Appl. Environ. Microbiol.* 72, 3685–95. doi:10.1128/AEM.72.5.3685-3695.2006.

Li, D., Liu, C. M., Luo, R., Sadakane, K., and Lam, T. W. (2015). MEGAHIT: An ultra-fast
single-node solution for large and complex metagenomics assembly via succinct de Bruijn
graph. *Bioinformatics* 31, 1674–1676. doi:10.1093/bioinformatics/btv033.

Li, H. (2015). BFC: Correcting Illumina sequencing errors. *Bioinformatics* 31, 2885–2887.
doi:10.1093/bioinformatics/btv290.

Loreau, M. (2004). Does functional redundancy exist? *Oikos* 104, 606–611. doi:10.1111/j.0030-
1299.2004.12685.x.

Louca, S., Polz, M., Mazel, F., Albright, M., Julie, H., O'Connor, M., et al. (2018). Function and
functional redundancy in microbial systems. *Nat. Ecol. Evol.* Accepted.
doi:10.1038/s41559-018-0519-1.

Loveland-Curtze, J., Miteva, V. I., and Brenchley, J. E. (2009). Herminiimonas glaciei sp. nov., a
novel ultramicrobacterium from 3042 m deep Greenland glacial ice. *Int. J. Syst. Evol.*
*Microbiol.* 59, 1272–1277. doi:10.1099/ijs.0.001685-0.

Loy, A., Duller, S., Baranyi, C., Mussmann, M., Ott, J., Sharon, I., et al. (2009). Reverse
dissimilatory sulfite reductase as phylogenetic marker for a subgroup of sulfur-oxidizing
prokaryotes. *Environ. Microbiol.* 11, 289–99. doi:10.1111/j.1462-2920.2008.01760.x.

Lu, H., Sato, Y., Fujimura, R., Nishizawa, T., Kamijo, T., and Ohta, H. (2011). Limnobacter
litoralis sp. nov., a thiosulfate-oxidizing, heterotrophic bacterium isolated from a volcanic
deposit, and emended description of the genus Limnobacter. *Int. J. Syst. Evol. Microbiol.*
61, 404–407. doi:10.1099/ijs.0.020206-0.

Lutz, S., Anesio, A. M., Field, K., and Benning, L. G. (2015). Integrated "Omics", Targeted
Metabolite and Single-cell Analyses of Arctic Snow Algae Functionality and Adaptability.
*Front. Microbiol.* 6, 1323. doi:10.3389/fmicb.2015.01323.

Madigan, M. T., Jung, D. O., Woese, C. R., and Achenbach, L. A. (2000). Rhodoferax
antarcticus sp. nov., a moderately psychrophilic purple nonsulfur bacterium isolated from an
Antarctic microbial mat. *Arch. Microbiol.* 173, 269–77. Available at:
http://www.ncbi.nlm.nih.gov/pubmed/10816045 [Accessed April 21, 2018].

Massé, M., Bourgeois, O., Le Mouélic, S., Verpoorter, C., Spiga, A., and Le Deit, L. (2012).
Wide distribution and glacial origin of polar gypsum on Mars. *Earth Planet. Sci. Lett.* 317–
318, 44–55. doi:10.1016/j.epsl.2011.11.035.

Matsen, F. A., Kodner, R. B., and Armbrust, E. V. (2010). pplacer: linear time maximum-
likelihood and Bayesian phylogenetic placement of sequences onto a fixed reference tree.
*BMC Bioinformatics* 11, 538. doi:10.1186/1471-2105-11-538.

Mattes, T. E., Nunn, B. L., Marshall, K. T., Proskurowski, G., Kelley, D. S., Kawka, O. E., et al.
(2013). Sulfur oxidizers dominate carbon fixation at a biogeochemical hot spot in the dark
ocean. *ISME J.* 7113, 2349–2360. doi:10.1038/ismej.2013.113.

Mikucki, J. A., Lee, P. A., Ghosh, D., Purcell, A. M., Mitchell, A. C., Mankoff, K. D., et al.
(2016). Subglacial Lake Whillans microbial biogeochemistry: A synthesis of current
knowledge. *Philos. Trans. R. Soc. A Math. Phys. Eng. Sci.* 374, 20140290.
doi:10.1098/rsta.2014.0290.

Mikucki, J. A., Pearson, A., Johnston, D. T., Turchyn, A. V, Farquhar, J., Schrag, D. P., et al.
(2009). A Contemporary Microbially Maintained Subglacial Ferrous "Ocean." *Science (80-.*
*).* 324, 397–400. doi:10.1126/science.1167350.

Mikucki, J. A., and Priscu, J. C. (2007). Bacterial diversity associated with blood falls, a
subglacial outflow from the Taylor Glacier, Antarctica. *Appl. Environ. Microbiol.* 73, 4029–
4039. doi:10.1128/AEM.01396-06.

Momper, L., Jungbluth, S. P., Lee, M. D., and Amend, J. P. (2017). Energy and carbon

metabolisms in a deep terrestrial subsurface fluid microbial community. *ISME J.* 11, 2319–
2333. doi:10.1038/ismej.2017.94.

Niederberger, T. D., Perreault, N. N., Lawrence, J. R., Nadeau, J. L., Mielke, R. E., Greer, C. W.,
et al. (2009). Novel sulfur-oxidizing streamers thriving in perennial cold saline springs of
the Canadian high Arctic. *Environ. Microbiol.* 11, 616–629. doi:10.1111/j.1462-
2920.2008.01833.x.

Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P., and Tyson, G. W. (2015).
CheckM: assessing the quality of microbial genomes recovered from isolates, single cells,
and metagenomes. *Genome Res.* 25, 1043–55. doi:10.1101/gr.186072.114.

Perner, M., Seifert, R., Weber, S., Koschinsky, A., Schmidt, K., Strauss, H., et al. (2007).
Microbial CO2 fixation and sulfur cycling associated with low-temperature emissions at the
Lilliput hydrothermal field, southern Mid-Atlantic Ridge (9°S). *Environ. Microbiol.* 9,
1186–1201. doi:10.1111/j.1462-2920.2007.01241.x.

Perreault, N. N., Andersen, D. T., Pollard, W. H., Greer, C. W., and Whyte, L. G. (2007).
Characterization of the prokaryotic diversity in cold saline perennial springs of the
Canadian high Arctic. *Appl. Environ. Microbiol.* 73, 1532–1543. doi:10.1128/AEM.01729-
06.

Perreault, N. N., Greer, C. W., Andersen, D. T., Tille, S., Lacrampe-Couloume, G., Lollar, B. S.,
et al. (2008). Heterotrophic and autotrophic microbial populations in cold perennial springs
of the high arctic. *Appl. Environ. Microbiol.* 74, 6898–6907. doi:10.1128/AEM.00359-08.

Pitcher, R. S., and Watmough, N. J. (2004). The bacterial cytochrome cbb3 oxidases. *Biochim.
Biophys. Acta - Bioenerg.* 1655, 388–399. doi:10.1016/j.bbabio.2003.09.017.

Poniecka, E. A., Bagshaw, E. A., Tranter, M., Sass, H., Williamson, C. J., Anesio, A. M., et al.
(2017). Rapid development of anoxic niches in supraglacial ecosystems. *Arctic, Antarct.
Alp. Res.* 50, 14. doi:10.1080/15230430.2017.1420859.

Purcell, A. M., Mikucki, J., Alekhina, I., Achberger, A. M., Barbante, C., Brent, C., et al. (2014).
Microbial sulfur transformations in sediments from Subglacial Lake Whillans. *Front.
Microbiol.* 5, 1–28. doi:10.3389/fmicb.2014.00594.

Richter, M., Rosselló-Móra, R., Oliver Glöckner, F., and Peplies, J. (2016). JSpeciesWS: a web
server for prokaryotic species circumscription based on pairwise genome comparison.
*Bioinformatics* 32, 929–931. doi:10.1093/bioinformatics/btv681.

Roalkvam, I., Drønen, K., Stokke, R., Daae, F. L., Dahle, H., and Steen, I. H. (2015).
Physiological and genomic characterization of Arcobacter anaerophilus IR-1 reveals new
metabolic features in Epsilonproteobacteria. *Front. Microbiol.* 6, 987.
doi:10.3389/fmicb.2015.00987.

Robertson, C. E., Spear, J. R., Harris, J. K., and Pace, N. R. (2009). Diversity and stratification
of archaea in a hypersaline microbial mat. *Appl. Environ. Microbiol.* 75, 1801–10.
doi:10.1128/AEM.01811-08.

Spear, J. R., Figueroa, L. A., and Honeyman, B. D. (1999). Modeling the Removal of Uranium
U(VI) from Aqueous Solutions in the Presence of Sulfate Reducing Bacteria. *Environ. Sci.*

785            *Technol.* 33, 2667–2675. doi:10.1021/es981241y.

786    Spear, J. R., Figueroa, L. A., and Honeyman, B. D. (2000). Modeling reduction of uranium
787            U(VI) under variable sulfate concentrations by sulfate-reducing bacteria. *Appl. Environ.*
788            *Microbiol.* 66, 3711–21. doi:10.1128/AEM.66.9.3711-3721.2000.

789    Takai, K., Campbell, B. J., Cary, S. C., Suzuki, M., Oida, H., Nunoura, T., et al. (2005).
790            Enzymatic and genetic characterization of carbon and energy metabolisms by deep-sea
791            hydrothermal chemolithoautotrophic isolates of Epsilonproteobacteria. *Appl. Environ.*
792            *Microbiol.* 71, 7310–20. doi:10.1128/AEM.71.11.7310-7320.2005.

793    Tatusov, R. L., Galperin, M. Y., Natale, D. A., and Koonin, E. V (2000). The COG database: a
794            tool for genome-scale analysis of protein functions and evolution. *Nucleic Acids Res.* 28,
795            33–6. Available at: http://www.ncbi.nlm.nih.gov/pubmed/10592175 [Accessed March 27,
796            2018].

797    Thullner, M., Dale, A. W., and Regnier, P. (2009). Global-scale quantification of mineralization
798            pathways in marine sediments: A reaction-transport modeling approach. *Geochemistry,*
799            *Geophys. Geosystems* 10, n/a-n/a. doi:10.1029/2009GC002484.

800    Trivedi, C. B., Lau, G. E., Grasby, S. E., Templeton, A. S., and Spear, J. R. (2018). Low-
801            temperature sulfidic-ice microbial communities, Borup Fiord Pass, Canadian high Arctic.
802            *Front. Microbiol.* 9, 1622. doi:10.3389/FMICB.2018.01622.

803    Tully, B. J., Wheat, C. G., Glazer, B. T., and Huber, J. A. (2018). A dynamic microbial
804            community with high functional redundancy inhabits the cold, oxic subseafloor aquifer.
805            *ISME J.* 12, 1–16. doi:10.1038/ismej.2017.187.

806    Wright, K. E., Williamson, C., Grasby, S. E., Spear, J. R., and Templeton, A. S. (2013).
807            Metagenomic evidence for sulfur lithotrophy by epsilonproteobacteria as the major energy
808            source for primary productivity in a sub-aerial arctic glacial deposit, Borup Fiord Pass.
809            *Front. Microbiol.* 4, 1–20. doi:10.3389/fmicb.2013.00063.

810    Zhang, J., Kobert, K., Flouri, T., and Stamatakis, A. (2014). PEAR: A fast and accurate Illumina
811            Paired-End reAd mergeR. *Bioinformatics* 30, 614–620. doi:10.1093/bioinformatics/btt593.

**Figures**

 **Figure 1. Borup Fiord Pass satellite view.** Image was collected on 2 July 2014. M- and A-
 samples from 2014 are marked with gold circles and red squares, respectively. The 2016 Spring
 region is highlighted with a dashed gold line, and the location of the spring is marked with a
 black triangle. Samples from 2017 were collected from within yellow dashed area. The inset line
 map shows the approximate location of Borup Fiord Pass (red dot) on Ellesmere Island,
 Nunavut, Canada. *Satellite image courtesy of the DigitalGlobe Foundation, and inset line map*
 *courtesy of Wikimedia Commons and the CC 3.0 license.*

828
829
830 **Figure 2. BFP sampling locations. A)** The "Sulfidic Aufeis" from Trivedi, et al. (2018) during
831 2014. Note: Two samples locations are to the southwest of this picture and their approximate
832 locations can be seen in Figure 1. **B)** The sulfidic aufeis on the left of the picture with the 2016
833 Spring a few hundred meters to the east (indicated by the orange dashed line; also see Figure S1).
834 **C)** The same location and focus of the sampling in 2017.

835
836

837
838
839 **Figure 3. Anvi'o diagram and gene presence/absence table. A)** This diagram shows all MAGs that are >50% completion and <10%
840 redundancy. Higher contribution to a given MAG by a sample is indicated by darker bars, while lower contribution is indicated by
841 lighter/no bars. **B)** This table shows the presence/absence of the specific genes queried in relation to the MAGs generated. Individual
842 assembly statistics are available in Table 1.

843
844
845

846
847

848
849
850 **Figure 4. Cladogram of MAGs.** Cladogram of metagenome assembled genomes (MAGs) and
851 nearest neighbors to confirm putative taxonomy. Created in the interactive tree of life (iTOL)
852 using the output tree from pparser after initial identification in CheckM.
853
854

855
856

857
858
**Figure 5. Observed gene sequences within queried MAG contigs.** Total number of select
metabolism-associated gene sequences found when queried against all 31 reportable MAGs.
Gene numbers are normalized to the total number of genes identified within the co-assembly
MAGs. *fcc*, sulfide dehydrogenase; *sqr*, sulfide-quinone reductase; *sorAB*, sulfite
dehydrogenase; soxXA, soxYZ, soxB, soxCD, proteins making up the Sox enzyme complex;
*aprBA*, SAT, *qmoABC*, gene operon; PAPS, 3'-phosphoadenosine 5'-phosphosulfate; *dsrAB*,
dissimilatory sulfite reductase; *phsABC*, thiosulfate reductase; *nap*, periplasmic nitrate
reductase; *nar*, nitrate reductase; *nrf*, nitrite reductase; *nir*, nitrite reductase; rCoA, reductive
acetyl-coenzyme A (Wood-Ljungdahl); *aclAB*, ATP citrate lyase; rTCA, phosphoenolpyruvate
carboxylase; *cbbL/cbbM*, ribulose 1,5-bisphosphate carboxylase/oxygenase; *cbb3*, cytochrome *c*
oxidase; *hydAB*, hydrogenase I; *hox*, NiFe hydrogenase; *mmo*, methane monooxygenase.
870

**Table 1. Metadata for all reported MAGs.** All MAGs shown are >50% completion with <10% redundancy via CheckM. Nearest neighbors were placed in a cladogram via CheckM/pplacer and confirmed via RAST (Figure 5).

| MAG # | Putative taxonomy | Length[a] | Contigs[b] | N50[a] | GC[c] | Comp.[c] | Redund.[c] |
|---|---|---|---|---|---|---|---|
| 1 | Epsilonproteobacteria; *Sulfurimonas* | 2064274 | 14 | 214847 | 38.19 | 99.59 | 0 |
| 2 | Gammaproteobacteria; *Shewanella* | 4979965 | 67 | 119878 | 41.35 | 99.45 | 1.12 |
| 3 | Alphaproteobacteria; *Rhodopseudomonas* | 7775429 | 338 | 70375 | 63.24 | 99.15 | 3.72 |
| 4 | Gammaproteobacteria; *Cellvibrio* | 3581784 | 67 | 99909 | 43.59 | 98.28 | 2.46 |
| 5 | Betaproteobacteria; *Limnobacter* | 3445427 | 282 | 19477 | 52.80 | 97.39 | 1.13 |
| 6 | Alphaproteobacteria; Erythrobacteraceae (f) | 3009949 | 326 | 11575 | 58.93 | 93.54 | 1.92 |
| 7 | Deltaproteobacteria; *Desulfocapsa* | 2805299 | 284 | 13203 | 51.33 | 92.84 | 2.28 |
| 8 | Actinobacteria; Coriobacteriales (o) | 2150593 | 263 | 13565 | 60.87 | 92.50 | 3.33 |
| 9 | Epsilonproteobacteria; *Sulfurovum* | 1833503 | 136 | 22712 | 38.75 | 92.21 | 1.84 |
| 10 | Alphaproteobacteria; *Loktanella* | 3281266 | 483 | 7930 | 60.60 | 89.02 | 2.30 |
| 11 | Gammaproteobacteria; *Arenimonas* | 2721939 | 304 | 11951 | 68.73 | 87.86 | 1.12 |
| 12 | Deltaproteobacteria; *Desulfocapsa* | 2954298 | 447 | 7903 | 50.14 | 84.94 | 2.08 |
| 13 | Cytophagia; *Algoriphagus* | 4253479 | 791 | 5920 | 41.69 | 80.28 | 9.29 |
| 14 | Betaproteobacteria; *Herminiimonas* | 2209716 | 410 | 5958 | 51.10 | 79.43 | 3.13 |
| 15 | Gammaproteobacteria; *Marinobacter* | 3818186 | 748 | 5456 | 54.16 | 78.56 | 5.04 |
| 16 | Gammaproteobacteria; *Pseudomonas* | 6369279 | 177 | 60989 | 58.13 | 77.54 | 0.88 |
| 17 | Chloroflexi; Chloroflexales (o) | 4795723 | 834 | 6374 | 49.05 | 74.65 | 3.67 |
| 18 | Sphingobacteriia; *Pedobacter* | 3604420 | 465 | 10926 | 33.46 | 73.33 | 5.44 |
| 19 | Betaproteobacteria; *Thiobacillus* | 2632478 | 497 | 5833 | 61.30 | 72.78 | 1.57 |
| 20 | Betaproteobacteria; *Rhodoferax* | 2881094 | 579 | 5077 | 59.36 | 72.38 | 3.83 |
| 21 | Deltaproteobacteria; *Geopsychrobacter* | 2948378 | 557 | 5659 | 51.99 | 69.53 | 0.65 |
| 22 | Elusimicrobia (p); Candidatus Endomicrobium | 2483169 | 437 | 6174 | 58.20 | 69.41 | 2.35 |
| 23 | Gammaproteobacteria; *Thiomicrospira* | 2196521 | 426 | 5432 | 42.35 | 68.46 | 7.66 |

| 24 | Alphaproteobacteria; *Sandarakinorhabdus* | 2926576 | 607 | 4990 | 63.80 | 67.43 | 1.12 |
|----|-------------------------------------------|---------|-----|------|-------|-------|------|
| 25 | Alphaproteobacteria; *Sandarakinorhabdus* | 2352194 | 538 | 4519 | 64.93 | 62.24 | 4.89 |
| 26 | Planctomycetia; *Isosphaera* | 2408829 | 457 | 5553 | 45.58 | 61.92 | 2.43 |
| 27 | Betaproteobacteria; *Hylemonella* | 2910411 | 640 | 4770 | 62.31 | 61.79 | 0.44 |
| 28 | Deinococci; *Deinococcus* | 3481023 | 613 | 6273 | 60.43 | 59.34 | 2.97 |
| 29 | Betaproteobacteria; *Methylotenera* | 1596490 | 423 | 3649 | 52.68 | 55.17 | 8.31 |
| 30 | Mollicutes; Acholeplasmataceae (f) | 974526 | 65 | 28355 | 33.20 | 51.75 | 4.55 |
| 31 | Flavobacteriia; *Flavobacterium* | 1930691 | 418 | 4726 | 34.69 | 51.45 | 2.36 |

877    [a] base pairs (bp)

878
879    [b] total number of contigs

880    [c] percentage (%)

    **Supplementary Materials**

884
885
886
887
888    **Table S1. Co-assembly metadata.** These data represent raw values from Megahit prior to
889    mapping through Bowtie2 and final processing via Anvi'o.
890

| Sample | Type | Total bp | Contigs | Max contig (bp) | Avg (bp) | N50 (bp) |
|---|---|---|---|---|---|---|
| BFP14_A4b | Aufeis | 85484280 | 30017 | 380944 | 2848 | 3571 |
| BFP14_A6 | Aufeis | 82535183 | 33359 | 128624 | 2474 | 2871 |
| BFP14_M2 | Melt pool | 10144305 | 4841 | 418178 | 2095 | 1977 |
| BFP14_M4b | Melt pool | 1.59E+08 | 71593 | 111481 | 2215 | 2359 |
| BFP16_Spring | Spring fluid | 60894340 | 18996 | 206795 | 3206 | 4748 |
| BFP17_14C | Aufeis surface | 6.38E+08 | 272035 | 297076 | 2347 | 2611 |
| BFP17_AS3b | Aufeis surface | 1.98E+08 | 78690 | 78857 | 2513 | 3028 |
| BFP17_AS4e | Aufeis surface | 89868307 | 27395 | 44699 | 3280 | 4468 |
| BFP17_AS6b | Aufeis surface | 1E+08 | 34932 | 60991 | 2866 | 4186 |
| **Co-Assembly** | | **1.16E+09** | **477394** | **406785** | **2440** | **2778** |

899
900 **Table S2. JSpeciesWS results.** MAGs 7 and 12 were compared to database genomes for
901 *Desulfocapsa sulfexigens* and *Desulfotalea psychrophila* LSv54. Average nucleotide identity was
902 calculated for BLAST and MUMmer as well as Tetra-nucleotide comparison.
903

| #ANI BLAST and aligned percentage | | | | |
|---|---|---|---|---|
| | Desulfotalea psychrophila LSv54 | Desulfocapsa sulfexigens DSM 10523 | MAG 12 | MAG 7 |
| Desulfotalea psychrophila LSv54 | * | 66.86 [20.05] | 67.07 [14.44] | 67.13 [14.59] |
| Desulfocapsa sulfexigens DSM 10523 | 66.75 [18.50] | * | 68.41 [25.07] | 68.16 [24.91] |
| MAG 12 | 67.28 [20.09] | 68.74 [35.44] | * | 81.71 [44.41] |
| MAG 7 | 67.17 [20.68] | 68.48 [36.64] | 82.11 [44.96] | * |

| #ANI MUMmer and aligned percentage | | | |
|---|---|---|---|---|
| | Desulfotalea psychrophila LSv54 | Desulfocapsa sulfexigens DSM 10523 | MAG 12 | MAG 7 |
| Desulfotalea psychrophila LSv54 | * | 87.49 [1.04] | 87.97 [0.04] | 85.18 [0.14] |
| Desulfocapsa sulfexigens DSM 10523 | 87.74 [0.47] | * | 82.75 [0.55] | 82.87 [0.26] |
| MAG 12 | 87.97 [0.02] | 82.75 [0.75] | * | 85.30 [33.12] |
| MAG 7 | 85.18 [0.17] | 82.84 [0.41] | 85.30 [34.60] | * |

| #Tetra results | | | | |
|---|---|---|---|---|
| | Desulfotalea psychrophila LSv54 | Desulfocapsa sulfexigens DSM 10523 | MAG 12 | MAG 7 |
| Desulfotalea psychrophila LSv54 | * | 0.90901 | 0.86371 | 0.84073 |
| Desulfocapsa sulfexigens DSM 10523 | 0.90901 | * | 0.92621 | 0.90605 |
| MAG 12 | 0.86371 | 0.92621 | * | 0.98997 |
| MAG 7 | 0.84073 | 0.90605 | 0.98997 | * |

906
907

908
909

910
911
912 **Figure S1. BFP16 Spring location.** Shown is the location of the 2016 spring. Its exact location
913 is at the northern-most portion of the yellow-colored area, approximately in the center of the
914 picture. Also shown is the area known as the "Sulfidic Aufeis" from Trivedi, et al. (2018)
915 towards the right edge of the picture. This is the location of sampling from 2014 and 2017 field
916 campaigns.

917
918

919
920
**Figure S2. Sample type examples.** Examples of sample types used for metagenomes. **A)** Melt
pool; **B)** Aufeis sample, where surface ice was removed, a pit dug and ice from this pit was
melted and filtered; **C)** Mineral precipitates from the 2017 BFP site.

924
925

926
927

**Figure S3. Metabolic gene grid (Figure 3b) with associated counts for each.** This grid, presented in the main text as Figure 3b, now with gene counts shown. Each instance of a specific gene found in each MAG was recorded here and then the presence/absence figure was generated. Counts were normalized to generate Figure 5.

43