1   **G-baToN: a versatile cell-cell interaction reporter system**

Rui Tang[1*], Christopher W. Murray[2], Ian Linde[4], Nicholas J. Kramer[1,3], Min K. Tsai[1], Leo Chen[1],
Hongchen Cai[1], Aaron D. Gitler[1,3], Edgar Engleman[2,4,5], and Monte M. Winslow[1,2,5*]

[1] Department of Genetics, Stanford University School of Medicine, Stanford, CA, USA
[2] Cancer Biology Program, Stanford University School of Medicine, Stanford, CA, USA
[3] Neuroscience Program, Stanford University School of Medicine, Stanford, CA, USA
[4] Immunology Program, Stanford University School of Medicine, Stanford, CA, USA
[5] Department of Pathology, Stanford University School of Medicine, Stanford, CA, USA

* Corresponding authors: Rui Tang: tangrui@stanford.edu and Monte M. Winslow: mwinslow@stanford.edu

ABSTRACT

Cell-cell interactions influence all aspects of development, homeostasis, and disease. Here, we

describe GFP-based Touching Nexus (G-baToN) which relies upon fluorescent protein transfer

driven by nanobody-mediated recognition to enable sensitive and specific labeling of cells after

cell-cell interactions. A suite of orthogonal baToN tools enables reciprocal cell-cell labeling,

interaction-dependent cargo transfer, and the identification of higher order cell-cell interactions

across a wide range of cell types.

 MAIN TEXT

Cell-cell interactions contribute to almost all physiological and pathological states.

Despite the explosion of interest in uncovering and understanding cell heterogeneity in tissues

and across disease states, the extent to which cell-cell interactions influence cell state, drive

heterogeneity, and enable proper tissue function remains poorly understood[1, 2]. Detailed analysis

of the impact of defined cell-cell interactions has illuminated critical aspects of biology, however,

these analyses have been limited to a small collection of signaling interfaces engaged by direct

cell-cell contact that are tractable to study. Molecular methods to profile cell state, including *in*

situ approaches within intact tissue architecture, largely fail to uncover the causal relationship of

cell-cell interactions with the underlying biology[3, 4]. Computational and experimental methods to

characterize cell-cell interactions yield additional layers of dimensionality, however, modalities

to capture cell-cell interactions are limited[5-7] (Supplementary Fig. 1). Much as diverse systems to

detect and quantify protein-protein interactions have revolutionized our biochemical

understanding of molecular systems, the development of novel systems to detect and quantify

cell-cell interactions should help to map the interaction networks of multicellular systems

(Supplementary Fig. 1).

Endogenous cell-cell interactions can result in the transfer of surface proteins between

cells, thus we sought to couple this phenomenon with fluorescent protein tagging as a means of

labeling cells that have undergone direct interaction with a neighboring cell[8-10]. Specifically, we

adapted an engineered surface protein/ligand system based on the expression of surface GFP

(sGFP) on sender cells and a cell surface anti-GFP (αGFP) nanobody on receiver cells[6, 11, 12]. Co-

culturing sGFP sender cells with αGFP receiver cells led to GFP transfer and labeling of the

receiver cells (Fig. 1a, b and Supplementary Fig. 1). Receiver cell labeling required direct cell-

cell contact, active membrane dynamics, and pairing between sGFP and its cognate αGFP

receptor (Fig. 1c, d and Supplementary Fig. 2b, c). Interestingly, sGFP transfer was accompanied

by reduced GFP on the sender cells, downregulation of αGFP from the surface of the receiver

cells and was partially blocked by chemical inhibitors of endocytosis – all consistent with active

and extensive GFP transfer from sender to receiver cells (Supplementary Fig. 2d-f). Thus, this

simple system (which we term GFP-based Touching Nexus or G-baToN) enables robust and

efficient labeling of predefined receiver cells that have interacted with sender cells.

To further characterize the interaction reporter system, we deconstructed the G-baToN

design into three functional modules. We found that both the αGFP and sGFP transmembrane

domains influence GFP transfer efficiency (Supplementary Fig. 3a-c, g-h). Furthermore, the

affinity of the GFP-αGFP pairing impacted GFP transfer (Supplementary Fig. 3d-f). Thus, the

analysis of different transmembrane domains on sGFP and αGFP as well as different αGFP

nanobodies enabled the identification of combinations that improve unidirectional receiver cell

labeling.

To characterize the kinetics of G-baToN-mediated receiver cell labeling, we performed

co-culture time course experiments with time-lapse imaging and flow cytometry readouts. Time-

lapse imaging showed rapid transfer and internalization of GFP by receiver cells (Fig. 1e and

Supplementary Movie 1). GFP transfer could be detected within five minutes of co-culture and

was half-maximal after six hours (Fig. 1f and Supplementary Fig. 2g). Importantly, GFP

fluorescence in receiver cells decayed rapidly after isolation of touched receiver cells from

sender cells, thus documenting the transient "real-time" labeling of receiver cells (Supplementary

Fig. 2h). To determine the sensitivity of this system, we co-cultured receiver cells with different

numbers of sender cells. The fraction of labeled receiver cells was proportional to the number of

sender cells and even the addition of very few sender cells (representing less than 1 sender

cell/10^5 receiver cells) was sufficient to label rare receiver cells (Fig. 1g, h). Thus, the transfer of

GFP to αGFP-expressing cells is a rapid and sensitive method to mark cells that have physically

interacted with a predefined sender cell type.

Given the high efficiency with which sGFP labels receiver cells upon interaction with

cognate sender cells, we tested whether other surface antigen/antibody pairs could lead to protein

transfer and labeling. Due to the cross reactivity of αGFP with BFP, co-culture of surface BFP

(sBFP) sender cells with αGFP receiver cells generated BFP-labeled receivers[11] (Supplementary

Fig. 4a, b). Orthogonal systems consisting of surface-mCherry/αmCherry (LaM4)[11] and surface-

GCN4-GFP/αGCN4 (single-chain variable fragment, scFV)[13] also led to efficient and specific

receiver cell labeling (Supplementary Fig. 4c-f).

We then integrated these orthogonal baToN systems to enable reciprocal labeling and

detection higher order multi-cellular interactions. Engineering cells with baToN systems in an

anti-parallel fashion should enable reciprocal labeling of both interacting cells. Co-culture of

cells expressing sGFP and αmCherry with cells expressing smCherry and αGFP resulted in

reciprocal labeling of both interacting cell types (Supplementary Fig. 5a-c). Using orthogonal

ligand-receptor pairs, we also created an AND gate dual labeling strategy. Specifically, co-

expression of αmCherry and αGFP on receiver cells enabled dual color labeling of receiver cells

that had interacted with smCherry-expressing, sGFP-expressing, or both sender cell types (Fig.

2a, b, Supplementary Fig. 5d). Analogously, we achieved dual color labeling of receiver cells by

leveraging the ability of αGFP to bind both sGFP and sBFP (Supplementary Fig. 5e, f). Thus,

derivatives of the G-baToN system allow for additional degrees of resolution of complex cell-

cell interactions.

Given the high efficiency of protein transfer using the G-baToN system, we also

investigated whether cargo proteins could be co-transferred with GFP from sender cells to

receiver cells. Initially, we generated sender cells with surface expression of a GFP-tdTomato

fusion protein (sGFP-Tom) and uncovered stoichiometric tdTomato and GFP transfer to αGFP

receiver cells (Supplementary Fig. 6a, b). We extended this system further by generating sender

cells expressing the HaloTag protein fused to sGFP (sHalo-GFP; Fig. 2c)[14]. Covalent attachment

of a fluorescent synthetic ligand to sHalo-GFP enabled specific loading onto sender cells

(Supplementary Fig. 7a). Co-culture of Alexa Fluor 660 (AF660)-loaded sHalo-GFP sender cells

with αGFP receiver cells enabled co-transfer of both GFP and AF660 (Fig. 2d and

Supplementary Fig. 7b). Compared to GFP, transfer of the chemical fluorophore using sHalo-

GFP-based labeling of receiver cells led to increased foreground to background and higher

sensitivity (Fig. 2d and Supplementary Fig. 7c). Importantly, changing from a protein (GFP) to a

chemical fluorophore also extended the half-life of labeling, thus enabling partially tunable

perdurance of labeling after touching (Supplementary Fig. 7d).

Next, we coupled the enhanced properties of chemical fluorophore-based labeling with

the generalizability of the GCN4-baToN system to assemble a robust and versatile system to

label receiver cells that have interacted with two or more different sender cell types (Fig. 2e, f).

Co-culturing αGCN4 receiver with AF488- and AF660-loaded sGCN4-Halo sender cells

generated receiver cells with varying degrees of AF488 and AF660 labeling (Supplementary Fig.

7e). Importantly, the ratio of AF488 to AF660 transferred to the dually labeled receiver cells

strongly correlates with the ratio of the two sGCN4-Halo sender populations within the co-

culture, suggesting that this system can quantitively measure higher order cell-cell interactions

(Fig. 2f).

Beyond fluorescent labels, we tested whether other cargo could be transferred to receiver

cells. We generated sGFP-PuroR-expressing sender cells and found that co-culture of sGFP-

PuroR sender cells with αGFP receiver cells led to moderate puromycin resistance of touched

receiver cells (Supplementary Fig. 6c-e). Furthermore, loading of sGCN4-HaloTag sender cells

with biotinylated ssDNA and subsequent staining with streptavidin-conjugated AF647 prior to

co-culture with αGCN4 receiver cells revealed successful co-transfer of fluorescently labeled

ssDNA (Fig. 2g, h and Supplementary Fig. 7f-g). Thus, baToN systems can achieve contact-

dependent transport of different macromolecules between cells.

To assess the generalizability of the G-baToN system across cell types, we expressed

αGFP on a panel of cell lines and primary cells. Each receiver cell type was able to uptake GFP

from sGFP-expressing KPT sender cells upon cell-cell contact (Fig. 3a). Furthermore, diverse

sender cancer cell lines and primary cell types expressing sGFP were able to transfer GFP to

αGFP-expressing 293 receiver cells (Fig. 3b, Supplementary Fig. 8). Cancer cells undergo

interactions with a variety of cell types within primary tumors as well as at metastatic sites. Co-

culturing sGFP-expressing lung cancer cells with primary human endothelial cells led to robust

endothelial cell labeling (Fig. 3c, d). As anticipated, receiver cell labeling required sGFP-

expression on the sender cell and αGFP expression on the receiver cells. Furthermore, co-

culturing sGFP-expressing cancer cells with primary cortical neurons expressing αGFP also led

to receiver cell labeling (Supplementary Fig. S9c, d). Thus, G-baToN-based labeling extends

beyond transformed cell types and can label diverse primary cell types.

To further test the generalizability of this system and determine whether primary cells

can serve as both sender and receiver cells, we assessed G-baToN-based GFP transfer in the

context of two well-established heterotypic cell-cell interactions: endothelial cells with smooth

muscle cells and astrocytes with neurons. Consistent with our previous results, we found that

sGFP-expressing primary cells (both endothelial cells and astrocytes) were able to transfer GFP

to neighboring αGFP-expressing receivers (Fig. 3e, f: smooth muscle cells; Fig. 3g and

Supplementary Fig. 9a, b: cortical neurons). Finally, given the importance of adaptive immune

cell interactions with cancer cells, we assessed whether the G-baToN could map the interactions

of primary human CD4 and CD8 T cells with lung cancer cells (Fig. 3h and Supplementary Fig.

10a, b). αGFP-expressing CD4 and CD8 T cells that interacted with sGFP-expressing lung

cancer cells were specifically labeled (Fig. 3h and Supplementary Fig. 10a). Inversely, αGFP-

expressing cancer cells that interacted with sGFP-expressing CD4 T cells were also specifically

identified by G-baToN labeling (Supplementary Fig. 10b).

Our data document the ability of diverse primary cell types to serve as both sender and

receiver cells, suggesting that baToN systems are not only simple, sensitive and rapid, but also

generalizable. Multicolor baToN systems will enable qualitative and quantitative analyses of

higher-order interactions. Finally, the ability of baToN systems to co-transfer protein, DNA or

chemical cargos suggests that this versatile platform could be leveraged to manipulate target cell

functions. The simplicity of this two-component system, combined with its generalizability

across cell types, excellent foreground to background, and rapid labeling, should enable facile

analysis of cell-cell interaction dynamics and have a broad impact on our ability to understand

the outputs of cell-cell interactions across various biological systems.

Figure Legends

Figure 1. GFP-based Touching Nexus (G-baToN) leads to cell-cell interaction-dependent receiver cell labeling.

a. Schematic of the G-baToN system. Surface GFP (sGFP) on a sender cell is transferred to a receiver cell expressing a cell surface anti-GFP nanobody (αGFP) leading to GFP labeling of the "touched" receiver.

b. GFP transfer from sGFP-expressing KPT lung cancer sender cells (marked by intracellular tdTomato) to αGFP-expressing 293 receiver cells. Receiver cell labeling is sGFP- and αGFP-dependent. Control sender cells do not express sGFP. Control receiver cells do not express αGFP. Cytoplasmic GFP (Cyto-GFP) is not transferred to receiver cells. Sender and receiver cells were seeded at a 1:1 ratio and co-cultured for 24 hours. Receivers were defined as $Tomato^{neg}PI^{neg}$ cells.

c. GFP transfer to 293 receiver cells requires direct cell-cell contact. Receiver cells separated from sender cells by a transwell chamber are not labeled. Sender and receiver cells were seeded in upper and lower chambers respectively at a 1:1 ratio and cultured for 24 hours. Receivers were defined as $Tomato^{neg}PI^{neg}$ cells.

d. GFP transfer to 293 receiver cells requires sGFP-αGFP interaction and is blocked by anti-GFP antibody in a dose-dependent manner. sGFP sender cells were pre-incubated with the indicated

concentration of anti-GFP antibody for 2 hours, washed with PBS, and then co-cultured with

receiver cells at a 1:1 ratio for 24 hours. Receivers were defined as Tomato^{neg}PI^{neg} cells.

e. Time-lapse imaging of GFP transfer from a sGFP-expressing sender cell to an αGFP-

expressing receiver cell. Time after contact is indicated. Receiver cell is outlined with white

dashed line. Scale bar: 10 μm.

f. Analysis of GFP Mean Fluorescence Intensity (MFI) of αGFP receiver cells (marked by

intracellular BFP) co-cultured with sGFP sender cells (marked by intracellular tdTomato) co-

cultured for the indicated amount of time. Sender and receiver cells were seeded at a 1:1 ratio.

Receivers were defined as Tomato^{neg}PI^{neg}BFP^{pos} cells.

g. Percentage of labeled αGFP receiver cells after co-culture with different numbers of sender

cells for 24 hours. Receivers were defined as Tomato^{neg}PI^{neg}BFP^{pos} cells.

h. Detection of rare labeled αGFP receiver cells after co-culture with sGFP sender cells at

approximately a 1:10^5 ratio for 24 hours. Receivers were defined as Tomato^{neg}PI^{neg}BFP^{pos} cells.

Figure 2. X-baToN systems enable recording of multicellular interactions and cell-cell

interaction-based cargo transfer.

a. Schematic of the AND gate-baToN system. sGFP and smCherry sender cells express either

sGFP or mCherry. Dual receiver cells express both αGFP (LaG17, tagged by Myc-tag) and

αmCherry (LaM4, tagged by intracellular BFP).

b. Representative FACS plots of dual receiver 293 cells cultured with the indicated 293 sender

cells at 1:1 (for single sender cell) or 1:1:1 (for dual sender cells) ratios. Percent of labeled

receiver cells (gated as BFP^{pos}) after 24 hours of co-culture is indicated as mean +/- SD of

triplicate cultures.

c. Diagram of HaloTag-baToN system. Sender cells (marked by intracellular 2A-mCherry)

express surface HaloTag-GFP fusion which can be loaded with HaloTag ligands (in this example

AF660). Receiver cells express αGFP (LaG17, tagged by intracellular BFP).

d. Representative plot of GFP and AF660 intensity in αGFP 293 receiver cells co-cultured with

HaloTag-GFP KP (lung adenocarcinoma) sender cells at a 1:1 ratio for 6 hours. Receivers were

defined as mCherry^{neg}PI^{neg}BFP^{pos} cells.

e. Diagram of dual color GCN4-HaloTag-baToN system. Sender cells (marked by intracellular

2A-mCherry) express surface 4XGCN4 associated with HaloTag, loaded with either AF488- or

AF660- conjugated HaloTag ligand. Receiver cells express αGCN4 (tagged by intracellular

BFP).

f. AF488/AF660 GCN4-HaloTag sender ratio in the co-culture directly proportional to

AF488/AF660 intensity (MFI) of αGCN4 receiver after 6 hours of co-culture. Receivers were

defined as mCherry^{neg}PI^{neg}BFP^{pos} cells.

g. Diagram of using GCN4-HaloTag sender to transfer ssDNA into αGCN4 receiver cells.

Sender cells (marked by intracellular 2A-mCherry) express surface 4XGCN4 associated with

HaloTag, loaded with 5' HaloTag ligand, 3' biotin dual conjugated ssDNA (21 nt), then stained

with Avidin-AF647. Receiver cells express αGCN4 (tagged by intracellular BFP).

h. Representative plot of AF647 intensity in αGCN4 293 receiver cells co-cultured with GCN4-

HaloTag 293 sender cells at a 1:1 ratio for 6 hours. Receivers were defined as

mCherry^{neg}PI^{neg}BFP^{pos} cells.

Figure 3. G-baToN is a generalized system that can be used for touching-based labeling

between various cell types.

a. Many different cell lines and primary cell types can serve as receiver cells. Receiver cells

expressing αGFP were co-cultured with sGFP-expressing KPT sender cells for 24 hours. MFI of

GFP on Tomato^{neg}PI^{neg}BFP^{pos} receiver cells was assessed by FACS analysis and is shown as fold

change relative to monocultured receiver cells. For lung epithelial cells and kidney epithelial

cells MFI of GFP on Tomato^{neg}PI^{neg}BFP^{pos}EpCAM^{pos} cells is shown as mean +/- SD of triplicate

cultures. SkMC: skeletal muscle cells. Astro: Astocytes. Hepato: hepatocytes. Spleno:

splenocytes.

b. Many different cell lines and primary cell types can serve as sender cells. Sender cells

expressing sGFP were co-cultured with αGFP 293 receiver cells for 24 hours. MFI of GFP on

mCherry^{neg}PI^{neg}BFP^{pos} receiver cells was assessed by flow cytometry and is shown as fold

change relative to monocultured 293 receiver cells. Mean +/- SD of triplicate cultures is shown.

c, d. G-baToN can detect cancer cell-endothelial cell (EC) interactions. HUVECs expressing

αGFP were co-cultured with or without Tomato^{pos} sGFP-expressing lung cancer sender cells at a

1:1 ratio for 24 hours. MFI of GFP on PI^{neg}Tomato^{neg}BFP^{pos}CD31^{pos} Receiver cells was assessed

by flow cytometry and is shown as mean +/- SD of triplicate cultures (**c**). Representative images

of Tomato^{pos} sGFP-expressing lung cancer sender cells co-cultured with either control HUVEC

receiver cells (HUVECs expressing BFP) or αGFP HUVEC receiver cells at a 1:1 ratio for 24

hours (**d**). Scale bars = 50 μm.

e,f. G-baToN can detect endothelial cell (EC)-smooth muscle cell (SMC) interactions. Primary

human umbilical artery smooth muscle cells (HUASMC) expressing αGFP were co-cultured

with or without sGFP-expressing HUVEC senders cells at a 1:1 ratio for 24 hours. MFI of GFP

on PI^{neg}BFP^{pos} receiver cells was assessed by flow cytometry and is shown as mean +/- SD of

triplicate cultures (**e**). Representative images of sGFP-expressing HUVEC sender cells co-

cultured with either control HUASMC receiver cells (expressing BFP) or αGFP HUASMC

receiver cells at a 1:1 ratio for 24 hours (**d**). Scale bars = 50 μm.

g. Representative images of sGFP-expressing astrocyte sender cells co-cultured with either

control neuron receivers or αGFP neuron receivers at a 1:2 ratio for 24 hours. Neurons were

stained with Microtubule Associated Protein 2 (Map2). Scale bars = 50 μm. Higher

magnification of the boxed areas are shown on the right.

h. G-baToN can detect cancer cell-T cell interactions. Primary human CD4^{pos} (left) or CD8^{pos}

(right) T cells were co-cultured with sGFP-expressing lung cancer sender cells (A549 cells) at a

2:1 ratio for 24 hours. Receivers were defined as mCherry^{neg}Near-IR^{neg}BFP^{pos}CD4^{pos}/CD8^{pos} T

cells.

Supplementary Figure 1. Features of the synNotch, LIPSTIC, and G-BaToN cell-cell

interaction reporter systems.

Overview of systems that enable labeling of cells after cell-cell contact.

Supplementary Figure 2. GFP transfer requires direct GFP-αGFP interaction.

a. GFP protein is transferred to receiver cells. Western blot analysis of FACS purified Control

and αGFP receiver 293 cells cultured in isolation or co-culture with sGFP Tomato-positive

sender cells (mouse KPT lung cancer cells) for 24 hours. sGFP but not Tomato is present in

touched αGFP receiver cells. Human mitochondrial antigen (hu-Mito) is a marker for receiver

cells. GAPDH shows loading. Rightmost lane is sGFP sender cells.

b. GFP cannot be transferred from fixed sGFP sender cells to live αGFP receiver cells. sGFP

sender cells were fixed in 1% PFA for 5 minutes and washed with PBS before co-cultured with

αGFP receiver cells at a 1:1 ratio for 24 hours. Receivers were defined as $Tomato^{neg}PI^{neg}BFP^{pos}$

cells.

c. GFP transfer to 293 receiver cells required sGFP-αGFP recognition. GFP is transferred from

sGFP sender cells to αGFP-receiver cells but not from sGFP sender cells to αmCherry-receiver

cells. Control receiver cells do not express any nanobody. Sender and receiver cells were co-

cultured at a 1:1 ratio for 24 hours. Receivers were defined as $Tomato^{neg}PI^{neg}BFP^{pos}$ cells.

d. GFP transfer to receiver cells is accompanied by a reduction of GFP on the sender cells. GFP

expression on sender cells after 24 hour co-culture with control or αGFP-expressing 293 receiver

cells at a 1:1 ratio. Co-culture with αGFP expressing but not control receiver cells reduced GFP

on sGFP sender cells. Senders were defined as $Tomato^{pos}DAPI^{neg}BFP^{neg}$ cells.

e. GFP transfer is accompanied with αGFP internalization on receiver cells. Analysis of surface

αGFP (Myc-tag) on 293 receiver cell co-cultured for 24 hours with sGFP sender cells. Receivers

were defined as $Tomato^{neg}PI^{neg}$ cells.

f. GFP transfer to 293 receiver cells is partially dependent on membrane dynamics of

endocytosis. Both a clathrin inhibitor (Pitstop, 20 μM) and a dynamin inhibitor (Dyngo 4a, 10

μM) partially inhibit GFP transfer from sGFP sender cells to αGFP receivers. Inhibitors were

present during the 24 hour co-culture. Receivers were defined as $Tomato^{neg}PI^{neg}$ cells.

g. GFP transfer to αGFP 293 receiver cells can be very rapid. A shift in GFP MFI was detected 5

minutes after mixing sGFP sender cells with αGFP receiver cell. Receivers were defined as

$Tomato^{neg}PI^{neg}BFP^{pos}$ cells. MFI mean +/- SD of triplicate cultures is shown.

h. Rapid GFP degradation in touched receiver cells after removal of the sGFP sender cells. After

6 hours co-culture, GFP positive receiver cells were purified by FACS followed by culture

without sender cells. Receivers were defined as $mCherry^{neg}PI^{neg}BFP^{pos}$ cells. GFP MFI in

receiver cells reduced rapidly ($T_{1/2}$ approximately 2 hours). MFI mean +/- SD of triplicate

cultures is shown.

Supplementary Figure 3. Transmembrane domains and the nanobody affinity impact

sGFP transfer and receiver cell labeling.

a. Schematic of the sender and receiver cells used to determine the impact of different αGFP

transmembrane (TM) domains. TM domains contain the TM domain itself as well as membrane

proximal regions from the indicated mouse (m) and human (h) proteins.

b. Different TM domains impact cell surface αGFP expression on 293 receiver cells. Membrane

αGFP was accessed by anti-Myc staining. Control receiver cells do not express any nanobody.

Mean +/- SD of Myc MFI in triplicate cultures is shown.

c. VEGFR2 TM domain on αGFP receivers enables highest GFP transfer efficiency. Receiver

cells expressing αGFP linked by different TM domains were co-cultured with sGFP sender cells

at a 1:1 ratio for 6 hours. Receivers were defined as Tomato^{neg}PI^{neg} cells.

d. Schematic of the sender and receiver cells used to determine the impact of different sGFP TM

domains and different αGFP nanobodies on G-baToN-based labeling. TM domains contain the

TM domain itself as well as membrane proximal regions from the indicated mouse (m) and

human (h) proteins.

e. Different nanobodies exhibit different levels of expression on 293 receiver cells. Total αGFP

expression was assessed by BFP intensity. Mean +/- SD of GFP MFI in triplicate cultures is

shown.

f. αGFP affinity influences transfer of GFP to touched 293 receiver cells. Receiver cells

expressing different αGFP nanobodies were co-cultured with sGFP sender cells at a 1:1 ratio for

6 hours. GFP transfer was assessed by flow cytometry. GFP intensity on $Tomato^{neg}PI^{neg}BFP^{pos}$

receiver cells is shown as mean +/- SD of triplicate cultures.

g. Different TM domains on sGFP impact its expression in 293 sender cells. sGFP expression in

sender cells was assessed by flow cytometry for GFP. Mean +/- SD of GFP MFI in triplicate

cultures is shown.

h. PDGFR TM domain on sGFP minimized retrograde transfer of αGFP from receiver cells to

293 sGFP sender cells. αGFP transfer to sGFP senders were determined as the percentage of

$mCherry^{pos}GFP^{pos}$ sender cells that were also BFP^{pos}. Cells were co-cultured for 6 hours at a 1:1

ratio. Mean +/- SD of triplicate cultures is shown.

Supplementary Figure 4. X-baToN systems enable fluorescent labeling via various antigen-

nanobody/scFV pairs.

a. Schematic of the surface BFP (sBFP)-baToN system. Sender cell (marked by intracellular

tdTomato) expresses sBFP and receiver cell expresses αGFP.

b. The cross reactivity of αGFP with BFP allow BFP to be transferred from sBFP sender to

αGFP receiver cells. sBFP senders and αGFP receivers were co-cultured at 1:1 ratio for 24 hours.

Receivers were defined as $toTomato^{neg}PI^{neg}$ cells. BFP MFI of receiver cells is shown as mean

+/- SD of triplicate cultures.

c. Schematic of the surface mCherry (smCherry)-baToN system. Sender cell (marked by

intracellular GFP) expresses smCherry and receiver cell expresses αmCherry (tagged by

intracellular BFP)

d. The pairing between mCherry and αmCherry enable mCherry transfer from smCherry 293

sender cells to αmCherry-expressing 293 receiver cells. smCherry senders are not able to transfer

mCherry to αGFP receiver cells. smCherry sender cells were co-cultured with αGFP or

αmCherry receiver cells at a 1:1 ratio for 24 hours. Receivers were defined as GFP^{neg}BFP^{pos}

cells. mCherry MFI of receiver cell is shown as mean +/- SD of triplicate cultures. smCherry 293

sender is not able to transfer mCherry to αGFP 293 receiver.

e. Schematic of the surface GCN4 (sGCN4)-baToN system. Sender cells (marked by intracellular

2A-mCherry) express cell surface 4X GCN4 peptide fused with GFP. Receiver cells express a

cell surface anti-GCN4 single-chain variable fragment (scFV; αGCN4, tagged by intracellular

BFP).

f. The pairing between GCN4 and αGCN4 enables co-transfer of GFP from 4X sGCN4-GFP

senders to αGCN4 receivers 24 hours after co-culture at a 1:1 ratio. Receivers were defined as

mCherry^{neg}PI^{neg}BFP^{pos} cells. GFP MFI of receiver cell is shown as mean +/- SD of triplicate

cultures. sGFP sender are not able to transfer GFP to αGCN4 receivers.

Supplementary Figure 5. Dual color-baToN systems enable labeling in complex cell-cell

interaction systems.

a. Diagram of the Reciprocal-baToN system. Cell A expresses sGFP and αmCherry (tagged by

intracellular BFP), Cell B expresses smCherry and αGFP (tagged by Myc-tag).

b. Representative FACS plots of cell A and cell B monocultures (left 2 panels) and after co-

culture at a 5:1 ratio for 24 hours. Percent of labeled cells is indicated as mean +/- SD of

triplicate cultures.

c. Representative image of reciprocal labeling of cell A and cell B co-cultured for 24 hours. Cells

are outlined with a white dashed line. Cell A is BFP and GFP double positive and Cell B is

mCherry positive. Touched cell A is BFP, GFP and mCherry triple positive and touched cell B is

GFP and mCherry double positive (pointed out by white letter). Scale bar: 10 μm.

d. Representative image of dual receiver cells co-cultured with indicated sender cells. Cells are

outlined with a white dashed line. Dual receiver cells are BFP positive and sender touched dual

receiver cells are pointed out by white arrows. Scale bar: 10 μm.

e. Diagram of the BFP/GFP AND gate-baToN system. sBFP sender cells express intracellular

Tomato and surface BFP, sGFP sender cells express intracellular Tomato and surface GFP.

Common receiver cells expressed αGFP.

f. Representative FACS plots of common receiver 293 cells cultured with the indicated

Tomato^{pos} sender cells at 1:1 (for single sender cell) or 1:1:1 (for dual sender cells) ratios.

Receiver cells were gated as Tomato^{neg}PI^{neg}. Percent of labeled common receiver cells after 24

hours of co-culture is indicated as mean +/- SD of triplicate cultures.

Supplementary Figure 6. Using the G-baToN system to co-transfer cargo protein into

touched receiver cells.

a. Diagram of surface tdTomato-GFP (sGFP-Tom) co-transfer into touched receiver cells.

Sender cell expresses sGFP-tdTomato and receiver cell expresses αGFP (tagged by intracellular

BFP).

b. Representative FACS plots of αGFP 293 receiver cells cultured with the indicated sender cells

at 1:1 ratio for 24 hours. Percent of GFP/Tomato dual labeled receiver cells is indicated as mean

+/- SD of BFP^{pos} cells from triplicate cultures.

c. Diagram of GFP-PuroR co-transfer system. Sender cell expresses surface GFP associated with

PuroR (marked by intracellular tdTomato), receiver cell expresses αGFP (tagged by intracellular

BFP). PuroR: Gcn5-related N-acetyltransferase.

d. 293 Sender cells expressing sGFP-PuroR are only modestly resistant to puromycin treatment.

293 cells were transfected with plasmids as indicated. 24 hours after transfection, 293 cells were

treated with different doses of puromycin for 24 hours. Cell number was assessed by measuring

spectrophotometrically at 570 nm (OD570) after crystal violet staining.

e. Co-transfer of GFP-PuroR from sGFP-PuroR sender cells to αGFP 293 receiver cells confers

puromycin resistance to receiver cells. Control, sGFP or sGFP-PuroR sender cells were co-

cultured with αGFP 293 receiver cells for 24 hours at a 4:1 ratio before treatment with

different dose of puromycin for 48 hours. $\text{tdTomato}^{neg}\text{PI}^{neg}\text{BFP}^{pos}$ cell numbers were counted via

FACS.

Supplementary Figure 7. The HaloTag-baToN system enables touching based co-transfer.

a. Labeling HaloTag-expressing sender cells with AF660 fluorophore. Representative FACS

plots of KP (lung adenocarcinoma) sender cells expressing either sGFP or sGFP-sHaloTag

incubated with AF660-conjugated HaloTag ligand for 5 minutes on ice. AF660 specifically

labeled sHaloTag-GFP sender cells but not sGFP sender cells.

b. Representative histogram of GFP and AF660 intensity in αGFP 293 receiver cells cultured

with HaloTag-GFP sender cells at a 1:1 ratio for 6 hours (complete experiment results for Figure

2d). Receivers were defined as $\text{mCherry}^{neg}\text{PI}^{neg}\text{BFP}^{pos}$ cells.

c. AF660 transfer to αGFP 293 receiver cell is rapid after cell-cell interaction. AF660 MFI shift

was detected after mixing sHalo-GFP senders and αGFP receivers and co-culture for 10 minutes.

AF660 MFI shift was more dramatic than GFP. Receivers were defined as

mCherry^{neg}PI^{neg}BFP^{pos} cells.

d. Slower AF660 quenching in touched receiver cells after removing sHalo-GFP senders. After 6

hours co-culture, GFP/AF660 positive receiver cells were purified via FACS. Analysis of

GFP/AF660 MFI in purified receiver cells showed rapid GFP degradation but slower AF660

quenching. Receivers were defined as mCherry^{neg}PI^{neg}BFP^{pos} cells.

e. Representative FACS plots of αGCN4 receiver cells co-cultured with the indicated sender

cells at 1:1 (for single sender cell) or 1:1:1 (for dual sender cells) ratios. Percent of labeled

receiver cells (gated as mCherry^{neg}PI^{neg}BFP^{pos}) after 6 hours of co-culture is indicated as mean

+/- SD of triplicate cultures.

f. Loading of sender cells with ssDNA. Representative FACS plots of 293 sender cells

expressing either sGCN4 or sGCN4-Halo were loaded with 5' HaloTag-ligand, 3' biotin dual

conjugated ssDNA (21nt), then stained with Avidin-AF647. AF647 specifically labeled loaded

sGCN4-Halo sender cells but not sGCN4 sender cells.

g. Representative histogram of AF647 intensity in control or αGCN4 293 receiver cells cultured

with sGCN4 or sGCN4-Halo sender cells at a 1:1 ratio for 6 hours (complete experiment results

for Figure 2h). Receivers were defined as mCherry^{neg}PI^{neg}BFP^{pos} cells.

Supplementary Figure 8. Diverse primary mouse cells expressing sGFP can transfer GFP

to αGFP 293 receivers.

Different primary sender cells were first sorted based on their cell surface markers, then

transduced with lentiviral vectors expressing sGFP before being co-cultured with αGFP 293

receiver cells for 24 hours. Percentage of $PI^{neg}mCherry^{neg}BFP^{pos}GFP^{pos}$ receiver cells was

assessed by flow cytometry.

For each primary sender, approximate $sGFP^{pos}$ sender : αGFP^{pos} receiver ratio is indicated: a.

Lung epithelial cells (1:180) (sorted $EpCam^{pos}$ cells from dissociated adult lung); b. Kidney

epithelial cells (1:20) (sorted $EpCam^{pos}$ cells from dissociated adult kidney); c. Splenocyte

(1:20); d. Cardiomyocyte (1:5) (sorted $Sirpa^{pos}$ cells from dissociated adult heart). Percent of

labeled cells is indicated as mean +/- SD of triplicate cultures.

Supplementary Figure 9. G-baToN can detect neuron-astrocyte, neuron-cancer cell

interactions.

a. Split channels for Figure 3g. Scale bars = 50 μm.

b. Quantification of Figure 3g using images from 10 different fields. Each dot represents a field.

The bar indicates the mean +/- SD. GFP^{pos} neurons were defined as $Map2^{pos}$ cells with GFP.

c. Representative image of neuron and cancer cell interactions. Scale bars = 50 μm.

d. Quantification of supplementary Figure 9c using images from 10 different fields. Each dot

represents a field. The bar indicates the mean +/- SD. GFP^{pos} neurons were defined as

$Map2^{pos}tdTomato^{neg}$ cells with GFP.

Supplementary Figure 10. G-baToN can detect CD4 T cell-A549 cancer cell interactions.

a. Representative image of A549 cell and CD4 T cell interactions. Scale bars = 10 μm.

b. Primary human $CD4^{pos}$ T cells expressing sGFP can transfer GFP to αGFP A549 cells after

co-culture at a 1:1 ratio for 24 hours. Receivers were defined as

$mCherry^{neg}NearIR^{neg}CD4^{neg}BFP^{pos}$ cells.

ACKNOWLEDGEMENTS

We thank the Stanford Shared FACS and Cell Sciences Imaging Facilities for technical

support; A. Orantes for administrative support, N. Kipniss and Y.T. Shue for critical reagents; D.

Feldser, J. Sage, Y. Chien, N. Kipniss and members of the Winslow laboratory for helpful

comments. R.T. was supported by a Stanford University School of Medicine Dean's Postdoctoral

Fellowship and a TRDRP Postdoctoral fellowship (27FT-0044). C.W.M. was supported by the

NSF Graduate Research Fellowship Program and an Anne T. and Robert M. Bass Stanford

Graduate Fellowship. This work was supported by NIH R01-CA175336 (to M.M.W), NIH R01-

CA207133 (to M.M.W) and NIH R01-CA230919 (to M.M.W) and in part by the Stanford

Cancer Institute support grant (NIH P30-CA124435), 1S10OD01227601 from the National

Center for Research Resources (NCRR) and S10RR027431-01 from National Institutes of Health

(NIH).

METHODS

Cells, Plasmids and Reagents:

HEK-293T, B16-F10, A549, H460 and HUVEC cells were originally purchased from ATCC;

HUASMC were purchased from PromoCell (C-12500); H82 cells were kindly provided by Julien

Sage (Stanford School of Medicine); KP (238N1) and KPT (2985T2) lung adenocarcinoma cells

were generated in the Winslow Lab. HEK-293T, 238N1, 2985T2 and B16-F10 cells were

cultured in DMEM containing 10% FBS, 100 units/mL penicillin and 100 μg/mL streptomycin.

A549, H460 and H82 cells were cultured in RPMI1640 media containing 10% FBS, 100

units/mL penicillin and 100 μg/mL streptomycin. HUVECs were cultured in Vascular Cell Basal

Medium (ATCC, PCS-100-030) with Endothelial Cell Growth Kit (ATCC, PCS-100-041);

HUASMC were cultured in Smooth Muscle Cell Growth Medium 2 (PromoCell, C-22062). All

cell lines were confirmed to be mycoplasma negative (MycoAlert Detection Kit, Lonza).

Pitstop (ab120687) and Dyngo 4a (ab120689) were purchased from Abcam.

All plasmids used in this study are listed in Supplementary Table 1 and will be available on

Addgene.

Antibodies:

Anti-GFP antibody was purchased from MyBioSource (MBS560494), anti-RFP antibody was

purchased from Rockland (600-401-379), anti-human mitochondria antibody was purchased

from Abcam (ab92824), anti-GAPDH antibody was purchased from Cell Signaling Technology

(5174S)

Lentiviral Vector Packaging:

Lentiviral vectors were produced using polyethylenimine (PEI)-based transfection of 293T cells

with the plasmids indicated in Supplementary Table 1, along with delta8.2 and VSV-G

packaging plasmids in 150-mm cell culture plates. Sodium butyrate (Sigma Aldrich, B5887) was

added 8 hours after transfection to achieve a final concentration of 20 mM. Medium was

refreshed 24 hours after transfection. 20 mL of virus-containing supernatant was collected 36,

48, and 60 hours after transfection. The three collections were then pooled and concentrated by

ultracentrifugation (25,000 rpm for 1.5 hours), resuspended overnight in 100 µL PBS, then

frozen at -80°C.

Generation of Stable Cell Lines

Parental cells were seeded at 50% confluency in a 6-well plate the day before transduction (day

0). The cell culture medium was replaced with 2 mL fresh medium containing 8 µg/mL

hexadimethrine bromide (Sigma Aldrich, H9268-5G), 20 µL ViralPlus Transduction Enhancer

(Applied Biological Materials Inc., G698) and 40 µL concentrated lentivirus and cultured

overnight (Day 1). The medium was then replaced with complete medium and cultured for

another 24 hours (Day 2). Cells were transferred into a 100 mm cell culture dish with appropriate

amounts of puromycin (Dose used: 293T: 2 µg/mL; 238N1: 3 µg/mL; 2985T2: 2 µg/mL) and

selected for 48 hours (Day 3). After selection, FACS analysis was performed using fluorescent

markers indicated in Supplementary Table 2 for validation of selection efficiency.

Transwell Co-culture Assay

The Corning® Transwell® polycarbonate membrane cell culture inserts were purchased from

Corning Inc (3422 : CS, Corning, NY). sGFP sender cells were seeded in the upper chamber

inserts of the transwell (1×10^5/insert). The inserts were then placed back into the plate pre-

seeded with 1×10^5/well αGFP receiver cells and cultured in a humidified incubator at 37 °C,

with 5% CO_2 for 24 hours. sGFP sender and αGFP receiver cells co-cultured in the same plate

under the same conditions were used as control. After 24 hours, the upper chamber inserts were

removed, cells in the lower chamber were trypsinized and analyzed by flow cytometry.

Live and Fixed Cell Imaging

For live cell microscopy, 2×10^4 sGFP sender and 2×10^4 αGFP receiver cells were seeded into

35-mm FluoroDish Cell Culture Dishes (World Precision Instruments, FD35-100) and

immediately imaged under a DeltaVision OMX (GE Healthcare) microscope with a 60x oil

objective lens (Olympus) in a humidified chamber at 37 °C with 5% CO_2. One image was taken

per minute for three hours. Images were collected with a cooled back-thinned EM-CCD camera

(Evolve; Photometrics).

For fixed cell microscopy, sender and receiver cells were seeded at the ratios indicated in

Supplementary Table 2 with a total number of 1×10^5 cells onto Neuvitro coated cover slips

(Thermo Fisher Scientific, NC0301187) in a 12-well cell culture plate. 24 hours after co-culture,

cells were fixed in 4% paraformaldehyde (PFA) PBS solution at room temperature for 10

minutes and washed with PBS and distilled water three times each, before mounting onto slides

using 50% glycerol. Images were captured using a Leica DMI6000B inverted microscope with

an 40x oil objective lens. For quantification, GFP-containing receiver cells were counted.

Multiple coverslips were analyzed across independent experiments (n=10).

Western Blot

5×10^6 sGFP sender and 5×10^6 αGFP receiver cells were co-cultured in a 100mm cell culture dish

for 24 hours. Cells were trypsinized, resuspended in FACS buffer containing propidium iodide

(PI) (PBS, 2% FBS, 1 mM EDTA, and 1.5 µM PI). tdTomato^{neg}PI^{neg} cells were sorted and lysed

in RIPA buffer (50 mM Tris-HCl (pH 7.4), 150 mM NaCl, 1% Nonidet P-40, and 0.1% SDS)

and incubated at 4 °C with continuous rotation for 30 minutes, followed by centrifugation at

$12,000 \times$ rcf for 10 minutes. The supernatant was collected, and the protein concentration was

determined by BCA assay (Thermo Fisher Scientific, 23250). Protein extracts (20–50 μg) were

dissolved in 10% SDS-PAGE and transferred onto PVDF membranes. The membranes were

blocked with 5% non-fat milk in TBS with 0.1% Tween 20 (TBST) at room temperature for one

hour, followed by incubation with primary antibodies diluted in TBST (1:1000 for anti-GFP,

anti-Tomato (RFP) and anti-human mitochondria (hu-Mito), 1:5000 for anti-GAPDH) at 4 °C

overnight. After three 10-minutes washes with TBST, the membranes were incubated with the

appropriate secondary antibody conjugated to HRP diluted in TBST (1:10000) at room

temperature for 1 hour. After three 10-minutes washes with TBST, Protein expression was

quantified with enhanced chemiluminescence reagents (Fisher Scientific, PI80196).

GFP/AF660 Stability

To assess the stability of GFP and AF660 in touched receiver cells, 1×10^7 sGFP or sHalo-GFP

sender cells were co-cultured with 1×10^7 αGFP receiver cells in a 150mm cell culture dish for 6

hours. Cells were then trypsinized and resuspended in FACS buffer containing propidium iodide.

mCherry^{neg}PI^{neg}BFP^{pos} cells were sorted and 1×10^5 cells were re-plated in 12-well plate and

cultured for 2, 4, 6, 8, 12, 24 and 48 hours in DMEM containing 10% FBS, 100 units/mL

penicillin and 100 μg/mL streptomycin. GFP or AF660 intensity was assessed via FACS analysis

of mCherry^{neg}PI^{neg}BFP^{pos} cells and shown as Mean +/- SD of GFP/AF660 MFI in triplicate

cultures.

Puromycin Resistant Protein Transfer Assay

To validate puromycin resistant protein (GCN5-Related N-Acetyltransferases, PuroR) function

in sender cells, 5×10^6 HEK-293T cells were transfected with sGFP-PuroR-PDGFR, sGFP-

PDGFR or sGFP-PDGFR-IRES-PuroR in a 100-mm cell culture dish for 12 hours before re-

plating into a 12-well plate (1×10^5 cells/well). 24 hours after transfection, cells were treated with

1, 2 or 5 μg/mL puromycin for 24 hours. After treatment, dead cells were washed away, and

viable cells were fixed by 4% PFA and stained with crystal violet. Stained cells were washed

with distilled water three times before extraction with 500 µL of methanol for 15 minutes on a

rocker. The absorbance was measured at 570 nm with a spectrophotometer.

To count the number of viable receiver cells co-cultured with sGFP-PuroR sender cells, 2×10^5

αGFP receiver cells were co-cultured with 8×10^5 sGFP or sGFP-PuroR sender cells in a 6-well

plate. 24 hours after co-culture, cells were treated with 2 µg/mL puromycin for 48 hours. Viable

tdTomato^{neg}PI^{neg}BFP^{pos} receiver cells were counted via FACS.

Primary Mouse Cell Isolation

Mouse (C57BL/6J, The Jackson Laboratory) lung, kidney, heart, hindlimb skeleton muscle,

spleen and liver tissue were dissected, cut into small pieces and digested in 5 mL tissue digest

media (3.5 mL HBSS-Ca2+ free, 0.5 mL Trypsin-EDTA (0.25%), 5mg Collagenase IV

(Worthington), 25 U Dispase (Corning) for 30 minutes in hybridization chamber at 37˚C with

rotation. Digestion is then neutralized by adding 5 mL ice cold Quench Solution (4.5 mL L15

media, 0.5 mL FBS, 94 µg DNase). Single cell suspensions were generated by filtering through a

40uM cell strainer, spinning down at 500 rcf for 5 minutes and washed with PBS twice

For primary mouse lung epithelial cells, kidney epithelial cells and cardiomyocyte isolation and

culture, the single cell pellets were resuspended in 1mL FACS buffer containing 1:300 dilution

of anti-EpCam-AF467 (Biolegend, 118211) (for lung and kidney epithelial cell) or anti-Sirpa-

AF467 (Biolegend, 144027) (for cardiomyocyte) antibody and incubated on ice for 20 minutes

before FACS sorting. DAPI^{neg}EpCam^{pos} or DAPI^{neg}Sirpa^{pos} cells were sorted and seeded onto a

100 mm culture dish precoated with 5 µg/cm2 Bovine Plasma Fibronectin (ScienCell, 8248).

For primary skeleton muscle cell, splenocyte and hepatocyte culture, the single cell pellets were

resuspended in DMEM containing 20% FBS, 200 units/mL penicillin and 200 μg/mL

streptomycin, amphotericin and cultured in 100-mm culture dish at 37^{o}C for 1 hour to remove

fibroblast cells. The supernatant containing primary skeletal muscle cells, splenocytes and

hepatocytes was then transferred into a new 100-mm culture dish precoated with 5 µg/cm2

Bovine Plasma Fibronectin (ScienCell, 8248).

Conjugation of HaloTag Ligand to Oligonucleotides

Oligonucleotides to be conjugated with the HaloTag ligand were synthesized with a 5' amino

modifier C12 and a 3' biotin group (IDT). Oligonucleotides were initially ethanol-precipitated

and subsequently resuspended to 1 mM in conjugation buffer (100 mM Na2HPO4 (Sigma-

Aldrich S9763), 150 mM NaCl (Thermo Fisher Scientific S271), pH 8.5). Resuspended oligos

were combined with an equal volume of the HaloTag ligand succinimidyl ester (O4) (Promega

P6751) resuspended in N,N- dimethylformamide (Sigma-Aldrich D4551) with a 30-fold molar

excess of the ligand. Conjugation reactions were conducted overnight at room temperature with

constant agitation prior to final cleanup via ethanol precipitation.

Loading of HaloTag-Expressing Sender Cells

Prior to loading with HaloTag-conjugated elements, sender cells were washed once in cold PBS

following detachment and subsequently resuspended in cold Cell Staining Buffer (BioLegend

42021). For loading of HaloTag-conjugated fluorophores, senders were stained at a density of

1.00E+07 cells/mL on ice for five minutes in the presence of either 1 µM HaloTag-Alexa Fluor

488 (Promega G1001) or 3.5 µM HaloTag-Alexa Fluor 660 (Promega G8471). Stained sender

cells were then washed twice in Cell Staining Buffer (500 rcf for 5 min at 4˚C) prior to

resuspension in growth media in preparation for co-culture.

For loading with HaloTag-conjugated oligonucleotides, sender cells were initially resuspended

and incubated with 100 µg/mL salmon sperm DNA (Thermo Fisher Scientific 15632011).

Sender cells were then incubated with 3.5 µM HaloTag-conjugated oligonucleotides

(5AmMC12/TCTAGGCGCCCGGAATTAGAT/3Bio) and subsequently washed once.

Oligonucleotide-loaded sender cells were then stained with 5 µg/mL streptavidin-conjugated

Alexa Fluor 647 (Thermo Fisher Scientific S32357) for 30 minutes on ice. The loaded, stained

senders were then washed twice and resuspended in growth media in preparation for co-culture.

Primary Neuron and Astrocyte Cultures

Primary cortical neurons were dissociated from mouse (C57BL/6J, The Jackson Laboratory)

E16.5 embryonic cortices into single cell suspensions with a papain dissociation system

(Worthington Biochemical Corporation). Tissue culture plates were coated with poly-L-lysine

(0.1% w/v) before seeding cells. Neurons were grown in Neurobasal media (Gibco)

supplemented with B-27 serum free supplement (Gibco), GlutaMAX (Gibco), and penicillin-

streptomycin (Gibco) in a humidified incubator at 37 °C, with 5% CO2. Half media changes

were performed every 4-5 days. Primary astrocytes were dissociated from P0-P1 mouse cortices

using the same papain dissociation methods as neurons, except the single cell suspensions were

then plated onto tissue culture plates without poly-L-lysine in DMEM with 10% FBS and

penicillin-streptomycin. Primary astrocyte cultures were passaged using Accutase (Stemcell

Technologies).

Primary Human T Cell Cultures

Blood from healthy donors collected in leukoreduction system (LRS) chambers was separated by

Ficoll-Paque density gradient to obtain peripheral blood mononuclear cells (PBMCs). CD4[pos]

and CD8[pos] T cells were isolated by negative selection using EasySep Human CD4+ T Cell

Isolation Kit and EasySep Human CD8+ T Cell Isolation Kit (STEMCELL Technologies),

respectively, according to the manufacturer's instructions. T cells were cultured for 3 days with

CD3/CD28 Dynabeads (ThermoFisher Scientific) with 40 IU/mL IL-2 and spinoculated with

lentivirus for 2 hours at 400 rcf in the presence of 8 µg/mL polybrene. T cells were expanded

following transduction for two days in the presence of CD3/CD28 Dynabeads and 300 IU/mL

IL-2 prior to use in assays. Transduced or untransduced CD4[pos] or CD8[pos] T cells were co-

cultured for 24 hours together with A549 cells. Following co-culture, cells were harvested,

stained with antibodies against CD45, CD4, or CD8 (BioLegend) and analyzed on a

LSRFortessa flow cytometer (BD Biosciences). Each condition was run in triplicate, and two

independent experiments were conducted using T cells from different donors. For microscopy,

A549 cells were co-cultured with CD4[pos] T cells in glass-bottom plates (MatTek Corporation)

and imaged on an LSM 700 confocal microscope (Zeiss).

REFERENCES

1.   Zhang, Y. & Liu, F. Multidimensional Single-Cell Analyses in Organ Development and Maintenance. *Trends in cell biology* (2019).
2.   Tsioris, K., Torres, A.J., Douce, T.B. & Love, J.C. A new toolbox for assessing single cells. *Annual review of chemical and biomolecular engineering* **5**, 455-477 (2014).
3.   Zhong, S. et al. A single-cell RNA-seq survey of the developmental landscape of the human prefrontal cortex. *Nature* **555**, 524 (2018).
4.   Zhou, F. et al. Reconstituting the transcriptome and DNA methylome landscapes of human implantation. *Nature* **572**, 660-664 (2019).
5.   Boisset, J.-C. et al. Mapping the physical network of cellular interactions. *Nature methods* **15**, 547-553 (2018).
6.   Morsut, L. et al. Engineering customized cell sensing and response behaviors using synthetic notch receptors. *Cell* **164**, 780-791 (2016).
7.   Pasqual, G. et al. Monitoring T cell–dendritic cell interactions in vivo by intercellular enzymatic labelling. *Nature* **553**, 496 (2018).
8.   Ovcinnikovs, V. et al. CTLA-4–mediated transendocytosis of costimulatory molecules primarily targets migratory dendritic cells. *Science immunology* **4**, eaaw0902 (2019).
9.   Langridge, P.D. & Struhl, G. Epsin-dependent ligand endocytosis activates Notch by force. *Cell* **171**, 1383-1396. e1312 (2017).
10.   Li, G. et al. T cell antigen discovery via trogocytosis. *Nature methods* **16**, 183 (2019).
11.   Fridy, P.C. et al. A robust pipeline for rapid production of versatile nanobody repertoires. *Nature methods* **11**, 1253 (2014).
12.   Lim, K.H., Huang, H., Pralle, A. & Park, S. Stable, high-affinity streptavidin monomer for protein labeling and monovalent biotin detection. *Biotechnology and bioengineering* **110**, 57-67 (2013).
13.   Tanenbaum, M.E., Gilbert, L.A., Qi, L.S., Weissman, J.S. & Vale, R.D. A protein-tagging system for signal amplification in gene expression and fluorescence imaging. *Cell* **159**, 635-646 (2014).
14.   Los, G.V. et al. HaloTag: a novel protein labeling technology for cell imaging and protein analysis. *ACS chemical biology* **3**, 373-382 (2008).

Figure 1. GFP-based Touching Nexus (G-baToN) leads to cell-cell interaction-dependent receiver cell labeling

a. Schematic of the G-baToN system. Surface GFP (sGFP) on a sender cell is transferred to a receiver cell expressing a cell surface anti-GFP nanobody (αGFP) leading to GFP labeling of the "touched" receiver.

b. GFP transfer from sGFP-expressing KPT lung cancer sender cells (marked by intracellular tdTomato) to αGFP-expressing 293 receiver cells. Receiver cell labeling is sGFP- and αGFP- dependent. Control sender cells do not express sGFP. Control receiver cells do not express αGFP. Cytoplasmic GFP (Cyto-GFP) is not transferred to receiver cells. Sender and receiver cells were seeded at a 1:1 ratio and co-cultured for 24 hours. Receivers were defined as $Tomato^{neg}PI^{neg}$ cells.

c. GFP transfer to 293 receiver cells requires direct cell-cell contact. Receiver cells separated from sender cells by a transwell chamber are not labeled. Sender and receiver cells were seeded in upper and lower chambers respectively at a 1:1 ratio and cultured for 24 hours. Receivers were defined as $Tomato^{neg}PI^{neg}$ cells.

d. GFP transfer to 293 receiver cells requires sGFP-αGFP interaction and is blocked by anti-GFP antibody in a dose-dependent manner. sGFP sender cells were pre-incubated with the indicated concentration of anti-GFP antibody for 2 hours, washed with PBS, and then co-cultured with receiver cells at a 1:1 ratio for 24 hours. Receivers were defined as $Tomato^{neg}PI^{neg}$ cells.

e. Time-lapse imaging of GFP transfer from a sGFP-expressing sender cell to an αGFP-expressing receiver cell. Time after contact is indicated. Receiver cell is outlined with white dashed line. Scale bar: 10 μm.

f. Analysis of GFP Mean Fluorescence Intensity (MFI) of αGFP receiver cells (marked by intracellular BFP) co-cultured with sGFP sender cells (marked by intracellular tdTomato) co-cultured for the indicated amount of time. Sender and receiver cells were seeded at a 1:1 ratio. Receivers were defined as $Tomato^{neg}PI^{neg}BFP^{pos}$ cells.

g. Percentage of labeled αGFP receiver cells after co-culture with different numbers of sender cells for 24 hours. Receivers were defined as $Tomato^{neg}PI^{neg}BFP^{pos}$ cells.

h. Detection of rare labeled αGFP receiver cells after co-culture with sGFP sender cells at approximately a $1:10^5$ ratio for 24 hours. Receivers were defined as $Tomato^{neg}PI^{neg}BFP^{pos}$ cells.

	synNotch (Morsut *et al.*, Cell, 2016)	**LIPSTIC** (Pasqual *et al.*, Nature, 2018)	**G-baToN** (This manuscript)
Cell-cell interaction	Sender Cell Surface GFP αGFP-nanobody Receiver Cell	Sender Cell CD40 with SorA G5 ●-LPETG CD40L Receiver Cell	Sender Cell Surface GFP αGFP-nanobody Receiver Cell
Receiver Cell after interaction	Reporter Receiver Cell	G5-TEPL-● CD40L Receiver Cell	Receiver Cell
Basis of the labeling	Notch intracellular cleavage leads to transcription factor release and activation of a reporter	Covalent labeling of receptor by ligand associated enzyme	Ligand internalization
Additional requirement for labeling	Engineered reporter in receiver cells	Labeling substrate	None
Determinants of label half-life after touching	Transcription factor half-life and reporter mRNA and protein half-life in the receiver cell	Labeled protein half-life on the receiver cell	Half-life of transferred protein or fluorophore in the receiver cell
Advantages	Generic sender and receivers Generalizable across cell type Compatable with AND gates Tf can drive expression of many genes of interest	Requirement for exogenous substrate adds temporal control of labelling Conceptually generalizable to other receptor ligand pairs	Simple two-component system Low background Single cell level sensitivity Generalizable across cell type Generic sender and receivers Complex cell-cell interactions
Potential disadvantages	Engineered ligand-receptor interactions could influence cell-cell contact Background activation or reporter could be leaky	Requires knowledge of receptor-ligand interactions between cell types of interest	Engineered ligand-receptor interactions could influence cell-cell contact

Supplementary Figure 1. Features of the synNotch, LIPSTIC, and G-BaToN cell-cell interaction reporter systems.
Overview of systems that enable labeling of cells after cell-cell contact.

Supplementary Figure 2. GFP transfer requires direct GFP-αGFP interaction.

a. GFP protein is transferred to receiver cells. Western blot analysis of FACS purified Control and αGFP receiver 293 cells cultured in isolation or co-culture with sGFP Tomato-positive sender cells (mouse KPT lung cancer cells) for 24 hours. sGFP but not Tomato is present in touched αGFP receiver cells. Human mitochondrial antigen (hu-Mito) is a marker for receiver cells. GAPDH shows loading. Rightmost lane is sGFP sender cells.

b. GFP cannot be transferred from fixed sGFP sender cells to live αGFP receiver cells. sGFP sender cells were fixed in 1% PFA for 5 minutes and washed with PBS before co-cultured with αGFP receiver cells at a 1:1 ratio for 24 hours. Receivers were defined as $Tomato^{neg}PI^{neg}BFP^{pos}$ cells.

c. GFP transfer to 293 receiver cells required sGFP-αGFP recognition. GFP is transferred from sGFP sender cells to αGFP-receiver cells but not from sGFP sender cells to αmCherry-receiver cells. Control receiver cells do not express any nanobody. Sender and receiver cells were co-cultured at a 1:1 ratio for 24 hours. Receivers were defined as $Tomato^{neg}PI^{neg}BFP^{pos}$ cells.

d. GFP transfer to receiver cells is accompanied by a reduction of GFP on the sender cells. GFP expression on sender cells after 24 hour co-culture with control or αGFP-expressing 293 receiver cells at a 1:1 ratio. Co-culture with αGFP expressing but not control receiver cells reduced GFP on sGFP sender cells. Senders were defined as $Tomato^{pos}DAPI^{neg}BFP^{neg}$ cells.

e. GFP transfer is accompanied with αGFP internalization on receiver cells. Analysis of surface αGFP (Myc-tag) on 293 receiver cell co-cultured for 24 hours with sGFP sender cells. Receivers were defined as $Tomato^{neg}PI^{neg}$ cells.

f. GFP transfer to 293 receiver cells is partially dependent on membrane dynamics of endocytosis. Both a clathrin inhibitor (Pitstop, 20 μM) and a dynamin inhibitor (Dyngo 4a, 10 μM) partially inhibit GFP transfer from sGFP sender cells to αGFP receivers. Inhibitors were present during the 24 hour co-culture. Receivers were defined as $Tomato^{neg}PI^{neg}$ cells.

g. GFP transfer to αGFP 293 receiver cells can be very rapid. A shift in GFP MFI was detected 5 minutes after mixing sGFP sender cells with αGFP receiver cell. Receivers were defined as $Tomato^{neg}PI^{neg}BFP^{pos}$ cells. MFI mean +/- SD of triplicate cultures is shown.

h. Rapid GFP degradation in touched receiver cells after removal of the sGFP sender cells. After 6 hours co-culture, GFP positive receiver cells were purified by FACS followed by culture without sender cells. Receivers were defined as $mCherry^{neg}PI^{neg}BFP^{pos}$ cells. GFP MFI in receiver cells reduced rapidly ($T_{1/2}$ approximately 2 hours). MFI mean +/- SD of triplicate cultures is shown.

Supplementary Figure 3. Transmembrane domains and the nanobody affinity impact sGFP transfer and receiver cell labeling.
a. Schematic of the sender and receiver cells used to determine the impact of different αGFP transmembrane (TM) domains. TM domains contain the TM domain itself as well as membrane proximal regions from the indicated mouse (m) and human (h) proteins.
b. Different TM domains impact cell surface αGFP expression on 293 receiver cells. Membrane αGFP was accessed by anti-Myc staining. Control receiver cells do not express any nanobody. Mean +/- SD of Myc MFI in triplicate cultures is shown.
c. VEGFR2 TM domain on αGFP receivers enables highest GFP transfer efficiency. Receiver cells expressing αGFP linked by different TM domains were co-cultured with sGFP sender cells at a 1:1 ratio for 6 hours. Receivers were defined as Tomato^negPI^neg cells.
d. Schematic of the sender and receiver cells used to determine the impact of different sGFP TM domains and different αGFP nanobodies on G-baTon-based labeling. TM domains contain the TM domain itself as well as membrane proximal regions from the indicated mouse (m) and human (h) proteins.
e. Different nanobodies exhibit different levels of expression on 293 receiver cells. Total αGFP expression was assessed by BFP intensity. Mean +/- SD of GFP MFI in triplicate cultures is shown.
f. αGFP affinity influences transfer of GFP to touched 293 receiver cells. Receiver cells expressing different αGFP nanobodies were co-cultured with sGFP sender cells at a 1:1 ratio for 6 hours. GFP transfer was assessed by flow cytometry. GFP intensity on Tomato^negPI^negBFP^pos receiver cells is shown as mean +/- SD of triplicate cultures.
g. Different TM domains on sGFP impact its expression in 293 sender cells. sGFP expression in sender cells was assessed by flow cytometry for GFP. Mean +/- SD of GFP MFI in triplicate cultures is shown.
h. PDGFR TM domain on sGFP minimized retrograde transfer of αGFP from receiver cells to 293 sGFP sender cells. αGFP transfer to sGFP senders were determined as the percentage of mCherry^posGFP^pos sender cells that were also BFP^pos. Cells were co-cultured for 6 hours at a 1:1 ratio. Mean +/- SD of triplicate cultures is shown.

Figure 2. X-baToN systems enable recording of multicellular interactions and cell-cell interaction-based cargo transfer.
a. Schematic of the AND gate-baToN system. sGFP and smCherry sender cells express either sGFP or mCherry. Dual receiver cells express both αGFP (LaG17, tagged by Myc-tag) and αmCherry (LaM4, tagged by intracellular BFP).
b. Representative FACS plots of dual receiver 293 cells cultured with the indicated 293 sender cells at 1:1 (for single sender cell) or 1:1:1 (for dual sender cells) ratios. Percent of labeled receiver cells (gated as BFPpos) after 24 hours of co-culture is indicated as mean +/- SD of triplicate cultures.
c. Diagram of HaloTag-baToN system. Sender cells (marked by intracellular 2A-mCherry) express surface HaloTag-GFP fusion which can be loaded with HaloTag ligands (in this example AF660). Receiver cells express αGFP (LaG17, tagged by intracellular BFP).
d. Representative plot of GFP and AF660 intensity in αGFP 293 receiver cells co-cultured with HaloTag-GFP KP (lung adenocarcinoma) sender cells at a 1:1 ratio for 6 hours. Receivers were defined as mCherrynegPInegBFPpos cells.
e. Diagram of dual color GCN4-HaloTag-baToN system. Sender cells (marked by intracellular 2A-mCherry) express surface 4XGCN4 associated with HaloTag, loaded with either AF488- or AF660- conjugated HaloTag ligand. Receiver cells express αGCN4 (tagged by intracellular BFP).
f. AF488/AF660 GCN4-HaloTag sender ratio in the co-culture directly proportional to AF488/AF660 intensity (MFI) of αGCN4 receiver after 6 hours of co-culture. Receivers were defined as mCherrynegPInegBFPpos cells.
g. Diagram of using GCN4-HaloTag sender to transfer ssDNA into αGCN4 receiver cells. Sender cells (marked by intracellular 2A-mCherry) express surface 4XGCN4 associated with HaloTag, loaded with 5' HaloTag ligand, 3' biotin dual conjugated ssDNA (21 nt), then stained with Avidin-AF647. Receiver cells express αGCN4 (tagged by intracellular BFP).
h. Representative plot of AF647 intensity in αGCN4 293 receiver cells co-cultured with GCN4-HaloTag 293 sender cells at a 1:1 ratio for 6 hours. Receivers were defined as mCherrynegPInegBFPpos cells.

Supplementary Figure 4. X-baToN systems enable fluorescent labeling via various antigen-nanobody/scFV pairs.
a. Schematic of the surface BFP (sBFP)-baToN system. Sender cell (marked by intracellular tdTomato) expresses sBFP and receiver cell expresses α GFP.
b. The cross reactivity of αGFP with BFP allow BFP to be transferred from sBFP sender to αGFP receiver cells. sBFP senders and αGFP receivers were co-cultured at 1:1 ratio for 24 hours. Receivers were defined as toTomato^{neg}PI^{neg} cells. BFP MFI of receiver cells is shown as mean +/- SD of triplicate cultures.
c. Schematic of the surface mCherry (smCherry)-baToN system. Sender cell (marked by intracellular GFP) expresses smCherry and receiver cell express-es αmCherry (tagged by intracellular BFP)
d. The pairing between mCherry and αmCherry enable mCherry transfer from smCherry 293 Sender cells to αmCherry-expressing 293 Receiver cells. smCherry senders are not able to transfer mCherry to αGFP receiver cells. smCherry sender cells were co-cultured with αGFP or αmCherry receiver cells at a 1:1 ratio for 24 hours. Receivers were defined as GFP^{neg}BFP^{pos} cells. mCherry MFI of receiver cell is shown as mean +/- SD of triplicate cultures. smCherry 293 sender is not able to transfer mCherry to αGFP 293 receiver.
e. Schematic of the surface GCN4 (sGCN4)-baToN system. Sender cells (marked by intracellular 2A-mCherry) express cell surface 4X GCN4 peptide fused with GFP. Receiver cells express a cell surface anti-GCN4 single-chain variable fragment (scFV; αGCN4, tagged by intracellular BFP).
f. The pairing between GCN4 and αGCN4 enables co-transfer of GFP from 4X sGCN4-GFP senders to αGCN4 receivers 24 hours after co-culture at a 1:1 ratio. Receivers were defined as mCherry^{neg}PI^{neg}BFP^{pos} cells. GFP MFI of receiver cell is shown as mean +/- SD of triplicate cultures. sGFP sender are not able to transfer GFP to αGCN4 receivers.

Tang *et al.*

Supplementary Figure 5. Dual color-baToN systems enable labeling in complex cell-cell interaction systems.

a. Diagram of the Reciprocal-baToN system. Cell A expresses sGFP and αmCherry (tagged by intracellular BFP), Cell B expresses smCherry and α GFP (tagged by Myc-tag).

b. Representative FACS plots of cell A and cell B monocultures (left 2 panels) and after co-culture at a 5:1 ratio for 24 hours. Percent of labeled cells is indicated as mean +/- SD of triplicate cultures.

c. Representative image of reciprocal labeling of cell A and cell B co-cultured for 24 hours. Cells are outlined with a white dashed line. Cell A is BFP and GFP double positive and Cell B is mCherry positive. Touched cell A is BFP, GFP and mCherry triple positive and touched cell B is GFP and mCherry double positive (pointed out by white letter). Scale bar: 10 μm.

d. Representative image of dual receiver cells co-cultured with indicated sender cells. Cells are outlined with a white dashed line. Dual receiver cells are BFP positive and sender touched dual receiver cells are pointed out by white arrows. Scale bar: 10 μm.

e. Diagram of the BFP/GFP AND gate-baToN system. sBFP sender cells express intracellular Tomato and surface BFP, sGFP sender cells express intracellular Tomato and surface GFP. Common receiver cells expressed αGFP.

f. Representative FACS plots of common receiver 293 cells cultured with the indicated Tomato^{pos} sender cells at 1:1 (for single sender cell) or 1:1:1 (for dual sender cells) ratios. Receiver cells were gated as Tomato^{neg}PI^{neg}. Percent of labeled common receiver cells after 24 hours of co-culture is indicated as mean +/- SD of triplicate cultures.

Supplementary Figure 6. Using the G-baToN system to co-transfer cargo protein into touched receiver cells.
a. Diagram of surface tdTomato-GFP (sGFP-Tom) co-transfer into touched receiver cells. Sender cell expresses sGFP-tdTomato and receiver cell expresses αGFP (tagged by intracellular BFP).
b. Representative FACS plots of αGFP 293 receiver cells cultured with the indicated sender cells at 1:1 ratio for 24 hours. Percent of GFP/Tomato dual labeled receiver cells is indicated as mean +/- SD of BFP^{pos} cells from triplicate cultures.
c. Diagram of GFP-PuroR co-transfer system. Sender cell expresses surface GFP associated with PuroR (marked by intracellular tdTomato), receiver cell expresses αGFP (tagged by intracellular BFP). PuroR: Gcn5-related N-acetyltransferase.
d. 293 Sender cells expressing sGFP-PuroR are only modestly resistant to puromycin treatment. 293 cells were transfected with plasmids as indicated. 24 hours after transfection, 293 cells were treated with different doses of puromycin for 24 hours. Cell number was assessed by measuring spectrophotometrically at 570 nm (OD570) after crystal violet staining.
e. Co-transfer of GFP-PuroR from sGFP-PuroR sender cells to αGFP 293 receiver cells confers puromycin resistance to receiver cells. Control, sGFP or sGFP-PuroR sender cells were co-cultured with αGFP 293 receiver cells for 24 hours at a 4:1 ratio before treatment with different dose of puromycin for 48 hours. tdTomato^{neg}PI^{neg}BFP^{pos} cell numbers were counted via FACS.

Supplementary Figure 7. The Halotag-baToN system enables touching based co-transfer.

a. Labeling HaloTag-expressing sender cells with AF660 fluorophore. Representative FACS plots of KP (lung adenocarcinoma) sender cells expressing either sGFP or sGFP-sHaloTag incubated with AF660-conjugated HaloTag ligand for 5 minutes on ice. AF660 specifically labeled sHaloTag-GFP sender cells but not sGFP sender cells.

b. Representative histogram of GFP and AF660 intensity in αGFP 293 receiver cells cultured with HaloTag-GFP sender cells at a 1:1 ratio for 6 hours (complete experiment results for Figure 2d). Receivers were defined as mCherry^{neg}PI^{neg}BFP^{pos} cells.

c. AF660 transfer to αGFP 293 receiver cell is rapid after cell-cell interaction. AF660 MFI shift was detected after mixing sHalo-GFP senders and αGFP receivers and co-culture for 10 minutes. AF660 MFI shift was more dramatic than GFP. Receivers were defined as mCherry^{neg}PI^{neg}BFP^{pos} cells.

d. Slower AF660 quenching in touched receiver cells after removing sHalo-GFP senders. After 6 hours co-culture, GFP/AF660 positive receiver cells were purified via FACS. Analysis of GFP/AF660 MFI in purified receiver cells showed rapid GFP degradation but slower AF660 quenching. Receivers were defined as mCherry^{neg}PI^{neg}BFP^{pos} cells.

e. Representative FACS plots of αGCN4 receiver cells co-cultured with the indicated sender cells at 1:1 (for single sender cell) or 1:1:1 (for dual sender cells) ratios. Percent of labeled receiver cells (gated as mCherry^{neg}PI^{neg}BFP^{pos}) after 6 hours of co-culture is indicated as mean +/- SD of triplicate cultures.

f. Loading of sender cells with ssDNA. Representative FACS plots of 293 sender cells expressing either sGCN4 or sGCN4-Halo were loaded with 5' HaloTag-ligand, 3' biotin dual conjugated ssDNA (21nt), then stained with Avidin-AF647. AF647 specifically labeled loaded sGCN4-Halo sender cells but not sGCN4 sender cells.

g. Representative histogram of AF647 intensity in control or αGCN4 293 receiver cells cultured with sGCN4 or sGCN4-Halo sender cells at a 1:1 ratio for 6 hours (complete experiment results for Figure 2h). Receivers were defined as mCherry^{neg}PI^{neg}BFP^{pos} cells.

Figure 3. G-baToN is a generalized system that can be used for touching-based labeling between various cell types.
a. Many different cell lines and primary cell types can serve as receiver cells. Receiver cells expressing αGFP were co-cultured with sGFP-expressing KPT sender cells for 24 hours. MFI of GFP on Tomato[neg]PI[neg]BFP[pos] receiver cells was assessed by FACS analysis and is shown as fold change relative to monocultured receiver cells. For lung epithelial cells and kidney epithelial cells MFI of GFP on Tomato[neg]PI[neg]BFP[pos]EpCAM[pos] cells is shown as mean +/- SD of triplicate cultures. SkMC: skeletal muscle cells. Astro: Astrocytes. Hepato: hepatocytes. Spleno: splenocytes.
b. Many different cell lines and primary cell types can serve as sender cells. Sender cells expressing sGFP were co-cultured with αGFP 293 receiver cells for 24 hours. MFI of GFP on mCherry[neg]PI[neg]BFP[pos] receiver cells was assessed by flow cytometry and is shown as fold change relative to monocultured 293 receiver cells. Mean +/- SD of triplicate cultures is shown.
c, d. G-baToN can detect cancer cell-endothelial cell (EC) interactions. HUVECs expressing αGFP were co-cultured with or without Tomato[pos] sGFP-expressing lung cancer sender cells at a 1:1 ratio for 24 hours. MFI of GFP on PI[neg]Tomato[neg]BFP[pos]CD31[pos] Receiver cells was assessed by flow cytometry and is shown as mean +/- SD of triplicate cultures (**c**). Representative images of Tomato[pos] sGFP-expressing lung cancer sender cells co-cultured with either control HUVEC receiver cells (HUVECs expressing BFP) or αGFP HUVEC receiver cells at a 1:1 ratio for 24 hours (**d**). Scale bars = 50 µm.
e,f. G-baToN can detect endothelial cell (EC)-smooth muscle cell (SMC) interactions. Primary human umbilical artery smooth muscle cells (HUASMC) expressing αGFP were co-cultured with or without sGFP-expressing HUVEC senders cells at a 1:1 ratio for 24 hours. MFI of GFP on PI[neg]BFP[pos] receiver cells was assessed by flow cytometry and is shown as mean +/- SD of triplicate cultures (**e**). Representative images of sGFP-expressing HUVEC sender cells co-cultured with either control HUASMC receiver cells (expressing BFP) or αGFP HUASMC receiver cells at a 1:1 ratio for 24 hours (**d**). Scale bars = 50 µm.
g. Representative images of sGFP-expressing astrocyte sender cells co-cultured with either control neuron receivers or αGFP neuron receivers at a 1:2 ratio for 24 hours. Neurons were stained with Microtubule Associated Protein 2 (Map2). Scale bars = 50 µm. Higher magnification of the boxed areas are shown on the right.
h. G-baToN can detect cancer cell-T cell interactions. Primary human CD4[pos] (left) or CD8[pos] (right) T cells were co-cultured with sGFP-expressing lung cancer sender cells (A549 cells) at a 2:1 ratio for 24 hours. Receivers were defined as mCherry[neg]Near-IR[neg]BFP[pos]CD4[pos]/CD8[pos] T cells.

Supplementary Figure 8. Diverse primary mouse cells expressing sGFP can transfer GFP to αGFP 293 receivers.
Different primary sender cells were first sorted based on their cell surface markers, then transduced with lentiviral vectors expressing sGFP before being co-cultured with αGFP 293 receiver cells for 24 hours. Percentage of mCherry[neg]PI[neg]BFP[pos]GFP[pos] receiver cells was assessed by flow cytometry. For each primary sender, approximate sGFP[pos] sender : αGFP[pos] receiver ratio is indicated: **a**. Lung epithelial cells (1:180) (sorted EpCam[pos] cells from dissociated adult lung); **b**. Kidney epithelial cells (1:20) (sorted EpCam[pos] cells from dissociated adult kidney); **c**. Splenocyte (1:20); **d**. Cardiomyocyte (1:5) (sorted Sirpa[pos] cells from dissociated adult heart). Percent of labeled cells is indicated as mean +/- SD of triplicate cultures.

Supplementary Figure 9. G-baToN can detect neuron-astrocyte, neuron-cancer cell interactions.

a. Split channels for Figure 3g. Scale bars = 50 μm.

b. Quantification of Figure 3g using images from 10 different fields. Each dot represents a field. The bar indicates the mean +/- SD. GFP[pos] neurons were defined as Map2[pos] cells with GFP.

c. Representative image of neuron and cancer cell interactions. Scale bars = 50 μm.

d. Quantification of supplementary Figure 9c using images from 10 different fields. Each dot represents a field. The bar indicates the mean +/- SD. GFP[pos] neurons were defined as Map2[pos]tdTomato[neg] cells with GFP.

Supplementary Figure 10. G-baToN can detect CD4 T cell-A549 cancer cell interactions.
a. Representative image of A549 cell and CD4 T cell interactions. Scale bars = 10 µm.
b. Primary human CD4^{pos} T cells expressing sGFP can transfer GFP to αGFP A549 cells after co-culture at a 1:1 ratio for 24 hours. Receivers were defined as mCherry^{neg}NearIR^{neg}CD4^{neg}BFP^{pos} cells.