Reciprocal transplant experiment reveals resolution dependent partial functional redundancy in aquatic microbiome

Kshitij Tandon[1,2,3], Min-Tao Wan[4], Chia-Chin Yang[1], Shan-Hua Yang[5], Bayanmunkh Baatar[1], Chih-Yu Chiu[1], Jeng-Wei Tsai[6], Wen-Cheng Liu[7], Chen Siang Ng[3] and Sen-Lin Tang[1,2]*

[1]Biodiversity Research Center, Academia Sinica, Taipei 11529, Taiwan;

[2]Bioinformatics Program, Institute of Information Science, Taiwan International Graduate Program, Academia Sinica, Taipei 11529, Taiwan;

[3]Institute of Molecular and Cellular Biology, National Tsing Hua University, Hsinchu 300, Taiwan

[4]EcoHealth Microbiology Laboratory, WanYu Co., Ltd., Chiayi 600, Taiwan;

[5]Institute of Fisheries Science, National Taiwan University, Taipei 10617, Taiwan.

[6]China Medical University, Department of Biological Science and Technology, Taichung, Taiwan

[7]Department of Civil and Disaster Prevention Engineering, National United University, Miao-Li, Taiwan

* Corresponding author

Corresponding author Email: sltang@gate.sinica.edu.tw

Abstract:

Microbial communities have long been assumed to have functional redundancy, but very few studies have tested this empirically. Here, we performed a time series *in situ* reciprocal transplant experiment in two lakes with disparate trophic states to empirically test for functional redundancy. We determined that the two lakes had different community compositions, and these did not change even after incubating in a different environment. However, functional attributes were different between read- and predicted gene-level analyses of the communities, highlighting the importance of resolution when delineating community functional relationships. Furthermore, there was a linear relationship between community composition and functional attributes, with a broad range of similarities in the former and a narrower range and overall more similarity in the latter, providing support for partial functional redundancy. Ecoplate analysis suggests that the total metabolic activity of a community is influenced by its local environment, even though broad functions like carbon metabolism are widespread. Together, our results support the presence of resolution-dependent partial functional redundancy in aquatic environments.

Introduction

Understanding the diversity and role of microbial communities is important because microbes are ubiquitous (**Hug LA et al, 2016**) and drive ecosystem processes by governing biogeochemical cycles (**Ren Z et al, 2017; Graham BE et al, 2016; Falkowski PG et al, 2008**). In recent years, researchers have studied community compositions and functions and attempted to understand the community function relationship in different ecosystems ranging from oceans (**Tringe S et al, 2005; Louca S et al, 2016; Galand PE et al, 2018**) to freshwater ecosystems (**Gasol MJ, 2002; Peter H et al, 2011; Langenheder S et al, 2005**), soil (**Griffiths BS et al, 2003; Bahram M et al, 2018**), plants (**Reich PB et al, 2012; Louca S et al, 2017**), and the human gut (**Turnbaugh PJ et al, 2009; Moya A and Ferrer M, 2016**). These studies have put forward two plausible scenarios delineating community function relationships: 1) community and function are coupled so that changes in the community composition will change microbial-mediated processes, and 2) where community and function appear decoupled, microbe-mediated processes are somewhat independent of changes in community composition (**Allison SD and Martiny JBH, 2008**). The latter is also known as functional redundancy.

Lately, functional redundancy has been categorized into strict and partial functional redundancy. Strict functional redundancy implies that different taxa or phylotypes can conduct the same set of metabolic processes and can readily replace each other (**Allison SD and Martiny JBH, 2008; Yin B et al, 2000; Fuhrman JA et al, 2015**). Partial redundancy implies that certain organisms can coexist while sharing specific functions, but differing in other functions or ecological requirements (**Jurburg SD et al, 2015**). The paradigm of strict functional redundancy does not hold true in all ecosystems, as shown in soil (**Strickland MS et al, 2009; Fierer N et al, 2013**) and, recently, the oceans (**Galand PE et al, 2018**). Furthermore, studies applying

different molecular approaches to understand the community function relationship have yielded different conclusions. Studies using taxonomic profiling with the 16S rRNA gene or housekeeping genes (**Jaspers E and Overamann E, 2004; Doolittle WD and Zhaxybayeva O, 2009; Maharajan R et al, 2006**) have reported a disconnect between community composition and function, whereas studies focused on ecology and microbial molecular evolution have reported a strong relationship between community composition and functional traits (**Allison SD and Martiny JBH, 2008; Gravel D et al, 2011; Martiny JBH et al, 2006; Martiny AC et al, 2013**).

Terrestrial plants and animal ecosystem studies have provided the most insights into the community function relationship (**Peter H et al, 2011**) in recent years. Unlike terrestrial ecosystems, aquatic ecosystems have different and unique features (**Giller PS et al, 2004**). Studies of aquatic ecosystems, especially rivers and lakes, in both the field (**Shade A et al, 2007; Jones SE et al, 2009; Nelson CE et al, 2009**) and laboratory (**Judd KE et al, 2006; Kritzberg ES et al, 2006; Newton RJ et al, 2011**) have been central for deducing community function relationships; some of these studies support functional redundancy (**Fernandez AS et al, 2000; Langenheder S et al, 2005; Cheaib B et al, 2018; Tseng CH et al, 2015**) and others challenge it (**Findlay SEG et al, 2003; Traving SJ et al, 2016; Delgado-Baqueriza M et al, 2016**).

Langenheder S et al (2005) used water samples from four lakes as an incubating medium and inocula of combinations of each inside controlled laboratory bioreactors to test the community function relationship in the aquatic environment. They reported that communities with different compositions can have the same functions, and hence there is strong functional redundancy in the aquatic environment. Whereas **Delgado-Baquerizo M et al (2016)**, in their independent microcosm experiment, reported that a decline in microbial diversity has significant

and direct consequences for both broad and specific ecosystem functions, concluding that there is

either little or no functional redundancy in lake ecosystems. It is important to note that these

studies and many others are confined to specific functions (biomass, respiration rate, growth rate,

etc.) **(Gasol MJ et al, 2002; Peter H et al, 2011; Comte J et al, 2013; Langenheder S et al,**

2005), and results of direct 16S rRNA gene sequencing rarely align with those of whole

metagenomic approaches (**Lima-Mendez G et al, 2015**). Although these studies have broadened

our understanding of community function relationships in aquatic ecosystems, none have

mimicked the *in vivo* dynamics of natural ecosystems.

Efforts have been made using *in vitro* reciprocal transplant experiments to test the

hypothesis of functional redundancy in aquatic environmental studies, but most of these studies

were performed for short durations of time and in artificially controlled laboratory environments

(**Gasol MJ et al, 2002; Kirchman DL et al, 2004; Langenheder S et al, 2005; Shade A et al,**

2010). *In situ* transplant experiments have been used to study shifts in bacterial community

composition as a response to nutrient availability (**Gasol MJ et al, 2002)** and link functional

redundancy with metabolic plasticity in aquatic environments (**Comte J et al, 2013**). These

studies are limited to measuring broad functions such as cumulative microbial respiration,

dissolved organic carbon, biomass duplication time, and specific growth rate as proxies for the

metabolic potential of a community when it is transplanted into a different environment. Hence,

empirical research is needed to test the hypothesis of functional redundancy in aquatic

ecosystems using high throughput approaches encompassing both community composition and

function.

In this study, we performed an *in situ* experiment in which water (surface and bottom)

from one lake was reciprocally transplanted into another lake. Inoculated water was maintained

inside dialysis tubes resembling "microbial cages" that are exposed to ambient nutrient

concentrations from the incubating lake (**Herndl GJ et al, 1993**). We performed a multi-

approach analysis using the 16S rRNA gene and gene transcripts for total and active community

compositions, respectively. We used whole metagenome sequencing and 16S rRNA-based

pseudo-metagenome for functional profiles and Biolog Ecoplates for metabolic activity of the

microbial communities in the two lakes. Community function comparisons were performed to

test the hypothesis of functional redundancy and also to study how changing the local

environment affects the metabolic activity of microbial communities.

Materials and methods

Sampling sites

Two close proximity (21 km), sub-alpine lakes in northern Taiwan were chosen as

experimental sites: Tsuei-Feng Lake (TFL) (24° 30' 52.5'', 121° 36' 24.8'' E) and Yuan-Yang

Lake (YYL) (24° 34' 33.6'' N, 121° 24' 7.2'' E), at an altitude of 1,840 and 1,670 m,

respectively. These lakes, despite being at similar altitudes and geographical locations, have

different trophic states, with TFL being oligotrophic (**Wang LC et al, 2013**) and YYL

mesotrophic (**Tsai JW et al, 2008**). Both lakes are within a protected area, and are therefore

introduced to minimal or no anthropogenic activity year-round. These lakes have been at the

focus of limnological and meteorological research in Taiwan over the past years.

Experimental setup, sample collection, and preparation

Water samples from the surface (0.5 m deep) and bottom (1 m above the sediment) of

both lakes were collected every two weeks from January-February 2015 (four times total)

(**Figure 1D**). These samples were collected at the center of the lakes (also the deepest point)

using a boat. Two L of lake water (BG) (**Figure 1A**) was filtered using a piece of gauze (to

remove large debris, e.g., leaves), then sand and dust particles were removed using an 11-µm

filter. Finally, 200 ml of water was filtered using a 0.22-µm filter (diameter: 47 mm, Advantec);

samples were kept at -20° C before DNA extractions in the laboratory. For each sample, three

replicates (n = 3) were collected.

To create an artificial disturbance (for both the surface and bottom), 6 L from each places

in the lakes (surface and bottom) was swapped and incubated in the same and different lakes

(e.g. TFL water was swapped and incubated in TFL (Self-swapped) and YYL (Cross-swapped))

(**Figure 1B, 1C**). The dialysis tube (Sigma-Aldrich, St. Louis, MO, USA) allows only ions (Mol.

wt. <14 KDa) to pass through. These dialysis tubes were placed in specially designed metal

cages.

Seventy-two dialysis tubes (36 surface and 36 bottom), each filled with water samples

(~500 ml each), were incubated for six weeks. Every two weeks, samples from 12 tubes from

each lake (6 self-swapped and 6 cross-swapped) were collected and the rest of the dialysis tube-

containing water samples were replaced by new tubes to avoid algal growth on their outer

surface. Collected water samples (~500 ml each tube) were filtered as described above and stored

at -20°C before DNA extraction.

Nucleic acid extraction, PCR amplification, and sequencing

Total genomic DNA from filtered water samples was extracted using the UltraClean Soil

DNA Kit (MoBio, Solana Beach, CA, USA) following the manufacturer's protocol. RNA from

water samples was extracted using UltraClean Soil RNA Kit (MoBio, Solana Beach, CA, USA),

and residual DNA was removed using a TURBO DNA-Free Kit (Invitrogen, Carlsbad, CA,

USA). The nucleic acid yield was determined by NanoDrop ND-1000 UV-Vis

Spectrophotometer (Nano-drop Technologies, Inc., Wilmington, DE, USA).

Complementary DNA (cDNA) was synthesized from purified RNA using the SuperScript

IV First-Strand Synthesis System for RT-qPCR (Invitrogen, Carlsbad, CA, USA) following the

manufacturer's protocol. To prepare 16S rRNA gene transcripts, PCR amplification was

conducted to target the V6-V8 hypervariable regions of the bacterial 16S rRNA gene using the

bacterial universal primers -968F (5'AACGCGAAGAACCTTAC-3') **(Chen CP et al, 2011)**

and 1391R (5'-ACGGGCGGTGWGTRC-3') (**Jorgenson SL et al, 2012**). The reaction mixture

contained 1 μl of 5 U TaKaRa *Ex Taq* HS (Takara Bio, Otsu, Japan), 5 μl of 10× *Ex Taq* buffer,

4 μl of 2.5 mM deoxynucleotide triphosphate mixture, 1 μl of each primer (10 μM), and 1-5 μl

(10–20 ng) of template DNA in a volume of 50 μl. PCR was performed under the following

conditions: 94°C for 5 min, 30 cycles of 94°C for 30 s, 52°C for 20 s, and 72°C for 45 s, with a

final extension of 72°C for 10 min.

DNA-tagging PCR was used to tag each PCR product of the bacterial 16S rRNA gene's

V6-V8 region (**Chen CP et al, 2011**). The tag primer was designed with four overhanging

nucleotides; this arrangement ensured 256 distinct tags—at the 5′ end of the 968F and 1391R

primers—for bacterial DNA. The tagging PCR conditions comprised an initial step of 94°C for 3

min, 5 cycles at 94°C for 20 s, 60°C for 15 s, 72°C for 20 s, and a final step of 72°C for 2 min.

Illumina sequencing was performed on pooled 40-ng lots of uniquely marked samples by

Yourgene Biosciences, Taiwan using the Miseq system. Six TruSeq DNA-PCR Free libraries (3:

16S rRNA gene and 3: 16S rRNA gene transcripts) were prepared for 2×300 bp paired-end reads

by Yourgene Biosciences. Raw reads were sorted and primers removed before further analyses.

For whole-metagenome sequencing, the total genomic DNA was extracted using the

UltraClean Soil DNA Kit (MoBio, Solana Beach, CA, USA) following the manufacturer's

protocol. Extracted DNAs from the 41 samples were sent to Yourgene Biosciences (Taipei,

Taiwan) for library preparation without amplification (PCR Free library) and sequencing by the

Illumina MiSeq system (USA).

Community and functional attribute analysis

Illumina sequenced 16S rRNA gene reads were sorted into samples using an in-house

script. Sequences in each sample were screened by MOTHUR v1.3.81 (**SchlossPD et al, 2009**)

to keep those 1) of lengths >390 bp and <450 bp, 2) with 0 ambiguous bases, and 3) with

homopolymers <8 bp. UCHIME with USEARCH v8 (**Edgar RC, 2010**) (parameters: reference

mode, rdp gold database, and mindiv of 3) was used to remove chimeric reads. After

dereplication and removal of singletons, the reads were clustered at a 97% identity threshold to

get cross-sample representative sequences as operational taxonomic units (OTUs) using the

UPARSE pipeline (**Edgar RC, 2013**). The taxonomic classification of OTUs was performed by

MOTHUR using SILVA 16S rRNA database v128 (**Quast CE et al, 2013; Yilmaz P et al,**

2014). 16S rRNA gene transcript read datasets were processed using the same approach outlined

above.

Metagenomic reads obtained from the Illumina MiSeq system were preprocessed and the

adapters removed using Cutadapt v1.4.2 (**Martin M, 2011**), quality trimming of reads was

performed with Seqtk v. 1.2.-r94 (phred score <20, short read cut off <35 bp) (**Shen W et al,**

2016), singlet reads were discarded. Metagenomes data were processed using two approaches

(**Figure S1**). 1) Predicted ORF level: reads were *de-novo* assembled individually for each sample

using CLC Genomics workbench version 1.10.1 (Qiagen) with default parameters, scaffolding

was not performed, ORF prediction was performed using Prodigal (**Hyatt D et al, 2010**)

wrapped in PROKKA on contigs >200 bp long (**Seemann T, 2014**) with metagenome mode

enabled. 2) Reads level: quality filtered paired reads were merged using FLASH version 1.2.11

(**Magoc T et al, 2011**). These merged reads were used for the functional analysis.

Predicted ORFs and merged reads from each metagenome were searched for similarity

using DIAMOND (**Buchfink B et al, 2014**), implementing blastx (**Altschul SF et al, 1990**)

against the conserved domain database (CDD) of NCBI (last downloaded: 2018) (**Marcher-**

Bauer A et al, 2014) with e-value <1e-5 and aligned length of 50 amino acids. Annotated ORFs

and reads were categorized at COG (Clusters of Orthologous Groups) Class and Family levels

from the blastx results using in-house scripts. At the reads level, COG-Family represented in

90% of samples with at least 10 reads each, were used for the downstream analysis. Furthermore,

previously-obtained 16S rRNA gene transcript abundance profiles were used to obtain imputed-

metagenome functional profiles (at the COG Family level only) with PICRUSt2

(**https://github.com/picrust/picrust2**).

Ecoplates as the proxy for metabolic activity

The carbon assimilation potential of the microbial communities in the two lakes was

assessed with Biolog EcoPlates[TM] (Biolog Inc., Hayward, CA, USA), hereafter written as

ecoplates. Undiluted water (150 μl) from each sample was added to ecoplate wells, each

containing 31 different carbon sources (and one control well with no carbon source). Plates were

incubated at 25°C and the absorbance of each well was measured at 590 nm using an automatic

optical density microplate reader (Spectramax M2, Molecular Devices, San Jose, CA, USA)

every 24 hours for 7 days. On day 7, absorbance values from ecoplates were used to calculate

average well color development (AWCD) in the plate, as expressed by the equation below.

$$AWCD = (\Sigma\,(R - C)/93)$$

215   where C and R represent the absorbance value in the blank well and substrate, respectively.

216   Negative values were converted to 0. The absorbance values were then treated and normalized

217   using AWCD.

Statistical analysis

All statistical analyses were performed in R (**R core Team, 2016**). Abundance profiles

from the 16S rRNA gene and 16S rRNA gene transcript amplicon datasets were calculated using

an in-house R script and processed with R packages phyloseq (**McMurdie PJ et al, 2013**), vegan

(**Okansen J et al, 2007**), and ggplot (**Wickham H, 2016**) for downstream analysis and

visualization, including ordination, alpha and beta diversity calculations. Alpha diversities

(Richness, Shannon, and Inv. Simpson) were measured using the estimate_richness function

from phyloseq in R, with subsampling libraries with replacement 100 times and averaging the

diversity estimates from each trial. Rarefaction curves were drawn using the ggrare function

from the ranacapa package (**Kandlikar GS et al, 2018**) on unrarefied samples. All the

abundance profiles were log (x+1)-transformed to achieve normality. Beta diversity was

performed based on an ordination analysis using the Bray-Curtis distance method unless

otherwise stated. Permutational multivariate analyses of variance (PERMANOVA) were

performed using the Adonis function in the vegan package. Ecoplate normalized absorbance

values were also log (x+1)-transformed for PCA analysis and functional clusters were

determined using gap statistics from the cluster (**Maechler M et al, 2013)** with the K-means

clustering method. Effects of the transplant experiment on carbon assimilation was tested using

unpaired Student's t-test. The Mantel's Pearson correlation test was performed with Bray-Curtis

distance for the community, functional data, and with Euclidean distance for ecoplate substrate

utilization profiles. Microbial community and function differences were quantified using *L*-

divergence (**Lin J et al, 1991**). The linear relation between community and function was

statistically tested with linear regression analysis and ANOVA.

Data Availability

All the sequencing data have been deposited in the NCBI database under the

BioProject PRJNA596577. Specifically, the amplicon 16S rRNA gene and gene-transcripts data

have been deposited in NCBI SRA database accession number ###, ### respectively. The whole-

metagenome reads for 41 samples have been deposited in NCBI SRA (). Other miscellaneous

information's are available from the corresponding author upon request.

Results

Diversity analysis and its change

We sampled water from the surface and bottom of TFL and YYL every two weeks from January-

February 2015. BG samples acted as the control for community composition and functional

attributes of undisturbed lake water in their natural setting. Unrarefied samples were applied to

generate rarefaction curves and determine how well the sequencing effort represented for the

microbial diversity. Most of the samples from the 16S rRNA gene and 16S rRNA amplicons

appeared to plateau, indicating that the sequencing adequately captured most of the microbial

diversity in the two lakes (**Figure S2, S3**).

Alpha diversity (Shannon and Inv. Simpson) was measured after rarifying 16S rRNA

gene samples at 27,689 (surface) and 26,238 (bottom) reads, and 16S rRNA samples at 25,513

(surface) and 23,131 (bottom) reads. YYL BG samples (Ys and Yb) had higher Shannon and Inv.

Simpson diversities than TFL BG samples (Ts and Tb) (**Table S1**). We observed a mixed pattern

in which diversity (Shannon and Inv. Simpson) in self-swap and cross-swap samples decreased

from two to four weeks, and later increased from four to six weeks (**Table S1**).

Community function relationship in BG samples: hints of resolution dependence

The total community composition of BG samples from the two lakes was significantly different

at the class and genus taxonomic levels. At the genus level, PCoA axis1 and axis2 explained

>75% of the total variance in the surface and bottom levels, with inoculum (lake water) being the

significant factor (Surface: $ADONIS_{inoculum}$: $F_{1, 6} = 16.01$, $p < 0.05$; Bottom: $ADONIS_{inoculum}$: $F_{1, 6}$

$= 15.71$, $p < 0.05$) (**Figure S4A**).

There was no significant difference between the COG-Class based functional profile

(ORF level) of BG samples from the two lakes (**Table S2**). Since COG-Class is a general

classification for functional attributes, we performed a functional profile comparison at the finest

resolution possible (i.e., COG-Family). At COG-Family (ORF-level), we observed that inoculum

had a weak but significant effect driving the functional difference between BG samples from the

lakes (Surface: $ADONIS_{inoculum}$: $F_{1, 6} = 2.85$, $p < 0.05$, Bottom: $ADONIS_{inoculum}$ $F_{1, 6}$: 2.69, $p <$

0.05) (**Table S2**).

We also performed a functional profile analysis on merged reads at the COG-Class and -

Family levels. With reads at the COG-Family level, inoculum (lake water) had a strong and

significant effect on the functional profile (COG-Family) of the BG samples (Surface:

ADONIS$_{inoculum}$ $F_{1,6}= 6.24$, $p < 0.05$; Bottom: ADONIS$_{incoculum}$: $F_{1,6}= 4.152$, $p < 0.05$) (**Figure**

S4B, **Table S2**). No significant difference was observed in reads at the COG-Class level (**Table**

S2).

The different functional profiling outcomes for BG samples from the two lakes, based on

the level of analysis (ORF vs Reads; COG-Class vs COG-Family), suggest a possible resolution

dependency when delineating community function relationships. Hence, both approaches were

used (Metagenome level: ORF and Read; COG: Class and Family level) to compare the

functional profiles and test the hypothesis that these lake microbial communities have functional

redundancy.

The community functional relationship is resolution-dependent

The *in situ* reciprocal transplant experiment was conducted to empirically test the community

function relationship in a temporal manner. The total community composition of the cross-swap

samples was similar to their self-swap counterparts for both the surface and bottom samples. At

the genus level, PCoA axis1 and axis2 explained >75% of the total variance (**Figure 2A**). Only

the inoculum (not the incubating lake) had a significant effect (Surface: ADONIS$_{inoculum*incubating-}$

$_{lake}$: $F_{1,8\ (inoculum)} =18.90$, $p < 0.05$; Bottom: ADONIS$_{inoculum*incubating-lake}$: $F_{1,8\ (inoculum)} = 20.44$, $p <$

0.05) (**Table S2**). This clustering of cross-swap samples with their self-swap counterparts spoke

to the resistance and/or resilience of the bacterial community to external perturbations over time.

There was no significant difference in the functional profile of swapped samples (cross-

and self-swap) between COG-Class and -Family when analyzed with ORF-level **(Table S2)**.

Nevertheless, when analyzed with reads-level at the COG-Family level, the inoculum had a

significant effect on the clustering of the samples (Surface: ADONIS$_{\text{inoculum*incubating-lake}}$: $F_{1, 8}$

$_{(\text{inoculum})}$ = 4.02, $p < 0.05$; Bottom: ADONIS$_{\text{inoculum*incubating-lake}}$: $F_{1, 8 \, (\text{inoculum})}$ = 5.41, $p < 0.05$)

(**Figure 2B**). No significant difference was observed at the COG-Class level for the surface

samples **(Table S2)**. Furthermore, we observed that cross-swap samples tended to shift away

from their self-swap counterparts with an increase in incubation time (**Figure 2B marked by**

arrows).

Differences in community compositions and the results of the functional profile for

swapped samples confirmed that community function relationships are resolution dependent.

Furthermore, shifts in the functional profile of cross-swap samples away from their self-swap

counterparts after two and four weeks of incubation in different lakes suggest that their

community functions are dynamic. These shifts also suggest that the local environment plays an

important role in shaping the community's functional profile.

Active community, its functional attributes, and partial functional redundancy

16S rRNA gene-based community composition and whole metagenome-based functional

profiles represents the entire bacterial community and overall functional profile of the

ecosystem, respectively. As not all bacteria are metabolically active at the same time, it is

important to determine the community function relationship based on active community profiles,

as metabolically active bacteria are one of the major driving forces in the relationship. Hence, we

analyzed community profiles with 16S rRNA gene transcript amplicons and derived imputed

pseudo-metagenome functional profiles (COG-Family level) from the active community composition using PICRUSt2.

The active community composition profile was positively correlated with the 16S rRNA gene-based total community profile ($R_{surface}$: 0.9318, $p < 0.01$; R_{bottom}: 0.9246, $p < 0.01$) (**Table S3**). Two lakes had significantly different community compositions at the genus level (**Figure S5**). PICRUSt2 derived a pseudo-metagenome-based functional profiles at the COG-Family level, and was also positively correlated with the whole metagenome-based functional profile (COG-Family: Reads level) ($R_{surface}$: 0.3430, $p < 0.01$; R_{bottom}: 0.2895, $p < 0.01$) (**Table S3**). PERMANOVA analysis showed that only the inoculum had a significant effect on sample clustering (Surface: $ADONIS_{inoculum*incubating-lake}$: $F_{1, 8 (inoculum)} = 16.65$, $p < 0.05$; Bottom: $ADONIS_{inoculum*incubating-lake}$: $F_{1, 8 (inoculum)} = 15.05$, $p < 0.05$) (**Table S2**).

We performed a linear regression analysis based on the Bray-Curtis similarity (1-Bray-Curtis distance) on the active community and the PICRUSt2-based functional profile to test whether there is a linear relationship between bacterial community composition and functional profile at any similarity level. We found that similarities in community composition profiles were strongly correlated with functional attributes ($R^2_{surface} = 0.65$, $R^2_{bottom} = 0.63$). This linear relationship between community and function was based on a broad range of similarities in the community (<20% to >70%) and a relatively narrow but higher range of similarity in functional attributes (>90% to <97%). This result supports the idea that functional redundancy and functional dissimilarity co-exist in this ecosystem (i.e., partial functional redundancy) (**Figure 3A**).

Quantifying functional redundancy

L-divergence, which scales from 0 (identical) to 2 (highly divergent), was used to

quantify functional redundancy. We identified a highly divergent community (*L*-divergence$_{surface}$:

1.02±0.48; *L*-divergence$_{bottom}$: 0.91±0.44) based on the 16S rRNA gene and a relatively similar

functional profile (*L*-divergence$_{surface}$: 0.04±0.1; *L*-divergence$_{bottom}$: 0.039±0.41) based on whole

metagenomes in the two lakes (**Figure 3B**). Furthermore, we also calculated *L*-divergence for

community and functional profiles using different datasets from this study and obtained similar

results (**Table S4**).

Metabolic activity is influenced by the local environment

We used ecoplates as a proxy to both measure the metabolic activity (in terms of carbon

metabolism) of the microbial communities in the two lakes and test temporal-physiological

changes in metabolic activity during the reciprocal transplant experiment. Mean AWCD was

used as an indicator of metabolic activity, and we observed that YYL lake BG samples had a

higher mean AWCD than did TFL lake samples. Different values of AWCD were observed at

different sampling time-points (**Figure S6**).

Swapped samples were used to test whether incubating the samples from the low nutrient

environment (TFL) in the high nutrient environment (YYL) and vice versa affected metabolic

activity. Two categories were used to compare samples. 1) Same inoculum in different

incubating lakes (surface: Ts→Y vs Ts→T, Ys→T vs Ys→Y; bottom: Tb→Y vs Tb→T, Yb→T

vs Yb→Y) 2) Different inoculum in the same incubating lake (surface: Ts→Y vs Ys→Y, Ys→T

vs Ts→T; bottom: Tb→Y vs Yb→Y, Yb→T vs Tb→T).

While comparing samples with the same inoculum in different incubating lakes, we

observed that the metabolic activity of Ts→Y samples was significantly higher than in Ts→T

samples (**Figure 4A**) across all time points. However, Ys→T samples had significantly lower

metabolic activity (except Ys3→T vs Ys3→Y) than Ys→Y samples (**Figure S7A**). In the

bottom samples, the metabolic activity of Tb1→Y samples was significantly higher than Tb1→T

samples (**Figure 4B**), and no significant difference was observed at other sampling time points.

Yb1→T samples had a lower metabolic activity than Yb1→Y samples, and no significant

difference was observed between Yb2→T and Yb2→Y. However, Yb3→T had significantly

higher metabolic activity than Yb3→Y (**Figure S7B**).

When the inoculum was different in the same incubating lake, we observed that Ys→T

samples had higher metabolic activity than Ts→T samples across all sampling time points

(**Figure 4C**), and Ys→Y samples had higher metabolic activity than Ts→Y samples (except

Ts3→Y vs Ys3→Y) (**Figure S7C**). Of the bottom samples, Yb→T metabolic activity was higher

than Tb→T (**Figure 4D**), and Yb→Y was higher than Tb→Y (except Tb3→Y) (**Figure S7D**).

Discussion

In recent years, the focus of microbial ecology studies has shifted from understanding the

community diversity to understanding what microbes are doing and how they respond to changes

in the environment, and delineating community function relationships in diverse ecosystems

(Tringe SG et al, 2005; Louca S et al, 2016; Mooshammer M et al, 2017; Louca S et al,

2018; Galand PE et al, 2018; Orland C et al, 2019). With advancements in sequencing

technology and relative improvements in metagenomic analyses, efforts have been made to

establish community function relationships in pristine and disturbed ecosystems. Recently,

laboratory and field studies from different environments testing community function

relationships have provided insightful results (**Louca S et al, 2016; Debroas D et al, 2009;**

Delgado-Baquerizo M et al, 2016; Galand PE et al, 2018, **Cheaib B et al, 2018**;

Langenheder S et al, 2005; Comte J et al, 2013; Shade A et al, 2010, 2011, 2012; Baho DL et

al, 2012). Yet, our understanding of community function relationships and their magnitudes in

natural environments remains poor because of the lack of *in situ* empirical time series studies

applying high-throughput sequencing technology alongside functional approaches. In this

setting, the main motivation of our study was to empirically test the hypothesis of functional

redundancy in the aquatic ecosystem with an *in situ* reciprocal transplant experiment. We used

high throughput amplicon, whole metagenome sequencing, and ecoplate analysis to determine

the community function relationship in the aquatic ecosystem. To the best of our knowledge, this

is the first study that examines the function redundancy hypothesis in surface and bottom lake

waters with time series reciprocal transplant experiments *in situ.*

The 16S rRNA gene-based bacterial community composition of BG samples was

significantly different between lakes **(Figure S4A)**, which was expected, as the two lakes have

different trophic states (**Tsai JW et al, 2008; Wang LC et al, 2013; Tandon K et al, 2018**).

However, results from the functional profile analysis have unexpected implications for resolution

dependency and the community function relationship. ORF-based functional profiles (BG

samples) for COG-Class and -Family were not significantly different between the two lakes,

suggesting that functional redundancy is present (**Table S2**). Functional profiles supported

functional redundancy in previous studies as well when analyzed at the ORF level (**Sunagawa S**

et al, 2015; Louca S et al, 2016). Read-based functional profiles, however, were significantly

different for the two lakes **(Figure S4B)**, challenging our conclusions about functional

redundancy and suggesting that resolution dependency helps determine the community function

relationship. Recently, **Galand PE et al, 2018,** using sequenced reads from a three-year survey,

determined a functional profile and identified a lack of strict functional redundancy in the

oceans. They argued that the functional profile, when examined at the predicted gene level, can

significantly undermine the overall functional profile of ecosystems, as predicted genes are

extrapolated from the similarity between cultured and taxonomic annotations of environmental

samples. Nonetheless, we propose that clustering reads directly to assign functions can also give

spurious results that have little significance without taxonomy. Thus, we used reads directly to

get functional profiles of the lake water community using *a priori* knowledge of operational

databases.

Studies in recent years have explored community responses and functional profiles in

different aquatic environments using reciprocal transplant experiments and other approaches

(**Langenheder S et al, 2005, 2006; Shade A et al, 2010; Peter H et al, 2011; Comte J et al,**

2013). We identified a significant difference in microbial communities in two lakes, even after

reciprocal transplantation and incubation for six weeks. Cross-swap samples clustered with their

self-swap counterparts for both surface and bottom samples (**Figure 2A**). Furthermore, inoculum

had a significant effect on the clustering of the samples (**Table S2**)**.** Our results are similar to

Langenheder S et al, 2006; thus, we suggest that the bacterial community was more or less

resistant to change, which is consistent with previous studies (**Allison SD et al, 2008; Shade A**

et al, 2011). We cannot, however, rule out the possibility that the bacterial communities are

resilient and can recover from a disturbance to an original or alternate stable state (**Pimm SL,**

1984) because our sampling interval was two weeks—which is longer than most sample time

point experiments—and also because of the shifts observed within the cluster of cross-swap

samples (**Figure 2A, Figure S5**). We believe that the community might have varied immediately

or very shortly after transplantation, but then recovered to an alternate stable state in less than

two weeks. Bacterial communities in a lake ecosystem was demonstrated to recover quickly after

natural disturbance (lake mixing) (**Shade A et al, 2012**). Earlier studies have shown bacterial

communities to be both resilient and resistant in various ecosystems (**Allison SD et al, 2008;**

Shade A et al, 2011).

We hypothesized that transplanting an inoculated sample from one lake into another lake

with an entirely different trophic state would influence the sample's functional profile. A similar

hypothesis was tested and confirmed in an earlier study (**Graham BE et al, 2016**). Interestingly,

we observed shifts in the functional profile of cross-swap samples away from their self-swap

counterparts (**Figure 2B, marked as dashed arrows**). This phenomenon was observed for

Ys$\rightarrow$T and Yb$\rightarrow$T samples, but was not as clearly for Ts$\rightarrow$Y and Tb$\rightarrow$Y samples (**Figure 2B**).

During our experiment, the Tb2$\rightarrow$Y sample (after four weeks of incubation) was lost, and hence

the shift in Tb$\rightarrow$Y samples cannot be determined accurately. Furthermore, transplant samples

behaved differently in the two lakes, suggesting that local environment influences the functional

profile of a community, maybe due to the nutrient availability in that environment and how it

differs from that of the original environment (**Tsai JW et al, 2008; Wang LC et al, 2013**).

Although these shifts are not quantified, they support our hypothesis that local environment

influences the functional profile of the swapped bacterial community, as was previously reported

in plants (**Louca S et al, 2017**) and animals (**Hou, D et al, 2017**). A long-time series analysis

would help determine whether communities in a new environment eventually settle into an

alternate stable state.

Partial functional redundancy has been reported in different ecosystems (**Langenheder S**

et al, 2006; Delgado-Baquerizo M et al, 2016; Galand PE et al, 2018). Therefore, we

speculated that there is a linear relationship between community composition and ecosystem

function. Regression analysis using active community composition and the imputed metagenome

functional profile identified a linear relationship between the community and its function in the

ecosystem of both lakes (**Figure 3A**). Our result contrasts with a study on the marine ecosystem,

where the range of similarities between community composition and function profile was of a

similar magnitude (**Galand PE et al, 2018**); this might be due to the aquatic environment studied

and the duration and frequency of sampling. Furthermore, a relatively highly divergent

community (L-divergence$_{surface}$: 1.02±0.48; L-divergence$_{bottom}$: 0.91±0.44) and relatively low

divergent functional profile (L-divergence$_{surface}$: 0.04±0.1; L-divergence$_{bottom}$: 0.039±0.41)

(**Figure 3B**) between the two lakes also confirms a partial functional redundancy, whose

magnitude varies based on the resolution of the analysis (**Table S4**). Significant high positive

correlations were obtained between the 16S rRNA gene and 16S rRNA gene transcript-based

community composition and also with whole metagenome and PICRUSt2 derived imputed

metagenome functional profiles (**Table S3**). This implies that either of the approaches can be

used to determine the community function relationship in our study, and their results will be not

differing significantly.

Natural communities can use diverse carbon and energy sources for growth and survival

(**Eiler A et al, 2006**). Microbes, especially bacteria, often overcome changes in the environment

by expressing a range of metabolic capabilities or mixotrophy (**Meyer AF et al, 2004; Swingly**

WD et al, 2007). Therefore, ecoplates containing carbon sources of different complexities were

used to test whether a change in the local environment has any consequence on the total

metabolic activity of the bacterial community; they identified a similar potential substrate utilization pattern to other studies (**Comte J and Del Giorgio PA, 2009; Comte J et al, 2013; Traving SJ et al, 2016**). Over seven days of incubation and observation, and using carbon metabolism as a proxy for overall metabolism, we determined that the total metabolic activity of Ts→Y samples were higher than Ts→T ones (**Figure 4A**), and that of Ys→T samples was lower (except Ys3→T) than Ys→Y ones (**Figure S7A**). Similar results were obtained for bottom samples (**Figure 4B** and **Fig S7B**). Furthermore, Ys→T and Yb→T samples had higher metabolic activity than Ts→T and Tb→T, respectively (**Figure 4C, 4D**), and Ts→Y and Tb→Y had lower metabolic activity than Ys→Y and Yb→Y, respectively (**Figure S7C, S7D**). These changes in total metabolic activity highlight the effect that local environment can have on the total metabolic activity of a bacterial community. These results also provide important evidence that metabolic activity differs distinctly between the surface and bottom waters of the lakes. We also demonstrate that inoculum has either a very weak or no effect on functional profiles in terms of carbon substrate utilization (**Figure S8, Table S5**), which is similar to findings from other studies (**Tranvik LJ et al, 1987; Gasol MJ et al, 2002; Findlay SEG et al, 2003; Langenheder S et al, 2005**). Lack of correlation between bacterial community functional attributes and composition has been interpreted as evidence of functional redundancy (**Rosenfeld SJ et al, 2002; Allison SD and Martiny JBH, 2008**). However, the significantly high correlation suggests that functional redundancy may be constrained. We observed no or very small positive correlation between total or active bacterial community and substrate utilization profiles based on ecoplates (**Table S3**). A similar result was observed in boreal freshwater bacterioplankton communities and coastal waters (**Ruiz-Gonzalez C et al, 2015; Traving SJ et al, 2016**). This may again suggest that some substrates cause functional redundancy.

In the present study, we used a reciprocal transplant experiment in which the bacterial

community from one lake was grown in another, very different lake. Dialysis tubing bags, which

acted as microbial cages, provided bacteria access to the nutrients in their transplant

environment. Although dialysis bags are not perfect cages, they are much more realistic than any

experimental bottles or reactors. Our sampling strategy did not determine whether the bacterial

communities in the two lakes were resilient to change; future experiments aiming to answer this

question should use shorter intervals between sampling. Functional redundancy is a difficult

concept to quantify, and we still do not know its magnitude and variability among different

ecosystems, so care should be taken when applying results from this study to other environments.

Conclusion:

Establishing the community function relationship has been central to understanding the

complexity of microbial diversity and their functions in different ecosystems. Laboratory and

field studies have made important findings on this subject in recent years. Yet, few *in situ* long

term experiments have been conducted empirically to test the hypothesis of functional

redundancy. Furthermore, most previous studies were limited to measuring specific functions,

and hence they could not make conclusions about overall ecosystem dynamics. This study is the

first to test the community function relationship through reciprocal transplant experiments in lake

waters (surface and bottom) using a time series, utilizing amplicon sequencing complemented

with whole metagenome sequencing and metabolic analysis using ecoplates. The insights offered

by this work deciphering the community function relationship of the freshwater aquatic

ecosystem will add much-needed knowledge in this field.

We found that bacterial community composition is resistant and/or resilient to change in

the local environment. Whereas, the functional profile is influenced by the local environment.

Further, the community function relationship is dependent on the resolution of the analysis. At

the predicted ORF level, we observed no significant difference in the community functional

profile of the two lakes. However, at the reads level, a significant difference in the community

functional profile of the two lakes was observed. Moreover, a significant correlation between

community composition (broad similarity range) and community functional attributes (small but

higher similarity range) supports the presence of partial functional redundancy in aquatic

environments. Metabolic activity analysis using ecoplates also provides strong support for the

presence of functional redundancy in the aquatic environments and also provides evidence for

the influence of the local environment on the metabolic activity. Further, the two lakes and their

surface and bottom waters show differing effects on the community and functional profile

behavior when disturbed artificially.

Acknowledgements

The authors would like to thank Dr. Ching-Hung Tseng, Dr. Jiun-Yan Ding, Mr. Cheng-Yu

Yang, Dr. Sonny T. M. Lee, and Dr. Carol Enumi Lee for their assistance with sampling. K.T

would like to acknowledge the Taiwan International Graduate Program (TIGP) for its fellowship.

The authors also express their gratitude to the anonymous reviewers for their insightful

comments and suggestions to improve the quality of the manuscript, and to Mr. Noah Last of

Third Draft Editing for his English language editing. This study was supported by the Thematic

Project at Academia Sinica (AS-103-TP-B15-3).

Contributions

S.L.T conceived of the idea for this study. K.T performed all the analyses and wrote and revised

the manuscript. M.W and C.Y performed sampling and experiments. S.Y played an active role in

the discussion, interpretation of results, and writing of the manuscript. B.B, C.C, and J.T helped

with sampling and provided logistical support for the experimental setup. C.S.N provided critical

comments on the manuscript. All authors read and approved the current version of the

manuscript. All authors also declare that they have no conflict of interest.

Competing Interests

All authors declare no competing interests.

References

1.   Hug, L. A. *et al.* A new view of the tree of life. *Nat. Microbiol.* **1**, 1–6 (2016).

2.   Ren, Z. *et al.* Taxonomic and functional differences between microbial communities in

Qinghai Lake and its input streams. *Front. Microbiol.* **8**, 1–14 (2017).

3.   Graham, E. B. *et al.* Microbes as engines of ecosystem function: When does community

structure enhance predictions of ecosystem processes? *Front. Microbiol.* **7**, 1–10 (2016).

4.   Falkowski, P. G., Fenchel, T. & Delong, E. F. The Microbial Engines That Drive Earth's

Biogeochemical Cycles. *Science (80-.).* **320**, (2008).

5.   Tringe, S. G. *et al.* Comparative metagenomics of microbial communities. *Science (80-.).*

308, 554–557 (2005).

6.   Louca, S., Parfrey, L. W. & Doebeli, M. Decoupling function and taxonomy in the global

ocean microbiome. *Science (80-.).* **353**, (2016).

7.   Galand, P. E., Pereira, O., Hochart, C., Auguet, J. C. & Debroas, D. A strong link between

marine microbial community composition and function challenges the idea of functional

redundancy. *ISME J.* **12**, 2470–2478 (2018).

8.   Gasol, J. M. *et al.* A transplant experiment to identify the factors controlling bacterial

abundance, activity, production, and community composition in a eutrophic canyon-

shaped reservoir. *Limnol. Oceanogr.* **47**, 62–77 (2002).

9.   Peter, H. *et al.* Function-specific response to depletion of microbial diversity. *ISME J.* **5**,

351–361 (2011).

10.   Langenheder, S., Lindström, E. S. & Tranvik, L. J. Weak coupling between community

composition and functioning of aquatic bacteria. *Limnol. Oceanogr.* **50**, 957–967 (2005).

11.   Griffiths, B. S. *et al.* The relationship between microbial community structure and

functional stability, tested experimentally in an upland pasture soil. *Microb. Ecol.* **47**,

104–113 (2004).

12.   Bahram, M. *et al.* Structure and function of the global topsoil microbiome. *Nature* **560**,

233–237 (2018).

13.   Reich, P. B. *et al.* Impacts of biodiversity loss escalate through time as redundancy fades.

Science (80-.). **336**, 589–592 (2012).

14.   Louca, S. *et al.* Functional structure of the bromeliad tank microbiome is strongly shaped

by local geochemical conditions. *Environ. Microbiol.* **19**, 3132–3151 (2017).

15.   Turnbaugh, PJ. *et al.* A core gut microbiome in obese and lean twins. *Nature* **457**, 222–

227 (2009).

16.   Moya, A. & Ferrer, M. Functional Redundancy-Induced Stability of Gut Microbiota

Subjected to Disturbance. *Trends Microbiol.* **24**, 402–413 (2016).

17.   Allison, S. D. & Martiny, J. B. H. Resistance, resilience, and redundancy in microbial

communities. *Light Evol.* **2**, 149–166 (2009).

18.   Yin, B., Crowley, D., Sparovek, G., De Melo, W. J. & Borneman, J. Bacterial functional

586   redundancy along a soil reclamation gradient. *Appl. Environ. Microbiol.* **66**, 4361–4365

587   (2000).

588   19.   Fuhrman, J. A., Cram, J. A. & Needham, D. M. Marine microbial community dynamics

589   and their ecological interpretation. *Nat. Rev. Microbiol.* **13**, 133–146 (2015).

590   20   Jurburg, S.D., & Salles, J.F. Functional Redundancy and Ecosystem Function- The Soil

591   Microbiota as a Case Study. *Ecosystems-Linking Strucuture and Function.* p29-42. InTech

592   Oen Science. Rijeka (2015)

593   21.   Strickland, M. S., Lauber, C., Fierer, N. & Bradford, M. A. Testing the functional

594   significance of microbial community composition. *Ecology* **90**, 441–451 (2009).

595   22.   Fierer, N. *et al.* Prairie Soils in the United States. *Science (80-.).* **342**, 621–625 (2013).

596   23.   Elke Jaspers and Jorg Overmann. Ecological Significance of Microdiversity: Identical 16S

597   rRNA Gene Sequences Can Be Found in Bacteria with Highly Divergent Genomes and

598   Ecophysiologies. *Inst. fur Chemie und Biol. des Meeres, Univ. Oldenburg, D-26111*

599   *Oldenburg, Ger.* **70**, 62 (2014).

600   24.   Doolittle, WF and Zhaxybayeva O. On the origion of prokaryotic species. *Genome Res*

601   **19**(5), 744-756 (2009).

602   25.   Maharjan, R., Seeto, S., Notley-McRobb, L. & Ferenci, T. Clonal adaptive radiation in a

603   constant environment. *Science (80-.).* **313**, 514–517 (2006).

26.   Gravel, D. *et al.* Experimental niche evolution alters the strength of the diversity-

productivity relationship. *Nature* **469**, 89–94 (2011).

27.   Martiny, J. B. H. *et al.* Microbial biogeography: putting microorganisms on the map. *Nat.*

Rev. Microbiol. **4**, 102–112 (2006).

28.   Martiny, A. C., Treseder, K. & Pusch, G. Phylogenetic conservatism of functional traits in

microorganisms. *ISME J.* **7**, 830–838 (2013).

29.   Giller, P. S. *et al.* Biodiversity effects on ecosystem functioning: Emerging issues and

their experimental test in aquatic environments. *Oikos* **104**, 423–436 (2004).

30.   Shade, A. *et al.* Interannual dynamics and phenology of bacterial communities in a

eutrophic lake. *Limnol. Oceanogr.* **52**, 487–494 (2007).

31.   Jones, S. E., Newton, R. J. & McMahon, K. D. Evidence for structuring of bacterial

community composition by organic carbon source in temperate lakes. *Environ. Microbiol.*

11, 2463–2472 (2009).

32.   Nelson, C. E. Phenology of high-elevation pelagic bacteria: The roles of meteorologic

variability, catchment inputs and thermal stratification in structuring communities. *ISME*

J. **3**, 13–30 (2009).

33.   Judd, K. E., Crump, B. C. & Kling, G. W. Variation in dissolved organic matter controls

bacterial production and community composition. *Ecology* **87**, 2068–2079 (2006).

34.   Kritzberg, E. S., Langenheder, S. & Lindström, E. S. Influence of dissolved organic matter

source on lake bacterioplankton structure and function - Implications for seasonal

dynamics of community composition. *FEMS Microbiol. Ecol.* **56**, 406–417 (2006).

35.   Newton, R. J. & Mcmahon, K. D. Seasonal differences in bacterial community

composition following nutrient additions in a eutrophic lake. *Environ. Microbiol.* **13**, 887–

899 (2011).

36.   Fernandez, A. S. *et al.* Flexible community structure correlates with stable community

function in methanogenic bioreactor communities perturbed by glucose. *Appl. Environ.*

Microbiol. **66**, 4058–4067 (2000).

37.   Cheaib, B., Boulch, M. Le, Mercier, P. L. & Derome, N. Taxon-function decoupling as an

adaptive signature of Lake Microbial metacommunities under a chronic polymetallic

pollution gradient. *Front. Microbiol.* **9**, 1–17 (2018).

38.   Tseng, C. H. *et al.* Microbial and viral metagenomes of a subtropical freshwater reservoir

subject to climatic disturbances. *ISME J.* **7**, 2374–2386 (2013).

39.   Findlay, S. E. G., Sinsabaugh, R. L., Sobczak, W. V. & Hoostal, M. Metabolic and

structural response of hyporheic microbial communities to variations in supply of

dissolved organic matter. *Limnol. Oceanogr.* **48**, 1608–1617 (2003).

40.   Traving, S. J. *et al.* Coupling bacterioplankton populations and environment to community

function in coastal temperate waters. *Front. Microbiol.* **7**, 1–13 (2016).

41.   Delgado-Baquerizo, M. *et al.* Lack of functional redundancy in the relationship between

microbial diversity and ecosystem functioning. *J. Ecol.* **104**, 936–946 (2016).

42.   Lima-Mendez, G. *et al.* (2015) Determinants of community structure in the grobal

plankton interactome.pdf. *Ocean Plankt.* **348**, 1–10 (2015).

43.   Kirchman, D. L., Dittel, A. I., Findlay, S. E. G. & Fischer, D. Changes in bacterial activity

and community structure in response to dissolved organic matter in the Hudson River,

647   New York. *Aquat. Microb. Ecol.* **35**, 243–257 (2004).

648 44. Shade, A., Chiu, C.-Y. & McMahon, K. D. Differential bacterial dynamics promote

649   emergent community robustness to lake mixing: an epilimnion to hypolimnion transplant

650   experiment. *Environ. Microbiol.* **12**, 455–66 (2010).

651 45. Comte, J., Fauteux, L. & Del Giorgio, P. A. Links between metabolic plasticity and

652   functional redundancy in freshwater bacterioplankton communities. *Front. Microbiol.* **4**,

653   1–11 (2013).

654 46. Herndl, G.J., Kaltenbok, E., Muller-Niklas, G. Dialysis bag incubation as a non

655   radiolabeling technique to estimate bacterioplankton production in situ. *Handbook of*

656   *Methods in Aquatic Microbial Ecology.* 495-503. Lewis Publisher, Florida (1993).

657 47. Wang, L. C. *et al.* Increased precipitation during the Little Ice Age in northern Taiwan

658   inferred from diatoms and geochemistry in a sediment core from a subalpine lake. *J.*

659   *Paleolimnol.* **49**, 619–631 (2013).

660 48. Tsai, J. W. *et al.* Seasonal dynamics, typhoons and the regulation of lake metabolism in a

661   subtropical humic lake. *Freshw. Biol.* **53**, 1929–1941 (2008).

662 49. Chen, C. P., Tseng, C. H., Chen, C. A. & Tang, S.-L. The dynamics of microbial

663   partnerships in the coral Isopora palifera. *ISME J.* **5**, 728–740 (2011).

664 50. Jorgenson, S.L. *et .al.* Correlating microbial community profiles with geochemical data in

665   highly stratified sediments from the Arctic Mid-Ocean Ridge. *Proc. Natl. Acad. Sci.*

666   **109**(42), 2846-2855. (2012).

667 51. Schloss, P. D. *et al.* Introducing mothur: Open-source, platform-independent, community

supported software for describing and comparing microbial communities. *Appl. Environ.*

Microbiol. **75**, 7537–7541 (2009).

52. Edgar, R. C. Search and clustering orders of magnitude faster than BLAST.

Bioinformatics **26**, 2460–2461 (2010).

53. Edgar, R. C. UPARSE: highly accurate OTU sequences from microbial amplicon reads.

Nat. Methods **10**, 996–998 (2013).

54. Quast, C. *et al.* The SILVA ribosomal RNA gene database project: improved data

processing and web-based tools. *Nucleic Acids Res.* **41**, D590–D596 (2013).

55. Yilmaz, P. *et al.* The SILVA and "All-species Living Tree Project (LTP)"

taxonomic frameworks. *Nucleic Acids Res.* **42**, D643-8 (2014).

56. Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads.

EMBnet.journal 17, 10–12 (2011).

57. Shen, W., Le, S., Li, Y. & Hu, F. SeqKit: A cross-platform and ultrafast toolkit for

FASTA/Q file manipulation. *PLoS One* **11**, 1–10 (2016).

58. Hyatt, D., Chen, G.L., Locascia, P.F., Land, M.L.,Larimer, F.W., & Hauser, L.J. Prodigal:

prokaryotic gene recognition and translation initiation site identificatin. *BMC*

Bioinformatics **11**, 119. (2010).

59. Seemann, T. Prokka: Rapid prokaryotic genome annotation. *Bioinformatics* **30**, 2068–

2069 (2014).

60. Magoč, T. & Salzberg, S. L. FLASH: Fast length adjustment of short reads to improve

genome assemblies. *Bioinformatics* **27**, 2957–2963 (2011).

61.   Buchfink, B., Xie, C. & Huson, D. H. Fast and sensitive protein alignment using

DIAMOND. *Nat. Methods* **12**, 59–60 (2014).

62.   Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. Basic local alignment

search tool. *J. Mol. Biol.* **215**, 403–410 (1990).

63.   Marchler-Bauer, A. *et al.* CDD: NCBI's conserved domain database. *Nucleic Acids Res.*

43, D222–D226 (2015).

64.   R Developmetn Core Team. R: A Language and Environment for Staistical Compting. R

Foundation for Statistical Computing. Vienna, Austria. (2016).

65.   McMurdie, P. J. & Holmes, S. Phyloseq: An R Package for Reproducible Interactive

Analysis and Graphics of Microbiome Census Data. *PLoS One* **8**, (2013).

66.   Okansen, J. *et al.* vegan: Community Ecology Package: Ordination, Diversity and

Dissimilarities. R package 2.5-5 (2011).

67.   Wickham, H. ggplot2: Elegant Graphics for Data Analysis. R package version 3.2.0.

Spinger-Verlag New York. (2016)

68.   Kandlikar, G. S. *et al.* Ranacapa: An R package and shiny web app to explore

environmental DNA data with exploratory statistics and interactive visualizations [version

1; referees: 1 approved, 2 approved with reservations]. *F1000Research* **7**, 1–19 (2018).

69.   Maechler, M. Rousseeuw, P., Struyf, A., Hubert, M., Hornik, K. cluster: Cluster Analysis

Basics and Extensions. R package version 2.1.0. (2019).

70.   Lin, J. Divergence Measures Based on the Shannon Entropy. *IEEE Trans. Inf. Theory* **37**,

145–151 (1991).

71.   Mooshammer, M. *et al.* Decoupling of microbial carbon, nitrogen, and phosphorus cycling
in response to extreme temperature events. *Sci. Adv.* **3**, 1–14 (2017).

72.   Orland, C. *et al.* Microbiome functioning depends on individual and interactive effects of
the environment and community structure. *ISME J.* **13**, 1–11 (2019).

73.   Debroas, D. *et al.* Metagenomic approach studying the taxonomic and functional diversity
of the bacterial community in a mesotrophic lake (Lac du Bourget - France). *Environ.*
Microbiol. **11**, 2412–2424 (2009).

74.   Shade, A. *et al.* Resistance, resilience and recovery: Aquatic bacterial dynamics after
water column disturbance. *Environ. Microbiol.* **13**, 2752–2767 (2011).

75.   Baho, D. L., Peter, H. & Tranvik, L. J. Resistance and resilience of microbial communities
- Temporal and spatial insurance against perturbations. *Environ. Microbiol.* **14**, 2283–
2292 (2012).

76.   Shade, A. *et al.* Lake microbial communities are resilient after a whole-ecosystem
disturbance. *ISME J.* **6**, 2153–2167 (2012).

77.   Tandon, K. *et al.* Bacterial community in water and air of two sub-alpine lakes in taiwan.
Microbes Environ. **33**, 120–126 (2018).

78.   Sunagawa, S. *et al.* Structure and function of the global ocean microbiome. *Science (80-.*
). **348**, 1261359–1261359 (2015).

79.   Langenheder, S., Lindström, E. S. & Tranvik, L. J. Structure and function of bacterial
communities emerging from different sources under identical conditions. *Appl. Environ.*
Microbiol. **72**, 212–220 (2006).

80.   Pimm, S. L. The complexity and stability of ecosystems. *Nature* **307**, 321–326 (1984).

81.   Hou, D. *et al.* Environmental factors shape water microbial community structure and

733   function in shrimp cultural enclosure ecosystems. *Front. Microbiol.* **8**, 1–12 (2017).

734   82.   Eiler, A. Evidence for the ubiquity of mixotrophic bacteria in the upper ocean:

735   Implications and consequences. *Appl. Environ. Microbiol.* **72**, 7431–7437 (2006).

736   83.   Meyer, A. F., Lipson, D. A., Martín, A. P., Schadt, C. W. & Schmidt, S. K. Molecular and

737   Metabolic Characterization of Cold-Tolerant Alpine Soil Pseudomonas Sensu Stricto.

738   *Appl. Environ. Microbiol.* **70**, 483–489 (2004).

739   84.   Swingley, W. D. *et al.* The complete genome sequence of Roseobacter denitrificans

740   reveals a mixotrophic rather than photosynthetic metabolism. *J. Bacteriol.* **189**, 683–690

741   (2007).

742   85.   Tranvik, L.J. & Hofle, M.G. bacterial Growth in Mixed Cultures on Dissoloved Organic

743   Carbon from Humic and Clear Waters. *Appl. Environ. Micrbiol.* **53**, 482-488 (1987).

744   85.   Rosenfeld, J. S. & Mall, M. Functional redundancy in ecology and conservation. *Fish.*

745   *Res.* **1**, (2002).

746   86.   Ruiz-Gonzalez, C., Nino-Garcia, J. P., Lapierre, J. F. & del Giorgio, P. A. The quality of

747   organic matter shapes the functional biogeography of bacterioplankton across boreal

748   freshwater ecosystems. *Glob. Ecol. Biogeogr.* **24**, 1487–1498 (2015).

 Figure Legends

 Figure 1: Schematic representation of the experiment. 1A) Sampling of background (BG)

 samples from the surface and bottom of Yuan-Yang and Tsuei-Feng Lakes (a, b, c, d). **1B)** Self-

 swap scheme with sample representation (e, f, g, h). **1C)** Cross-Swap scheme with sample

 representation (i, j, k, l). **1D)** Timeline for sample collection and the overall duration of the

 experiment.

 Figure 2: Principal Coordinate Analysis (PCoA) plots. 2A) 16S rRNA gene-based bacterial

 community composition was significantly different between the two lakes, and cross-swap

 samples clustered tightly with their self-swap counterparts**. 2B)** Functional profile (read-based

 COG-Family) derived from whole-metagenome sequencing was also significantly different

 between the swapped samples from the two lakes. Dotted grey arrows represent the shift in

 cross-swap samples away from their self-swap counterparts, indicating that the local

 environment had an effect on the functional profile.

 Figure 3: Community function relationship and *L*-divergence as a measure to quantify the

 functional redundancy. 3A) The relationship between the similarities in communities' (16S

 rRNA gene transcript-based) overall functional attributes (PICRUSt2-derived imputed

 metagenome) and community composition of surface ($R^2 = 0.65$, $p < 0.01$, F test for the overall

 significance of the linear regression) and bottom samples ($R^2 = 0.63$, $p < 0.01$, F test for the

 overall significance of the linear regression). **3B)** Heat maps based on *L*-divergence

 measurements for community and overall functional attributes of surface and bottom samples.

 Figure 4: Effect of average well color development (AWCD) as a measure of metabolic activity

 of the community with a change in the local environment**. A)** and **B)** TFL self-swap samples

 (Ts(n)→T; Tb(n)→T) compared with cross-swap samples (Ts(n)→Y; Tb(n)→Y). **C)** and **D)**

775   TFL self-swap samples (Ts(n)→T; Tb(n)→T) compared with cross-swap samples (Ys(n)→T;

776   Yb(n)→T). Significant difference tested with Student's t-test $p < 0.05$. Error bars indicate

777   standard deviation.

A

Surface

Bottom

$R^2 = 0.65$, P <0.01

$R^2 = 0.63$, P <0.01

Similarity in communitie's functional attributes

Similarity in communtie's functional attributes

Similarity in community composition

Similarity in community composition

B

Community

Function