Challenging a preconception: Optoacoustic spectrum differs from the absorption spectrum of proteins and dyes for molecular imaging.

Juan Pablo Fuenzalida Werner,[†¥] Yuanhui Huang,[†,§ ¥] Kanuj Mishra,[†,§] Andriy Chmyrov,[†,§,‖] Vasilis Ntziachristos,[†,§,‖] and Andre C. Stiel[,†]*

[†] Institute of Biological and Medical Imaging (IBMI), Helmholtz Zentrum München, D-85764 Neuherberg, Germany
[§] Chair of Biological Imaging, Technische Universitat München, D-81675 Munich, Germany
[‖] Center for Translational Cancer Research (TranslaTUM), Technische Universitat Müchen, D-81675 Munich, Germany
[¥] *These authors contributed equally.*

Abstract

Despite the tremendous technological advances in detection and data analysis in optoacoustic (OA) imaging, there is no detailed knowledge and understanding of the photophysics of OA signal generation of commonly used contrast agents, such as dyes and chromoproteins. This gap blocks the further development of dedicated labels for optoacoustics. To close it, we developed a multi-modal laser spectrometer (MLS) to enable the simultaneous measurement of OA, absorbance, and fluorescence spectra. MLS provides reproducible, high-quality OA spectra by using correction and referencing workflow. Herein, we employ MLS to analyze several common dyes (Methylene Blue, Rhodamine 800, Alexa Fluor 750, IRDye 800CW and Indocyanine green) and proteins (sfGFP, mCherry, mKate, HcRed, iRFP720 and smURFP) and shed light on their internal conversion properties. Our data shows that the absorption spectra do not correlate with the OA spectra for the majority of the analytes. We determine that for dyes, the transition underlying the high energy shoulder, which mostly correlates with an aggregation state of the dyes, has significantly more OA generation efficiency than the monomer transition. Our analyses for proteins point to a favored vibrational relaxation and OA signal generation that stems from the neutral or zwitterionic chromophores. Such data is highly relevant for the engineering of tailored contrast agents for OA imaging. Furthermore, discrepancies between absorption and OA spectra underline the importance of correct spectral information as a prerequisite for the spectral-unmixing schemes that are often required for *in vivo* OA imaging. Finally, OA-spectra recorded on our MLS of some of the most commonly used proteins and dyes in optical imaging reveal previously unknown photophysical characteristics, such as unobserved photoswitching behavior.

Introduction

Optoacoustic (OA, also termed photoacoustic) imaging combines optical contrast with ultrasound resolution, enabling resolved, real-time *in vivo* imaging well-beyond the 1 mm penetration depth of typical optical methods [1]. Therefore, it is emerging as a particularly appealing *in vivo* imaging method to study tumor biology [2], inflammatory diseases [3], developmental processes [4], or brain functioning in mammals *in vivo* [5].

However, in the majority of such studies, OA imaging relies entirely on endogenous absorbing-agents, like blood-hemoglobin, melanin, or lipids. In contrast, targeted labels enable longitudinal, cell-specific *in vivo* imaging, drastically expanding the potential applications of OA imaging [6]. Several potential agents have been tested for OA imaging, including nanoparticles and targeted dyes (reviewed in [7]), as well as transgenic labels, such as fluorescent proteins, bacteriophytochromes, and their switchable variants (reviewed in [8]).

In some cases, these labels were specifically developed for OA applications [9,10]. However, despite these successes, most label choices and the strategies for their development are still based on OA performance and photophysics inferred from independent transmission-mode absorption and fluorescence spectroscopy data. As shown by our previous work and others, low quantum yield and high molar absorption coefficient do not automatically translate to high optoacoustic signal[11].

In this work, we sought to develop a method of spectral analysis that combines absorption and fluorescence with OA spectral data to enable the full characterization of photophysical parameters of OA labels. Such a method would help to identify competing transitions (e.g., triplet states, long-lived dark states, or molecular isomerization; see **Figure 1a**). This data will become increasingly important to accelerate the tailored development of OA contrast agents. Furthermore, setting standards and building a reliable database of spectral information could guide researchers in their choice of suitable labels, similar to the "molecular brightness" standard used in fluorescence imaging. Lastly, a combined method of spectral analysis would enable the study of various photophysical phenomena, for example, dye aggregation, exciton coupling, and energy conversion.

We have developed and present herein a multi-modal laser spectrometer (MLS) that allows us to measure optoacoustic, absorption, and fluorescence spectra simultaneously with high precision and spectral resolution. Key characteristics of this system are i) illumination with pulsed lasers, similar to OA imaging; ii) homogenization of fluence for all measured wavelengths; iii) in-line reference and correction to overcome laser instabilities; iv) simultaneous detection of fluorescence excited by the laser pulse, and v) simultaneous absorbance measurement. In contrast to other devices for recording OA spectra, which required samples with high concentrations and/or large volumes [12–14], our MLS requires only 200 µL of minimum 0.2 optical density (OD) concentration for high-quality, reproducible spectra ($R^2 > 0.95$). This high spectral quality allowed us the study of a range of dyes and chromoproteins, namely: Methylene Blue, Rhodamine 800, Alexa Fluor 750, IRDye 800CW and Indocyanine green, as well as sfGFP, mCherry, mKate, HcRed, iRFP720 and smURFP. In most cases, we show significant differences between absorbance and OA spectra. We correlated these discrepancies either to an aggregation state of the dyes or to the isomeric state of the chromophores. Finally, we discuss how such high-quality MLS spectra can empower future optoacoustic contrast agent development and spectral unmixing to achieve high detection sensitivity in molecular imaging.

Results & Discussion

Multi-modality laser spectrometer (MLS)
In order to fully characterize the photophysical parameters underlying the OA signal of a particular sample, it is essential to obtain exact measurements of absorbance, fluorescence, and OA spectra under the same light fluency to assess the relative contributions of the different electronic transitions. Based on expertise gained measuring the OA kinetic parameters of photo-controllable proteins [15], we developed a multi-modal laser spectrometer (MLS, **Figure 1b**) that can record all three modalities simultaneously in the wavelength range between 420 - 1000 nm. In brief (**Figure 1b**), for the OA and fluorescence spectroscopic measurements, we used an optical parametric oscillator (OPO) laser with a 50 Hz repetition rate and average 7 ns pulse width as the light source. The laser energy at different wavelengths (420-1000 nm) was kept constant by controlling a motor-controlled half-wave plate (HWP) in combination with a polarizing beam-splitter (PBS) according to a lookup table of pulse energies at all wavelengths. The laser energy was measured by an in-line power meter before every measurement. The light was delivered to the sample submerged in water through a fiber bundle. The light travels first through a 200 μm thick in-line reference chamber filled with Indian ink (0.2-0.7 OD) before reaching a similar chamber containing the actual sample. The absorbance of ink was determined relative to the absorbance of the sample in order to optimize the dynamic range of our data acquisition system. The laser excited optoacoustic wave is recorded using a cylindrically focused ultrasound transducer with a central frequency of 3.5 MHz. The signal is amplified by a variable-gain amplifier (Amp) and then digitized by a 12-bit digitizer (DAQ) and PC. The fluorescence signal, which is excited by the same laser pulse, is recorded simultaneously by a fiber-coupled diode-array spectrometer oriented 45° to the laser beam path. The absorption is recorded using the same spectrometer an instant after the OA and fluorescence measurements by blocking the laser beam with an electronically controlled shutter (E. Shutter) and opening the shutter of a halogen lamp to illuminate the sample with a fiber opposite to the spectrometer.

An essential part of the MLS performance is a stepwise correction procedure enabling highly accurate OA spectral measurements. In brief (**Figure 1c**, Supplementary **Figure 1-3**): An initial laser energy look-up table is computed to correct for the major wavelength-dependent changes in laser energy. This table is then used to control the HWP and beam splitter to maintain the same energy at each wavelength. We adopted a measure-and-fit routine to measure the lookup table and designed an iteration-based lookup table update method. This afforded a fast lookup table measurement and precise control of the laser energy. However, the response time of the power meter is too low to ensure the correction of pulse-to-pulse energy fluctuations. Furthermore, while the peak power variations might be correctible with a fast-response power-meter, the variation in pulse width is harder to measure and correct with high-precision. Thus, for pulse-to-pulse correction, we rely on the signal from the in-line ink-reference to deliver a time-shifted signal from the same excitation light as the actual sample. This method of correction necessitates the subtraction of the ink absorbance spectrum from the fluctuation corrected spectrum. We further corrected for the spectral coloring induced

by in-line reference material (ink) and the water path of laser, to reduce possible fluence alterations.

Characterization of the MLS
To validate the optics setup and the correction steps, we analyzed the spectra of $NiCl_2$. $NiCl_2$ is known to have no apparent photo-fatigue and a complete transfer of absorbed energy into acoustic waves via non-radiative decay channels, so its absorption spectrum should correlate with its optoacoustic spectrum [16]. The OA spectra of $NiCl_2$ measured without any correction (R^2 0.988 and 7.9%) and using the full correction protocol of our MLS (R^2 0.997 and 1.9%) demonstrate the increase of spectral quality, with each correction step (**Figure 1c**). The improvement of spectral quality through our measurement protocol is significant over a range of concentrations, with an R^2 of 0.85 for a $NiCl_2$ sample OD as low as 0.05 (**Figure 1d to f**). In order to test the wavelength accuracy of our setup, we measured OA and absorbance spectra of potassium permanganate (**Figure 1g**), and we were able to match its 6 absorption peaks with 2 nm accuracy correctly. To further characterize our MLS system, we similarly measured different concentrations of Brilliant Black N dissolved in PBS (BBN, **Figure 1h, and i**). BBN, similar to $NiCl_2$, has a negligible fluorescence quantum yield (QY_{fluo}) and very high photostability, and thus undergoes fully non-radiative deexcitation. Importantly, BBN, in contrast to $NiCl_2$, is soluble and stable in aqueous buffer solutions. BBN thus shares a common Grüneisen parameter [16–18] with labels for *in vivo* imaging, which are frequently dissolved in aqueous buffer systems. The combination of the excellent solubility of BBN and its stable generation of OA signals over a range of concentrations and excitation energies makes it well suited to serve as a universal standard for precise and reproducible OA measurements. Such allows reproducible characterization of dyes and other labels eventually allowing the determination and reporting of universally meaningful OA signal strength values.

OA signal generation
Vibrational relaxation (non-radiative energy decay) from an excited state produces a change in pressure, resulting in an optoacoustic signal (OAS, **Figure 1a**). This change in pressure is generally described by equation (2) [19].

$$p_0 = \Gamma \mu_a \Phi \left(1 - \Phi_{fl}\right) \qquad (2)$$

In equation 2, μ_a is the local optical absorption coefficient, Φ is the light fluence, and Φ_{fl} is the fluorescence quantum yield. Γ is the dimensionless Grüneisen parameter, which represents the thermoelastic properties of the medium. Equation 2 dismisses all other competitive channels, like conformational changes, photochemical transformations, intramolecular charge transfer, electron transfer, intersystem crossing, proton transfer, exciplex formation, excimer formation, and energy transfer [20] (**Figure 1a**). All of these processes reduce the number of available electrons that can undergo vibrational relaxation and produce a change in pressure in the time scale that the transducers and the illumination pulse length used in imaging devices can detect [21–23]. Thus, in order to accurately compare different optoacoustic capacities, the OAS has to be normalized by the absorbance, so that the slope of the linear relation between the

OAS and absorbance yields the photoacoustic generation efficiency (PGE), which is a measure of how much of the absorbed energy is converted to pressure [12,17,18,24,25].

$$\text{PGE} = \frac{p_0}{\mu_a} = \Phi \, \Gamma \, (1 - \Phi_{fl}) \, (1 - X) \qquad (3)$$

As is shown in equation 3, the PGE is attenuated by fluorescence (Φ_{fl}) and any other competitive pathway (X). As mentioned before, photostability and lack of any fluorescence make BBN an ideal reference for photoacoustic measurements; accordingly, we assign it a PGE value of 1.0, assuming that no other decay channels take place [17,18].

Characterization of dyes

The sensitivity of some natural chromophores and organic dyes to aggregation and other environmental changes has been exploited for various sensing applications in OA imaging [26–30]. However, the photophysics underlying the changes in the OA signals of these chromophores have never been thoroughly understood or described, precluding efficient and rational exploitation of such effects for the design of next-generation functional OA labels. Methylene Blue (MB) and Rhodamine 800 (Rh800) are two well-known synthetic xanthene dyes. MB has been commonly employed as a label in OA due to its lack of fluorescence [31]. Rh800 would be even more favorable for OA imaging due to its more pronounced red absorption (682 nm vs. 660 nm); however, its PGE is attenuated by strong fluorescence at 712 nm [32]. Both dyes have the tendency to form aggregates through strong aromatic interactions. The aggregates have photophysical characteristics that are clearly distinguishable from those of the monomers and can be distinguished based on additional spectral bands next to the monomers' absorbance.

The H-aggregate, with the dye molecules stacked in the same orientation, shows a blue-shifted absorbance, while the J-aggregate, in which the molecules are shifted with respect to each other, shows a more red-shifted absorbance [33]. The formation of H-type aggregates commonly reduces fluorescence significantly, while J-type aggregates can fluoresce [33–35]. Intersystem crossing (ISC) rates are also determined by the supramolecular architecture and orientation geometry of the aggregates [33,35]. In general, the aggregation behavior is strongly depending on environmental conditions like pH, ionic strength, and crowding agents [36,37]. In order to create a common defined medium for our measurements that resemble the cellular environment, we dissolved the dyes in 10% fetal bovine serum (FBS) in phosphate buffer saline (PBS) [38]. In this environment, all dyes are in a mixed aggregation state, which allows us to study the different characteristics of the aggregate related transitions. Even more critically, this provides stable environments that enable us to observe the spectra without changes over time.

For solutions of MB with concentrations ranging from 17 to 29 µM, we found fluorescence emission only from excitation of the red-shifted spectral band, corresponding to the weakly fluorescent monomer, while the blue-shifted band,

associated predominantly with the H-aggregates, did not show fluorescent transitions (**Figure 2a**). Conversely, the OA spectrum shows a comparably stronger signal for the H-aggregate band (610 nm), reflecting a higher OA signal generation (photoacoustic generation efficiency, PGE) for this transition compared to the monomer (665nm, 0.61 vs. 0.37, **Figure 2b)**. It is generally accepted that the loss of fluorescence in highly symmetric H-aggregates, like the one formed by MB, stems from a favored ISC [27,33] (**Supplementary Figure 4**). This is corroborated by our analysis of the photo-fatigue of the different transitions, which shows faster bleaching for the H-aggregate at 610 nm (**Supplementary Figure 5**) than for the monomer at 665 nm. However, the vibronic relaxation following ISC, regularly occurring in the µs time frame, is too slow to contribute to OA signals in our observation time window of 5 ns, and thus would not explain the higher PGE of the H-aggregate. Interestingly, the ISC for MB is very fast (ns vs. µs) [27], suggesting that we can indeed detect the resultant vibronic relaxations. Consequently, we can show here that the formation of H-aggregates by MB results in fast ISC, which contributes to the observed strengthening of the OA signal.

Because the dye Rh800 fluoresces, it has thus far not been considered for use in OA applications; however, the increase in the PGE of MB that we observed upon its aggregation prompted us also to study the photophysics of Rh800. We measured Rh800 in a concentration range from 30 to 49 µM. Rh800 shows a similar combination of monomer (695nm) and predominantly H-aggregate (635 nm) bands, similar to MB (**Figure 2c**). However, in contrast to MB, the PGEs of the two bands are only slightly different (1.14 and 1.08, **Figure 2d**). To our surprise, the PGE of the monomer form is high, despite its fluorescence, even surpassing our standard BBN (**Figure 1h and i**). Moreover, the bleaching rates of both transitions are moderate and suggest no strong ISC, as also observed for MB (**Supplementary Table 1, Supplementary Figure 5**). Upon closer inspection of the spectral relations, it becomes apparent that the fluorescence excitation spectrum is shifted in respect to the absorbance band at 695 nm, hinting at a third transition, presumably a non-fluorescent J-aggregate, in the flank of the peak primarily attributed to monomer (720 nm, arrow). Further inspection of the spectra reveals an additional shoulder in the blue region (580 nm, **Figure 2c arrow**). Such high energy transitions in xanthene dyes are often correlated with higher-order aggregates and have been previously observed [39]. This is additionally underpinned by a pronounced H-aggregate band suggesting that most of the dye is in an aggregate state. The acoustic wave for Rh800 at 695nm and 635nm differ from each and differ from the one of BBN,(**Supplementary Figure 6**). it is likely that this behavior is because the formation of these high-order aggregates alters the size of the acoustic emitter and that replacing the aqueous surrounding of the monomeric dye molecules alter Grüneisen parameter. This favors the generation of OA signals upon deexcitation, thus explaining the unusually high PGE. Moreover, the low bleaching rate (10% signal loss after 18 exaphotons,(**Supplementary Figure 5**) suggests only minor contributions from transitions competing with the non-radiative decay (like fluorescence or ISC), further explaining the high PGE.

Alexa Fluor 750 (Al750) and IRDye 800CW (800CW) show absorbance peaks centered in the favorable blood window (749 nm and 774 nm, **Figure 2e, and g**), which is why

they have been developed for *in vivo* fluorescence imaging applications. Despite their relatively high fluorescence (QY_{fluo} = 0.12 for both AL750 and 800CW), both dyes have already been used in OA imaging [40,41]. Analysis of the spectra of Al750 shows a shift for the excitation maximum as well as the PGE maximum of 10 nm relative to the absorbance spectrum. This points to a transition at ca. 698 nm (**Figure 2e**), blue shoulder, arrow A). Additionally, this shoulder bleaches considerably less than the main peak, with only 40% loss versus 80% after 60000 light pulses for the shoulder and main peak, respectively (**Supplementary Figure 5**). This low bleaching and effective deexcitation in comparison to the main peak of Al750 points to a strong triplet tendency of the main peak leaving the transition in the shoulder as the more effective one for imaging. Similar to the xanthene dyes, the absorbance spectrum 800CW has two peaks (**Figure 2g**) with one centered at 710 nm and the second one centered at 770 nm. The fluorescence excitation spectrum mirrors the primary absorption peak (770 nm), with a stronger PGE in the blue shoulder (710nm), attributable to H-aggregates. Despite evidence of 800CW showing excitation coupling with other aromatic dyes [42,43], there is no description in the literature of aggregation of 800CW molecules. The observed behavior, spectrally similar to MB and Rh800 described above, might point to the blue shoulder resulting from H-type aggregation for 800CW. Intuitively, the formation of self-aggregates in CW800 seems to be unfavorable due to CW800s' strong negative character (four negatively charged sulfonate groups). However, it has been noted that several other similarly charged dyes undergo aggregation and exciton coupling in the presence of macromolecules, such as polymers, HSA, or BSA [44,45] Thus, the FBS that is present in our solutions could lead to the formation of H-aggregates of CW800, explaining the observed exciton coupling behavior.

Indocyanine green

Indocyanine green (ICG) is a dye widely used in OA due to its relatively strong signal intensity, favorable absorbance at 800 nm, and FDA approval[46–49]. The zwitterionic and hydrophobic character of ICG allows its aggregation in an aqueous solvent at very low concentrations[50,51]. Furthermore, it has been shown that proteins, such as albumin, can modify the aggregation state of ICG[38,51]. Similar to the previous dyes, we prepared and measured solutions of ICG with concentrations ranging from 2 to 26 µM, in 10% FBS. OA and absorbance spectra show a constant mismatch (**Figure 3 a and b**, arrow), with the whole spectral shape exhibiting a redshift and an increase of the 745 nm band with increasing concentration. The PGE for the OA shoulder centered at 745 is 0.80. Interestingly the absorbance at 795 nm does not correlate linearly with the OAS. To compensate for this non-linear behavior, we used two windows for the calculation of the PGEs, allowing a linear fit (**Figure 3c**). The PGE at 795 nm (0.62) is significantly lower at concentrations below 10 µM than it is above 10 µM (0.87), suggesting a higher PGE for aggregates of ICG. Thus, the difference between the dimer/monomer ratio from absorbance and OA represents the aggregation tendency of ICG (**Figure 3d**). The inflection point of the sigmoidal curve at 9.2 µM is between the two linear regimes chosen for PGE calculation. The PGE increases at the monomer peak (795 nm) with increasing concentration because the fraction of ICG aggregates that also absorb at this wavelength increases. The discrepancies between absorbance and OA spectra observed could arise from different species of ICG present primarily at low concentrations, such as monomer, monomer bound to proteins (FBS), dimer, and dimer bound to proteins, as

previously suggested [51]. At higher concentrations, aggregated ICG interacting with proteins becomes the predominant species, the acoustic and absorption spectra of which do not vary significantly. Our results for ICG prove clearly that aggregation increases optoacoustic signal generation. Beyond encouraging the production of nanoparticles with highly aggregated ICG for optoacoustic imaging, this also suggests the use of the aggregation states as ratiometric readout.

Characterization of GFP-type chromophore bearing proteins
Genetically encode labels are a prerequisite for longitudinal imaging and one of the reasons for fluorescence imaging becoming a standard tool in life sciences. GFP-like labels are prominent in fluorescence imaging and, despite their mostly visible absorbance profiles, have also been studied and employed as OA labels[6,12,52–54]. However, biliverdin bearing chromoproteins like Bacteriophytochromes (BphPs) or Phycobiliproteins (Pbp) are advantageous due to their near-infrared absorbance (NIR) and have already been employed prominently in OA imaging[10,11,54–57]. The hydroxybenzylidene imidazolidone chromophores of GFP-like proteins share complex photophysics involving different protonation states of the chromophore (neutral, zwitterionic, and anionic) and a plethora of additional effects like excited-state proton transfer (ESPT) and long-lived dark states (often involving isomerization along the methine bridge). For a review, see [58].

We focused our spectral analysis on sfGFP (an advanced variant of the original *Aequorea victoria* GFP [59]), red mKate2 [60], mCherry [61], tdTomato [61], and far-red non-fluorescent HcRed [62]. Comparing the OA and absorption spectra of all these proteins reveals that in almost all cases (except for tdTomato), the blue shoulder of the peak is pronounced in OA, sometimes even leading to a blue-shift of the OA maxima compared to the absorption. Specifically, sfGFP shows a maximum for absorbance and OA at 480 nm; however, the shoulder in the blue region of the OA spectra is more prominent, including an identifiable slope rising at ~425 nm for the neutral chromophore (**Figure 4a and b**). This is also reflected in a higher PGE for the blue transition compared to the main peak (425 nm: 0.75; 480 nm: 0.34). mKate2 has an optoacoustic spectrum that is 5 nm blue-shifted for the absorption spectra (585 nm vs. 580 nm) and matching PGEs of 550 nm: 0.64 and 580 nm: 0.51(**Figure 4c and d**). For mCherry, the maximum is similarly blue-shifted compared to the absorbance (580 nm vs. 585 nm), but with the corresponding PGE only slightly higher for the blue shoulder than for the main peak (580 nm: 0.54; 585 nm: 0.53) (**Figure 5e and f**). For the far-red and non-fluorescent tetrameric protein HcRed, the optoacoustic spectrum is blue-shifted by 20 nm with respect to the absorption spectra, with the peak center at 570 nm instead of 590 nm (**Figure 5g and h**). Accordingly, the PGEs at 570 nm and 590 nm are 0.97 and 0.80, respectively. In general, except again for tdTomato, the PGEs of the main peaks are in good agreement with reported fluorescent quantum yields (**Supplementary Table 1**). tdTomato, on the other hand, a protein related to mCherry, is the only example of a protein measured in our study that shows an OA spectrum that is highly displaced to the red. tdTomato has a maximum in the OA spectrum at 565 nm, rather than 555 nm, with a PGE of 0.79 at 485 nm and of 0.83 at 565 nm (**Figure 5i and j**).

Several explanations are possible for the higher PGE of the blue shoulder observed in most of the proteins. i) The blue shift could point to an involvement of the neutral state of the chromophore (commonly called an A state, as opposed to the anionic B form). For several FPs, fast (ps) ESPT from the excited state of the neutral form (A*) to an excited intermediate state (I*) are possible [63,64]. It is possible that deexcitation from this state results in vibronic relaxation and a higher PGE. However, the excitation spectra for sfGFP, mKate, and mCherry have shapes that are similar to the optoacoustic spectra, all with significant pronunciation in the shorter wavelength range. Additionally, the fluorescent lifetimes for sfGFP exciting at 420 and 490 nm show only a slightly longer lifetime at 490 nm (2.77 vs. 2.37 ns), suggesting similar deexcitation compared to the B form. ii) For HcRed, our data is in agreement with the theory of a mixture of cis-trans isomers, with a nonfluorescent, neutral trans-isomer centered at around 570 nm and a slightly fluorescent cis-isomer centered at 590 nm [65,66]. The high PGE from the peak at 570 nm could arise from the non-fluorescent properties of the trans-isomer and, independently, the tetrameric character of HcRed promoting exciton coupling between nearby chromophores, as observed in other multimeric proteins[67]. iii) Beyond our previous explanation, discrepancies between OA and absorption spectra have been observed before and explained by ground state depopulation [12]. In short, when a fluorescent molecule is excited by early arriving photons with the pulsed illumination used in OA (5 - 7 ns), and if the fluorescence lifetime of the molecule is long enough, later arriving photons have a lower number of ground state electrons for excitation. This effect primarily come into play for the strongly absorbing central transitions of an absorption spectra, leading to the "flattened" absorption peaks observed by Laufer and colleagues [12]. Since in our study the discrepancies are regularly found at the flanks of the spectra or only for certain peaks (e.g., aggregate peaks or protonated chromophores), we believe the phenomenon of ground state depopulation does not explain our results. Interestingly, however, we can see that excitation spectra in figure 4 show ground state depopulation and loss of the spectral shape with a difficult to differentiate main peak. This might be possible since electrons that decay trough the radiative pathway have slower rate constants than those that decay through internal conversion (ns vs. ps) [22]. Thus, if ground state depopulation due to the intense short pulse irradiation is present, it will primarily affect the radiative transitions and thus the excitation spectra.

Another peculiarity, which we observed by recording spectra under conditions regularly used for OA imaging, is that several proteins showed a level of photoconversion or photo-switching that was previously unknown. This was initially observed when comparing forward (420 to 900 nm) and reverse (900 to 420 nm) modes of collecting spectral data. For mCherry and tdTomato, the forward and reverse measurements produced different results, which suggests long-lived states induced by the illumination. The effects are different for the two proteins, but always show a long-lived change of the OA spectra. For mCherry, measuring from 420 to 900 nm results in a stronger blue shoulder than measuring in the opposite direction (**Figure 5a**). Surprisingly, the absorption spectrum of mCherry does not change accordingly (**Figure 5b**), resulting in an increase of the PGE of the shoulder from 0.35 to 0.45. mCherry does not show evidence of phototransformation after continued illumination with 550 or 580 nm light (**Figure 5b**). For tdTomato, in contrast to mCherry, the shoulder increases when imaged from 900 to 420 nm with a change in PGE for both peaks (**Figure 5d and 4j**).

Surprisingly, the maxima of the OA spectra also shift. Alternating illumination between 480 nm and 550 nm shows how the absorption spectra changed with an increase of the band at 480 nm, which can be attributed to the neutral chromophore in those proteins (**Figure 5e and 5f**). Both mCherry, as well as tdTomato, are related to the photo-switchable proteins rsCherry and rsCherryRev [68], which primarily show fluorescence based photo-switching, but also some level of photochromism (more so for rsCherry). It could be that the effects in mCherry and tdTomato under strong pulsed laser illumination are similar to photo-switching events, although we see no dark-relaxation, typical of reversible photo-switching.

For the far-red protein, HcRed, continuous illumination at 570 nm did not produce a decrease in intensity (bleaching) as expected, but an increase in signal – an effect also observed for rsCherry [68]. However, in contrast to rsCherry, we did not observe an inflection point, even after 60,000 pulses. Alternating illumination between 565 nm and 590 nm results in a reduction in the signal at the fluorescence peak at 590, but an increase at the optoacoustic peak at 565nm; the absorption spectra show the same kinetics (**Figure 5g-i**). Trans to cis-isomerization of HcRed has been previously induced by heating [66], but until now, there was no report of light-induced changes in this protein. While such phenomena are interesting from a perspective of FP photophysics, they can also be useful for OA imaging, since the transitions can be used similar to photo-switching for locked-in OA detection [23]. In this regard, tdTomato is of especially high interest, since a vast number of established lines and transgene animals exist with this protein due to its excellent properties for fluorescence imaging. It is all the more astonishing because in our measurements, tdTomato showed a high PGE of 0.8, which does not agree with its long fluorescence lifetime of around 3.6 ns or its reported quantum yield of 0.66. This discrepancy remains to be elucidated, but is favorable in this particular case for both OA and fluorescence imaging.

Biliverdin binding proteins are of critical importance in optoacoustic imaging, as their absorption in the near-infrared region gives them superior penetration depth and spectral separation for endogenous contrast. Until now, only two classes of near-infrared biliverdin binding proteins have been reported: bacteriophytochromes and phycobiliproteins [69,70]. IRFP720 and smURFP are examples of both classes. In both cases, the optoacoustic spectra match the absorption spectra, with PGEs of 0.56 and 0.86 for smURFP and IRFP720 at their primary OA peaks, respectively (**Fig. 4k and m, respectively**). The lower PGE of smURFP correlates with its higher quantum yield 0.2, versus 0.1 for IRFP. Additional reductions in PGE could come from faster bleaching and enhanced transient isomerization, a phenomenon already reported for biliverdin [71,72]. The excitation spectrum of smURFP shows evidence of ground state depopulation; even more so than for the classical proteins, ground state depopulation is not apparent in the optoacoustic spectrum (**Figure 4k**). Instead, IRFP shows a shoulder whose transition is less excitable and has a higher PGE (0.97) (**Figure 4n**).

Conclusion

Our first results using the MLS showed a wealth of insight into the behavior of a number of well-used dyes and proteins. Describing the exact changes in the photophysics of OA signal generation upon the aggregation of MB, Rh800, and ICG will afford new applications for sensors that exploit their environment-dependent aggregation. For example ratiometric analysis of dye aggregation under different physiological conditions. A similar strategy is used in measurements of blood oxygenation with two wavelengths [73].

Furthermore, determining which transitions are most effective for OA imaging enables dyes like Alexa Fluor 750 to be employed more effectively for this modality. Finally, our results on ICG shed light on the complex behavior of this molecule, which is so regularly used in OA studies [46,47]. Cognizance of the strong dependence of PGE on the aggregation of dyes could help researchers to analyze concentration-dependent measurements of ICG more accurately. On the protein side, the switching or conversion behavior of mCherry, tdTomato, and HcRed is extremely interesting for GFP-like protein research and warrants further study; however, it also has an immediate impact on employing those proteins in OA applications. In particular, tdTomato – for which there are numerous animal models from fluorescence studies – can be used more effectively for OA measurements by exploiting the transition from 550 nm to 480 nm and using a dual-wavelength ratio analysis to identify the protein-label in tissue with a strongly absorbing background.

Lastly, we could show that accurate measurements for OA labels are a prerequisite for their understanding, their future design, and especially their comparison. Regarding this, the use of standards like BBN for novel OA contrast agents to come is highly important for comparability, and for empowering researchers to make strategic choices on what labels to use in their experiments.

Material and Methods:

Preparation of dyes
Dyes were dissolved at 5 mg/ml in DMSO and frozen; previous to measurement, a stock solution of the dyes 1% v/v was prepared in 10% FBS. The dye was further diluted to provide different concentrations. The absorbance spectra of each concentration were measured several times during one hour to assure no change in the absorption spectra over time.

Preparation of Protein
Proteins were expressed in Escherichia coli BL21 and purified by Ni-NTA affinity chromatography, followed by gel filtration on a HiLoad 26/600 Superdex 75pg column (Amersham Biosciences) in PBS buffer. Purified proteins were frozen immediately in liquid nitrogen and stored at −80 °C. Thawed proteins were centrifuged at 14,000 rpm for 45 min at 4 °C, and the supernatant was used for the measurements. *Measurements*

Optoacoustic Measurements
200uL samples placed in IBDI ship were illuminated by a nanosecond excitation pulses were generated by an optical parametric oscillator (OPO) laser (Spitlight-DPSS 250 ZHG-OPO, InnoLas). Constant pulse energy was ensured by use of a half-wave plate in a motorized rotation stage (PRM1Z8, Thorlabs) and a polarizing beam splitter; using a lookup table and adapting

the polarization with the half-wave plate, we kept the power constant at 1.2 mJ for the proteins measurements and 1 mJ for the dyes, over the whole illumination spectrum. Optoacoustic signals were detected with a cylindrically focused single-element transducer (V382-SU, 3.5 MHz, Olympus) and averaged over 3-10 laser pulses to improve the signal-to-noise ratio (SNR). We confirmed the linearity and stability of the optoacoustic signal generated by BBN and NiCl2 in the range of fluences used in our experiments. The optoacoustic signals were amplified by 60 dB via a wide-band voltage amplifier (DHPVA-100, Femto) and digitized at 100 MS/s with a data acquisition card (RZE-002 400, GaGe). Absorption spectra were recorded after each excitation wavelength by fast interruption of the laser pulse illumination with an electronic shutter (SHB1, Thorlabs) and opening the shutter of a fiber-coupled broadband white-light illumination lamp (DH-2000, Ocean Optics). Triggering for both shutters was provided by microcontroller (Arduino Uno). Detection of the transmission spectra was provided by a commercial UV–vis spectrometer (USB 4000, Ocean Optics). The setup has a time delay for the absorption measurement of at least 110 ms after the laser pulses due to shutter responses. Both the illumination and the detection fiber were attached to a collimator rotated by 45° to the optoacoustic illumination-detection axis.

Since fluorescence and optoacoustic signal was excited by the same laser pulses, both can be recorded simultaneously. We set the UV-vis spectrometer in external triggering mode for excitation/emission spectrum measurement and shared the trigger as optoacoustic recording. Due to the coincidence of sweeping excitation laser and fluorescence emission, there was overlapping region of the two in a direct recording. To extract the excitation spectrum only, for example, from the recordings, we used the normalized readings of three different emission wavelengths at each excitation wavelengths. When the excitation wavelength was overlapping with one of the selected emission wavelengths, we ignored the current emission reading and used the extrapolated value averaged from the other two emission wavelengths. Similar method was adopted to extract the emission spectrum from such overlapping recordings.

OA laser spectrometer

Building upon previous art [15], we developed a multi-modal laser spectrometer (MLS, Figure 1b) that allows us to record all three modalities simultaneously in the wavelength range between 420 - 1000 nm. Figure 1b schematized the optical system consisting of 1. A diode-pumped solid-state optical parametric oscillator (DPSS OPO, SpitLight DPSS250 OPO, Innolas Laser GmbH, Germany) laser covering wavelength range 420 - 2000 nm with a 50 Hz pulse repetition rate (PRR), a pulse width at full width at half maximum τ_{FWHM} =7 ns, and a ~10 nm linewidth of specific wavelength output. The OPO signal beam of 420-709 nm is output with horizontal polarization and idler beam 710-2000 nm with vertical polarization (switching controlled by a stepper motor and a microcontroller Arduino Uno); 2. A pulse energy control to maintain constant pulse energy, which is measured using pyroelectric sensor head PE10-C and USB readout power meter Juno (Ophir-Spiricon LLC, USA), using a combination of a motorized half-wave plate (HWP, AHWP05M-600; mounted on motorized rotation stage PRM1Z8; Thorlabs Inc., USA), a polarizing beam splitter (PBS052, Thorlabs Inc., USA), and a beam dump; 3. An optical filter wheel that switches between long-pass (LP, cut-off wavelength at 700 nm for OPO idler beam) and short-pass (SP, cut-off at 800 nm for signal beam) edge filters that are switched by an Arduino Uno via a stepper motor, in order to remove the possible leakage of signal/idler at output beam; 4. An electrically controlled diaphragm shutter (SHB05T, Thorlabs Inc., USA) that is triggered by an Arduino Uno; And 5. a fiber bundle and its coupling lens (numerical aperture, NA 0.22; 430 fibers bundled in ϕ4.6 mm; CeramOptec GmbH, Germany) to deliver light to a deionized water chamber which is determined for optimal coupling of optoacoustic wave.

In the water chamber, we custom-built a holder to fix the relative position of all components, including 1. The Ibidi chips (μ-Slide I Luer, 200 μm channel thickness, ibidi GmbH, Germany)

for sample and reference materials (water solution of Indian ink). The in-line reference chip is inserted in the beam path to remove the influence of laser pulse energy fluctuation of the sample spectrum. 2. Multi-mode fibers and collimating lens that delivers light from a deuterium tungsten lamp (DH-2000-BAL, Ocean Optics, USA) and collects the transmitted light through the sample to a Vis-NIR spectrometer (USB4000-VIS-NIR, Ocean Optics, USA), and an ultrasound transducer (V382-SU, central frequency 3.5 MHz, focus length 38.1 mm, diameter 13 mm) that simultaneously detects optoacoustic wave generated from reference and sample chips. The detected temporal optoacoustic signals were amplified by a 10-60 dB variable-gain amplifier (DHPVA, FEMTO Messtechnik GmbH, Germany) and sampled by a 12-bit digitizer (DAQ, Gage CSE1222, DynamicSignals LLC, USA). At each wavelength 10 sequences are acquired for signal averaging. All data acquisition, processing and setup synchronization are controlled by our custom-built programs using Matlab 2015b (The MathWorks Inc., USA). Empowered by the MLS spectrometer, four spectra of a sample can be obtained simultaneously: optoacoustics, absorbance, fluorescence excitation, and emission. When measuring optoacoustic spectrum and fluorescence spectra, the electrical shutter is opened, while the shutter to the lamp of Vis-NIR spectrometer is closed, and the ultrasound transducer, DAQ, and spectrometer (configured to external triggering mode and fixed integration time of 10 ms) are triggered by the laser to acquire data. When measuring absorbance, only the spectrometer acquires data (configured to circular-buffered acquisition mode using integration time that uses 90% of the full pixel well depth when making dark/no sample measurement), and laser light is blocked by the closed shutter and lamp shutter is open to deliver white light to the sample. One challenge in measuring with high-fidelity on common dyes/proteins based chromophores is that they suffer from photobleaching and such processes are usually wavelength dependent.

Lookup table measurement for laser pulse energy
To ensure a controlled constant delivery of pulse energy to the sample across the 420-1000 nm range, the operation of MLS starts with a pulse energy lookup table measurement. Supplementary Figure 1 shows a complete flow chart on our high-precision laser pulse energy lookup table measurement method. The lookup table is intended to reflect the sinusoidal oscillation relation between the angle of HWP and the pulse energy of OPO at certain wavelengths. Therefore, we used a measure-and-fit scheme (Eq. 1) to fasten the speed of lookup table measurement.

$$E(\lambda, \theta) = A_0(\lambda) \cdot \left[\frac{1}{2} + sin(\frac{2\pi}{180^0} \theta + \theta_0) \right] (1)$$

where, $E(\lambda, \theta)$ is the pulse energy measured at laser wavelength when HWP is at angle θ; $A_0(\lambda)$ is the maximum pulse energy that is achievable only when HWP angle is at θ_0 that coincides with the transmission polarization of PBS.
According to Eq. 1 (shown in Supplementary Figure 1), we first measured the energy at one OPO signal wavelength, e.g., at 420 nm with HWP angle rotating with step 2° across 72° angles (in order to include one peak and one trough). Then we fit the energy-angle curve to the parametric sinusoidal model as indicated by Eq. 1, to find the phase angle θ_0 where the peak energy appears. Fixing HWP at the peak phase angle θ_0, we measure the peak energy $A_0(\lambda)$ of the rest signal wavelengths, and based on which we extrapolate the sinusoids model. We show in Supplementary Figure 2a an exemplary laser pulse energy lookup table measured using our measure-and-fit based scheme. The same procedure is applied to OPO idler wavelengths (≥ 710 nm) to obtain a crude estimate of the lookup table. But due to the fluctuation of OPO laser pulse, especially at wavelengths around 710 nm where signal/idler beam are separated, the extrapolated lookup table required still correction. Aiming at a pulse energy of 1.0 mJ at all wavelengths, we measure the pulse energy back using the angles suggested by the lookup table targeting at 1.0 mJ, and find the ratio of the targeted energy and the new measurements. Then we multiply this ratio by the amplitude $A_0(\lambda)$ to scale the lookup table proportionally. Usually,

less than 5 iterations of this lookup table correction enable a root mean square error (RMSE) of ~2% at a specified pulse energy, as shown in Supplementary Figure 2b.

Principle of in-line correction in optoacoustic spectrometer

Another challenge resides with the pulse-to-pulse fluctuation at given wavelength, which can range from a few percent to more than 30% as at the transition wavelengths of signal/idler output (Supplementary Figure 1b). To minimize the influence of such fluctuation, we designed an in-line reference using water solution of Indian ink in optoacoustic sensing. The resulting OA signals from both reference and sample are collected by the ultrasound transducer in the same temporal sequence. As shown in Figure 1b, laser light fluence φ_0 with fluctuation Δfl traveling a water path d_{FR} from fiber to reference chip pass through the low concentration of ink with a light fluence φ_R according to Beer-Lambert's law:

$$\varphi_R = e^{-d_{FR} \cdot \mu_{H2O}} \cdot \varphi_0 (1 + \Delta fl) = e^{-d_{FR} \cdot \mu_{H2O}} \cdot \varphi_0 \cdot \frac{(1+\Delta E) \cdot E_0}{[(1+\Delta\sigma) \cdot \sigma_0]^2} \quad (2)$$

where μ_{H2O} is water absorbance. It is important to note that the pulse-to-pulse light fluence fluctuation Δfl manifests as both a peak power (E_0) variation (ΔE) and a pulse width (σ_0) variation ($\Delta\sigma$), assuming Gaussian temporal profile. The pulse width variation (sub-nanosecond for 7 ns pulse width we used) is a term cannot be corrected using a slow-responding power meter for optoacosutics, but only possible by using high-speed photodiode (<100 picosecond) and fast acquisition system (>20 GHz), which is a costly. Instead, we used optoacoustic reference. The transmitted light after reference chip with absorbance μ_{aR} and path d same to channel thickness of Ibidi chip then travels a water path d_{RS} from reference chip to sample chip. The incidence light fluence to the sample, φ_S, is:

$$\varphi_S = e^{-d \cdot \mu_{aR}} \cdot e^{-d_{RS} \cdot \mu_{H2O}} \cdot \varphi_R \quad (3)$$

The photoacoustic signal generation induced by optical absorption and the subsequent thermal expansion can be described by a simplified model [74], $p_0 = \gamma\mu_a\varphi$, assuming a linear relation between absorption coefficient and light fluence in low absorbance and low scattering sample [75], Γ is the Grüneisen coefficient indicating the conversion efficiency from heat to pressure wave. Therefore, the optoacoustic wave generated by the reference chip is:

$$p_R = \Gamma_R \cdot \mu_{aR} \cdot \varphi_R \quad (4)$$

Similarly, the sample generated optoacoustic wave is:

$$p_S = \Gamma_S \cdot \mu_{aS} \cdot \varphi_S \quad (5)$$

Dividing sample optoacoustic signal by reference can cancel the influence of laser pulse-to-pulse fluctuation if substituting in the light fluence to reference (Eq. 2) and sample (Eq. 3).

$$\frac{p_S}{p_R} = \frac{\Gamma_S \cdot \mu_{aS} \cdot \varphi_S}{\Gamma_R \cdot \mu_{aR} \cdot \varphi_R} = \Gamma_{S/R} \cdot \frac{\mu_{aS}}{\mu_{aR}} \cdot e^{-d \cdot \mu_{aR}} \cdot e^{-d_{RS} \cdot \mu_{H2O}} \quad (6)$$

As a consequence, Eq. 6 indicates that fluctuation correction renders a mixed spectrum of the reference and sample. The first term on the right-hand side of Eq.6 is a ratio of Grüneisen coefficients of sample over reference. The second term is absorption coefficient ratio of sample over reference. The third term indicates the spectral coloring effect [76] (light fluence alteration) by the in-line reference. The fourth term represents the spectral coloring by the water path between reference and sample chips.

There are ways to account for the effect of above-mentioned terms. The absorption coefficient μ_{aR} of reference solution can be obtained by measuring its absorbance A (cm^{-1}) and converting using $\mu_{aR} = ln(10) \cdot A$, assuming low scattering by a low concentration/absorbance of ink. Thus multiplying the spectrometer measured μ_{aR} and Eq.6 would remove the influence by ink spectra. Furthermore, although the effect is minor, the spectral coloring effect by ink can be removed, given the thickness, d Ibidi chip channel thickness. Similarly, given water path d_{RS} between reference and sample chip and water absorption coefficient (data adapted from http://www.spectra.arizona.edu/), the spectral coloring effect by water can be removed. The corrected optoacoustic signal at wavelength as a result is:

$$p_S(\lambda) = \Gamma_{S/R} \cdot \mu_{aS}(\lambda) \qquad (7)$$

where the Grüneisen coefficient ratio $\gamma_{S/R}$ is not supposed to be wavelength dependent thus a scaling factor of optoacoustic signal amplitude. For reproducible comparison across various samples, if we use ink as reference for all the samples, the Grüneisen coefficient ratio still reflects the heat conversion efficiency of samples.

Data acquisition and process for multi-modal laser spectroscopic measurement

Supplementary Figure 3 shows the program for data acquisition and data process in multi-modal laser spectroscopy. Due to our in-line referencing to correct for the laser pulse-to-pulse fluctuation, we applied similar process to fluorescence emission data as to optoacoustic data.

The computation of the spectra (blue shading part showed in Supplementary Figure 3) is done as described in the following sequential steps: taking a sample of e.g. NiCl$_2$ with 0.5 OD at 720 nm. The results are shown in Figure 1c, providing an examination on the improvement brought by each of the processes in comparison with the raw spectrum (Raw (Max) in Figure 1c) which is acquired by taking simply the maximum value measured at each wavelength. Matching the absorbance spectrum, the Raw spectra showed a coefficient of determination R^2 of 0.988 and a standard deviation 7.9% between repeated recording, owing to the 2%-RMSE laser energy control (Supplementary Figure 2). The temporal signal at 720 nm is shown in Fig. 1e indicating the improvement by the correction process on signal level, in which due to Hilbert transform-based envelope detection, the corrected signal showed only absolute value.

1. *'Filtered.'* We firstly filtered the acquire temporal sequence by using a 4th order digital bandpass filter in temporal frequency band 0.1-10 MHz respecting the bandwidth of our transducer, a 1-D digital wavelet transform (decomposition level=5) based denoising method to reduce thermal noise level, a 3rd order Savitzky-Golay filter with smoothing length corresponding to 20 MHz in our DAQ (200 MSamples/second) to reduce digitization error.

2. *'Hilbert'*. Hilbert transform is used to detect the peak envelope of temporal sequence. The optoacoustic spectrum relies on the maximum signal in the sequence at position (time-of-flight) of sample and reference. We detected the peak signals by summing up all sequences measured at all wavelengths, then finding the locations of peaks of sample and reference respectively, and using the peak locations as center point to take averaged values of adjacent 3-5 sample points from sequence of each wavelength. A better signal-to-noise ratio (SNR) of optoacoustic spectra for low concentration/absorbance sample was achieved using this way than taking area under curve of signal envelope.

3. *'- Dark'*. We subtracted from the raw signal with the root mean square of a dark background signal that was measured with no laser irradiation before each spectral measurement to reduce the noise floor and thus improve the detection sensitivity of low concentration sample.

4. *'Pulse Fl.'* To reduce the influence of laser pulse-to-pulse energy fluctuation, we divided the sample spectrum acquired at each laser pulse by the corresponding reference spectrum. But this step introduces spectra of ink into the sample spectra.

5. *'- Ink Spect.'* To remove the ink spectra, we multiplied the fluctuation corrected spectra with normalized spectra of ink to recover the sample spectra.

6. *'- Coloring'*. Notwithstanding that it is minor effect in 420-900 nm range induced by the spectral coloring by ink (short absorption path) and water path between reference and sample (water absorbance at 900 nm is 0.025 cm-1), we corrected for spectral colorings. The improvement was not reflected in the coefficient of determination R^2 or standard deviation of measurements, but was instead shown as a higher fidelity in the peak value at 420 nm.

Acknowledgement

YH wants to address special thanks to Hong Yang, to Dr. Ara Ghazaryan, and to Dr. Jan G. Laufer for inspiring discussions.

References

(1)   Ntziachristos, V. Going Deeper than Microscopy: The Optical Imaging Frontier in Biology. *Nature Methods*. 2010. https://doi.org/10.1038/nmeth.1483.

(2)   Omar, M.; Schwarz, M.; Soliman, D.; Symvoulidis, P.; Ntziachristos, V. Pushing the Optical Imaging Limits of Cancer with Multi-Frequency-Band Raster-Scan Optoacoustic Mesoscopy (RSOM). *Neoplasia* **2015**. https://doi.org/10.1016/j.neo.2014.12.010.

(3)   Aguirre, J.; Schwarz, M.; Garzorz, N.; Omar, M.; Buehler, A.; Eyerich, K.; Ntziachristos, V. Precision Assessment of Label-Free Psoriasis Biomarkers with Ultra-Broadband Optoacoustic Mesoscopy. *Nat. Biomed. Eng.* **2017**. https://doi.org/10.1038/s41551-017-0068.

(4)   Vetschera, P.; Koberstein-Schwarz, B.; Schmitt-Manderbach, T.; Dietrich, C.; Hellmich, W.; Chekkoury, A.; Symvoulidis, P.; Reber, J.; Westmeyer, G.; López-Schier, H.; et al. Beyond Early Development: Observing Zebrafish over 6 Weeks with Hybrid Optical and Optoacoustic Imaging. *BioRxiv* **2019**. https://doi.org/10.1101/586933.

(5)   Ovsepian, S. V.; Olefir, I.; Westmeyer, G.; Razansky, D.; Ntziachristos, V. Pushing the Boundaries of Neuroimaging with Optoacoustics. *Neuron*. 2017. https://doi.org/10.1016/j.neuron.2017.10.022.

(6)   Razansky, D.; Distel, M.; Vinegoni, C.; Ma, R.; Perrimon, N.; Köster, R. W.; Ntziachristos, V. Multispectral Opto-Acoustic Tomography of Deep-Seated Fluorescent Proteins in Vivo. *Nat. Photonics* **2009**, *3* (7), 412–417. https://doi.org/10.1038/nphoton.2009.98.

(7)   Gujrati, V.; Mishra, A.; Ntziachristos, V. Molecular Imaging Probes for Multi-Spectral Optoacoustic Tomography. *Chem. Commun.* **2017**, *53* (34), 4653–4672. https://doi.org/10.1039/C6CC09421J.

(8)   Brunker, J.; Yao, J.; Laufer, J.; Bohndiek, S. E. Photoacoustic Imaging Using Genetically Encoded Reporters: A Review. *J. Biomed. Opt.* **2017**, *22* (7), 70901.

(9)   Li, Y.; Forbrich, A.; Wu, J.; Shao, P.; Campbell, R. E.; Zemp, R. Engineering Dark Chromoprotein Reporters for Photoacoustic Microscopy and FRET Imaging. *Sci. Rep.* **2016**, *6* (February), 22129. https://doi.org/10.1038/srep22129.

(10)   Chee, R. K. W.; Li, Y.; Zhang, W.; Campbell, R. E.; Zemp, R. J. In Vivo Photoacoustic Difference-Spectra Imaging of Bacteria Using Photoswitchable Chromoproteins. *J. Biomed. Opt.* **2018**, *23* (10), 1–11. https://doi.org/10.1117/1.JBO.23.10.106006.

(11)   Fuenzalida Werner, J. P.; Mishra, K.; Huang, Y.; Vetschera, P.; Glasl, S.; Chmyrov,

A.; Richter, K.; Ntziachristos, V.; Stiel, A. C. Structure-Based Mutagenesis of Phycobiliprotein SmURFP for Optoacoustic Imaging. *ACS Chem. Biol.* **2019**. https://doi.org/10.1021/acschembio.9b00299.

(12)   Laufer, J.; Jathoul, A.; Pule, M.; Beard, P. In Vitro Characterization of Genetically Expressed Absorbing Proteins Using Photoacoustic Spectroscopy. *Biomed. Opt. Express* **2013**, *4* (11), 2477–2490. https://doi.org/10.1364/BOE.4.002477.

(13)   Teng, Y. C.; Royce, B. S. H. Absolute Optical Absorption Coefficient Measurements Using Photoacoustic Spectroscopy Amplitude and Phase Information. *J. Opt. Soc. Am.* **1980**. https://doi.org/10.1364/josa.70.000557.

(14)   Schneider, S.; Coufal, H. Influence of Intermolecular Energy Transfer on Photoacoustic Spectra. *J. Chem. Phys.* **1982**. https://doi.org/10.1063/1.443374.

(15)   Vetschera, P.; Mishra, K.; Fuenzalida-Werner, J. P.; Chmyrov, A.; Ntziachristos, V.; Stiel, A. C. Characterization of Reversibly Switchable Fluorescent Proteins in Optoacoustic Imaging. *Anal. Chem.* **2018**. https://doi.org/10.1021/acs.analchem.8b02599.

(16)   Laufer, J.; Zhang, E.; Beard, P. Evaluation of Absorbing Chromophores Used in Tissue Phantoms for Quantitative Photoacoustic Spectroscopy and Imaging. *IEEE J. Sel. Top. Quantum Electron.* **2010**, *16* (3), 600–607. https://doi.org/10.1109/JSTQE.2009.2032513.

(17)   Brown, J. E.; Diaz, L.; Christoff-Tempesta, T.; Nesbitt, K. M.; Reed-Betts, J.; Sanchez, J.; Davies, K. W. Characterization of Nitrazine Yellow as a Photoacoustically Active PH Reporter Molecule. *Anal. Chem.* **2015**, *87* (7), 3623–3630. https://doi.org/10.1021/ac503515k.

(18)   Abbruzzetti, S.; Viappiani, C.; Murgida, D. H.; Erra-Balsells, R.; Bilmes, G. M. Non-Toxic, Water-Soluble Photocalorimetric Reference Compounds for UV and Visible Excitation. *Chem. Phys. Lett.* **1999**, *304* (3–4), 167–172. https://doi.org/10.1016/S0009-2614(99)00306-1.

(19)   Deán-Ben, X. L.; Gottschalk, S.; Mc Larney, B.; Shoham, S.; Razansky, D. Advanced Optoacoustic Methods for Multiscale Imaging of in Vivo Dynamics. *Chem. Soc. Rev.* **2017**, *46* (8), 2158–2198. https://doi.org/10.1039/C6CS00765A.

(20)   Land, E. J. THE TRIPLET EXCITED STATES OF BILIVERDIN AND BILIVERDIN DIMETHYL ESTER. *Photochem. Photobiol.* **1979**. https://doi.org/10.1111/j.1751-1097.1979.tb07079.x.

(21)   Schaberle, F. A.; Rego Filho, F. D. A. M. G.; Reis, L. A.; Arnaut, L. G. Assessment of Lifetime Resolution Limits in Time-Resolved Photoacoustic Calorimetry vs. Transducer Frequencies: Setting the Stage for Picosecond Resolution. *Photochem. Photobiol. Sci.* **2016**. https://doi.org/10.1039/c5pp00397k.

(22)   S. P. McGlynn, T. A. and M. K. *Molecular Spectroscopy of the Triplet State*; Prentice-Hall: New Jersey, 1969.

(23)   Mishra, K.; Fuenzalida-Werner, J. P.; Ntziachristos, V.; Stiel, A. C. Photocontrollable Proteins for Optoacoustic Imaging. *Anal. Chem.* **2019**.

https://doi.org/10.1021/acs.analchem.9b01048.

(24) Rosencwaig, A. Photoacoustic Spectroscopy. *Annu. Rev. Biophys. Bioeng.* **1980**, *9* (1), 31–54. https://doi.org/10.1146/annurev.bb.09.060180.000335.

(25) Braslavsky, S. E.; Ellul, R. M.; Weiss, R. G.; Al-Ekabi, H.; Schaffner, K. Photoprocesses in Biliverdin Dimethyl Ester in Ethanol Studied by Laser-Induced Optoacoustic Spectroscopy (Lioas). *Tetrahedron* **1983**. https://doi.org/10.1016/s0040-4020(01)88704-4.

(26) Jiang, Y.; Pu, K. Advanced Photoacoustic Imaging Applications of Near-Infrared Absorbing Organic Nanoparticles. *Small (Germany)* **2017**, *13* (30). https://doi.org/10.1002/smll.201700710.

(27) Morgounova, E.; Shao, Q.; Hackel, B. J.; Thomas, D. D.; Ashkenazi, S. Photoacoustic Lifetime Contrast between Methylene Blue Monomers and Self-Quenched Dimers as a Model for Dual-Labeled Activatable Probes. *J. Biomed. Opt.* **2013**, *18* (5), 56004. https://doi.org/10.1117/1.JBO.18.5.056004.

(28) Peters, L.; Weidenfeld, I.; Klemm, U.; Loeschcke, A.; Weihmann, R.; Jaeger, K.-E.; Drepper, T.; Ntziachristos, V.; Stiel, A. C. Phototrophic Purple Bacteria as Optoacoustic in Vivo Reporters of Macrophage Activity. *Nat. Commun.* **2019**, *10* (1), 1191. https://doi.org/10.1038/s41467-019-09081-5.

(29) Wang, J.; Jeevarathinam, A. S.; Humphries, K.; Jhunjhunwala, A.; Chen, F.; Hariri, A.; Miller, B. R.; Jokerst, J. V. A Mechanistic Investigation of Methylene Blue and Heparin Interactions and Their Photoacoustic Enhancement. *Bioconjug. Chem.* **2018**, *29* (11), 3768–3775. https://doi.org/10.1021/acs.bioconjchem.8b00639.

(30) Lovell, J. F.; Jin, C. S.; Huynh, E.; Jin, H.; Kim, C.; Rubinstein, J. L.; Chan, W. C. W.; Cao, W.; Wang, L. V; Zheng, G. Porphysome Nanovesicles Generated by Porphyrin Bilayers for Use as Multimodal Biophotonic Contrast Agents. *Nat. Mater.* **2011**, *10* (4), 324–332. https://doi.org/10.1038/nmat2986.

(31) Taruttis, A.; Ntziachristos, V. Advances in Real-Time Multispectral Optoacoustic Imaging and Its Applications. *Nat Phot.* **2015**, *9* (4), 219–227. https://doi.org/10.1038/nphoton.2015.29.

(32) Alessi, A.; Salvalaggio, M.; Ruzzon, G. Rhodamine 800 as Reference Substance for Fluorescence Quantum Yield Measurements in Deep Red Emission Range. *J. Lumin.* **2013**, *134*, 385–389. https://doi.org/10.1016/j.jlumin.2012.08.017.

(33) Kasha, M.; Rawls, H. R.; Ashraf El-Bayoumi, M. The Exciton Model in Molecular Spectroscopy. *Pure Appl. Chem.* **1965**, *11* (3–4). https://doi.org/10.1351/pac196511030371.

(34) Hestand, N. J.; Spano, F. C. Expanded Theory of H- and J-Molecular Aggregates: The Effects of Vibronic Coupling and Intermolecular Charge Transfer. *Chem. Rev.* **2018**, *118* (15), 7069–7163. https://doi.org/10.1021/acs.chemrev.7b00581.

(35) Hestand, N. J.; Spano, F. C. Molecular Aggregate Photophysics beyond the Kasha Model: Novel Design Principles for Organic Materials. *Acc. Chem. Res.* **2017**, *50* (2), 341–350. https://doi.org/10.1021/acs.accounts.6b00576.

(36)   Moreno-Villoslada, I.; Fuenzalida, J. P.; Tripailaf, G.; Araya-Hermosilla, R.; Pizarro, G. D. C.; Marambio, O. G.; Nishide, H. Comparative Study of the Self-Aggregation of Rhodamine 6G in the Presence of Poly(Sodium 4-Styrenesulfonate), Poly(N - Phenylmaleimide- Co -Acrylic Acid), Poly(Styrene- Alt -Maleic Acid), and Poly(Sodium Acrylate). *J. Phys. Chem. B* **2010**. https://doi.org/10.1021/jp104340k.

(37)   Krieger, A.; Fuenzalida Werner, J. P.; Mariani, G.; Gröhn, F. Functional Supramolecular Porphyrin-Dendrimer Assemblies for Light Harvesting and Photocatalysis. *Macromolecules* **2017**, *50* (9). https://doi.org/10.1021/acs.macromol.6b02435.

(38)   Bongsu Jung; Vullev, V. I.; Anvari, B. Revisiting Indocyanine Green: Effects of Serum and Physiological Temperature on Absorption and Fluorescence Characteristics. *IEEE J. Sel. Top. Quantum Electron.* **2014**, *20* (2), 149–157. https://doi.org/10.1109/JSTQE.2013.2278674.

(39)   Moreno-Villoslada, I.; Torres-Gallegos, C.; Araya-Hermosilla, R.; Nishide, H. Influence of the Linear Aromatic Density on Methylene Blue Aggregation around Polyanions Containing Sulfonate Groups. *J. Phys. Chem. B* **2010**, *114* (12), 4151–4158. https://doi.org/10.1021/jp909105r.

(40)   Lakshman, M.; Needles, A. Screening and Quantification of the Tumor Microenvironment with Micro-Ultrasound and Photoacoustic Imaging. *Nat. Methods* **2015**, *12* (4), iii–v. https://doi.org/10.1038/nmeth.f.381.

(41)   Chen, Z.; Deán-Ben, X. L.; Gottschalk, S.; Razansky, D. Performance of Optoacoustic and Fluorescence Imaging in Detecting Deep-Seated Fluorescent Agents. *Biomed. Opt. Express* **2018**, *9* (5), 2229–2239. https://doi.org/10.1364/BOE.9.002229.

(42)   Joseph, J.; Baumann, K. N.; Koehler, P.; Zuehlsdorff, T. J.; Cole, D. J.; Weber, J.; Bohndiek, S. E.; Hernández-Ainsa, S. Distance Dependent Photoacoustics Revealed through DNA Nanostructures. *Nanoscale* **2017**, *9* (42), 16193–16199. https://doi.org/10.1039/c7nr05353c.

(43)   Zhang, J.; Smaga, L. P.; Satyavolu, N. S. R.; Chan, J.; Lu, Y. DNA Aptamer-Based Activatable Probes for Photoacoustic Imaging in Living Mice. *J. Am. Chem. Soc.* **2017**, *139* (48), 17225–17228. https://doi.org/10.1021/jacs.7b07913.

(44)   Hoffmann, S.; Fuenzalida Werner, J. P.; Moreno-Villoslada, I.; Goycoolea, F. M. New Insights into the Nature of the Cibacron Brilliant Red 3B-A - Chitosan Interaction. In *Pure and Applied Chemistry*; Walter de Gruyter GmbH, 2016; Vol. 88, pp 891–904.

(45)   Liang-Schenkelberg, J.; Fieg, G.; Waluga, T. Molecular Insight into Affinity Interaction between Cibacron Blue and Proteins. *Ind. Eng. Chem. Res.* **2017**, *56* (34), 9691–9697. https://doi.org/10.1021/acs.iecr.7b01556.

(46)   Beziere, N.; Lozano, N.; Nunes, A.; Salichs, J.; Queiros, D.; Kostarelos, K.; Ntziachristos, V. Dynamic Imaging of PEGylated Indocyanine Green (ICG) Liposomes within the Tumor Microenvironment Using Multi-Spectral Optoacoustic Tomography (MSOT). *Biomaterials* **2015**, *37*, 415–424. https://doi.org/10.1016/j.biomaterials.2014.10.014.

(47)   Wilson, K. E.; Bachawal, S. V; Willmann, J. K. Intraoperative Resection Guidance

with Photoacoustic and Fluorescence Molecular Imaging Using an Anti-B7-H3 Antibody-Indocyanine Green Dual Contrast Agent. *Clin. Cancer Res.* **2018**, *24* (15), 3572–3582. https://doi.org/10.1158/1078-0432.CCR-18-0417.

(48) Weber, J.; Beard, P. C.; Bohndiek, S. E. Contrast Agents for Molecular Photoacoustic Imaging. *Nat. Methods* **2016**, *13* (8), 639–650. https://doi.org/10.1038/nmeth.3929.

(49) Kim, C.; Song, K. H.; Gao, F.; Wang, L. V. Sentinel Lymph Nodes and Lymphatic Vessels: Noninvasive Dual-Modality in Vivo Mapping by Using Indocyanine Green in Rats--Volumetric Spectroscopic Photoacoustic Imaging and Planar Fluorescence Imaging. *Radiology* **2010**, *255* (2), 442–450. https://doi.org/10.1148/radiol.10090281.

(50) Zweck, J.; Penzkofer, A. Microstructure of Indocyanine Green J-Aggregates in Aqueous Solution. *Chem. Phys.* **2001**, *269* (1–3), 399–409. https://doi.org/10.1016/S0301-0104(01)00368-8.

(51) Philip, R.; Penzkofer, A.; B?umler, W.; Szeimies, R. M.; Abels, C. Absorption and Fluorescence Spectroscopic Investigation of Indocyanine Green. *J. Photochem. Photobiol. A Chem.* **1996**, *96* (1–3), 137–148. https://doi.org/10.1016/1010-6030(95)04292-X.

(52) Liu, C.; Gong, X.; Lin, R.; Liu, F.; Chen, J.; Wang, Z.; Song, L.; Chu, J. Advances in Imaging Techniques and Genetically Encoded Probes for Photoacoustic Imaging. *Theranostics* **2016**, *6* (13), 2414–2430. https://doi.org/10.7150/thno.15878.

(53) Scheibe, K.; Kersten, C.; Schmied, A.; Vieth, M.; Primbs, T.; Carlé, B.; Knieling, F.; Claussen, J.; Klimowicz, A. C.; Zheng, J.; et al. Inhibiting Interleukin 36 Receptor Signaling Reduces Fibrosis in Mice with Chronic Intestinal Inflammation. *Gastroenterology* **2019**, *156* (4), 1082-1097.e11. https://doi.org/10.1053/j.gastro.2018.11.029.

(54) Filonov, G. S.; Krumholz, A.; Xia, J.; Yao, J.; Wang, L. V.; Verkhusha, V. V. Deep-Tissue Photoacoustic Tomography of a Genetically Encoded near-Infrared Fluorescent Probe. *Angew. Chem. Int. Ed. Engl.* **2012**, *51* (6), 1448–1451. https://doi.org/10.1002/anie.201107026.

(55) Yao, J.; Wang, L.; Yang, J.-M.; Maslov, K. I.; Wong, T. T. W.; Li, L.; Huang, C.-H.; Zou, J.; Wang, L. V. High-Speed Label-Free Functional Photoacoustic Microscopy of Mouse Brain in Action. *Nat. Methods* **2015**, *12* (5), 407–410. https://doi.org/10.1038/nmeth.3336.

(56) Märk, J.; Dortay, H.; Wagener, A.; Zhang, E.; Buchmann, J.; Grötzinger, C.; Friedrich, T.; Laufer, J. Dual-Wavelength 3D Photoacoustic Imaging of Mammalian Cells Using a Photoswitchable Phytochrome Reporter Protein. *Commun. Phys.* **2018**, *1* (1), 3. https://doi.org/10.1038/s42005-017-0003-2.

(57) Li, L.; Shemetov, A. A.; Baloban, M.; Hu, P.; Zhu, L.; Shcherbakova, D. M.; Zhang, R.; Shi, J.; Yao, J.; Wang, L. V.; et al. Small Near-Infrared Photochromic Protein for Photoacoustic Multi-Contrast Imaging and Detection of Protein Interactions in Vivo. *Nat. Commun.* **2018**, *9* (1), 2734. https://doi.org/10.1038/s41467-018-05231-3.

(58) van Thor, J. J. Photoreactions and Dynamics of the Green Fluorescent Protein. *Chem. Soc. Rev.* **2009**, *38* (10), 2935–2950. https://doi.org/10.1039/b820275n.

(59)   Pédelacq, J. D.; Cabantous, S.; Tran, T.; Terwilliger, T. C.; Waldo, G. S. Engineering and Characterization of a Superfolder Green Fluorescent Protein. *Nat. Biotechnol.* **2006**. https://doi.org/10.1038/nbt1172.

(60)   Shcherbo, D.; Murphy, C. S.; Ermakova, G. V.; Solovieva, E. A.; Chepurnykh, T. V.; Shcheglov, A. S.; Verkhusha, V. V.; Pletnev, V. Z.; Hazelwood, K. L.; Roche, P. M.; et al. Far-Red Fluorescent Tags for Protein Imaging in Living Tissues. *Biochem. J.* **2009**. https://doi.org/10.1042/BJ20081949.

(61)   Shaner, N. C.; Campbell, R. E.; Steinbach, P. A.; Giepmans, B. N. G.; Palmer, A. E.; Tsien, R. Y. Improved Monomeric Red, Orange and Yellow Fluorescent Proteins Derived from Discosoma Sp. Red Fluorescent Protein. *Nat. Biotechnol.* **2004**. https://doi.org/10.1038/nbt1037.

(62)   Gurskaya, N. G.; Fradkov, A. F.; Terskikh, A.; Matz, M. V.; Labas, Y. A.; Martynov, V. I.; Yanushevich, Y. G.; Lukyanov, K. A.; Lukyanov, S. A. GFP-like Chromoproteins as a Source of Far-Red Fluorescent Proteins. *FEBS Lett.* **2001**. https://doi.org/10.1016/S0014-5793(01)02930-1.

(63)   Cotlet, M.; Hofkens, J.; Maus, M.; Gensch, T.; Van der Auweraer, M.; Michiels, J.; Dirix, G.; Van Guyse, M.; Vanderleyden, J.; Visser, A. J. W. G.; et al. Excited-State Dynamics in the Enhanced Green Fluorescent Protein Mutant Probed by Picosecond Time-Resolved Single Photon Counting Spectroscopy. *J. Phys. Chem. B* **2002**. https://doi.org/10.1021/jp003813i.

(64)   Miyawaki, A. Fluorescent Proteins in a New Light. *Nat. Biotechnol.* **2004**, *22* (11), 1374–1376. https://doi.org/10.1038/nbt1104-1374.

(65)   Wilmann, P. G.; Petersen, J.; Pettikiriarachchi, A.; Buckle, A. M.; Smith, S. C.; Olsen, S.; Perugini, M. A.; Devenish, R. J.; Prescott, M.; Rossjohn, J. The 2.1A Crystal Structure of the Far-Red Fluorescent Protein HcRed: Inherent Conformational Flexibility of the Chromophore. *J. Mol. Biol.* **2005**, *349* (1), 223–237. https://doi.org/10.1016/j.jmb.2005.03.020.

(66)   Mudalige, K.; Habuchi, S.; Goodwin, P. M.; Pai, R. K.; De Schryver, F.; Cotlet, M. Photophysics of the Red Chromophore of HcRed: Evidence for Cis-Trans Isomerization and Protonation-State Changes. *J. Phys. Chem. B* **2010**, *114* (13), 4678–4685. https://doi.org/10.1021/jp9102146.

(67)   Visser, N. V; Hink, M. A.; Borst, J. W.; van der Krogt, G. N. M.; Visser, A. J. W. G. Circular Dichroism Spectroscopy of Fluorescent Proteins. *FEBS Lett.* **2002**, *521* (1–3), 31–35. https://doi.org/10.1016/s0014-5793(02)02808-9.

(68)   Stiel, A. C.; Andresen, M.; Bock, H.; Hilbert, M.; Schilde, J.; Schönle, A.; Eggeling, C.; Egner, A.; Hell, S. W.; Jakobs, S. Generation of Monomeric Reversibly Switchable Red Fluorescent Proteins for Far-Field Fluorescence Nanoscopy. *Biophys. J.* **2008**, *95* (6), 2989–2997. https://doi.org/10.1529/biophysj.108.130146.

(69)   Shcherbakova, D. M.; Verkhusha, V. V. Near-Infrared Fluorescent Proteins for Multicolor in Vivo Imaging. *Nat. Methods* **2013**, *10* (8), 751–754. https://doi.org/10.1038/nmeth.2521.

(70)   Rodriguez, E. A.; Tran, G. N.; Gross, L. A.; Crisp, J. L.; Shu, X.; Lin, J. Y.; Tsien, R.

Y. A Far-Red Fluorescent Protein Evolved from a Cyanobacterial Phycobiliprotein. *Nat. Methods* **2016**, *13* (9), 763–769. https://doi.org/10.1038/nmeth.3935.

(71)   Braslavsky, S. E.; Ellul, R. M.; Weiss, R. G.; Al-Ekabi, H.; Schaffner, K.; Silvia E. Braslavsky, Raymongd M. Ellul, Richard G. Weiss, Hussain Al-Ekabi, K. S. Photoprocesses in Biliverdin Dimethyl Ester in Ethanol Studied by Laser-Induced Optoacoustic Spectroscopy (Lioas). *Tetrahedron* **1983**, *39* (11), 1909–1913. https://doi.org/https://doi.org/10.1016/S0040-4020(01)88704-4.

(72)   Braslavsky, S. E.; Holzwarth, A. R.; Schaffner, K. Solution Conformations, Photophysics, and Photochemistry of Bile Pigments; Bilirubin and Biliverdin, Dimethyl Esters and Related Linear Tetrapyrroles. *Angewandte Chemie International Edition in English*. 1983. https://doi.org/10.1002/anie.198306561.

(73)   Choi, S. soo S.; Mandelis, A.; Guo, X.; Lashkari, B.; Kellnberger, S.; Ntziachristos, V. Wavelength-Modulated Differential Photoacoustic Spectroscopy (WM-DPAS) for Noninvasive Early Cancer Detection and Tissue Hypoxia Monitoring. *J. Biophotonics* **2016**, *9* (4), 388–395. https://doi.org/10.1002/jbio.201500131.

(74)   Beard, P. Biomedical Photoacoustic Imaging. *Interface Focus* **2011**, *1* (4), 602–631. https://doi.org/10.1098/rsfs.2011.0028.

(75)   Cox, B.; Laufer, J. G.; Arridge, S. R.; Beard, P. C. Quantitative Spectroscopic Photoacoustic Imaging: A Review. *J. Biomed. Opt.* **2012**, *17* (6), 061202. https://doi.org/10.1117/1.JBO.17.6.061202.

(76)   Tzoumas, S.; Nunes, A.; Olefir, I.; Stangl, S.; Symvoulidis, P.; Glasl, S.; Bayer, C.; Multhoff, G.; Ntziachristos, V. Eigenspectra Optoacoustic Tomography Achieves Quantitative Blood Oxygenation Imaging Deep in Tissues. *Nat. Commun.* **2016**, *7* (May), 12121. https://doi.org/10.1038/ncomms12121.

Figure 1 (a) Photophysics of OA signal generation. Jablonski energy diagram. Plain lines represent radiative transitions, dashed lines represent nonradiative processes. Abs: absorbance S_0: ground state, S_1: singlet excited state, T_1: primary triplet excited state, VR: vibrational relaxation ISC: intersystem crossing, IC: internal conversion, Pho: phosphorescence, Fluo Fluorescence. **(b)** Setup diagram (explanations and abbreviations can be found in the text). **(c)** Improvement in optoacoustic spectrum of $NiCl_2$ brought by each of the correction steps. The coefficient of determination R^2 and standard deviation (Std) between measurements are shown to indicate the improvement by each of steps. 'Max (Raw)' shows a spectrum that can be obtained using simply the maximum values in sample optoacoustic signals. 'Filtered' shows the spectrum can be obtained using filtering techniques on signals. 'Hilbert' shows the spectrum formed by maximum values of the signal envelops using Hilbert transform. '- Dark' shows the spectrum that is without DC bias by subtracting values measured in dark (no laser). 'Pulse Fl.' shows the spectrum after pulse-to-pulse fluctuation correction enabled by our in-line reference. '- Ink Spec.' shows the spectrum after removing the ink spectrum which is induced when performing fluctuation correction. '- Coloring' shows the spectrum after the correction for the light fluence coloring effect induced by the reference Ink and US-couplant water. 'Corrected' shows the spectrum that can be obtained after performing all the afore-mentioned steps (further explanations can be found in the text). **(d)** Raw optoacoustic spectrum of $NiCl_2$ at different concentrations and **(e)** after correction. **(f)** The coefficients of determination of linearity between absorbance and optoacoustic spectrum of $NiCl_2$ at decreasing concentrations. **(g)** OA and absorbance spectrum for potassium permanganate at 2 nm steps. **(h)** Raw and corrected optoacoustic spectrum of BBN. **(i)** Linear relation between absorbance and OA signal for BBN.

Figure 2 Normalized optoacoustic signal (OAS), absorption, emission and excitation spectra for **(a)** Methylene blue, **(c)** Rhodamine 800, **(e)** Alexa Fluor 750 and **(g)** IRDye 800CW. **(b,d,f,h)** The linear relationship of OAS and absorbance are given for the main peaks and shoulders, respectively.

Figure 3 (a and b) Absorption and optoacoustic spectrum of ICG at increasing concentration. **(c)** Linear relation between absorbance and optoacoustic signal for ICG. **(d)** Difference between the dimer/monomer ratio from absorbance and OA at increasing ICG concentration.

Figure 4 Normalized optoacoustic signal (OAs), absorption, emission and excitation spectra and the linear relationship of optoacoustic signal and absorbance for **(a, b)** sfGFP, **(c, d)** mKate2, **(e, f)** mCherry, **(g, h)** HcRed, **(i, j)** tdTomato, **(k, l)** smURFP and (m, n) IRFP720. Representation similar to figure 2

Figure 5 Photoswitching, -conversion or bleaching effects in mCherry, tdTomato and HcRed. Shown are the OA spectra measured from 900 to 420 nm (reverse, red) or 420 to 900 nm (forward, blue) along with the absorption spectra in the non illuminated state (a,d and g). mCherry Absorption spectra before and after spectral measurements (c) and after multiple pulses of 550/480nm and 590/565nm light for tdTomato and HcRed, respectively (f and g). The temporal development of the OA signals at the two main peak (b, e and h)