1 **Extreme Genomic Makeover: Evolutionary History of Maternally-transmitted**
2 **Clam Symbionts**
3
4 Short title:
5 Evolution of maternally-transmitted symbiont genomes
6

7 M Perez[1], C Breusing[2], B Angers[1], YJ Won[3], and CR Young[4]
8
9 [1] Department of Biological Sciences, Université de Montréal, Montreal, Canada
10 [2] Graduate School of Oceanography, University of Rhode Island, Narragansett, USA
11 [3] **Division of EcoScience, Ewha Womans University, Seoul, South Korea**
12 [4] National Oceanography Centre, Southampton, UK

13
14
15 Classification:
16 Evolution, Genetics, Microbiology, Selection, Recombination, Symbiosis, Genomics,
17 Vesicomyidae
18
19
20
21 **Abstract**
22

Given their recent switch to a vertically-transmitted intracellular lifestyle, the
chemosynthetic bacteria associated with deep-sea vesicomyid clams are an excellent
model system to study the processes underlying reductive genome evolution. In this
study, we provide the first estimates of the relative contributions of drift,
recombination and selection in shaping the ongoing reductive genome evolution in
these symbionts. To do so, we compared the genomes of endosymbionts associated
with 11 vesicomyid clam species to that of closely related free-living bacteria and
their respective hosts' mitochondria. Our investigation confirmed that neutral
evolutionary processes were the dominant driver of reductive genome evolution in
this group and highlighted the important role of horizontal gene transfer in mitigating
genome erosion. Finally, a genome-wide screen for episodic positive selection across
the symbiont phylogeny revealed the pervasive role of selective processes in
maintaining symbiont functional integrity.

Introduction

The evolution of biological complexity includes many examples of symbiotic associations. For example, the early evolution of the eukaryotic cell involved multiple endosymbiotic events leading to mitochondria and plastids [1,2]. More recent examples include associations of metazoans with intracellular bacteria[3–6], including the well-studied associations of insects and *Buchnera* proteobacterial symbionts [7]. These associations have profound consequences for both host and symbiont, ranging from alterations of sex-ratio in insect hosts to providing nutrients that are otherwise unavailable in the host's habitat. Some intracellular symbionts are transmitted from parent to offspring of hosts through the germline (i.e. vertical transmission), while others are acquired from the environment every generation [6]. The mode of transmission strongly affects the evolution of the microbial partner in these symbioses, as the genomes of vertically transmitted symbionts all seem to follow the same process of reductive genome evolution (RGE) regardless of their phylogenetic origin, host, or habitat. Compared to their free-living counterparts, the genomes of host-restricted symbionts are smaller, contain fewer genes, and are enriched in AT [8,9]. A prime example is the genomes of cellular organelles such as mitochondria and plastids which are extremely streamlined compared to their bacterial cousins [10]. Symbiont genome evolution is thought to follow two main stages [11]. Following host restriction, symbionts undergo rapid genome erosion as they lose non-essential genes through pseudogenization and deletions [12,13]. Then, symbionts enter a "stabilizing phase". At this point, their genomes are streamlined, redundant genes and functions are lost [14], and the effective rate of deletion diminishes [15]. This process might be largely neutral due to the reduced effective population size of host-restricted taxa.

The pea aphid/*Buchnera* symbiosis and several other insect/bacteria models support the neutral hypothesis. Captured symbionts experience successive bottleneck events during their transmission that reduce their effective population size and increase genetic clonality. As a consequence, genetic drift increases relative to selection in these taxa [16–18]. Under these circumstances, elevated mutation load (i.e. the Muller's ratchet [19]) and genetic erosion might lead to the functional death of the symbiont lineage [17,20–22] unless compensating mechanisms such as gene transfer to the host nucleus or compensatory mutations alleviate the genetic load. Likewise, deep-sea taxa exhibit evidence of nearly neutral processes affecting evolutionary rates due to reduced population sizes in vertically transmitted symbionts [23]. Other metazoan/microbial symbioses highlight the importance of selection in shaping reductive genome evolution. For instance, symbiont traits that are beneficial for the host are likely to experience increased selective pressures, while selection may be relaxed on genes that are functionally redundant [8]. Red Queen dynamics are expected to occur in obligate symbioses to maintain the host-symbiont specificity and the functioning of cyto-nuclear interactions through speciation [20]. Unfortunately, the role of positive selection has often been ignored in studies of symbiont genome evolution and broad screens for positive selection have almost never been performed.

The intracellular sulfur-oxidizing bacteria associated with deep-sea vesicomyid clams (Bivalvia: Vesicomyidae: Pliocardiinae) represent an ideal model to address the neutral and selective processes driving reductive genome evolution. The symbionts are found within the epithelial cells of their host's gills and provide them with chemosynthetically derived food. They are vertically transmitted to the next generation through the eggs [24,25] and generally show co-speciation with their hosts

[26,27]. It is assumed that symbiont capture in these animals was a single event that,
based on fossil and molecular information, happened before their radiation about 45
Mya [28], an acquisition that is much more recent than that of other well-studied models
such as the aphid/*Buchnera* (~ 200 Mya [29]) and nematode/*Wolbachia* (~100Mya [30])
symbioses. Today, the hosts represent the most diverse group of deep-sea bivalves [31],
with 173 described species present in a variety of reducing habitats worldwide from
hydrocarbon seeps on continental margins to hydrothermal vents on mid-ocean ridges
[32–34]. A comparative study of the first two sequenced vesicomyid symbiont genomes
[35,36] indicated that they possessed intermediate genome sizes and level of AT
enrichment compared to other host-restricted symbionts [11]. The symbionts of deep-sea
vesicomyid clams group into two divergent clades: Clade I (associated with hosts of
the *gigas* group), and Clade II (associated with all other lineages of vesicomyid hosts)
[37]. The genomic characteristics of Clade I symbionts indicate that this group is in an
advanced state of reductive genome evolution compared to Clade II. However, in
contrast to the well-studied pea aphid/*Buchnera* association, which has been in a state
of stasis for 50 Myrs [38], the evolutionary processes responsible for remodeling the
genomes of vertically transmitted symbionts appear to be still operating in the
vesicomyid clam symbiosis. Conspicuous bottlenecks during transmission [25] and loss
of DNA repair genes in several lineages [37] suggest that neutral processes and
mutational pressures are driving RGE in vesicomyid symbionts, although this
hypothesis has not been formally tested.
108
In this study, we aim to assess the relative contribution of neutral and selective
processes to genome evolution in the maternally transmitted symbionts of deep-sea
vesicomyid clams. Specifically, we test the hypotheses that genetic drift is the main
driver of RGE in these symbionts and that diversifying selection has shaped their
genome to maintain host-symbiont epistasis throughout the evolutionary history of the
symbiosis. To do so, we applied comparative methods to the symbiont genomes of 11
vesicomyid deep-sea clam taxa representative of the diversity of Clade I and Clade II,
the mitochondrial genomes of their respective hosts, and two of their close free-living
relatives: the environmentally acquired gill symbiont of the hydrothermal vent mussel
Bathymodiolus thermophilus and the free-living bacteria of the SUP05 group, which
are marine chemoautotrophic Gammaproteobacteria found in hypoxic waters [39,40].
120
121

Results

Host mitochondrial and symbiont phylogenies

Host mitochondrial genomes from the lineages examined in this study possess
identical gene orders and contents as previously published mitochondrial genomes
[41,42]. The phylogeny constructed with mitochondrial genome data (Figure 1A) is
congruent with the known host phylogenetic relationships based on multilocus
sequence data and the *COI* phylogeny [31]. Structural variation is, however, present. We
observe the previously described noncoding structural variation, hypothesized to be
the control region, between the *tRNA^Trp* or *tRNA^His_2* and *ND6* loci [41–43] but we were
unable to resolve this region with the current sequence data. We also found the *COX2*
gene varies in length among taxa (range: 1005-1452bp). All protein-coding genes in
the mitochondrial genomes were screened for selection using the adaptive branch-site
random effects likelihood method. Interestingly, the *COX2* gene exhibited evidence
for episodic diversifying selection on multiple branches of the phylogeny.

Genome size and GC content for the 11 symbiont assemblies in our study varied from
1.02Mb to 1.59 Mb and 31% to 37% GC, respectively (Table 1). The number CDS
ranged from 939 in *Ca.* V okutanii to 2210 in *Ca.* R. phaseoliformis. Following initial
nomenclature, the symbiont lineages are referred to by the previously erected genera
for this group, *Candidatus* Vesicomyosocius for Clade I, and *Candidatus* Ruthia for
Clade II symbionts, followed by host species names [35,36,44]. This classification at the
genus level is coherent with both the phylogenetic definition based on 16S identity
(inter-genus identity < 95% [45]) and functional definition based on criteria of genetic
isolation [46] (see Symbiont genome structure and recombination)

Examination of the mitochondrial and symbiont phylogenies (Figure 1) shows good
concordance for all lineages except one. The symbiont lineages of *Ca.* V. diagonalis
and *Ca.* V. extenta are nearly identical whereas their respective host mitochondrial
lineages are divergent. The donor lineage in this recent symbiont replacement appears
to be *A. diagonalis*. It is noteworthy that these clams were both collected from sites in
Monterey Canyon. Pairwise comparison of mitochondrial and symbiont genome-wide
synonymous divergence indicates faster evolutionary rates in the mitochondria
compared to the symbionts in almost every holobiont pair (Figure 2). Within the
symbionts, we detect signatures of elevated substitution rates on the branch leading to
Clade I: the symbiont pairs across the Clade I- Clade II bipartition have significantly
higher divergence than the others even when controlled for host divergence ($1 < dS_{mito}$
<2).

Symbiont genome structure and recombination

Free living bacteria associated with *B. thermophilus* and *Ca.* T. autotrophicus shared
about 1 Mbp of their genomes with the clam symbionts. Permutation analysis of
locally collinear blocks (i.e. long fragments of aligned genomes) with GRIMM
(http://grimm.ucsd.edu/cgi-bin/grimm.cgi) showed that at least 18 inversion events
occurred between the genome of the *B. thermophilus* symbiont and that of the *Ca.* R.
magnifica reference. Fewer rearrangements (3 inversions) were observed between
SUP05 and *Ca.* R. magnifica.

Genome structure among the clam symbionts was also variable (Figure 1B). The
previously reported *Ca.* V. okutanii genome [36] possesses one inversion compared to
that of *Ca.* R. magnifica [35] but that of *Ca.* V. okutanii's closest relative, *Ca.* V. soyoae,
does not. The genomes of *Ca.* R. pacifica and *Ca.* R. rectimargo share a single
inversion distinct to that of *Ca.* V. okutanii. Two other inversions were found in the
Ca. V. gigas genome. Finally, read-mapping to the consensus assemblies for *Ca.* R.
phaseoliformis and *Ca.* R. southwardae suggested the presence of intra-host structural
variation in these symbionts.
177
Applying Bayesian concordance analysis to all core protein-coding genes, we detect a
large amount of recombination among symbiont lineages, though recombination is not
randomly distributed. We observe no recombination between members of Clade I and
II, but recombination is occurring within these genera (Figure 1B). Strikingly, much
less topological concordance was found in Clade II – more than 40 different
topologies were necessary to fully represent the diversity of conflicting phylogenetic
signals – compared to that of Clade I whose phylogeny was fully represented by 5
different trees. Within Clade I, conflict originates from the uncertainty of the position
of *Ca.* V. gigas. Only 50% of the genes support its position in the phylogenetic tree
issued from the concatenated core genome alignment (Figure 1B). Other well
supported positions for this species are at the base of the clade (supported by 27% of
genes) and closer to the group composed of *Ca.* V. soyoae and *Ca.* V. okutanii
(supported by 20% of genes). Within Clade II, only the grouping of the sister species
Ca. R. rectimargo and *Ca.* R. pacifica is supported by the topologies of all genes
while the positions of other species have low support.
193

Gene conservation across symbionts and free-living bacteria

Genes of free-living and horizontally-transmitted bacteria missing in vesicomyid
symbionts
The genomes of the free-living bacteria contained many large (> 5kb) contiguous
sections that were not found in the symbionts. These genomic islands were mostly
composed of unannotated genes and mobile elements (transposases, integrases,
prophage genes) (Table S1). We found more selfish genetic elements in the genome
of the *Bathymodiolus* symbiont than in that of SUP05. The genomic islands found in
the two genomes also encoded several gene clusters of particular functional interest
described below.
204
Unsurprisingly for a bacterium living in a metal-rich hydrothermal environment, the *B.*
thermophilus symbiont genome possesses genes for resistance against heavy-metal
toxicity such as a multi-copper oxidase (*mmcO*), a copper ion exporting ATPase
(*copB*), cobalt-zinc-cadmium resistance proteins (*czcD* and c*zcCBA*), and a chromate
transport protein (*chrA*). The genomic islands of the mussel symbiont also carried full
operons for three different defense systems; a type I restriction and modification
system (*hsdRMS*), a CRISPR-Cas type II system (*cas9*, *cas1*, *cas2*, *cas4*), and a type
II toxin-antitoxin system (*vapCB*). Finally, this genome possesses a 23kb hydrogenase
operon that has 83% and 82% identity to that of the symbionts of *Bathymodiolus*
septemdierum [47] and *B. puteoserpentis* [48], respectively. The representative SUP05
genome contained a 21kb motility locus, comprising a type IV pilus biogenesis
operon (*pilA*, *pilB*, *pilC*, *pilT*, *pilQ*, *pilY1*), and a toxin-antitoxin locus (*higAB*), that
was not found in the other genomes. Furthermore, this genome possessed two

additional smaller genomic islands (6kb and 15 kb) encoding a nitric oxide reductase
(*norCBQD*), and a periplasmic nitrate reductases (*napAB*), respectively, which
clustered with sulfur covalently binding protein genes (*soxYZ*).
221
Gene content in vesicomyid symbionts
The symbionts of Clade I and Clade II possessed essentially a subset of the genes
found in the free-living lineages. Indeed, sequence-based comparisons of free-living
lineages to the symbionts revealed that many genes present only within the symbiont
lineages are hypothetical genes with unknown function resulting from the
degeneration of ancestral genes, as indicated by premature stop codons, frameshifts,
and loss of neighboring genes (Table S1). These pseudogenes were more prevalent in
the genomes of Clade I than Clade II symbionts. In many instances, homologous
regions within the Clade I symbiont genomes were instead characterized by large
deletions. In general, patterns of gene decay were more variable within Clade II than
Clade I. Genes were overall more conserved within *Ca.* R. southwardae, *Ca.* R.
phaseoliformis and *Ca.* R. pliocardia than in other lineages. Among the *Ca.* Ruthia
symbionts, gene degeneration was most pronounced in *Ca.* R. magnifica, which
possessed a conservation pattern closer to that of the Clade I lineages (Figure S1B).

Genome-wide pattern of relaxed selection

Codon usage bias was reduced in the symbiont lineages compared to their free-living
relatives. Furthermore, symbionts in Clade I showed reduced bias and variance
compared to Clade II (Figure 3A). The CDC values of core protein-coding genes were
significantly correlated between lineage pairs both at the clade and species level
(Pearson's test p-value <0.001; Figure 3B, Table S2), suggesting that the reduction in
codon usage bias in the vertically transmitted symbionts result from a genome-wide
reduction of the efficacy of purifying selection.
RELAX analysis revealed intensified selection in the vesicomyid symbionts
compared to free-living bacteria for less than 5% of the core orthologous genes, while
relaxed selection was detected in more than half of the core gene set (Figure 3C,
Table S3). The magnitude of relaxation (k<1) was in the range of that observed in
insect endosymbionts [49] but was not correlated to codon bias. Genes exhibiting
intensified and relaxed selection represented a multitude of metabolic functions, but
genes under relaxed selection were enriched in the protein metabolism, nucleoside
and nucleotides, and DNA metabolism categories while genes under intensifying
selection were more likely to be associated with respiration, cell wall and capsule, and
sulfur metabolism. However, we did not find increased relaxation in the symbionts of
Clade I compared to Clade II. Indeed, fewer genes exhibited significant change in
selection pressure (intensified or relaxed) between these groups than between
symbionts and free-living bacteria, and about the same proportion of genes under
relaxed and intensified selection was found in both clades.
258

Genome-wide screen for positive selection

The symbiont genes that passed the inclusion criteria to be screened for selection (see
methods) included 652 loci. The application of the adaptive BS-REL method yielded
223 genes with significant evidence for episodic diversifying selection along branches
in the phylogeny. Selection is distributed throughout the evolutionary history of the
group (Figure S1A, and Table S4) with most selection occurring on the branches
discriminating free-living bacteria, Clade I, and Clade II (branches a, b, and c in

Figure 4), as well as within the *B. thermophilus* symbiont and SUP05 lineage (43 and
37 genes, respectively). Eighty-five percent of the loci that exhibited unequivocal
evidence of selection was assigned to SEED categories (Figure 4, Table S5). Within
each clade and along each of the main branches, these selected loci were not equally
represented amongst cellular functions of the core genome (hypergeometric tests p-
values < 0.001). Genes in overrepresented functional categories are presented in Table
2. The complete list of selected genes is available in Table S4.
273
Selection within free-living bacteria
Amongst the free-living lineages, a larger than expected number of genes associated
with protein metabolism, respiration, and sulfur metabolism were under selection
(Fisher tests p-value <0.05). These included genes involved in ribosome assembly, t-
RNA biogenesis, protein folding, oxidative phosphorylation, sulfur oxidation, and
dissimilatory sulfate reduction. On the bipartition between the free-living and
symbiont groups, additional genes associated with protein metabolism were positively
selected, including ribosomal protein genes, and the t-RNA ligase genes.
282
Selection within symbionts
Many genes coding for chaperones, ribosomal proteins, and t-RNA ligases were under
selection within the symbiont phylogeny. In addition, we found evidence for selection
in metabolic genes that are central to the chemosynthetic role of these symbionts.
Several genes involved in sulphur metabolism (i.e. *dsrA, dsrP, soxB, cobB-cbiA/dsrN*)
and electron donating/accepting reactions were under selection. Two genes involved
in ammonia assimilation (*gltB*, and *glnD*) also exhibited evidence of selection within
both symbiont clades. Within Clade II and along the branch partitioning this group,
there was an over-representation of selected genes involved in *de novo* purine and
pyrimidine biosynthesis, carbon fixation, and DNA recombination and repair.
Selection within Clade I favored additional genes broadly associated with DNA
metabolism. Notably, 60 genes showed evidence for positive selection in multiple
branches of the phylogeny, including 44 genes within the symbiont phylogeny. These
genes were mostly associated with protein metabolism.
297
Discussion

Reductive genome evolution is still ongoing in the clam symbionts and is driven by
neutral processes

Comparative analyses of the first two reference genomes of vesicomyid clam
endosymbionts revealed variation in genome structure, genome characteristics, and
genome composition between distantly-related symbiont species [11] suggesting that
RGE might still be ongoing in this group. Our results confirm these early findings and
reveal additional genomic variation among the deeply diverging lineages. These
findings expand the ranges of genome size, genome content and GC% considerably.
307
As in other models of recently acquired bacteria [22,50], gene content differed greatly
between vesicomyid symbiont genomes indicating that the different lineages are
independently losing genes. The presence of structural variation and putative
pseudogenes (Figure S2) within the vesicomyid symbiont genomes suggest that these
symbionts have not yet reached a stable streamlined state as those of the *Buchnera* or
Paulinella symbionts [15,38]. Comparing the clam symbionts to their free-living relatives
revealed reduced GC%, a reduction in codon usage bias, pseudogenization, and

evidence for reduced purifying selection in the vast majority of genes. Taken together,
these observations support the nearly neutral theory of RGE, driven by a reduction of
effective population size in these taxa.
318
Finally, in agreement with the findings of Stewart et al. [27,51], Decker et al. [52], and
Ozawa et al. [53], we detected no recombination between Clade I and II symbionts even
though some of the host taxa co-occur [54–56]. These findings imply that there is enough
molecular and ecological divergence between the two clades for clonal interference
and/or strong host-symbiont epistatic interactions to constrain symbiont exchange [20,52].
Thus, our results support the nomenclature initially put forward by Newton *et al.* [35]
and Kuwahara *et al.* [36] classifying the symbionts from Clade I and II into two distinct
bacterial genera, *Ca.* Vesicomyosocius and *Ca.* Ruthia. For clarity, we will keep
referring to these two genera as Clade I and Clade II in the rest of the discussion.

Reductive genome evolution is exacerbated in non-recombining symbionts

Clade I symbionts are in a more advanced state of RGE than the others. Indeed,
compared to Clade II, their genomes are smaller and lower in GC%, possess fewer
genes and pseudogenes, and exhibit less codon usage bias. The genomes of Clade I
symbionts are also more homogeneous. Patterns of gene conservation suggest that
much of the loss in this group happened after its speciation but before its radiation, a
period of roughly 20Mys [26,31]. Together with increased substitution rate on its
diverging branch these results show that the ancestral Clade I lineage experienced an
episodic acceleration of reductive genome evolution. It is likely that the increased
level of genome reduction in Clade I results from a reduction of homologous
recombination in the ancestor of the group exacerbating Muller's ratchet [57]. Drift-
driven loss of recombination machinery may have strongly reduced the rate of genetic
exchange among the symbionts in this genus. Indeed, essential genes of the RecF and
RecBCD pathways for homologous recombination appear to be lost in all of the Clade
I symbionts [37] and while horizontal transfer of genetic material is widespread among
symbionts within Clade II it is almost absent in Clade I.
344
Strong linkage disequilibrium forces whole genomes to sweep in populations that lack
genetic exchange capabilities. Hence, the loss of homologous recombination genes
should favor symbiont replacement in cases where the divergence between "native"
and foreign symbionts is low (i.e. when the foreign symbionts are not too easily
outcompeted by those that have co-evolved with the host). In fact, we find multiple
examples of symbiont replacement among symbionts of Clade I. For instance,
individual clams of the species *P. extenta* have acquired the symbionts of the
sympatric species *A. diagonalis*. Likewise, Breusing et al. [56] found a population of *A.*
gigas carrying the symbionts of the host species *P. soyoae*. Symbiont replacement
occurs in several vertically transmitted symbioses [58–61] and is speculated to constitute
a mechanism for escaping the evolutionary rabbit hole caused by Muller's ratchet
[20,58,62]. The present data support this notion, and future population genomic studies
could determine the prevalence of symbiont replacement and relative rates of
recombination in these taxa on more recent time scales.
359
Despite the lack of recombining machinery in Clade I, one lineage in this genus, *Ca.*
V. gigas, showed evidence for recombination. It is puzzling how recombination might
be occurring in this species. Breusing et al. [56] recently found evidence of
unidirectional introgression from *P. soyoae* into *A. gigas*. This mechanism might

enable *A. gigas* symbionts to come into contact with other symbionts. Perhaps the
recombination in this species is enabled via host-encoded proteins [63]. Transfer of
symbiont genes to the host nuclear genome is possible and should be investigated in
future studies. Indeed, evidence for such transfer was recently found by Ip *et al.* [44]
who identified *Bathymodiolus* symbiont gene homologs in the genome of the *A.*
marissinica.

Putative ecological and evolutionary consequences of RGE

The Muller's ratchet has been hypothesized to lead to a progressive loss of fitness in
host restricted symbionts [20]. Sympatric populations of symbionts from Clade I and II
represent an excellent model to test this hypothesis because of their contrasting
reductive stages. For instance, comparisons of the sulfide physiology of the host
species *P. soyoae* and *C. pacifica*, which occupy different micro-niches in the same
habitat, reveal that *P. soyoae* individuals have lower sulfide oxidation capacities than
those of *C. pacifica* [55]. This could be the consequence of a less efficient sulfide
metabolism in *Ca.* V. soyoae resulting from a more advanced state reductive genome
evolution in this species compared to *Ca.* R. pacifica. If RGE in the symbionts can
restrict their host's ecological range, contrasting degrees of RGE may put constraints
on the potential for genetic exchange across different holobiont species and even
promote speciation [20]. Future observational and experimental studies could help
define the evolutionary constraints imposed by both host and symbiont physiology
and clarify the role of reductive genome evolution in niche partitioning and speciation.
385

Selective processes in the evolutionary history of the symbionts

Contrasting patterns of gene conservation between the symbionts and their free-living
relatives are caused by a shift in selective regime in the host-associated bacteria.
Genes enabling bacteria to face the challenges of a free-living environment, such as
detoxification, anti-viral defense and inter-species competition, were not conserved in
the vesicomyid clam symbionts. Furthermore, different patterns of pseudogenization
in Clade I and Clade II likely translate to different physiological adaptations at the
level of the holobiont. For example, Breusing et al. [in review] found that the two
vesicomyid symbiont clades show enzymatic differences related to sulfide oxidation
and nitrate reduction and have contrasting dependencies on nickel and vitamin B12 in
accordance with adaptations to different ecological niches. In addition, episodes of
diversifying selection on genes associated with respiration, ammonia assimilation, and
chemosynthesis might reflect the constraints imposed by the diverse selective
pressures of host physiology throughout their radiation and niche expansion.
400

Selective constrains are expected to affect genes involved in host-symbiont
interactions. Interspecific communication between eukaryotes and microbes generally
involve molecules with distinct motifs produced by the symbiont (e.g., Nod factors,
lipopolysaccharides, or peptidoglycans) that are sensed by special receptor in the host
[64,65]. These molecular pathways must experience reciprocal adaptations to persist
through speciation and niche expansion. Diversifying selection acting on genes
involved in the mediation of host-symbiont interactions such as lipopolysaccharides
and peptidoglycans was observed in divergent clades of *Wolbachia* [66] and many
facultative endosymbionts [67]. In a recent study, Chong et al. [68] performed a genome-
wide screen for selection in the *Buchnera* symbionts from the aphid subfamily
Aphidinae. Of the 371 protein-coding genes tested, the authors detected 29 positively

selected genes representing a variety of metabolic functions including two outer
membrane porins (OmpF and OmpA), which are assumed to be important for host
interaction.
415
Surprisingly, in the clam symbionts, we did not detect selection on proteins associated
with host-symbiont interactions but found instead a pervasive pattern of diversifying
selection that affected many loci related to housekeeping functions such as DNA and
RNA metabolism, transcription and translation. Many ribosomal proteins and
chaperones showed evidence for episodic positive selection repeatedly throughout the
symbiont phylogeny. These results could indicate that the accumulation of slightly
deleterious mutations in the symbiont genomes initiates a selective pressure for
compensatory mutations [69,70]. Evidence for such mutations exist in several organelles
and symbiont models [70–73]. For instance, in insect endosymbionts, positively selected
loci of the chaperonin GroEL are suspected to permit better protein binding and allow
proper protein folding despite mutations affecting their conformation [71]. Alternatively,
these signatures of selection might be in response to other generalized selection
pressures such as differences in host habitat (e.g., depth). However, the host
mitochondria do not overall seem to be similarly affected making this alternative less
likely. Regardless, the pervasive nature of episodic diversifying selection at the level
of amino acids in the symbiont genomes suggests that increased drift due to effective
size reduction is not the sole driver of molecular evolution in these taxa.
433

Conclusion

The vertically transmitted symbionts of deep-sea vesicomyid clams are an ideal model
to study the processes of reductive genome evolution, as they constitute a highly
diverse group of host-restricted bacteria with varying degrees of genomic reduction.
We show that both neutral and selective processes have played a role in the
evolutionary history of these symbiont and that factors affecting their clonality have
strongly influenced the rate of genome evolution. While the vesicomyid clams have
yet to be successfully bred in aquaria, significant progress has been made towards
their cultivation [74]. Examination of the symbionts at the population-level, both within
and across individual hosts, will help to decipher the contributions of host physiology,
genetic drift, symbiont fitness, cytonuclear incompatibilities, and horizontal gene
transfer to their evolution. Additionally, experimental studies on host-symbiont
interactions and holobiont metabolism will shed further light onto the role of these
symbionts in the ecological partitioning of their hosts.
448

Acknowledgements
The Monterey Bay Aquarium Research Institute kindly provided samples from the
Vrijenhoek collection for this study. We thank the ships' crews and submersible pilots
involved in the collections for this study, without whose efforts this work would not
have been possible. N. Pratt and A. Baylay contributed to sequencing at the National
Oceanography Centre Genomics Facility. This work was supported by NERC
National Capability funding. MP acknowledges the support of the National Science
and Engineering Research Counsil (NSERC) of Canada's Alexander Graham Bell
graduate scholarship and Michael Smith Foreign Study Supplements. CB's
contribution was supported through a postdoctoral fellowship of the German Research
Foundation (BR 5488/1-1) and a grant from the United States National Science
Foundation awarded to CB's mentor Roxanne Beinart at the University of Rhode

Island (OCE-1736932). BA acknowledges support from NSERC research grant
#238600. The genome of *Bathymodiolus thermophilus* was sequenced as part of a
project titled 'Understanding the deep-sea biosphere on seafloor hydrothermal vents
in the Indian Ridge (No. 20170411)' funded to YJW by the Ministry of Oceans and
Fisheries, Korea.

Materials and Methods

Sample collection and sequencing

Host taxa examined in this study were chosen from the deepest diverging lineages within the Vesicomyidae that are distributed globally in the northern hemisphere (Figure S3) and are representative of the known host diversity [31]. Specimens of nine clam species were collected between 1996 and 2004 over eight research expeditions (Table 3, Figure S3). Depths of sampling locations ranged from 650–3550m. Samples were dissected aboard ship and then frozen at -80C or were frozen whole at -80C. DNA was extracted from symbiont bearing gill tissue using the DNeasy Blood & Tissue extraction kit (Qiagen, Hilden, Germany) following the manufacturer's protocol. Host species identification was initially confirmed by sequencing the host mitochondrial *COI* gene using vesicomyid-specific primers [28].

Mixed host and symbiont DNA samples were sequenced in-house on a MiSeq instrument. Genomic DNA libraries were prepared using the KAPA Hyperplus Library Preparation kit (KAPA Biosystems, Wilmington, MA, US) according to kit instructions. Read quality of genomic data was assessed using FastQC [38].

Mitochondrial and symbiont genome reconstruction and annotation

Initial symbiont and mitochondrial assemblies were constructed from the same metagenomic libraries (Table 3) using Velvet [76], manually optimizing for *k*-mer size distribution and read depth. Some assemblies were also constructed using the read mapping and assembly functions in Geneious version 10.1.3 [77].

Scaffolding and circularization of the symbiont genomes were performed by mapping, extracting and reassembling reads mapping to the extremities of contigs using Bowtie2 [78], Samtools [79] and SPAdes [80], respectively. Mitochondrial genomes were assembled de novo with MITObim [81] using as seed a set of initial contigs constructed using the read mapping and assembly functions in Geneious version 10.1.3 [77]. Mitochondrial genome annotations were produced by the GeSeq application [82] using ARWEN v1.2.3 for tRNA prediction, and manually curated with the aid of previously annotated mitochondrial genomes [41,42] in Geneious. Mitogenome assembly statistics are presented in Table S6. The symbiont genomes were annotated in RAST [83].

Structural variation and phylogenomic analyses

Host mitochondrial and symbiont genomes were aligned with Progressive Mauve [84]. Progressive Mauve and GRIMM (http://grimm.ucsd.edu/cgi-bin/grimm.cgi [85]) were used to identify large-scale structural differences among genomes. Locally collinear blocks (LCBs) longer than 100bp and found in all genomes were extracted with Mauve's stripSubsetLCBs program, aligned with Mafft [86] and concatenated into host mitochondrial and bacterial core genomes. Phylogenetic trees were produced from these core genomes using the GTR model and 100 bootstraps in PhyML-3.1 [87].

We compared host and symbiont evolutionary rates by estimating the divergence at synonymous sites for each host pair. Using the Biopython toolkit [88], we extracted the nucleic and amino acid sequences of 13 conserved mitochondrial and 718 bacterial core protein-coding genes (see below). Amino acid sequences were then aligned with Muscle [89] and reverse translated into codon alignments using the "build" function

from the Biopython codonalign package. The mitochondrial and bacterial codon-
based alignments were then each concatenated into two genome-wide alignments with
complete (no gaps, no N) lengths of 10417 bp and 662118 bp, respectively. We
assessed substitution saturation by plotting transitions and transversions against
adjusted genetic distance. Pairwise synonymous (dS) substitution rates were
computed using the Maximum-Likelihood method [90] implemented in the Biopython
codonalign package. The source code was slightly modified to accommodate for
ambiguous bases in the mitochondrial genomes.

Identification of bacterial core genes

Because of low structural differences among genomes, orthologous genes could be
inferred based on homology and position [91]. A list of positional homologs with a
minimum identity of 30% and a minimum coverage of 60% was exported from the
Mauve alignments. Additional maps with a stricter identity criterion (60% identity, 80%
coverage) were produced from the alignments of multiple subsets of symbiont
genomes. The consensus of these orthologous maps yielded 749 core genes (Figure
S2, Table S1) including 718 core protein-coding genes ranging from 138 bp to 4554
bp (average 975bp) (Table S3).
531

Bayesian concordance analyses

We used Bucky v.1.2 [92] to estimate the proportion of core protein-coding genes
supporting each topology. Putative recombination breakpoints within the 718 core
protein-coding genes previously found were identified with GARD and the KH test
[93,94]. Using a false positive discovery rate threshold of 5%, recombination was found
in 66 genes which were thus split into multiple contiguous non-recombining gene
segments at the inferred breakpoints prior to phylogenetic inference.
Bucky takes as input the posterior distribution of topologies for each gene (or gene
segment). These distributions were each obtained from 800 trees generated in
MrBayes v.3.2.7a [95] using a Gamma + I rate variation across sites. These trees
represented a well-mixed sample of the tree space after convergence of four
independent Markov Chain Monte Carlo (MCMC) chains which were each run for
2,000,000 generations after an initial 100,000 generations burn-in period. Trees were
sampled every 10,000 generations to avoid autocorrelation. Parameter optimization
for the MCMCs was performed by assessing convergence and mixing of both
continuous parameters of the model and tree topologies using the R package RWTY
v.1.0.2 [96].
In Bucky, two independent MCMC runs were carried out using the prior assumption
that all genes shared the same topology (alpha=0). MCMC runs performed 1,000,000
updates after an initial 10% burn-in period. One cold and three heated chains
(swapping frequency =10) were used to improve mixing and convergence of all of the
MCMC runs.
554

Relaxed and positive selection detection

Relaxation of the strength of selection was detected in the symbiont genomes by two
independent methods. First we use the Codon Deviation Coefficient (CDC) [97] to
quantify codon usage bias on all protein-coding genes (Table S2) because this index
does not require a priori knowledge of gene expression and is not biased by GC
content. Second, we used RELAX [49] on individual core genes. RELAX detects

change in the strength of selection between two groups by observing change in the
distribution of ω (dN/dS ratio) classes in a branch-site random effects likelihood (BS-
REL) framework between a set of test and reference branches. We compared *Ca.*
Vesicomyosocius, *Ca.* Ruthia, and both clades together to the group composed of the
free-living lineages.
To reduce false positives in phylogenetic selection tests [98], genes with significant
evidence of recombination (see Bayesian concordance analyses) were excluded from
these analyses. Episodic diversifying selection on individual lineages was identified
on the remaining non-recombining 652 protein-coding genes using the adaptive
Branch-site Random Effects Likelihood method (aBSRel [99]). The Holm-Bonferroni
correction for multiple testing was applied and threshold for detection was set to 10%
false positive discovery rate. We used the hypergeometric test function dmvhyper
from extraDistr v.1.8.11 [100] to test whether the genes under relaxed or positive
selection represented a random subsample of all core genes according to SEED
categories [83]. The Fisher test [101] was applied to find SEED categories that were over-
represented in the genes under relaxed or positive selection.
577

Data availability

Symbiont genomes and Sequence Read Archives (SRAs) are available at the National
Center for Biotechnology Information (NCBI) under the BioProject PRJNA641445.
The mitochondrial genomes were deposited in GenBank under the references
MT947381-MT947391.

Genome alignment files and Rmarkdown scripts of downstream analyses are available
at https://github.com/maepz/VesicSymb_Evolution

Authors contributions
CRY designed the study; CRY and YJW contributed to data collection; MP, CB, and
CRY performed analysis, BA contributed to data interpretation. And all authors co-
wrote the manuscript.

Competing interests
The authors declare no competing interests.

References

1. Roger, A. J., Muñoz-Gómez, S. A. & Kamikawa, R. The Origin and Diversification of Mitochondria. *Current Biology* **27**, R1177–R1192 (2017).

2. McFadden, G. I. Origin and Evolution of Plastids and Photosynthesis in Eukaryotes. *Cold Spring Harb Perspect Biol* **6**, a016105 (2014).

3. Cavanaugh, C. M. Microbial Symbiosis: Patterns of Diversity in the Marine Environment. *American Zoologist* **34**, 79–89 (1994).

4. Clavijo, J. M., Donath, A., Serôdio, J. & Christa, G. Polymorphic adaptations in metazoans to establish and maintain photosymbioses. *Biological Reviews* **93**, 2006–2020 (2018).

5. Garate, L., Sureda, J., Agell, G. & Uriz, M. J. Endosymbiotic calcifying bacteria across sponge species and oceans. *Scientific Reports* **7**, 43674 (2017).

6. Bright, M. & Bulgheresi, S. A complex journey: transmission of microbial symbionts. *Nat Rev Micro* **8**, 218–230 (2010).

7. Baumann, P. *et al.* Genetics, physiology, and evolutionary relationships of the genus Buchnera: intracellular symbionts of aphids. *Annu. Rev. Microbiol.* **49**, 55–94 (1995).

8. Wernegreen, J. J. Endosymbiont evolution: predictions from theory and surprises from genomes. *Annals of the New York Academy of Sciences* **1360**, 16–35 (2015).

9. Fisher, R. M., Henry, L. M., Cornwallis, C. K., Kiers, E. T. & West, S. A. The evolution of host-symbiont dependence. *Nature Communications* **8**, 15973 (2017).

10. Husnik, F. & Keeling, P. J. The fate of obligate endosymbionts: reduction, integration, or extinction. *Current Opinion in Genetics & Development* **58–59**, 1–8 (2019).

11. Kuwahara, H. *et al.* Reductive genome evolution in chemoautotrophic intracellular symbionts of deep-sea Calyptogena clams. *Extremophiles* **12**, 365–374 (2008).

12. Itoh, T., Martin, W. & Nei, M. Acceleration of genomic evolution caused by enhanced mutation rate in endocellular symbionts. *PNAS* **99**, 12944–12948 (2002).

13. Moran, N. A., McLaughlin, H. J. & Sorek, R. The Dynamics and Time Scale of Ongoing Genomic Erosion in Symbiotic Bacteria. *Science* **323**, 379–382 (2009).

14. Mendonça, A. G., Alves, R. J. & Pereira-Leal, J. B. Loss of Genetic Redundancy in Reductive Genome Evolution. *PLOS Computational Biology* **7**, e1001082 (2011).

15. Lhee, D. *et al.* Evolutionary dynamics of the chromatophore genome in three photosynthetic Paulinella species. *Scientific Reports* **9**, 2560 (2019).

16. Wernegreen, J. J. & Moran, N. A. Evidence for genetic drift in endosymbionts (Buchnera): analyses of protein-coding genes. *Mol Biol Evol* **16**, 83–97 (1999).

17. Moran, N. A. Accelerated evolution and Muller's rachet in endosymbiotic bacteria. *PNAS* **93**, 2873–2878 (1996).

18. Kuo, C.-H., Moran, N. A. & Ochman, H. The consequences of genetic drift for bacterial genome complexity. *Genome Res.* **19**, 1450–1454 (2009).

19. Muller, H. J. The relation of recombination to mutational advance. *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis* **1**, 2–9 (1964).

20. Bennett, G. M. & Moran, N. A. Heritable symbiosis: The advantages and perils of an evolutionary rabbit hole. *PNAS* **112**, 10169–10176 (2015).

21. Andersson, S. G. E. & Kurland, C. G. Reductive evolution of resident genomes. *Trends in Microbiology* **6**, 263–268 (1998).

22. Andersson, J. O. & Andersson, S. G. Genome degradation is an ongoing process
in Rickettsia. *Mol Biol Evol* **16**, 1178–1191 (1999).
23. Peek, A. S., Vrijenhoek, R. C. & Gaut, B. S. Accelerated evolutionary rate in
sulfur-oxidizing endosymbiotic bacteria associated with the mode of symbiont
transmission. *Mol Biol Evol* **15**, 1514–1523 (1998).
24. Cary, S. C. & Giovannoni, S. J. Transovarial inheritance of endosymbiotic
bacteria in clams inhabiting deep-sea hydrothermal vents and cold seeps. *PNAS*
90, 5695–5699 (1993).
25. Ikuta, T. *et al.* Surfing the vegetal pole in a small population: extracellular vertical
transmission of an 'intracellular' deep-sea clam symbiont. *Royal Society Open*
Science **3**, 160130 (2016).
26. Peek, A. S., Feldman, R. A., Lutz, R. A. & Vrijenhoek, R. C. Cospeciation of
chemoautotrophic bacteria and deep sea clams. *PNAS* **95**, 9962–9966 (1998).
27. Stewart, F. J., Young, C. R. & Cavanaugh, C. M. Evidence for homologous
recombination in intracellular chemosynthetic clam symbionts. *Mol. Biol. Evol.*
26, 1391–1404 (2009).
28. Peek, A. S., Gustafson, R. G., Lutz, R. A. & Vrijenhoek, R. C. Evolutionary
relationships of deep-sea hydrothermal vent and cold-water seep clams (Bivalvia:
Vesicomyidae): results from the mitochondrial cytochrome oxidase subunit I.
Marine Biology **130**, 151–161 (1997).
29. Moran, N. A., Munson Mark A., Baumann Paul & Ishikawa Hajime. A molecular
clock in endosymbiotic bacteria is calibrated using the insect hosts. *Proceedings*
of the Royal Society of London. Series B: Biological Sciences **253**, 167–171
(1993).
30. Ferri, E. *et al.* New Insights into the Evolution of Wolbachia Infections in Filarial
Nematodes Inferred from a Large Range of Screened Species. *PLOS ONE* **6**,
e20843 (2011).
31. Johnson, S. B., Krylova, E. M., Audzijonyte, A., Sahling, H. & Vrijenhoek, R. C.
Phylogeny and origins of chemosynthetic vesicomyid clams. *Systematics and*
Biodiversity **15**, 346–360 (2017).
32. Krylova, E. M. & Sahling, H. Vesicomyidae (Bivalvia): Current Taxonomy and
Distribution. *PLoS ONE* **5**, e9957 (2010).
33. Audzijonyte, A., Krylova, E. M., Sahling, H. & Vrijenhoek, R. C. Molecular
taxonomy reveals broad trans-oceanic distributions and high species diversity of
deep-sea clams (Bivalvia: Vesicomyidae: Pliocardiinae) in chemosynthetic
environments. *Systematics and Biodiversity* **10**, 403–415 (2012).
34. MolluscaBase. MolluscaBase. Vesicomyidae Dall & Simpson, 1901.
http://www.marinespecies.org/aphia.php?p=taxdetails&id=23140#distributions
(2019).
35. Newton, I. L. G. *et al.* The Calyptogena magnifica Chemoautotrophic Symbiont
Genome. *Science* **315**, 998–1000 (2007).
36. Kuwahara, H. *et al.* Reduced Genome of the Thioautotrophic Intracellular
Symbiont in a Deep-Sea Clam, Calyptogena okutanii. *Current Biology* **17**, 881–
886 (2007).
37. Kuwahara, H. *et al.* Loss of genes for DNA recombination and repair in the
reductive genome evolution of thioautotrophic symbionts of Calyptogena clams.
BMC Evolutionary Biology **11**, 285 (2011).
38. Tamas, I. *et al.* 50 Million Years of Genomic Stasis in Endosymbiotic Bacteria.
Science **296**, 2376–2379 (2002).

39. Roeselers, G. *et al.* Complete genome sequence of Candidatus Ruthia magnifica.
Stand Genomic Sci **3**, 163–173 (2010).
40. Anantharaman, K., Breier, J. A., Sheik, C. S. & Dick, G. J. Evidence for hydrogen
oxidation and metabolic plasticity in widespread deep-sea sulfur-oxidizing
bacteria. *PNAS* **110**, 330–335 (2013).
41. Liu, H., Cai, S., Zhang, H. & Vrijenhoek, R. C. Complete mitochondrial genome
of hydrothermal vent clam Calyptogena magnifica. *Mitochondrial DNA Part A* **27**,
4333–4335 (2016).
42. Ozawa, G. *et al.* Updated mitochondrial phylogeny of Pteriomorph and
Heterodont Bivalvia, including deep-sea chemosymbiotic Bathymodiolus mussels,
vesicomyid clams and the thyasirid clam Conchocele cf. bisecta. *Mar Genomics*
31, 43–52 (2017).
43. Yang, M., Gong, L., Sui, J. & Li, X. The complete mitochondrial genome of
Calyptogena marissinica (Heterodonta: Veneroida: Vesicomyidae): Insight into
the deep-sea adaptive evolution of vesicomyids. *PLoS One* **14**, (2019).
44. Ip, J. C.-H. *et al.* Host-Endosymbiont Genome Integration in a Deep-Sea
Chemosymbiotic Clam. *Molecular Biology and Evolution* (2020)
doi:10.1093/molbev/msaa241.
45. Stackebrandt, E. & Goebel, B. M. Taxonomic Note: A Place for DNA-DNA
Reassociation and 16S rRNA Sequence Analysis in the Present Species Definition
in Bacteriology. *International Journal of Systematic and Evolutionary
Microbiology,* **44**, 846–849 (1994).
46. Cohan, F. M. Sexual Isolation and Speciation in Bacteria. *Genetica* **116**, 359–370
(2002).
47. Fujiwara, Y. *et al.* Phylogenetic characterization of endosymbionts in three
hydrothermal vent mussels: influence on host distributions. *Marine Ecology
Progress Series* **208**, 147–155 (2000).
48. Petersen, J. M. *et al.* Hydrogen is an energy source for hydrothermal vent
symbioses. *Nature* **476**, 176–180 (2011).
49. Wertheim, J. O., Murrell, B., Smith, M. D., Kosakovsky Pond, S. L. & Scheffler,
K. RELAX: Detecting Relaxed Selection in a Phylogenetic Framework. *Mol Biol
Evol* **32**, 820–832 (2015).
50. Burke, G. R. & Moran, N. A. Massive Genomic Decay in Serratia symbiotica, a
Recently Evolved Symbiont of Aphids. *Genome Biol Evol* **3**, 195–208 (2011).
51. Stewart, F. J., Young, C. R. & Cavanaugh, C. M. Lateral Symbiont Acquisition in
a Maternally Transmitted Chemosynthetic Clam Endosymbiosis. *Mol Biol Evol*
25, 673–687 (2008).
52. Decker, C., Olu, K., Arnaud-Haond, S. & Duperron, S. Physical Proximity May
Promote Lateral Acquisition of Bacterial Symbionts in Vesicomyid Clams. *PLoS
One* **8**, (2013).
53. Ozawa, G. *et al.* Ancient occasional host switching of maternally transmitted
bacterial symbionts of chemosynthetic vesicomyid clams. *Genome biology and
evolution* **9**, 2226–2236 (2017).
54. Kojima, S. The distribution and the phylogenies of the species of genus
Calyptogena and those of vestimentiferans around Japan. *JAMSTEC Journal of
Deep-Sea Research* **11**, 243–248 (1995).
55. Goffredi, S. K. & Barry, J. P. Species-specific variation in sulfide physiology
between closely related Vesicomyid clams. *Marine Ecology Progress Series* **225**,
227–238 (2002).

56. Breusing, C., Johnson, S. B., Vrijenhoek, R. C. & Young, C. R. Host
hybridization as a potential mechanism of lateral symbiont transfer in deep-sea
vesicomyid clams. *Molecular Ecology* **28**, 4697–4708 (2019).
57. Naito, M. & Pawlowska, T. E. Defying Muller's Ratchet: Ancient Heritable
Endobacteria Escape Extinction through Retention of Recombination and
Genome Plasticity. *mBio* **7**, e02057-15 (2016).
58. Pérez-Brocal, V. *et al.* A small microbial genome: the end of a long symbiotic
relationship? *Science* **314**, 312–313 (2006).
59. Koga, R. & Moran, N. A. Swapping symbionts in spittlebugs: evolutionary
replacement of a reduced genome symbiont. *ISME J* **8**, 1237–1246 (2014).
60. Sudakaran, S., Kost, C. & Kaltenpoth, M. Symbiont Acquisition and Replacement
as a Source of Ecological Innovation. *Trends in Microbiology* **25**, 375–390 (2017).
61. Chong, R. A. & Moran, N. A. Evolutionary loss and replacement of Buchnera, the
obligate endosymbiont of aphids. *The ISME Journal* **12**, 898 (2018).
62. Latorre, A. & Manzano Marín, A. Dissecting genome reduction and trait loss in
insect endosymbionts. *Annals of the New York Academy of Sciences* **1389**, 52–75
(2017).
63. Saki, M. & Prakash, A. DNA damage related crosstalk between the nucleus and
mitochondria. *Free Radical Biology and Medicine* **107**, 216–227 (2017).
64. Chu, H. & Mazmanian, S. K. Innate immune recognition of the microbiota
promotes host-microbial symbiosis. *Nature immunology* **14**, 668–675 (2013).
65. Oldroyd, G. E. Speak, friend, and enter: signalling systems that promote
beneficial symbiotic associations in plants. *Nature Reviews Microbiology* **11**,
252–263 (2013).
66. Brownlie, J. C., Adamski, M., Slatko, B. & McGraw, E. A. Diversifying selection
and host adaptation in two endosymbiont genomes. *BMC Evolutionary Biology* **7**,
68 (2007).
67. Dale, C. & Moran, N. A. Molecular Interactions between Bacterial Symbionts and
Their Hosts. *Cell* **126**, 453–465 (2006).
68. Chong, R. A., Park, H. & Moran, N. A. Genome Evolution of the Obligate
Endosymbiont Buchnera aphidicola. *Mol Biol Evol* (2019)
doi:10.1093/molbev/msz082.
69. Wagner, G. P. & Gabriel, W. Quantitative Variation in Finite Parthenogenetic
Populations: What Stops Muller's Ratchet in the Absence of Recombination?
Evolution **44**, 715–731 (1990).
70. Rand, D. M., Haney, R. A. & Fry, A. J. Cytonuclear coevolution: the genomics of
cooperation. *Trends in Ecology & Evolution* **19**, 645–653 (2004).
71. Fares, M. A., Moya, A. & Barrio, E. Adaptive evolution in GroEL from distantly
related endosymbiotic bacteria of insects. *Journal of Evolutionary Biology* **18**,
651–660 (2005).
72. Howe, D. K. & Denver, D. R. Muller's Ratchet and compensatory mutation in
Caenorhabditis briggsae mitochondrial genome evolution. *BMC Evolutionary*
Biology **8**, 62 (2008).
73. Castillo, D. M. & Pawlowska, T. E. Molecular Evolution in Bacterial
Endosymbionts of Fungi. *Mol Biol Evol* **27**, 622–636 (2010).
74. Ohishi, K. *et al.* Long-term Cultivation of the Deep-Sea Clam Calyptogena
okutanii: Changes in the Abundance of Chemoautotrophic Symbiont, Elemental
Sulfur, and Mucus. *The Biological Bulletin* **230**, 257–267 (2016).
75. Andrews, S. *et al. FastQC*. (2012).

76. Zerbino, D. R. & Birney, E. Velvet: algorithms for de novo short read assembly
using de Bruijn graphs. *Genome Res.* **18**, 821–829 (2008).
77. Kearse, M. *et al.* Geneious Basic: An integrated and extendable desktop software
platform for the organization and analysis of sequence data. *Bioinformatics* **28**,
1647–1649 (2012).
78. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat*
Meth **9**, 357–359 (2012).
79. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics*
25, 2078–2079 (2009).
80. Bankevich, A. *et al.* SPAdes: A New Genome Assembly Algorithm and Its
Applications to Single-Cell Sequencing. *Journal of Computational Biology* **19**,
455–477 (2012).
81. Hahn, C., Bachmann, L. & Chevreux, B. Reconstructing mitochondrial genomes
directly from genomic next-generation sequencing reads—a baiting and iterative
mapping approach. *Nucleic Acids Res* **41**, e129–e129 (2013).
82. Tillich, M. *et al.* GeSeq – versatile and accurate annotation of organelle genomes.
Nucleic Acids Res **45**, W6–W11 (2017).
83. Overbeek, R. *et al.* The SEED and the Rapid Annotation of microbial genomes
using Subsystems Technology (RAST). *Nucleic Acids Res* **42**, D206–D214 (2014).
84. Darling, A. E., Mau, B. & Perna, N. T. progressiveMauve: Multiple Genome
Alignment with Gene Gain, Loss and Rearrangement. *PLoS ONE* **5**, e11147
(2010).
85. Tesler, G. GRIMM: genome rearrangements web server. *Bioinformatics* **18**, 492–
493 (2002).
86. Katoh, K. & Standley, D. M. MAFFT Multiple Sequence Alignment Software
Version 7: Improvements in Performance and Usability. *Mol Biol Evol* **30**, 772–
780 (2013).
87. Guindon, S. *et al.* New Algorithms and Methods to Estimate Maximum-
Likelihood Phylogenies: Assessing the Performance of PhyML 3.0. *Syst Biol* **59**,
307–321 (2010).
88. Cock, P. J. A. *et al.* Biopython: freely available Python tools for computational
molecular biology and bioinformatics. *Bioinformatics* **25**, 1422–1423 (2009).
89. Edgar, R. C. MUSCLE: multiple sequence alignment with high accuracy and high
throughput. *Nucleic Acids Res.* **32**, 1792–1797 (2004).
90. Goldman, N. & Yang, Z. A codon-based model of nucleotide substitution for
protein-coding DNA sequences. *Mol Biol Evol* **11**, 725–736 (1994).
91. Lemoine, F., Lespinet, O. & Labedan, B. Assessing the evolutionary rate of
positional orthologous genes in prokaryotes using synteny data. *BMC Evol Biol* **7**,
237 (2007).
92. Larget, B. R., Kotha, S. K., Dewey, C. N. & Ané, C. BUCKy: Gene tree/species
tree reconciliation with Bayesian concordance analysis. *Bioinformatics* **26**, 2910–
2911 (2010).
93. Kishino, H. & Hasegawa, M. Evaluation of the maximum likelihood estimate of
the evolutionary tree topologies from DNA sequence data, and the branching
order in hominoidea. *J Mol Evol* **29**, 170–179 (1989).
94. Kosakovsky Pond, S. L., Posada, D., Gravenor, M. B., Woelk, C. H. & Frost, S. D.
W. GARD: a genetic algorithm for recombination detection. *Bioinformatics* **22**,
3096–3098 (2006).

95. Ronquist, F. *et al.* MrBayes 3.2: Efficient Bayesian Phylogenetic Inference and
Model Choice Across a Large Model Space. *Systematic Biology* **61**, 539–542
(2012).
96. Warren, D. L., Geneva, A. J. & Lanfear, R. RWTY (R We There Yet): An R
package for examining convergence of Bayesian phylogenetic analyses.
Molecular Biology and Evolution msw279 (2017) doi:10.1093/molbev/msw279.
97. Zhang, Z. *et al.* Codon Deviation Coefficient: a novel measure for estimating
codon usage bias and its statistical significance. *BMC Bioinformatics* **13**, 43
(2012).
98. Pond, S. L. K., Frost, S. D. W. & Muse, S. V. HyPhy: hypothesis testing using
phylogenies. *Bioinformatics* **21**, 676–679 (2005).
99. Smith, M. D. *et al.* Less Is More: An Adaptive Branch-Site Random Effects
Model for Efficient Detection of Episodic Diversifying Selection. *Mol Biol Evol*
32, 1342–1353 (2015).
100.   Wołodźko, T. *extraDistr: Additional Univariate and Multivariate*
Distributions. (2019).
101.   Fisher, S. R. A. Confidence Limits for a Cross-Product Ratio. *Australian*
Journal of Statistics **4**, 41–41 (1962).
856

857 **Tables**
858 Table 1 Annotation statistics for symbiont and free-living genomes in this study.
859

Sample name	# of contigs (N50 Mbp)	Mean coverage	Genome size (Mbp)	GC %	# of CDS	# of tRNA	# of rRNA	% non-annotated CDS	NCBI Accession number	Reference
Ca. V. okutanii	1	9	1.02	32	939	35	3	7	AP009247	Kuwahara et al. (2007)
Ca. V. soyae (kilmeri)	1	110	1.02	32	983	36	3	11	CP060686	this paper
Ca. V. extenta	1	137	1.02	31	995	36	3	9	CP060685	this paper
Ca. V. diagonalis	1	110	1.02	31	1005	36	3	10	CP060680	this paper
Ca. V. gigas	1	153	1.04	31	979	36	3	10	CP060682	this paper
Ca. R. magnifica	1	14	1.16	34	976	36	3	7	CP000488	Newton et al. (2007)
Ca. R. pliocardia	1	113	1.23	37	1642	36	3	31	CP060688	this paper
Ca. R. southwardae	39 (0.63)	159	1.59	37	2035	36	3	28	JACRUS00	this paper
Ca. R. phaseoliformis	8 (0.37)	118	1.53	37	2210	36	3	39	JACRUR00	this paper
Ca. R. rectimargo	1	91	1.23	37	1476	37	3	29	CP060684	this paper
Ca. R. pacifica	1	140	1.18	37	1456	35	3	30	CP060683	this paper
Ca. B. thermophilus	1	126	2.83	39	2067	36	3	43	CP024634	
Ca. T. autotrophicus (SUP05)	1	106	1.51	39	1506	35	3	32	CP010552	Shah and Morris (2015)

860   Table 2 Overrepresented functional categories for genes exhibiting significant
861   evidence for episodic diversifying selection

	Overrepresented function	gene	reference locus_tag
Within free-living	**t-RNA biogenesis**	*pheS*	Rmag_0643
		tyrS	Rmag_0132
		valS	Rmag_0464
		fmt	Rmag_0785
	ribosome assembly	*rplB*	Rmag_0168
		rplO	Rmag_0184
		rpsM	Rmag_0187
		rpsS	Rmag_0169
	protein folding	*dnaJ*	Rmag_0352
		dnaK	Rmag_0353
		htpG	Rmag_0493
		clpB	Rmag_0787
	oxidative phosphorylation	*ccmE*	Rmag_0659
		ccmF	Rmag_0272
		CYTB/petB	Rmag_0010
		nhd	Rmag_0224
	sulfur oxidation	*soxB*	Rmag_0156
		soxY	Rmag_0807
	dissimilatory sulfate reduction	*aprM*	Rmag_0086
		aprAB	Rmag_0088, Rmag_0087
		dsrAB	Rmag_0870, Rmag_0869
bipartition FL-SYMB	**ribosomal proteins**	*rplJ*	Rmag_0813
		rpsA	Rmag_0592
		rpsC	Rmag_0171
		rpsH	Rmag_0179
	t-RNA ligases	*ileS*	Rmag_0340
		cysS	Rmag_0097
		thrS	Rmag_0648
		glyS	Rmag_0721
within symbionts	**chaperones, ribosomal proteins**	*rplC*	Rmag_0165
		rplD	Rmag_0166
		rplF	Rmag_0180
		rplL	Rmag_0812
		rpsC	Rmag_0171
		rpsP	Rmag_0990
	and t-RNA ligases	*glyS*	Rmag_0721
		argS	Rmag_0079
		aspS	Rmag_0396
		gltX	Rmag_0051
		ileS	Rmag_0340
		metG	Rmag_0570
		trpS	Rmag_0338
	sulphur metabolism	*dsrA*	Rmag_0870
		dsrP	Rmag_0859
		soxB	Rmag_0156

		cobB-cbiA	Rmag_0858
	electron donating/accepting reactions	nuoFG	Rmag_0242, Rmag_0243
		rnfABC	Rmag_0139, Rmag_0140, Rmag_0141
		rnfE	Rmag_0788
	ammonia assimilation	gltB	Rmag_0333, Rmag_1018
		glnD	Rmag_0475
within Clade II	de novo purine and pyrimidine biosynthesis	purO	Rmag_0969
		purL	Rmag_0837
		purA	Rmag_0531
		purC	Rmag_0392
		carB	Rmag_0875
		pyrDII	Rmag_0963
	carbon fixation	shmt/glyA	Rmag_0632
		rbcL	Rmag_0701
		cbbOQ	Rmag_0699, Rmag_0700
	DNA recombination and repair	ihfB	Rmag_0591
		yebC	Rmag_0394
		pcrA/uvrD	Rmag_0080, Rmag_0320
		recJ	Rmag_0649
		uvrA	Rmag_0263
within Clade I	DNA metabolism	parC	Rmag_0302
		dnaX	Rmag_0466
		ihfB	Rmag_0591
		exoI	Rmag_0946

862
863

864   Table 3 Sampling information and genome accession numbers for taxa in this study

Species	Accession	Locality[a]	Dive #	Lat	Long	Depth (m)	Year
Clade I							
Ca. Vesicomyosocius okutanii	AP009247.1	Sagami Bay		35.2	139.5	1157	2004
Phreagena okutanii (mtDNA)	AP014555	Sagami Bay		35.0150	139.222	852	2007
Ca. Vesicomyosocius soyae (kilmeri)	CP060686	Monterey Canyon (s)	V2059	36.7762	-122.084	985	2001
Phreagena soyoae (mtDNA)	MT947390						
Ca. Vesicomyosocius extenta	CP060685	Monterey Canyon (s)	T406	36.6088	-122.437	2889	2002
Phreagena extenta (mtDNA)	MT947388						
Ca. Vesicomyosocius diagonalis	CP060680	Monterey Canyon (s)	T488	36.2254	-122.885	3455	2002
Archivesica diagonalis (mtDNA)	MT947381						
Ca. Vesicomyosocius gigas	CP060682	Guaymas Basin (v)	T548	27.3400	-111.270	1754	2003
Archivesica gigas (mtDNA)	MT947383						
Clade II							
Ca. Ruthia magnifica	CP000488.1	East Pacific Rise (v)		9.8505	-104.300	2500	2004
"Calyptogena" magnifica (mtDNA)	KR862368	East Pacific Rise (v)		20.8305	-109.103	2601	2003
Ca. Ruthia pliocardia	CP060688	Blake Spur (s)	A3710	32.4948	-76.185	2155	2001
Pliocardia sp. Blake Ridge (mtDNA)	MT947391						
Ca. Ruthia southwardae	JACRUS00	Logatchev, MAR (v)	A3133	14.7532	-44.980	3038	1997
Abyssogena southwardae (mtDNA)	MT947385						
Ca. Ruthia phaseoliformis	JACRUR00	Aleutian Trench (s)	TVG	54.3050	-157.213	3550	1996
Abyssogena phaseoliformis (mtDNA)	MT947384						
Abyssogena phaseoliformis (mtDNA)	AP014557	Japan trench		39.1052	143.893	5347	2009
Ca. Ruthia rectimargo	CP060684	Monterey Canyon (s)	V2338	36.6816	-122.120	1540	2003
Calyptogena rectimargo (mtDNA)	MT947387						
Ca. Ruthia pacifica	CP060683	Monterey Canyon (s)	V2555	36.7739	-122.049	650	2004
Calyptogena pacifica (mtDNA)	MT947386						
Abyssogena mariana (mtDNA)	LC126311	Mariana trench		11.6569	143.049	5633	2013
"free-living"							
Ca. B. thermophilus	CP024634	East Pacific Rise (v)		9.82	-104.30	2518	2000
Ca. T autotrophicus	CP010552	Effingham Inlet		49.0369	-125.208	60	2013

 Table S6 Assembly and annotation statistics for mitochondrial genomes in this study.

Sample name	# of contigs (N50 Mbp)	Mean coverage	Genome size (Mbp)	GC %	# of CDS	# of tRNA	# of rRNA	NCBI Accession number	Reference
Phreagena okutanii	1	n.a.	16336	34	13	23	2	AP014555	Ozawa et al (2017)
Phreagena soyoae	1	25	19254	34	13	23	2	MT947390	this paper
Phreagena extenta	1	6	18098	33	13	22	2	MT947388	this paper
Archivesica diagonalis	1	8	20322	33	13	22	2	MT947381	this paper
Archivesica gigas	1	7	15625	35	13	21	2	MT947383	this paper
"Calyptogena" magnifica*	1	n.a.	19738	32	13	22	2	KR862368	Liu et al (2016)
Pliocardia sp.	1	20	18885	28	13	22	2	MT947391	this paper
Abyssogena southwardae	1	15	19082	29	13	24	2	MT947385	this paper
Abyssogena phaseoliformis	1	10	17997	31	13	23	2	MT947384	this paper
Abyssogena phaseoliformis*	1	n.a.	19424	30	13	24	2	AP014557	Ozawa et al (2017)
Calyptogena rectimargo	1	22	19326	32	13	25	2	MT947387	this paper
Calyptogena pacifica	1	18	19897	31	13	23	2	MT947386	this paper
Abyssogena mariana	1	n.a.	15927	30	13	23	2	LC126311	Ozawa et al (2017)

*Complete mitogenome

Figures and captions

868
869
Figure 1 Host and symbiont phylogenomic estimates. A) Neighbor-joining phylogeny
based on genetic distance (GTR model) between genome-wide alignments of
mitochondrial genomes (15272 bp). Numbers in black are bootstrap values. B)
Neighbor-joining phylogeny based on genetic distance (GTR model) between
genome-wide alignments of symbiont (Clade I; *Ca.* Vesicomyosocius in blue, Clade
II; *Ca.* Ruthia in green) and free-living (in red) genomes (761866 bp). Chromosome
schemes showing genome inversions and assembly fragmentation are displayed at the
end of the branches. Refer to text for a description of the genome structures. Numbers
in red are the genome-wide mean covariance factors; they represent the percentage of
protein-coding genes supporting each split of the phylogeny.

880
881

882
883 Figure 2 Relationship between symbiont and mitochondrial divergence. For each
884 holobiont pair, host and symbiont divergences are expressed as pairwise synonymous
885 substitutions rates (dS) in their respective genomes.

Figure 3 Codon bias in symbionts and free-living. A) Codon Deviation Coefficient
(CDC) spectra for each genome (all protein-coding genes). Within the free-living,
yellow: *Ca.* B. thermophilus; red: *Ca.* T. autotrophicus. B) Correlation between the
average CDC of free-living, *Ca.* Ruthia (green) and *Ca.* Vesicomyosocius (blue) core
genes. Linear regressions are shown. CDC values vary from 0 (no bias) to 1
(maximum bias). C) Selection parameter (k) spectra of genes for which a significant
change in selection was detected by RELAX. Note that k is on a log scale.

897
Figure 4 SEED category distribution of core genes under episodic diversifying
selection within phylogenetic clades (A, B, C, D), and on partitioning branches (C, D,
E, F). A) Distribution of all non-recombining core genes (dark grey, 652 loci) and loci
under selection within the free-living, *Ca.* Ruthia, and *Ca.* Vesicomyosocius clades
(light grey, 168 loci). The number of loci selected within each clade is represented in
the inset. B) Genes under selection within the free-living. C) Genes under selection
within *Ca.* Ruthia. D) Genes under selection within the *Ca.* Vesicomyosocius. E)
Distribution of all non-recombining core genes (dark grey, 652 loci) and loci under
selection on all partitioning branches (light grey, 80 loci). The number of loci selected
on each branch is represented in the inset. F) Genes under selection on branch a. G)
Genes under selection on branch b. H) Genes under selection on branch c. Note that
genes may be represented in multiple functional categories and multiple clades or
branches. SEED categories significantly overrepresented (in red) and
underrepresented (in blue) in the groups compared to the core genome are highlighted.
Refer to text for further breakdown of these categories. NA: no functional annotation.

A) Core gene sequences

B) Gene conservation pattern

913
Figure S1 Distance-based neighbour-joining trees established from A) a concatenated
alignment of 652 non-recombining core protein-coding genes sequences (618342 bp,
HKY nucleotide substitution model). In red are the number of genes under episodic
diversifying selection in each branch; B) the presence/absence of positionally
orthologous genes (Jaccard distance on 6110 genes). Numbers above branches are
bootstrap support values.

921
Figure S2 Venn diagram representing the 6110 unique and shared putative genes
amongst the free-living, *Ca.* Vesicomyosocius and *Ca.* Ruthia. The outer circles
represent the pan-genome while the inner circles represent the core-genome of the
groups. Free-living: *Ca.* B. thermophilus and *Ca.* T. autotrophicus); *Ca.* Ruthia: *Ca.* R.
magnifica, *Ca.* R. phaseoliformis, *Ca.* R. pacifica, *Ca.* R. rectiomargo, *Ca.* R.
pliocardia, and *Ca.* R. southwardae; *Ca.* Vesicomyosocius: *Ca.* V. okutanii, *Ca.* V.
soyoae, *Ca.* V. diagonalis-extenta, and *Ca.* V. gigas. Groups in which more than 50%
of the genes are unannotated are identified in red. The complete orthology is available
in Table S1.

Figure S3 Sampling locations. Inset depicts samples collected from varying depths in
Monterey Bay.

934