Similarities in the behaviour of dance followers among honey bee species suggests a conserved mechanism of dance communication

Ebi Antony George[1*], Smruti Pimplikar[1], Neethu Thulasi[1,], Axel Brockmann[1]

1 National Centre for Biological Sciences, Bangalore, India – 560065

ORCID

E.A.G.: 0000-0002-5533-5428

S.P.: 0000-0002-6683-5946

N.T.: 0000-0003-4702-8900

A.B.: 0000-0003-0201-9656

* corresponding author:

Ebi Antony George

Email ID: ebiantonyg@ncbs.res.in, ebi.antony.george@gmail.com

Abstract

Group living organisms rely on intra-group communication to adjust individual and collective behavioural decisions. Complex communication systems are predominantly multimodal and combine modulatory and information bearing signals. The honey bee waggle dance, one of the most elaborate form of communication in invertebrates, stimulates nestmates to search for food and communicates symbolic information about the location of the food source. Previous studies on the dance behaviour in diverse honey bee species demonstrated distinct differences in the combination of visual, auditory, olfactory, and tactile signals produced by the dancer. We now studied the behaviour of the receivers of the dance signals, the dance followers, to explore the significance of the different signals in the communication process. In particular, we ask whether there are differences in the behaviour of dance followers between the 3 major Asian honey bee species, *A. florea, A. dorsata and A. cerana*, and whether these might correlate with the differences in the signals produced by the dancing foragers. Our comparison demonstrates that the behaviour of the dance followers is highly conserved across all 3 species despite the differences in the dance signals. The highest number of followers was present lateral to the dancer throughout the waggle run, and the mean body orientation of the dance followers with respect to the waggle dancer was close to 90° throughout the run for all 3 species. These findings suggest that dance communication might be more conserved than implied by the differences in the signals produced by the dancer. Along with studies in *A. mellifera*, our results indicate that all honey bee species rely on tactile contacts between the dancer and follower to communicate spatial information. The cues and signals that differ between the species may be involved in attracting the followers towards the dancer in the different nest environments.

Keywords

Apis florea, Apis cerana, Apis dorsata, waggle dance evolution, spatial information transfer, tactile hypothesis

Introduction

Organisms have evolved various ways to communicate amongst themselves [1]. Communication involves indirect cues and direct signals and varies in its complexity [2,3]. Complexity in communication systems correlates with social group complexity [4, but see 5]. Social communication mechanisms consist of multiple signal channels which can be of the same modality, e.g. ant pheromone trails, or different modalities, as in the case of ritualised courtship signals in birds [6,7]. In these communication systems, the different signals are either equally informative, or one of the signals contains the information and the others act as modulators, enhancing the effect or spread of the signal [8]. Finally, environmental factors and plasticity in the signal can lead to divergence in signals across closely related species [9–11].

One of the most elaborate types of social communication in invertebrates is the honey bee waggle dance used by foragers returning from profitable food sources to recruit nestmates [12]. The waggle dance motivates foragers to fly out and in addition encodes spatial information about the food source [13,14]. Each waggle dance consists of multiple circuits and one circuit contains two phases; a straight walking phase in which the dancer waggles its abdomen back and forth (the waggle run or the waggle phase) and a circular walking phase which brings the dancer back towards the point of origin of the first phase (the return phase). The duration of the waggle run corresponds to the distance to the food source [12,14]. In *Apis mellifera*, in which foragers dance in the dark on vertical combs, the body orientation of the dancer with respect to the vertical (gravity) axis during the waggle run corresponds to the direction of the food source from the hive with respect to the sun's azimuth [12]. Further, the duration of the return phase corresponds to the reward value of the food source as perceived by the forager [15]. For food sources very close to the hive, the dance circuit becomes nearly circular with a very short waggle run [16].

Cues and signals from the environment and nestmates can modulate the probability and intensity of dance behaviour [15,17–19]. Interactions with nestmates in the hive inform nectar foragers about the colony food stores and the nectar influx into the colony [18,20–22]. In addition, interactions with other foragers provide information

about predation and overcrowding at the food source [23–26]. An individual forager's

dance activity is modulated by the perceived reward value of the food source along

with information from these interactions [19]. This in turn drives recruitment to each

food source proportional to its relative reward value, which leads to an efficient

distribution of the colony's foraging force [27]. Thus, the waggle dance acts as the

primary regulatory mechanism of the colony's recruitment activity in addition to its

role in the efficient spatial distribution of the colony's foragers.

Although extensive research has been done on the honey bee waggle dance behaviour,

the mechanism underlying the transfer of spatial information has remained elusive

[28,29]. The experimental difficulty lies in determining which of the multiple dances

followed is used to obtain information [27] and in tracking whether the follower

visited the indicated food source. Moreover, recent studies showed that followers can

choose to rely on either the information from the dancer or their own memory [30,31].

Currently, there are two major hypotheses on which signals dance followers use to

obtain spatial information from the dancer. The "tactile hypothesis" proposes that

dance followers use tactile signals, associated with physical contact between the

dancer and follower [32,33] or even mechanosensory signals, associated with the air

flow caused by a dancer's vibrating wings [34,35]. The "follow hypothesis" suggests

that followers obtain information from dancers by following the path of the dancer

from behind [36]. In this case, followers receive the dance information from their own

body positions and walking paths to calculate the direction and distance of the food

source being advertised. Studies on the mechanism of spatial information transfer in

A. mellifera offered preliminary evidence for both hypotheses [32,36]. Instead,

comparative studies on dance communication including the behaviour of the dance

followers in different honey bee species might help to decide the controversy [37].

Interestingly, there are characteristic differences between the species in the

combination of signals generated by the dancer [37–42]. Dances of the dwarf honey

bees (e.g. *A. florea*), which take place on a horizontal surface exposed to the sun,

include a conspicuous visual signal (raised abdomens) but no auditory signals [43].

The giant honey bees (e.g. *A. laboriosa* and *A. dorsata*) usually dance on the vertical

surface of the bee curtain exposed to the sun [41]. Dances of the diurnal *A. laboriosa*

are silent [44], whereas those of the crepuscular *A. dorsata* contain auditory signals

[45]. All cavity nesting species investigated (e.g. *A. mellifera, A. cerana* and *A.*

nigrocincta), which perform dances in the hive, produce auditory signals [41].

However, it is unclear whether these differences in dance signals imply the evolution

of different mechanisms for information transfer across the genus or whether the

mechanism is the same with the different signals and cues serving to attract followers

[46,47].

We performed a comparative study to explore the dance follower behaviour of Asian

honey bees and to test if possible differences in dance follower behaviour might

correlate with differences in the signals and cues produced by the dancer. Parallel

changes in the behaviour of the dancer and dance follower would be a strong

argument for the evolution of different mechanisms of information transfer across the

genus *Apis*. On the other hand, a high degree of similarity in the behaviour of the

dance follower would suggest that the major mechanism of information transfer is

conserved. Further, we monitored the spatial positions of the followers to obtain

evidence for either of the two proposed hypotheses regarding the mechanism of

information transfer. If followers arrange themselves towards the side of the dancer,

then it is likely that they use tactile cues ("tactile hypothesis") to obtain information

about the food source [32]. In contrast, if followers orient themselves behind the

dancer, they are more likely using their own body orientation ("follow hypothesis") to

determine the spatial position of the food source [36].

Methods

Colonies

Experiments were done with two *A. florea* and *A. cerana* colonies each and one *A.*

dorsata colony. For *A. florea*, we obtained colonies which were nesting on the

National Centre for Biological Sciences (NCBS) campus, Bangalore. The *A. cerana*

colonies was obtained from Karnataka Apiaries, Bangalore. For *A. dorsata*, we

observed a colony which had made a nest in the Naik Bhavan building adjacent to our

experimental location in the University of Agricultural Sciences (UAS), Gandhi Krishi

Vignana Kendra (GKVK), Bangalore. The first set of colonies from all 3 species was

observed in January - March 2017, and the second *A. cerana* and *A. florea* colony was

observed in February - April 2018. The experiments were part of a larger study on

dance dialects in Asian honey bees and are described in detail in section 2.2 of Kohl *et*

al., 2020 (in preparation). A brief description of the same experimental protocol is

provided below.

Experimental Location

Our experiments were performed in the Botanical Garden at GKVK (latitude: 13.07,

longitude: 77.57). The garden covers an area of 26 hectares and is home to around

2000 plant species including trees and shrubs. This provided dense vegetation cover

and hence good optic flow for the foragers (fig S1 *a*). We obtained a straight line

transect of at least 500m from the colony location to train foragers along.

Colony Preparation

We cut the branch holding the *A. florea* colonies and shifted them into a wooden box

which had one opening at the top (fig S1 *b*). This opening was used to record the

activity on the crown area of the colony, where most of the recruitment activity

happens [43]. In the case of *A. dorsata,* we did not shift the colony from the building

where it had made its hive. We used a feeder box (fig S1 *c*) to train foragers from the

colony to the ground first, and then along the transect. Four frame *A. cerana* colonies

were housed in a specially constructed box, with glass walls on one side (fig S1 *d*).

This allowed us to observe the waggle dances and the dance follower behaviour in a

non-invasive manner and without having to move the colony into a traditional 2

frame vertical observation hive.

Distance training

Apis florea and Apis cerana

We used a similar protocol for distance training in *A. florea* and *A. cerana*. We trained

foragers to an artificial feeder near the colony filled with sucrose solution scented with

star anise (*Illicium verum*) extract. The sucrose concentration at the feeder was

adjusted between 1 and 2.5 M depending on the number of foragers visiting our food

source. Once 5-10 foragers were trained to the artificial feeder, we shifted it away from

the hive, along the transect. At 25 m we individually paint marked foragers at the

feeder using Uni POSCA Paint markers (Uni Mitsubishi Pencil, UK). Once all foragers

coming to the feeder were marked, the feeder was shifted further along the transect.

We recorded the dance activity of the marked foragers for one hour each at 100m,

200m, 300m, 400m and 500m using a Sony HDR CX260V Handycam (Sony

Corporation, Tokyo). The videos were recorded at 1080p and 50 frames per second and

covered the dance floor area (crown area in *A. florea* and the frame facing the glass

wall of the box in *A. cerana*).

Apis dorsata

Since the *A. dorsata* colony was on the side of a building, we used a different protocol

for training the foragers. First, we got foragers to come to a long rod with a piece of

comb filled with sucrose solution which was kept next to the bee curtain. We then

transferred these foragers carefully into a feeder box (fig S1 *c*), closed the box and

transported it to an artificial feeder at the ground level. We released the foragers at the

feeder and repeated the process till foragers came to the feeder on their own. After

this, the same protocol of paint marking, feeder shifts, and video recordings were done

as described earlier. We recorded the dance activity of *A. dorsata* foragers at distances

of 100m, 200m, 300m and 400m. Since we recorded dances for the *A. dorsata* colony

during a time of plentiful flowering, we could not get foragers to come to a feeder

further than 400m from the hive. The transect used for *A. dorsata* feeder training was

in the opposite direction of the ones used for *A. florea* and *A. cerana* as the colony had

nested on a building at the opposite end of the botanical garden.

Video Analysis

For the video analysis, individual foragers were first shortlisted based on whether they

were active at the feeder at multiple distances. We then analysed each dance circuit in

the dances by these individuals to determine the duration of the waggle run. The

videos were observed frame-by-frame in Virtual Dub 1.10.4

(http://www.virtualdub.org). The first frame in which a focal bee clearly moved its

abdomen laterally or dorsoventrally was defined as the start of the waggle run in that

circuit. The frame in which the bee stopped waggling its abdomen and started turning

to the left or right was defined as the end of the waggle run. The time between the
start and the end frames was calculated as the duration of the waggle run.

Follower Behaviour

We analysed dance follower behaviour for each waggle run in each waggle dance and
did not observe the behaviour of dance attendants. Dance followers were defined as
those bees which positioned themselves within one bee length of the dancer while we
excluded dance attendants, who were beyond this distance threshold and not
following the dancer [33]. In each run, we focussed on 3 phases (time-points); the
start, middle and end, based on the waggle run duration we had calculated (fig 1 *a*).
This was done to look at whether there was a change in the number of followers as the
run progressed [32]. At each time-point, the number of followers present in the
following three zones (fig 1 *b*) around the dancer was counted; the anterior zone
around the head region of the dancer, the lateral zone near the thorax and abdomen,
and the posterior zone behind the abdomen [32].

Figure 1

211
212

Schematic of the analysis done on the dance followers. (a) Each waggle run was
divided into 3 phases, the start, middle and end and the number of followers around
the dancer was quantified. (b) Followers around the dancer were grouped into 3 zones
(Anterior, Lateral and Posterior) based on the position they occupied around the
dancer (area of the zones in the figure are representative). (c) The orientations of the
followers were then quantified by obtaining the angle made by a vector representing
the follower with respect to the vector representing the dancer in the clockwise
direction.

The body angles of the follower with respect to the angle of the dancer were also
quantified. We used OnScreenProtractor v0.3 (http://osprotractor.sourceforge.net), to
do this manually. We first used the program to make a vector representing the dancer,
pointing in the direction of the dancer's head, away from its abdomen. We then made
a similar vector for the follower, but from its head to abdomen. Finally, the body angle
of the follower was quantified as the angle subtended by the follower vector with
respect to the dancer vector in the clockwise direction (fig 1 *c*). In total, we calculated
5036 follower positions and body angles (from 330 waggle runs of 41 waggle dances) in
A. florea, 1363 (from 119 waggle runs of 7 waggle dances) in *A. dorsata* and 4989 (from
411 waggle runs of 35 waggle dances) in *A. cerana* (table S1). Since followers were not
individually identifiable, it is possible that some of the followers we counted were the
same across multiple runs and dances.

Statistical Analysis

Number of Followers

We used a model comparison approach to analyse the variables which affected the
number of dance followers. The dataset of the number of followers in each zone
during each of the waggle run phase was zero inflated, with 21.17 % (1639 out of 7740)
of the observations being zero. To take this into account, we fit zero-inflated Poisson
models [48] with bee ID as a random effect. In total, 27 models were built (table S2)
with a combination of our 4 predictors for the conditional part of the model: i) zone of
dance follower (a categorical variable of 3 levels; Anterior, Lateral and Posterior), ii)
phase of waggle run (a categorical variable of 3 levels; Start, Middle and End), iii)
species (a categorical variable of 3 levels; *A. florea, A. cerana* and *A. dorsata*) and iv)
distance (a continuous variable which was scaled with a mean of zero and a standard
deviation of 1). For the zero-inflated part of the model, we only included the 3
categorical variables. Although we tried to include the same predictors for the zero-
inflated part of the model in all 27 models, we had to fit a reduced number of
predictors in 5 of them, to prevent errors with model convergence (see table S2). We
then compared the models based on their AIC values and shortlisted the models
within a cut-off of 0.95 cumulative Akaike weights [49]. The list of predictors, effect
sizes and confidence intervals from the shortlisted model(s) was then obtained.

Further, we performed multiple comparisons (with Tukey corrections for multiple

comparisons) of the estimated mean number of followers. We focussed on: i)

comparison between the 3 species, ii) comparisons between the different zones within

each waggle run phase and iii) comparisons between the same zones across the waggle

run phases. We did these specific comparisons as in our shortlisted model, the

important predictors were the species and an interaction term between the waggle run

phase and the zone of the follower (table S2 and S3).

Orientation of followers

We used circular statistics to analyse the orientation of the dance followers. We first

constrained the body angles of all the dance followers to lie between 0° and 180°, by

converting all the angles greater than 180° to their mirror images in the 0° - 180° range

(e.g., 210° was converted to 150°, 270° to 90° and 300° to 60°). We based this on the

assumption that occupying either the left or the right side of the dancer provided

similar access to information for the follower. In addition, this prevented us from

obtaining biased estimates of the circular mean and length due to potential bimodality

in the circular distributions to the left and the right of the dancer. We focussed the

statistical analysis on the median orientation of the dance follower in each of the three

waggle run phases within each species. First, we checked whether the circular

distribution of the dance follower orientation angles during each phase for each

species showed uniformity using the Rayleigh test for unimodal departures from

uniformity [50]. Then we checked whether the distributions showed reflective

symmetry [51]. Finally, we used Fishers Circular Nonparametric test to compare the

median angles of the circular distributions [52,53]. We compared the circular

distributions for each of the 3 pairs of the waggle run phase to determine which were

different from each other (Start – Middle, Start – End and Middle – End).

All the models and the plots were made in R [54] using the RStudio IDE [55]. The

GLMMs were fit using the glmmTMB package [56], model selection and averaging

were done using the MuMIn package [57] and the model assumptions were checked

using the DHARMa package [58]. Multiple comparisons were done using the emmeans

package [59]. Circular statistics were done using the circular package [60] and code

found in Pewsey et al. 2013.

Results

Number of Followers

We found only one model at the 0.95 cut-off level for cumulative sum of Akaike weights based on our model comparisons (table S3). In this model, the important predictors were the species and an interaction between the waggle run phase and the zone of the follower, but not the distance (table S4). In the zero-inflated part of this model, none of the predictors significantly correlated with the number of absences (i.e., in the number of observations where there were no followers), and these results are provided in the supplementary information (table S4).

Figure 2

Estimated number of followers and the 95% confidence intervals (circles and error bars) for the different predictors present in the conditional part of the final shortlisted model (model 18, see Tables 2 and 3). (a) Waggle run phase and the zone of the follower had an interactive effect on the number of followers. The number of followers in the anterior zone were similar throughout the waggle run. The number of followers in the lateral zone showed an increasing trend, with significantly higher numbers than the other two zones in the start of the run. The number of followers in the posterior

zone was similar to the number of followers in the anterior at the beginning of the run
but increased to numbers similar to the lateral zone in the middle and the end of the
run. The circles and error bars are coloured according to the waggle run phase with
blue representing the start, cyan representing the middle and green representing the
end of the waggle run. The alphabets above each circle represents results from the
multiple comparisons done (estimates with different alphabets were significantly
different from each other at the p < 0.05 level). Upper case alphabets represent
differences in the number of followers present in the same waggle run phase across
zones. Lower case alphabets represent differences in the number of followers present
in the same zone across different waggle run phases. (b) The species had an effect on
the number of followers present in the waggle run. The number of followers per run
for *A. florea* and *A. dorsata* were much higher than the number of followers for *A.
cerana* (different roman numerals above the estimates represent significant differences
at the p < 0.05 level). The circles and error bars are coloured according to the species,
with red for *A. florea*, orange for *A. dorsata* and purple for *A. cerana*.

Effect of species

The species had an effect on the number of followers in the conditional model, but

this effect was the same across the different zones, species and distance, i.e., there was

no interaction between species and any of the other predictors (table S4 and fig 2 and

S2). There were fewer followers per run in *A. cerana* as compared to both *A. florea* and

A. dorsata (*A. cerana* vs *A. florea*: estimated mean number of followers = 1.188 vs 1.512, t

ratio = 10.705, *p* < 0.001; *A. cerana* vs *A. dorsata*: estimated mean = 1.188 vs 1.549, t ratio

= 7.082, *p* < 0.001). The number of followers in *A. florea* and *A. dorsata* were not

significantly different (*A. florea* vs *A. dorsata*: stimated mean = 1.512 vs 1.549, t ratio = -

0.621, *p* = 0.809).

Effect of waggle run phase and zone of the followers

The waggle run phase and the zone around the dancer had an interactive effect on the

number of followers in the conditional model and hence their main effects are not

considered (table S4 and fig 2 and S2). Within each waggle run phase, there were

differences in the number of followers in the different zones. At the Start of the waggle

run, there were more followers in the lateral zone as compared to both the anterior

and the posterior zone (Anterior vs Lateral: estimated mean number of followers =

1.142 vs 1.521, t ratio = -6.360, *p* < 0.001; Lateral vs Posterior: estimated mean = 1.521 vs

1.209, t ratio = 5.578, *p* < 0.001). The number of followers in the anterior and posterior

zone were similar (Anterior vs Posterior: estimated mean = 1.142 vs 1.209, t ratio = -

1.213, p = 0.446).

In the middle of the waggle run phase, there were fewer followers in the anterior zone

as compared to both the lateral and the posterior zone (Anterior vs Lateral: estimated

mean = 1.057 vs 1.690, t ratio = -11.226, p < 0.001; Anterior vs Posterior: estimated mean

= 1.057 vs 1.748, t ratio = -12.110, p < 0.001). The number of followers in the lateral and

posterior zone were similar (Lateral vs Posterior: estimated mean = 1.690 vs 1.748, t

ratio = -0.927, p = 0.623).

In the end of the waggle run phase, there were fewer followers in the anterior zone as

compared to both the lateral and the posterior zone (Anterior vs Lateral: estimated

mean = 1.025 vs 1.838, t ratio = -14.046, p < 0.001; Anterior vs Posterior: estimated mean

= 1.025 vs 1.743, t ratio = -12.656, p < 0.001). The number of followers in the lateral and

the posterior zone were similar (Lateral vs Posterior: estimated mean = 1.838 vs 1.743, t

ratio = 1.478, p = 0.301).

There were differences between each zone in the number of followers across the

waggle run phase. In the Anterior zone, there were similar number of followers in all 3

phases of the waggle run (Start vs Middle: estimated mean = 1.142 vs 1.057, t ratio =

1.597, p = 0.247; Start vs End: estimated mean = 1.142 vs 1.025, t ratio = 2.212, p = 0.069;

Middle vs End: estimated mean = 1.057 vs 1.025, t ratio = 0.654, p = 0.790).

In the Lateral zone, the number of followers increased as the waggle run progressed.

There were fewer followers in the Start of the run as compared to both the Middle and

the End (Start vs Middle: estimated mean = 1.521 vs 1.690, t ratio = -2.802, p = 0.014;

Start vs End: estimated mean = 1.521 vs 1.838, t ratio = -5.125, p < 0.001). Although there

was an increase in the number of followers from the Middle to the End of the waggle

run, this difference was non-significant (Middle vs End: estimated mean = 1.690 vs

1.838, t ratio = -2.333, p = 0.051).

In the Posterior zone, the number of followers changed with the waggle run phase

There were fewer followers in the Start of the run as compared to both the Middle and

the End (Start vs Middle: estimated mean = 1.209 vs 1.748, t ratio = -9.24, p < 0.001;

Start vs End: estimated mean = 1.209 vs 1.743, t ratio = -9.169, p < 0.001). There were no

significant differences in the number of followers between the Middle and the End of the waggle run (Middle vs End: estimated mean = 1.748 vs 1.743, t ratio = 0.072, p = 0.997).

Effect of distance

Distance had no effect on the number of followers (fig S3). Distance was not a predictor that was present in the short-listed models (table S4). Thus, the main effects of the waggle run phase and the interactive effects of zone and species were similar across all distances.

Orientation of followers

The species differed in the change of the median circular orientation of the dance followers from the start to the end of the waggle run, even though all the median orientations were close to 90° (table 1, fig 3 and S4).

Apis florea

The mean of the circular distribution for the dance follower angles during each of the 3 phases in *A. florea* were all close to 90° (mean (95 % confidence interval) - Start: 88.293 (85.2671 - 91.3188); Middle: 105.152 (102.449 - 107.8549); End: 98.2651 (95.5425 - 100.9876); see table 1, fig 3 and S4). None of the 3 circular distributions were uniform (Rayleigh test statistic – Start: 0.668, p < 0.001; Middle: 0.669, p < 0.001; End: 0.656, p < 0.001) or reflectively symmetric (test statistic – Start: 2.879, p = 0.004; Middle: 8.557, p < 0.001; End: 4.529, p < 0.001). The medians of the 3 circular distributions were significantly different from each other (median – Start: 86.69; Middle: 108.43; End: 100.33; Fisher test statistic – Start vs Middle: 58.898, p < 0.001; Start vs End: 25.878, p < 0.001; Middle vs End: 10.109, p = 0.001).

Table 1

Species	Waggle Run Phase	μ (CI)	ρ (CI)	β_2 (CI)	$\bar{\alpha_2}$ (CI)	Median
Apis florea	Start	88.293 (85.2671 - 91.3188)	0.6676 (0.6834 - 0.6518)	87.466 (85.7509 - 89.1812)	85.6183 (83.4845 - 87.752)	86.69
	Middle	105.152 (102.449 - 107.8549)	0.6688 (0.6836 - 0.654)	96.8049 (95.282 - 98.3277)	84.6392 (82.6064 - 86.6719)	108.43

		μ	ρ	β_2	$\overline{\alpha_2}$	median
	End	*98.2651 (95.5425 - 100.9876)*	*0.6561 (0.6703 - 0.6419)*	*93.662 (92.0888 - 95.2352)*	*86.8135 (84.9144 - 88.7126)*	*100.33*
Apis dorsata	Start	85.6489 (80.1198 - 91.178)	0.6755 (0.7044 - 0.6466)	0	84.8189 (80.9594 - 88.6785)	86.36
	Middle	89.3173 (84.1632 - 94.4714)	0.6701 (0.6973 - 0.6428)	0	85.1766 (81.5321 - 88.821)	91.925
	End	92.2348 (86.8583 - 97.6112)	0.6576 (0.6858 - 0.6293)	0	86.5634 (82.8405 - 90.2862)	90.665
Apis cerana	*Start*	*83.3729 (80.3027 - 86.4432)*	*0.6471 (0.6626 - 0.6316)*	*85.7728 (83.9274 - 87.6182)*	*88.2383 (86.1242 - 90.3524)*	*79.555*
	Middle	*94.5799 (91.966 - 97.1939)*	*0.6858 (0.6999 - 0.6718)*	*91.4155 (90.0163 - 92.8147)*	*83.1709 (81.2688 - 85.0729)*	*94.075*
	End	*100.9708 (98.2417 - 103.6999)*	*0.6695 (0.6842 - 0.6548)*	*94.9577 (93.4502 - 96.4652)*	*84.8524 (82.8413 - 86.8636)*	*104.4*

List of circular summary statistics for the dance follower angles in each of the 3 waggle
run phase for each of the 3 species. The mean direction (μ), mean resultant length (ρ),
second central sine movement (β_2), second central cosine movement ($\overline{\alpha_2}$) along with
the estimated 95% confidence intervals is provided. In addition, the median of the
circular distribution is also provided. The circular distributions for each of the 3
phases for *A. dorsata* showed reflective symmetry and hence the central sine
movements are set to 0. The waggle run phases within each species which significantly
differed in their medians at the p < 0.05 level are highlighted in italics.

Apis dorsata

The mean of the circular distribution for the dance follower angles during each of the

3 phases in *A. dorsata* were all close to 90° (mean and 95 % confidence interval - Start:

85.6489 (80.1198 - 91.178); Middle: 89.3173 (84.1632 - 94.4714); End: 92.2348 (86.8583 -

97.6112); see table 1, fig 3 and S4). None of the 3 circular distributions were uniform

(Rayleigh test statistic – Start: 0.676, *p* < 0.001; Middle: 0.671, *p* < 0.001; End: 0.658, *p* <

0.001) but all of them were reflectively symmetric (test statistic – Start: 0.107, *p* = 0.915;

Middle: 0.193, *p* = 0.847; End: 0.105, *p* = 0.916). The medians of the 3 circular

distributions did not significantly differ from each other (median – Start: 86.36;

Middle: 91.92; End: 90.66; Fisher test statistic – Start vs Middle: 0.404, *p* = 0.525; Start

vs End: 0.194, *p* = 0.659; Middle vs End: 0.017, *p* = 0.896).

Figure 3

Relative circular density plots of the constrained dance follower angle for each of the
different waggle run phase for (a) *A. florea*, (b) *A. dorsata* and (c) *A. cerana*. The
relative circular density is obtained by normalising the density in each bin to the
highest density value, such that the relative circular density for each bin lies within 0
and 1. The arrows in the plot represent the mean of the constrained circular
distribution with the shaded region around the arrow representing the 95%
confidence interval. The length of the arrow corresponds to the mean resultant length
of the distribution (ρ). The lines representing the density plot as well as the arrows are
coloured based on the waggle run phase, with blue representing the start, cyan
representing the middle and green representing the end of the waggle run.

Apis cerana

The mean of the circular distribution for the dance follower angles during each of the

3 phases in *A. cerana* were all close to 90° (mean and 95 % confidence interval - Start:

83.3729 (80.3027 - 86.4432); Middle: 94.5799 (91.966 - 97.1939); End: 100.9708 (98.2417

- 103.6999); see table 1, fig 3 and S4). None of the 3 circular distributions were uniform

(Rayleigh test statistic – Start: 0.647, *p* < 0.001; Middle: 0.686, *p* < 0.001; End: 0.669, *p* <

0.001) or reflectively symmetric (test statistic – Start: 4.449, *p* < 0.001; Middle: 1.978, *p*

= 0.048; End: 6.358, *p* < 0.001). The medians of the 3 circular distributions were

significantly different from each other (median – Start: 79.55; Middle: 94.07; End:

104.4; Fisher test statistic – Start vs Middle: 24.725, *p* < 0.001; Start vs End: 61.418, *p* <

0.001; Middle vs End: 17.150, *p* = 0.001).

Discussion

The results of our study demonstrate that dance followers in three Asian honey bee species, *A. florea*, *A. dorsata* and *A. cerana* behave similarly throughout the waggle run. At the start of the run, most followers positioned themselves laterally to the dancer. Then, in the middle and the end of the run, the number of followers in the lateral and posterior positions around the dancer was similar. Further, in all three species, the mean orientation of the dance follower was close to 90° throughout the waggle run. The species differed in the number of followers per run. Dances by *A. florea* and *A. dorsata* foragers attracted larger number of followers than those by *A. cerana* foragers. The distance of the food source, and hence the duration of the waggle phase had no effect on the average number of followers present per waggle run in all 3 species.

The waggle run is hypothesized to represent a ritualization of the initiation of flight towards the food source [37–39,42]. In the open nesting and phylogenetically ancestral honey bee ,*A. florea*, dances are indeed oriented in direction of the food source and there is no transposition of the direction information to a vertical plane, unlike in the giant and cavity nesting honey bees [43]. With respect to these two traits, one would predict that there should be strong differences in the dance follower behaviour between *A. florea* and the other species. Specifically, dance followers should align themselves behind the dancers as this would allow them to most easily obtain the direction of the food source. However, in contrast to these two predictions *A. florea* dance followers neither aligned themselves behind the dancer nor showed any other major differences in their behavior in comparison with followers in the other species'. This finding supports the idea of an evolutionarily conserved mechanism of spatial information transfer in the dance behavior in all honey bees species [for *A. mellifera*, see 34].

Regarding the question of which sensory signals the dance followers use to obtain the spatial information of the dance, our results provide two arguments for the "tactile hypothesis". The first is the higher number of followers in the lateral positions around the dancer throughout the run, which is similar to the pattern observed in *A. mellifera* [28,32,61]. The second line of evidence comes from the median body angle of the

dance followers, which was close to 90° throughout the waggle run for all 3 species.

Thus, followers preferred arranging themselves perpendicular to the dancer, likely

using the same signals associated with this position, to obtain the spatial information

in all 3 species.

Regarding the "follow hypothesis", the pattern of the number of followers in the

posterior position seen in our study provides an argument against it. The number of

followers in the posterior position at the start of the run was not significantly different

from the number of followers in the anterior position and was lower than the number

of followers in the lateral position. Since the entire run encodes spatial information,

the number of followers in the posterior position should have been high throughout

the dance if following from this position was important for the information transfer.

Similar to previous studies, the number of followers in the posterior position increased

as the run progressed [32,62], certainly a direct consequence of the dancers forward

movement during the run [32,61]. As the dancer moves forward, followers are shifted

from the lateral to the posterior position. However, there was no decrease in the

number of followers in the lateral position in the middle and the end of the waggle run

in our observations. This suggests that either some of the followers are able to actively

maintain their lateral positions or that vacated positions to the lateral side of the

dancer are immediately occupied. Both possibilities would support the idea that the

lateral position is more important than the posterior.

If the "tactile hypothesis" is correct, tactile cues associated with the lateral position

around the dancer are the mechanism by which spatial information is transferred

during the waggle dance. Dance followers who are laterally positioned experience a

regular pattern of antennal deflections which correlate strongly with the number of

abdomen waggles [32,62,63]. Since the frequency of waggling of the abdomen is

physically constrained to be similar amongst bees of the same species [64], followers

can use this to estimate the duration of the waggle phase. At the same time, the dance

followers can obtain the orientation of the waggle run by using their own body

position with respect to gravity as a reference. The Johnston's organ may play a major

role in sensing information about the direction of the waggle phase and hence the

direction of the food source [65–67]. The similarity in dance follower behaviour across

four species of the genus *Apis* [this study, 34] suggests that the mechanism of spatial
information transfer in the waggle dance is likely through these tactile cues.

To further substantiate the "tactile hypothesis", detailed high-speed video recordings
of the antennal contacts between the dancers and the followers in all 3 species during
the waggle run would be needed. In addition, the follower's flight patterns after
exiting the hive should also be observed to identify whether the information is
transmitted. Even though our study provides strong support for the tactile hypothesis,
we can not completely rule out the possibility that followers can also obtain relevant
information from orienting themselves behind the dancer [68]. Recent studies which
have focused on the information transfer during the dance show that the transfer does
not depend on the follower position around the dancer [36,68,69]. However, these
studies only quantified the number of followers in the various zones around the
dancer and did not compare the mean body orientation of the followers while
following the dance. Combining detailed observations of the follower behaviour using
a high-speed camera with tracking of their foraging trips [31] is essential to gain a
better understanding of the mechanism underlying spatial information transfer in the
waggle dance and confirm the tactile hypothesis.

Figure 4

Signals produced by the dancer in different honey bee species. Signals can either attract followers to the dancer or contain information. While the former is expected to be different amongst species depending on nesting and foraging conditions, the latter should be highly conserved amongst the species. Signals with an asterisk (*) next to them have only been studied in *A. mellifera* so far. Visual and acoustic signals are only present in some species, olfactory signals can not contain any spatial information and vibrational signals are not expected to play a role in the open nesting species as the dances often happen over other bees [37]. Tactile signals are the most likely to contain spatial information about the food source in all the closely related species of the genus *Apis.*

Differences in other signals associated with the waggle dance in the various *Apis* species is linked to the modality best suited to attract followers to the dancer according to the nest environment of the species (fig 4). In our study, the two open nesting species, *A. florea* and *A. dorsata*, had higher numbers of dance followers throughout the waggle run as compared to the cavity nesting *A. cerana*. Additionally, the median body orientation did not change significantly throughout the run in the case of *A. dorsata*, while it increased from the start to the end of the run in both *A. florea* and *A. cerana*. Dancers in *A. dorsata* produce both a visual and an acoustic signal, while dancers in *A. florea* and *A. cerana* are known to use only one additional signal modality, visual and acoustic respectively [39,43,70]. Further work will be needed to tease out the exact modulatory effect of the additional signals in the waggle dance in these species.

In conclusion, the behavioural responses of the followers to the differing dance signals in Asian honey bees provides new insights into the evolution of this complex communication system. The difficulty in incorporating a mechanism for the transfer of spatial information into any communication system is evident from the fact that such a mechanism has evolved only rarely in animals [12]. The symbolic communication of navigational information in *Apis* likely evolved on the basis of a less complicated and non-error prone modulatory communication as seen in the closely related stingless bees and bumblebees [37–39,71]. Given the difficulty in evolving such a symbolic communication system, it can be expected that there will be very little variation in the mechanism for information transfer within a small group of closely related species. In line with this, we found that the behaviour of the dance followers, who receive the

spatial information, is highly conserved across the genus. Additional signals in the

waggle dance of the different species may be involved in attracting followers to the

dancer (fig 4). Thus, our study highlights the usefulness of comparative studies to

understand complex communication systems like the honey bee waggle dance and

provides a foundation for future studies exploring the dancer and follower behaviour

in the genus *Apis*.

Data Accessibility

557 All the raw data obtained from these experiments is available online at Figshare.

Funding

559 E.A.G. was supported by the NCBS Graduate school. N.T. was supported by ICAR-JRF

560 (PGS). A.B. was supported by National Centre for Biological Sciences – Tata Institute

561 of Fundamental Research institutional funds No. 12P4167.

Author Contributions

E.A.G. participated in the conception and design of the study, participated in the

experiments, performed the data analysis and drafted the manuscript. S. P.

participated in the video and data analysis and critically revised the manuscript. N. T.

participated in the experiments and the video analysis and critically revised the

manuscript. A. B. participated in the conception and design of the study, coordinated

the study and helped draft the manuscript. !6P2B%z^#7qg

Acknowledgments

570 We would like to thank Patrick Kohl and Benjamin Rutschmann from the University

571 of Würzburg for help with the experiments.

References

1. Searcy WA, Nowicki S. 2005 *The Evolution of Animal Communication*. New Jersey, U.S.A:
Princeton University Press. See
https://press.princeton.edu/books/paperback/9780691070957/the-evolution-of-animal-
communication.

2. Conradt L, Roper TJ. 2003 Group decision-making in animals. *Nature* **421**, 155–158.
(doi:10.1038/nature01294)

3. Chittka L, Leadbeater E. 2005 Social learning: Public information in insects. *Curr. Biol.* **15**, 869–
871. (doi:10.1016/j.cub.2005.10.018)

4. Freeberg TM, Dunbar RIM, Ord TJ. 2012 Social complexity as a proximate and ultimate factor in
communicative complexity. *Philos. Trans. R. Soc. B Biol. Sci.* **367**, 1785–1801.
(doi:10.1098/rstb.2011.0213)

5. Ord TJ, Garcia-Porta J. 2012 Is sociality required for the evolution of communicative complexity?
Evidence weighed against alternative hypotheses in diverse taxonomic groups. *Philos. Trans. R.*
Soc. B Biol. Sci. **367**, 1811–1828. (doi:10.1098/rstb.2011.0215)

6. Czaczkes TJ, Grüter C, Ratnieks FLW. 2015 Trail Pheromones: An Integrative View of Their Role in
Social Insect Colony Organization. *Annu. Rev. Entomol.* **60**, 581–599. (doi:10.1146/annurev-ento-
010814-020627)

7. Mitoyen C, Quigley C, Fusani L. 2019 Evolution and function of multimodal courtship displays.
Ethology **125**, 503–515. (doi:10.1111/eth.12882)

8. Hölldobler B. 1999 Multimodal signals in ant communication. *J. Comp. Physiol. A* **184**, 129–141.
(doi:10.1007/s003590050313)

9. Andersson M. 1994 *Sexual Selection*. New Jersey, U.S.A: Princeton University Press. See
https://press.princeton.edu/books/paperback/9780691000572/sexual-selection.

10. Ligon RA, Diaz CD, Morano JL, Troscianko J, Stevens M, Moskeland A, Laman TG, Iii ES. 2018
Evolution of correlated complexity in the radically different courtship signals of birds-of-
paradise. *PLOS Biol.* **16**, e2006962. (doi:10.1371/journal.pbio.2006962)

11. Ord TJ, Stamps JA, Losos JB. 2010 Adaptation and Plasticity of Animal Communication in
Fluctuating Environments. *Evolution* **64**, 3134–3148. (doi:10.1111/j.1558-5646.2010.01056.x)

12. von Frisch K. 1967 *The dance language and orientation of bees*. Harvard University Press.

13. Jarau Stefan, Hrncir Michael. 2009 *Food Exploitation by Social Insects : Ecological, Behavioral,*
and Theoretical Approaches. Taylor & Francis.

14. Seeley TD. 1995 *The Wisdom of the Hive*.

15. Seeley TD, Mikheyev AS, Pagano GJ. 2000 Dancing bees tune both duration and rate of waggle-
run production in relation to nectar-source profitability. *J. Comp. Physiol. - Sens. Neural Behav.*
Physiol. **186**, 813–819. (doi:10.1007/s003590000134)

16. Gardner KE, Seeley TD, Calderone NW. 2008 Do honeybees have two discrete dances to
advertise food sources? *Anim. Behav.* **75**, 1291–1300. (doi:10.1016/j.anbehav.2007.09.032)

17. De Marco RJ, Farina WM. 2001 Changes in food source profitability affect the trophallactic and
dance behavior of forager honeybees (Apis mellifera L.). *Behav. Ecol. Sociobiol.* **50**, 441–449.
(doi:10.1007/s002650100382)

18. Seeley TD. 1989 Social foraging in honey bees: how nectar foragers assess their colony's
nutritional status. *Behav. Ecol. Sociobiol.* **24**, 181–199. (doi:10.1007/BF00292101)

19. George EA, Brockmann A. 2019 Social modulation of individual differences in dance
communication in honey bees. *Behav. Ecol. Sociobiol.* **73**, 41. (doi:10.1007/s00265-019-2649-0)

20. Grüter C, Farina WM. 2009 Past experiences affect interaction patterns among foragers and
hive-mates in honeybees. *Ethology* **115**, 790–797. (doi:10.1111/j.1439-0310.2009.01670.x)

21. Farina WM, Núñez JA. 1991 Trophallaxis in the honeybee, Apis mellifera (L.) as related to the
profitability of food sources. *Anim. Behav.* **42**, 389–394. (doi:10.1016/S0003-3472(05)80037-5)

22. De Marco RJ. 2006 How bees tune their dancing according to their colony's nectar influx: re-
examining the role of the food-receivers' `eagerness'. *J. Exp. Biol.* **209**, 421–432.
(doi:10.1242/jeb.02025)

23. Nieh JC. 2010 A negative feedback signal that is triggered by peril curbs honey bee recruitment.
Curr. Biol. **20**, 310–5. (doi:10.1016/j.cub.2009.12.060)

24. Sen Sarma M, Fuchs S, Werber C, Tautz J. 2002 Worker piping triggers hissing for coordinated
colony defence in the dwarf honeybee Apis florea. *Zoology* **105**, 215–223.

25. Tan K, Dong S, Li X, Liu X, Wang C, Li J, Nieh JC. 2016 Honey Bee Inhibitory Signaling Is Tuned to
Threat Severity and Can Act as a Colony Alarm Signal. *PLOS Biol.* **14**, e1002423.
(doi:10.1371/journal.pbio.1002423)

26. Thom C. 2003 The tremble dance of honey bees can be caused by hive-external foraging
experience. *J. Exp. Biol.* **206**, 2111–2116. (doi:10.1242/jeb.00398)

27. Seeley TD, Towne WF. 1992 Tactics of dance choice in honey bees: do foragers compare dances?
Behav. Ecol. Sociobiol. **30**, 59–69. (doi:10.1007/BF00168595)

28. Landgraf T, Bierbach D, Kirbach A, Cusing R, Oertel M, Mar RO. 2018 Dancing Honey bee Robot
Elicits Dance-Following and Recruits Foragers. *ArXiv* , 1–25.

29. Michelsen A, Andersen B. 1992 How honeybees perceive communication dances, studied by
means of a mechanical model. *Behav. Ecol. Sociobiol.* **30**, 143–150.

30. Chatterjee A, George EA, Prabhudev MV, Basu P, Brockmann A. 2019 Honey bees flexibly use
two navigational memories when updating dance distance information. *J. Exp. Biol.* , jeb.195099.
(doi:10.1242/jeb.195099)

31. Menzel R *et al.* 2011 A Common Frame of Reference for Learned and Communicated Vectors in
Honeybee Navigation. *Curr. Biol.* **21**, 645–650. (doi:10.1016/j.cub.2011.02.039)

32. Gil M, De Marco RJ. 2010 Decoding information in the honeybee dance: Revisiting the tactile
hypothesis. *Anim. Behav.* **80**, 887–894. (doi:10.1016/j.anbehav.2010.08.012)

33. Božič J, Valentinčič T. 1991 Attendants and followers of honey bee waggle dances. *J. Apic. Res.*
30, 125–131. (doi:10.1080/00218839.1991.11101246)

34. Michelsen A. 2003 Signals and flexibility in the dance communication of honeybees. *J. Comp.*
Physiol. A **189**, 165–174. (doi:10.1007/s00359-003-0398-y)

35. Michelsen A, Kirchner WH, Lindauer M. 1986 Sound and vibrational signals in the dance
language of the honeybee, Apis mellifera. *Behav. Ecol. Sociobiol.* , 207–212.

36. Judd TM. 1994 The waggle dance of the honey bee: which bees following a dancer successfully
acquire the information? *J. Insect Behav.* **8**, 343–354. (doi:10.1007/BF01989363)

37. Dyer FC. 2002 The biology of the dance language. *Annu. Rev. Entomol.* **47**, 917–949.
(doi:10.1146/annurev.ento.47.091201.145306)

38. Barron AB, Plath JA. 2017 The evolution of honey bee dance communication: a mechanistic
perspective. *J. Exp. Biol.* **220**, 4339–4346. (doi:10.1242/jeb.142778)

39. I'Anson Price R, Grüter C. 2015 Why, when and where did honey bee dance communication
evolve? *Front. Ecol. Evol.* **3**, 1–7. (doi:10.3389/fevo.2015.00125)

40. Tautz J, Casas J, Sandeman D. 2001 Phase reversal of vibratory signals in honeycomb may assist
dancing honeybees to attract their audience. *J. Exp. Biol.* **204**, 3737–3746.

41. Towne WF. 1985 Acoustic and visual cues in the dances of four honey bee species. *Behav. Ecol.*
Sociobiol. **16**, 185–187. (doi:10.1007/BF00295154)

42. Lindauer M. 1956 Über die verständigung bei indischen bienen. *Z. Für Vgl. Physiol.* **38**, 521–557.
(doi:10.1007/BF00341108)

43. Dyer FC. 1985 Mechanisms of dance orientation in the Asian honey bee Apis florea L. *J. Comp.*
Physiol. A , 183–198.

44. Kirchner WH, Dreller C, Grasser A, Baidya D. 1996 The silent dances of the Himalayan honeybee,
Apis laboriosa. *Apidologie* **27**, 331–339. (doi:10.1051/apido:19960501)

45. Kirchner WH, Dreller C. 1993 Acoustical Signals in the Dance Language of the Giant Honeybee ,
Apis dorsata. *Behav. Ecol. Sociobiol.* **33**, 67–72.

46. Dyer FC, Seeley TD. 1991 Dance Dialects and Foraging Range in 3 Asian Honey-Bee Species.
Behav. Ecol. Sociobiol. **28**, 227–233. (doi:10.1007/BF00175094)

47. Sen Sarma M, Esch HE, Tautz J. 2004 A comparison of the dance language in Apis mellifera
carnica and Apis florea reveals striking similarities. *J. Comp. Physiol. A* **190**, 49–53.
(doi:10.1007/s00359-003-0470-7)

48. Johnson NL, Kemp AW, Kotz S. 2005 *Univariate Discrete Distributions*. Third. Hoboken, NJ, USA:
John Wiley & Sons, Inc. (doi:10.1002/0471715816)

49. Wagenmakers E-J, Farrell S. 2004 AIC model selection using Akaike weights. *Psychon. Bull. Rev.*
11, 192–196. (doi:10.3758/BF03206482)

50. Watson GS, Williams EJ. 1956 On the Construction of Significance Tests on the Circle and the
Sphere. *Biometrika* **43**, 344. (doi:10.2307/2332913)

51. Pewsey A. 2002 Testing circular symmetry. *Can. J. Stat.* **30**, 591–600. (doi:10.2307/3316098)

52. Fisher NI. 1993 *Statistical Analysis of Circular Data*. Cambridge: Cambridge University Press.
(doi:10.1017/CBO9780511564345)

53. Pewsey Arthur, Neuhäuser Markus, Ruxton GD. 2013 *Circular Statistics in R*. Oxford University
Press.

54. R Core Team. 2018 R: A Language and Environment for Statistical Computing.

55. RStudio Team. 2016 RStudio: Integrated Development Environment for R.

56. Brooks M E, Kristensen K, Benthem K J ,van, Magnusson A, Berg C W, Nielsen A, Skaug H J,
Mächler M, Bolker B M. 2017 glmmTMB Balances Speed and Flexibility Among Packages for
Zero-inflated Generalized Linear Mixed Modeling. *R J.* **9**, 378. (doi:10.32614/RJ-2017-066)

57. Barton K. 2017 MuMIn: Multi-Model Inference.

58. Hartig F. 2018 DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed) Regression
Models.

59. Lenth R. 2019 emmeans: Estimated Marginal Means, aka Least-Squares Means.

60. Agostinelli C, Lund U. 2017 R package 'circular' : Circular Statistics (version 0.4-93).

61. Lam C, Li Y, Landgraf T, Nieh J. 2017 Dancing attraction: followers of honey bee tremble and
waggle dances exhibit similar behaviors. *Biol. Open* **6**, 810–817. (doi:10.1242/bio.025445)

62. Rohrseitz K, Tautz J. 1999 Honey bee dance communication: waggle run direction coded in
antennal contacts? *J. Comp. Physiol. [A]* **184**, 463–470. (doi:10.1007/s003590050346)

63. Michelsen A. 2012 How Do Honey Bees Obtain Information About Direction by Following
Dances? In *Honeybee Neurobiology and Behavior*, pp. 65–76. Dordrecht: Springer Netherlands.
(doi:10.1007/978-94-007-2099-2_6)

64. Hrncir M, Maia-Silva C, Mc Cabe SI, Farina WM. 2011 The recruiter's excitement - features of
thoracic vibrations during the honey bee's waggle dance related to food source profitability. *J.
Exp. Biol.* **214**, 4055–4064. (doi:10.1242/jeb.063149)

65. Ai H, Kai K, Kumaraswamy A, Ikeno H, Wachtler T. 2017 Interneurons in the Honeybee Primary
Auditory Center Responding to Waggle Dance-Like Vibration Pulses. *J. Neurosci.* **37**, 10624–
10635. (doi:10.1523/jneurosci.0044-17.2017)

66. Ai H, Okada R, Sakura M, Wachtler T, Ikeno H. 2019 Neuroethology of the Waggle Dance: How
Followers Interact with the Waggle Dancer and Detect Spatial Information. *Insects* **10**, 336.
(doi:10.3390/insects10100336)

67. Brockmann A, Robinson GE. 2007 Central projections of sensory systems involved in honey bee
dance language communication. *Brain. Behav. Evol.* **70**, 125–36. (doi:10.1159/000102974)

68. Tanner D, Visscher K. 2008 Does the body orientation of waggle dance followers affect the
accuracy of recruitment? *Apidologie* **40**, 55–62. (doi:10.1051/apido:2008074)

69. Kietzman PM, Visscher PK. 2019 Follower Position Does Not Affect Waggle Dance Information
Transfer. *Psyche N. Y.* **2019**, 8–13. (doi:10.1155/2019/4939120)

70. Beekman M, Makinson JC, Couvillon MJ, Preece K, Schaerf TM. 2015 Honeybee linguistics - a
comparative analysis of the waggle dance among species of Apis. *Front. Ecol. Evol.* **3**, 1–11.
(doi:10.3389/fevo.2015.00011)

71. Dornhaus A, Chittka L. 2001 Food alert in bumblebees (Bombus terrestris): possible mechanisms
and evolutionary implications. *Behav. Ecol. Sociobiol.* **50**, 570–576.

725