A geometric basis for surface habitat complexity and biodiversity

Damaris Torres-Pulliza[1,2], Maria A. Dornelas[3], Oscar Pizarro[4], Michael Bewley[4], Shane A.

Blowes[5,6], Nader Boutros[4], Viviana Brambilla[3], Tory J. Chase[7], Grace Frank[7], Ariell Friedman[4,8],

Mia O. Hoogenboom[7], Stefan Williams[4], Kyle J. A. Zawada[3], Joshua S. Madin[1]*

[1]Hawai'i Institute of Marine Biology, University of Hawai'i, Kaneohe, HI, United States.

[2]Department of Biological Sciences, Macquarie University, Sydney, NSW, Australia.

[3]Centre for Biological Diversity, Scottish Oceans Institute, University of St Andrews, St Andrews

KY16 9TH, UK.

[4]Australian Centre for Field Robotics, University of Sydney, Sydney, NSW, Australia.

[5]German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Deutscher Platz

5e, Leipzig 04103, Germany.

[6]Department of Computer Science, Martin Luther University Halle-Wittenberg, Am Kirchtor 1,

Halle (Salle) 06108, Germany.

[7]ARC Centre of Excellence for Coral Reef Studies and College of Science and Engineering, James

Cook University, Townsville, Queensland 4811, Australia.

[8]Greybits Engineering, Sydney, NSW, Australia.

*Correspondence to: jmadin@hawaii.edu.

Abstract

Structurally complex habitats tend to contain more species and higher total abundances than simple habitats. This ecological paradigm is grounded in first principles: species richness scales with area, and surface area and niche density increase with three-dimensional complexity. Here we present a geometric basis for surface habitats that unifies ecosystems and spatial scales. The theory is framed by fundamental geometric constraints among three structure descriptors—surface height, rugosity and fractal dimension—and explains 98% of surface variation in a structurally complex test system: coral reefs. We then show how coral biodiversity metrics (species richness, total abundance and probability of interspecific encounter) vary over the theoretical structure descriptor plane, demonstrating the value of the theory for predicting the consequences of natural and human modifications of surface structure.

Main text

Most habitats on the planet are surface habitats—from the abyssal trenches to the tops of mountains, from coral reefs to the tundra. These habitats exhibit a broad range of structural complexities, from relatively simple, planar surfaces to highly complex three-dimensional structures. Currently, human and natural disturbances are changing the complexity of habitats faster than at any time in history[1-4]. Therefore, understanding and predicting the effects of habitat complexity changes on biodiversity is of paramount importance[5]. However, empirical relationships between commonly-used descriptors of structural complexity and biodiversity are

variable, often weak or contrary to expectation[6-10]. Moreover, there are no standards for

quantifying structural complexity, precluding general patterns in the relationship between

structure and diversity from being identified in different habitats. We therefore propose a new

geometric basis for surface habitats that integrates and standardises existing surface

descriptors[8,10].

In theory, species richness scales with surface area according to a power law[11]. Island

biogeography theory articulates that this relationship arises out of extinction and colonization,

as larger areas provide larger targets for species to colonize and a greater variety of habitats

allowing species to coexist[12]. Our geometric theory builds on these ideas by exploring the

notion that habitat surfaces with the same rugosity (defined here as surface area per planar

area) can exhibit a range of different forms (Fig. 1). Total surface area is the integration of

component areas at the smallest scale (i.e., resolution), but it does not explain how these

component areas fold and fill the three-dimensional spaces they occupy. Rather, fractal

dimension quantifies space-filling at different scales[13]. Space-filling promotes species co-

existence by dividing surface area into a greater variety of structural elements[14], microhabitats

and niches[15] (e.g., high and low irradiance; small and large spaces; fast and slow flow). This

variety of niches allows species to coexist (e.g. different competitors, or predator and prey[16])

and therefore enhances biodiversity[17,18]. We posit that there is a fundamental geometric

constraint between surface rugosity and fractal dimension: for a given surface rugosity, an

increase in fractal dimension will result in a reduction of the surface's mean height (Fig. 1). As

the basis for a geometric theory, we mathematically derived the trade-off between surface

rugosity (R), fractal dimension (D) and surface height range (ΔH) as (see Methods for

derivation):

$\frac{1}{2} log(R^2 - 1) + log\left(\frac{L}{L_0}\right)(3 - D) = log\left(\frac{\Delta H}{\sqrt{2}L_0}\right)$ Eq. 1

Where L is the surface linear extent and L_0 is the resolution (i.e., the smallest scale of

observation). R and D are both dimensionless, with $R \geq 1$ and $2 \leq D \leq 3$; ΔH is dimensionless

when standardised by resolution L_0, with $\frac{\Delta H}{\sqrt{2}L_0} \geq 0$. When rugosity is expressed as R^2-1 (with R^2-1

≥ 0) and height range as $\frac{\Delta H}{\sqrt{2}L_0}$, Eq. 1 is a plane equation. Moreover, it is clear that any one of the

surface descriptors can easily be expressed in terms of the other two, highlighting that any of

the three variables is required, but not sufficient alone, to describe the structural complexity of

a surface habitat.

76   Results

To test the theory, we examined associations among surface rugosity, fractal dimension and

height range across coral reef habitat patches. Coral reefs are ideal ecosystems for testing a

theory of surface habitats, because they are structurally complex surface habitats constructed

in large part by the reef-building scleractinian corals that, in turn, live upon the habitat (i.e.,

corals are autogenic ecosystem engineers[19]). Structural complexity affects biodiversity in

general[20] and of coral reefs in particular[21]. Using Structure from Motion (SfM), we estimated

surface rugosity (expressed as the *log* of R^2-1), fractal dimension (D) and height range (as the

log of $\frac{\Delta H}{\sqrt{2}L_0}$) from digital elevation models (DEMs) for 591 reef patches of 4 m^2 at 21 reef sites

encircling Lizard Island on the Great Barrier Reef, Australia (see Methods). Analyses of the

structure of these patches reveal that while rugosity, fractal dimension and surface height

range are not independent, they have substantial independent variation (r^2 for pairwise

relationships between surface descriptors ranging between 3% and 30%, Fig. 2a-c). However,

when framed together, the three descriptors formed a plane, whereupon the trivially measured

surface descriptors, rugosity and height range, captured 98% of the variation in *D* (Fig. 2d). The

remaining 2% of the variation occurs because real surfaces do not necessarily behave like

fractals (i.e., are self-similar) across a wide range of scales (Extended Data Fig. 5). The

observation that the structure of nearly all measured reef patches fell upon a plane delineated

by three simple surface descriptors highlights the fundamental geometric constraints of surface

habitats. If fractal dimension increases, then either rugosity increases, or height range

decreases, or both. All three descriptors are essential for capturing structural complexity

because they explain different elements of surface geometry: height range captures patch scale

variation, rugosity captures fine scale variation (which sums to surface area), and fractal

dimension captures degree of space filling when transitioning from broad to fine scales

(Extended Data Fig. 1a).

Different reef locations, with different ecological and environmental histories, occupied

different regions on the surface descriptor plane (Fig. 3). For example, one site that was

stripped of living coral during back-to-back tropical cyclones[22] largely occupied the region of the

plane where rugosity, fractal dimension and surface height range are all low (Fig. 3a); that is,

the patches at this site were closest to a theoretical flat surface. Another site also impacted by

the cyclones but left littered with dead coral branches, had similar levels of rugosity and height

range, but fractal dimension was relatively high (Fig. 3b). In contrast, a site containing several

large colonies of living branching coral had patches with the highest fractal dimension and

rugosity, yet the height range of these patches was low (Fig. 3c) reflecting the approximately

uniform height of living branching corals in shallow waters where water depth and tidal range

constrains colony growth. Meanwhile, a site containing large hemispherical *Porites* corals had

patches with large height ranges and high rugosity but lower fractal dimension (Fig. 3d). Three

sites contained patches with similar distributions of rugosities (Fig. 3b,d,f), and therefore similar

surface areas. However, these sites ranged from smooth reef surfaces with large holes (Fig. 3e)

to highly bumpy surfaces with no holes (Fig. 3b), demonstrating why rugosity alone does not

capture structural complexity and how varying mixtures of structural components dictate

habitat complexity[14].

Finally, to connect the geometric variables to biodiversity, we examined how species richness,

total abundance and diversity (measured as the probability of interspecific encounter[23]) varied

across the surface descriptor plane. Strong ecological feedbacks occur between coral reef

habitat structure and coral biodiversity metrics. Coral reef structures are largely created by

corals, but their structure is mechanistically affected by environmental conditions such as tidal

range, currents, storm impacts and wave exposure. For instance, coral larvae are poor

swimmers and are more likely to settle in reef patches with small-scale complexity, because

they get entrapped by micro-eddies[24]. At the same time, more intricate coral structures (with

higher fractal dimension, *D*) are more likely to be damaged or dislodged during storms that

flatten reef patches[25,26]. Species-area theory predicts that species richness and abundances

should be highest in patches with the greatest surface area[11] (i.e., highest rugosity). We

predicted that higher fractal dimension would also enhance species richness and abundance,

because of niche diversity (i.e., increases in surface area at different scales), and that this effect

would be additional to overall surface area. The surface descriptor plane allows estimating the

combined effects of not just area, but also niche differentiation associated with fractal

dimension and height range[10,15].

We examined geometric-biodiversity coupling for a large plot, containing 261 of the 4 m^2 reef

patches, in which 9,264 coral colonies of 171 species were recorded (see Methods). Contrary to

expectation, we found that all biodiversity metrics considered peaked in reef patches with

intermediate surface rugosities (Fig. 4a shows diversity, and Extended Data Table 2 includes

species richness and abundance). Indeed, several recent studies have argued that the

relationship should be unimodal because, as complexity increases, the amount of area available

for individuals to live declines[27,28]. However, biodiversity metrics also tended to increase

monotonically in association with patches with higher fractal dimension and smaller height

range (Extended Data Fig. 7). The explanatory power of reef geometry on biodiversity metrics

was over 50% (Extended Data Table 1)—5 to 45% higher than any surface descriptor alone.

Explaining this much variation in biodiversity is striking, given the number of other, non-

geometric processes that govern coral biodiversity, including environmental filtering, dispersal

and species interactions[29]. Because corals are autogenic ecosystem engineers, reciprocal

causality is likely to strengthen and shape geometric-biodiversity coupling. For instance, high

rugosity is often generated by large hemispherical corals (e.g., Fig. 3d) that reduce the number

of individuals, and hence species, per area[14]. Subsequently, geometric-biodiversity coupling

may be weaker for other surface-associated taxa, such as fishes and invertebrates, and should

be tested. Nonetheless, our findings have implications for resilience following disturbances and

for restoration efforts that aim to maximise biodiversity[30], specifically identifying the reef

structural characteristics that should be maintained (or built) to maximize biodiversity.

Discussion

A general, scale-independent geometric basis for surface habitats provides a much-needed way

to quantify habitat complexity across ecosystems and spatial scales. Meanwhile, creating three-

dimensional habitat surfaces is becoming increasingly accessible and cost effective, for example

using Structure from Motion[31,32], both underwater and on land. The importance of surface

complexity as a determinant of habitat condition, biodiversity, and ecosystem function is well

recognised[33], yet different metrics are typically used for different ecosystems, or different taxa

within the same ecosystem[10]. The general quantitative approach we propose is applicable

across surface habitats in both marine and terrestrial environments, allowing formal

comparisons examining whether geometric-biodiversity couplings differ among systems in

terms of both pattern and strength. The surface descriptor plane uncovered here clearly

defines the fundamental geometric constraints acting to shape surface habitats, and

consequently, how changes in surface geometry affect biodiversity. Nonetheless, there remain

several unknowns about the surface descriptor plane and its associations with biodiversity

metrics that require further exploration. These unknowns range from technical limitations (e.g.,

how does the theory translate from digital elevation models that exclude overhanging surfaces

to 3D surface meshes?) to ecological patterns (e.g., how do different types of structural

components, such as different mixtures of branching and hemispherical corals or live and dead

elements[14,34], mediate geometric-biodiversity coupling?).

As powerful ecosystem engineers, humans are modifying the planet through the structures we

destroy, both physically and indirectly via environmental change[4], and those we construct.

Indeed, human-modified structures differ significantly in their geometry from nature-built

structures[35]. Determining how biodiversity, conservation status and recovery rates relate to

habitat complexity measures is paramount in the Anthropocene. The approach we propose

here allows for predictions of the biodiversity consequences of these structural changes across

land and seascapes.

Methods

Geometric theory for surface habitats. The variation method for calculating fractal dimension

D measures the mean height range of a surface at different scales[36,37]. At the broadest scale,

the linear extent L, the surface height range is ΔH (Extended Data Fig. 1a). At the finest scale,

the resolution L_0, the height range (ΔH_0) is the mean of height ranges of all the component

areas at that scale. The slope S of the resulting log-log relationship (shown in Extended Data Fig.

1a) is:

$S = \dfrac{log(\Delta H) - log(\Delta H_0)}{log(L) - log(L_0)}$ Eq. 2

Where fractal dimension is[37]:

$D = 3 - S$ Eq. 3

Rearranging Eq. 2 gives:

$S = \dfrac{log(\Delta H / \Delta H_0)}{log(L / L_0)}$ Eq. 4

Surface area A can be estimated by summing areas A_0 at the finest grain L_0. Given the mean

height range ΔH_0 at L_0, we assume any finer scale detail is not observable, and we calculate A_0

from the minimal surface consistent with ΔH_0 (Extended Data Fig. 1b) as:

$A_0 = \dfrac{L_0}{2}\sqrt{2\Delta H_0^2 + 4L_0^2}$ Eq. 5

And then multiply by the number of component areas ($\frac{L^2}{L_0^2}$) giving:

$A = \dfrac{L^2}{2L_0}\sqrt{2\Delta H_0^2 + 4L_0^2}$ Eq. 6

Surface rugosity is[32]:

$R = \dfrac{A}{L^2}$ Eq. 7

Substituting A for Eq. 6 and rearranging gives:

$R = \sqrt{\dfrac{\Delta H_0^2}{2L_0^2} + 1}$ Eq. 8

Rearranging for ΔH_0 gives:

$\Delta H_0 = \sqrt{2}L_0\sqrt{R^2 - 1}$ Eq. 9

And substituting into Eq. 4 gives:

$S = \dfrac{log\left(\dfrac{\Delta H}{\sqrt{2}L_0\sqrt{R^2 - 1}}\right)}{log\left(\dfrac{L}{L_0}\right)}$ Eq. 10

Leading to:

$$D = 3 - \frac{log\left(\frac{\Delta H}{\sqrt{2}L_0\sqrt{R^2-1}}\right)}{log\left(\frac{L}{L_0}\right)}$$ Eq. 11

Further rearranging gives a plane as Eq. 1 in the main text. The boundaries equations for the

limits of fractal dimension D are:

$\Delta H_{D=2} = \sqrt{2}L\sqrt{R^2-1}$ Eq. 12

$\Delta H_{D=3} = \sqrt{2}L_0\sqrt{R^2-1}$ Eq. 13

Coral reef surface field study. Twenty-one reef flat sites were selected approximately 1 km

apart and encircling Lizard Island on the Great Barrier Reef, Australia (Extended Data Fig. 2). The

spatial arrangement of the sites captured a broad range of habitats that were shaped

predominantly by wave exposure generated by prevailing southeast trade winds[22]. Mean water

depth across all study sites range between 2 to 3.5 meters. In 2014, at the Trimodal site, we

used an Iver2 Autonomous Underwater Vehicle[38] to collect 45,000 georeferenced overlapping,

stereo-pair images of an approximately 30 m by 50 m section of the reef crest (Fig. 4c). In 2016,

at all 21 sites, we used the spiral method[39], which involves swimming a camera rig that

unspools from a central point to capture approximately 3000 overlapping, stereo-pair images of

approximately 130 m^2 of reef crest (Extended Data Fig. 3). We used a simultaneous localisation

and mapping approach[40] fusing GPS, stereo imagery and altitude information to provide an

initial pose estimate for the cameras. We used Agisoft Metashape software to process the

images and produce a 3D dense cloud from which we derived a gridded digital elevation model

(DEM) and orthographic mosaic for coral annotation per site. The output resolution of all DEMs

was 0.002 m. We used DEMs in order to exclude overhanging surfaces (i.e., only one height for

each xy combination), because the degree to which overhangs are captured from plan view

photographic surveys is biased by the changing lighting conditions of the environment[41] (e.g.,

the sun angle, cloud cover, water turbidity, etc.). On the other hand, plan view surveys were

preferred in order to reduce the time costs associated with capturing stereo pairs from multiple

view angles over large areas. The use of DEMs will underestimate surface rugosity and fractal

dimension; i.e., the reason why D tended to range below 2.6. However, given that overhanging

structures were rare at our study sites, R and D measures are likely to exhibit the correct rank

order for patches.

Given the lack of coral cover following the 2016 mass bleaching event on the GBR[22], we used

the 2014 Trimodal large plot to quantify geometric-biodiversity relationships (Extended Data

Fig. 2). The plot was divided into a contiguous grid of 2 by 2 m reef patches (Fig. 4c, black

squares). Patches of the orthographic mosaic were printed on underwater paper and used as

reference maps for *in situ* identification of all coral colonies of diameter >5 cm to species by a

team of six researchers over four weeks. We focused on the reef crest and flat (shallower areas

in Fig. 4c) but also included reef edge and deeper reef. Colonies of unknown or hard to identify

species were photographed and identified in consultation with guide books and other

observers. Hemispherical *Porites* colonies were identified to genus due to the difficulty

differentiating among the few known species without collecting samples for microscopy. Colony

annotations were digitized over the orthographic mosaic using QGIS software (e.g., Fig. 4c,

white points). Only scleractinian corals were included for analyses. In total, 9,264 coral colonies

of 171 species were observed within the 255 reef patches censused. Diversity was calculated as

the probability of interspecific encounter (PIE), or 1 – Simpson diversity[23].

Each of the circular DEMs had a central point, from which an 8 by 8 m square was centred

(Extended Data Fig. 3) and divided into 16 contiguous reef patches of 2 by 2 m. DEMs for each 2

by 2 m patch from both the Trimodal large plot (where corals were censused) and the 21

circular plots were cropped to calculate surface rugosity R, fractal dimension D and the height

range of the patch ΔH; the latter being the difference between the deepest and shallowest

point in a patch. D was calculated using the variation method[37], where each patch was divided

into squares with sides lengths (L) of 2, 1, 0.5, 0.25, 0.125, 0.0625 and 0.03125 m capturing

approximately two orders of magnitude[42]. The resolution L_0 for the theory is the smallest scale

(i.e., 0.03125 m). The height range within each grid at each scale were calculated, and then

averaged for that scale to avoid weighting the many estimates at smaller scales more than the

fewer estimates at larger scales when calculating the slope S. S was calculated for each patch by

fitting a linear model to the log of scale (i.e., grid sizes) versus the log of mean height. D was

then calculated according to Eq. 3. R was calculated according to Eq. 8. There are many ways to

estimate the surface area of a DEM, so we compared surface rugosity calculated from theory

(Eq. 8) with estimates based on surface area calculations using the *surfaceArea* function in the

package *sp*[43]. The theory underestimated surface rugosity by approximately 5% (Extended Data

Fig. 4), because of the minimal area assumption (Extended Data Fig. 1b), but this disparity was

consistent across the range of rugosities.

Analyses. Surface rugosity (expressed as R^2-1) and standardised height range (expressed as

$\frac{\Delta H}{\sqrt{2}L_0}$) were log-transformed (base 10) as per the plane equation (Eq. 1). Species richness and

abundance were sqrt-transformed, and diversity arcsin-transformed, for all analyses to improve

model residuals. Coefficients of determination (r square) of pairwise associations of the three

geometric variables were estimated by squaring Pearson correlation coefficients. Reef surfaces

were not perfectly fractal: mean height ranges at L and L_0 anchor the theory (Extended Data Fig.

1a), but mean height ranges at scales intermediate to L and L_0 could shift the overall

relationship, albeit subtlety. Therefore, we calculated the r square for the surface descriptor

plane based on the deviances of empirically derived D from theory derived D (Extended Data

Fig. 5) (i.e., by dividing the residual sums of squares by the total sums of squares, and then

subtracting this value from one).

We quantified geometric-biodiversity relationships for the large plot at the Trimodal site using

both generalised additive models (GAMs) and linear models (LMs). We applied the default

smoother term to each surface descriptor for the GAMs and second-order polynomials for the

LMs, to allow for non-linear relationships among predictor and response variables. We

quantified the effect of each geometric variable separately on species richness, total abundance

and diversity (PIE), and then all together to assess improvement in explained variation as

adjusted r square values (Extended Data Fig. 7; Extended Data Table 1). We included the three

reef patches with no living coral, but also confirmed that removing these points had no

discernible influence on the geometric-biodiversity relationships. We also ran analyses

following the removal of the 5-6 highest rugosity patches that appeared to be largely

responsible for producing the hump-shaped rugosity-biodiversity relationships (red curves,

Extended Data Fig. 7). Smooth terms for rugosity and height range were significant, with

reference degrees of freedom much greater than one for all biodiversity metrics, suggesting

significant non-linear effects for these surface descriptors[44]. Fractal dimension showed a linear

effect for richness and diversity, and so the smoother term was removed for these analyses.

Residuals for all models were approximately normal and were homogeneous when plotted

against predictor variables. The linear models with second-order polynomial terms gave the

same overall results as GAMs (Extended Data Fig. 7). That is, the polynomial term was

significant for the same terms that retained the smoother function in the GAMs. However, the

LMs had lower adjusted r square values and so we presented the final results using GAMs (Fig.

4a; Extended Data Table 2).

All analyses, including model selection and diagnostics and figure creation, were conducted in

the statistical program language R[45] and can be downloaded or cloned at GitHub

(https://github.com/jmadin/surface_geometry).

Data and code availability

316   Source data and code for data preparation, statistical analyses and figures are available at

317   https://github.com/jmadin/surface_geometry.

References

1. S. L. Pimm, C. N. Jenkins, R. Abell, T. M. Brooks, J. L. Gittleman, L. N. Joppa, P.H. Raven, C. M.

Roberts, J. O. Sexton, The biodiversity of species and their rates of extinction, distribution,

and protection. *Science* **344**, 1246752 (2014).

2. T. Newbold, L. N. Hudson, S. L. L. Hill, S. Contu, I. Lysenko, R. A. Senior, L. Börger, D. J.

Bennett, A. Choimes, B. Collen, J. Day, A. De Palma, S. Díaz, S. Echeverria-Londoño, M. J.

Edgar, A. Feldman, M. Garon, M. L. K. Harrison, T. Alhusseini, D. J. Ingram, Y. Itescu, J.

Kattge, V. Kemp, L. Kirkpatrick, M. Kleyer, D. L. P. Correia, C. D. Martin, S. Meiri, M.

Novosolov, Y. Pan, H. R. P. Phillips, D. W. Purves, A. Robinson, J. Simpson, S. L. Tuck, E.

Weiher, H. J. White, R. M. Ewers, G. M. Mace, J. P. W. Scharlemann, A. Purvis, Global effects

of land use on local terrestrial biodiversity. *Nature* **520**, 45-50 (2015).

3. Alvarez-Filip, L., Dulvy, N. K., Gill, J. A., Côté, I. M. & Watkinson, A. R. Flattening of Caribbean

coral reefs: region-wide declines in architectural complexity. *Proc. R. Soc. B.* **276**, 3019–3025

(2009).

4. Millennium Ecosystem Assessment. 2005. Ecosystems and human well-being: biodiversity

synthesis. World Resources Institute, Washington, D.C., USA.

http://www.unep.org/maweb/en/index.aspx

5. E. D. Schulze, H. A. Mooney, *Biodiversity and Ecosystem Function* (Springer, New York 1993).

6. S. L. Pimm, The complexity and stability of ecosystems. *Nature* **307**, 321–326 (1984).

7. D. R. Morse, J. H. Lawton, M. M. Dodson, M. H. Williamson, Fractal dimension of vegetation

and the distribution of arthropod body lengths. *Nature* **314**, 731–733 (1985).

8. E. D. McCoy, S. S. Bell, Habitat structure: The evolution and diversification of a complex

topic. in *Habitat Structure: The physical arrangement of objects in space* (eds. Bell S.S.,

McCoy E.D. & Mushinsky H.R.) 3 – 27 (Springer, 1991).

9.   M. W. Beck, Separating the elements of habitat structure: independent effects of habitat

complexity and structural components on rocky intertidal gastropods. *J. Exp. Mar. Biol. Ecol.*

249, 29–49 (2000).

10. K. E. Kovalenko, S. M. Thomaz, D. M. Warfe, Habitat complexity: approaches and future

directions. *Hydrobiologia* **685**, 1–17 (2012).

11. O. Arrhenius, Species and Area. *J. Ecol.* **9**, 95-99 (1921).

12. R. H. MacArthur, E. O. Wilson, *The Theory of Island Biogeography* (Princeton Univ. Press,

1967).

13. B. B. Mandelbrot, *The Fractal Geometry of Nature* (W. H. Freeman and Co., 1983).

14. Tokeshi, M. & Arakaki, S. Habitat complexity in aquatic systems: fractals and beyond.

Hydrobiologia **685**, 27–47 (2012).

15. M. P. Johnson, N. J. Frost, M. W. J. Mosley, M. F. Roberts, S. J. Hawkins, The area-

independent effects of habitat complexity on biodiversity vary between regions. *Ecol. Lett.*

6, 126–132 (2003).

16. P. Chesson, Mechanisms of maintenance of species diversity. *Annu. Rev. Ecol. Syst.* **31**, 343–

366 (2000).

17. E. R. Pianka, *Evolutionary Ecology* (Harper and Row, 1988).

18. G. Sugihara, R. M. May, Applications of fractals in ecology. *Trends Ecol. Evol.* **5**, 79–86

(1990).

19. C. G. Jones, J. H. Lawton, M. Shachak, Positive and negative effects of organisms as physical

ecosystem engineers. *Ecology* **78**, 1946-1957 (1997).

20. J. H. Brown, V. K. Gupta, B. L. Li, B. T. Milne, C. Restrepo, G. B. West, The fractal nature of

nature: power laws, ecological complexity and biodiversity. *Philos. T. R. Soc. B.* **357**, 619–

626 (2002).

21. N. A. J. Graham, K. L. Nash, The importance of structural complexity in coral reef

ecosystems. *Coral Reefs* **32**, 315–326 (2012).

22. J. S. Madin, A. H. Baird, T. C. L. Bridge, S. R. Connolly, K. J. A. Zawada, M. Dornelas,

Cumulative effects of cyclones and bleaching on coral cover and species richness at Lizard

Island. *Mar. Ecol. Prog. Ser.* **604**, 263–268 (2018).

23. Hurlbert, S. H. The Nonconcept of Species Diversity: A Critique and Alternative Parameters.

Ecology **52**, 577–586 (1971).

24. T. Hata, J. S. Madin, V. R. Cumbo, M. Denny, J. Figueiredo, S. Harii, C. J. Thomas, A. H. Baird,

Coral larvae are poor swimmers and require fine-scale reef structure to settle. *Sci. Rep.* **7**,

2249 (2017).

25. J. S. Madin, S. R. Connolly, Ecological consequences of major hydrodynamic disturbances on

coral reefs. *Nature* **444**, 477–480 (2006).

26. L. Alvarez-Filip, J. A. Gill, N. K. Dulvy, A. L. Perry, A. R. Watkinson, I. M. Côté, Drivers of

region-wide declines in architectural complexity on Caribbean reefs. *Coral Reefs* **30**, 1051-

1060 (2011).

27. Allouche, O., Kalyuzhny, M., Moreno-Rueda, G., Pizarro, M. & Kadmon, R. Area-

heterogeneity tradeoff and the diversity of ecological communities. *Proc. Nat. Acad. Sci.*

109, 17495–17500 (2012).

28. Paxton, A. B., Pickering, E. A., Adler, A. M., Taylor, J. C. & Peterson, C. H. Flat and complex

temperate reefs provide similar support for fish: Evidence for a unimodal species-habitat

relationship. *PLoS ONE* **12**, e0183906 (2017).

29. Huston, M. A. Patterns of species diversity on coral reefs. *Ann. Rev. Ecol. Syst.* **16**, 149-177

(1985).

30. Loke, L. H. L., Ladle, R. J., Bouma, T. J. & Todd, P. A. Creating complex habitats for

restoration and reconciliation. *Ecological Engineering* **77**, 307–313 (2015).

31. Young, G. C., Dey, S., Rogers, A. D. & Exton, D. Cost and time-effective method for multi-

scale measures of rugosity, fractal dimension, and vector dispersion from coral reef 3D

models. *PLoS ONE* **12**, e0175341 (2017).

32. A. Friedman, O. Pizarro, S. B. Williams, M. Johnson-Roberson, Multi-scale measures of

rugosity, slope and aspect from benthic stereo image reconstructions. *PLoS ONE* **7**, e50440

(2012).

33. E. Weiher, P. Keddy, *Ecological Assembly Rules: Perspectives, Advances, Retreats.*

(Cambridge Univ. Press, 2004).

34. Bartholomew, A., Diaz, R. & Cicchetti, G. New dimensionless indices of structural habitat

complexity: predicted and actual effects on a predator's foraging success. *Mar. Ecol. Prog.*

Ser. **206**, 45–58 (2000).

35. Strain, E. M. A. *et al.* Eco-engineering urban infrastructure for marine and coastal

biodiversity: Which interventions have the greatest ecological benefit? *Journal of Applied*

Ecology **55**, 426–441 (2018).

36. B. Dubuc, S. W. Zucker, C. Tricot, J. F. Quiniou, D. Wehbi, Evaluating the fractal dimension of

surfaces. *P. Roy. Soc. A-Math. Phy.* **425**, 113–127 (1989).

37. G. Zhou, N. S-N. Lam, A comparison of fractal dimension estimators based on multiple

surface generation algorithms. *Comput. Geosci.* **31**, 1260–1269 (2005).

38. M. Johnson-Roberson, M. Bryson, A. Friedman, O. Pizarro, G. Troni, P. Ozog, J. C.

Henderson, High-resolution underwater robotic vision-based mapping and three-

dimensional reconstruction for archaeology. *J. Field Robot.* **34**, 625-643 (2017).

39. O. Pizarro, A. Friedman, M. Bryson, S. B. Williams, J. Madin, A simple, fast, and repeatable

survey method for underwater visual 3D benthic mapping and monitoring. *Ecol. Evol.* **7**,

1770-1782 (2017).

40. I. Mahon, S. B. Williams, O. Pizarro, M. Johnson-Roberson, Efficient view-based SLAM using

visual loop closures. *IEEE T. Robot.* **24**, 1002-1014 (2008).

41. M. Bryson, R. Ferrari, W. Figueira, O. Pizarro, J. Madin, S. Williams, M. Byrne,

Characterization of measurement errors using structure-from-motion and photogrammetry

to measure marine habitat structural complexity. *Ecol. Evol.* **7**, 5669-5681 (2017).

42. D. G. Zawada, J. C. Brock, A Multiscale Analysis of Coral Reef Topographic Complexity Using

Lidar-Derived Bathymetry. *J. Coastal Res.* **10053**, 6–15 (2009).

43. R. S. Bivand, E. Pebesma, V. Gómez-Rubio, *Applied Spatial Data Analysis with R. Use R!, 2 ed*

(Springer, 2013).

44. S. N. Wood, N. Pya, B. Saefken, Smoothing parameter and model selection for general

smooth models (with discussion). *J. Am. Stat. Assoc.* **111**, 1548-1575 (2016).

45. R Core Team. R: A language and environment for statistical computing. R Foundation for

Statistical Computing, (Vienna, Austria 2019). https://www.R-project.org/.

Acknowledgments

431   We thank Dr. Anne Hoggett and Dr. Lyle Vail of the Lizard Island Research Station for their

432   support.

Funding

435   This work was supported by an Australian Research Council Future Fellowship (JM), the John

436   Templeton Foundation (MD, JM), a Royal Society research grant and a Leverhulme fellowship

437   (MD), an International Macquarie University Research Excellence Scholarship (DTP), two Ian

438   Potter Doctoral Fellowships at Lizard Island (DTP and VB), and an Australian Endeavour

439   Scholarship (TC).

Author contributions

The study was conceptualized by JSM, DTP, MD and OP. All authors collected the data. JSM and

OP developed the theory and JSM ran the analyses. JSM, DTP and OP developed the software

pipeline for data and produced the visualizations. The investigation was led by JSM, DTP, MD

and OP. JSM and MD led and funded the broader project, with additional field robotics

resources provided by OP and SW. JSM wrote the first draft of the paper and all authors

reviewed at least one draft.

Competing interests

450   Authors declare no competing interests.

Figure captions

Fig. 1 | Increasing fractal dimension (i.e., space filling) while keeping surface rugosity

constant results in a decline in a surface's mean height range. A two-dimensional

representation of three hypothetical surface habitats with the same surface rugosities (**a**, **b** and

c). That is, the lengths of the lines **a**, **b** and **c** are the same and occur over the same planar

extent (black points). However, line **a** fills less of its two-dimensional space (black rectangle)

than does line **c**, and therefore has a lower fractal dimension.

Fig. 2 | Comparison of the geometric theory with field data. (**a**-**c**) Pairwise relationships

between the descriptors that frame the geometric theory for n=595 reef patches: surface

rugosity (as R^2-1); fractal dimension D; and surface height range (as $\frac{\Delta H}{\sqrt{2}L_0}$). Coefficients of

determination (r^2) show the variance explained in the y-axis variable by the x-axis variable. (**d**)

When combined the three descriptors explain more than 98% of the variation in fractal

dimension D despite reef surfaces not being perfectly fractal (see Methods). Field data are

points, and the surface descriptor plane is coloured by fractal dimension.

Fig. 3 | The geometric diversity of coral reef habitats. Reef patches (*n*=16) from a subset of six sites are superimposed onto a two-dimensional representation of the surface descriptor plane (colour used here to delineate sites). (**a**) North Reef; (**b**) Osprey; (**c**) Lagoon-2; (**d**) Resort; (**e**) South Island; and (**f**) Horseshoe. Patch height range is greater in the top left corner and decreases towards the bottom right corner. The corresponding DEMs and orthographic mosaics show selected patches at each site to help visualise geometric differences.

Fig. 4 |Geometric-biodiversity coupling of coral reef habitats. (**a**) Predicted coral species diversity (represented as probability of interspecific encounter) when plotted upon the surface descriptor plane given by rugosity and fractal dimension (height range is greater in the top left and decreases towards the bottom right, as per Eq. 1). Prediction contours are from the general additive model summarised in Extended Data Table 2. (**b**) A digital elevation model of the large plot with *n*=255 contiguous 2 x 2 m reef patches (black squares) capturing 9,264 coral colony annotations (white points) representing 171 species.

a Diversity (probability of interspecific encounter) $r^2 = 0.524$

Greater height range

Smaller height range

Rugosity ($R^2 - 1$)

D

b