A geometric basis for surface habitat complexity and biodiversity

Damaris Torres-Pulliza[1,2], Maria A. Dornelas[3], Oscar Pizarro[4], Michael Bewley[4], Shane A. Blowes[5,6], Nader Boutros[4], Viviana Brambilla[3], Tory J. Chase[7], Grace Frank[7], Ariell Friedman[4,8], Mia O. Hoogenboom[7], Stefan Williams[4], Kyle J. A. Zawada[3], Joshua S. Madin[1]*

[1]Hawai'i Institute of Marine Biology, University of Hawai'i, Kaneohe, HI, United States.

[2]Department of Biological Sciences, Macquarie University, Sydney, NSW, Australia.

[3]Centre for Biological Diversity, Scottish Oceans Institute, University of St Andrews, St Andrews KY16 9TH, UK.

[4]Australian Centre for Field Robotics, University of Sydney, Sydney, NSW, Australia.

[5]German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Deutscher Platz 5e, Leipzig 04103, Germany.

[6]Department of Computer Science, Martin Luther University Halle-Wittenberg, Am Kirchtor 1, Halle (Salle) 06108, Germany.

[7]ARC Centre of Excellence for Coral Reef Studies and College of Science and Engineering, James Cook University, Townsville, Queensland 4811, Australia.

[8]Greybits Engineering, Sydney, NSW, Australia.

*Correspondence to: jmadin@hawaii.edu.

Abstract

Structurally complex habitats tend to contain more species and higher total abundances than simple habitats. This ecological paradigm is grounded in first principles: species richness scales with area, and surface area and niche density increase with three-dimensional complexity. Here we present a geometric basis for surface habitats that unifies ecosystems and spatial scales. The theory is framed by fundamental geometric constraints among three structure descriptors—surface height, rugosity and fractal dimension—and explains 98% of surface variation in a structurally complex test system: coral reefs. We then show how coral

29   biodiversity metrics vary over the theoretical structure descriptor plane, demonstrating the

30   value of the new theory for predicting the consequences of natural and human modifications of

31   surface structure.

33   **One Sentence Summary**

34   Surface habitats fall upon a plane of geometric possibilities that mediate biodiversity.

Main text

Most habitats on the planet are surface habitats—from the abyssal trenches to the tops of
mountains, from coral reefs to the tundra. These habitats exhibit a broad range of structural
complexities, from relatively simple, planar surfaces to highly complex three-dimensional
structures. Currently, human and natural disturbances are changing the complexity of habitats
faster than at any time in history (*1*, *2*). Therefore, understanding and predicting the effects of
habitat complexity changes on biodiversity is of paramount importance (*3*). However, empirical
relationships between commonly-used descriptors of structural complexity and biodiversity are
variable, often weak or contrary to expectation (*4-8*). Moreover, there are no standards in how
structural complexity is quantified, precluding general patterns in the relationship between
structure and diversity from being identified. We therefore propose a new geometric basis for
surface habitats that integrates and standardises existing surface descriptors (*6*, *8*).

The geometric basis arises from quantifying the distribution of surface area in habitat surfaces.
In theory, species richness scales with surface area according to a power law (*9*). Island
biogeography theory articulates that this relationship arises out of extinction and colonization,
as larger areas provide larger targets for species to colonize and a greater variety of habitats
allowing species to coexist (*10*). Our geometric theory builds on these ideas by exploring the
notion that habitat surfaces with the same total surface (and hence rugosity) can exhibit a
range of different forms (Fig. 1). Total surface area is the integration of component areas at the
smallest scale (i.e., resolution), but it does not explain how these surfaces fold and fill the
three-dimensional spaces they occupy. Rather, fractal dimension quantifies space-filling at
different scales (*11*). Space-filling promotes species co-existence by dividing surface area into a
greater variety of microhabitats and niches (e.g., high and low irradiance; small and large
spaces; fast and slow flow) (*12*). This variety of niches allows species to coexist (e.g. different
competitors, or predator and prey [*13*]) and therefore enhances biodiversity (*14*, *15*). There is a
fundamental geometric constraint between surface rugosity and fractal dimension: for a given
surface rugosity, an increase in fractal dimension will result in a reduction of the surface's mean
height (Fig. 1). As the basis for a geometric theory, we derived the trade-off between surface

rugosity (R), fractal dimension (D) and surface height range (ΔH) as (see Supplementary

Methods for mathematical derivation):

$\frac{1}{2}log(R^2 - 1) + log\left(\frac{L}{L_0}\right)(3 - D) = log\left(\frac{\Delta H}{\sqrt{2}L_0}\right)$ Eq. 1

Where L is the surface extent and L_0 is the resolution (i.e., the smallest scale of observation). R

and D are both dimensionless, with $R \geq 1$ and $2 \leq D \leq 3$; ΔH is dimensionless when standardised

by resolution L_0, with $\frac{\Delta H}{\sqrt{2}L_0} \geq 0$. When rugosity is expressed as R^2-1 (with R^2-1 $\geq$ 0) and height

range as $\frac{\Delta H}{\sqrt{2}L_0}$, Eq. 1 is a plane equation. Moreover, it is clear than any one of the surface

descriptors can easily be expressed in terms of the other two, highlighting that any of the three

variables is required, but not sufficient alone, to describe the structural complexity of a surface

habitat.

To test the theory, we examined associations among surface rugosity, fractal dimension and

height range across coral reef habitat patches. Coral reefs are ideal ecosystems for testing a

theory of surface habitats, because they are structurally complex surface habitats constructed

in large part by the reef-building scleractinian corals that, in turn, live upon the habitat (i.e.,

corals are autogenic ecosystem engineers [16]). Structural complexity affects biodiversity in

general (17) and of coral reefs in particular (18). We estimated surface rugosity (as the *log* of

R^2-1), fractal dimension (D) and height range (as the *log* of $\frac{\Delta H}{\sqrt{2}L_0}$) for 591 reef patches of 4 m^2 at

21 reef sites encircling Lizard Island on the Great Barrier Reef, Australia (see Supplementary

Methods). Analyses of the structure of these patches reveal that while rugosity, fractal

dimension and surface height range are not independent, they have substantial independent

variation (r^2 for pairwise relationships between variables ranging between 3% and 30%, Fig. 2A-

C). However, when framed together, the three variables formed the surface descriptor plane

that captured 98% of the variation in D (Fig. 2D). The remaining 2% of the variation can be

explained by the model assuming perfectly fractal surfaces (Eq. S2), while the digital

representation of natural reef patches often deviated from statistical self-similarity (Fig. S3B).

The observation that the structure of nearly all measured reef patches fell upon a plane

delineated by three simple surface descriptors highlights the fundamental geometric

constraints of surface habitats. All three descriptors are essential for capturing structural

complexity because they explain different elements of surface geometry: height range captures

large scale variation, rugosity captures small scale variation, and fractal dimension captures

topographic entropy when transitioning from large to small scales (Fig. S1A).

Different reef locations, with different ecological and environmental histories, occupied

different regions on the surface descriptor plane (Fig. 3). For example, one site that was

stripped of living coral during back-to-back tropical cyclones (*19*) largely occupied the region of

the plane where rugosity, fractal dimension and surface height range are all low (Fig. 3A); that

is, the patches at this site were closest to a theoretical flat surface. Another site also impacted

by the cyclones but left littered with dead coral branches, had similar levels of rugosity and

height range, but fractal dimension was relatively high (Fig. 3B). In contrast, a site containing

several large colonies of living branching coral had patches with the highest fractal dimension

and rugosity, yet the height range of these patches was low (Fig. 3C) reflecting the

approximately uniform height of living branching corals in shallow waters where water depth

and tidal range constrains colony growth. Meanwhile, a site containing large hemispherical

Porites corals had patches with large height ranges and high rugosity but lower fractal

dimension (Fig. 3D).

Finally, to connect the geometric variables to biodiversity, we examined how species richness

and abundance varied across the surface descriptor plane. Strong ecological feedbacks occur

between coral reef habitat structure and coral biodiversity. Coral reef structures are largely

created by corals, but their structure is mechanistically affected by environmental conditions

such as tidal range, currents, storm impacts and wave exposure. For instance, coral larvae are

poor swimmers and are more likely to settle in reef patches with small-scale complexity,

because they get entrapped by micro-eddies (*20*). At the same time, more intricate coral

structures (with higher fractal dimension, *D*) are more likely to be damaged or uprooted during

storms that flatten reef patches (*21*, *22*). Species-area theory predicts that species richness and

abundances should be highest in patches with the greatest surface area (i.e., highest rugosity)

(*9*). We predicted that higher fractal dimension would also enhance species richness and

abundance, because of niche diversity (i.e., increases in surface area at different scales), and

that this effect would be additional to overall surface area. The surface descriptor plane allows

estimating the combined effects of not just area, but also niche differentiation associated with

fractal dimension and depth range (*8*, *12*).

We examined geometric-biodiversity coupling for a large plot, containing 261 of the 4 m^2 reef

patches, in which 9,264 coral colonies of 171 species were recorded (see Supplementary

Methods). Contrary to expectation, we found that coral species richness and abundance peaked

in reef patches with intermediate surface rugosities (Fig. 4A, B and table S2). Richness and

abundance also tended to be associated with patches with higher fractal dimension and smaller

height range. (The consistency of patterns for both species richness and abundance was not

surprising, because the two biodiversity metrics are strongly associated in our data.) The

explanatory power of reef geometry on both biodiversity metrics was more than 40% (Table

S1)—10 to 15% higher than any surface descriptor alone. The geometric plane explaining this

much variation in biodiversity is striking, given the number of other, non-geometric processes

that govern coral biodiversity, including environmental filtering, dispersal and species

interactions (*23*). Because corals are autogenic ecosystem engineers, reciprocal causality is

likely to strengthen and shape geometric-biodiversity coupling. For instance, high rugosity is

often generated by large hemispherical corals (e.g., Fig. 3D) that reduce the number of

individuals, and hence species, per area. Subsequently, geometric-biodiversity coupling may be

weaker for other surface-associated taxa, such as fishes and invertebrates, and should be

tested. Nonetheless, our findings have implications for resilience following disturbances and for

restoration efforts that aim to maximise biodiversity, specifically identifying the reef structural

characteristics that should be maintained (or built) to maximize biodiversity.

A general, scale-independent geometric basis for surface habitats provides a much-needed way

to quantify habitat complexity across ecosystems and spatial scales. The importance of surface

complexity as a determinant of habitat condition, biodiversity, and ecosystem function is well

recognised (*24*), yet different metrics are typically used for different ecosystems, or different

taxa within the same ecosystem (*8*). The general approach we propose is applicable across

surface habitats in both marine and terrestrial environments, allowing formal comparisons

examining whether geometric-biodiversity couplings differ among systems in terms of both

pattern and strength. The surface descriptor plane uncovered here clearly defines the

fundamental geometric constraints acting to shape surface habitats, and consequently how

changes in surface geometry affect biodiversity. As powerful ecosystem engineers, humans are

modifying the planet, through the structures we destroy, as well as those we construct.

Typically, human-modified structures differ significantly in their geometry from nature-built

structures. Determining biodiversity, conservation, and recovery rates through habitat

complexity measures is paramount in the Anthropocene. The approach we propose here allows

for predictions of the biodiversity consequences of these structural changes across land and

seascapes.

References

1. S. L. Pimm, C. N. Jenkins, R. Abell, T. M. Brooks, J. L. Gittleman, L. N. Joppa, P.H. Raven, C. M.

Roberts, J. O. Sexton, The biodiversity of species and their rates of extinction, distribution,

and protection. *Science* **344**, 1246752 (2014).

2. T. Newbold, L. N. Hudson, S. L. L. Hill, S. Contu, I. Lysenko, R. A. Senior, L. Börger, D. J.

Bennett, A. Choimes, B. Collen, J. Day, A. De Palma, S. Díaz, S. Echeverria-Londoño, M. J.

Edgar, A. Feldman, M. Garon, M. L. K. Harrison, T. Alhusseini, D. J. Ingram, Y. Itescu, J.

Kattge, V. Kemp, L. Kirkpatrick, M. Kleyer, D. L. P. Correia, C. D. Martin, S. Meiri, M.

Novosolov, Y. Pan, H. R. P. Phillips, D. W. Purves, A. Robinson, J. Simpson, S. L. Tuck, E.

Weiher, H. J. White, R. M. Ewers, G. M. Mace, J. P. W. Scharlemann, A. Purvis, Global effects

of land use on local terrestrial biodiversity. *Nature* **520**, 45-50 (2015).

3. E. D. Schulze, H. A. Mooney, *Biodiversity and Ecosystem Function* (Springer, New York 1993).

4. S. L. Pimm, The complexity and stability of ecosystems. *Nature* **307**, 321–326 (1984).

5. D. R. Morse, J. H. Lawton, M. M. Dodson, M. H. Williamson, Fractal dimension of vegetation

and the distribution of arthropod body lengths. *Nature* **314**, 731–733 (1985).

6. E. D. McCoy, S. S. Bell, Habitat structure: The evolution and diversification of a complex

topic. in *Habitat Structure: The physical arrangement of objects in space* (eds. Bell S.S.,

McCoy E.D. & Mushinsky H.R.) 3 – 27 (Springer, 1991).

7.   M. W. Beck, Separating the elements of habitat structure: independent effects of habitat

complexity and structural components on rocky intertidal gastropods. *J. Exp. Mar. Biol. Ecol.*

249, 29–49 (2000).   added

8. K. E. Kovalenko, S. M. Thomaz, D. M. Warfe, Habitat complexity: approaches and future

directions. *Hydrobiologia* **685**, 1–17 (2012).

9. O. Arrhenius, Species and Area. *J. Ecol.* **9**, 95-99 (1921). supplementary

10. R. H. MacArthur, E. O. Wilson, *The Theory of Island Biogeography* (Princeton Univ. Press,

1967).

11. B. B. Mandelbrot, *The Fractal Geometry of Nature* (W. H. Freeman and Co., 1983).

12. M. P. Johnson, N. J. Frost, M. W. J. Mosley, M. F. Roberts, S. J. Hawkins, The area-

independent effects of habitat complexity on biodiversity vary between regions. *Ecol. Lett.*

6, 126–132 (2003). additional

13. P. Chesson, Mechanisms of maintenance of species diversity. *Annu. Rev. Ecol. Syst.* **31**, 343–

366 (2000).

14. E. R. Pianka, *Evolutionary Ecology* (Harper and Row, 1988).

15. G. Sugihara, R. M. May, Applications of fractals in ecology. *Trends Ecol. Evol.* **5**, 79–86

(1990).

16. C. G. Jones, J. H. Lawton, M. Shachak, Positive and negative effects of organisms as physical

ecosystem engineers. *Ecology* **78**, 1946-1957 (1997).

17. J. H. Brown, V. K. Gupta, B. L. Li, B. T. Milne, C. Restrepo, G. B. West, The fractal nature of

nature: power laws, ecological complexity and biodiversity. *Philos. T. R. Soc. B.* **357**, 619–

626 (2002).

18. N. A. J. Graham, K. L. Nash, The importance of structural complexity in coral reef
ecosystems. *Coral Reefs* **32**, 315–326 (2012).

19. J. S. Madin, A. H. Baird, T. C. L. Bridge, S. R. Connolly, K. J. A. Zawada, M. Dornelas,
Cumulative effects of cyclones and bleaching on coral cover and species richness at Lizard
Island. *Mar. Ecol. Prog. Ser.* **604**, 263–268 (2018).

20. T. Hata, J. S. Madin, V. R. Cumbo, M. Denny, J. Figueiredo, S. Harii, C. J. Thomas, A. H. Baird,
Coral larvae are poor swimmers and require fine-scale reef structure to settle. *Sci Rep* **7**,
2249 (2017).

21. J. S. Madin, S. R. Connolly, Ecological consequences of major hydrodynamic disturbances on
coral reefs. *Nature* **444**, 477–480 (2006).

22. L. Alvarez-Filip, J. A. Gill, N. K. Dulvy, A. L. Perry, A. R. Watkinson, I. M. Côté, Drivers of
region-wide declines in architectural complexity on Caribbean reefs. *Coral Reefs* **30**, 1051-
1060 (2011).

23. M. A. Huston, Patterns of species diversity on coral reefs. *Ann Rev Ecol Syst* **16**, 149-177
(1985).

24. E. Weiher, P. Keddy, *Ecological Assembly Rules: Perspectives, Advances, Retreats.*
(Cambridge Univ. Press, 2004).

25. B. Dubuc, S. W. Zucker, C. Tricot, J. F. Quiniou, D. Wehbi, Evaluating the fractal dimension of
surfaces. *P. Roy. Soc. A-Math. Phy* **425**, 113–127 (1989).

26. G. Zhou, N. S-N. Lam, A comparison of fractal dimension estimators based on multiple
surface generation algorithms. *Comput. Geosci.* **31**, 1260–1269 (2005).

27. A. Friedman, O. Pizarro, S. B. Williams, M. Johnson-Roberson, Multi-scale measures of
rugosity, slope and aspect from benthic stereo image reconstructions. *PLoS ONE* **7**, e50440
(2012).

28. M. Johnson-Roberson, M. Bryson, A. Friedman, O. Pizarro, G. Troni, P. Ozog, J. C.
Henderson, High-resolution underwater robotic vision-based mapping and three-
dimensional reconstruction for archaeology. *J. Field Robot.* **34**, 625-643 (2017).

29. O. Pizarro, A. Friedman, M. Bryson, S. B. Williams, J. Madin, A simple, fast, and repeatable

survey method for underwater visual 3D benthic mapping and monitoring. *Ecol. Evol.* **7**,

1770-1782 (2017).

30. I. Mahon, S. B. Williams, O. Pizarro, M. Johnson-Roberson, Efficient view-based SLAM using

visual loop closures. *IEEE T. Robot.* **24**, 1002-1014 (2008).

31. M. Bryson, R. Ferrari, W. Figueira, O. Pizarro, J. Madin, S. Williams, M. Byrne,

Characterization of measurement errors using structure-from-motion and photogrammetry

to measure marine habitat structural complexity. *Ecol. Evol.* **7**, 5669-5681 (2017).

32. D. G. Zawada, J. C. Brock, A Multiscale Analysis of Coral Reef Topographic Complexity Using

Lidar-Derived Bathymetry. *J Costal Res.* **10053**, 6–15 (2009).

33. R. S. Bivand, E. Pebesma, V. Gómez-Rubio, *Applied Spatial Data Analysis with R. Use R!, 2 ed*

(Springer*,* 2013).

34. S. N. Wood, N. Pya, B. Saefken, Smoothing parameter and model selection for general

smooth models (with discussion). *J. Am. Stat. Assoc.* **111**, 1548-1575 (2016).

35. R Core Team. R: A language and environment for statistical computing. R Foundation for

Statistical Computing, (Vienna, Austria 2019). https://www.R-project.org/.

Acknowledgments

We thank the Lizard Island Research Station staff for their support.

Funding

This work was supported by an Australian Research Council Future Fellowship (JM), the John

Templeton Foundation (MD, JM), a Royal Society research grant and a Leverhulme fellowship

(MD), an International Macquarie University Research Excellence Scholarship (DTP), two Ian

Potter Doctoral Fellowships (DTP and VB), and an Australian Endeavour Scholarship (TC).

Author contributions

The study was conceptualized by JSM, DTP, MD and OP. All authors curated the data. JSM and

OP developed the theory and JSM ran the analyses. JSM, DTP and OP developed the software

pipeline for data and produced the visualizations. The investigation was led by JSM, DTP, MD

and OP. JSM and MD lead and fund the broader project, with additional field robotics resources

from OP and SW. JSM wrote the first draft of the paper and all authors reviewed at least one

draft.

Competing interests

Authors declare no competing interests.

Data and materials availability

Source data and code for data preparation, statistical analyses and figures are available at

https://github.com/jmadin/surface_geometry (currently private until published; but access can

be granted if requested for reviewing purposes).

Fig. 1. Increasing fractal dimension (i.e., space filling) while keeping surface rugosity constant

results in a decline in a surface's mean height range. A two-dimensional representation of three

hypothetical surface habitats with the same surface rugosities (**A**, **B** and **C**). That is, the lengths

of the lines **A**, **B** and **C** are the same and occur over the same planar extent (black points).

However, line **A** fills less of its two-dimensional space (black rectangle) than does line **C**, and

therefore has a lower fractal dimension.

Fig. 2. Comparison of the geometric theory with field data. Relationships between the

descriptors that frame the geometric theory for n=595 reef patches: (**A**) surface rugosity (as R^2-

1); (**B**) fractal dimension D; and (**C**) surface height range (as $\frac{\Delta H}{\sqrt{2}L_0}$). Coefficients of determination

(r^2) show the variance explained in the y-axis variable by the x-axis variable. (**D**) When

combined the three descriptors explain more than 98% of the variation in fractal dimension D.

Field data are points, and the surface descriptor plane is coloured by fractal dimension.

Fig. 3. The geometric diversity of coral reef habitats. Reef patches (*n*=16) from a subset of six

sites are superimposed onto a two-dimensional representation of the surface descriptor plane

(colour used here to delineate sites). (**A**) North Reef; (**B**) Osprey; (**C**) Lagoon-2; (**D**) Resort; (**E**)

South Island; and (**F**) Horseshoe. The corresponding DEMs and orthographic mosaics show

selected patches at each site to help visualise geometric differences. Patch height range is

greater in the top left corner and decreases towards the bottom right corner.

Fig. 4. Geometric-biodiversity coupling of coral reef habitats. Predicted coral species (**A**)

richness and (**B**) abundance within reef patches as contours and plotted upon the surface

descriptor plane, where the colour-scale shows the gradient of height range according to Eq. 1.

Prediction contours are from general additive models (Table S2). (**C**) A digital elevation model of

300   the large plot with contiguous reef patch delineations (black squares) and the 9,264 coral

301   colony annotations (white points) capturing 171 species.